Paclitaxel causes *de novo* induction of invasive breast cancer cells by repolarizing tumor-associated macrophages

Madeline Friedman-DeLuca,[1,2,3] George S. Karagiannis,[2,3,4,5,6] Camille L. Duran,[1,2,3] Suryansh Shukla,[1,2,3,5] Jiufeng Li,[1,3] John S. Condeelis,[2,3,5,6,7,8] Maja H. Oktay,*[1,2,3,5,6,7] David Entenberg*[1,2,3,5,6]

[1]Department of Pathology, Albert Einstein College of Medicine/ Montefiore Medical Center, Bronx, NY, USA; [2]Integrated Imaging Program for Cancer Research, Albert Einstein College of Medicine/ Montefiore Medical Center, Bronx, NY, USA; [3]Cancer Dormancy Institute, Albert Einstein College of Medicine/ Montefiore Medical Center, Bronx, NY, USA; [4]Department of Microbiology and Immunology, Albert Einstein College of Medicine/ Montefiore Medical Center, Bronx, NY, USA; [5]Gruss-Lipper Biophotonics Center, Albert Einstein College of Medicine/ Montefiore Medical Center, Bronx, NY, USA; [6]Tumor Microenvironment and Metastasis Program, Albert Einstein College of Medicine/ Montefiore Medical Center, Bronx, NY, USA; [7]Department of Surgery, Albert Einstein College of Medicine/ Montefiore Medical Center, Bronx, NY, USA [8]Department of Cell Biology, Albert Einstein College of Medicine/ Montefiore Medical Center, Bronx, NY, USA.

*Corresponding authors

Keywords: Breast cancer, macrophages, MenaINV, NF-κB, Notch1

Abstract

Background: Metastasis is the leading cause of breast cancer (BC) death, and tumor cells must migrate and invade to metastasize. BC cells that express the pro-metastatic actin regulatory protein MenaINV have an enhanced ability to migrate and intravasate within the primary tumor and extravasate at secondary sites. Though chemotherapy improves patient survival, treatment with paclitaxel leads to upregulation of MenaINV and an increase in metastasis in mice. MenaINV expression can be induced in BC cells through cooperative NF-κB/ Notch1 signaling with macrophages, which are often increased in tumors in response to chemotherapy. MenaINV-expressing cells are also resistant to paclitaxel, begging the question of whether paclitaxel increases MenaINV by *de novo* induction or by selectively killing non-MenaINV-expressing cells. We hypothesized that paclitaxel causes *de novo* MenaINV induction by increasing macrophage-tumor cell NF-κB/ Notch1 signaling. Understanding this pro-metastatic effect of chemotherapy is crucial to refining treatment strategies.

Findings: Paclitaxel-treated tumors expressed significantly more MenaINV than vehicle-treated tumors. This effect was dependent upon both macrophages and NF-κB/ Notch signaling. This indicates that chemotherapy increases MenaINV expression by induction. Mechanistically, paclitaxel causes macrophages to take on a pro-inflammatory phenotype and increase NF-κB/ Notch1 signaling with tumor cells.

Conclusion: Paclitaxel causes *de novo* MenaINV induction. Mechanistically, paclitaxel repolarizes tumor-associated macrophages towards a pro-inflammatory phenotype. These pro-inflammatory macrophages then participate in enhanced NF-κB/ Notch1 signaling with tumor cells, which leads to MenaINV induction in the tumor cells. These results lay the groundwork for novel microenvironment-based therapies to alleviate the pro-metastatic effects of chemotherapy in BC.

Introduction

Breast cancer (BC) is the second leading cause of cancer death among women in the United States, and the majority of these deaths are due to metastasis (1). To metastasize, tumor cells must be able to migrate and intravasate within the primary tumor and extravasate at secondary sites. Tumor cells perform these tasks using invadopodia – actin-rich protrusions with proteolytic activity that degrade surrounding extracellular matrix, forging a path for the cell. In BC cells, the actin regulatory protein MenaINV (Mammalian enabled, invasive variant) supports invadopodium stability and degradative activity (2), making cells more migratory and metastatic. Furthermore, tumor cells that express MenaINV are more sensitive to chemotactic gradients in the tumor microenvironment, which facilitate invadopodium formation and directional migration towards blood vessels where they intravasate (3, 4). Indeed, BC cells that express MenaINV have an enhanced ability to migrate, intravasate, and extravasate (4-6), and high MenaINV expression is associated with poor outcomes in both mouse and human BC (4, 7).

Though chemotherapy improves patient survival, treatment with neoadjuvant paclitaxel increases the proportion of MenaINV-expressing cells in both mouse and human residual breast tumors, as well as lung metastasis in mice (8). MenaINV-expressing cells are resistant to paclitaxel (9), so the observed increase in MenaINV after paclitaxel treatment may be due to the selective killing of non-MenaINV-expressing cells. However, we have also shown that MenaINV is induced in tumor cells via cooperative NF-κB/ Notch1 signaling with macrophages (10). Chemotherapy has been shown to cause a macrophage influx into tumors in response to tissue damage, potentially increasing macrophage-tumor cell NF-κB/ Notch1 signaling and MenaINV induction. This means that, in addition to the selective killing of non-MenaINV-expressing cells, paclitaxel may also cause *de novo* induction of MenaINV in cells that did not previously express it. We hypothesized that paclitaxel induces MenaINV expression by increasing macrophage infiltration and macrophage-tumor cell NF-κB/ Notch1 signaling. Here, we find that paclitaxel

causes macrophages to adopt a pro-inflammatory phenotype that promotes NF-κB/ Notch1

signaling and leads to *de novo* induction of MenaINV in tumor cells.

Results

Paclitaxel causes *de novo* induction of MenaINV

To test whether paclitaxel causes *de novo* MenaINV induction, we treated FVB-PyMT mice

bearing spontaneous breast tumors with paclitaxel and co-administered either DAPT – to prevent

MenaINV induction (10, 11) – or vehicle control (**Fig. 1A**). DAPT is a γ-secretase inhibitor that is

traditionally used to block Notch signaling. However, given the significant crosstalk between the

Notch and NF-κB pathways, we (10) and others (12) have found that DAPT effectively inhibits

NF-κB activation as well. If paclitaxel increases MenaINV only by selection, we would expect that

dual DAPT/ paclitaxel treatment would result in the same increase in MenaINV as treatment with

paclitaxel alone through the selective killing of non-MenaINV-expressing cells.

Following treatment, mice were sacrificed, and their tumors were analyzed for MenaINV

expression, Notch1 activation (measured by the expression of cleaved Notch1 intracellular

domain), NF-κB activation (measured by p65 nuclear localization), and macrophage infiltration.

As previously observed, tumors from mice treated with paclitaxel expressed significantly more

MenaINV than the control group (**Fig. 1B**), and this was accompanied by a significant increase in

Notch1 and NF-κB activation (**Fig. 1C, D**). When MenaINV induction was inhibited with DAPT,

paclitaxel no longer caused an increase in MenaINV expression (**Fig. 1E**). This indicates that

paclitaxel causes *de novo* MenaINV induction.

We originally hypothesized that the paclitaxel-mediated MenaINV induction was due to an

increase in macrophage infiltration following paclitaxel treatment, but no difference in macrophage

infiltration was observed in any treatment group (**Fig. 1F**). This led us to question the necessity

of macrophages in paclitaxel-mediated MenaINV induction.

Macrophages are required for paclitaxel-mediated MenaINV induction

To clarify the role of macrophages in paclitaxel-mediated MenaINV induction, we treated

FVB-PyMT mice bearing spontaneous breast tumors with paclitaxel and co-administered

clodronate liposomes – to deplete macrophages – or PBS control liposomes (**Fig. 2A**). Following

the treatment period, the mice were sacrificed, and their tumors were analyzed for MenaINV

expression, Notch1 and NF-κB activation, and macrophage infiltration. Paclitaxel again caused a

significant induction of MenaINV, and this induction was prevented when macrophages were

depleted (**Fig. 2B, C**). This indicates that macrophages *are* required for paclitaxel to induce

MenaINV. Notch1 and NF-κB signaling were also significantly decreased upon macrophage

depletion (**Fig. 2D, E**), indicating that macrophages are the primary source of additional Notch1

and NF-κB stimulation in the tumor microenvironment following paclitaxel treatment. Our data

indicate that for paclitaxel to induce MenaINV, macrophages must be present, but not necessarily

in higher proportions than in untreated conditions. This led us to ask whether paclitaxel changes

the phenotype of macrophages to promote NF-κB/ Notch1 signaling.

Paclitaxel induces pro-inflammatory phenotype in macrophages that promotes NF-κB/ Notch1

signaling

To investigate how paclitaxel affects the ability of macrophages to participate in NF-κB/

Notch1 signaling, we cultured RAW264.7 murine macrophages with paclitaxel (0 μM, 10 μM, and

30 μM) for 24 h and measured the levels of secreted TNFα, IL-1β, and TGF-β1, which are known

to directly activate NF-κB. TGF-β1 has also been shown to enhance Notch signaling in several

contexts (13, 14). Treatment with paclitaxel significantly increased macrophage secretion of all

three cytokines (**Fig. 3A, B, C**). These effects were not dose-dependent at the tested

concentrations.

Paclitaxel repolarizes macrophages in patients treated with neoadjuvant chemotherapy

We next sought to establish supporting evidence that chemotherapy causes a

repolarization of primary tumor macrophages in patients, and to confirm the clinical relevance of

this finding. To do this, we identified a previously published dataset of single-cell RNA sequencing

reads from biopsies from human BC patients taken either before receiving any treatment

(treatment naïve) or after receiving neoadjuvant chemotherapy (NAC), as described in the original

publication (15). We analyzed the macrophage compartment of this dataset and found that

macrophages cluster into 4 major groups (0-3), and one minor group (4) (**Fig. 3D**) which express

gene signatures that are associated with either NF-κB activation/inflammatory response (e.g.

TNFα, TGF-β, and IL-6) (referred to here as M1), or anti-inflammatory phenotypes with markedly

lower inflammatory pathway activation (referred to here as M2), or a mixture of the two (**Fig. 3E**),

as previously published (16). By including these three categories, our analysis accounts for a

spectrum of macrophage polarizations, rather than imposing a dichotomy. Under chemotherapy,

clusters 1 & 3 (M1 and mixed, respectively) dominate, with the M1 cluster accounting for 75% of

macrophages. Meanwhile, clusters 0 & 2 (mixed and M2, respectively) dominate under naïve

conditions (**Fig. 3F**). Population analysis of the clusters shows a dramatic increase in the M1

cluster 1 and mixed cluster 3 after chemotherapy (**Fig. 3G**). Pathway analysis shows that,

compared to clusters 0 & 2, cluster 1 is higher in TNFα and TGF-β signaling, and cluster 3 is

higher in TGF-β signaling (**Fig. 3H**). These results confirm that NAC promotes an inflammatory

phenotype in macrophages in human BC, which increases their ability to participate in NF-κB/

Notch1 signaling, and complement our previous finding that NAC increases MenaINV expression

in human BC (8).

Taken together, our results reveal that chemotherapy skews macrophages from an anti-

inflammatory/M2 phenotype towards a pro-inflammatory/M1 phenotype, increasing their ability to

participate in NF-κB/ Notch1 signaling and subsequent MenaINV induction (**Fig. 4**).

Discussion

Discussion

In this study, we discovered that neoadjuvant paclitaxel induces the expression of the pro-

metastatic protein MenaINV in BC cells by promoting NF-κB/ Notch1 signaling with macrophages.

We previously found that paclitaxel causes an increase in MenaINV (8, 10). However, given that

MenaINV expression confers resistance to paclitaxel (9), it was unknown whether the observed

increase was a result of *de novo* induction or the selective killing of non-MenaINV-expressing

cells. Here, we show that the increase in MenaINV following paclitaxel treatment is the result of

de novo induction. When the pathways involved in MenaINV induction were blocked during

paclitaxel treatment, no increase in MenaINV expression was observed (**Fig. 1E**). These results

indicate that paclitaxel induces MenaINV expression.

We have previously shown that Notch1 signaling between macrophages and tumor cells

is crucial for robust MenaINV induction (10, 11). The Notch1 receptor and its ligands are

membrane-bound, so Notch1 signaling only occurs between adjacent cells. Given this, and our

previous finding that macrophages play a key role in the paclitaxel-mediated increase in MenaINV

(10), we hypothesized that paclitaxel causes a macrophage influx into the tumor, increasing the

potential for macrophage-tumor cell Notch1 signaling by increasing the number of tumor cells in

direct contact with macrophages. Interestingly, we found that, while macrophages are necessary

for paclitaxel to induce MenaINV expression (**Fig. 2B**), an increase in macrophage proportion is

not (**Fig. 1E, F**). We conclude that, rather than increasing macrophage-tumor cell NF-κB/ Notch1

signaling by increasing macrophage infiltration, paclitaxel repolarizes macrophages, allowing

them to participate in such signaling more readily (**Fig. 3**). This is consistent with previous reports

that paclitaxel activates TLR4 receptors on macrophages, leading to an increase in inflammatory

signaling (17).

Chemotherapy has long been known to exert both pro- and anti-tumoral effects on

macrophages, and an existing body of evidence shows that chemotherapy-induced pro-

inflammatory macrophages are part of the anti-tumoral response (18). Indeed, an increase in pro-

inflammatory macrophages following chemotherapy has been linked with an elevated innate and

adaptive anti-tumoral immune response (18). In the present study, we provide evidence that

chemotherapy-induced pro-inflammatory macrophages also have the pro-tumoral effect of

increasing the expression of the pro-metastatic protein MenaINV. As drug discovery platforms and

the ability to target specific proteins improve, it is crucial to continue deeper investigations into

the exact mechanisms behind both the pro- and anti-tumoral responses to chemotherapy to

continue refining treatment strategies.

Materials & Methods

Animal models

All procedures were performed in accordance with National Institutes of Health regulations and approved by the Albert Einstein College of Medicine Institutional Animal Care and Use Committee. Transgenic mice expressing the polyoma virus middle-T (PyMT) antigen under the control of the mouse mammary tumor virus long terminal repeat (MMTV-LTR) (19) were bred in house and have palpable mammary tumors at approximately 6 weeks of age. Only female mice were used in experiments.

In vivo treatments

Treatments began when mice were 8-10 weeks of age. Paclitaxel (Sigma) was reconstituted to a stock concentration of 10 mg/mL in 1:1 ethanol/Cremophor-EL (Millipore Sigma 238470) and stored in aliquots at -20 °C. Immediately before treatments, the stock solution was further diluted in PBS and given at a dose of 10 mg/kg intravenously (total injected volume 200 μL) for a total of 3 doses 5 days apart. Mice in the vehicle control groups received the same volume of ethanol/Cremophor-EL diluted in PBS on the same schedule.

DAPT (Sigma D5942) was reconstituted in 100% ethanol to a stock concentration of 20 mg/mL and stored in aliquots at -20 °C. Immediately before treatments, the stock solution was further diluted in corn oil and given at a dose of 10 mg/kg intraperitoneally (total injected volume 100 μL) daily for 14 days. Mice in the vehicle control groups received the same volume of ethanol in corn oil on the same schedule.

Clodronate liposomes (Encapsula Nano Sciences, SKU# CLD-8901) or PBS control liposomes were stored at 4 °C. Mice received 200 μL liposomes (concentration 18.4 mM) intraperitoneally every 2 days for 14 days.

<u>Tissue fixing and staining</u>

Following treatments, mice were sacrificed, and mammary tumors were removed and

fixed in 10% formalin for 24-48 hours (tumor to formalin volume ratio of 1:7). Tissues were then

paraffin-embedded and cut in 5 μm thick sections. Slides were deparaffinized by melting the

paraffin at 60 °C for 1 h, followed by 2 x 10-min incubations in xylene. Slides were then rehydrated

in ethanol and water, and antigen retrieval was performed in 1x citrate buffer (pH 6.0, Vector

H3300) at 97 °C for 20 min in a steamer.

For cleaved Notch1 intracellular domain (cN1ICD) and p65 staining, slides were then

incubated in 0.3% hydrogen peroxide in water for 10 min to quench endogenous peroxidase and

then blocked in a buffer containing 1% bovine serum albumin, 1% normal goat serum, and 10%

fetal bovine serum (FBS) in 0.05% PBS-Tween20 for 1 h at room temperature. Slides were then

stained using the multiplex tyramide signal amplification (TSA) immunofluorescence assay

(Akoya Biosciences) according to the manufacturer's instructions with the following primary

antibodies: rabbit anti-cN1ICD (1:200, CST4147), rabbit anti-p65 (1:4000, CST8242). Briefly,

slides were incubated in the appropriate primary antibody diluted in blocking buffer overnight at 4

°C. The next day, slides were washed 3 x 5 min in 0.5% PBS-Tween20 and then incubated in

goat anti-rabbit-HRP diluted 1:2000 in blocking buffer for 1 h at room temperature. After washing

3 x 5 min with 0.5% PBS-Tween20, TSA was performed by incubating slides in Opal reagents

(Akoya FP1487001KT and FP1496001KT) diluted 1:100 in amplification buffer for exactly 9 min.

The process of antigen retrieval, blocking, antibody incubation, and TSA was repeated for each

primary antibody. After the final wash, slides were incubated in DAPI for 5 min and mounted with

ProLong Diamond Antifade Mountant (Invitrogen).

For MenaINV and IBA1 staining, following antigen retrieval, slides were incubated in

blocking buffer (same as above) for 1 h at room temperature. The tissues were then incubated in

the following primary antibodies diluted in blocking buffer overnight at 4 °C: chicken anti-MenaINV

(0.25 μg/mL, generated by Covance as previously described (11)), rat anti-IBA1 (1:800, Invitrogen

MA5-38266). The next day, slides were washed 3 x 5 min in 0.05% PBS-Tween20 and incubated

in the following secondary antibodies diluted in blocking buffer for 1 h at room temperature: goat

anti-chicken 555 (1:250, Invitrogen A32932), goat anti-rat 647 (1:1000, Invitrogen A21247). After

washing 3 x 5 min in 0.05% PBS-Tween20, slides were incubated in DAPI for 5 minutes and

mounted using ProLong Diamond Antifade Mountant (Invitrogen). All stained slides were imaged

on the Pannoramic 250 Flash II digital whole slide scanner using a 20 x 0.75NA objective lens.

<u>Image analysis</u>

To analyze the % MenaINV+ area and % IBA1+ area, the digital whole slide scans of slides

stained for MenaINV and IBA1 were uploaded into CaseViewer (3DHISTECH), and 10 different

40X fields of view representative of the entire tumor were chosen for each mouse from invasive,

non-necrotic areas of the tumor. Tif files of each field of view were acquired and uploaded to FIJI

(NIH). In FIJI, images were thresholded based on the negative staining control, and the area

fraction (area of signal/total area of field) was calculated in each channel for each field of view.

The average value of the 10 fields per mouse is reported.

To analyze Notch1 activation, the digital whole slide scans of slides stained for cN1ICD

were uploaded into CaseViewer, and representative fields of view were chosen and uploaded to

FIJI as described above. In FIJI, images were thresholded based on the negative staining control.

The area of cN1ICD signal was normalized to the total DAPI area in each field, and the average

value of the 10 fields per mouse is reported.

To analyze NF-κB activation via p65 nuclear localization, the digital whole slide scans of

slides stained for p65 were uploaded into CaseViewer, and representative fields of view were

chosen and uploaded to FIJI as described above. In FIJI, images were thresholded based on the

negative staining control, and two masks were made - one of the total p65 signal and one of the

DAPI signal. P65 was considered nuclear if it overlapped with a DAPI-stained nucleus. Nuclear

p65 was normalized to the total DAPI signal in the field, and the average value of the 10 fields per

mouse is reported.

Enzyme-linked immunosorbent assay (ELISA)

Paclitaxel used in the cell culture (ThermoFisher J62734.MC) was reconstituted in DMSO

at a stock concentration of 0.01 M and stored in aliquots at -20 °C.

In the morning, RAW264.7 cells were seeded at a density of 7.5 x 10^5 cells/well in a 6-well

plate in DMEM containing 10% FBS and 1% penicillin/streptomycin. That evening, the media was

replaced with serum-starving DMEM (1% FBS, 1% penicillin/streptomycin). The cells were serum-

starved overnight. The next morning, the media was replaced with fresh serum-starving media

with the indicated concentrations of paclitaxel. Conditioned media from the cultured cells were

collected after 24 h and stored at -80 °C until use. Levels of TNFα, IL-1β, and TGF-β1 were

determined using ELISA kits (R&D Systems) according to the manufacturer's instructions. The

concentrations of secreted proteins were interpolated from the standard curve measurements.

Single-cell RNA sequencing analysis

Single-cell RNA sequencing data was obtained from the publicly available site

http://biokey.lambrechtslab.org. Raw sequencing reads can also be found in the European

Genome-phenome Archive under study number EGAS00001004809 (15). Only cells taken from

biopsies before anti-PD1 therapy ("Pre" condition) were considered in our analysis.

Quality control and dimensionality reduction: As the archived dataset had already been

subjected to rigorous quality control measures by the original authors, no additional quality control

measures needed to be performed. Raw counts were normalized using the Seruat

"NormalizeData" function, and principal component analysis (PCA) was performed to reduce the

dimensionality of the data and mitigate potential batch effects. Harmony integration was

subsequently applied using the PCA embeddings as input.

Cell type annotation: Cell types were annotated using the SingleR package, as previously

described (16).

Clustering: Cells were clustered using the Seurat framework. PCA and Harmony reduction

were employed to derive a low-dimensional representation of the data, followed by the

identification of neighbors and clusters using Seurat's FindNeighbors and FindClusters functions.

A resolution of 0.3 was used to balance cluster granularity. Clusters were visualized using Uniform

Manifold Approximation and Projection (UMAP) for high-dimensional data.

Gene set variation analysis (GSVA): GSVA was performed using the GSVA R package to

evaluate pathway activity across the dataset. Gene sets were sourced from the Molecular

Signatures Database (MSigDB). GSVA was parallelized using the BiocParallel package. Pathway

activity was summarized for each cluster by calculating the mean enrichment scores for each

pathway across cells in the cluster. Differential gene expression analysis was performed between

the identified clusters.

Macrophage marker analysis: Macrophages were categorized into three classes: M1, M2,

or mixed M1/M2 based on literature-reported markers, as previously described (16). The M1

markers used were CD80, CD86, IRF3, IRF5, STAT1, and STAT5. The M2 markers used were

STAT3, STAT6, IRF4, KLF4, JMJD3, CD163, CD206, and CD209.

Figure Legends

Figure 1: (**A**) Experimental design of paclitaxel (ptx) and DAPT treatments in PyMT mice. (**B**) % MenaINV+ area in control (ctrl)- and paclitaxel-treated tumors. (**C**) Notch1 activation in control- and paclitaxel-treated tumors. (**D**) NF-kB activation in control- and paclitaxel-treated tumors. (**E**) % MenaINV+ area in tumors treated with control, paclitaxel, and paclitaxel + DAPT. (**F**) Macrophage infiltration (% IBA1+ area) in tumors treated with control, paclitaxel, and paclitaxel + DAPT.

Figure 2: (**A**) Experimental design of paclitaxel and clodronate (cld) treatments in PyMT mice. (**B**) % MenaINV+ area in tumors treated with control, paclitaxel, and paclitaxel + clodronate. (**C**) Macrophage infiltration (# IBA1+ cells per high power field (HPF) in tumors treated with control, paclitaxel, and paclitaxel + clodronate. (**D**) Notch1 activation in tumors treated with control, paclitaxel, and paclitaxel + clodronate. (**E**) NF-kB activation in tumors treated with control, paclitaxel, and paclitaxel + clodronate.

Figure 3: Secretion of (**A**) TNAα, (**B**) IL-1β, and (**C**) TGF-β1 from RAW264.7 macrophages treated with paclitaxel. (**D-H**) scRNAseq analysis of macrophages in the tumors of breast cancer patients who were treatment naïve or had undergone neoadjuvant chemotherapy. (**D**) Clustering analysis showing 4 major groups (0-3), and one minor group (4). (**E**) Clusters correlated with signatures indicative of M1, M2 or mixed types. (**F**) M1 cluster 1 & mixed cluster 3 dominate under chemotherapy; mixed cluster 0 & M2 cluster 2 dominate under naïve conditions. (**G**) Chemotherapy results in a dramatic increase in M1 cluster 1 & mixed cluster 3. (**H**) Pathway analysis shows that, compared to clusters 0 & 2, cluster 1 is higher in TNFα & TGF-β signaling (green boxes), and cluster 3 is higher in TGF-β signaling.

321   **Figure 4**: Paclitaxel induces MenaINV expression in tumor cells by skewing macrophages from

322   a mixed/M2 (anti-inflammatory) phenotype to a mixed/M1 (pro-inflammatory) phenotype that

323   increasingly activates Notch1 and NF-kB signaling in tumor cells, leading to increased MenaINV

324   expression.

Declarations

Ethics approval and consent to participate

All procedures were conducted in accordance with National Institutes of Health regulations and approved by the Albert Einstein College of Medicine Institutional Animal Care and Use Committee.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This study was supported by grants from the NIH (R01 CA255153, K99/R00 CA237851), SIG OD019961, the Gruss-Lipper Biophotonics Center, the Integrated Imaging Program for Cancer Research, The Evelyn Gruss-Lipper Charitable Foundation, and The Helen & Irving Spatz Foundation.

Authors' contributions

MFD, MHO, and DE performed conceptualization. MFD, GSK, CLD, SS, and JL provided methodology. MFD and JL carried out formal analysis. SS & DE provided software and computational image analysis. MFD, GSK, and CLD performed investigation. MFD wrote the

manuscript. GSK, JSC, MHO, and DE performed funding acquisition. MHO and DE supervised

the study. All authors read and approved the final manuscript.

<u>Acknowledgments</u>

We thank the Histopathology Core Facility for their assistance with sample processing and the

Analytical Imaging Facility for their imaging services. We thank Robert Eddy for the development

and validation of the MenaINV antibody. We thank Luis Rivera Sanchez for assistance with

macrophage quantification and Ved Sharma for contributing to sample generation. We thank

Shanye Yin and Erqiang Hu for providing their scRNAseq macrophage analysis code.

References

1.   Siegel RL, Giaquinto AN, Jemal A. Cancer statistics, 2024. CA Cancer J Clin. 2024;74(1).

2.   Weidmann MD, Surve CR, Eddy RJ, Chen X, Gertler FB, Sharma VP, et al. MenaINV dysregulates cortactin phosphorylation to promote invadopodium maturation. Sci Rep. 2016;6:36142.

3.   Hughes SK, Oudin MJ, Tadros J, Neil J, Del Rosario A, Joughin BA, et al. PTP1B-dependent regulation of receptor tyrosine kinase signaling by the actin-binding protein Mena. Mol Biol Cell. 2015;26(21):3867-78.

4.   Philippar U, Roussos ET, Oser M, Yamaguchi H, Kim HD, Giampieri S, et al. A Mena invasion isoform potentiates EGF-induced carcinoma cell invasion and metastasis. Dev Cell. 2008;15(6):813-28.

5.   Goswami S, Philippar U, Sun D, Patsialou A, Avraham J, Wang W, et al. Identification of invasion specific splice variants of the cytoskeletal protein Mena present in mammary tumor cells during invasion in vivo. Clin Exp Metastasis. 2009;26(2):153-9.

6.   Borriello L, Coste A, Traub B, Sharma VP, Karagiannis GS, Lin Y, et al. Primary tumor associated macrophages activate programs of invasion and dormancy in disseminating tumor cells. Nat Commun. 2022;13(1):626.

7.   Oudin MJ, Jonas O, Kosciuk T, Broye LC, Guido BC, Wyckoff J, et al. Tumor Cell-Driven Extracellular Matrix Remodeling Drives Haptotaxis during Metastatic Progression. Cancer Discov. 2016;6(5):516-31.

8.   Karagiannis GS, Pastoriza JM, Wang Y, Harney AS, Entenberg D, Pignatelli J, et al. Neoadjuvant chemotherapy induces breast cancer metastasis through a TMEM-mediated mechanism. Sci Transl Med. 2017;9(397).

9.   Oudin MJ, Barbier L, Schafer C, Kosciuk T, Miller MA, Han S, et al. MENA Confers Resistance to Paclitaxel in Triple-Negative Breast Cancer. Mol Cancer Ther. 2017;16(1):143-55.

10.   Duran CL, Karagiannis GS, Chen X, Sharma VP, Entenberg D, Condeelis JS, et al.

Cooperative NF-kappaB and Notch1 signaling promotes macrophage-mediated MenaINV

expression in breast cancer. Breast Cancer Res. 2023;25(1):37.

11.   Pignatelli J, Bravo-Cordero JJ, Roh-Johnson M, Gandhi SJ, Wang Y, Chen X, et al.

Macrophage-dependent tumor cell transendothelial migration is mediated by Notch1/MenaINV-

initiated invadopodium formation. Sci Rep. 2016;6:37874.

12.   Li LH, Hsu DZ, Chandrasekaran VRM, Liu MY. Inhibiting CD44-ICD Attenuates LPS-

Induced Initiation of Hepatic Inflammation in Septic Mice. Int J Mol Sci. 2024;25(16).

13.   Wang Y, Shen RW, Han B, Li Z, Xiong L, Zhang FY, et al. Notch signaling mediated by

TGF-beta/Smad pathway in concanavalin A-induced liver fibrosis in rats. World J Gastroenterol.

2017;23(13):2330-6.

14.   Blokzijl A, Dahlqvist C, Reissmann E, Falk A, Moliner A, Lendahl U, et al. Cross-talk

between the Notch and TGF-beta signaling pathways mediated by interaction of the Notch

intracellular domain with Smad3. J Cell Biol. 2003;163(4):723-8.

15.   Bassez AA-O, Vos H, Van Dyck L, Floris GA-O, Arijs IA-OX, Desmedt CA-O, et al. A single-

cell map of intratumoral changes during anti-PD1 treatment of patients with breast cancer. (1546-

170X (Electronic)).

16.   Huang Y, Wu G, Bi G, Cheng L, Liang J, Li M, et al. Unveiling chemotherapy-induced

immune landscape remodeling and metabolic reprogramming in lung adenocarcinoma by scRNA-

sequencing. Elife. 2024;13.

17.   Wanderley CW, Colón DF, Luiz JPM, Oliveira FF, Viacava PR, Leite CA, et al. Paclitaxel

Reduces Tumor Growth by Reprogramming Tumor-Associated Macrophages to an M1 Profile in

a TLR4-Dependent Manner. Cancer Res. 2018;78(20):5891-900.

18.   Friedman-DeLuca M, Karagiannis GS, Condeelis JS, Oktay MH, Entenberg D.

Macrophages in tumor cell migration and metastasis. Front Immunol. 2024;15:1494462.

19.   Guy CT, Cardiff RD, Muller WJ. Induction of mammary tumors by expression of

polyomavirus middle T oncogene: a transgenic mouse model for metastatic disease. Mol Cell Biol.

1992;12(3):954-61.