PLANT DOMESTICATION DOES NOT REDUCE DIVERSITY IN ROOT

MICROBIOMES

Alejandra Hernández-Terán[1]*, Ana E. Escalante[2], Maria Rebolleda-Gómez[1]*

[1]Ecology and Evolutionary Biology Department, COMMONS Lab, University of California Irvine.

Irvine, California, 92697.

[2]Laboratorio Nacional de Ciencias de la Sostenibilidad, Instituto de Ecología, Universidad Nacional

Autónoma de México, Mexico City, Mexico.

*Corresponding authors: Alejandra Hernández-Terán, alejh18@uci.edu, María Rebolleda-Gómez,

mreboll1@uci.edu

Keywords (5-8): Community assembly, host control, microbial diversity, rhizosphere, root

microbiome, and plant domestication.

SUMMARY

- Domestication has profoundly shaped the genetic makeup of numerous plant and animal species. While the effects of plant domestication at the genetic and phenotypic levels are well-documented, its impact on plant microbiome remains less understood.

- Two primary hypotheses have been proposed: 1) the reduction in microbial diversity resulting from the domestication process, and 2) the diminished ability of host plants to control their microbiomes.

- We conducted a meta-analysis of multiple crops, comparing the root microbiomes of domesticated plants and their wild relatives. Our results indicate that the effects of domestication are species-specific and context-dependent, with most domesticated plants exhibiting increased microbial diversity and more structured communities.

- Overall, this study provides evidence that plant domestication does not lead to a uniform reduction in microbial diversity or a consistently diminished ability of plants to influence their microbiomes.

- Based on these findings, we discuss new perspectives and the need for future studies incorporating native soils and host genetic variation in such experiments, analyzing diversity and microbiome function, and considering how root morphology might affect microbiome recruitment.

INTRODUCTION

Domestication, the artificial selection of phenotypic traits useful to humans, has profoundly shaped human civilization and the genetic makeup of numerous plant and animal species (Purugganan & Fuller, 2009). The consequences of domestication at the genetic and phenotypic level are well-documented (Meyer *et al.*, 2012). Typically, domestication events involve propagating only selected plant seeds, leading to genetic bottlenecks and, thus, a reduction in genetic diversity across the genome (Doebley *et al.*, 2006). Phenotypically, convergent changes in certain traits, known as 'domestication syndrome', have been observed across diverse domesticated species (Alam & Purugganan, 2024). In plants, this syndrome is characterized by enlarged fruit sizes, increased apical dominance, and the loss of seed dispersal mechanisms and dormancy, among other traits (Doebley *et al.*, 2006). Many of these traits are the target of selection, but many others are not. Those non-target traits can evolve through correlations (genetic linkages or pleiotropy) with the target traits (e.g. reduction of seed shattering, uniform flowering, and ripening) and can greatly influence the ecology of the plant (Purugganan & Fuller, 2009; Meyer *et al.*, 2012). The extent to which domestication-induced phenotypic changes affect ecological interactions remains an open question.

It is often proposed that domestication leads to reduced investment in other ecological interactions. In the case of plant-herbivore interactions, a commonly proposed hypothesis suggests that domestication has modified plant traits in ways that decrease resistance to herbivores (Benrey *et al.*, 1998). Nonetheless, a meta-analysis (Whitehead *et al.*, 2017) conducted in 2017 assessed the overall impact of crop domestication on plant-herbivore interactions and found that, while domestication appears to have a generally negative effect on herbivore resistance, it did not significantly alter plant defense traits. Similarly, a recent study (Glasser *et al.*, 2023) exploring the consequences of domestication on plant-pollinator interactions in members of the *Cucurbita* genus revealed that domestication in this group may have promoted greater resource investment in floral traits. This increased investment likely enhances attractiveness to pollinators, potentially increasing reproductive success.

Roots are an important locus for the evolution of non-target traits and species interactions. Root phenotypes have often significantly diverged from their wild relatives due to adaptation to the agricultural environment or as non-target modifications associated with selected traits above ground. Domesticated plants have a different root system architecture, altered root exudate compositions, and significant changes in secondary metabolites across a wide range of species

(Meyer *et al.*, 2012; Singh & van der Knaap, 2022) which necessarily has an impact in plant
interactions with other organisms, including symbiotic microbial communities associated with
plant roots (the root or rhizosphere microbiome).

Plants rely on their microbiomes for several functions, including nutrient acquisition, protection
against pathogens, and tolerance to biotic stress (Pérez-Jaramillo *et al.*, 2018). Often, the root
microbiome promotes plant health and growth, contributing to plant development (Singh *et al.*,
2020). Given the pivotal importance of root microbiomes for plants and considering the known
effects of plant domestication on root systems, in recent years, efforts have been undertaken to
decipher the impact of plant domestication on the diversity and assembly of these microbial
communities. Despite these efforts, it is still unclear if, or to what extent, plant domestication has
modified the ability of the plants to interact with the root microbiome.

Two main hypotheses have been proposed regarding the consequences of domestication: 1) the
reduction in microbial diversity resulting from the domestication process and 2) the diminished
ability of host plants to shape the assembly of their microbiomes. These hypotheses arise from
the idea that the genetic bottlenecks accompanying domestication may lead to the loss of alleles
crucial for plant-microbiome interactions, that alterations in root exudates metabolic profiles might
further influence microbiome assembly in domesticated plants (Gutierrez & Grillo, 2022), and
because of a potential reduction in the benefits of mutualism for domesticated plants. After
growing in environments with important nutrient inputs, domesticated plants could reduce the
investment in symbiosis, leading to the evolution of reduced symbiont cooperation (Porter &
Sachs, 2020). To evaluate these hypotheses, we conducted a meta-analysis comparing studies
inoculating domesticated plants and their wild relatives with the same bulk soil in common garden
conditions. We found that plant domestication does not result in a uniform reduction in microbial
diversity or a consistently diminished ability of host plants to influence their microbiomes. The
effects of domestication are species-specific and context-dependent, with some domesticated
plants showing increased microbial diversity and structured communities, while others do not.
Despite this variation, we found consistent losses and enrichments of microbial taxa in
domesticated crops relative to their wild progenitors.

MATERIALS AND METHODS

Data collection

To investigate the impact of the domestication process on the diversity and composition of the root microbiome across various crop species, we collected 16S rRNA sequence data from common garden experiments reported in scientific publications. We conducted a comprehensive search in the Scopus and Google Scholar databases using the Boolean search string: Wild [AND] domesticated [AND] microbiome [OR] rhizosphere [AND] crop. Our search centered on contemporary crops with well-identified crop wild relatives, including but not limited to maize, potato, tomato, squash, sorghum, bean, coffee, barley, soybean, cotton, and sunflower. Eligibility for inclusion in our meta-analysis was based on the following selection criteria: 1) the experiments must have been conducted under controlled environmental conditions typical of common garden studies, 2) wild and domesticated variants of the same crop species must have been cultivated concurrently, and inoculated with the same soil, and 3) sequencing must have been performed using Illumina platforms. We excluded studies that assessed the impact of extreme environmental stressors, such as herbivory, but we incorporated those that utilized different soil types (e.g., forest or agricultural soils) as inoculum. Applying these criteria, we narrowed down the pool to 15 publications, representing eight crops: barley, soybean, cotton, tomato, potato, sunflower, maize, and bean. A detailed overview of the number of crop accessions, experimental replicates, and analyzed publications per crop is available in Table S1.

Additionally, due to the large number of papers we could not include in our meta-analysis because of missing raw sequences or metadata, we selected a subset of studies that met our inclusion criteria and reported diversity data and/or ASV tables. A detailed summary of the crop accessions, experimental replicates, and publications analyzed in this secondary dataset is provided in Table S2.

With these datasets, we investigated two primary hypotheses regarding the impact of plant domestication on root microbiomes: 1) the reduction in microbial diversity resulting from the domestication process, and 2) the diminished ability of host plants to control their microbiomes. Since this data did not experimentally assess plant "control", hereafter we refer as "host influence" to the contribution of plants in shaping root microbiome assemblies. Our analysis focused on the comparison between wild and domesticated categories. To mitigate the potential

bias introduced by merging amplicon sequence data from disparate studies, our comparative
analyses were restricted to within-study wild and domesticated pairs (W-D pairs). We defined
W-D pairs as plants cultivated concurrently and inoculated with the same soil. Typically, each
paper contributed a single W-D pair to our analysis. However, in instances where a study
investigated more than one soil type, the number of W-D pairs was equivalent to the number of
different soil treatments applied.
160
161   *Data processing*
162
16S rRNA raw sequences and metadata were downloaded from public repositories, including
NCBI, MG-Rast, and Github, and were analyzed with the Quantitative Insights Into Microbial
Ecology (QIIME2) platform (Bolyen *et al.*, 2019) version 2023.5. For each dataset corresponding
to a specific publication, paired-end sequences were analyzed independently. Sequence
denoising and quality filtering, including primers and adapters trimming, were performed with the
DADA2 plug-in (Callahan *et al.*, 2016). Given that all selected studies utilized primers targeting
the V4 16S rRNA gene region (e.g., V4, V3-V4, V4-V5), we standardized the analysis by
extracting the V4 segment using the 515F–806R primer pairs (Caporaso *et al.*, 2011). This
approach minimized discrepancies arising from taxonomic assignments across different regions
of the 16S rRNA gene. The length of the V4 region sequences was standardized to a range
between 120 and 150 nucleotides, based on sequence quality and primer positioning. This
criterion was consistently applied across all datasets. Taxonomic classification of the Amplicon
Sequence Variants (ASVs) for each dataset was conducted using the "sklearn-base" classifier
within QIIME2, referencing the *Greengenes* 2 database (McDonald *et al.*, 2024) with a 99%
sequence identity threshold. Mitochondrial, chloroplast, and ASVs that were not assignable at the
kingdom level were excluded from the analysis. Comprehensive details on the total number of
reads (before and after processing) and the count of ASVs for each study have been compiled in
Table S3.
181
182   *Data Analysis*
183
Data analysis was performed on R software (v.4.3.0) using custom scripts and employing the
following packages: Phyloseq (v. 1.46.0) (McMurdie & Holmes, 2013), ggplot2 (v. 3.5.1) (Valero-
Mora, 2010), metafor (v. 4.2-0) (Viechtbauer, 2025), vegan (v.2.6-4) (Oksanen, 2017), DESEq2
(v.1.40.2) (Love *et al.*, 2014), Source Tracker 2 (Knights *et al.*, 2011), and breakaway (v.4.8.4)

(Willis & Bunge, 2015). Prior to amplicon analysis, each of the 15 datasets, corresponding to the
individual studies included in our meta-analysis, was rarified to a standardized depth of 15,000
reads per sample to ensure comparability across datasets. Specifics on the number of samples
and ASVs retained after rarification are available in Table S3. As described previously, all
analyses were performed using W-D pairs (pairs of wild and domesticated plants from the same
study, species, and treatment).
194
Alpha diversity analyses: To assess the effect of plant domestication on the diversity of the
microbiome we calculated Chao1 and Shannon alpha diversity indexes using the *estimate*
richness function in the phyloseq package. In addition, we estimated species richness using
frequency ratios with the breakaway package. This method applies statistical modeling to predict
the number of unobserved taxa, providing flexible modeling of rare species. To test for differences
in these indexes we applied the Log Ratio of Means (LogRoM) analysis which compensates for
the non-independence of the data by calculating the effect size and variance for each study
individually (Friedrich *et al.*, 2008). We implemented this test with the *metafor* package, using W-
D pairs from all crops and the diversity of the wild plants as reference.
204
205   Also, we calculated Rank Abundances Distributions (RADs) for all samples using vegan package,
206   following the methods reported in (Dal Bello *et al.*, 2021). From the RAD data, we computed log-
207   linear models to fit linear regressions and compare the slopes between wild and domesticated
208   microbiomes. Absolute slope values were plotted and contrasted; in general, more even
209   communities display smaller slopes.
210
211   Additionally, from our secondary dataset (see *Data collection*), we obtained Chao1 and Shannon
212   diversity data either from supplementary information provided in the papers or by downloading
213   ASV tables and using the *estimate richness* function in the phyloseq package. Using these data,
214   we constructed separated databases to test for differences in wild and domesticated microbiomes
215   using the LogRM analysis as previously described.
216
Beta diversity analyses: To explore potential differences in beta diversity between plant
categories and crops, we calculated the Bray-Curtis distance for every W-D pair in the dataset.
Dissimilarity matrices were analyzed with Permutational Multivariate Analysis of Variance
(PERMANOVA) using the *Adonis* function in the vegan package. From the PERMANOVA result,
we extracted the R^2 values to represent the proportion of variation (in terms of beta diversity)

explained by category for all W-D pairs and all crops. A dot plot was constructed with this
information using ggplot2.

Furthermore, from the dissimilarity matrices we calculated, for all crops, the mean dissimilarity to
the bulk of both wild and domesticated plants, and the mean distance to centroid between groups
using the *betadisper* function in the vegan package. We plotted both metrics for all W-D pairs in
the different crops using ggplot2. Finally, a Bayesian microbial source tracker was performed with
the SourceTracker2 program to calculate the contribution of the bulk soil microbial community to
the final root microbiome. All samples were rarefied to a depth of 15,000 reads per sample before
the analysis.

Differentially enriched microbes between categories: We identified differentially enriched
microbes between W-D pairs across all crops using the DESeq2 test, implemented in the DESeq2
package. This test was applied exclusively to W-D pairs (excluding bulk soil sequences), with a
P-value cutoff of <0.01 and a False Discovery Rate (FDR) correction. Subsequently, we merged
the results from all W-D comparisons with differentially abundant features at the family level and
selected only features with a log2Fold change of ≤-5 or ≥5, or those present in two or more crops.
The results were visualized by crop using forest plots constructed in ggplot2. Additionally, we
plotted the individual abundances of microbial families enriched in the same category across
multiple crop species (Fig. S6).

RESULTS

We conducted a systematic meta-analysis to assess the influence of plant domestication on root microbiomes across different plant species. After applying our inclusion criteria and filters of data quality, 15 independent studies were retained to perform the full analyses. These studies encompass eight different crop species and included 947 individual plants belonging to maize (n = 128), wheat (n= 172), barley (n= 55), sunflower (n= 127), tomato (n= 169), cotton (n= 31), bean (n= 52), and soybean (n= 160) (Table S1). Except for soybean, and sunflower, all crops were represented by more than one accession of wild and domesticated plants (Table S1).

Additionally, we included diversity metrics from studies that met our inclusion criteria but were excluded from the full analyses due to missing data. This secondary dataset included eight independent studies from six different crop species, including 758 individual plants belonging to: tomato (n= 28), potato (n= 210), maize (n= 264), sorghum (n= 100), rice (n= 115), and agave (n= 41) (Table S2). All crops in this dataset were represented by more than one accession of wild and domesticated plants, increasing the host genetic diversity analyzed in this study.

In total, we analyzed 40 different W-D pairs (pairs of wild and domesticated plants from the same study, species, and treatment). This included 23 W-D pairs from the main dataset, used for the full analyses, and 17 W-D pairs from the secondary dataset, used exclusively for diversity comparisons.

Domesticated plants show increased microbiome diversity

To test for differences in microbial diversity between wild and domesticated plants we performed multiple diversity, richness, and evenness analyses. Contrary to the common expectation that domestication reduces microbiome diversity, we found, on average, domesticated plants exhibited slightly higher diversity in terms of Chao1 index and species richness calculated using frequency ratios. This pattern was consistent across both the main dataset that included the raw sequences (LogRoM: 0.01, 95% CI: -0.06-0.7, $p < 0.000$, Fig. 1 - Chao1; LogRoM: 0.01, 95% CI: -0.01-0.8, $p < 0.000$, Fig. S1 - Frequency ratios) and the secondary dataset (LogRoM: 0.04, 95% CI: -0.08-0.17, $p < 0.000$, Fig. S3b - Chao1 index). Despite these differences in estimated richness, wild and domesticated species tended to have similar diversity using the Shannon index (main dataset: LogRoM: -0.01, 95% CI: -0.01-0.0, $p= 0.1$, Fig. S2; secondary dataset: LogRoM: 0.00, 95% CI: -0.02-0.02, $p= 0.1$ (without agave data - figure not shown)). The main exception to

this trend was the comparison using the Shannon index for Agave samples in the secondary
dataset, where we found higher diversity in wild plants, influencing the overall model toward
greater diversity in wild plants (LogRoM: -0.01, 95% CI: -0.02-0.01, p < 0.00, Fig. S3a).

Most pairwise comparisons (pairs of wild and domesticated plants from the same study, species,
and treatment) showed differences in diversity, but the effects were highly variable across species
(Fig.1). Among the 23 Wild-Domesticated (W-D) pairs evaluated, only three (13%) exhibited no
statistical difference. Of the 19 (82%) pairs that showed statistically significant differences, 12
(63%) displayed increased diversity in domesticated plants, whereas eight (42%) had higher
diversity in their wild counterparts. Whereas for the secondary dataset (Fig. S3b), we found that
77% of the W-D pairs analyzed showed significant differences, from those, 71% showed
increased diversity in domesticated plants, while 29% showed higher diversity in wild plants.
Notably, the effect of domestication appears to be species-specific and related to functional
groups. Crops like wheat, tomato, and cotton showed increased diversity in their domesticated
forms compared to their wild forms, whereas maize and legumes (beans and soybeans)
presented greater diversity in their wild forms. Despite these overall patterns, we observed
different outcomes, even within the same species. The only exception to this inconsistency was
observed in tomatoes, where all four comparisons consistently revealed higher diversity in
domesticated plants. The most variable crop species was soybean, where of the six total W-D
pairs analyzed, two resulted in higher diversity in domesticated versions, three in higher diversity
in wild plants, and one showed no statistically significant difference. Furthermore, we did not find
differences in microbiome evenness between wild and domesticated plants; when comparing
absolute slope values calculated from the RADs data, we found that all analyzed microbial
communities had small slopes, indicating highly even communities regardless of domestication
category (Fig. S4).

Host species effect is more important than domestication in shaping microbiome
composition

We then evaluated the effect of domestication on microbiome composition using PERMANOVA.
We found that the domestication category accounted for a relatively small fraction of the variation
across most of the crop species studied (Fig. 2 and Fig. S5), suggesting a potential species-
specific effect. For several species, including maize, wheat, barley, sunflower, tomato, and bean,
domestication explained approximately 10% of the observed variance in microbial community

composition. Notably, cotton and soybean (except for two pairs) stood out, with domestication

accounting for a larger portion of the variance, ranging from 12% to 25%. Additionally, we

leveraged experiments that used multiple soil types to inoculate plants, allowing us to test the

contribution of the inoculum to the final microbial community by separating wild and domesticated

plants independently (Table S4). Although only a small set of papers had different soil types, for

barley, different initial soils accounted for a minor portion of the total variation (9.8% for wild plants

and 10.9% for domesticated plants). In contrast, for soybeans, using soils from different origins

accounted for a significant percentage of the variation (~75% for wild and ~78% for domesticated

plants). Taken together, these results suggested that domestication could affect the degree of

host influence but that there might be variation across species.

Wild relatives do not display systematic evidence of increased host influence of their

microbiomes

To evaluate the extent of host influence exerted by wild and domesticated plants over their

microbiomes, we leveraged beta diversity metrics, comparing the microbial communities

associated with plants to those from bulk soils, which served as a reference for the initial microbial

community (Fig. 3). We hypothesized that if plants have a relatively strong influence in the

assembly of their microbiomes, then we would expect to find: a) significant dissimilarity between

plant-associated microbiomes and bulk soil microbiomes (Fig. 3a), suggesting that plants are

selecting specific microbial assemblages distinct from the initial soil; b) low intra-group dispersion

(Fig. 3b), indicating that plant replicates of the same genotype consistently select for similar

microbial compositions; and c) less proportion of the final community explained by the initial bulk

soil (Fig. 3c), implying that the host plays a decisive role in shaping its microbiome, potentially

steering it away from the original soil community composition.

Overall, we did not identify a consistent pattern linking host influence to the category of

domestication. Most comparisons were not statistically significant. Among the significant

comparisons, we observed greater dissimilarity to bulk in wild barley and cotton and in

domesticated wheat and soybean (Fig. 3a). Similarly, we found a diverse range of outcomes in

the intra-group dispersion analyses (Fig. 3b). Domesticated plants generally exhibited more intra-

group dispersion in barley and soybean, whereas we found the opposite effect (more intra-group

dispersion in wild varieties) in wheat, cotton, and bean. Maize, sunflower, and tomato showed no

differences. We didn't find a clear pattern in the contribution of bulk soil microbiomes to the final

plant-associated communities across different levels of domestication (Fig. 3c).

We explored if some other plant attribute could better account for our results. Grouping our results

by functional plant group rather than domestication status provided increased insight.

Grasses (maize, wheat, barley) consistently demonstrated a more distinct divergence from their

initial bulk soil microbiomes (Fig. 3a), with less than 50% of the final community composition

attributable to the bulk soil (Fig. 3c). Whereas the legume soybean presented final communities

closely aligned with the initial bulk soil, reflecting less than 50% differentiation.

Some microbial taxa differentially associate with plants across species according to the

domestication status

Finally, we investigated if, despite all the variation, we could identify microbial groups differentially

present in wild and domesticated crops (see methods; Fig. 4). We identified a few families

consistently enriched or depleted from domesticated crops (relative to their wild counterparts)

across different crops. Two members of the *Rhizobiaceae* family

(*Rhizobiaceae_A_500471* and *Rhizobiaceae_A_500279*) were significantly enriched in the

domesticated variants of wheat, tomato, and soybean. Whereas two members of the

Burkholderiaceae family (*Burkholderiaceae_A_592522* and *Burkholderiaceae_A_574758)* were

consistently enriched in the wild counterparts of wheat, tomato, bean, and soybean. We found

these *Burkholderiaceae* taxa in both the domesticated and the wild rhizosphere. In contrast, the

Rhizobiaceae members appeared almost absent in the wild versions of wheat and tomato

(minimum absolute abundance < 50 reads; Fig. S6).

DISCUSSION

Our study provides new insights into the relationship between plant domestication and microbial diversity and assembly, challenging the widely held belief that domestication always leads to a decrease in microbial diversity and impairs the plant's influence in the assembly of its microbiome. Our results indicate that while domestication does have an effect on microbial diversity and community composition, these effects vary widely across species and experimental conditions.

Contrary to the hypothesis that domestication generally reduces microbial diversity, we found a small but significant reduction of microbiome diversity in the rhizosphere of wild relatives (Fig. 1, Log Ratio of Means with Chao1 index, $p < 0.001$). In fact, for the Chao1 index as well as for the frequency ratios test, most of the significant comparisons indicated that domesticated plants exhibited higher microbial diversity than wild plants (Fig. 1 and Fig. S1). When using the Shannon diversity index, we observed fewer differences and mostly saw no change in diversity across comparisons. The hypothesis that domestication leads to a reduction in microbiome diversity is largely based on the idea that domestication reduces genetic diversity, potentially leading to the loss of alleles important for plant-microbe interactions through selective bottlenecks (Gutierrez & Grillo, 2022) and that in domesticated plants, a reduced investment in symbiosis could promote the evolution of reduced symbiont cooperation (Porter & Sachs, 2020).

It is not entirely clear why the reduction of genetic diversity at the level of populations would lead to reduced microbiome diversity at the level of individuals. Instead, we might expect a decrease in microbiome diversity across hosts (less dispersion) in domesticated varieties compared to their wild counterparts. However, our results highlight that this is only the case in some pairs –mostly wheat, barley, and cotton. The opposite is true in some other pairs, including many of the soybean comparisons (Fig. 3). One of the limitations in making clear conclusions about the impact of host diversity on the microbiome is that host diversity is rarely controlled in these studies (for example, many domesticated accessions might be compared against a single population of wild relatives).

Even the loss of alleles important for plant-microbe interactions does not have to be associated with decreased microbiome diversity. It has been proposed, for example, that a reduction in genetic diversity might lead to less efficient microbiome filtering mechanisms, allowing domesticated plants to be less selective and thus associate with a greater diversity of microbes

(Soldan *et al.*, 2021; Favela *et al.*, 2022; Gutierrez & Grillo, 2022). This hypothesis has gained
attention due to its potential implications for agriculture and resilience to climate change. The
variation in microbial communities explained by host genotype represents the substrate upon
which natural selection can act and, therefore, it is also the variation that could be manipulated
for agricultural improvement (Brown *et al.*, 2020). Thus, a reduction in host influence resulting
from domestication could lead to decreased evolutionary potential within microbial communities,
potentially limiting the adaptive resources of plants in response to environmental stresses (Soldan
et al., 2021). This hypothesis of less efficient filtering mechanisms in domesticated crops is
consistent with our alpha (within-plant) diversity results, where we observed a small increase in
microbiome diversity in domesticated varieties. But, contrary to this expectation, we found little
support for decreased host influence in domesticated plants. In most pairs, the microbial
community of the rhizosphere is equally distant to the bulk soil, with soybean being the only
consistent example of increased distance to bulk soil in the domesticated varieties (even if most
are not significant) (Fig. 3).

Until recently (Doebley *et al.*, 2006), the existence of a domestication syndrome had promoted
the idea that most plant species that undergo domestication suffer the same phenotypic
consequences. Likewise, this could have also promoted the assumption that, for all crop species,
domestication would always negatively affect microbial diversity. In addition to finding, on
average, increased microbial diversity in domesticated plants, our analysis revealed significant
variability inside species. For many of the crops analyzed, there were instances where wild plants
exhibited higher microbial diversity, while in other cases, domesticated plants showed greater
diversity. We found this variability in different studies using the same plant accessions and even
within the same study when different initial soils were used, implying a significant role of genotype-
by-environment (GxE) interactions in shaping microbiome assembly.

It is known that microbial communities from different habitats (i.e., different initial soils) can exhibit
distinct responses to the same host genotype. It has been proposed that environmental
heterogeneity can decouple host genotypes from their microbiomes by altering genotype-microbe
relationships (Bradshaw, 1965; Wagner *et al.*, 2016), potentially promoting mixed results even
when controlling for host genotypes. Our results suggest that domestication's effects on microbial
diversity are not universally negative and may vary depending on the specific crop species and
environmental conditions involved. This observation is further supported when examining the
impact of domestication on microbial community composition. Although most comparisons

showed differences in beta diversity (Figure S5), domestication status accounted for a relatively
minor fraction of the microbial variation across the analyzed species (Fig. 2). Instead, much of the
variation, at least in studies using more than one soil type, is explained by the soil origin (Table
S4). This finding aligns with the results of individual studies on rice (Edwards *et al.*, 2015), poplar
(Bonito *et al.*, 2019), soybean (Liu *et al.*, 2019), and the legume *Medicago truncatula* (Brown *et*
al., 2020), where soil origin is a stronger predictor of community composition than host genotype.
Some of these studies suggest that endophytes are more sensitive to host genotype signals, while
rhizosphere communities are more affected by the environment (Brown *et al.*, 2020). Future
studies looking at the effect of domestication should aim to separate the effects of domestication
on different microbial functional groups.
462
Despite all the variation found between crops, we detected two members of the *Rhizobiaceae*
family consistently enriched in domesticated wheat, tomato, and soybean, and two members of
the *Burkholderiaceae* family enriched in wild wheat, tomato, bean, and soybean (Fig. 4). In some
crop species, domesticated plants exhibit increased photosynthetic capacity compared to wild
relatives, which is thought to support the development of larger grains or fruits (Koester *et al.*,
2016; Brestic *et al.*, 2018). This greater carbon fixation could be increasing the root exudation in
domesticated plants, potentially attracting more rhizobia and other copiotrophs in the rhizosphere
(Richards, 2000). Additionally, prolonged nitrogen fertilization in agricultural systems is proposed
to promote the evolution of less-cooperative rhizobia in domesticated cultivars, potentially leading
to higher root abundance of *Rhizobiaceae* members, many of which may not contribute
significantly to plant nutrition (Weese *et al.*, 2015; Liu *et al.*, 2019). Particularly, rhizobia with poor
or null symbiotic effectiveness are more prevalent in agricultural soils and are predominantly
associated with cultivated crops (Sachs *et al.*, 2010). Thus, a high abundance of those microbes
does not necessarily indicate a higher nitrogen fixation in the rhizosphere of domesticated crops.
477
On the other hand, the consistent enrichment of *Burkholderiaceae* family members in the wild
versions of multiple species aligns with previous research on common beans and rice (Pérez-
Jaramillo *et al.*, 2019; Wu *et al.*, 2022). Although the mechanisms behind wild plants harboring an
increased abundance of those particular microbes remain unknown, it has been shown that
members of the *Burkholderiaceae* family are known for their ability to promote plant growth and
suppress soil-borne pathogens, supporting plant health overall (Estrada-De Los Santos *et al.*,
2001). One potential explanation is that compared to domesticated plants, wild relatives produce
a greater diversity of secondary metabolites, which may promote the recruitment of these

beneficial microbes (Ku *et al.*, 2020). Nonetheless, for the case of the common bean, positive

associations with *Burkholderiaceae* family members were found only when using native soils (e.g.

versus agricultural soils) (Pérez-Jaramillo *et al.*, 2019), suggesting potential local adaptation to

microbes found within the natural distribution of such species.

CONCLUSIONS AND FUTURE DIRECTIONS

In this study, we challenged accepted hypotheses regarding the consequences of plant

domestication on root microbiomes. Contrary to the expectation that domestication generally

reduces microbial diversity, we found a small but significant reduction of microbiome diversity in

the rhizosphere of wild relatives. Furthermore, by analyzing multiple crop species, we did not find

sufficient evidence to support the hypothesis that domestication negatively affects host influence

on the assembly of the microbiome. The consistent enrichment of certain bacterial families in both

wild or domesticated versions of different crop species (*Rhizobiaceae* for domesticated and

Burkholderiaceae for wild plants) could be suggesting, at least for the data analyzed in this work,

that the effects of domestication may be stronger at the level of recruiting specific microbes rather

than affecting the entire community. Overall, our results suggest that the impacts of domestication

on the root microbiomes are not universally negative in terms of microbial diversity. Instead, they

appear to be species-specific and contingent upon environmental factors.

Despite our findings, much work remains to fully understand the ecological and evolutionary

consequences of plant domestication on root microbiomes. First, diversity is not everything;

characterizing functional traits of microbiomes associated with wild and domesticated plants is

crucial to better understanding the consequences of domestication for plant-microbe interactions.

Second, it is possible that the effect of domestication would be more pronounced with close

symbionts and endophytes, more work is necessary to understand how domestication can alter

these interactions. Third, it is well-documented that domestication significantly impacts root

system architecture, but there is limited research on how such changes affect the microbes

inhabiting the roots. How changes in root size, phenotype, and physical properties of the roots

impact the recruitment of microbes remains as open questions. Similarly, although exudates have

been studied more than the physical structure of roots for microbiome recruitment, this is still an

area where more research is needed to identify general patterns.

Critical directions for future research are to incorporate plant diversity in the design or analysis of

the experiments to start uncovering the connection between genetic changes of domesticated

crops and their impact on the microbiome. In particular, controlling for genetic diversity between

plants is important to be able to attribute differences between groups to the process of

domestication itself. Similarly, to evaluate the role of domestication on the loss of host influence,

it will be important to incorporate native soils into common garden experiments. Local adaptation

is important for wild populations, and relying exclusively on agricultural soils (or other soils outside

the natural range of wild plants) may hinder significant microbial interactions that occur only with

the specific set of microbes to which wild plants are naturally adapted.

Finally, introgression between wild and domesticated plants is a common phenomenon,

especially at the centers of origin of the species (Ellstrand *et al.*, 2002). The evolutionary

consequences of this have been extensively studied from a genetic perspective (Janzen *et al.*,

2019), and as a source for key agronomic traits (Le Corre *et al.*, 2020). Nonetheless, even though

there is increasing evidence showing that hybridization impacts root microbiomes (Bouffaud *et*

al., 2014; Zhang *et al.*, 2023), the consequences of introgression between wild and domesticated

plants remain unclear. Recently, studies on rice (Zhang *et al.*, 2023) and maize (Wagner *et al.*,

2021) have found that both root and seed microbiomes could contribute to heterosis ("hybrid

vigor") in hybrids, providing evidence of the important role of microbiomes in the fitness of inbreds.

Studying the potential adaptive and maladaptive consequences of microbiome introgression is a

new and exciting area of study, with significant implications for plant breeding and conservation

biology.

ACKNOWLEDGMENTS AND FUNDING SOURCES

We would like to thank Brandon S. Gaut, Eduardo Choreño-Parra, Santiago Rosas-Plaza, Celia Symons, and members of the COMMONS Lab for productive discussions about this work. AHT was supported by the Human Frontiers Science Program Long-Term Postdoctoral Fellowship (LT 0047/2023-L) and the University of California President's Postdoctoral Fellowship. AEH was supported by UNAM-PAPIIT: IN214124. MRG was supported by the School of Biological Sciences at the University of California, Irvine (UCI).

AUTHOR CONTRIBUTIONS

AHT and MRG designed the research with input from AE. AHT performed all statistical analyses. AHT wrote the manuscript with substantial feedback from MRG and AE.

COMPETING INTERESTS

The authors declare no competing interests.

DATA AVAILABILITY STATEMENT

Data and R code for analyses is available on Github at:

https://github.com/AlejandraHT/Plant_domestication_microbiomes

FIGURES

Figure 1. Effect of Domestication on Microbial Alpha Diversity Across Different Crop Species.

Forest plot showing the log Ratio of Means (LogRoM) using Chao1 diversity index for all 23 Wild-

Domesticated (W-D) pairs. The mean microbial diversity of wild plants was used as reference. Dashed

lines denote the separation between different plant species. Bars represent the mean response ratios

with 95% confidence intervals. Colors indicate the results: dark green represents higher diversity in wild

plants, light green represents higher diversity in domesticated plants, and grey indicates no statistical

difference.

594 **Figure 2. Effect of Domestication on Microbial Community Composition Across Different Crop**

595 **Species.** Dot plot showing the percentage of variation in microbial community composition explained by

596 domestication category (wild vs. domesticated), according to the Bray-Curtis distance measure. Numbers

597 in X axis represent the number of Wild-Domesticated pairs (pairs of wild and domesticated plants from the

598 same study, species, and treatment) analyzed per crop.

Figure 3. Impact of Domestication on Host Influence of the Microbiome

A) Bray-Curtis dissimilarity to bulk soil for wild-domesticated (W-D) pairs across different crop

species. Statistically significant differences are indicated by an asterisk (Wilcoxon rank-sum test,

$p < 0.05$). This analysis includes only experiments that included bulk soil sequences. **B)** Intra-

group dispersion for all W-D pairs analyzed using Bray-Curtis distance. Statistically significant

differences are indicated by an asterisk (Wilcoxon rank-sum test, $p < 0.05$). **C)** Source Tracker

analysis estimating the contribution of bulk soil to the final microbial community of wild and

domesticated plants across different species.

631 **Figure 4. Differentially enriched microbial families between wild and domesticated plants across**

632 **various crop species.** DESeq analysis at the family level displaying microbes either enriched in two or

633 more crop species or exhibiting a fold change greater than +5 or less than -5.

REFERENCES

Alam O, **Purugganan MD**. **2024**. Domestication and the evolution of crops: variable syndromes, complex
genetic architectures, and ecological entanglements. *The plant cell* **36**: 1227–1241.

Benrey B, **Callejas A**, **Rios L**, **Oyama K**, **Denno RF**. **1998**. The effects of domestication
ofBrassicaandPhaseoluson the interaction between phytophagous insects and parasitoids. *Biological*
control: theory and applications in pest management **11**: 130–140.

Bolyen E, **Rideout JR**, **Dillon MR**, **Bokulich NA**, **Abnet CC**, **Al-Ghalith GA**, **Alexander H**, **Alm EJ**,
Arumugam M, **Asnicar F**, *et al.* **2019**. Reproducible, interactive, scalable and extensible microbiome
data science using QIIME 2. *Nature biotechnology* **37**: 852–857.

Bonito G, **Benucci GMN**, **Hameed K**, **Weighill D**, **Jones P**, **Chen K-H**, **Jacobson D**, **Schadt C**,
Vilgalys R. **2019**. Fungal-bacterial networks in the Populus rhizobiome are impacted by soil properties
and host genotype. *Frontiers in microbiology* **10**: 481.

Bouffaud M-L, **Poirier M-A**, **Muller D**, **Moënne-Loccoz Y**. **2014**. Root microbiome relates to plant host
evolution in maize and other Poaceae: Poaceae evolution and root bacteria. *Environmental microbiology*
16: 2804–2814.

Bradshaw AD. **1965**. Evolutionary significance of phenotypic plasticity in plants. In: Caspari EW, Thoday
JM, eds. Advances in genetics. Advances in Genetics. Elsevier, 115–155.

Brestic M, **Zivcak M**, **Hauptvogel P**, **Misheva S**, **Kocheva K**, **Yang X**, **Li X**, **Allakhverdiev SI**. **2018**.
Wheat plant selection for high yields entailed improvement of leaf anatomical and biochemical traits
including tolerance to non-optimal temperature conditions. *Photosynthesis research* **136**: 245–255.

Brown SP, **Grillo MA**, **Podowski JC**, **Heath KD**. **2020**. Soil origin and plant genotype structure distinct
microbiome compartments in the model legume Medicago truncatula. *Microbiome* **8**: 139.

Callahan BJ, **McMurdie PJ**, **Rosen MJ**, **Han AW**, **Johnson AJA**, **Holmes SP**. **2016**. DADA2: High-
resolution sample inference from Illumina amplicon data. *Nature methods* **13**: 581–583.

Caporaso JG, **Lauber CL**, **Walters WA**, **Berg-Lyons D**, **Lozupone CA**, **Turnbaugh PJ**, **Fierer N**,
Knight R. **2011**. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample.
Proceedings of the National Academy of Sciences of the United States of America **108 Suppl 1**: 4516–
4522.

Dal Bello M, **Lee H**, **Goyal A**, **Gore J**. **2021**. Resource-diversity relationships in bacterial communities
reflect the network structure of microbial metabolism. *Nature ecology & evolution* **5**: 1424–1434.

Doebley JF, **Gaut BS**, **Smith BD**. **2006**. The molecular genetics of crop domestication. *Cell* **127**: 1309–
1321.

Edwards J, **Johnson C**, **Santos-Medellín C**, **Lurie E**, **Podishetty NK**, **Bhatnagar S**, **Eisen JA**,
Sundaresan V. **2015**. Structure, variation, and assembly of the root-associated microbiomes of rice.
Proceedings of the National Academy of Sciences of the United States of America **112**: E911-20.

Ellstrand NC, **Prentice HC**, **Hancock JF**. **2002**. Gene flow and introgression from domesticated plants
into their wild relatives. *Horizontal gene transfer*.

Estrada-De Los Santos P, **Bustillos-Cristales R**, **Caballero-Mellado J**. **2001**. Burkholderia, a genus
rich in plant-associated nitrogen fixers with wide environmental and geographic distribution. *Applied and*
environmental microbiology **67**: 2790–2798.

Favela A, **Bohn M**, **Kent A**. **2022**. N-cycling microbiome recruitment differences between modern and
wild *Zea mays*. *Phytobiomes journal* **6**: 151–160.

Friedrich JO, **Adhikari NKJ**, **Beyene J**. **2008**. The ratio of means method as an alternative to mean
differences for analyzing continuous outcome variables in meta-analysis: a simulation study. *BMC*
medical research methodology **8**: 32.

Glasser SK, **de Santiago-Hernández MH**, **Delgado-Carrillo O**, **Espino LAV**, **Pérez AC**, **González-**
Rodríguez A, **Lira-Saade R**, **Quesada M**. **2023**. Influence of plant domestication on plant-pollinator
interactions: Floral attributes and floral visitor communities in wild and cultivated squash plants. *American*
journal of botany **110**: e16170.

Gutierrez A, **Grillo MA**. **2022**. Effects of domestication on plant-microbiome interactions. *Plant & cell*
physiology **63**: 1654–1666.

Janzen GM, **Wang L**, **Hufford MB**. **2019**. The extent of adaptive wild introgression in crops. *The new*
phytologist **221**: 1279–1288.

Knights D, **Kuczynski J**, **Charlson ES**, **Zaneveld J**, **Mozer MC**, **Collman RG**, **Bushman FD**, **Knight R**,
Kelley ST. **2011**. Bayesian community-wide culture-independent microbial source tracking. *Nature*
methods **8**: 761–763.

Koester RP, **Nohl BM**, **Diers BW**, **Ainsworth EA**. **2016**. Has photosynthetic capacity increased with 80
years of soybean breeding? An examination of historical soybean cultivars: Photosynthesis in historical
soybean cultivars. *Plant, cell & environment* **39**: 1058–1067.

Ku Y-S, **Contador CA**, **Ng M-S**, **Yu J**, **Chung G**, **Lam H-M**. **2020**. The effects of domestication on
secondary metabolite composition in legumes. *Frontiers in genetics* **11**: 581357.

Le Corre V, **Siol M**, **Vigouroux Y**, **Tenaillon MI**, **Délye C**. **2020**. Adaptive introgression from maize has
facilitated the establishment of teosinte as a noxious weed in Europe. *Proceedings of the National*
Academy of Sciences of the United States of America **117**: 25618–25627.

Liu F, **Hewezi T**, **Lebeis SL**, **Pantalone V**, **Grewal PS**, **Staton ME**. **2019**. Soil indigenous microbiome
and plant genotypes cooperatively modify soybean rhizosphere microbiome assembly. *BMC microbiology*
19: 201.

Love MI, **Huber W**, **Anders S**. **2014**. Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. *Genome biology* **15**: 550.

McDonald D, **Jiang Y**, **Balaban M**, **Cantrell K**, **Zhu Q**, **Gonzalez A**, **Morton JT**, **Nicolaou G**, **Parks DH**,
Karst SM, *et al.* **2024**. Greengenes2 unifies microbial data in a single reference tree. *Nature*
biotechnology **42**: 715–718.

McMurdie PJ, **Holmes S**. **2013**. phyloseq: an R package for reproducible interactive analysis and
graphics of microbiome census data. *PloS one* **8**: e61217.

Meyer RS, **DuVal AE**, **Jensen HR**. **2012**. Patterns and processes in crop domestication: an historical
review and quantitative analysis of 203 global food crops. *The new phytologist* **196**: 29–48.

Oksanen J. **2017**. *Vegan: Community Ecology Package*.

Pérez-Jaramillo JE, **Carrión VJ**, **de Hollander M**, **Raaijmakers JM**. **2018**. The wild side of plant
microbiomes. *Microbiome* **6**: 143.

Pérez-Jaramillo JE, **de Hollander M**, **Ramírez CA**, **Mendes R**, **Raaijmakers JM**, **Carrión VJ**. **2019**.
Deciphering rhizosphere microbiome assembly of wild and modern common bean (Phaseolus vulgaris) in
native and agricultural soils from Colombia. *Microbiome* **7**: 114.

Porter SS, **Sachs JL**. **2020**. Agriculture and the disruption of plant-microbial symbiosis. *Trends in*
ecology & evolution **35**: 426–439.

Puruganan MD, **Fuller DQ**. **2009**. The nature of selection during plant domestication. *Nature* **457**: 843–
848.

Richards RA. **2000**. Selectable traits to increase crop photosynthesis and yield of grain crops. *Journal of*
experimental botany **51 Spec No**: 447–458.

Sachs JL, **Ehinger MO**, **Simms EL**. **2010**. Origins of cheating and loss of symbiosis in wild
Bradyrhizobium. *Journal of evolutionary biology* **23**: 1075–1089.

Singh J, **van der Knaap E**. **2022**. Unintended consequences of plant domestication. *Plant & cell*
physiology **63**: 1573–1583.

Singh BK, **Trivedi P**, **Egidi E**, **Macdonald CA**, **Delgado-Baquerizo M**. **2020**. Crop microbiome and
sustainable agriculture. *Nature reviews. Microbiology* **18**: 601–602.

Soldan R, **Fusi M**, **Cardinale M**, **Daffonchio D**, **Preston GM**. **2021**. The effect of plant domestication on
host control of the microbiota. *Communications biology* **4**: 936.

Valero-Mora PM. **2010**. Ggplot2:Elegant graphics for data analysis. *Journal of statistical software* **35**: 1–
3.

Viechtbauer W. **2025**. Meta-Analysis Package for R [R package metafor version 4.8-0].

Wagner MR, **Lundberg DS**, **Del Rio TG**, **Tringe SG**, **Dangl JL**, **Mitchell-Olds T**. **2016**. Host genotype
and age shape the leaf and root microbiomes of a wild perennial plant. *Nature communications* **7**: 12151.

Wagner MR, **Tang C**, **Salvato F**, **Clouse KM**, **Bartlett A**, **Vintila S**, **Phillips L**, **Sermons S**, **Hoffmann**
M, **Balint-Kurti PJ**, *et al.* **2021**. Microbe-dependent heterosis in maize. *Proceedings of the National*
Academy of Sciences of the United States of America **118**: e2021965118.

Weese DJ, **Heath KD**, **Dentinger BTM**, **Lau JA**. **2015**. Long-term nitrogen addition causes the evolution
of less-cooperative mutualists. *Evolution; international journal of organic evolution* **69**: 631–642.

Whitehead SR, **Turcotte MM**, **Poveda K**. **2017**. Domestication impacts on plant-herbivore interactions: a
meta-analysis. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*
372.

Willis A, **Bunge J**. **2015**. Estimating diversity via frequency ratios. *Biometrics* **71**: 1042–1049.

Wu Y, **Sun J**, **Yu P**, **Zhang W**, **Lin Y**, **Ma D**. **2022**. The rhizosphere bacterial community contributes to
the nutritional competitive advantage of weedy rice over cultivated rice in paddy soil. *BMC microbiology*
22: 232.

Zhang M, **Wang Y**, **Hu Y**, **Wang H**, **Liu Y**, **Zhao B**, **Zhang J**, **Fang R**, **Yan Y**. **2023**. Heterosis in root
microbiota inhibits growth of soil-borne fungal pathogens in hybrid rice. *Journal of integrative plant biology*
65: 1059–1076.

773