PLANT DOMESTICATION DOES NOT REDUCE DIVERSITY IN ROOT MICROBIOMES

Alejandra Hernández-Terán[1], Ana E. Escalante[2], Maria Rebolleda-Gómez[1]

[1]Ecology and Evolutionary Biology Department, COMMONS Lab, University of California Irvine.

Irvine, California, 92697.

[2]Laboratorio Nacional de Ciencias de la Sostenibilidad, Instituto de Ecología, Universidad Nacional

Autónoma de México, Mexico City, Mexico.

Classification: Biological Sciences - Plant Biology

Keywords: Plant domestication, root microbiome, rhizosphere

ABSTRACT

Domestication has profoundly shaped human civilization and the genetic makeup of numerous plant and animal species. While the effects of plant domestication at the genetic and phenotypic levels are well-documented, its impact on plant microbiome remains less understood. Root microbiomes play crucial roles in nutrient acquisition, pathogen defense, and biotic stress tolerance, yet the influence of domestication on their diversity and assembly is still debated. Two primary hypotheses have been proposed: 1) the reduction in microbial diversity resulting from the domestication process, and 2) the diminished ability of host plants to control their microbiomes. To evaluate these hypotheses, we conducted a meta-analysis of multiple crops, comparing the root microbiomes of domesticated plants and their wild relatives. Our results indicate that the effects of domestication are species-specific and context-dependent, with most domesticated plants exhibiting increased microbial diversity and more structured communities, while others show no significant change. Overall, this study provides evidence that plant domestication does not lead to a uniform reduction in microbial diversity or a consistently diminished ability of plants to influence their microbiomes. Based on these findings, we discuss new perspectives and the need for future studies incorporating native soils and host genetic variation in such experiments, analyzing not only diversity but also microbiome function, and considering how root morphology might affect microbiome recruitment. Finally, we highlight the need for research on the potential adaptive or maladaptive consequences that introgression between wild and domesticated plants could have from a microbiome perspective.

SIGNIFICANCE STATEMENT

Plant domestication has had profound impacts on plant traits and their genetic makeup. These traits are likely to influence microbial interactions with the plants, but the overall effect of domestication on plant microbiomes is still subject to debate. In this work, we challenge two primary hypotheses regarding the effects of domestication on root microbiomes; we compared published microbiomes of wild and domesticated plants and found that domestication does not reduce microbial diversity or the plant's contribution to their associated microbial communities. Instead, these effects vary between plant species and environmental conditions. This work provides insights into how domestication shapes plant-microbe relationships and discusses new approaches that are critical for advancing this area of research.

INTRODUCTION

Domestication, the artificial selection of phenotypic traits useful to humans, has profoundly shaped human civilization and the genetic makeup of numerous plant and animal species [1]. The consequences of domestication at the genetic and phenotypic level are well-documented [2]. Typically, domestication events involve propagating only selected plant seeds, leading to genetic bottlenecks and thus, a reduction in genetic diversity across the genome [3]. Phenotypically, convergent changes in certain traits, known as 'domestication syndrome', have been observed across diverse domesticated species [4]. In plants, this syndrome is characterized by enlarged fruit sizes, increased apical dominance, and the loss of seed dispersal mechanisms and dormancy, among other traits [3]. Many of these traits are the target of selection, but many others are not. Those non-target traits can evolve through correlations (genetic linkages or pleiotropy) with the target traits (e.g. reduction of seed shattering, uniform flowering, and ripening) and can greatly influence the ecology of the plant [1, 2]. The extent to which domestication-induced phenotypic changes affect ecological interactions remains an open question.

It is often proposed that domestication leads to reduced investment in other ecological interactions. In the case of plant-herbivore interactions, a commonly proposed hypothesis suggests that domestication has modified plant traits in ways that decrease resistance to herbivores [9]. Nonetheless, a meta-analysis [10] conducted in 2017 assessed the overall impact of crop domestication on plant-herbivore interactions and found that, while domestication appears to have a generally negative effect on herbivore resistance, it did not significantly alter plant defense traits. Similarly, a recent study [11] exploring the consequences of domestication on plant-pollinator interactions in members of the *Cucurbita* genus revealed that domestication in this group may have promoted greater resource investment in floral traits. This increased investment likely enhances attractiveness to pollinators, potentially increasing reproductive success.

Roots are an important case for the evolution of non-target traits and species interactions. Root

phenotypes have often significantly diverged from their wild relatives due to adaptation to the

agricultural environment or as non-target modifications associated with selected traits above

ground. Domesticated plants often have a different root system architecture, altered root exudate

compositions, and significant changes in secondary metabolites across a wide range of species

(2, 12), which necessarily has an impact in plant interactions including one of the most critical

plant interactions, the symbiotic relationship between microbial communities and plant roots (the

root or rhizosphere microbiome).

Plants rely on their microbiomes for several functions, including nutrient acquisition, protection

against pathogens, and tolerance to biotic stress (13). Often, the root microbiome promotes plant

health and growth, contributing to plant development (14). Given the pivotal importance of root

microbiomes for plants and considering the known effects of plant domestication on root systems,

in recent years efforts have been undertaken to decipher the impact of plant domestication on the

diversity and assembly of these microbial communities. Despite these efforts, it is still unclear if,

or to what extent, plant domestication has modified the ability of the plants to interact with the root

microbiome.

Two main hypotheses have been proposed regarding the consequences of domestication: 1) the

reduction in microbial diversity resulting from the domestication process, and 2) the diminished

ability of host plants to shape the assembly of their microbiomes. These hypotheses arise from

the idea that the genetic bottlenecks accompanying domestication may lead to the loss of alleles

crucial for plant-microbiome interactions and that alterations in root exudates metabolic profiles

might further influence microbiome assembly in domesticated plants (15), and because of a

potential reduction in the benefits of mutualism for domesticated plants. Specifically, domesticated

plants, after growing in environments with important nutrient inputs, could potentially reduce the investment in symbiosis, leading to the evolution of reduced symbiont cooperation (16). To evaluate these hypotheses, we conducted a meta-analysis comparing studies inoculating domesticated plants and their wild relatives with the same bulk soil in common garden conditions. We found that plant domestication does not result in a uniform reduction in microbial diversity or a consistently diminished ability of host plants to influence their microbiomes. The effects of domestication are species-specific and context-dependent, with some domesticated plants showing increased microbial diversity and structured communities, while others do not. Despite this variation, we found consistent losses and enrichments of microbial taxa in domesticated crops relative to their wild progenitors.

RESULTS

We conducted a systematic meta-analysis to assess the influence of plant domestication on root microbiomes across different plant species. After applying our inclusion criteria and filters of data quality, 14 independent studies were retained. These studies encompass eight different crop species and included 882 individual plants belonging to maize (n = 129), wheat (n= 172), barley (n= 55), sunflower (n= 127), tomato (n= 169), cotton (n= 31), bean (n= 39), and soybean (n= 160) (Table S1). Except for bean, soybean, and sunflower, all crops were represented by more than one accession of wild and domesticated plants (Table S1). With this dataset, we investigated two primary hypotheses regarding the impact of plant domestication on root microbiomes: 1) the reduction in microbial diversity resulting from the domestication process, and 2) the diminished ability of host plants to control their microbiomes. This data did not experimentally assess plant "control", hereafter we refer as "host influence" to the contribution of plants in shaping root microbiome assemblies. Our analysis focused on the comparison between wild and domesticated categories. To mitigate the potential bias introduced by merging amplicon sequence data from disparate studies, our comparative analyses were restricted to within-study wild and domesticated pairs (W-D pairs). We defined W-D pairs as plants cultivated concurrently and inoculated with the same soil. Typically, each paper contributed a single W-D pair to our analysis. However, in instances where a study investigated more than one soil type, the number of W-D pairs was equivalent to the number of different soil treatments applied (we analyzed a total of 22 different W-D pairs).

Domesticated plants show increased microbiome diversity

Contrary to the common expectation of reduced diversity in the microbiomes as a result of domestication, on average, domesticated plants had higher alpha diversity according to the Shannon diversity index (Fig. 1 - Shannon index, $p < 0.001$, Fig. S1 - Chao1 index, $p = 0.134$). Most pairwise comparisons (pairs of wild and domesticated plants from the same study, species, and treatment) showed differences in alpha diversity, but the effects were highly variable across species. Among the 22 Wild-Domesticated (W-D) pairs evaluated, only four (18%) exhibited no statistical difference. Of the 18 (81%) pairs that showed statistically significant differences, 11 (61%) displayed increased diversity in domesticated plants, whereas seven (38.8%) had higher diversity in their wild counterparts. Notably, the effect of domestication appears to be species-specific and related to functional groups. Crops like wheat, tomato, and cotton showed increased diversity in their domesticated forms compared to their wild forms, whereas maize and legumes (beans and soybeans) presented greater diversity in their wild forms. Despite these overall patterns, we observed different outcomes, even within the same species. The only exception to this inconsistency was observed in tomatoes, where all four comparisons consistently revealed higher diversity in domesticated plants. The most variable crop species was soybean, where from the six total W-D pairs analyzed, two resulted in higher diversity in domesticated versions, three in higher diversity in wild plants, and one showed no statistically significant difference.

Host species effect is more important than domestication in shaping microbiome composition

We then evaluated the effect of domestication on microbiome composition using PERMANOVA. We found that the domestication category accounted for a relatively small fraction of the variation across most of the crop species studied (Fig. 2 and Fig. S2), suggesting a potential species-

specific effect. For several species, including maize, wheat, barley, sunflower, tomato, and bean,

domestication explained approximately 10% of the observed variance in microbial community

composition. Notably, cotton and soybean (except for two pairs) stood out, with domestication

accounting for a larger portion of the variance, ranging from 12% to 25%. Additionally, we

leveraged experiments that used multiple soil types to inoculate plants, allowing us to test the

contribution of the inoculum to the final microbial community by separating wild and domesticated

plants independently (Table S3). Although only a small set of papers had different soil types, for

barley, different initial soils accounted for a minor portion of the total variation (9.8% for wild plants

and 10.9% for domesticated plants). In contrast, for soybeans, using soils from different origins

accounted for a significant percentage of the variation (~75% for wild and ~78% for domesticated

plants). Taken together, these results suggested that domestication could affect the degree of

host influence, but that there might be variation across species.

Wild relatives do not display systematic evidence of increased host influence of their

microbiomes

To evaluate the extent of host influence exerted by wild and domesticated plants over their

microbiomes, we leveraged beta diversity metrics, comparing the microbial communities

associated with plants to those from bulk soils, which served as a reference for the initial microbial

community (Fig. 3). We hypothesized that if plants have a relatively strong influence in the

assembly of their microbiomes, then we would expect to find: a) significant dissimilarity between

plant-associated microbiomes and bulk soil microbiomes (Fig. 3A), suggesting that plants are

selecting specific microbial assemblages distinct from the initial soil; b) low intra-group dispersion

(Fig. 3B), indicating that plant replicates of the same genotype consistently select for similar

microbial compositions; and c) less proportion of the final community explained by the initial bulk

soil (Fig. 3C), implying that the host plays a decisive role in shaping its microbiome, potentially

steering it away from the original soil community composition.

Overall, we did not identify a consistent pattern linking host influence to the category of

domestication. In general, we found greater dissimilarity to bulk in wild barley and cotton and in

domesticated wheat and soybean (Fig. 3A). Similarly, we found a diverse range of outcomes in

the intra-group dispersion analyses (Fig. 3B). Domesticated plants generally exhibited more intra-

group dispersion in barley and soybean, whereas we found the opposite effect (more intra-group

dispersion in wild varieties) in wheat, cotton, and bean. Maize, sunflower, and tomato showed no

differences. We didn't find a clear pattern in the contribution of bulk soil microbiomes to the final

plant-associated communities across different levels of domestication (Fig. 3C).

We explored if some other plant attribute could better account for our results. Grouping our results

by functional plant group rather than domestication status provided increased insight.

Grasses (maize, wheat, barley) consistently demonstrated a more distinct divergence from their

initial bulk soil microbiomes (Fig. 3A), with less than 50% of the final community composition

attributable to the bulk soil (Fig. 3C). Whereas the legume soybean presented final communities

closely aligned with the initial bulk soil, reflecting less than 50% differentiation.

Some microbial taxa differentially associate with plants across species according to the

domestication status

Finally, we investigated if*,* despite all the variation, we could identify microbial groups differentially

present in wild and domesticated crops (see methods; Fig. 4). We identified a few families

consistently enriched or depleted from domesticated crops (relative to their wild counterparts)

across different crops. Two members of the *Rhizobiaceae* family

(*Rhizobiaceae_A_500471* and *Rhizobiaceae_A_500279*) were significantly enriched in the

domesticated variants of wheat, tomato, and soybean. Whereas two members of the

Burkholderiaceae family (*Burkholderiaceae_A_592522* and *Burkholderiaceae_A_574758)* were

consistently enriched in the wild counterparts of wheat, tomato, bean, and soybean. We found

these *Burkholderiaceae* taxa in both the domesticated and the wild rhizosphere. In contrast, the

Rhizobiaceae members appeared almost absent in the wild versions of wheat and tomato

(minimum absolute abundance < 50 reads; Fig. S3).

DISCUSSION

Our study provides new insights into the relationship between plant domestication and microbial diversity and assembly, challenging the widely held belief that domestication always leads to a decrease in microbial diversity and impairs the plant's influence in the assembly of its microbiome. Our results indicate that while domestication does have an effect on microbial diversity and community composition, these effects vary widely across species and experimental conditions.

Contrary to the hypothesis that domestication generally reduces microbial diversity, we found a small but significant reduction of microbiome diversity in the rhizosphere of wild relatives (Fig. 1, Log Ratio of Means with Shannon index, $p < 0.001$). In fact, for the Shannon index, most of the significant comparisons indicated that domesticated plants exhibited higher microbial diversity than wild plants (Fig. 1). When using the Chao1 diversity index, we observed fewer differences and mostly saw no change in diversity across comparisons. The hypothesis that domestication leads to a reduction in microbiome diversity is largely based on the idea that domestication reduces genetic diversity, potentially leading to the loss of alleles important for plant-microbe interactions through selective bottlenecks [15] and from the hypothesis in domesticated plants, a reduced investment in symbiosis, could be promoting the evolution of reduced symbiont cooperation [16].

It is not entirely clear why the reduction of genetic diversity at the level of populations would lead to reduced microbiome diversity at the level of individuals. Instead, we might expect a decrease in microbiome diversity across hosts (less dispersion) in domesticated varieties compared to their wild counterparts. However, our results highlight that this is only the case in some pairs –mostly wheat, barley, and cotton. The opposite is true in some other pairs, including many of the soybean comparisons (Fig. 3). One of the limitations in making clear conclusions about the impact of host

diversity on the microbiome is that host diversity is rarely controlled in these studies (many

domesticated accessions might be compared against a single population of wild relatives).

Even the loss of alleles important for plant-microbe interactions does not have to be associated

with decreased microbiome diversity. It has been proposed, for example, that a reduction in

genetic diversity might lead to less efficient microbiome filtering mechanisms, allowing

domesticated plants to be less selective and thus associate with a greater diversity of microbes

(15, 17, 18). This hypothesis has gained attention due to its potential implications for agriculture

and resilience to climate change. The variation in microbial communities explained by host

genotype represents the substrate upon which natural selection can act and, therefore, it is also

the variation that could be manipulated for agricultural improvement (19). Thus, a reduction in

host influence resulting from domestication could lead to decreased evolutionary potential within

microbial communities, potentially limiting the adaptive resources of plants in response to

environmental stresses (18). This hypothesis of less efficient filtering mechanisms in

domesticated crops is consistent with our alpha (within-plant) diversity results, where we observed

a small increase in microbiome diversity in domesticated varieties. But, contrary to this

expectation, we found little support for decreased host influence in domesticated plants. In most

pairs, the microbial community of the rhizosphere is equally distant to the bulk soil with soybean

being the only consistent example of increased distance to bulk soil in the domesticated varieties

(even if most are not significant) (Fig. 3).

Until recently (4), the existence of a domestication syndrome had promoted the idea that most

plant species that undergo domestication suffer the same phenotypic consequences. Likewise,

this could have also promoted the assumption that, for all crop species, domestication would

always negatively affect microbial diversity. In addition to finding, on average, increased microbial

diversity in domesticated plants, our analysis revealed significant variability inside species. For

many of the crops analyzed, there were instances where wild plants exhibited higher microbial

diversity, while in other cases, domesticated plants showed greater diversity. We found this

variability in different studies using the same plant accessions and even within the same study

when different initial soils were used, implying a significant role of genotype-by-environment (GxE)

interactions in shaping microbiome assembly.

It is known that microbial communities from different habitats (i.e., different initial soils) can exhibit

distinct responses to the same host genotype. It has been proposed that environmental

heterogeneity can decouple host genotypes from their microbiomes by altering genotype-microbe

relationships (20, 21), potentially promoting mixed results even when controlling for host

genotypes. Our results suggest that domestication's effects on microbial diversity are not

universally negative and may vary depending on the specific crop species and environmental

conditions involved. This observation is further supported when examining the impact of

domestication on microbial community composition. Although most comparisons showed

differences in beta diversity (Fig. S2), domestication status accounted for a relatively minor

fraction of the microbial variation across the analyzed species (Fig. 2). Instead, much of the

variation, at least in studies using more than one soil type, is explained by the soil origin (Table

S3). This finding aligns with the results of individual studies on rice (22), poplar (23), soybean

(24), and the legume *Medicago truncatula* (19), where soil origin is a stronger predictor of

community composition than host genotype. Some of these studies suggest that endophytes are

more sensitive to host genotype signals, while rhizosphere communities are more affected by the

environment (19). Future studies looking at the effect of domestication should aim to separate the

effects of domestication on different microbial functional groups.

Despite all the variation found between crops, we detected two members of the *Rhizobiaceae*

family consistently enriched in domesticated wheat, tomato, and soybean, and two members of

the *Burkholderiaceae* family enriched in wild wheat, tomato, bean, and soybean (Fig. 4). In some crop species, domesticated plants exhibit increased photosynthetic capacity compared to wild relatives, which is thought to support the development of larger grains or fruits (25, 26). This greater carbon fixation could be increasing the root exudation in domesticated plants, potentially attracting more rhizobia and other copiotrophs in the rhizosphere (27). Additionally, prolonged nitrogen fertilization in agricultural systems is proposed to promote the evolution of less-cooperative rhizobia in domesticated cultivars, potentially leading to higher root abundance of *Rhizobiaceae* members, many of which may not contribute significantly to plant nutrition (24, 28). Particularly, rhizobia with poor or null symbiotic effectiveness are more prevalent in agricultural soils and are predominantly associated with cultivated crops (29). Thus, a high abundance of those microbes does not necessarily indicate a higher nitrogen fixation in the rhizosphere of domesticated crops.

On the other hand, the consistent enrichment of *Burkholderiaceae* family members in the wild versions of multiple species aligns with previous research on common beans and rice (30, 31)). Although the mechanisms behind wild plants harboring an increased abundance of those particular microbes remain unknown, it has been shown that members of the *Burkholderiaceae* family are known for their ability to promote plant growth and suppress soil-borne pathogens, supporting plant health overall (32). One potential explanation is that compared to domesticated plants, wild relatives produce a greater diversity of secondary metabolites, which may promote the recruitment of these beneficial microbes (33). Nonetheless, for the case of the common bean, positive associations with *Burkholderiaceae* family members were found only when using native soils (e.g. versus agricultural soils) (30), suggesting potential local adaptation to microbes found within the natural distribution of such species.

CONCLUSIONS AND FUTURE DIRECTIONS

In this study, we challenged accepted hypotheses regarding the consequences of plant domestication on root microbiomes. Contrary to the expectation that domestication generally reduces microbial diversity, we found a small but significant reduction of microbiome diversity in the rhizosphere of wild relatives. Furthermore, by analyzing multiple crop species, we did not find sufficient evidence to support the hypothesis that domestication negatively affects host influence on the assembly of the microbiome. The consistent enrichment of certain bacterial families in both wild or domesticated versions of different crop species (*Rhizobiaceae* for domesticated and *Burkholderiaceae* for wild plants) could be suggesting, at least for the data analyzed in this work, that the effects of domestication may be stronger at the level of recruiting specific microbes rather than affecting the entire community. Overall, our results suggest that the impacts of domestication on the root microbiomes are not universally negative in terms of microbial diversity. Instead, they appear to be species-specific and contingent upon environmental factors.

Despite our findings, much work remains to fully understand the ecological and evolutionary consequences of plant domestication on root microbiomes. First, diversity is not everything; characterizing functional traits of microbiomes associated with wild and domesticated plants is crucial to better understanding the consequences of domestication for plant-microbe interactions. Second, it is possible that the effect of domestication would be more pronounced with close symbionts and endophytes, more work is necessary to understand how domestication can alter these interactions. Third, it is well-documented that domestication significantly impacts root system architecture, but there is limited research on how such changes affect the microbes inhabiting the roots. How changes in root size, phenotype, and physical properties of the roots impact the recruitment of microbes remains as open questions. Similarly, although exudates have

been studied more than the physical structure of roots for microbiome recruitment, this is still an

area where more research is needed to identify general patterns.

Critical directions for future research are to incorporate plant diversity in the design or analysis of

the experiments to start uncovering the connection between genetic changes of domesticated

crops and their impact on the microbiome. In particular, controlling for genetic diversity between

plants is important to be able to attribute differences between groups to the process of

domestication itself. Similarly, to evaluate the role of domestication on the loss of host influence

it will be important to incorporate native soils into common garden experiments. Local adaptation

is important for wild populations, and relying exclusively on agricultural soils (or other soils outside

the natural range of wild plants) may hinder significant microbial interactions that occur only with

the specific set of microbes to which wild plants are naturally adapted.

Finally, introgression between wild and domesticated plants is a common phenomenon,

especially at the centers of origin of the species (34). The evolutionary consequences of this have

been extensively studied from a genetic perspective (35), and as a source for key agronomic traits

(36). Nonetheless, even though there is increasing evidence showing that hybridization impacts

root microbiomes (37, 38), the consequences of introgression between wild and domesticated

plants remain unclear. Recently, studies on rice (38) and maize (39) have found that both root

and seed microbiomes could contribute to heterosis ("hybrid vigor") in hybrids, providing evidence

of the important role of microbiomes in the fitness of inbreds. Studying the potential adaptive and

maladaptive consequences of microbiome introgression is a new and exciting area of study, with

significant implications for plant breeding and conservation biology.

MATERIALS AND METHODS

Data collection

To investigate the impact of the domestication process on the diversity and composition of the root microbiome across various crop species, we collected 16S rRNA sequence data from common garden experiments reported in scientific publications. We conducted a comprehensive search in the Scopus and Google Scholar databases using the Boolean search string: Wild [AND] domesticated [AND] microbiome [OR] rhizosphere [AND] crop. Our search centered on contemporary crops with well-identified crop wild relatives, including but not limited to maize, potato, tomato, squash, bean, coffee, barley, soybean, cotton, and sunflower. Eligibility for inclusion in our meta-analysis was based on the following selection criteria: 1) the experiments must have been conducted under controlled environmental conditions typical of common garden studies, 2) wild and domesticated variants of the same crop species must have been cultivated concurrently, and inoculated with the same soil, and 3) sequencing must have been performed using Illumina platforms. We excluded studies that assessed the impact of extreme environmental stressors, such as herbivory, but we incorporated those that utilized different soil types (e.g., forest or agricultural soils) as inoculum.

Applying these criteria, we narrowed down the pool to 14 publications, representing eight crops: barley, soybean, cotton, tomato, potato, sunflower, maize, and bean. A detailed overview of the number of crop accessions, experimental replicates, and analyzed publications per crop is available in Supplementary Table 1.

Data processing

16S rRNA raw sequences and metadata were downloaded from public repositories including

NCBI, MG-Rast, and Github, and were analyzed with the Quantitative Insights Into Microbial

Ecology (QIIME2) platform (40) version 2023.5. For each dataset corresponding to a specific

publication, paired-end sequences were analyzed independently. Sequence denoising and

quality filtering, including primers and adapters trimming, were performed with the DADA2 plug-

in (41). Given that all selected studies utilized primers targeting the V4 16S rRNA gene region

(e.g., V4, V3-V4, V4-V5), we standardized the analysis by extracting the V4 segment using the

515F–806R primer pairs (42). This approach minimized discrepancies arising from taxonomic

assignments across different regions of the 16S rRNA gene. The length of the V4 region

sequences was standardized to a range between 120 and 150 nucleotides, based on sequence

quality and primer positioning. This criterion was consistently applied across all datasets.

Taxonomic classification of the Amplicon Sequence Variants (ASVs) for each dataset was

conducted using the "sklearn-base" classifier within QIIME2, referencing the Greengenes 2

database (43) with a 99% sequence identity threshold. Mitochondrial, chloroplast, and ASVs that

were not assignable at the kingdom level were excluded from the analysis. Comprehensive details

on the total number of reads (before and after processing) and the count of ASVs for each study

have been compiled in Supplementary Table 1.

Data Analysis

Data analysis was performed on R software (v.4.3.0) using custom scripts and employing the

following packages: Phyloseq (v. 1.46.0) (44), ggplot2 (v. 3.5.1) (45), metafor (v. 4.2-0) (REF),

Vegan (v.2.6-4) (46), DESEq2 (v.1.40.2) (47), Source Tracker 2 (48). Prior to amplicon analysis,

each of the 14 datasets, corresponding to the individual studies included in our meta-analysis,

was rarified to a standardized depth of 15,000 reads per sample to ensure comparability across

datasets. Specifics on the number of samples and ASVs retained after rarification are available

in Supplementary Table 1. Our analysis focused on the comparison between wild and

domesticated categories. To mitigate the potential bias introduced by merging amplicon sequence

data from disparate studies, our comparative analyses were restricted to within-study wild and

domesticated pairs (W-D pairs). We defined W-D pairs as plants cultivated concurrently and

inoculated with the same soil. Typically, each paper contributed a single W-D pair to our analysis.

However, in instances where a study investigated more than one soil type, the number of W-D

pairs was equivalent to the number of different soil treatments applied.

Alpha diversity analyses: To assess the effect of plant domestication on the diversity of the

microbiome we calculated Chao1 and Shannon alpha diversity indexes using the *estimate*

richness function in the phyloseq package. To test for differences in these indexes we applied the

Log Ratio of Means (LogRM) analysis which compensates for the non-independence of the data

by calculating the effect size and variance for each study individually (49). We implemented this

test with the *metafor* package, using W-D pairs from all crops and the diversity of the wild plants

as a reference.

Beta diversity analyses: To explore potential differences in beta diversity between plant

categories and crops, we calculated the Bray-Curtis distance for every W-D pair in the dataset.

Dissimilarity matrices were analyzed with Permutational Multivariate Analysis of Variance

(PERMANOVA) using the *Adonis* function in the vegan package. From the PERMANOVA result,

we extracted the R^2 values to represent the proportion of variation (in terms of beta diversity)

explained by category for all W-D pairs and all crops. A dot plot was constructed with this

information using ggplot2.

Furthermore, from the dissimilarity matrices we calculated, for all crops, the mean dissimilarity to

the bulk of both wild and domesticated plants, and the mean distance to centroid between groups

using the *betadisper* function in the vegan package. We plotted both metrics for all W-D pairs in

the different crops using ggplot2. Finally, a Bayesian microbial source tracker was performed with

the SourceTracker2 to calculate the contribution of the bulk soil microbial community to the final

root microbiome. All samples were rarefied to a depth of 15,000 reads per sample before the

analysis.

Differentially enriched microbes between categories: We identified differentially enriched

microbes between W-D pairs across all crops using the DESeq2 test, implemented in the DESeq2

package. This test was applied exclusively to W-D pairs (excluding bulk soil sequences), with a

P-value cutoff of <0.01 and a False Discovery Rate (FDR) correction. Subsequently, we merged

the results from all W-D comparisons with differentially abundant features at the family level and

selected only features with a log2Fold change of ≤-5 or ≥5, or those present in two or more crops.

The results were visualized by crop using forest plots constructed in ggplot2. Additionally, we

plotted the individual abundances of microbial families enriched in the same category across

multiple crop species (Supplementary Figure 3).

ACKNOWLEDGMENTS AND FUNDING SOURCES

We would like to thank Brandon Gaut, Eduardo Choreño-Parra, Santiago Rosas-Plaza, Celia

Symons, and members of the COMMONS Lab for productive discussions about this work. AHT

was supported by the Human Frontiers Science Program Long-Term Postdoctoral Fellowship

(LT 0047/2023-L) and the University of California President's Postdoctoral Fellowship. AEH was

supported by UNAM-PAPIIT: IN214124. MRG was supported by the School of Biological

Sciences at the University of California, Irvine (UCI).

AUTHOR CONTRIBUTIONS

AHT and MRG designed the research with input from AE. AHT performed all statistical

analyses. AHT wrote the manuscript with substantial feedback from MRG and AE.

COMPETING INTERESTS

The authors declare no competing interests.

DATA AVAILABILITY STATEMENT

Data and R code for analyses is available on Github at:

https://github.com/AlejandraHT/Plant_domestication_microbiomes

Figure 1. Effect of Domestication on Microbial Alpha Diversity Across Different Crop Species. Forest

plot showing the log Ratio of Means (LROM) using Chao1 diversity index for all 22 Wild-Domesticated (W-

D) pairs. The mean microbial diversity of wild plants was used as reference. Dashed lines denote the

separation between different plant species. From top to down: Maize, Wheat, Barley, Sunflower, Tomato,

Cotton, Bean, and Soybean. In the color code: red colors represent grasses and blue colors represent

legumes. Bars represent the mean response ratios with 95% confidence intervals. Colors of the bars

indicate the results: dark green represents higher diversity in wild plants, light green represents higher

diversity in domesticated plants, and grey indicates no statistical difference.

Figure 2. Effect of Domestication on Microbial Community Composition Across Different Crop

Species. Dot plot showing the percentage of variation in microbial community composition explained by

domestication category (wild vs. domesticated), according to the Bray-Curtis distance measure. Numbers

in X axis represent the number of Wild-Domesticated pairs (pairs of wild and domesticated plants from the

same study, species, and treatment) analyzed per crop.

Figure 3. Impact of Domestication on Host Influence of the Microbiome

A) Bray-Curtis dissimilarity to bulk soil for wild-domesticated (W-D) pairs across different crop species.

Statistically significant differences are indicated by an asterisk (Wilcoxon rank-sum test, p < 0.05). This

analysis includes only experiments that included bulk soil sequences. **B)** Intra-group dispersion for all W-

D pairs analyzed using Bray-Curtis distance. Statistically significant differences are indicated by an

asterisk (Wilcoxon rank-sum test, p < 0.05). **C)** Source Tracker analysis estimating the contribution of bulk

soil to the final microbial community of wild and domesticated plants across different species.

Figure 4. Differentially enriched microbial families between wild and domesticated plants across various crop species. DESeq analysis at the family level displaying microbes either enriched in two or more crop species or exhibiting a fold change greater than +5 or less than -5.

References

1. M. D. Purugganan, D. Q. Fuller, The nature of selection during plant domestication. *Nature* **457**,
843–848 (2009).

2. R. S. Meyer, A. E. DuVal, H. R. Jensen, Patterns and processes in crop domestication: an
historical review and quantitative analysis of 203 global food crops. *New Phytol.* **196**, 29–48
(2012).

3. J. F. Doebley, B. S. Gaut, B. D. Smith, The molecular genetics of crop domestication. *Cell* **127**,
1309–1321 (2006).

4. O. Alam, M. D. Purugganan, Domestication and the evolution of crops: variable syndromes,
complex genetic architectures, and ecological entanglements. *Plant Cell* **36**, 1227–1241 (2024).

5. M. Chacón-Fuentes, *et al.*, Domestication in murtilla (Ugni molinae) reduced defensive flavonol
levels but increased resistance against a native herbivorous insect. *Environ. Entomol.* **44**, 627–
637 (2015).

6. M. M. Turcotte, N. E. Turley, M. T. J. Johnson, The impact of domestication on resistance to two
generalist herbivores across 29 independent domestication events. *New Phytol.* **204**, 671–681
(2014).

7. A. R. Fernandez, A. Sáez, C. Quintero, G. Gleiser, M. A. Aizen, Intentional and unintentional
selection during plant domestication: herbivore damage, plant defensive traits and nutritional
quality of fruit and seed crops. *New Phytol.* **231**, 1586–1598 (2021).

8. Y. H. Chen, R. Gols, B. Benrey, Crop domestication and its impact on naturally selected trophic
interactions. *Annu. Rev. Entomol.* **60**, 35–58 (2015).

9. B. Benrey, A. Callejas, L. Rios, K. Oyama, R. F. Denno, The effects of domestication
ofBrassicaandPhaseoluson the interaction between phytophagous insects and parasitoids. *Biol.*
Control **11**, 130–140 (1998).

10. S. R. Whitehead, M. M. Turcotte, K. Poveda, Domestication impacts on plant-herbivore
interactions: a meta-analysis. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **372** (2017).

11. S. K. Glasser, *et al.*, Influence of plant domestication on plant-pollinator interactions: Floral
attributes and floral visitor communities in wild and cultivated squash plants. *Am. J. Bot.* **110**,
e16170 (2023).

12. J. Singh, E. van der Knaap, Unintended consequences of plant domestication. *Plant Cell Physiol.*
63, 1573–1583 (2022).

13. J. E. Pérez-Jaramillo, V. J. Carrión, M. de Hollander, J. M. Raaijmakers, The wild side of plant
microbiomes. *Microbiome* **6**, 143 (2018).

14. B. K. Singh, P. Trivedi, E. Egidi, C. A. Macdonald, M. Delgado-Baquerizo, Crop microbiome and
sustainable agriculture. *Nat. Rev. Microbiol.* **18**, 601–602 (2020).

15. A. Gutierrez, M. A. Grillo, Effects of domestication on plant-microbiome interactions. *Plant Cell*
Physiol. **63**, 1654–1666 (2022).

16. S. S. Porter, J. L. Sachs, Agriculture and the disruption of plant-microbial symbiosis. *Trends Ecol.*
Evol. **35**, 426–439 (2020).

17. A. Favela, M. Bohn, A. Kent, N-cycling microbiome recruitment differences between modern and
wild *Zea mays*. *Phytobiomes J.* **6**, 151–160 (2022).

18. R. Soldan, M. Fusi, M. Cardinale, D. Daffonchio, G. M. Preston, The effect of plant domestication
on host control of the microbiota. *Commun. Biol.* **4**, 936 (2021).

19. S. P. Brown, M. A. Grillo, J. C. Podowski, K. D. Heath, Soil origin and plant genotype structure
distinct microbiome compartments in the model legume Medicago truncatula. *Microbiome* **8**, 139
(2020).

20. M. R. Wagner, *et al.*, Host genotype and age shape the leaf and root microbiomes of a wild
perennial plant. *Nat. Commun.* **7**, 12151 (2016).

21. A. D. Bradshaw, "Evolutionary significance of phenotypic plasticity in plants" in *Advances in*
Genetics, Advances in genetics., E. W. Caspari, J. M. Thoday, Eds. (Elsevier, 1965), pp. 115–
155.

22. J. Edwards, *et al.*, Structure, variation, and assembly of the root-associated microbiomes of rice.
Proc. Natl. Acad. Sci. U. S. A. **112**, E911-20 (2015).

23. G. Bonito, *et al.*, Fungal-bacterial networks in the Populus rhizobiome are impacted by soil
properties and host genotype. *Front. Microbiol.* **10**, 481 (2019).

24. F. Liu, *et al.*, Soil indigenous microbiome and plant genotypes cooperatively modify soybean
rhizosphere microbiome assembly. *BMC Microbiol.* **19**, 201 (2019).

25. M. Brestic, *et al.*, Wheat plant selection for high yields entailed improvement of leaf anatomical
and biochemical traits including tolerance to non-optimal temperature conditions. *Photosynth.*
Res. **136**, 245–255 (2018).

26. R. P. Koester, B. M. Nohl, B. W. Diers, E. A. Ainsworth, Has photosynthetic capacity increased
with 80 years of soybean breeding? An examination of historical soybean cultivars:
Photosynthesis in historical soybean cultivars. *Plant Cell Environ.* **39**, 1058–1067 (2016).

27. R. A. Richards, Selectable traits to increase crop photosynthesis and yield of grain crops. *J. Exp.*
Bot. **51 Spec No**, 447–458 (2000).

28. D. J. Weese, K. D. Heath, B. T. M. Dentinger, J. A. Lau, Long-term nitrogen addition causes the
evolution of less-cooperative mutualists. *Evolution* **69**, 631–642 (2015).

29. J. L. Sachs, M. O. Ehinger, E. L. Simms, Origins of cheating and loss of symbiosis in wild
Bradyrhizobium. *J. Evol. Biol.* **23**, 1075–1089 (2010).

30. J. E. Pérez-Jaramillo, *et al.*, Deciphering rhizosphere microbiome assembly of wild and modern
common bean (Phaseolus vulgaris) in native and agricultural soils from Colombia. *Microbiome* **7**,
114 (2019).

31. Y. Wu, *et al.*, The rhizosphere bacterial community contributes to the nutritional competitive
advantage of weedy rice over cultivated rice in paddy soil. *BMC Microbiol.* **22**, 232 (2022).

32. P. Estrada-De Los Santos, R. Bustillos-Cristales, J. Caballero-Mellado, Burkholderia, a genus
rich in plant-associated nitrogen fixers with wide environmental and geographic distribution.
Appl. Environ. Microbiol. **67**, 2790–2798 (2001).

33. Y.-S. Ku, *et al.*, The effects of domestication on secondary metabolite composition in legumes.
Front. Genet. **11**, 581357 (2020).

34. N. C. Ellstrand, H. C. Prentice, J. F. Hancock, Gene flow and introgression from domesticated
plants into their wild relatives. *Horizontal gene transfer* (2002).

35. G. M. Janzen, L. Wang, M. B. Hufford, The extent of adaptive wild introgression in crops. *New
Phytol.* **221**, 1279–1288 (2019).

36. V. Le Corre, M. Siol, Y. Vigouroux, M. I. Tenaillon, C. Délye, Adaptive introgression from maize
has facilitated the establishment of teosinte as a noxious weed in Europe. *Proc. Natl. Acad. Sci.*
U. S. A. **117**, 25618–25627 (2020).

37. M.-L. Bouffaud, M.-A. Poirier, D. Muller, Y. Moënne-Loccoz, Root microbiome relates to plant
host evolution in maize and other Poaceae: Poaceae evolution and root bacteria. *Environ.*
Microbiol. **16**, 2804–2814 (2014).

38. M. Zhang, *et al.*, Heterosis in root microbiota inhibits growth of soil-borne fungal pathogens in
hybrid rice. *J. Integr. Plant Biol.* **65**, 1059–1076 (2023).

39. M. R. Wagner, *et al.*, Microbe-dependent heterosis in maize. *Proc. Natl. Acad. Sci. U. S. A.* **118**,
e2021965118 (2021).

40. E. Bolyen, *et al.*, Reproducible, interactive, scalable and extensible microbiome data science
using QIIME 2. *Nat. Biotechnol.* **37**, 852–857 (2019).

41. B. J. Callahan, *et al.*, DADA2: High-resolution sample inference from Illumina amplicon data. *Nat.*
Methods **13**, 581–583 (2016).

42. J. G. Caporaso, *et al.*, Global patterns of 16S rRNA diversity at a depth of millions of sequences
per sample. *Proc. Natl. Acad. Sci. U. S. A.* **108 Suppl 1**, 4516–4522 (2011).

43. D. McDonald, *et al.*, Greengenes2 unifies microbial data in a single reference tree. *Nat.*
Biotechnol. **42**, 715–718 (2024).

44. P. J. McMurdie, S. Holmes, phyloseq: an R package for reproducible interactive analysis and
graphics of microbiome census data. *PLoS One* **8**, e61217 (2013).

45. P. M. Valero-Mora, Ggplot2:Elegant graphics for data analysis. *J. Stat. Softw.* **35**, 1–3 (2010).

46. J. Oksanen, *Vegan: Community Ecology Package* (2017).

47. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

48. D. Knights, *et al.*, Bayesian community-wide culture-independent microbial source tracking. *Nat.*
Methods **8**, 761–763 (2011).

49. J. O. Friedrich, N. K. J. Adhikari, J. Beyene, The ratio of means method as an alternative to mean
differences for analyzing continuous outcome variables in meta-analysis: a simulation study.
BMC Med. Res. Methodol. **8**, 32 (2008).

757