1

Title:

Strand swapping in the LolA-like protein GerS promotes the allosteric activation of a bacterial

amidase in *Clostridioides difficile*

Authors:

Jacob M. Bouchier[1] and Aimee Shen[1]

Affiliations:

[1]Department of Molecular Biology and Microbiology, Tufts University School of Medicine,

Boston, MA, USA

Running head: Strand-swapping in GerS promotes the allosteric activation of CwlD

#**Corresponding author:**

aimee.shen@tufts.edu

(617)636-3792

Abstract:

GerS is a key lipoprotein regulator of *Clostridioides difficile* spore germination that shares

structural, but not sequence, similarity to LolA in Gram-negative bacteria. GerS and LolA have

vastly different biological roles: LolA functions in the trafficking of lipoproteins across the

periplasmic space to the outer membrane of Gram-negative bacteria, while GerS allosterically

activates the germination-specific amidase CwlD in endospore-forming Gram-positive bacteria.

Here, we explore the diversity of LolA-like proteins across bacteria and reveal that GerS

homodimer formation is a conserved feature of Peptostreptococcaceae family orthologs. Since

dimerization of *C. difficile* GerS occurs via strand-swapping, we sought to investigate the

functional importance of its strand swap by identifying mutations that alter the GerS

dimer:monomer equilibrium. Our analyses reveal that GerS^{K71R} and GerS^{N74L} substitutions in the

strand-swap hinge of GerS stabilize the homodimer, while a GerS^{T72P} substitution promotes

monomer formation. Conversely, substituting the highly conserved proline at the equivalent site

in LolA for threonine converts LolA from a monomer to a homodimer. Since our data indicate

that destabilizing the GerS dimer impairs both GerS:CwlD binding *in vitro* and CwlD function in

C. difficile, strand-swapped dimer formation in LolA-like proteins enhances their affinity for their

binding partners. These data are consistent with analyses of equivalent LolA mutants in *E. coli*

and thus provide new insight into the evolution and specialization of this intriguing group of

proteins across the bacterial domain.

Introduction

The InterPro homologous superfamily "Prokaryotic lipoproteins and lipoprotein

localization factors" (PLLLF; SSF89392) encompasses a group of structurally similar proteins

that are widely distributed across the bacterial domain. While some proteins of this group

function as anti-sigma factors [1,2], most function as chaperones for transporting proteins or

glycolipids across the periplasmic space of diderm bacteria. For example, LolA, the archetypal

PLLLF protein, functions within the Lol system (LolABCDE) to facilitate lipoprotein trafficking

across the periplasm in diderm bacteria [3–9], while LppX [10], LprG [11], and LprF [12]

transport various glycolipids across the periplasm of *Mycobacterium* spp. In contrast, PLLLF

proteins in endospore-forming monoderm bacteria seem to have evolved specialized roles in

sporulation and germination. For example, SsdC from *Bacillus subtilis* forms a ring-like structure

in the forespore that functions in the development of spore shape [13], and GerS regulates

spore germination in *Clostridioides difficile* [14].

Spore germination depends on the enzymatic degradation of the cortex, a protective

layer of modified peptidoglycan that maintains metabolic dormancy [15,16]. The cortex lytic

enzymes that degrade the cortex recognize spore-specific peptidoglycan modifications found

exclusively within the cortex, namely muramyl-δ-lactam (MAL) [17]. The specificity of these

enzymes for MAL limits their degradative activities to the cortex layer. MAL is made by the

coordinated activities of the germination-specific amidase CwlD and the peptidoglycan

deacetylase PdaA [18–21]. While *B. subtilis* CwlD has intrinsic activity, *C. difficile* CwlD requires

an additional factor to license its activity, the GerS lipoprotein [22,23]. GerS forms a 1:1 complex

with CwlD and allosterically activates CwlD's amidase activity by stabilizing CwlD's binding to its

Zn^{2+} cofactor [23]. This interaction is essential for germination because deletion of *gerS*

abolishes spore germination in *C. difficile* [14] and disruption of a salt bridge formed between

aspartate 106 of GerS and arginine 169 of CwlD reduces germination [23]. However, the

reduction in *C. difficile* spore germination caused by the GerS D106R substitution is relatively

subtle, even though it significantly reduces binding to CwlD *in vitro* [23]. These data indicate that

additional factors beyond the salt bridge regulate the GerS:CwlD interaction and thus the

amidase activity of CwlD.

In the crystal structure of the GerS:CwlD complex, GerS forms a homodimer via a

strand-swap mechanism [23]. Strand swapping, a specific form of domain swapping [24–31], is

a dimerization mechanism in which one or more β-strands are exchanged between two proteins

[32]. Dimerization via strand swapping is observed in the dimeric structures found in the light

chain subunits of immunoglobulins [33] and cystatins [34], and it is thought to be associated with

pathogenic amyloid fibril formation in humans [35–37]. Strand swapping can also regulate

protein function [38]. For example, monomeric human cystatin C inhibits cathepsin B, but its

strand-swapped dimer loses this ability [39], and strand swapping in cadherins facilitates cell-

cell adhesion, forming weak, reversible interactions [28,40]. However, the functional significance

of strand swapping in GerS on the allosteric activation of CwlD by GerS remains unclear.

Notably, *C. difficile* GerS is the only known PLLLF protein that naturally dimerizes via a

strand swap mechanism, whereas LolA from *Escherichia coli* can be induced to dimerize when

Phe68 is substituted with Glu, which induces a similar strand swap [41,42]; the F68E

substitution was designated F47E based on the sequence after lipoprotein signal peptide

processing. This mutation is toxic to *E. coli* because LolA[F68E] binds LolC too tightly and prevents

proper lipoprotein trafficking to the outer membrane [42]. Here, we test the impact of strand

swapping in GerS on CwlD function. By identifying mutations that shift the GerS dimer:monomer

equilibrium, we demonstrate that strand-swapping in GerS stabilizes its interaction with CwlD

and promotes the allosteric activation of this important amidase. Our analyses provide

molecular insight into the mechanisms that allow for strand swapping and highlight the general

role of these structures in regulating adhesive interactions.

Results

PLLLF proteins maintain low sequence identity but high sequence similarity and

structural similarity.

With the goal of identifying the underlying and shared features between GerS and the

rest of the PLLLF superfamily, we analyzed the evolutionary relatedness of proteins within the

group. We found that heuristic search methods, such as BLASTp, were inadequate for

identifying members of this group. Instead, we employed probabilistic searches using InterPro

and structural searches using FoldSeek to identify which major bacterial classes contain

bacteria that encode PLLLF proteins [43–46]. We then generated a phylogenetic tree using core

genes across species based on the major bacterial classes that have species encoding PLLLF

proteins (Supplemental Figure 1). Remarkably, we found that PLLLF proteins are widespread

across the bacterial domain. Further, we found that the presence of PLLLF proteins is

independent of whether the bacteria are monoderms or diderms (Supplemental Figure 1),

suggesting that not all PLLLF proteins function in a similar way to LolA.

Using AlphaFold 3 [47], we predicted the structure of 17 representative PLLLF proteins

and aligned them using DALI [48]. Using this structural alignment, we compared sequence

identity and sequence similarity across the proteins. Despite sharing little sequence identity, the

sequence similarity across PLLLF group members is high (Figure 1A). Together, these

observations indicate that PLLLF proteins are highly conserved and suggest that they share

evolutionary ancestry, although they likely evolved specialized roles within diderms vs.

monoderms.

GerS strand swapping is unique among characterized PLLLF proteins.

While AlphaFold 3 [47] predicts that GerS adopts a structure remarkably similar to other

PLLLF proteins, the GerS-CwlD crystal structure shows that GerS forms a homodimer via a

strand-swap mechanism (Figure 1B)[23]. Crystal structures of other PLLLF proteins, ranging

across several bacterial classes, suggest that PLLLF proteins exist either as monomers or

heterodimers with other proteins but rarely as homodimers [3,5,10,11,49,50].

While LolA does not form homodimers naturally, a single substitution of phenylalanine to

glutamate at residue 68 (F68E, previously published as F47E based on numbering from the

lipobox) is sufficient to induce homodimerization [42]. This homodimerization is achieved by

converting a β-turn into a strand-swap hinge, which is remarkably similar to the strand swap

observed in GerS (Figure 1C, Supplemental Figure 2). Notably, the LolA[F68E] variant binds its

interacting partner LolC with increased affinity, but this enhanced affinity interferes with LolA's

ability to bind the lipoprotein substrates that it normally transports across the periplasm,

rendering the LolA[F68E] variant toxic to *E. coli* [41,42]. Based on these observations, we

hypothesized that strand swap-mediated dimerization of GerS stabilizes GerS binding to CwlD

and promotes CwlD binding to its Zn^{2+} co-factor.

132   Dimerization is conserved in GerS orthologs of the Peptostreptococcaceae family.

Since CwlD is widely conserved in endospore-forming bacteria, we first assessed the

conservation of *gerS* among the Firmicutes, the only known class to contain endospore-forming

bacteria like *C. difficile* and *Bacillus subtilis*. To avoid issues associated with identifying PLLLF

proteins through heuristic methods like BLASTp, we instead used synteny-based searches such

as GeCoViz [51] and BioCyc genome alignment [52] to find GerS orthologs; we confirmed their

structural homology using AlphaFold3-predicted structures and ChimeraX (Figure 2A;

Supplemental Figure 3) [47,53].

Six orthologs were selected for further characterization from Firmicutes species that

sporulate and encode a GerS ortholog. To investigate dimerization, each ortholog was

expressed in *E. coli* and purified for analysis via size-exclusion chromatography (SEC). The

SEC trace of GerS from *Clostridioides difficile* revealed two distinct peaks, which represent the

monomeric and homodimeric forms of the protein ([23], Figure 2B; Supplemental Figure 4).

Notably, two other Firmicutes, *Paraclostridium bifermentans* and *Paraclostridium sordellii*

(formerly *Paeniclostridium sordellii* [54]), exhibited similar SEC profiles, indicating that they also

form homodimers. In contrast, GerS orthologs from *Clostridium aceticum*, *Gottschalkia acidurici*,

and *Clostridium perfringens* displayed a single peak corresponding to the monomeric form.

Since *P. bifermentans* and *P. sordellii* are members of the Peptostreptococcaceae family like *C.*

difficile, unlike the other species tested, these data suggest that GerS dimerization is specific to

the Peptostreptococcaceae family.

We also identified GerS orthologs in the Bacilli class, but they are distinguished by an

additional domain of unknown function (DUF4367). We purified the *Bacillus subtilis* ortholog,

previously published as SsdC [13], with and without the DUF4367 domain and analyzed the two

variants via SEC. Although both traces had two peaks, Coomassie staining of earlier eluting

fractions revealed proteins larger than the expected size for SsdC that are likely chaperones

(Supplemental Figure 4B-C), indicating that SsdC primarily exists as a monomer *in vitro* and

that the DUF4367 domain does not prevent SsdC dimerization. Thus, GerS dimerization is

limited to the Peptostreptococcaceae family.

Mutations in the strand-swap hinge alter GerS dimer stability.

To determine the role of the strand swap in regulating GerS dimerization and function,

we aimed to identify mutations that destabilize the GerS dimer. Previous studies have shown

that mutations that disrupt the 'hinge' region of the swapped strand can prevent dimerization

(Figure 3A) [28,29,55–57]. While these studies have also shown that it can be difficult to predict

which residue will alter a given dimer:monomer equilibrium, we reasoned that replacing the

hinge of GerS with the equivalent sequence found in a monomeric PLLLF protein like LolA

might prevent GerS dimerization. To this end, we used AlphaFold3 to predict the monomeric

structure of GerS and aligned it with the LolA crystal structure (PDB 1IWL). This alignment

revealed that the GerS strand-swap hinge, Lys71-Thr72-Asp73-Asn74 (KTDN), lies in a similar

position to LolA Arg64-Pro65-Asn66-Leu67 (RPNL; Figure 3B). We then tested the effect of

replacing individual amino acids in the *C. difficile* GerS hinge with the analogous residues in the

E. coli LolA region. Intriguingly, replacing Thr72 with Pro increased the proportion of GerS in the

monomeric form, while replacing Lys71 with Arg and Asn74 with Leu increased the dimeric form

(Figure 3C; Supplemental Figure 5A). In contrast, the GerS^{D73N} variant precipitated quickly after

purification and co-purified with the chaperones from *E. coli* (Supplemental Figure 5B)[58–60],

strongly suggesting that the substitution destabilized GerS.

Since the GerS^{K71R} and GerS^{N74L} variants favored the dimeric form relative to WT GerS

(Figure 3C-D), we purified the dimeric fractions and re-analyzed them by SEC to compare the

relative stabilities of the GerS homodimers formed by each variant. The re-purified GerS^{K71R} or

GerS^{N74L} variants again showed a higher dimer-to-monomer ratio than WT GerS (Supplemental

Figure 6A-B), indicating that the K71R and N74L substitutions enhance the stability of the GerS

homodimer structure.

185   Proline substitutions in the strand-swap hinge destabilize the GerS homodimer.

Since GerS^{T72P} favors the monomeric form compared to WT GerS (Figure 3C-D), we

next tested the effect of changing the other hinge residues to proline. Previous studies have

found that introducing prolines into the strand-swap hinge affects dimerization in a wide range of

proteins across all domains of life [28,29]. For example, highly conserved prolines in the loops

of *Mycobacterium tuberculosis* BlaC maintain BlaC in a monomeric and functional form,

whereas substitutions of these residues result in dimerization and loss of function [61]. When

the other hinge residues of GerS were replaced with proline (K71P, D73P, and N74P), all three

substitutions appeared to destabilize GerS (Figure 4A), and the resulting variants were quick to

crash out of solution. The D73P and N74P substitutions resulted in low protein yields, while

SEC analyses of the GerS^{K71P} variant revealed that it copurified with *E. coli* proteins likely to be

chaperones (Supplemental Figure 7B). Thus, the location of the proline substitution is critical for

regulating the GerS dimer-to-monomer equilibrium.

Consistent with this finding, analysis of the conservation of hinge prolines among PLLLF

proteins using the DALI alignment revealed that three proline residues are highly conserved

among this superfamily, and each of these prolines correspond to beta-turns in PLLLF proteins

(Figure 4B). Notably, GerS contains two of these conserved prolines but not the conserved

proline at residue 72 (Figure 4C). Together, these observations highlight how a proline at this

location destabilizes the GerS homodimer and promotes GerS monomer formation.

To test whether the conserved proline in the analogous position of other monomeric

PLLLF proteins regulates dimerization, we substituted the *E. coli* LolA Pro65 with a Thr (P65T).

Remarkably, while wild-type LolA eluted as a monomer, LolA[P65T] eluted primarily as a dimer

(Figure 4D). Taken together, these data highlight how a proline at this site promotes β-turn

formation and thus the monomeric form, consistent with past studies of β-turns [62,63].

While the T72P substitution effectively shifted the dimer-to-monomer equilibrium toward

the monomer state, it did not completely prevent the formation of GerS homodimers (Fig 3C,

4A). To further destabilize the dimer, we introduced a second proline at the fourth residue,

creating the hinge sequence Lys71-Pro72-Asp73-Pro74 (KPDP). Since prolines in β-turns are

often paired with flexible glycine residues to counteract their rigidity [63–65], we also replaced

the first and third residues in the hinge with glycines, yielding a Gly71-Pro72-Gly73-Pro74

(GPGP) hinge sequence. The GerS[KPDP] variant eluted as a single peak at a volume that falls

between the dimer and monomer peaks, suggesting that the variant interconverts between the

monomer and dimer forms as it is resolved on the column, whereas the GerS[GPGP] variant

exclusively eluted as a monomer (Figure 4E). These results indicate that the addition of a

second proline in the strand-swap hinge further destabilizes the GerS homodimer—favoring the

monomeric form—especially when glycines are introduced to provide greater conformation

flexibility.

Decreasing GerS dimer formation reduces its affinity for CwlD and impairs *C. difficile* spore germination

Since we originally hypothesized that strand-swapping in GerS enhances its ability to bind CwlD based on analyses of other strand-swapped proteins [42], we next tested if destabilizing the homodimer affects its affinity for CwlD. To this end, we co-affinity purified His$_6$-tagged CwlD with untagged GerS variants and analyzed their interactions by SEC. Untagged GerS^{WT}, GerS^{K71R}, and GerS^{N74L} co-purified efficiently with His$_6$-tagged CwlD and eluted as the GerS:CwlD complex as well as the GerS and CwlD monomers in SEC analyses (Figures 5A,B, Supplementary Figure 8). It should be noted that the GerS:CwlD complex eluted as several peaks due to the higher resolution SEC performed relative to our prior work [23].

Since the K71R and N74L substitutions stabilized GerS homodimerization, we considered the possibility that these variants may bind CwlD more tightly than wild-type GerS. To test this, we resolved co-purifications of GerS^{K71R}:CwlD-His$_6$ and GerS^{N74L}:CwlD-His$_6$ using SEC, collected the dimer fractions, and re-resolved them through SEC again (Supplemental Figure 9). To our surprise, the GerS^{K71R} and GerS^{N74L} variants formed weaker interactions with CwlD-His$_6$ than wild-type GerS. However, the differences between GerS^{K71R} and GerS^{WT} were subtle. We next attempted to further stabilize the GerS dimer by replacing the GerS hinge with the hinge (QVVAG) from the strand-swapping cystatin, Stefin B, because this hinge was previously reported to reliably form domain-swapped homodimers [66]. However, substituting GerS's hinge region for Stefin B's hinge region (GerS^{QVVAG}) destabilized GerS, causing it to quickly crash out of solution; indeed, the SEC trace matched that of the unstable D73N substitution (Supplemental Figure 10), and untagged GerS^{QVVAG} failed to copurify with CwlD-His$_6$ (Supplemental Figure 10B). Therefore, the effect of stabilizing the GerS homodimer on CwlD remains unclear.

In contrast to GerS^{K71R} and GerS^{N74L}, considerably less GerS^{T72P} co-purified with His$_6$-tagged CwlD (Figure 5A, Supplementary Figure 7). In addition, when the co-purification was

resolved by SEC, no GerS^{T72P}:CwlD-His$_6$ complex was observed, indicating that the co-affinity

purified complex was not stable during SEC (Supplementary Figure 8). These latter data

indicate that destabilizing the GerS homodimer decreases its binding to CwlD. Consistent with

this conclusion, the GerS^{KPDP} and GerS^{GPGP} variants, which favor the monomeric form (Figure

4E), had even lower affinity for CwlD-His$_6$ (Figure 5B).

To determine the impact of preventing GerS dimerization on GerS function during *C.*

difficile germination, we complemented our Δ*gerS* mutant with constructs encoding dimer-

breaking GerS mutations and measured their effects on GerS stability and spore germination

efficiency. While GerS^{K71R} and GerS^{N74L} interacted with CwlD at the same levels as wild-type

GerS *in vitro*, both variants were detected at lower levels in sporulating cells compared to the

Δ*gerS* wild-type complement. Additionally, both mutants were almost undetectable in purified

spores (Figure 5C). Consistent with our previous findings that interaction with CwlD stabilizes

GerS [23], very low levels of the GerS^{GPGP} variant were detected in sporulating cells, and we

were unable to detect this variant in purified spores (Figure 5C). However, the GerS^{T72P} variant

and GerS^{KPDP} variant were readily detectable in both sporulating cells and purified spores,

implying that their ability to interact with CwlD is sufficient to stabilize the GerS variants in

spores (Figure 5C).

Notably, even though the dimer-disrupting strand-swap mutations decreased GerS

levels in mature spores, the GerS dimerization mutants exhibited only an ~2-fold decrease in

germination ($p < 0.00005$, Figure 5D). This finding is consistent with our prior work indicating

that a GerS mutation, D106R, which disrupts a salt-bridge at the interaction interface of GerS

and CwlD, results in a modest but significant decrease in germination [23]. Our results also

suggest that relatively GerS is needed in spores to license CwlD amidase activity.

Interaction with CwlD requires both strand swapping and a salt bridge.

Since our analyses revealed that GerS mutants defective in dimerization still allow for *C.*

difficile germination to proceed, additional features of GerS beyond strand-swapping allow GerS

to bind CwlD. Indeed, our *in vitro* analyses revealed that GerS dimerization mutants can still co-

purify with CwlD-His$_6$, albeit at greatly reduced levels (Figure 5B). Since we previously showed

that the interaction between GerS and CwlD is partially dependent on a salt bridge that forms

between arginine 169 of CwlD and aspartate 106 of GerS [23], we reasoned that GerS binding

to CwlD may depend on both the salt bridge and GerS strand swap (Figure 6A). To test this

possibility, we combined the salt-bridge and strand-swapping mutations to generate

GerS^{T72P,D106R}, GerS^{KPDP,D106R}, and GerS^{GPGP,D106R} variants. Remarkably, these combined

mutations fully disrupted GerS binding to CwlD, since the untagged GerS variants were not

detectable in the co-elutions with His$_6$-tagged CwlD (Figure 6B). Consistent with this finding,

purified spores from Δ*gerS* strains complemented with constructs encoding the combined salt-

bridge and strand-swap mutations failed to germinate (Figure 6C), presumably due to their

inability to bind CwlD and allosterically activate its amidase activity during cortex modification

[23]. In addition, the combined mutations greatly decreased GerS levels even in sporulating

cells (Figure 6D). Taken together, our data highlight the importance of both the strand swap and

salt bridge on GerS binding to CwlD and, by extension, regulation of *C. difficile* spore

germination*.*

Discussion

Our prior discovery that the *C. difficile* GerS lipoprotein binds the CwlD amidase as a

strand-swapped dimer led to the question of whether strand-swapping by GerS enhances its

ability to allosterically activate the CwlD amidase [23]. By identifying mutations that shift the

GerS dimer:monomer equilibrium to the monomeric form (Figure 3), we show that monomeric

GerS has greatly reduced affinity for its binding partner, CwlD, which in turn reduces, but does

not abolish, *C. difficile* spore germination (Figure 5). Since combining monomer-promoting

mutations with a mutation that disrupts a salt-bridge at the GerS:CwlD interface abrogates

GerS's ability to activate the CwlD amidase, our data reveal a critical role for GerS's hinge

region in regulating the allosteric activation of CwlD.

While our findings are consistent with prior work highlighting that strand-swapping can

stabilize adhesive interactions, such as in cadherins and the LolA[F68E] variant, they also raise the

possibility that the dynamic instability of the strand-swapped dimer, which exists in near-equal

proportions with the monomeric form *in vitro* (Figure 3), contributes to GerS function. Strand

swapping is sometimes the result of localized unfolding around the hinge. While the rest of the

protein folds, the strand to be swapped unfolds, leading to monomeric and dimeric

conformations [32,67]. Like GerS, the yeast cell-cycle regulation protein p13[suc1] has both a

native monomer and dimer state [29]. Two prolines in the hinge generate tension in each state,

functioning as a sort of spring between both the monomer and dimer conformation.

Substitutions in the hinge of the swapped structure shift the equilibrium of the dimer and

monomer populations [29]; in a similar manner, the K71R and N74L substitutions in GerS

stabilize the dimeric conformation, while the T72P and GPGP substitutions stabilize the

monomeric conformation. The persistence of the two stable populations in near equal

proportions suggests that the GerS monomer may be functionally important, perhaps to ensure

that GerS transiently interacts with CwlD during MAL formation. Alternatively, additional

regulatory factors may influence or stabilize one conformation over the other during this

process.

Our finding that replacing the GerS strand-swap hinge with the corresponding region

from Stefin B [66] destabilizes the protein (Supplemental Figure 10) suggests that dimerization

in these proteins depends on distinct, non-interchangeable hinge features. While most domain-

swapped proteins have hinge loops of four to five residues, they otherwise vary widely in

sequence and structure. Nonetheless, a common feature is that these hinges originate from β-

turns in their monomeric structures [38,68,69]. Consistent with previous findings that conserved

prolines in β-turns can prevent deleterious strand-swapping in *M. tuberculosis* BlaC [61], we

found that the introduction of prolines into the GerS hinge destabilizes the strand-swapped

dimer (T72P, GPGP; Figure 4)[61]. Yet, there are also examples where introducing prolines into

β-turns instead promotes domain swapping [70–72]. Together, these findings underscore that

there is no universal "syntax" for strand swapping and that structural states must be empirically

determined on a case-by-case basis.

The fact that a single threonine-to-proline substitution in the GerS hinge dramatically

alters its conformational stability implies that PLLLF proteins may have evolved inherent

instability at this site, potentially enabling their structural flexibility. Although *C. difficile* and *E.*

coli diverged billions of years ago [73], a reciprocal substitution at the same hinge position in

LolA (LolA^{P65T}) is sufficient to shift the protein from a monomer to a strand-swapped dimer

(Figure 4). This evidence, along with the similarly strand-swapped LolA^{F68E} (published F47E)

[41,42] structure, supports the idea that the LolA monomeric structure around these sites is

intrinsically unstable. These findings also highlight the ease with which PLLLF proteins can

evolve strand-swapped dimerization. This finding, along with the grouping of both proteins into

the same superfamily by SCOP [74], also raises the intriguing possibility of a shared ancestry

between LolA and GerS. During the evolution of monoderm bacteria, which involved the loss of

the ancestral outer membrane [75–77], there would no longer be a need to transport

lipoproteins across the periplasmic space, facilitating a specialization of the ancestral proteins

during sporulation in endospore-forming monoderms [13,23]. This hypothesis may also clarify

why PLLLF proteins are absent in certain non-spore-forming monoderms such as

Staphylococcus aureus and the seemingly convergent roles of LppX [10] and Lpr proteins

[11,12] in transporting glycolipids across the mycobacterial periplasm.

Much work remains to be done to understand precisely why GerS strand swapping

stabilizes its interaction with CwlD. Notably, CwlD binding to its Zn^{2+} co-factor also stabilizes the

GerS:CwlD interaction, so it is possible that Zn^{2+}-bound CwlD adopts a conformation that

similarly promotes GerS strand-swapping. Identifying the allosteric circuit that allows GerS

binding to CwlD to be communicated to CwlD's active site and vice versa will likely provide

insight into the role of GerS strand-swapping in the allosteric activation of CwlD. Another

question that arises from our identification of GerS orthologs in other clostridial species that do

not form dimers is whether these GerS orthologs have additional or different conserved

functions beyond regulating CwlD activity during cortex modification? The requirement of CwlD

for GerS processing and incorporation into the spore may indicate that GerS functions in a very

specialized way in the Peptostreptococcaceae family [22]. In contrast, it is possible that GerS

orthologs in other Clostridia function similarly to SsdC in *B. subtilis*, which is required for proper

spore morphology through its impacts on spore cortex assembly [13]. Nevertheless, how these

orthologs are incorporated into the spore if they do not interact with CwlD as well as whether

GerS and CwlD can independently localize to the forespore remain open questions and will

require analyses in organisms other than *C. difficile*.

Materials and Methods

Structural alignment, sequence analysis, and phylogenetics. 17 PLLLF proteins were

selected from the InterPro [44] SuperFamily SSF89392, representing the taxonomic diversity of

the PLLLF protein family. AlphaFold3 [47] was used to predict the structure of each protein. The

17 proteins were aligned using DALI [48], which generates an amino-acid sequence alignment

based on the closest amino acids in each aligned structure. The structural alignment was imported

into Geneious Prime 2024.0.5. to calculate the %identity and %similarity between each sequence

(see Table S1). For %similarity calculations, the Blosum45 matrix was used with a threshold of 0.

WebLogo [78] was used to generate the sequence logo of the structural alignment.

The phylogenetic tree of the major bacterial classes, with Archaea and Eukarya as used

as outgroups, was constructed using genomes publicly available on KBase [79]. Five genomes

for each class were selected, although there were several instances when five were not available

for a given class. The complete list of these genomes can be found in Table S2. The KBase app

"Insert Genome into SpeciesTree - v2.2.0" was used to identify, concatenate, and align a set of

49 core, universal genes defined by COG (Clusters of Orthologous Groups) gene families across

all selected genomes. The app generates the phylogenetic tree using FastTree2 [80]. The tree

was visualized and prepared into a figure using iTOL [81]. Identification of PLLLF proteins in each

class was done by searching for PLLLF proteins encoded by members of each class in the

InterPro [44] SuperFamily SSF89392 and by running FoldSeek [43] using the LolA crystal

structure (1IWL).

Bacterial strains and growth conditions. *C. difficile* strains 630Δ*erm*Δ*cwlD*Δ*pyrE* and

630Δ*erm*Δ*gerS*Δ*pyrE* were used as the parental strains for *pyrE*-based allelic-coupled exchange

(ACE) [82], which allows for single-copy complementation from the ectopic *pyrE* locus. The *C.*

difficile strains used are listed in Table S3. They were grown on brain heart infusion media (BHIS)

supplemented with taurocholate (TA, 0.1% wt/vol; 1.9 mM), thiamphenicol (10-15 µg /mL),

kanamycin (50 µg/ml), cefoxitin (8 µg/ml) and L-cysteine (0.1% w/v; 8.25 mM) as needed. For

ACE, *C. difficile* was grown on *C. difficile* defined medium (CDDM) [83]. Cultures were grown at

37°C under anaerobic conditions using a gas mixture containing 85% N_2, 5% CO_2 and 10% H_2.

Escherichia coli strains used for BL21(DE3)-based protein production and HB101/pRK24-

based conjugations are listed in Table S3. *E. coli* strains were grown at 37°C with shaking at 225

rpm in Luria-Bertani broth (LB). The media was supplemented with chloramphenicol (20 µg/ml),

ampicillin (50 µg/ml), or kanamycin (30 µg/ml) as indicated.

***E. coli* strain construction.** Plasmid constructs were confirmed by sequencing using Genewiz

or Plasmidsaurus (see the Benchling links in Table S3) and transformed into either the

HB101/pRK24 conjugation strain (used with *C. difficile*) or BL21(DE3) expression strain.

***C. difficile* strain construction and complementation**. Allele-couple exchange (ACE) was used

to construct all complementation strains as previously described by conjugating HB101/ pRK24

carrying pMTL-YN1C plasmids into Δ*pyrE*-based strains.

Protein purification. Starter cultures were grown in 20 mL LB broth with 30 µg/mL of

kanamycin and, in some cases, 100 µg/mL ampicillin. Terrific broth (TB) with ampicillin and

kanamycin was inoculated with the starter culture (1:1000) and incubated for ~60 hr at 20°C

with 225 rpm shaking. The purifications were carried out in two steps: nickel bead-based (co-

)affinity purification followed by size exclusion chromatography (SEC). Affinity purification was

carried out as stated above: the cells were pelleted, resuspended in lysis buffer (500 mM NaCl,

50 mM Tris [pH 7.5], 15 mM imidazole, 10% [vol/vol] glycerol, 2 mM ß-mercaptoethanol), flash

frozen in liquid nitrogen, thawed and then sonicated. The insoluble material was pelleted, and

the soluble fraction was incubated with Ni-NTA agarose beads (Qiagen) for 2 hours to capture

the His$_6$-tagged proteins. Washes were carried out five times with low imidazole buffer (500mM

NaCl, 10mM Tris-HCl pH 7.5, 10% (v/v) glycerol, 15 mM imidazole, 2 mM ß -mercaptoethanol)

to decrease non-specific binding to the beads. Proteins were eluted using high-imidazole buffer

(500 mM NaCl, 50 mM Tris [pH 7.5], 200 mM imidazole, 10% [vol/vol] glycerol, 2 mM ß-

mercaptoethanol) after nutating the sample for 5 to 10 min. Pooled protein elutions were

concentrated to 20 mg/ml or less in a size exclusion chromatography buffer consisting of 200

mM NaCl, 10 mM Tris HCl pH 7.5, 5% glycerol, and 1 mM DTT.

Size exclusion chromatography (SEC). SEC was carried out using a Superdex 200 Increase

10/300 GL (GE Healthcare) column. A high-resolution protocol was used to resolve affinity-

purified protein elutions by SEC. Specifically, purified proteins were diluted to a final

concentration of 10 mg/mL in the SEC buffer listed above, and then 100 µL of the 10 mg/mL

diluted protein was resolved using a 0.2 mL/min flow rate. The resulting fractions were resolved

using SDS-PAGE, and the gels were Coomassie stained using GelCode Blue according to the

manufacturer's instructions (ThermoFisher Scientific). Protein sequencing of non-GerS bands

was performed by automated Edman degradation on an ABI Procise 494 HT Protein/Peptide

Sequencer at the Tufts University Core Facility (TUCF).

Western blot analysis. Samples for immunoblotting were prepared by resuspending sporulating

cell pellets in 100 µl of PBS, and freeze-thawing a 50 µL aliquot of this resuspension for three

cycles. The pellet was then resuspended in 100 µL EBB buffer (8 M urea, 2 M thiourea, 4% [wt/vol]

SDS, 2% [vol/vol] ß-mercaptoethanol). *C. difficile* spores (~1 x 10^6) were resuspended in EBB

buffer, which extracts proteins in all layers of the spore, including the core. Both sporulating cells

and spores were incubated at 95°C for 20 min with vortex mixing. Samples were centrifuged for

5 min at 15,000 rpm, and 4 x sample buffer (40% [vol/vol] glycerol, 1 M Tris [pH 6.8], 20% [vol/vol]

ß-mercaptoethanol, 8% [wt/vol] SDS, 0.04% [wt/vol] bromophenol blue) was added. Samples

were incubated again at 95°C for 5 to 10 min with vortex mixing followed by centrifugation for 5

min at 15,000 rpm. The samples were resolved by the use of 14% SDS-PAGE gels and

transferred to a Millipore Immobilon-FL polyvinylidene difluoride (PVDF) membrane. The

membranes were blocked in Odyssey blocking buffer with 0.1% (vol/vol) Tween 20. Polyclonal

rabbit anti-GerS and anti-CwlD antibodies were used at a 1:500 and 1:750 dilution, respectively.

Polyclonal mouse anti-SleC antibodies were used at a 1:5000 dilution. IRDye 680CW and 800CW

infrared dye-conjugated secondary antibodies were used at 1:30,000 dilutions. An Odyssey LiCor

CLx was used to visualize secondary antibody infrared fluorescence emissions.

Spore purification. Sporulation was induced on 70:30 agar plates for 3-5 days, and the cultures

were harvested for spore purifications. Spores were washed 6 times in ice-cold water, incubated

overnight in water on ice, treated with DNase I (New England Biolabs) at 37°C for 60 min, and

purified on a HistoDenz (Sigma- Aldrich) gradient. Phase-contrast microscopy was used to assess

spore purity (>95% pure); the optical density of the spore stock was measured at OD_{600}; and

spores were stored in water at 4°C. Four 70:30 plates were used to induce sporulation for each

strain; purified spores were resuspended in 200 µL, and the total optical density of this mixture

was determined for each strain. At least three independent spore preparations were performed.

Spore purity as assessed by phase-contrast microscopy. Only spore suspensions that were >95%

pure were used.

Optical density-based germination assay. For OD_{600}-based germination assays, germination

was induced with taurocholate in 96-well plates. Specifically, ~1.6 x 10^7 spores (0.55 OD_{600} units)

for each condition tested were resuspended in BHIS and then 180 µL of this suspension was

aliquoted in duplicate into a 96 well flat-bottom tissue culture plate (Falcon) for each condition

tested. The spores were exposed to 1% taurocholate (19 mM), or water (untreated) in a final

volume of 200 µL. The OD_{600} of the sample was measured every 3 minutes in a Synergy H1

microplate reader (Biotek) at 37°C with constant shaking between readings. The change in OD_{600}

over time was calculated as the ratio of the OD_{600} at each time point to the OD_{600} at time zero.

Relative germination was calculated by measuring the area between the drop in OD_{600} and 1.0

after 39 minutes (13 reads). These assays were performed on 3-6 independent spore

preparations depending on the condition. Data shown are averages from these replicates.

Comparison of each mean to WT was performed using a one-way ANOVA followed by Dunnett's

multiple comparisons test, with a single-pooled variance.

References

[1] A. De Las Peñas, L. Connolly, C.A. Gross, The σE-mediated response to extracytoplasmic stress in *Escherichia coli* is transduced by RseA and RseB, two negative regulators of σE, Mol Microbiol 24 (1997) 373–385. https://doi.org/10.1046/J.1365-2958.1997.3611718.X.

[2] M.J. Schurr, H. Yu, J.M. Martinez-Salazar, J.C. Boucher, V. Deretic, Control of AlgU, a member of the sigma E-like family of stress sigma factors, by the negative regulators MucA and MucB and *Pseudomonas aeruginosa* conversion to mucoidy in cystic fibrosis, J Bacteriol 178 (1996) 4997–5004. https://doi.org/10.1128/JB.178.16.4997-5004.1996.

[3] D. Jaiman, R. Nagampalli, K. Persson, A comparative analysis of lipoprotein transport proteins: LolA and LolB from *Vibrio cholerae* and LolA from *Porphyromonas gingivalis*, Scientific Reports 2023 13:1 13 (2023) 1–10. https://doi.org/10.1038/s41598-023-33705-y.

[4] S. Okuda, H. Tokuda, Model of mouth-to-mouth transfer of bacterial lipoproteins through inner membrane LolC, periplasmic LolA, and outer membrane LolB, Proc Natl Acad Sci U S A 106 (2009) 5877–5882. https://doi.org/10.1073/PNAS.0900896106/SUPPL_FILE/0900896106SI.PDF.

[5] V. Furlanetto, C. Divne, LolA and LolB from the plant-pathogen *Xanthomonas campestris* forms a stable heterodimeric complex in the absence of lipoprotein, Front Microbiol 14 (2023) 1216799. https://doi.org/10.3389/FMICB.2023.1216799/BIBTEX.

[6] K.L. May, M. Grabowicz, Outer membrane lipoproteins: late to the party, but the center of attention, J Bacteriol (2024). https://doi.org/10.1128/JB.00442-24.

[7] E.D. LoVullo, L.F. Wright, V. Isabella, J.F. Huntley, M.S. Pavelka, Revisiting the gram-negative lipoprotein paradigm, J Bacteriol 197 (2015) 1705–1715. https://doi.org/10.1128/JB.02414-14/SUPPL_FILE/ZJB999093599SO4.XLSX.

[8] S. ichiro Narita, H. Tokuda, Sorting of bacterial lipoproteins to the outer membrane by the Lol system, Methods Mol Biol 619 (2010) 117–129. https://doi.org/10.1007/978-1-60327-412-8_7.

[9] S.Y. Tanaka, S.I. Narita, H. Tokuda, Characterization of the *Pseudomonas aeruginosa* Lol system as a lipoprotein sorting mechanism, Journal of Biological Chemistry 282 (2007) 13379–13384. https://doi.org/10.1074/jbc.M611840200.

[10] G. Sulzenbacher, S. Canaan, Y. Bordat, O. Neyrolles, G. Stadthagen, V. Roig-Zamboni, J. Rauzier, D. Maurin, F. Laval, M. Daffé, C. Cambillau, B. Gicquel, Y. Bourne, M. Jackson, LppX is a lipoprotein required for the translocation of phthiocerol dimycocerosates to the surface of *Mycobacterium tuberculosis*, EMBO Journal 25 (2006) 1436–1444. https://doi.org/10.1038/SJ.EMBOJ.7601048/SUPPL_FILE/EMBJ7601048-SUP-0001.PDF.

[11] M.G. Drage, H.C. Tsai, N.D. Pecora, T.Y. Cheng, A.R. Arida, S. Shukla, R.E. Rojas, C. Seshadri, D.B. Moody, W.H. Boom, J.C. Sacchettini, C. V. Harding, *Mycobacterium tuberculosis* lipoprotein LprG (Rv1411c) binds triacylated glycolipid agonists of Toll-like receptor 2, Nature Structural & Molecular Biology 2010 17:9 17 (2010) 1088–1095. https://doi.org/10.1038/nsmb.1869.

[12] J.S. Kim, L. Jiao, J. Il Oh, N.C. Ha, Y.H. Kim, Crystal structure and functional implications
of LprF from *Mycobacterium tuberculosis* and *M. bovis*, Acta Crystallogr D Biol Crystallogr
70 (2014) 2619–2630. https://doi.org/10.1107/S1399004714016599.

[13] J. Luhur, H. Chan, B. Kachappilly, A. Mohamed, C. Morlot, M. Awad, D. Lyras, N. Taib, S.
Gribaldo, D.Z. Rudner, C.D.A. Rodrigues, A dynamic, ring-forming MucB / RseB-like
protein influences spore shape in *Bacillus subtilis*, PLoS Genet 16 (2020) e1009246.
https://doi.org/10.1371/JOURNAL.PGEN.1009246.

[14] K.A. Fimlaid, O. Jensen, M.L. Donnelly, M.B. Francis, J.A. Sorg, A. Shen, Identification of a
Novel Lipoprotein Regulator of *Clostridium difficile* Spore Germination, PLoS Pathog 11
(2015) e1005239. https://doi.org/10.1371/JOURNAL.PPAT.1005239.

[15] D.A. Burns, J.T. Heap, N.P. Minton, SleC is essential for germination of *Clostridium difficile*
spores in nutrient-rich medium supplemented with the bile salt taurocholate, J Bacteriol 192
(2010) 657–664. https://doi.org/10.1128/JB.01209-09/ASSET/F0B6A88D-0D10-4FE5-
ADDE-C2E652FF1E28/ASSETS/GRAPHIC/ZJB9990993070005.JPEG.

[16] D. Paredes-Sabja, P. Setlow, M.R. Sarker, SleC Is Essential for Cortex Peptidoglycan
Hydrolysis during Germination of Spores of the Pathogenic Bacterium *Clostridium*
perfringens, J Bacteriol 191 (2009) 2711–2720. https://doi.org/10.1128/JB.01832-08.

[17] D. Gutelius, K. Hokeness, S.M. Logan, C.W. Reid, Functional analysis of SleC from
Clostridium difficile: an essential lytic transglycosylase involved in spore germination,
Microbiology (N Y) 160 (2014) 209. https://doi.org/10.1099/MIC.0.072454-0.

[18] T. Fukushima, H. Yamamoto, A. Atrih, S.J. Foster, J. Sekiguchi, A polysaccharide
deacetylase gene (pdaA) is required for germination and for production of muramic δ-
lactam residues in the spore cortex of *Bacillus subtilis*, J Bacteriol 184 (2002) 6007–6015.
https://doi.org/10.1128/JB.184.21.6007-6015.2002/ASSET/281D2E2F-2AB6-4943-9460-
50EB1BA91E0B/ASSETS/GRAPHIC/JB2120422006.JPEG.

[19] H. Coullon, A. Rifflet, R. Wheeler, C. Janoir, I.G. Boneca, T. Candela, N-Deacetylases
required for muramic-lactam production are involved in *Clostridium difficile* sporulation,
germination, and heat resistance, Journal of Biological Chemistry 293 (2018) 18040–
18054. https://doi.org/10.1074/JBC.RA118.004273/ATTACHMENT/A0930DBF-1D55-
4CDE-96DE-AEFCB897DBF5/MMC1.ZIP.

[20] M.J. Tobin, S.Y. Cho, W. Profy, T.M. Ryan, D.H. Le, C. Lin, E.Z. Yip, J.L. Dorsey, B.R.
Levy, J.D. Rhodes, M.A. Welsh, Reconstituting Spore Cortex Peptidoglycan Biosynthesis
Reveals a Deacetylase That Catalyzes Transamidation, (2023).
https://doi.org/10.1021/acs.biochem.3c00100.

[21] C. Kim, M. Lee, B.T. Birhanu, D. Hesek, M. Chang, S. Mobashery, Synthesis of Muramyl-δ-
Lactam in Spore Peptidoglycan of *Clostridioides difficile*, ChemBioChem 24 (2023)
e202300282. https://doi.org/10.1002/CBIC.202300282.

[22] O.R. Diaz, C. V. Sayer, D.L. Popham, A. Shen, *Clostridium difficile* Lipoprotein GerS Is
Required for Cortex Modification and Thus Spore Germination, MSphere 3 (2018).
https://doi.org/10.1128/MSPHERE.00205-18/SUPPL_FILE/SPH003182575SF7.TIF.

[23] C.A. Feliciano, B.E. Eckenroth, O.R. Diaz, S. Doublié, A. Shen, A lipoprotein allosterically
activates the CwlD amidase during *Clostridioides difficile* spore formation, PLoS Genet 17
(2021) e1009791. https://doi.org/10.1371/JOURNAL.PGEN.1009791.

[24] P.R. Gouldson, C.R. Snell, R.P. Bywater, C. Higgs, C.A. Reynolds, Domain swapping in G-
protein coupled receptor dimers., Protein Engineering, Design and Selection 11 (1998)
1181–1193. https://doi.org/10.1093/PROTEIN/11.12.1181.

[25] L. Vitagliano, S. Adinolfi, F. Sica, A. Merlino, A. Zagari, L. Mazzarella, A potential allosteric
subsite generated by domain swapping in bovine seminal ribonuclease, J Mol Biol 293
(1999) 569–577. https://doi.org/10.1006/JMBI.1999.3158.

[26] R. Janowski, M. Kozak, E. Jankowska, Z. Grzonka, A. Grubb, M. Abrahamson, M.
Jaskolski, Human cystatin C, an amyloidogenic protein, dimerizes through three-
dimensional domain swapping, Nature Structural Biology 2001 8:4 8 (2001) 316–320.
https://doi.org/10.1038/86188.

[27] A.D. Cameron, B. Olin, M. Ridderström, B. Mannervik, T.A. Jones, Crystal structure of
human glyoxalase I—evidence for gene duplication and 3D domain swapping, EMBO J 16
(1997) 3386–3395. https://doi.org/10.1093/EMBOJ/16.12.3386.

[28] J. Vendome, S. Posy, X. Jin, F. Bahna, G. Ahlsen, L. Shapiro, B. Honig, Molecular design
principles underlying β-strand swapping in the adhesive dimerization of cadherins, Nature
Structural & Molecular Biology 2011 18:6 18 (2011) 693–700.
https://doi.org/10.1038/nsmb.2051.

[29] F. Rousseau, J.W.H. Schymkowitz, H.R. Wilkinson, L.S. Itzhaki, Three-dimensional domain
swapping in p13suc1 occurs in the unfolded state and is controlled by conserved proline
residues, Proceedings of the National Academy of Sciences 98 (2001) 5596–5601.
https://doi.org/10.1073/PNAS.101542098.

[30] K. Shivaji Paithankar, M. Enderle, D.C. Wirthensohn, A. Miller, M. Schlesner, F. Pfeiffer, A.
Rittner, M. Grininger, D. Oesterhelt, Structure of the archaeal chemotaxis protein CheY in a
domain-swapped dimeric conformation, (n.d.).
https://doi.org/10.1107/S2053230X19010896.

[31] M.J. Bennett, S. Choe, D. Eisenberg, Domain swapping: Entangling alliances between
proteins, Proc Natl Acad Sci U S A 91 (1994) 3127–3131.
https://doi.org/10.1073/PNAS.91.8.3127;WEBSITE:WEBSITE:PNAS-
SITE;WGROUP:STRING:PUBLICATION.

[32] N.M. Mascarenhas, S. Gosavi, Understanding protein domain-swapping using structure-
based models of protein folding, Prog Biophys Mol Biol 128 (2016) 113.
https://doi.org/10.1016/J.PBIOMOLBIO.2016.09.013.

[33] B. Brumshtein, S.R. Esswein, M. Landau, C.M. Ryan, J.P. Whitelegge, M.L. Phillips, D.
Cascio, M.R. Sawaya, D.S. Eisenberg, Formation of Amyloid Fibers by Monomeric Light
Chain Variable Domains, Journal of Biological Chemistry 289 (2014) 27513–27525.
https://doi.org/10.1074/JBC.M114.585638.

[34] B. Gerhartz, I. Ekiel, M. Abrahamson, Two stable unfolding intermediates of the disease-causing L68Q variant of human cystatin C, Biochemistry 37 (1998) 17309–17317. https://doi.org/10.1021/BI980873U/ASSET/IMAGES/LARGE/BI980873UF00007.JPEG.

[35] E. Zerovnik, V. Stoka, A. Mirtič, G. Gunčar, J. Grdadolnik, R.A. Staniforth, D. Turk, V. Turk, Mechanisms of amyloid fibril formation–focus on domain-swapping, Wiley Online Library 278 (2011) 2263–2282. https://doi.org/10.1111/J.1742-4658.2011.08149.X.

[36] M.J.-A.B. Polonica, undefined 2001, 3D domain swapping, protein oligomerization, and amyloid formation., Frontierspartnerships.Org 48 (2001) 807–827. https://doi.org/10.18388/ABP.2001_3849/PDF.

[37] P. Jurczak, P. Groves, A. Szymanska, S. Rodziewicz-Motowidlo, Human cystatin C monomer, dimer, oligomer, and amyloid structures are related to health and disease, FEBS Lett 590 (2016) 4192–4201. https://doi.org/10.1002/1873-3468.12463;WEBSITE:WEBSITE:FEBS;PAGE:STRING:ARTICLE/CHAPTER.

[38] J.H. Ha, J.M. Karchin, N. Walker-Kopp, C.A. Castañeda, S.N. Loh, Engineered domain swapping as an on/off switch for protein function, Chem Biol 22 (2015) 1384–1393. https://doi.org/10.1016/j.chembiol.2015.09.007.

[39] J.M. Modenbach, C. Möller, S. Asgarbeik, N. Geist, N. Rimkus, M. Dörr, H. Wolfgramm, L. Steil, A. Susemihl, L. Graf, O. Schmöker, D. Böttcher, E. Hammer, J. Glaubitz, M. Lammers, M. Delcea, U. Völker, A.A. Aghdassi, M.M. Lerch, F.U. Weiss, U.T. Bornscheuer, M. Sendler, Biochemical analyses of cystatin-C dimers and cathepsin-B reveals a trypsin-driven feedback mechanism in acute pancreatitis, Nature Communications 2025 16:1 16 (2025) 1–18. https://doi.org/10.1038/s41467-025-56875-x.

[40] S. Posy, L. Shapiro, B. Honig, Sequence and structural determinants of strand swapping in cadherin domains: Do all cadherins bind through the same adhesive interface?, J Mol Biol 378 (2008) 954. https://doi.org/10.1016/J.JMB.2008.02.063.

[41] A. Miyamoto, S. ichi Matsuyama, H. Tokuda, Dominant negative mutant of a lipoprotein-specific molecular chaperone, LolA, tightly associates with LolCDE, FEBS Lett 528 (2002) 193–196. https://doi.org/10.1016/S0014-5793(02)03305-7.

[42] E. Kaplan, N.P. Greene, A. Crow, V. Koronakis, Insights into bacterial lipoprotein trafficking from a structure of LolA bound to the LolC periplasmic domain, Proc Natl Acad Sci U S A 115 (2018) E7389–E7397. https://doi.org/10.1073/PNAS.1806822115/SUPPL_FILE/PNAS.1806822115.SM03.MOV.

[43] M. van Kempen, S.S. Kim, C. Tumescheit, M. Mirdita, J. Lee, C.L.M. Gilchrist, J. Söding, M. Steinegger, Fast and accurate protein structure search with Foldseek, Nature Biotechnology 2023 42:2 42 (2023) 243–246. https://doi.org/10.1038/s41587-023-01773-0.

[44] M. Blum, A. Andreeva, L. Cavalcanti Florentino, S. Rocio Chuguransky, T. Grego, E. Hobbs, B. Lazaro Pinto, A. Orr, T. Pa ysan-Laf osse, I. Ponamareva, G.A. Salazar, N. Bordin, P. eer Bor, A. Br idg, L. Colw ell, J. Gough, D.H. Haft, I. Letunic, F. Llinares-López, A. Marchler-Bauer, L. Meng-Papaxanthos, H. Mi, D.A. Natale, C.A. Orengo, A.P. Pandur ang an, D. Pio esan, C. ine Rivoire, C.J. istian Sigr ist, N. mada Thanki, F. ançoise Thibaud-Nissen, P.D. Thomas, S.C. Tosatto, C.H. W, A. Bat eman, InterPro: the protein

sequence classification resource in 2025, Nucleic Acids Res 1 (2013) 13–14.
https://doi.org/10.1093/NAR/GKAE1082.

[45] C. Camacho, G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos, K. Bealer, T.L. Madden,
BLAST+: Architecture and applications, BMC Bioinformatics 10 (2009) 1–9.
https://doi.org/10.1186/1471-2105-10-421/FIGURES/4.

[46] S.F. Altschul, W. Gish, W. Miller, E.W. Myers, D.J. Lipman, Basic local alignment search
tool, J Mol Biol 215 (1990) 403–410. https://doi.org/10.1016/S0022-2836(05)80360-2.

[47] J. Abramson, J. Adler, J. Dunger, R. Evans, T. Green, A. Pritzel, O. Ronneberger, L.
Willmore, A.J. Ballard, J. Bambrick, S.W. Bodenstein, D.A. Evans, C.C. Hung, M. O'Neill,
D. Reiman, K. Tunyasuvunakool, Z. Wu, A. Žemgulytė, E. Arvaniti, C. Beattie, O. Bertolli,
A. Bridgland, A. Cherepanov, M. Congreve, A.I. Cowen-Rivers, A. Cowie, M. Figurnov, F.B.
Fuchs, H. Gladman, R. Jain, Y.A. Khan, C.M.R. Low, K. Perlin, A. Potapenko, P. Savy, S.
Singh, A. Stecula, A. Thillaisundaram, C. Tong, S. Yakneen, E.D. Zhong, M. Zielinski, A.
Žídek, V. Bapst, P. Kohli, M. Jaderberg, D. Hassabis, J.M. Jumper, Accurate structure
prediction of biomolecular interactions with AlphaFold 3, Nature 2024 630:8016 630 (2024)
493–500. https://doi.org/10.1038/s41586-024-07487-w.

[48] L. Holm, A. Laiho, P. Törönen, M. Salgado, DALI shines a light on remote homologs: One
hundred discoveries, Protein Science 32 (2023) e4519. https://doi.org/10.1002/PRO.4519.

[49] K. Remans, K. Pauwels, P. van Ulsen, L. Buts, P. Cornelis, J. Tommassen, S.N. Savvides,
K. Decanniere, P. Van Gelder, Hydrophobic Surface Patches on LolA of *Pseudomonas*
aeruginosa Are Essential for Lipoprotein Binding, J Mol Biol 401 (2010) 921–930.
https://doi.org/10.1016/J.JMB.2010.06.067.

[50] K. Takeda, H. Miyatake, N. Yokota, S.I. Matsuyama, H. Tokuda, K. Miki, Crystal structures
of bacterial lipoprotein localization factors, LolA and LolB, EMBO J 22 (2003) 3199–3209.
https://doi.org/10.1093/EMBOJ/CDG324.

[51] J. Botas, Á. Rodríguez Del Río, J. Giner-Lamia, J. Huerta-Cepas, GeCoViz: genomic
context visualisation of prokaryotic genes from a functional and evolutionary perspective,
Nucleic Acids Res 50 (2022) W352–W357. https://doi.org/10.1093/NAR/GKAC367.

[52] J. Herson, M. Krummenacker, A. Spaulding, P. O'Maille, P.D. Karp, The Genome Explorer
genome browser, MSystems 9 (2024). https://doi.org/10.1128/MSYSTEMS.00267-
24/ASSET/D092CD98-EDF4-4F91-B5B5-
0333B26B6D93/ASSETS/IMAGES/LARGE/MSYSTEMS.00267-24.F007.JPG.

[53] E.C. Meng, T.D. Goddard, E.F. Pettersen, G.S. Couch, Z.J. Pearson, J.H. Morris, T.E.
Ferrin, UCSF ChimeraX: Tools for structure building and analysis, Protein Science 32
(2023) e4792. https://doi.org/10.1002/PRO.4792.

[54] S. Bello, S. McQuay, B. Rudra, R.S. Gupta, Robust demarcation of the family
Peptostreptococcaceae and its main genera based on phylogenomic studies and taxon-
specific molecular markers, Int J Syst Evol Microbiol 74 (2024) 006247.
https://doi.org/10.1099/IJSEM.0.006247/CITE/REFWORKS.

[55] J.W. O'Neill, D.E. Kim, K. Johnsen, D. Baker, K.Y.J. Zhang, Single-site mutations induce
3D domain swapping in the B1 domain of Protein L from *Peptostreptococcus magnus*,

Structure 9 (2001) 1017–1027. https://doi.org/10.1016/S0969-2126(01)00667-0/ASSET/72670EC8-3DCE-47A6-B0BE-0F27AFC6C8EF/MAIN.ASSETS/FX3.JPG.

[56] K.H. Miller, J.R. Karr, S. Marqusee, A Hinge Region cis-Proline in Ribonuclease A Acts as a Conformational Gatekeeper for C-Terminal Domain Swapping, J Mol Biol 400 (2010) 567–578. https://doi.org/10.1016/J.JMB.2010.05.017.

[57] M. Orlikowska, E. Jankowska, R. Kołodziejczyk, M. Jaskólski, A. Szymańska, Hinge-loop mutation can be used to control 3D domain swapping and amyloidogenesis of human cystatin C, J Struct Biol 173 (2011) 406–413. https://doi.org/10.1016/J.JSB.2010.11.009.

[58] G. Calloni, T. Chen, S.M. Schermann, H.C. Chang, P. Genevaux, F. Agostini, G.G. Tartaglia, M. Hayer-Hartl, F.U. Hartl, DnaK Functions as a Central Hub in the *E. coli* Chaperone Network, Cell Rep 1 (2012) 251–264. https://doi.org/10.1016/j.celrep.2011.12.007.

[59] M. Kovermann, F.X. Schmid, J. Balbach, Molecular function of the prolyl cis/trans isomerase and metallochaperone SlyD, Biol Chem 394 (2013) 965–975. https://doi.org/10.1515/HSZ-2013-0137/ASSET/GRAPHIC/HSZ-2013-0137_CV3.JPG.

[60] S. Hottenrott, T. Schumann, A. Plückthun, G. Fischer, J.U. Rahfeld, The *Escherichia coli* SlyD is a metal ion-regulated peptidyl-prolyl cis/trans-isomerase, Journal of Biological Chemistry 272 (1997) 15697–15701. https://doi.org/10.1074/jbc.272.25.15697.

[61] A. Chikunova, M.P. Manley, C.N. Heijjer, C.S. Drenth, A.J. Cramer-Blok, M.U.D. Ahmad, A. Perrakis, M. Ubbink, Conserved proline residues prevent dimerization and aggregation in the β-lactamase BlaC, Protein Science 33 (2024) e4972. https://doi.org/10.1002/PRO.4972.

[62] H. Fu, G.R. Grimsley, A. Razvi, J.M. Scholtz, C.N. Pace, Increasing protein stability by improving beta-turns, Proteins: Structure, Function, and Bioinformatics 77 (2009) 491–498. https://doi.org/10.1002/PROT.22509.

[63] C.M. Wilmot, J.M. Thornton, Analysis and prediction of the different types of β-turn in proteins, J Mol Biol 203 (1988) 221–232. https://doi.org/10.1016/0022-2836(88)90103-9.

[64] E.G. Hutchinson, J.M. Thornton, A revised set of potentials for β-turn formation in proteins, Protein Science 3 (1994) 2207–2216. https://doi.org/10.1002/PRO.5560031206.

[65] M.W. MacArthur, J.M. Thornton, Influence of proline residues on protein conformation, J Mol Biol 218 (1991) 397–412. https://doi.org/10.1016/0022-2836(91)90721-H.

[66] N. Nandwani, P. Surana, H. Negi, N.M. Mascarenhas, J.B. Udgaonkar, R. Das, S. Gosavi, A five-residue motif for the design of domain swapping in proteins, Nature Communications 2019 10:1 10 (2019) 1–13. https://doi.org/10.1038/s41467-019-08295-x.

[67] F. Ding, K.C. Prutzman, S.L. Campbell, N. V. Dokholyan, Topological determinants of protein domain swapping, Structure 14 (2006) 5–14. https://doi.org/10.1016/J.STR.2005.09.008.

[68] J.W. O'Neill, D.E. Kim, K. Johnsen, D. Baker, K.Y.J. Zhang, Single-site mutations induce 3D domain swapping in the B1 domain of Protein L from *Peptostreptococcus magnus*,

Structure 9 (2001) 1017–1027. https://doi.org/10.1016/S0969-2126(01)00667-0/ASSET/72670EC8-3DCE-47A6-B0BE-0F27AFC6C8EF/MAIN.ASSETS/FX3.JPG.

[69] A. Cámara-Artigas, S. Martínez-Rodríguez, E. Ortiz-Salmerón, J.M. Martín-García, 3D domain swapping in a chimeric c-Src SH3 domain takes place through two hinge loops, J Struct Biol 186 (2014) 195–203. https://doi.org/10.1016/J.JSB.2014.02.007.

[70] A. Szymánska, E. Jankowska, M. Orlikowska, I. Behrendt, P. Czaplewska, S. Rodziewicz-Motowidło, Influence of point mutations on the stability, dimerization, and oligomerization of human cystatin C and its L68Q variant, Front Mol Neurosci 5 (2012) 82–82. https://doi.org/10.3389/FNMOL.2012.00082/PDF<IMG.

[71] M. Bergdoll, M. Remy, C. Cagnon, J.M.- Structure, undefined 1997, Proline-dependent oligomerization with arm exchange, Cell.ComM Bergdoll, MH Remy, C Cagnon, JM Masson, P DumasStructure, 1997•cell.Com (n.d.). https://www.cell.com/AJHG/fulltext/S0969-2126(97)00196-2 (accessed May 14, 2025).

[72] A. Di Donate, V. Cafaro, G. D'Alessio, Ribonuclease A can be transformed into a dimeric ribonuclease with antitumor activity, Journal of Biological Chemistry 269 (1994) 17394–17396. https://doi.org/10.1016/S0021-9258(17)32451-1.

[73] F.U. Battistuzzi, A. Feijao, S.B. Hedges, A genomic timescale of prokaryote evolution: Insights into the origin of methanogenesis, phototrophy, and the colonization of land, BMC Evol Biol 4 (2004). https://doi.org/10.1186/1471-2148-4-44,.

[74] T.J.P. Hubbard, A.G. Murzin, S.E. Brenner, C. Chothia, SCOP: A Structural Classification of Proteins database, Nucleic Acids Res 25 (1997) 236–239. https://doi.org/10.1093/NAR/25.1.236,.

[75] D. Megrian, N. Taib, J. Witwinowski, C. Beloin, S. Gribaldo, One or two membranes? Diderm Firmicutes challenge the Gram-positive/Gram-negative divide, Mol Microbiol 113 (2020) 659–671. https://doi.org/10.1111/MMI.14469.

[76] N. Taib, D. Megrian, J. Witwinowski, P. Adam, D. Poppleton, G. Borrel, C. Beloin, S. Gribaldo, Genome-wide analysis of the Firmicutes illuminates the diderm/monoderm transition, Nature Ecology & Evolution 2020 4:12 4 (2020) 1661–1672. https://doi.org/10.1038/s41559-020-01299-7.

[77] J. Witwinowski, A. Sartori-Rupp, N. Taib, N. Pende, T.N. Tham, D. Poppleton, J.M. Ghigo, C. Beloin, S. Gribaldo, An ancient divide in outer membrane tethering systems in bacteria suggests a mechanism for the diderm-to-monoderm transition, Nature Microbiology 2022 7:3 7 (2022) 411–422. https://doi.org/10.1038/s41564-022-01066-3.

[78] G.E. Crooks, G. Hon, J.M. Chandonia, S.E. Brenner, WebLogo: A Sequence Logo Generator, Genome Res 14 (2004) 1188–1190. https://doi.org/10.1101/GR.849004.

[79] A.P. Arkin, R.W. Cottingham, C.S. Henry, N.L. Harris, R.L. Stevens, S. Maslov, P. Dehal, D. Ware, F. Perez, S. Canon, M.W. Sneddon, M.L. Henderson, W.J. Riehl, D. Murphy-Olson, S.Y. Chan, R.T. Kamimura, S. Kumari, M.M. Drake, T.S. Brettin, E.M. Glass, D. Chivian, D. Gunter, D.J. Weston, B.H. Allen, J. Baumohl, A.A. Best, B. Bowen, S.E. Brenner, C.C. Bun, J.M. Chandonia, J.M. Chia, R. Colasanti, N. Conrad, J.J. Davis, B.H. Davison, M. Dejongh, S. Devoid, E. Dietrich, I. Dubchak, J.N. Edirisinghe, G. Fang, J.P.

Faria, P.M. Frybarger, W. Gerlach, M. Gerstein, A. Greiner, J. Gurtowski, H.L. Haun, F. He,
R. Jain, M.P. Joachimiak, K.P. Keegan, S. Kondo, V. Kumar, M.L. Land, F. Meyer, M. Mills,
P.S. Novichkov, T. Oh, G.J. Olsen, R. Olson, B. Parrello, S. Pasternak, E. Pearson, S.S.
Poon, G.A. Price, S. Ramakrishnan, P. Ranjan, P.C. Ronald, M.C. Schatz, S.M.D. Seaver,
M. Shukla, R.A. Sutormin, M.H. Syed, J. Thomason, N.L. Tintle, D. Wang, F. Xia, H. Yoo,
S. Yoo, D. Yu, KBase: The United States Department of Energy Systems Biology
Knowledgebase, Nature Biotechnology 2018 36:7 36 (2018) 566–569.
https://doi.org/10.1038/nbt.4163.

[80] M.N. Price, P.S. Dehal, A.P. Arkin, FastTree 2 – Approximately Maximum-Likelihood Trees
for Large Alignments, PLoS One 5 (2010) e9490.
https://doi.org/10.1371/JOURNAL.PONE.0009490.

[81] I. Letunic, P. Bork, Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic
tree display and annotation tool, Nucleic Acids Res 52 (2024) W78–W82.
https://doi.org/10.1093/NAR/GKAE268.

[82] Y.K. Ng, M. Ehsaan, S. Philip, M.M. Collery, C. Janoir, A. Collignon, S.T. Cartman, N.P.
Minton, Expanding the Repertoire of Gene Tools for Precise Manipulation of the
Clostridium difficile Genome: Allelic Exchange Using pyrE Alleles, PLoS One 8 (2013)
e56051. https://doi.org/10.1371/JOURNAL.PONE.0056051.

[83] T. Karasawa, S. Ikoma, K. Yamakawa, S. Nakamura, A defined growth medium for
Clostridium difficile, Microbiology (N Y) 141 (1995) 371–375.
https://doi.org/10.1099/13500872-141-2-371,.

[84] M. Lauren Donnelly, W. Li, Y.Q. Li, L. Hinkel, P. Setlow, A. Shen, A *Clostridium difficile*-
specific, gel-forming protein required for optimal spore germination, MBio 8 (2017).
https://doi.org/10.1128/MBIO.02085-16,.

[85] R.A. Laskowski, J. Jabłońska, L. Pravda, R.S. Vařeková, J.M. Thornton, PDBsum:
Structural summaries of PDB entries, Protein Science 27 (2018) 129–134.
https://doi.org/10.1002/PRO.3289.

Figure 1. PLLLF proteins are structurally similar but share low sequence identity. A)
Comparisons of sequence identity and sequence similarity between 17 representative PLLLF
proteins. Gradients represent percentages from 0 to 100. B) Ribbon representation of the crystal
structure of the GerS-CwlD complex (PBD 7RAG) with the GerS strand swap colored yellow.
The other GerS is colored in transparent blue, and both CwlD proteins are colored in
transparent pink. C) Ribbon representation of the crystal structures of the *Escherichia coli*
LolA^WT (PDB 1IWL) and LolA^F68E (PDB 6FHM). The strand swap in LolA^F68E structure is shown
in yellow; the corresponding β-strand in LolA^WT is shown in yellow. The other LolA^F68E protomer
in the homodimer is colored in transparent blue. Residue 68, which is a phenylalanine in LolA^WT
and a glutamate in LolA^F68E, is colored green.

Figure 2. GerS dimerization is conserved in the Peptostreptococcaceae family despite being distributed throughout the Bacilliota phylum. A) Synteny analysis of the conserved *gerS* locus across the orders of the Bacilliota phylum. Peptostreptococcales (blue), Tissierelliales (turquoise), Lachnospirales (purple), Eubacterales (green), Selenomonadales (pink), Vellionellales (orange), Bacillales (red), and the outgroup Enterbacterales (yellow) classes are shown. Colored stars correspond to representative GerS orthologs selected for SEC analysis. B) Size exclusion chromatography (SEC) analysis of GerS orthologs tagged with His$_6$ for purification. Peaks are labeled with "D" for dimer or "M" for monomer.

Figure 3. GerS hinge mutations guided by the LolA structure modulate its dimer:monomer equilibrium. A) Ribbon representation of the crystal structure of the GerS-CwlD complex (PDB 7RAG). The hinge is colored by the residue shown in the inset. B) Ribbon representation of the AlphaFold3 GerS structure and LolA crystal structure (PDB - 1IWL). Hinge of GerS and analogous residues on LolA are colored by aligned residues. C) Size exclusion chromatography (SEC) traces of each GerS variant tagged with His$_6$. Peaks are labeled with "D" for dimer or "M" for monomer. D) Ratio of the areas under each peak as a percentage of the total area measured. "D" is the area under the dimer peak, and "M" is the area under the monomer peak.

Figure 4. Prolines in the hinge region of PLLLF proteins reduce dimerization. A) Size
exclusion chromatography (SEC) traces of His$_6$-tagged GerS variants carrying single proline
substitutions in the hinge region. B) Sequence logo depicting the frequency of amino acids
found at a given position based on DALI structural alignments of 17 representative PLLLF
proteins [48]. Prolines are highlighted in blue. (C) Amino acids aligned to each conserved
proline in GerS and LolA are displayed in the ribbon representations of the crystal structures
shown (PDB 7RAG and 1IWL, respectively). D) SEC trace of LolA^{WT} and LolA^{P65T} with area
under the curve calculations. E) SEC trace of dual proline GerS variants. Peaks are labeled with
either "D" for dimer or "M" for monomer.

Figure 5. GerS substitutions that reduce dimerization decrease binding to CwlD and spore germination. A) SEC traces of GerS hinge region variants copurified with His_6-tagged CwlD. Peaks labeled with "GerS:CwlD-His_6", correspond to the complex formed between GerS and CwlD-His_6. Peaks labeled with "GerS" and "CwlD-His_6", represent the individual proteins that elute outside of complex. B) Coomassie stain of untagged GerS variants copurified with His_6-tagged CwlD. C) Western blot analyses of sporulating cell or purified spore lysates prepared from Δ*gerS* complemented with constructs encoding GerS hinge region variants. SleC was used as a loading control. D) Relative germination based on the change in optical density (OD_{600}) of purified spores over time following the addition of 1% (19 mM) taurocholate in rich media. Germination in arbitrary units was calculated using the area below inverted OD_{600} curves. Each *gerS* mutant is expressed by its native promoter in the *pyrE* locus in a Δ*gerS* background. A one-way ANOVA followed by Dunnett's multiple comparisons test was used to

842   compare the mean of each *gerS* mutant to the Δ*gerS*/*gerS* complement strain based on a

843   minimum of three biological replicates. Significance relative to WT is shown. (**** $p < 0.00005$).

Figure 6. GerS interaction with CwlD is dependent on both strand-swapping and a salt bridge. A) GerS:CwlD structure (7RAG). SB, Salt bridge; H, Hinge. B) Coomassie-stained eluates of untagged homodimer-disrupting GerS variants combined with the salt bridge mutation D106R copurified with His$_6$-tagged CwlD. C) Relative germination calculated by area under the percent change in OD600 curve of spores in BHIS in the presence of 1% taurocholate over 1.5 hours. Each *gerS* mutant is expressed by its native promoter in the *pyrE* locus in a Δ*gerS* background. A one-way ANOVA followed by Dunnett's multiple comparisons test was used to compare each *gerS* mutant mean to the mean Δ*gerS*/*gerS* complement strain. Each mean is calculated from at least three biological replicates. Germination assay and statistics were run with Figure 5D, therefore Wild Type, Δ*gerS*, Δ*cwlD,* and Δ*gerS*/*gerS* complement are the same between both panels. The significance relative to WT is shown. (**** $p < 0.00005$). D) Detection of GerS variants by western blotting of sporulating cell or purified spore lysates. Polyclonal antibodies against GerS, CwlD, and SleC were used.