The Power of Three: Dynactin associates with three dyneins under load for greater force production

Authors:

Lu Rao[1,#], Xinglei Liu[1], Mirjam Arnold[4], Rick McKenney[3], Kristy Stengel[4], Simone Sidoli[5], Florian Berger[2], and Arne Gennerich[1,#]

[1]Department of Biochemistry and Gruss Lipper Biophotonics Center, Albert Einstein College of Medicine, Bronx, NY 10461
[2]Department of Biology, Utrecht University, 35845 Utrecht, The Netherlands
[3]Department of Molecular and Cellular Biology, University of California Davis, Davis, CA 95616
[4]Department of Cell Biology, Albert Einstein College of Medicine, Bronx, NY 10461
[5]Department of Biochemistry, Albert Einstein College of Medicine, Bronx, NY 10461

[#] Correspondence: lu.rao@einsteinmed.edu and arne.gennerich@einsteinmed.edu

Abstract

Cytoplasmic dynein is an essential microtubule motor protein involved in organelle transport and mitotic spindle assembly. Its activity depends on forming dynein-dynactin-cargo adaptor complexes, such as dynein-dynactin-BicD2 (DDB), which typically operate with two dynein motors. We discovered that mechanical tension recruits a third dynein motor via an auxiliary BicD adaptor binding the light intermediate chain of the third dynein. This second BicD molecule stabilizes multi-dynein assemblies, enabling high-force generation. Lis1 further facilitates this adaptation by alleviating dynein autoinhibition. Single-dynein DDB generates ~2.5 pN of force, increasing to ~4.5 pN with Lis1, while complexes with two or three motors produce ~7 pN and ~9 pN, respectively, consistent with a staggered motor arrangement. Under load, DDB complexes primarily take ~8 nm steps, challenging current models of dynein coordination. These findings reveal how DDB complexes dynamically sense and adapt to mechanical challenges, ensuring robust intracellular transport under varying force demands.

INTRODUCION

Cytoplasmic dynein is the primary microtubule (MT) minus-end-directed motor in eukaryotic cells, playing essential roles in processes such as organelle transport, mitotic spindle positioning, and neuronal development[1-6]. Its activity is tightly regulated by cofactors, including lissencephaly-1 (Lis1), a protein critical for brain development[7-13]. Lis1 enables dynein to generate motion and force along MTs, and disturbances in Lis1 function—due to mutations or deletions—lead to type I lissencephaly[14,15], a severe developmental brain disorder[16-18]. Despite its significance, the precise mechanisms by which Lis1 controls dynein function remain unclear, posing challenges for therapeutic advancements.

Dynein function is further modulated by interactions with dynactin, its largest co-factor. Dynactin not only enhances dynein's processivity[1,4,19-29]—the ability to take consecutive steps without detaching—but also links dynein to cargoes and positions it within specific cellular domains[3,30,31]. Mutations in dynactin are associated with neurodegenerative diseases[32-34] such as distal spinal bulbar muscular atrophy (dSBMA)[33] and Perry syndrome[34], highlighting its physiological importance. The coiled-coil cargo adaptor Bicaudal-D2 (BicD2) bridges the dynein tail and the dynactin shoulder[35,36] (**Fig. 1A**), converting dynein from a diffusive[37] or weakly processive[38,39] entity into an ultraprocessive motor[24,25,40]. Recent studies have revealed that cargo adaptors like BicD2 can recruit two dynein motors to the dynein-dynactin-BicD (DDB-2) complex, forming a configuration of four parallel motor domains[41,42] (**Fig. 1A**). This arrangement enhances force generation via load-sharing and improves processivity by reducing detachment from MTs[42].

The role of Lis1 within the DDB complex remains poorly understood. Initial studies suggested that Lis1 inhibits dynein motion by uncoupling ATPase activity from MT-binding dynamics[43-45]. More recent evidence indicates that Lis1 activates dynein by preventing its auto-

inhibited "phi" state[46-53], facilitating the recruitment of a second dynein dimer to the DDB complex[54,55]. Supporting this, Lis1 is thought to detach from dynein upon MT binding[56], suggesting its primary role is motor recruitment rather than the regulation of active dynein motion.

Despite recent structural and mechanistic insights into the function of dynein-dynactin-cargo adaptor complexes, key questions remain unresolved: How does force output scale with the number of dyneins in a complex? Studies of dynein-dynactin-BICDR1 (DDR) complexes have identified a second auxiliary BICDR1 molecule bound to dynactin as of yet unknown function. Furthermore, the interplay between Lis1-mediated dynein activation and motor recruitment under load is poorly understood, leaving significant gaps in our understanding of how these complexes dynamically adapt to varying intracellular force demands. Addressing these questions is essential to uncovering the principles governing dynein regulation and intracellular transport.

Here, we combined optical trapping, three-color single-molecule fluorescence, and structural analyses to uncover new principles governing dynein regulation within DDB complexes under mechanical tension. We reveal that Lis1 plays a dual role in activating DDB bound to a single dynein (DDB-1) by relieving dynein's autoinhibited "phi" conformation and facilitating its transition to a fully active state, thereby increasing force generation from ~2.5 pN to ~4.5 pN. Notably, we discovered that the DDB complex adapts to increasing load demands by recruiting a third dynein motor (DDB-3), a process mediated by an auxiliary BicD adaptor that binds the light intermediate chain (LIC) of the third dynein. This finding highlights a previously unrecognized role of a second BicD molecule in stabilizing multi-dynein assemblies and enabling higher-force generation. Furthermore, the distinct forces generated by DDB-2 (~7 pN) and DDB-3 (~9 pN)—incremental increases of only ~2 pN and ~2.5 pN from fully activated DDB-1—suggest a staggered arrangement of dyneins within the complex, with load distributed unevenly among

motors. Fully assembled DDB complexes predominantly take load-independent steps of ~8 nm under tension, with smaller steps emerging near stalling forces, suggesting a compact yet adaptable motor configuration. Our results provide new mechanistic insights into Lis1-mediated dynein activation, the functional role of auxiliary cargo adaptors in motor assembly, and the dynamic, tension-driven recruitment of dynein motors, offering a framework for understanding intracellular force generation and transport in complex cellular environments.

RESULTS

"Trains" of DDB Force Generation Reveal Distinct Stalling Forces

Previous studies on DDB force generation have primarily utilized non-native axonemes—160-nm thick MT-containing structures purified from sea urchin[57]—and sub-stoichiometric dynein-dynactin-BicD concentrations (molar ratios of dynein-dynactin-BicD of 1:5:2[58], 1:5:20[42,59] and 1:2:10[54]). These conditions disfavored the formation of DDB complexes with two dyneins and limited the detection of distinct force-generation states.

For example, when DDB is tethered to trapping beads via a motor-less dynein tail, which permits only one dynein to bind while the second dynein-binding site is occupied by the tail fragment, the complex generates a force of 3.7 ± 0.2 pN (mean $\pm$ SEM) on axonemes[54]. This force matches that generated by DDB linked to beads via BicD[42], suggesting that these studies primarily involved complexes with only a single dynein. Similarly, *S. cerevisiae* dynein alone generates 3.6 ± 0.2 pN on axonemes[58] and 4.50 ± 0.04 pN on MTs[39,60], indicating potential differences in DDB behavior on MTs.

To investigate the function of dynein-dynactin-adaptor (DDA) complexes with two dynein motors (DDA-2) (**Fig. 1A**), previous studies assembled motor complexes using the cargo adaptors HOOK3 and BICDR1, which are more likely than BicD to recruit two dyneins to dynactin[42]. As dynein-dynactin-HOOK3 (DDH, 4.9 ± 0.2 pN[42]) and dynein-dynactin-BICDR1 (DDR, 6.5 ± 0.3 pN[42]) generate more force than DDB (3.7 ± 0.2 pN[42]), it was reasoned that DDH and DDR recruit two dyneins. However, these studies were conducted on axonemes with a 1:5 molar dynein-dynactin ratio[42,59], conditions under which, on average, only one in five dynactin complexes can bind a dynein. Thus, these complexes were unlikely to be fully saturated with two dyneins, leaving the source of their higher force generation unclear.

To address these limitations and gain insights into force production, we conducted optical trapping experiments using DDB complexes purified from rat brain lysate with a BicD-sfGFP fragment (amino acids 25–400)[24] (**Fig. 2A**). Since neither dynein nor dynactin alone binds to BicD[24], we anticipated that these complexes would contain dynein and dynactin in a 1:1 or 2:1 ratio. We then employed 500-nm trapping beads covalently linked to anti-GFP antibodies and used MTs instead of axonemes. Strikingly, we observed extended "trains" of force-generation events characterized by rapid force increases followed by stalling of movement (**Fig. 2C-F**). These force trains revealed four distinct stall forces for DDB: ~2.5, ~4.5, ~7, and ~9 pN (**Fig. 2B-F**). This discovery contrasts with earlier studies that reported broad detachment forces encompassing the plateaus we identified in our analyses (2–7 pN for DDB, 2.5–9 pN for DDH, and 3–11 pN for DDR)[51,55]. We speculate that the detection of these distinct plateaus in our experiments is due to the use of smaller beads, which, due to lower vertical forces, are likely to increase the DDB-MT on-rate and the likelihood of observing stalling (it's worth noting that the effects of varying vertical forces on human dynein have not been tested, but higher vertical forces enhance MT detachment of kinesin-1 and prevent motor stalling[61,62]).

DDB-1 Stalls at ~2.5 or ~4.5 pN Depending on Its Partial or Full Activation

Since DDB complexes purified from mouse brain lysate demonstrated a myriad of generated forces (**Fig. 2C-F**), we sought to have a more controlled ratio of the assembling components to further investigate each force profile. Additionally, because we frequently observed only 1–2 events per bead for forces of ~9 pN (**Fig. 2F**), we reasoned the binding strength between the anti-GFP antibody and GFP might not be strong enough to sustain large forces over time. To address this issue, we switched to using streptavidin-coated beads instead of anti-GFP-antibody-

coated beads, taking advantage of the robust streptavidin-biotin linkage known to withstand forces up to 60 pN in force-extension experiments[63].

We purified recombinant full-length human dynein, co-expressed with its five accessory chains, as previous desribed[25]. Biotinylated BicD (amino acids 25–400) was expressed in *E. coli* and purified (**Fig. 1B**). For dynactin, traditionally purified from bovine, porcine, rodent brain tissue[19,24,25], we employed a CRISPR-based approach to tag the DCTN4 subunit with a HaloTag-3×FLAG at its C-terminus in a human suspension culture cell line (HEK293sus), following established protocols[64]. This tagging strategy, recently used to generate a stable adherent HEK292 cell line[65], enabled efficient recombinant expression of dynactin (**Fig. 1B**). With each component—dynein, BicD2, and dynactin—expressed and purified to high purity and concentration, we successfully assembled DDB complexes in defined stoichiometric ratios.

To investigate the force generation of DDB when associated with a single dynein, we assembled the complexes in a 1:1:1 ratio of dynein, dynactin, and BicD. Our previous studies have indicated that isolated human dynein can transition from slow movement toward ~2.5 pN to a rapid movement toward ~4 pN (**Fig. 3A**, reproduced from Fig. S3 in ref[39]). We observed a similar behavior in DDB complexes, transitioning from slow movement toward ~2.5 pN followed to rapid movement toward ~4.5 pN (**Fig. 3B**). Interestingly, for human dynein alone, we only observed instances of transitioning from ~2.5 to ~4 pN under increased trap stiffnesses (~0.06 pN/nm); at a lower trap stiffness of 0.01 pN/nm, it stalls at ~1 pN (**Fig. 3A**, inset, and Fig. S3 in ref[39]), suggesting that a higher force per unit displacement facilitates the full activation of dynein. This indicates that the two distinct forces are generated by a single dynein altering its behavior depending on its activation state.

Recent studies have shown that mammalian dynein often adopts an auto-inhibited conformation, commonly referred to as the "phi" particle[48-52], which hinders efficient interactions with MTs. Unlike mammalian dynein, *S. cerevisiae* dynein moves processively along MTs by itself[66] and a mutant form unable to assume the phi conformation exhibits an increased run length along MTs[46], indicating that the auto-inhibited state limits the movement of wild-type *S. cerevisiae* dynein on MTs. To investigate whether a similar auto-inhibited state restricts in human dynein's force generation to ~2.5 pN, we assembled DDB complexes at a 1:1:1 ratio with a dynein mutant (K1610E and R1567E double mutant[48]) incapable of adopting this inhibitory conformation, referred to here as D(phi)DB. Contrary to expectations, D(phi)DB complexes predominantly stalled at ~7 pN (**Fig. 3C** and **Suppl. Fig. 1A**), even when assembled at a molar ratio of 1:6:6 or in the presence of excess dynein tail (**Suppl. Fig. 1B&C**), suggesting that dynein in its open conformation tends to cluster and form DDB complexes with two dyneins per complex (discussed further in the following section). However, this dynein mutant prevents stalling at ~2.5 pN in the lower force regime, with only ~4.5 pN forces observed (**Fig. 3C** and **Suppl. Fig. 1A&B**). Moreover, the regulatory protein Lis1, which prevents dynein's auto-inhibited state[46,47,53-55], significantly reduces the occurrence of stalling at ~2.5 pN (**Fig. 3C**). These results confirm that a DDB complex bound to a single dynein stalls at either ~2.5 or ~4.5 pN depending on whether it is partially or fully activated.

Fully active DDB-2 stalls 7 pN

D(phi)DB complexes generate forces of ~4.5, ~7, or ~9 pN (**Suppl. Fig. 1A**), suggesting that fully active DDB-1 generates ~4.5 pN, while a DDB complex with two dyneins (DDB-2)

generates ~7 pN. This observation aligns with that of DDR, which binds two dyneins preferentially and is reported to generate 6.5 pN[42].

A key question is what accounts for the observed stalling at 9 pN. If fully active DDB-1 generates ~4.5 pN, then logically, fully active DDB-2 might be expected to reach up to ~9 pN if both motors contribute equally to force generation. However, DDR assembled with the dynein mutant and in the presence of Lis1—both intended to prevent the phi conformation—has been observed to generate only 6.1 pN[54], which aligns more closely with the ~7 pN force we measure. This discrepancy suggests that the ~7 pN stall force does not arise from a collaboration between a fully active dynein and a partially active motor. Instead, it likely results from uneven load distribution between the motors, with the leading motor bearing a greater load than the lagging dynein, contributing only ~2.5 pN.

To further support the hypothesis that the ~7 pN force is generated by two actively engaged dyneins, we conducted experiments by introducing 5 nM dynein into a slide chamber containing pre-assembled 1:1:1 ratio DDB complexes bound to trapping beads. This resulted in ~60% of the moving beads displaying forces of ~7 pN (**Fig. 4E** and **Suppl. Fig. 2**). Notably, the proportion of beads showing force generation remained consistent with that observed in DDB-only complexes (~10%), despite the addition of free dynein (**Fig. 4D**). This indicates that the freely diffusing dyneins selectively bind to DDB complexes anchored via the biotinylated BicD to the bead surfaces, rather than non-specifically binding to the bead surfaces, which would have increased the fraction of force-generating beads[39].

Auxiliary BicD Enables Third Dynein Recruitment and Enhanced Force Generation

Given the observed force trains at ~9 pN and the likely contribution of two fully active dyneins to the ~7 pN force, we hypothesized that the ~9 pN force results from the functional recruitment of a third dynein to the DDB complex. Although this possibility has not been previously explored, the cryo-EM structure of two dyneins bound to dynactin and BICDR1[42] suggests that the dynactin shoulder provides sufficient space for a third dynein tail to bind (**Fig. 4F&G**).

To test this hypothesis, we examined whether dynactin can accommodate a third dynein by generating two truncated BicD constructs, BicD(25–276) and BicD(25–180), based on the cryo-EM structure of DDB[42]. BicD(25–276) spans the full shoulder of dynactin, encompassing the putative third dynein-binding site (**Fig. 4A&F**), whereas BicD(25–180) extends only to the position occupied by the middle dynein (dynein-A[42]) (**Fig. 4A&F**). Using single-molecule fluorescence and optical-tweezers assays, we assessed the functionality of these constructs. BicD(25–276) formed DDB complexes capable of motion and force generation (**Suppl. Fig. 3A**), similar to BicD(25–400) (**Fig. 2**). In contrast, the shortest construct, BicD(25–180), not only failed to form DDB complexes with two dyneins (DDB-2) but also formed significantly fewer complexes with one dynein (DDB-1) (**Suppl. Fig. 4**), generating forces of only ~2.5 or ~4.5 pN (**Suppl. Fig. 3B**). These results indicate that BicD's ability to span the full dynactin shoulder is essential for forming and stabilizing DDB complexes, although the terminal segment (amino acids 277–400) appears unnecessary for dynein recruitment. Consequently, the essential role of the middle segment (amino acids 180–276) leaves open the question of whether it participates in the binding of the third dynein.

A recent cryo-EM study revealed that a second BicDR1 molecule (BICDR-B) can stabilize the MT-bound DDR complex by acting as an auxiliary adaptor[67] (**Fig. 4A**). While a second BicD

molecule has not yet been visualized in DDB cryoEM structures, we investigated the possibility that a second auxiliary BicD adaptor facilitates the recruitment of the third dynein by employing the shortest construct, BicD(25–180), without a biotinylated C-terminus. This modified construct, incapable of binding to streptavidin beads or forming functional DDB complexes on its own, retains the segment capable of interacting with pre-assembled DDB complexes that include BicD(25–400) anchored to streptavidin beads. When 5 nM free dynein and 5 nM BicD(25–180) were added to DDB complexes assembled at a 1:1:1 ratio (which on their own generate only ~2.5 or ~4.5 pN), we observed dominant ~9 pN forces in ~60% of all force-generating beads (**Fig. 4E** and **Suppl. Fig. 5**). These results strongly suggest that a second auxiliary BicD adaptor plays a critical role in recruiting a third dynein.

Further insights into this interaction came from cryo-EM structures of DDR complexes with two BicDR1 molecules[68] (**Fig. 4A**). The CC1-Box (AAxxG; "x" marks any amino acid) of the N-terminus of the primary BicDR1 (BICDR-A) interacts with α-helix-1 of the C-terminus of the light intermediate chain (LIC) from dynein B (B1)[68] and with the LIC of dynein-A (A2) (**Fig. 4A&B**). Indeed, removing the N-terminus of BicD completely abolished the formation of DDB complex (**Suppl. Fig. 4**). In contrast, the auxiliary BicDR1 (BICDR-B) N-terminus interacts exclusively with the LIC of dynein-A2[68]. Although the LIC of dynein-A2 interacts with both BICDR-A and BICDR-B[68], the latter's unoccupied surface may bind the LIC of the third dynein. To directly assess whether disrupting the interaction between the auxiliary BicD and the third dynein's LIC impairs recruitment, we introduced a single-point mutation (Y46D) in BicD(25–180). This mutation disrupts the hydrophobic interaction between the BicD N-terminus and the dynein LIC[69]. When 5 nM mutated BicD(25–180, Y46D) and 5 nM free dynein were added to pre-assembled DDB complexes, we observed a significant reduction in force generation, with less than

13% of beads producing ~9 pN forces (**Fig. 4E**). These findings demonstrate that recruitment of the third dynein critically depends on the interaction between the second auxiliary BicD and the third dynein's LIC.

Tension-Driven Recruitment of a Third Dynein to DDB Complexes

A potential question arises: why have cryo-EM and cryo-ET studies not observed the presence of a third dynein, despite its capability to associate with DDB? The explanation likely lies in the experimental conditions. Unlike structural studies, our optical-trapping assay applied mechanical load to DDB complexes (**Fig. 2A**), and this mechanical tension revealed a step-like increase in stalling forces (**Fig. 2G**). This intriguing phenomenon suggests that mechanical tension facilitates the recruitment of additional dynein motors to DDB complexes, enabling adaptation to increased force demands.

To investigate this concept, we engineered a kinesin-DDB chimera by covalently linking a DDB complex through BicD to the C-terminal tail domain of a kinesin-1 rigor mutant[70] using the SpyTag-SpyCatcher system[71]. Specifically, we inserted the SpyCatcher (SpyCatcher003[72]) at the C-terminus of BicD—typically bound to cargo—and fused the SpyTag (SpyTag003[72]) to the C-terminus of the kinesin mutant (**Fig. 5A&B**). This rigor kinesin mutant is characterized by its strong binding to MTs and lack of motility[70]. We hypothesized that if the kinesin-DDB chimera attaches to the MT with the kinesin motor domains and the DDB complex bound to the MT, the DDB complex must generate enough force to detach the kinesin motor domains from the MT, allowing it to move toward the MT minus-end. Once the generated force reaches a certain threshold, a third dynein could potentially bind to the DDB complex (**Fig. 5A**).

To test this hypothesis, we pre-assembled DDB complexes with two differently labeled dyneins (Alexa647 and TMR dyes) and introduced these dual-labeled DDB complexes—expected to display two different colors in 50% of cases—into the slide chamber with free dynein labeled with a third color (Alexa488). Using three-color, single-molecule total internal reflection fluorescence (smTIRF) microscopy, we observed that moving chimeras were frequently associated with three dyneins (**Fig. 5C**). In contrast, DDB complexes lacking the kinesin did not recruit a third dynein. The average velocity of ~30 nm/s (**Fig. 5D**) aligns with the movement of a DDB-3 complex under a load above ~8 pN (**Suppl. Fig. 6**), further supporting the role of tension in motor recruitment.

In conclusion, our findings demonstrate for the first time that mechanical tension applied to a DDB complex facilitates the recruitment of a third dynein. This load-induced adaptation leads to augmented force generation and increased velocities, enabling the DDB complex to meet escalating mechanical demands.

DDB takes largely load-independent steps of ~8 nm along MTs

Cryo-EM studies suggest that the motor domains of the two dyneins engaged within DDB complexes adopt a relatively compact and parallel arrangement[41], which would predict small center-of-mass steps along MTs. In contrast, previous single-molecule fluorescence studies have reported highly variable step sizes for DDB complexes in the absence of load, ranging from 4 to 48 nm[54,59]. Optical trapping experiments further revealed that the average forward step size of DDB complexes decreases with increasing load, from ~15 nm (step sizes ranging from 4 to 32 nm) at a 0.4 pN load to ~10 nm (step sizes ranging from 2 to 28 nm) at a 3.6 pN load[59]. Given the

structural arrangement of the dynein motor domains[41], steps sizes as large 48 nm would require substantial rearrangements of the motor domains, making such displacements unlikely.

In our experiments, DDB complexes bound to a single dynein predominantly advanced in ~8-nm center-of-mass steps under loads exceeding ~1 pN, even after accounting for the complex's compliance (**Suppl. Fig. 7**). At lower loads, consecutive forward steps occasionally extended to ~16 nm (**Fig. 6B**, right top, red). For DDB complexes associated with two dyneins, the step sizes remained largely consistent at ~8 nm under loads above ~2 pN, with occasional steps reaching up to ~24 nm under low-load conditions (**Fig. 6B**, right middle, red). These findings align with previous step-size measurements in the absence of load[54,59]. In contrast, when DDB complexes were associated with three dyneins, the step sizes were smaller than those of DDB-1 and DDB-2 (**Fig. 6B**, right bottom, red), suggesting that the third dynein may sterically hinder the movements of the two leading motors, thereby reducing the center-of-mass displacements.

For consecutive backward steps (**Fig. 6B**, right, blue), DDB-1's exhibited a backward stepping behavior similar to its forward stepping (**Fig. 6B**, right top, blue). In contrast, DDB-2 displayed larger backward step sizes under loads below 4 pN (**Fig. 6B**, right middle, blue). This observation suggests that while DDB-1's backward step size is constrained by dimerization, the two dynein dimers of DDB-2 can separate by greater distance at low load. Such large backward movements may facilitate synchronization between the dynein dimers if the leading dimer advances too far ahead. Conversely, DDB-3 showed almost no backward stepping under low-load conditions (**Fig. 6B**, right bottom, blue), likely because, in this configuration, at least two dynein dimers remain bound to the MT at a given time, minimizing backward stepping.

These findings suggest that dynein motor domains within DDB and DDR complexes[67] maintain a compact arrangement under load, with some degree of splaying occurring only under

low-load conditions. Additionally, dwell times between steps were largely unaffected by sub-stall loads but increased significantly for both forward and backward movements as the motors approached their respective stall forces (**Fig. 6B**, left).

We identified two distinct stepping modes in DDB complexes: consecutive stepping (**Fig. 6A**) and oscillating forward-backward stepping (**Fig. 6C**). Both modes were observed across all DDB complexes. While the step sizes in the two modes are comparable (**Fig. 6B & D**, right), their dwell time profiles differ significantly (**Fig. 6B & D**, left). In the oscillating stepping mode, a backward step is often immediately followed by a forward step (**Fig. 6C**), with shorter dwell times preceding the steps and reduced force dependency compared to consecutive stepping. We propose that consecutive stepping reflects a hand-over-hand mechanism for DDB, similar to kinesin-1[73]. In contrast, oscillatory stepping likely represents an inchworm mechanism, where the leading motor domain detaches and rapidly rebinds. This suggest that, unlike kinesin-1, the stepping order of DDB's motor domains is more stochastic with similarities to the stepping of *S. cerevisiae* dynein[66,74-76].

Discussion

Our study reveals a remarkable mechanosensitive property of DDB motor complexes: under increasing load, additional dyneins can be recruited, enhancing both force production and processivity. By integrating recombinant protein expression, structural analyses, optical trapping, and multi-color single-molecule fluorescence imaging, we demonstrate how mechanical tension triggers the higher-order assembly of DDB complexes. These findings expand our understanding of how dynein adapts to the mechanical demands of intracellular transport and unveil novel mechanisms of load-dependent motor recruitment and coordination.

Dynein Force Generation Is Limited by Auto-Inhibition

For nearly two decades, optical trapping experiments have indicated that mammalian dynein, when isolated from cofactors, produces forces of ~1 pN or less when measured with a weak optical trap (stiffness ~0.01 pN/nm)[39,52,77]. Forces of ~7–8 pN have been reported[78,79], but the basis for these outliers remain unclear. At higher trap stiffness (0.1 pN/nm) —conditions where dynein's weak processivity has less impact—its force increases to ~2 pN[39], consistent with a link between its weak processivity and limited force generation[39]. By contrast, *S. cerevisiae* dynein exhibits high processivity and a trap-stiffness-independent stall force of 4.5 pN[39].

Our data reveal that mammalian DDB complexes with a single dynein (DDB-1) generate ~2.5 pN when the dynein is not fully activated (at a trap stiffness of 0.06 pN/nm), closely matching the ~2 pN stall force observed for isolated dynein[39,58]. This limited force likely reflects dynein's propensity to adopt an auto-inhibitory "phi" conformation, which reduces motor engagement with the MT. Notably, DDB-1 complexes carrying mutations that prevent auto-inhibition generate

exclusively ~4.5 pN forces. These findings indicate that auto-inhibition, rather than the absence of cofactors, is the primary constraint on dynein's force output.

Lis1 acts as a critical regulator by counteracting auto-inhibition and stabilizing dynein in an active state. In the presence of Lis1, the occurrence of ~2.5 pN stalls drops significantly, and DDB-1 complexes predominantly generate ~4.5 pN forces. Interestingly, higher trap stiffness also promotes this transition, further supporting the role of load in activating dynein[39]. Additionally, D(phi)DB complexes, which are incapable of auto-inhibition[48], predominantly form two-dynein assemblies (DDB-2) that stall at ~7 pN, even under sub-molar assembly conditions. This suggests that the open conformation of dynein aligns favorably with an engaged dynein, facilitating the recruitment of additional motors. Interactions between the leading dynein's LIC and the trailing dynein's motor domain[68] (**Fig. 1B**) may further stabilize multi-dynein assemblies. Collectively, our findings underscore how Lis1 and dynein recruitment overcome auto-inhibition to enhance force generation and processivity.

Tension-Driven Dynein Recruitment

We demonstrate that DDB complexes respond to increasing mechanical load by recruiting a third dynein, raising their stall force to ~9 pN. This tension-induced recruitment depends on the auxiliary BicD adaptor, which interacts with the LIC of the third dynein. Mutation of the LIC-binding region in the auxiliary BicD (BicD Y46D) significantly reduces third-dynein recruitment, highlighting the importance of this interaction in higher-order DDB assembly.

Structural studies using Cryo-EM[80] and cryo-ET[41] have documented DDB assemblies containing one BicD (BICD2) and one dynein, with occasional recruitment of a second dynein under specific conditions[41,42]. Our findings emphasize the pivotal role of mechanical tension in

facilitating the recruitment of a third dynein. This enhanced force generation is likely necessary for overcoming cytoplasmic resistance during demanding tasks such as organelle and nuclear transport[81]. Supporting this model, we observed that DDB-2 complexes recruit a third dynein when opposing a MT-bound rigor kinesin, but not in its absence. The measured velocity of ~30 nm/s for kinesin-DDB-3 complexes matches the velocity of DDB-3 under loads exceeding 8 pN (**Suppl. Fig. 6**). While the precise loading rate at which DDB-3 complexes enhance the force on the kinesin-MT bond remains unclear, the measured velocity suggests that the DDB-3 complex generates sufficient force (~9 pN) to unbind kinesin from the MT[82]. Furthermore, the constant velocity of the kinesin-DDB-3 complex implies rapid re-binding of kinesin to the MT as the DDB-3 complex continues moving (**Fig. 5C**). These findings strongly support a model in which the DDB complex adapts to increasing loads by recruiting a third dynein—a mechanism consistent with *in vivo* observations of tension-induced dynein recruitment to lipid droplets[83].

Earlier studies have also shown that Lis1 enhances DDB's ability to compete with kinesin-1 under load[54]. Without Lis1, only ~20% of DDB complexes contain two dyneins, resulting in a stall force of ~4.1 pN, which is insufficient to counter kinesin-1's stall force of ~5–6 pN[39,84-86]. Lis1 increases the proportion of two-dynein DDB complexes to ~40%, raising the stall force to 5.4 pN[54]. Under these conditions, ~20% of kinesin-DDB complexes exhibit dynein-directed movement, likely due to the recruitment of a third dynein[54]. These findings highlight Lis1's critical role in enhancing dynein function under load by promoting high-order assembly

A recent study on KIF1C-DDB motor complexes reported a 50% increase in DDB processivity when KIF1C was tethered to the complex[87]. Given that KIF1C can generate forces up to ~7 pN[88], our observations that DDB-2 complexes stall at ~7 pN suggests that recruitment of a third dynein, increasing the stall force to ~9 pN, is required to overcome KIF1C. Together, these

results underscore how mechanical tension and Lis1-mediated activation drive the assembly of higher-order dynein complexes, enabling robust force generation and processivity under challenging conditions.

Dyneins Share Load Unequally Through a Staggered Arrangement

Our results suggest that dyneins within DDB complexes do not share load uniformly. The leading dynein typically slows near its stall force (~4.5 pN), triggering the engagement of additional motors and rapid increases in force to ~7 or ~9 pN. This behavior implies that dyneins are often engaged sequentially as load increases (**Fig. 2G**).

Although cryo-EM tomography shows all four dynein motor domains arranged in parallel[41], the angled alignment of the dynactin shoulder (~40° relative to the MT axis) suggests that dyneins bind MTs at offset positions in physiologically relevant conditions[41,42]. Such a "staggered" arrangement would enforce unequal load sharing, consistent with our observed force increments of ~2.5 pN for additional dyneins. This model is further supported by DNA-origami scaffolds designed to bind two or three DDB-1 complexes in a staggered configuration (with the DNA scaffold aligned to the long MT axis), yielding stall forces of ~7 pN and ~9 pN, respectively[59]. Thus, while multiple dyneins contribute to force generation, load sharing among them is inherently unequal.

DDB Primarily Moves in 8-nm Steps That Slow New Stall

We found that DDB complexes predominantly advance with ~8-nm center-of-mass steps under loads exceeding 2 pN. This step size remains largely load-independent for DDB-1 and DDB-2, while DDB-3 exhibits smaller forward steps (~4 nm) as stall forces are approached. These

findings suggest a "forward inching" mechanism in which dyneins incrementally shift their center of mass to sustain motion under high loads.

Previous studies reported larger average step sizes (~10 nm) and highly variable displacements (up to 32 nm)[59]. However, such results may stem from step-detection algorithms that overfit data, inflating reported step sizes while missing smaller increments. Additionally, the use of axonemes rather than MTs in earlier studies may have influenced stepping behavior.

By employing an unbiased, automated step-detection algorithm, we accurately measured dynein steps and uncovered a conserved pattern of ~8-nm displacements under physiological loads. This stepping behavior, reminiscent of kinesin-1[89], provides new insights into dynein's load-adaptive stepping and coordination mechanisms.

Conclusion

In summary, we demonstrate that DDB complexes adapt to increasing mechanical tension by recruiting additional dyneins, thereby boosting force output and processivity. Our findings reveal a novel tension-driven assembly mechanism mediated by the interaction between the auxiliary BicD and dynein's LIC, linking structural and functional regulation to mechanical adaptation. These insights advance our understanding of intracellular transport and highlight potential therapeutic targets for disorders involving impaired dynein function. Future studies should explore how these mechanisms operate *in vivo* and contribute to cellular homeostasis under diverse mechanical challenges.

Materials and Methods

Plasmids

The plasmids for human dynein complexes were gifts from the laboratory of Dr. Andrew P. Carter (Medical Research Council (MRC) Laboratory of Molecular Biology, Cambridge, UK). The plasmids for Lis1 constructs were gifts from the laboratory of Dr. Samara Reck-Peterson (University of California, San Diego, CA). BicD2 constructs were generated based on a mouse BicD2 plasmid (Addgene, #64205[1]) and a backbone for *E. coli* expression (pSNAP-tag(T7)-2 vector (New England Biolabs, #N9181S))[2] or pET28b. The shorter BicD2 constructs were generated via Q5 mutagenesis (New England Biolabs, # E0554S). A single mutation Y46D was introduced into a BicD2(25-180) construct. The kinesin (Cys-light) S43C construct[3] was a gift from the laboratory of Dr. Ronald D. Vale (University of California, San Francisco, CA). A single mutation T92N was introduced in the construct to generate the rigor kinesin, and the SpyTag003 was inserted at the C-terminus of the kinesin via Q5 mutagenesis. SpyCatcher003 tag was amplified from SpyCatcher003-S49C plasmid[4] (Addgene, #133448) and inserted to the C-terminus of BicD2. Plasmids for Tev protease[5] (Addgene, #171782) and BirA (Addgene, #20857)[6] were received from Addgene. The plasmids generated for this study were confirmed by either Sanger sequencing (Genomics Core Facility, Albert Einstein College of Medicine, Bronx, NY) or full plasmid sequencing (Azenta). The constructs used in this study is list in Suppl. Table 1.

Tagging dynactin in HEK 293T/17 SF cells using CRISPR

A HaloTag-3×FLAG-P2A-mCherry tag was inserted at the C-terminus of DCTN4 (p62) subunit of dynactin using CRISPR based on a previous published method[7]. The HaloTag-3×FLAG has

been used successfully in a stable adherent HEK293 cell line for dynactin purification[8]. DCTN4 gene with 1 kb downstream was retrieved from UCSC Genome Browser (https://genome.ucsc.edu/), and the guide RNA was designed using CRISPOR online tool (https://crispor.gi.ucsc.edu/)[9]. The guide RNA (5'-TCCTTAAAAGGTTCCACTGG) was ordered as Alt-R® crRNA from IDTDNA. Alt-R® tracrRNA (IDTDNA, #1072532) and Alt-R® Cas9 nuclease (IDTDNA, #1081058) were also ordered from IDTDNA. The RNAs were dissolved in IDTDNA nuclease-free duplex buffer to final 100 µM. 303 bp before the stop codon of DCTN4 was amplified from HEK 293T/17 SF and stitched with HaloTag-P2A-mCherry, followed by 611 bp of DCTN4 downstream sequences. The DNA was then inserted into a backbone plasmid, with its sequence confirmed by full plasmid sequencing (Azenta).

1 mL of HEK 293T/17 SF (ATCC, #ACS-4500) was recovered following vendor's instruction in 10 mL complete medium (9.8 mL of BalanCD HEK293 (Fujifilm Irvine Scientific, #91165-1L), 0.1 mL of 100×GlutaMAX™ supplement (Gibco, #35050061), 0.1 mL of 100×ITS (Insulin-Transferrin-Selenium) (Corning, #25-800-CR), and 20 µL of anti-clumping agent (Gibco, #0010057DG)). The culture was shaken at 37°C and 125 rpm with 5% CO_2 (Eppendorf, New Brunswick S41i CO2 Incubator Shaker) and maintained at a density of $< 2×10^6$ cells/mL.

3 µL of crRNA, 3 µL of tracrRNA, and 4 µL of Neon™ NxT resuspension R buffer (R buffer) were mixed and the RNAs were annealed by being heated at 95°C for 5 min and cooled to room temperature (r.t.) gradually. 1.5 µL of the 10 mg/mL Cas9 was added to the RNA and incubated at r.t. for 15 min. Afterwards, 6 µg of the plasmid described previously were added into the mixture. $5×10^6$ cells were pelleted at 140×g for 5 min, and washed once with 10 mL PBS. The

cells were then resuspended in 100 µL of R buffer and added to the prepared Cas9-RNA-plasmid solution. The cells were electroporated at 1100 V for 20 ms with 2 pulses using the Neon™ NxT Electroporation System (ThermoFisher, # NEON1S), as recommended by the vendor's instruction. The cells were dispersed into 2 mL media in a 6-well plate, and recovered for 2 days at 37ºC without shaking. After recovering and growth, 10^7 cells were sorted via BD Aria cell sorter using mCherry. The sorted cells were then recovered, grown, frozen, and stored in liquid nitrogen.

Dynactin purification

800 mL of HEK 293T/17 SF with p62 tagged with HaloTag-3×FLAG ($2×10^6$ cells/mL) were harvested via centrifugation at 600×g for 10 min at 4°C. The pellet (~10 mL) was resuspended in 10 mL of 2×dynactin lysis buffer (DLB) (60 mM HEPES, pH 7.2, 200 mM NaCl, 4 mM $MgCl_2$, 2 mM EGTA, 20% glycerol, 0.2 mM ATP, 2 mM DTT, 0.4% (v/v) Triton X-100, 2 EDTA-free protease inhibitor cocktail tablets (Roche, #11836170001)) and nutated at 4°C for 15 min. The lysate was cleared via centrifugation at 80,000 rpm (260,000×g, k-factor=28) for 10 minutes in an TLA-110 rotor using a Beckman Tabletop Optima TLX Ultracentrifuge.

1 mL of anti-FLAG® M2 affinity gel (Sigma, #A2220) was washed with 5 mL of 1×DLB buffer, and the cleared lysate was added to the resin and nutated overnight at 4ºC. The resin was washed with 40 mL of wash buffer (30 mM HEPES, pH 7.2, 250 mM KCl, 2 mM $MgCl_2$, 1 mM EGTA, 10% glycerol, 0.1 mM ATP, 1 mM DTT, 0.2% (w/v) Pluronic F-127, 0.5 mM Pefabloc). The resin was then transferred to a 2-mL tube with 200 µL of wash buffer and 100 µL of 5 mg/ml FLAG peptide, and nutated at 4ºC for 30 min. The supernatant was collected and concentrated in an Amicon Ultra-0.5 mL centrifugal filter (100-kDa MWCO) at 5000×g at 4ºC. The concentration of

dynactin was determined via Bradford assay (ThermoFisher, #23200). The subunits of dynactin were confirmed by mass spectrometry.

Dynactin from porcine brain

Porcine brain dynactin was a generous gift from Dr. Andrew P. Carter (Medical Research Council (MRC) Laboratory of Molecular Biology, Cambridge, UK).

Sf9 expression of cytoplasmic dynein-1 and Lis1

The full cytoplasmic dynein-1 complex, the tail complex, the phi-mutant complex, and Lis1 were expressed in Sf9 cells as previously published papers[8,10]. Briefly, a bacmid was generated by transforming the plasmid of the desired construct into MAX Efficiency™ DH10Bac competent cells (Gibco, # 10361012), and selected via blue-white screening. The insertion was further confirmed by PCR.

1 mL of Sf9 cells in Sf-900™ II SFM (ThermoFisher, #11496015) was recovered and grown in 10 mL of Sf-900™ II SFM medium (ThermoFisher, #10902104) at 27ºC with shaking at 135 rpm. The culture was maintained at a density of $< 2 \times 10^6$ cells/mL. To transfect cells, 2 mL of 0.5×10^6 cells/mL culture was seeded in a 6-well plate. 2 µg of bacmid was added to 200 µL of medium, and then 6 µL of FuGENE® HD transfection reagent (Promega, #E2311) was added to the mixture. After the mixture was incubated at r.t. for 15 min, it was added to the cells dropwise. The culture was then incubated at 27ºC without shaking for 4 days to generate P1 virus. Afterwards, the P1 virus was transferred to a 15-mL tube and stored at 4ºC in the dark. To generate P2 virus, 0.5 mL of the P1 virus was added to 50 mL of 1.5×10^6 cells/mL culture, and shaken at 27ºC at 135 rpm

for 3 days. The P2 virus was harvested by centrifugation at 2000×g for 10 min at 4ºC and stored at 4ºC in the dark. To express protein, 5 mL of the P2 virus was added to 500 mL of $2×10^6$ cells/mL culture, which was shaken at 27ºC at 135 rpm for 60–72 h. The cells were harvested by centrifugation at 2000×g for 15 min at 4ºC. The pellet was washed with 40 mL of cold PBS and centrifuged at 2000×g for 10 min at 4ºC. The supernatant was discarded and the pellet was flash frozen in liquid nitrogen and stored at –80ºC until further usage.

Cytoplasmic dynein-1 and Lis1 purification

The pellet (~5 mL) was resuspended in 5 mL of 2×lysis buffer (60 mM HEPES, pH 7.2, 300 mM KCl, 4 mM $MgCl_2$, 2 mM EGTA, 20% (v/v) glycerol, 0.4 mM ATP, 4 mM DTT, 0.2% (w/v) Pluronic F-127, 100 µL of 2.5 U/µL Dnase I, 2 EDTA-free protease inhibitor cocktail tablets), and dounced for 30 strokes on ice. The lysate was cleared as for dynactin purification. 4 mL of IgG Sepharose 6 Fast Flow affinity resin (Cytiva, #17096901) was washed with 6 mL of lysis buffer, and cleared lysate was added to the resin to nutate at 4ºC for 4 h. The resin was then washed with 50 mL of wash buffer (50 mM HEPES, pH 7.2, 150 mM KCl, 2 mM $MgCl_2$, 1 mM EGTA, 10% (v/v) glycerol, 0.1 mM ATP, 1 mM DTT, 0.1% (w/v) Pluronic F-127, 1 mM PMSF). To label the SNAP-tag on the N-terminus of dynein heavy chain, 15 µL of 1 mM SNAP-tag dye was added to the resin which was nutated at 4ºC overnight. The resin was washed with 50 mL of tev-release buffer (50 mM HEPES, pH 7.2, 150 mM KCl, 2 mM $MgCl_2$, 1 mM EGTA, 10% (v/v) glycerol, 0.1 mM ATP, 1 mM DTT, 0.1% (w/v) Pluronic F-127), and then transferred to a 5-mL tube with tev-release buffer to a final volume of 5 mL. 50 µL of 200 µM MBP-superTevProtease was added to the resin, and the resin was nutated at 4ºC overnight in the dark. 1 mL of amylose resin (New England Biolabs, #E8021S) was added to remove the MBP-superTevProtease. The supernatant

was then concentrated as for dynactin purification. The purity and concentration of the complex was determined by 4-12% bis-tris polyacrylamide gel and Bradford assay, respectively.

E. coli-based expression

BicD2 constructs were expressed in *E. coli*. Each plasmid was transformed into BL21-CodonPlus(DE3)-RIPL competent cells (Agilent Technologies, #230280), and a single colony was picked and inoculated in 1 mL of terrific broth (TB) with 50 µg/mL carbenicillin and 50 µg/mL chloramphenicol. The 1-mL culture was shaken at 37°C overnight, and then inoculated into 400 mL of TB with 2 µg/mL carbenicillin and 2 µg/mL chloramphenicol. The culture was shaken at 37°C for 5 hours and cooled to 16ºC for 1 hour. Protein expression was induced by 0.1 mM IPTG overnight at 16ºC. The cells were harvested by centrifugation at 3,000×g for 10 minutes at 4°C. The supernatant was discarded, and 5 mL of B-PER™ Complete Bacterial Protein Extraction Reagent (ThermoFisher Scientific, #89821) supplemented with 2 mM $MgCl_2$, 1 mM EGTA, 1 mM DTT, 0.1 mM ATP, and 2 mM PMSF was added to fully resuspended the pellet. The resuspension was flash frozen in liquid nitrogen and stored at –80 °C.

E. coli-expressed construct purification

To purify *E. coli* expressed protein, the frozen cell pellet was thawed at 37 °C and nutated at r.t. for 20 minutes to lyse the cells. The cell lysate was cleared as for dynactin purification. The supernatant was flown through 500 µL of Ni-NTA Roche cOmplete™ His-Tag purification resin (Millipore Sigma, #5893682001) for His-tag tagged proteins, or Strep-Tactin® 4Flow® high capacity resin (IBA Lifesciences GmbH, #2-1250-010) for strep-II tagged proteins. The resin was washed with 10 mL of wash buffer (50 mM HEPES, pH 7.2, 300 mM KCl, 2 mM $MgCl_2$, 1 mM

EGTA, 1 mM DTT, 1 mM PMSF, 0.1 mM ATP, 0.1% (w/v) Pluronic F-127, 10% glycerol), and then eluted with elution buffer (50 mM HEPES, pH7.2, 150 mM KCl, 2 mM MgCl$_2$, 1 mM EGTA, 1 mM DTT, 1 mM PMSF, 0.1 mM ATP, 0.1% Pluronic F-127 (w/v), 10% glycerol, pH 7.2; for His-tagged protein: 150 mM imidazole; for strep-II tagged protein: 5 mM desthiobiotin). The elute was either flash frozen and stored at –80 °C, or concentrated using an Amicon Ultra-0.5 mL centrifugal filter unit (30-kDa MWCO) (Sigma, #UFC503024) before flash freezing. The purity and concentration of the proteins was determined on 4-12% bis-tris polyacrylamide gels. To label BicD2 with biotin, 50 µL of 10 µM BicD2 mixed with 1 µL of 100 mM ATP, 2 µL of 50 µM BirA, and 1 µL of 10 mM biotin was incubated at 30ºC for 2 h, and then BirA was removed by Ni-NTA resin. The solution was then buffer exchanged to a buffer (30 mM HEPES, pH 7.2, 50 mM KCl, 2 mM MgCl$_2$, 1 mM EGTA, 1 mM DTT, 0.1% (w/v) Pluronic F-127, 10% glycerol) to remove free biotin.

BicD2-kinesin constructs purification

The purification of BicD2-kinesin followed the same protocol as the purification for *E. coli*-expressed constructs, except after the cells were thawed, the BicD2 and kinesin solutions were combined to nutate at r.t. for 20 min, and then further nutated at 4ºC for 2 h to allow the binding between SpyCatcher003 and SpyTag003. Afterwards, the lysate was first flown through the Ni-NTA resin, and then the elution from the Ni-NTA was flown through the Strep-Tactin resin to capture the full-formed complex. The elute from the Strep-Tactin resin was further concentrated as in the purification for dynactin.

Optical tweezers assay

Polystyrene beads

Anti-GFP polystyrene beads were generated by covalently binding antiGFP antibody and BSA to carboxyl polystyrene beads (0.52 µm, Polysciences, #09836-15) via NHS-EDAC chemistry[11]. Streptavidin beads (0.55 µm) were ordered from Spherotech (#SVP-05-10).

Microtubule polymerization

2 µL of 10 mg/mL tubulin (Cytoskeleton, #T240-B) was mixed with 2 µL of 1 mg/mL biotinylated tubulin (Cytoskeleton, # T333P-A) and 1 µL of 10 mM GTP. The mixture was incubated at 37ºC for 20 min, and subsequently 0.5 µL of 0.2 mM paclitaxel in DMSO was added to the mixture and the incubation was continued for another 20 min. The solution was then carefully laid on top of 100 µL of glycerol cushion (80 mM PIPES, pH 6.8, 2 mM $MgCl_2$, 1 mM EGTA, 60% (v/v) glycerol, 1 mM DTT, 10 µM paclitaxel) in a 230-µL TLA100 tube (Beckman Coulter, #343775), and centrifuged at 80,000×rpm (250,000×g, *k*-factor=10) for 5 min at r.t.. The supernatant was carefully removed, and the pellet was resuspended in 11 µL of BRB80G10 (80 mM PIPES, pH 6.8, 2 mM $MgCl_2$, 1 mM EGTA, 10% (v/v) glycerol, 1 mM DTT, 10 µM paclitaxel). The microtubule solution was stored at r.t. in the dark for further usage.

Flow chamber preparation

A flow chamber was assembled using a glass slide (Fisher Scientific, #12-550-123) and an ethanol-cleaned coverslip (Zeiss, #474030-9000-000) with two thin strips of parafilm. 10 µL of 0.5 mg/ml BSA-biotin (ThermoScientific, #29130) was flown into the chamber with 10 min incubation. The chamber was then washed with 2×20 µL blocking buffer (80 mM PIPES, pH 6.8, 2 mM $MgCl_2$, 1 mM EGTA, 10 µM paclitaxel, 1% (w/v) Pluronic F-127, 2 mg/mL BSA, 1 mg/mL α-casein) and

incubated for 30 min to block the surface. Afterwards, 10 µL of 0.25 mg/ml streptavidin (Promega, #Z7041) was introduced into the chamber with 10 min incubation. The chamber was washed with 2×20 µL blocking buffer, and 10 µL of 0.02 mg/ml biotin-labeled MTs in the blocking buffer was flown into the chamber with 1 min incubation. The chamber was washed with 2×20 µL blocking buffer, and kept in a humid chamber until further usage.

Sample preparation

Dynein (dimer), dynactin, and BicD2 (dimer) were mixed in a 1:1:1 ratio to a final of 200 nM each. The solution was incubated at 4ºC overnight. 1 µL of the beads was mixed with 1 µL of motility buffer (60 mM HEPES, pH 7.2, 50 mM KCl, 2 mM MgCl$_2$, 1 mM EGTA, 10 µM paclitaxel, 0.5% (w/v) Pluronic F-127, 5 mg/mL BSA, 0.2 mg/mL α-casein) and 1 µL of appropriately diluted DDB complex, and incubated on ice for 30 min. Afterwards, 40 µL of motility buffer (motility buffer with 2 mM ATP, 2 mM biotin, gloxy oxygen scavenger system) was added to the beads-protein solution, and 2×20 µL of the solution was flown into the chamber. The chamber was sealed with vacuum grease.

Data acquisition

Optical tweezers experiments were performed using a C-trap/TIRF/IRM (C-trap combined with Total Internal Reflection Fluorescence (TIRF) and Interference Reflection Microscopy (IRM)) (Lumicks) at r.t.. Microtubules were visualized with IRM. Trap stiffness was adjusted to ~ 0.08–0.1 pN/nm. The optical trapping data were initially sampled at a rate of 20 MHz and then digitally down-sampled by a factor of 256, resulting in a final sample rate of 78.125 kHz. An anti-aliasing

filter was applied to achieve a passband of 31.25 kHz, and the resulting data were fitted to the Lorentzian power spectrum.

Data analysis

The data was processed using a home-built MATLAB program. A MATLAB built-in step finding algorithm was used to find steps in traces which were down-sampled by a factor of 80 via averaging. Steps with forces < 0.2 pN was discarded. For each step, the combined stiffness of the motor and bead-linkage, $k_{m\text{-}link}$, is calculated using the original data:

$$k_{m-link} = \frac{k_B T}{\sigma^2(x_b)} - k_{trap}$$

The steps for each trace were binned based on 0.5-pN force window, and the mean value of the 25^{th} to 75^{th} percentiles for each force window, such as dwell time, step sizes and velocity, was calculated as the representation in this force window. Traces with similar forces were grouped, and the mean and standard error of the mean were calculated and visualized using Prism 10 (GraphPad). To compensate for the link compliance, the difference between the motor stretch ($F/k_{m\text{-}link}$) for each force window in each group was calculated and added to the steps.

Total internal reflection fluorescence (TIRF) assay

Sample preparation

Microtubule polymerization and flow chamber preparation are the same as for optical tweezers assay. After the microtubule was bound on the surface, the assembled complex was diluted in the motility buffer (60 mM HEPES, pH 7.2, 75 mM KCl, 2 mM MgCl$_2$, 1 mM EGTA, 1 mM DTT,

100 µM ATP or 2 mM ATP, 0.5% Pluronic F-127 (w/v), 10% glycerol). For three-color experiment, 10 nM dynein labeled with Alexa488 was added to the motility buffer.

Data acquisition

The images were acquired using BioVis software (BioVision Technologies), with 500 ms acquisition time.

Data analysis

The kymographs were generated using Fiji[12], and the velocity was analyzed using a home-built MATLAB program.

AlphaFold2 prediction

The interaction between mouse BicD2(25-180) and human LIC2(417-444) was predicted using ColabFold (v1.5.5: AlphaFold2 using MMseqs2)[13].

Graphic visualization

The molecular structures in the figures were visualized using UCSF ChimeraX[14]. The model graphics in Figures 2 and 7 were generated via BioRender (https://www.biorender.com/).

Acknowledgments

We acknowledge Albert Einstein College of Medicine Flow Cytometry Core Facility (Einstein National Cancer Institute's Cancer Center support grant P30CA013330), especially Ming Liu, for help sorting the HEK293 cells, and Jan O. Wirth (Abberior) for discussions on step-finding algorithms. L. Rao and A. Gennerich were supported by National Institutes of Health grants R01GM098469 and would like to acknowledge the use of LUMICKS C-Trap, which was funded by the NIH grant S10OD034445-01. R. McKenney was supported by NIGMS National Institutes of Health grant GM124889. M. Arnold and K. Stengel were supported by National Institutes of Health grant R35GM147213. S. Sidoli gratefully acknowledges the funding from the Hevolution Foundation (AFAR), the Einstein-Mount Sinai Diabetes Center, and the NIH Office of the Director (S10OD030286). Molecular graphics and analyses performed with UCSF ChimeraX, developed at the University of California, San Francisco, was supported by National Institutes of Health R01-GM129325 and the Office of Cyber Infrastructure and Computational Biology, National Institute of Allergy and Infectious Diseases.

Author Contributions

L. Rao produced and purified all proteins; L. Rao and A. Gennerich designed the research; L. Rao and X. Liu expressed and purified proteins; L. Rao, F. Berger, and A. Gennerich analyzed the experimental data; L. Rao and A. Gennerich wrote the manuscript. A. Gennerich secured funding; R. McKenney generated and provided proteins purified from rat brain lysate for initial studies. M. Arnold and K. Stengel assisted in CRISPR-based dynactin construct generation; S. Sidoli performed the mass spectrometry assay for dynactin.

Competing Interests

The authors declare no further competing financial interests.

Data availability

Data supporting the findings of this manuscript are available from the corresponding author upon

reasonable request.

References

1. Abe, T.K. et al. Dynactin is essential for growth cone advance. *Biochem Biophys Res Commun* **372**, 418-22 (2008).
2. Grabham, P.W., Seale, G.E., Bennecib, M., Goldberg, D.J. & Vallee, R.B. Cytoplasmic dynein and LIS1 are required for microtubule advance during growth cone remodeling and fast axonal outgrowth. *J Neurosci* **27**, 5823-34 (2007).
3. Kardon, J. & Vale, R. Regulators of the cytoplasmic dynein motor. *Nat Rev Mol Cell Biol* **10**, 854-865 (2009).
4. Tai, C.Y., Dujardin, D.L., Faulkner, N.E. & Vallee, R.B. Role of dynein, dynactin, and CLIP-170 interactions in LIS1 kinetochore function. *J Cell Biol* **156**, 959-68 (2002).
5. Vallee, R.B., Williams, J.C., Varma, D. & Barnhart, L.E. Dynein: An ancient motor protein involved in multiple modes of transport. *J Neurobiol* **58**, 189-200 (2004).
6. Yamada, M. et al. LIS1 and NDEL1 coordinate the plus-end-directed transport of cytoplasmic dynein. *EMBO J* **27**, 2471-2483 (2008).
7. Vallee, R.B., Tai, C. & Faulkner, N.E. LIS1: cellular function of a disease-causing gene. *Trends Cell Biol* **11**, 155-160 (2001).
8. Feng, Y. & Walsh, C.A. Protein-protein interactions, cytoskeletal regulation and neuronal migration. *Nat Rev Neurosci* **2**, 408-416 (2001).
9. Cahana, A. et al. Targeted mutagenesis of Lis1 disrupts cortical development and LIS1 homodimerization. *Proc Natl Acad Sci U S A* **98**, 6429-6434 (2001).
10. Feng, Y. & Walsh, C.A. Mitotic spindle regulation by Nde1 controls cerebral cortical size. *Neuron* **44**, 279-293 (2004).
11. Pawlisz, A.S. et al. Lis1-Nde1-dependent neuronal fate control determines cerebral cortical size and lamination. *Hum Mol Genet* **17**, 2441-2455 (2008).
12. Reiner, O. et al. Lissencephaly gene (LIS1) expression in the CNS suggests a role in neuronal migration. *J Neurosci* **15**, 3730-3738 (1995).
13. Tsai, J.W., Chen, Y., Kriegstein, A.R. & Vallee, R.B. LIS1 RNA interference blocks neural stem cell division, morphogenesis, and motility at multiple stages. *J Cell Biol* **170**, 935-945 (2005).
14. Mochida, G.H. Genetics and biology of microcephaly and lissencephaly. *Semin Pediatr Neurol* **16**, 120-126 (2009).
15. Leventer, R. Lissencephaly type I. in *Handbook of clinical neurology*, Vol. 87 (ed. Bruyn, P.J.V.a.G.W.) 205-218 (2008).
16. Kato, M. & Dobyns, W.B. Lissencephaly and the molecular basis of neuronal migration. *Hum Mol Genet* **12 Spec No 1**, R89-96 (2003).
17. Wynshaw-Boris, A. Lissencephaly and LIS1: insights into the molecular mechanisms of neuronal migration and development. in *Clin Genet* Vol. 72 296-304 (2007).
18. Tanaka, T. et al. Lis1 and doublecortin function with dynein to mediate coupling of the nucleus to the centrosome in neuronal migration. *J Cell Biol* **165**, 709-721 (2004).
19. King, S.J. & Schroer, T.A. Dynactin increases the processivity of the cytoplasmic dynein motor. *Nat Cell Biol* **2**, 20-24 (2000).
20. Ross, J.L., Wallace, K., Shuman, H., Goldman, Y.E. & Holzbaur, E.L.F. Processive bidirectional motion of dynein-dynactin complexes in vitro. *Nat Cell Biol* **8**, 562-570 (2006).
21. Schroer, T.A. Dynactin. *Annu Rev Cell Dev Biol* **20**, 759-779 (2004).

22. Nirschl, J.J., Magiera, M.M., Lazarus, J.E., Janke, C. & Holzbaur, E.L. α-Tubulin Tyrosination and CLIP-170 Phosphorylation Regulate the Initiation of Dynein-Driven Transport in Neurons. *Cell Rep* **14**, 2637-52 (2016).

23. McKenney, R.J., Huynh, W., Vale, R.D. & Sirajuddin, M. Tyrosination of alpha-tubulin controls the initiation of processive dynein-dynactin motility. *EMBO J* **35**, 1175-1185 (2016).

24. McKenney, R.J., Huynh, W., Tanenbaum, M.E., Bhabha, G. & Vale, R.D. Activation of cytoplasmic dynein motility by dynactin-cargo adapter complexes. *Science* **345**, 337-341 (2014).

25. Schlager, M.A., Hoang, H.T., Urnavicius, L., Bullock, S.L. & Carter, A.P. In vitro reconstitution of a highly processive recombinant human dynein complex. *EMBO J* **33**, 1855-1868 (2014).

26. Ayloo, S. et al. Dynactin functions as both a dynamic tether and brake during dynein-driven motility. *Nat Commun* **5**, 4807 (2014).

27. Kardon, J.R., Reck-Peterson, S.L. & Vale, R.D. Regulation of the processivity and intracellular localization of Saccharomyces cerevisiae dynein by dynactin. *Proc Natl Acad Sci USA* **106**, 5669-5674 (2009).

28. Tripathy, S.K. et al. Autoregulatory mechanism for dynactin control of processive and diffusive dynein transpor. *Nat Cell Biol* **in press**(2014).

29. Moughamian, A.J. & Holzbaur, E.L.F. Dynactin Is Required for Transport Initiation from the Distal Axon. *Neuron* **74**, 331-343 (2012).

30. Cianfrocco, M.A., DeSantis, M.E., Leschziner, A.E. & Reck-Peterson, S.L. Mechanism and regulation of cytoplasmic dynein. *Annu Rev Cell Dev Biol* **31**, 83-108 (2015).

31. Qiu, R., Zhang, J. & Xiang, X. p25 of the dynactin complex plays a dual role in cargo binding and dynactin regulation. *J Biol Chem* **293**, 15606-15619 (2018).

32. Munch, C. et al. Point mutations of the p150 subunit of dynactin (DCTN1) gene in ALS. *Neurology* **63**, 724-6 (2004).

33. Puls, I. et al. Mutant dynactin in motor neuron disease. *Nat Genet* **33**, 455-456 (2003).

34. Farrer, M.J. et al. DCTN1 mutations in Perry syndrome. *Nat Genet* **41**, 163-165 (2009).

35. Urnavicius, L. et al. The structure of the dynactin complex and its interaction with dynein. *Science* **347**, 1441-6 (2015).

36. Chowdhury, S., Ketcham, S.A., Schroer, T.A. & Lander, G.C. Structural organization of the dynein-dynactin complex bound to microtubules. *Nat Struct Mol Biol* **22**, 345-7 (2015).

37. Trokter, M., Mücke, N. & Surrey, T. Reconstitution of the human cytoplasmic dynein complex. *Proc Natl Acad Sci USA* **109**, 20895-20900 (2012).

38. Nicholas, M.P. et al. Control of cytoplasmic dynein force production and processivity by its C-terminal domain. *Nat Commun* **6**, 6206 (2015).

39. Brenner, S., Berger, F., Rao, L., Nicholas, M.P. & Gennerich, A. Force production of human cytoplasmic dynein is limited by its processivity. *Sci Adv* **6**, eaaz4295 (2020).

40. Olenick, M.A., Tokito, M., Boczkowska, M., Dominguez, R. & Holzbaur, E.L. Hook Adaptors Induce Unidirectional Processive Motility by Enhancing the Dynein-Dynactin Interaction. *J Biol Chem* **291**, 18239-51 (2016).

41. Grotjahn, D.A. et al. Cryo-electron tomography reveals that dynactin recruits a team of dyneins for processive motility. *Nat Struct Mol Biol* **25**, 203-207 (2018).

42. Urnavicius, L. et al. Cryo-EM shows how dynactin recruits two dyneins for faster movement. *Nature* **554**, 202-206 (2018).

43. Toropova, K. et al. Lis1 regulates dynein by sterically blocking its mechanochemical cycle. *Elife* **3**(2014).

44. Huang, J., Roberts, A.J., Leschziner, A.E. & Reck-Peterson, S.L. Lis1 Acts as a "Clutch" between the ATPase and Microtubule-Binding Domains of the Dynein Motor. *Cell* **150**, 975-986 (2012).

45. DeSantis, M.E. et al. Lis1 Has Two Opposing Modes of Regulating Cytoplasmic Dynein. *Cell* **170**, 1197-1208 e12 (2017).

46. Marzo, M.G., Griswold, J.M. & Markus, S.M. Pac1/LIS1 stabilizes an uninhibited conformation of dynein to coordinate its localization and activity. *Nat Cell Biol* **22**, 559-569 (2020).

47. Qiu, R., Zhang, J. & Xiang, X. LIS1 regulates cargo-adapter-mediated activation of dynein by overcoming its autoinhibition in vivo. *J Cell Biol* **218**, 3630-3646 (2019).

48. Zhang, K. et al. Cryo-EM Reveals How Human Cytoplasmic Dynein Is Auto-inhibited and Activated. *Cell* **169**, 1303-1314 (2017).

49. Amos, L.A. Brain dynein crossbridges microtubules into bundles. *J Cell Sci* **93**, 19-28 (1989).

50. Toropova, K., Mladenov, M. & Roberts, A.J. Intraflagellar transport dynein is autoinhibited by trapping of its mechanical and track-binding elements. *Nat Struct Mol Biol* **24**, 461-468 (2017).

51. Jordan, M.A., Diener, D.R., Stepanek, L. & Pigino, G. The cryo-EM structure of intraflagellar transport trains reveals how dynein is inactivated to ensure unidirectional anterograde movement in cilia. *Nat Cell Biol* **20**, 1250-1255 (2018).

52. Torisawa, T. et al. Autoinhibition and cooperative activation mechanisms of cytoplasmic dynein. *Nat Cell Biol* **16**, 1118-24 (2014).

53. Indigo C. Geohring et al. A nucleotide code governs Lis1's ability to relieve dynein autoinhibition. *bioRxiv* **2024.12.30.630615**(2024).

54. Elshenawy, M.M. et al. Lis1 activates dynein motility by modulating its pairing with dynactin. *Nat Cell Biol* **22**, 570-578 (2020).

55. Htet, Z.M. et al. LIS1 promotes the formation of activated cytoplasmic dynein-1 complexes. *Nat Cell Biol* **22**, 518-525 (2020).

56. Ton, W.D. et al. Microtubule-binding-induced allostery triggers LIS1 dissociation from dynein prior to cargo transport. *Nat Struct Mol Biol* (2023).

57. Nicastro, D. et al. Cryo-electron tomography reveals conserved features of doublet microtubules in flagella. *Proc Natl Acad Sci U S A* **108**, E845-53 (2011).

58. Belyy, V. et al. The mammalian dynein-dynactin complex is a strong opponent to kinesin in a tug-of-war competition. *Nat Cell Biol* **18**, 1018-24 (2016).

59. Elshenawy, M.M. et al. Cargo adaptors regulate stepping and force generation of mammalian dynein-dynactin. *Nat Chem Biol* **15**, 1093-1101 (2019).

60. Liu, X., Rao, L. & Gennerich, A. The regulatory function of the AAA4 ATPase domain of cytoplasmic dynein. *Nat Commun* **11**, 5952 (2020).

61. Khataee, H. & Howard, J. Force Generated by Two Kinesin Motors Depends on the Load Direction and Intermolecular Coupling. *Phys. Rev. Lett.* **122**, 188101 (2019).

62. Pyrpassopoulos, S., Shuman, H. & Ostap, E.M. Modulation of Kinesin's Load-Bearing Capacity by Force Geometry and the Microtubule Track. *Biophys J* **118**, 243-253 (2019).

63. Sedlak, S.M. et al. Monodisperse measurement of the biotin-streptavidin interaction strength in a well-defined pulling geometry. *PLoS One* **12**, e0188722 (2017).

64. Layden, H.M., Eleuteri, N.A., Hiebert, S.W. & Stengel, K.R. A protocol for rapid degradation of endogenous transcription factors in mammalian cells and identification of direct regulatory targets. *STAR Protoc* **2**, 100530 (2021).

65. Redwine, W.B. et al. The human cytoplasmic dynein interactome reveals novel activators of motility. *Elife* **6**(2017).

66. Reck-Peterson, S.L. et al. Single-molecule analysis of dynein processivity and stepping behavior. *Cell* **126**, 335-348 (2006).

67. Sami Chaaban & Carter, A.P. Structure of dynein–dynactin on microtubules shows tandem adaptor binding. *Nature* **610**, 212-216 (2023).

68. Chaaban, S. & Carter, A.P. Structure of dynein-dynactin on microtubules shows tandem adaptor binding. *Nature* **610**, 212-216 (2022).

69. Lee, I.G., Cason, S.E., Alqassim, S.S., Holzbaur, E.L.F. & Dominguez, R. A tunable LIC1-adaptor interaction modulates dynein activity in a cargo-specific manner. *Nat Commun* **11**, 5695 (2020).

70. Nakata, T. & Hirokawa, N. Point mutation of adenosine triphosphate-binding motif generated rigor kinesin that selectively blocks anterograde lysosome membrane transport. *J Cell Biol* **131**, 1039-1053 (1995).

71. Zakeri, B. et al. Peptide tag forming a rapid covalent bond to a protein, through engineering a bacterial adhesin. *Proc Natl Acad Sci U S A* **109**, E690-7 (2012).

72. Keeble, A.H. et al. Approaching infinite affinity through engineering of peptide-protein interaction. *Proc Natl Acad Sci U S A* **116**, 26523-26533 (2019).

73. Yildiz, A., Tomishige, M., Vale, R.D. & Selvin, P.R. Kinesin walks hand-over-hand. *Science* **303**, 676-8 (2004).

74. Gennerich, A., Carter, A.P., Reck-Peterson, S.L. & Vale, R.D. Force-induced bidirectional stepping of cytoplasmic dynein. *Cell* **131**, 952-965 (2007).

75. Qiu, W. et al. Dynein achieves processive motion using both stochastic and coordinated stepping. *Nat Struct Mol Biol* **19**, 193-200 (2012).

76. DeWitt, M.A., Chang, A.Y., Combs, P.A. & Yildiz, A. Cytoplasmic dynein moves through uncoordinated stepping of the AAA+ ring domains. *Science* **335**, 221-225 (2012).

77. Mallik, R., Carter, B.C., Lex, S.A., King, S.J. & Gross, S.P. Cytoplasmic dynein functions as a gear in response to load. *Nature* **427**, 649-52 (2004).

78. Toba, S., Watanabe, T.M., Yamaguchi-Okimoto, L., Toyoshima, Y.Y. & Higuchi, H. Overlapping hand-over-hand mechanism of single molecular motility of cytoplasmic dynein. *Proc Natl Acad Sci U S A* **103**, 5741-5 (2006).

79. Walter, W.J., Koonce, M.P., Brenner, B. & Steffen, W. Two independent switches regulate cytoplasmic dynein's processivity and directionality. in *Proc Natl Acad Sci USA* Vol. 109 5289-5293 (2012).

80. Urnavicius, L. et al. The structure of the dynactin complex and its interaction with dynein. *Science* **347**, 1441-1446 (2015).

81. Hu, D.J.-K. et al. Dynein Recruitment to Nuclear Pores Activates Apical Nuclear Migration and Mitotic Entry in Brain Progenitor Cells. *Cell* **154**, 1300-1313 (2013).

82. Uemura, S. et al. Kinesin-microtubule binding depends on both nucleotide state and loading direction. in *Proc Natl Acad Sci USA* Vol. 99 5977-5981 (2002).

83. Reddy, B.J. et al. Load-induced enhancement of Dynein force production by LIS1-NudE in vivo and in vitro. *Nat Commun* **7**, 12259 (2016).

84. Budaitis, B.G. et al. Pathogenic mutations in the kinesin-3 motor KIF1A diminish force generation and movement through allosteric mechanisms. *J Cell Biol* **220**, e202004227 (2021).

85. Svoboda, K., Schmidt, C.F., Schnapp, B.J. & Block, S.M. Direct observation of kinesin stepping by optical trapping interferometry. *Nature* **365**, 721-727 (1993).

86. Svoboda, K. & Block, S.M. Force and velocity measured for single kinesin molecules. *Cell* **77**, 773-784 (1994).

87. Ferdos Abid Ali et al. KIF1C activates and extends dynein movement through the FHF cargo adaptor. *bioRxiv* **2023.10.26.56424**(2024).

88. Siddiqui, N. et al. Force generation of KIF1C is impaired by pathogenic mutations. *Curr Biol* **32**, 3862-3870 e6 (2022).

89. Sudhakar, S. et al. Germanium nanospheres for ultraresolution picotensiometry of kinesin motors. *Science* **371**(2021).

90. Zhang, R., LaFrance, B. & Nogales, E. Separating the effects of nucleotide and EB binding on microtubule structure. *Proc Natl Acad Sci U S A* **115**, E6191-E6200 (2018).

91. Nishida, N. et al. Structural basis for two-way communication between dynein and microtubules. *Nat Commun* **11**, 1038 (2020).

92. Mirdita, M. et al. ColabFold: making protein folding accessible to all. *Nat Methods* **19**, 679-682 (2022).

Figures

Figure 1. (A) Structural representation of dynein-dynactin-BicDR (DDR) complex (PDB 7Z8F[67]). The dynein heavy chain (B1) from the DDR complex was aligned with microtubules (PDB 6DPV, EMDB 7974[90]), connected via a high-affinity dynein microtubule-binding domain structure bound to microtubules (PDB 6KIQ[91]). **(B)** Polyacrylamide gel analysis of purified human dynactin and dynein complexes. Protein bands were identified based on molecular weight, with the dynactin subunits further confirmed by mass spectrometry.

Figure 2. (A) Schematic of optical tweezers assay (not to scale). A ~0.5 µm-diameter bead, coated with either anti-GFP antibody (as shown) or streptavidin, is tethered to DDB complexes via an sfGFP or biotin tag fused to the C-terminus of BicD2. The bead is positioned on a surface-attached microtubule. When the DDB binds to and moves along the MT in the presence of ATP, it pulls the attached bead with it. The optical trap resists this motion, generating a measurable force. **(B)** Stall forces of DDB complexes bound to anti-GFP antibody-coated beads. **(C–G)** Examples of DDB's force generation.

Figure 3. (A) Force generation of human dynein in a stiff optical trap (adapted from[39]). **(B)** Example of force generation by DDB-1, showing a transition from slow movement toward ~2.5 pN to rapid movement toward ~5 pN. **(C)** Distribution of stall forces (~2.5, ~4.5, ~7 and ~9 pN) across different experiments: DDB-1(semi-inhibited), DDB-1 (fully activated), DDB-2, and DDB-3. The percentages represent the fraction of beads exhibiting each stall force, with sample sizes and distributions as follows: DDB: n=41; 17%, 27%, 46%, 10%. D(phi)DB: n=31, 0%, 13%, 68%, 19%. DDB+Lis1: n=29; 3%, 17%, 59%, 21%. **(D)** The measured fraction of force-generating beads. The bars represent the mean value with SD. DDB: n=4, 0.16 ± 0.05 (mean $\pm$ SD). D(phi)DB: n=4, 0.19 ± 0.10; DDB + Lis1: n=3, 0.30 ± 0.06.

Figure 4. (A) Structure of DDR complex with two dyneins (light blue and pink) and two BicDR dimers (blue and magenta) (PDB 7Z8F). Dynactin is not shown in the representation. The interaction between the C-terminal helix of the LICs of dynein A2 (light green) and B1 (green) with the N-terminal CC1-box of BicDR-A (blue) highlighted with the dashed red circle. The span of BicDR across the dynactin shoulder is indicated with corresponding colors. **(B)** (Top) ColabFold-predicted[92] structure of the interactions between mouse BicD2(25–180) and human dynein LIC(417–444). (Middle) A close-up of the predicted structure, showing Tyr46 of one BicD2 as sphere style (colored by elements). The hydrophobic surface of one LIC is depicted in light yellow (hydrophobic) and cyan (hydrophilic). (Bottom) Crystal structure of human BicD2(1–98) binding to human dynein LIC(433–458), displayed with hydrophobic surface (PDB 6PSE[69]). **(C)** Schematic representation of the BicD constructs. Y46D mutation in BicD2(25–180) is marked in red. **(D)** The moving fraction of the beads per experiment. The bars represent the mean value with SD. DDB: n=4, 0.16 ± 0.05 (mean ± SD). DDB+dyn: n=3, 0.13 ± 0.01. DDB+dyn+BicD: n=6, 0.12 ± 0.03. DDB+dyn+BicD(Y46D): n=4, 0.13 ± 0.04. DDB+dyn tail: n=4, 0.14 ± 0.05. **(E)** The percentage for each DDB complex. The percentages represent DDB-1(semi-inhibited), DDB-1 (fully activated), DDB-2, and DDB-3, respectively. DDB: n=41; 17%, 27%, 46%, 10%. DDB+dyn: n=14, 7%, 14%, 64%, 15%. DDB+dyn+BicD: n=26, 0%, 8%, 35%, 57%. DDB+dyn+BicD(Y46D): n=25, 16%, 32%, 40%, 12%. DDB+dyn tail: n=15, 13%, 67%, 20%, 0%. **(F)** Hypothetical structure of a third dynein dimer (gray, semi-transparent) recruited to a DDB complex. The tail was manually added to the DDR structure. **(G)** Full DDR complex, including a microtubule, shown for context.

Figure 5. (A) Schematic of a rigor kinesin-1 covalently attached to BicD in the DDB complex via SpyTag-SpyCatcher system. **(B)** Polyacrylamide gel analysis of the rigor kinesin–BicD construct. (*) denote side products with truncated kinesin tails, which do not interfere with the experiments. **(C)** Representative kymograph of a kinesin-DDB complex containing three different dye-labeled dynein moelcules. **(D)** The velocity of the three-color colocalized complexes at 100 µM ATP. The bars indicate mean with SD (0.031 ± 0.018 µm/s).

Figure 6. (A) Example trace of DDB-2 taking consecutive steps. Red segments indicate the dwell time before consecutive forwards steps, while blue segments indicate the dwell time before consecutive backward steps. **(B)** Relationship between dwell time, step sizes of consecutive steps, and applied forces for DDB complexes. Forward steps are shown in red, and backward steps in blue. Data were binned using a 0.5-pN force window. Bars represent the mean ± SEM. **(C)** Example trace of DDB-2 taking oscillatory steps. Red segments indicate the dwell time before forward steps, while blue segments indicate the dwell time before backward steps. **(D)** Relationship between the dwell time, step sizes of oscillatory steps, and applied forces for DDB complexes. Forward steps are shown in red, and backward steps in blue. Data were binned using a 0.5-pN force window. Bars represent the mean ± SEM.

Figure 7. Proposed model of a third dynein dimer being recruited to a DDB complex through a second BicD molecule under applied force.