Multi-omics profiling of cross-resistance between ceftazidime-avibactam and meropenem identifies common and strain-specific mechanisms in *Pseudomonas aeruginosa* clinical isolates

Bartosz J. Bartmanski*[1,] Anja Bösch*[2,8], Steven Schmitt[3], Niranjan R. Ireddy[2,8,10], Qun Ren[4], Jacqueline Findlay[5], Adrian Egli[7], Maria Zimmermann-Kogadeeva[¶1], Baharak Babouee Flury[¶2,8,9]

[1]*Genome Biology Unit, European Molecular Biology Laboratory, Heidelberg, Germany*

[2]*HOCH, Cantonal Hospital St. Gallen, Medical Research Center, St. Gallen, Switzerland*

[3]*Department of Biosystems Science and Engineering, ETH Zürich, Basel, Switzerland*

Current address: Myria Biosciences AG, Basel, Switzerland

[4]*Empa, Swiss Federal Laboratories for Materials Science and Technology, St. Gallen, Switzerland*

[5]*Medical and Molecular Microbiology, Faculty of Science and Medicine, University of Fribourg, Fribourg, Switzerland*

[7]*Institute of Medical Microbiology, University of Zurich, Zurich, Switzerland*

[8]*HOCH, Cantonal Hospital St. Gallen,Division of Infectious Diseases, Infection Prevention and Travel Medicine, St. Gallen, Switzerland,*

[9]*Department of Infectious Diseases, Inselspital, Bern University Hospital, University of Bern*

[10]*Graduate School for Cellular and Biomedical Sciences, University of Bern, Bern, Switzerland*

* equal contribution

¶ co-corresponding authors:

Maria Zimmermann-Kogadeeva, Genome Biology Unit, European Molecular Biology Laboratory, Heidelberg, Germany, maria.zimmermann@embl.de

25 Baharak Babouee Flury, HOCH, Cantonal Hospital St. Gallen, Division of Infectious Diseases,
26 Infection Prevention and Travel Medicine, St. Gallen, Switzerland baharak.baboueeflury@h-
27 och.ch
28

Abstract

Pseudomonas aeruginosa is a highly versatile and resilient pathogen that can infect different tissues and

rapidly develop resistance to multiple drugs. Ceftazidime-avibactam (CZA) is an antibiotic often used to

treat multidrug resistant infections, however, the knowledge on the CZA resistance mechanisms in *P.*

aeruginosa is limited. Here we performed laboratory evolution of eight clinical isolates of *P. aeruginosa*

exposed to either CZA or meropenem (MEM) in sub-inhibitory concentrations, and used multi-omics

profiling to investigate emerging resistance mechanisms. The majority of strains exposed to MEM

developed high resistance (83%, 20/24 strains from eight clinical isolates), with only 17% (4/24)

acquiring cross-resistance to CZA. The rate of resistance evolution to CZA was substantially lower (21%,

5/24), while 38% (9/24) acquired cross-resistance to MEM. Whole-genome sequencing revealed strain

heterogeneity and different evolutionary paths, with three genes mutated in three or more strains: *dacB* in

CZA-treated strains and *oprD* and *ftsI* in MEM-treated strains. Transcriptomic and proteomic analysis

underlined heterogeneous strain response to antibiotic treatment with few commonly regulated genes and

proteins. To identify genes potentially associated with antibiotic resistance, we built a machine learning

model that could separate CZA- and MEM-resistant from sensitive strains based on gene expression and

protein abundances. To test some of the identified associations, we performed CRISPR/Cas9 genome

editing that demonstrated that mutations in *dacB, ampD,* and to a lesser extent in *mexR* directly affected

CZA resistance. Overall, this study provides novel insights into the strain-specific molecular mechanisms

regulating CZA resistance in *Pseudomonas aeruginosa.*

Importance

Pseudomonas aeruginosa is one of the most difficult-to-treat pathogens in the hospital, which often

acquires resistance to multiple antibiotics. Ceftazidime-avibactam (CZA) is an essential antibiotic used to

treat multidrug resistant infections, but its resistance mechanisms are not well understood. Here we

investigated the evolution of resistance to CZA and meropenem (MEM) in eight clinical bacterial isolates

from patients' blood, urine, and sputum. While the rate of resistance evolution to MEM was higher than to

CZA, MEM-resistant strains rarely acquired cross-resistance towards CZA. To identify changes at the

genome, transcriptome and proteome levels during antibiotic exposure, we performed multi-omics

profiling of the evolved strains, and confirmed the effect of several genes on antibiotic resistance with

genetic engineering. Altogether, our study provides insights into the molecular response of *P. aeruginosa*

to CZA and MEM and informs therapeutic interventions, suggesting that CZA could still be effective for

patients infected with MEM-resistant pathogens.

Keywords: *Pseudomonas aeruginosa*, ceftazidime-avibactam, meropenem, resistance

mechanisms, genomics, transcriptomics, proteomics, machine learning, crispr/cas9, dacB

Introduction

Pseudomonas aeruginosa is an opportunistic pathogen causing serious infections in hospitalized patients,

including chronic respiratory infections in patients with cystic fibrosis (1). This pathogen can adapt to

diverse ecological niches (2,3), and is considered a difficult-to-treat pathogen because of its versatile

antibiotic resistance mechanisms, hypermutable strains, and antibiotic tolerant cells (4–7).

Given a high prevalence of multidrug resistance (MDR) in *P. aeruginosa*, appropriate antibiotic selection

is essential. Acute, life-threatening infections must be treated immediately, while antibiotic susceptibility

testing results of clinical isolates often takes up to three days to obtain (8). Empiric treatment often

includes beta-lactam antibiotics, such as piperacillin-tazobactam, broad spectrum cephalosporins or

carbapenems such as meropenem (MEM) (9). Despite their notable efficacy against *P. aeruginosa* (8,10)*,*

multiple resistance mechanisms against these antibiotics have emerged (11–14). Ceftazidime-avibactam

(CZA), which combines ceftazidime with a non-β-lactam β-lactamase inhibitor, became a promising

treatment option with a broad spectrum activity against MDR pathogens (15,16). The introduction of

CZA has not been without setbacks. Soon after its release, resistance was described in *Klebsiella*

*pneumonia*e isolates (17,18). Resistance to CZA has also been described *in vitro* in *P. aeruginosa* that

were incubated in the presence of increasing concentrations of CZA. All resistant variants exhibited

mutations in their *ampC,* mainly in the Omega-loop region (19). In addition, cross-resistance to CZA was

revealed in *P. aeruginosa* with single nucleotide polymorphisms (SNPs) in *ampC* and *ampR*, which led to

ceftolozane-tazobactam resistance (20,21).

Multi-omics, including transcriptomics or proteomics, are increasingly used to study antibiotic resistance

mechanisms across molecular layers (22–24). Complex antibiotic responses reported in *P. aeruginosa* and

other pathogens range from changes in energy metabolism, ribosomal activity, and DNA metabolism, to

peptidoglycan biosynthesis, and stress response (25,26). While CZA responses include de-repression of

cephalosporinase AmpC, efflux pump upregulation, porin-loss/modification, and deficiencies in DNA

mismatch repair (27), the exact resistance mechanisms remain unclear (28–30). Another important aspect

of managing *P. aeruginosa* infections is the understanding of cross-resistance evolution, particularly in

cases previously treated with meropenem (MEM). This information is essential for rapid clinical

decision-making, but relevant data on this topic is currently lacking.

This study aims to elucidate resistance and cross-resistance mechanisms to MEM and CZA in *P.*

aeruginosa through *in vitro* evolution experiments, multi-omics profiling and genome engineering,

offering valuable insights for physicians selecting empirical treatments for *P. aeruginosa* infections pre-

treated with meropenem. Additionally, it highlights *P. aeruginosa* strategies for MEM and CZA

adaptation and resistance, which could guide the development of new antibiotics and combination

therapies.

Results

Experimental evolution of antimicrobial resistance to MEM and CZA in clinical isolates

To investigate the evolution of resistance and cross-resistance to MEM and CZA in *P. aeruginosa*, we

collected eight clinical isolates from three different infection locations (blood, sputum, and urine),

confirmed their sensitivity to the drugs (Supplementary Table 1), and exposed them to sub-inhibitory

concentrations of either drug for 18 days in triplicates (Figure 1a). We observed that among the strains

exposed to sub-inhibitory levels of CZA 21% (5/24) became resistant to CZA, while 38% (9/24)

demonstrated the development of cross-resistance to MEM in accordance with the European Committee

on Antimicrobial Susceptibility Testing (EUCAST) resistance breakpoint guidelines of MIC>8 mg/L (31)

(Figure 1b, Supplementary Figure 1a). In contrast, most strains (20/24, 83%) subjected to sub-inhibitory

concentrations of MEM acquired MEM resistance, with only a minority of 17% (4/24) concurrently

displaying cross-resistance to CZA (Figure 1c, Supplementary Table 2). This observation also holds true

if we consider a subset of five isolates which had a low initial MEM MIC (MIC<=1 mg/L)

(Supplementary Figure 1b). The increase of MIC was much higher for MEM compared to CZA (median

increase of 32-fold and 5.3-fold versus 2.7-fold and 3-fold upon exposure to MEM and CZA,

correspondingly, Supplementary Table 2).

Figure 1. Experimental evolution of antimicrobial resistance to MEM and CZA in clinical isolates. (A)

Experimental design. (B) Resistance evolution to MEM and CZA in strains exposed to CZA. (C)

Resistance evolution to MEM and CZA in strains exposed to MEM. (D) Percentage of MEM- and CZA-

resistant strains (MIC>8 mg/L) among strains exposed to either MEM or CZA. In (B) and (C), gray

horizontal dotted line corresponds to the MIC=8 mg/L threshold. Boxplots represent median and

interquartile range (between the first and the third quartile), while the whiskers extend from the box to

the farthest data point lying within 1.5x the interquartile range from the box for n=24 points (n=8 strains

in triplicates). MIC - minimal inhibitory concentration. In (A), genomesequencer-2 icon by DBCLS

MEM and CZA induce distinct molecular responses across multi-omics layers

Given that only a minority, constituting 21% of the strains exposed to CZA, had exceeded the clinically

defined resistance breakpoint of MIC>8 mg/L (31) after 18 days, we extended the evolutionary

experiment for a longer period of time to capture the molecular mechanisms underlying the development

of resistance. Specifically, one representative strain evolved from each of the parental isolates was

subjected to sub-inhibitory concentrations of the respective antibiotics for a maximum of 42 days of

continuous passaging. For technical reasons, we were able to collect the data from six strains evolved in

CZA and five strains evolved in MEM out of eight parental strains. After this prolonged period, four out

of six strains acquired CZA resistance (MIC>8 mg/L), with two of them being cross-resistant to MEM

(MIC>8 mg/L), and one having the intermediate resistance. One of the CZA-sensitive strains also had

intermediate MEM resistance. While all five MEM-exposed strains acquired high resistance (MIC>96

mg/L), only two of them acquired CZA cross-resistance (Figure 2a, Supplementary Table 3). Notably,

during exposure to MEM and CZA, strains also evolved resistance to other beta-lactams, cephalosporins

and other antibiotics (Supplementary Table 4).

For all the parent and the evolved strains, we performed whole genome sequencing, and transcriptomics

and proteomics analysis with three technical replicates to investigate acquired mutations and changes in

gene expression and protein abundance upon antibiotic treatment.

Whole genome sequencing revealed that the six strains carried five sequence types (ST) and were

genetically diverse, representing different clades of *P. aeruginosa* phylogenetic tree (Supplementary

Figures 2 and 3). Most evolved strains accumulated a similar number of mutations when exposed to

MEM or CZA (between three and twenty nine), while one of the strains (1_B) was a hypermutable strain,

accumulating more than 150 mutations (Figure 2b, Supplementary Table 5). Several genes acquired

mutations independently in several strains or in the same strain exposed to either of the drugs. The most

frequently mutated genes were penicillin binding proteins *ftsI* (in two strains exposed to CZA and four

strains exposed to MEM) and *dacB* (in four strains exposed to CZA). These were followed by outer

membrane protein *oprD* (in three strains exposed to MEM), two-component system sensor *phoQ* (in two

strains exposed to MEM and one strain exposed to CZA), and multi-drug efflux pump *mexB* and its

regulator *mexR* (in one strain exposed to CZA and two strains exposed to MEM) (Supplementary Table

5). However, none of the mutations could be unambiguously attributed to the levels of acquired CZA or

MEM resistance.

Exposure to MEM or CZA induced large changes in gene expression (up to 1390 differentially expressed

genes across strains, abs(log2(fold change)) > 1, FDR-adjusted p-value < 0.05) and protein abundance (up

to 888 differentially abundant proteins, abs(log2(fold change))>1, FDR-adjusted p-value < 0.05) (Figure

2c, Supplementary Tables 6 and 7, Supplementary Figures 3 and 4). While gene and protein responses

were distinct between the two drugs (up to 1329 differentially expressed genes and 888 differentially

abundant proteins per strain between MEM and CZA conditions), gene ontology enrichment analysis

revealed several common processes affected by either of the antibiotic treatments. These included

membrane transport, protein secretion and cellular localization. Furthermore, exposure to CZA and MEM

affected amino acid, lipid and nonribosomal peptide metabolic processes in different strains (Figure 2d,

Supplementary Table 8).

While we observed some common responses to MEM and CZA exposure between strains, there was no

process that could be unambiguously attributed to antimicrobial resistance. For example, while several

genes involved in protein transport, localization and secretion were regulated in CZA-resistant strains 1_B

and 2_C upon CZA exposure, they were not enriched among genes regulated by the two other resistant

strains 4_C and 5_B. The response to CZA of the highly CZA-resistant strain 5_B was unique to this strain and

characterised mainly by the changes in biosynthetic and metabolic processes. Therefore, next we set out

to compare molecular changes across strains in more detail to identify potential antibiotic-resistance

features.

Figure 2. Molecular responses to MEM and CZA exposure across clinical isolates. (A) MIC values for strains exposed to either CZA or MEM after the initial 18 days of passaging and after 24-42 days of passaging. (B) Number of mutations induced by MEM or CZA exposure (depicts the sum of all types of detected mutations, such as SNP, insertion, deletion and complex rearrangements). (C) Number of differentially expressed genes and differentially abundant proteins induced by MEM or CZA exposure compared to parent strains (PAR) (log2(fold change)>=1, FDR<=0.05 determined using Wald test and unpaired two-sided moderated t-test for gene expression and protein abundance, respectively, and adjusted with Benjamini–Hochberg procedure). (D) GO enrichment analysis of differentially abundant

genes and proteins. Color highlights significantly enriched GO terms (FDR<=0.05). For visualization

purposes, only GO terms that pass a significance threshold of FDR<=0.001 in at least two strains for

transcriptomics or in at least one strain for proteomics are shown.

Molecular signatures of evolved strains are largely specific to the parental strain

To compare molecular signatures between strains and conditions we performed dimensionality reduction

of the normalised transcriptomics and proteomics profiles with principal component analysis (PCA). This

analysis revealed that the evolved strains exposed to either MEM or CZA grouped together with the

parental strain, underlining that the parental strain determined the global transcriptional and protein

abundance profiles of the evolved strains more than the antibiotic exposure (Figure 3a). We note that due

to insufficient sample quality, we had to exclude the MEM-evolved 2_C strain from our analysis.

Indeed, while MEM or CZA treatment induced large transcriptional (1895 or 2178 out of 5563 detected

genes) and protein (984 or 809 out of 4028 detected proteins) changes across strains, most of these

changes were strain-specific (Figure 3b, Supplementary tables 6, 7 and 9). MEM induced more common

transcriptional and protein changes across strains than CZA. About 41% of proteins changing upon MEM

exposure (401 out of 984) and 40% of proteins changing upon CZA exposure (320 out of 809) were also

changing on the transcript levels. They were mainly enriched in the nucleic acid metabolic processes,

phenazine biosynthesis, and nitrogen compound transport (GO:0090304, GO:0002047 and GO:0071705)

in MEM, and pyruvate family amino acid and branched-chain amino acid catabolic processes in CZA

(GO:0009080 and GO:0009083, Supplementary Table 8). The only genes significantly upregulated by

four strains upon MEM treatment were *mexA* and *mexB,* and the corresponding proteins were upregulated

as well (Figure 3b, Supplementary Table 9). While no proteins were differentially abundant in more than

three strains cultured in CZA, there were seven genes that were significantly up- or down-regulated in

five strains cultured in CZA: porin B *oprB* (PA3186), beta-lactamase *ampC* (PA4110) and ribosomal

protein S3AE (PA4111), as well as four genes belonging to an operon of ABC transporter genes PA3187-

PA3190 (Supplementary Table 9). Since these genes were differentially expressed in both CZA-resistant

and sensitive strains, they are likely not unambiguously determining resistance, but reflecting the stress

response and bacterial attempts to extrude the antibiotic. To put the findings observed for the six clinical

isolates into a broader perspective, we leveraged publicly available datasets on *P. aeruginosa* PAO1

strain exposed to MEM (32,33), as well as transcriptomic profiles of large clinical isolate collections

tested for MEM or CZA resistance (34,35). While *mexA* was highly expressed in CZA-resistant strains

and not differentially expressed by MEM-treated PAO1 strain, *ampC* and PA4111 were highly expressed

both in MEM-resistant strains and in PAO1 treated with MEM. Three more genes out of the nine genes

discussed above were also upregulated in PAO1 strain in response to MEM (PA3186-88, Supplementary

Table 10), supporting that they are rather markers of MEM-response but not unambiguous resistance

determinants. *P. aeruginosa* PAO1 response to MEM shared up to 40% of differentially expressed genes

with the clinical isolates, providing an additional reference for general drug response of an antibiotic-

sensitive strain (Supplementary Figure 6, Supplementary Table 11).

Figure 3. Molecular signatures of evolved strains are largely specific to the parental strain. *(A) PCA*

plots of normalised transcriptomics and proteomics data for the parent strains and strains evolved in

MEM or CZA. (B) Number of differentially expressed transcripts, differentially abundant proteins, or

both, shared between each number of strains (log2(fold change)>=1, FDR<=0.05 determined using

Wald test and unpaired two-sided t-test for gene expression and protein abundance, respectively, and

adjusted with Benjamini–Hochberg procedure). PAR refers to parental strains.

A machine learning model identifies genes and proteins predictive of MEM and CZA resistance

To investigate whether MEM- or CZA-resistant strains have distinctive molecular signatures, we used a

machine learning method Partial Least Squares-Discriminant Analysis (PLS-DA), which is widely

applied for discriminative analysis and classification in high dimensional datasets (36,37). PLS-DA aims

to maximise the covariance between the sample class (defined as high resistance to CZA or MEM,

MIC>=32 mg/L) and the predictor features (gene expression or protein abundance), and extracts latent

features (latent factors) that capture the essential information for discriminating between the two classes.

First, we built four PLS-DA models (for each of the two antibiotics and each of the two omics data types)

using the full dataset, including parent and CZA- and MEM-evolved strains (51 samples per model). All

four models could separate resistant from sensitive strains with high accuracy (accuracy 0.96 and higher)

(Figure 4a, Supplementary Table 12). To test the generalizability of the models, we performed a leave-

one-strain-out analysis, where each of the models was trained on the data from five strains (42-45

samples) and tested on the data from the sixth strain not included in the training (6-9 samples) (Figure

4b). Reflecting the heterogeneity between strains, resistance prediction accuracy substantially varied

depending on which strain was left out (Figure 4c). Both MEM- and CZA-resistance models trained on

proteomics data achieved higher accuracy than models trained on transcriptomics data (median accuracies

0.72 and 0.67 compared to 0.67 and 0.36, Supplementary Table 12).

We then took advantage of the latent factor projections calculated by PLS-DA to identify genes and

proteins most informative for the prediction of each model. For each of the gene or protein features, we

calculated how many leave-one-strain-out models included this feature among the top 100 features sorted

by the latent factor projection coefficients (out of 5563 total features for transcriptomics and 1923 total

features for proteomics). While most of the top predictive features were model-specific, we identified

small subsets of features that were repeatedly picked by five or all six leave-one-strain-out models of each

type (Figure 4d, Supplementary Table 12). For MEM resistance prediction, the models based on

transcriptomics picked genes involved in membrane functions, while most frequently picked proteins

were involved in metabolic processes, specifically carbohydrate derivative, carnitine and dTDP-rhamnose

metabolism (Figure 4e, Supplementary Table 12). For CZA resistance prediction, the selected gene

features were attributed to aminoglycan and peptidoglycan catabolic processes, while protein features

were involved in gene expression, protein folding and cysteinyl-tRNA aminoacylation (Figure 4e,

Supplementary Table 12). We then focused on the features selected by five or more MEM- or CZA-

predicting models (40 and 24 features correspondingly, Figure 4d), and filtered them further by

calculating whether they were significantly differentially expressed between CZA-sensitive and resistant

strains (FDR<=0.01, Supplementary Table 13). This resulted in 9 features for MEM-resistance and 12

features for CZA-resistance (Supplementary Figures 5 and 6, Supplementary Table 14). Among the 12

CZA-resistance features, three genes were reported to affect resistance to different antibiotics in a

genome-wide transposon mutant study (30) (Supplementary Table 14). Two of these genes are associated

with alginate biosynthesis (PA0460 and PA0764 (*mucB*)), while the third one, glutathione hydrolase

proenzyme *ggt* (PA1338) was also reported to be downregulated in a gentamicin-resistant clinical isolate

(39). Alginate and other exopolysaccharides are known to affect biofilm formation, pathogenicity and

antibiotic resistance (40–43). The other genes from the list are involved in cell wall biosynthesis (PA4421

(*mraZ*)) or metabolic functions (PA2197, PA3945, PA4847 (*accB*) PA3082 (*gbt*) PA5556 (*atpA*)) and

have been associated with virulence, antibiotic response or resistance (44–48), while the upregulated

selenide, water dikinase *selD* (PA1642) was proposed as a drug target in *Klebsiella* (49) (Supplementary

Table 14). Overall, PLS-DA enabled identification of transcriptomics and proteomics features that are

associated with a higher resistance to MEM or CZA, which may include antibiotic resistance

mechanisms, as well as stress response to the higher antibiotic levels in the antibiotic-resistant group, or

general characteristics of highly pathogenic strains, such as biofilm formation capacities.

Figure 4. A machine learning model identifies genes and proteins predictive of MEM and CZA resistance. *(A) PLS-DA plots for models built for transcriptomics (left) or proteomics (right) data to predict high CZA (top) or MEM (bottom) resistance (threshold at MIC>=32 mg/L). (B) Scheme of the*

leave-one-strain-out approach to identify genes and proteins predictive of CZA or MEM resistance. (C)

Accuracy of the six leave-one-strain-out models on the leave-one-strain-out subset (calculated based on

n=6-9 samples for each left out strain) for each of the omics data types and antibiotics. (D) Number of

predictive features selected by one to six of the leave-one-strain-out models (only top 100 features per

model were considered for comparison). (E) GO enrichment analysis of features selected by >=5 leave-

one-strain-out models. The top five GO terms per model are shown sorted by the p-value. Only

GO:1901135 passes the FDR<=0.1 threshold.

Antibiotic exposure-associated features span from membrane and transport processes to energy and lipopolysaccharide metabolism

Antibiotic exposure-associated features span from membrane and transport processes to energy and

lipopolysaccharide metabolism

To summarise antibiotic exposure-associated processes, we combined genes and proteins identified with

mutation analysis, differential transcriptomic and proteomic analyses as well as the PLS-DA models.

From mutation analysis, we selected genes in which mutation occurred at least two times independently

in our assay (19 genes); from differential analysis we selected genes and proteins significantly up- or

downregulated (abs(log2(fold change))>=1 in either of the omics, FDR<0.05 in both omics) by at least

five strains from transcriptomics, four strains from proteomics and three strains from both (17

genes/proteins). Lastly, from PLS-DA we selected features that were highly important in at least five

leave-one-strain-out models and additionally differentially abundant between sensitive and resistant stains

(FDR<0.01, 21 genes/proteins, Supplementary Table 14). Since some genes or proteins were selected by

multiple of these analyses, there were 56 features selected in total (Supplementary Table 15).

Mutations most frequently occurred in penicillin binding proteins *dacB* (only upon CZA treatment) and

ftsI (both upon MEM and CZA treatment), while the expression of these genes was unchanged or

decreased (Figure 5, Supplementary Table 15). The most pronounced upregulation both on the transcript

and the protein levels was observed in *mexAB-oprM* efflux system in the majority of strains exposed to

MEM and two of the three strains which acquired high resistance to CZA, however, this upregulation was

not coinciding with cross-resistance. Mutations were also detected in the *mexB* gene upon both CZA and

MEM exposure, in the *nalD* gene that was reported to contribute to the increase of *mexAB-oprM*

expression in *P. aeruginosa* (50), and in the *mexR* gene, recognized as a repressor gene of the *mexAB-*

oprM Multidrug Efflux Operon (51) (Figure 5, Supplementary Table 15). The expected antibiotic

resistance determinant beta-lactamase *ampC* was upregulated in the majority of the strains exposed to

CZA, however, it did not exclusively determine CZA resistance or cross-resistance, since both CZA-

sensitive and resistant strains had high *ampC* expression levels according to the transcriptomic profiling.

Antibiotic exposure-associated genes and proteins belonging to metabolism and energy production

category were selected either by differential analysis or PLS-DA and were mainly downregulated,

indicating that antibiotic exposure impaired growth and metabolism of resistant strains more than

sensitive strains, likely due to the differences in the administered antibiotic doses.

The two categories of genes/proteins identified by all three types of analysis were membrane proteins and

outer membrane channels and polysaccharide and lipopolysaccharide metabolism (Figure 5,

Supplementary Tables 15). Mutations occurring within *oprD* were uniquely observed in strains exposed

to MEM, but not coinciding with cross-resistance to CZA. The expression of *oprD* as well as the outer

membrane protein *oprE* were generally downregulated (Figure 5, Supplementary Table 15). Genes

involved in lipopolysaccharide metabolism were mutated mainly upon MEM exposure (the *phoP-phoQ*

two-component regulatory system, murein peptide ligase *mpl*, and cell division protein *ftsZ*), while the

expression of these and other genes in this category was generally increasing both upon MEM and CZA

exposure.

To summarise, three types of analysis point towards mutations and expression changes in penicillin

binding proteins, efflux pumps membrane proteins, as well as lipopolysaccharide metabolism genes as the

most generally associated with antibiotic exposure, while highlighting drug-specific responses, such as

mutations in *dacB* upon CZA exposure or *oprD* upon MEM exposure. These affected processes likely

reflect not only antimicrobial resistance strategies, but also bacterial stress response and general virulence

factors.

Figure 5. Overview of the antibiotic exposure-associated features identified by different types of analyses. *The heatmap depicts transcriptomics log fold changes on the left and proteomics log fold changes on the right within each box for a given strain and gene. Red; up-regulated, blue; down-*

regulated. Mutations found for each gene in a given strain are depicted by a circle (non-synonymous

SNP), a triangle (synonymous SNP) or a cross (deletion). The top two rows show the MIC values for CZA

and MEM after the specified number of days. The panels on the right indicate the type of analysis through

which each gene was identified (mutations, differential analysis, PLS-DA), and whether the gene was

followed up in a CRISPR-Cas9, a transposon mutagenesis assay, or an expression vector cloning assay.

Note that due to technical reasons we removed MEM-exposed 2_C strain from all the analyses.

CRISPR-Cas9 genome editing reveals that dacB mutation has a strong effect on CZA resistance

To test whether specific identified mutations can affect CZA or MEM resistance in *P. aeruginosa*, we set

out to perform CRISPR-Cas9 genome editing of the *P. aeruginosa* mPAO1 strain using gene sequence

templates from the highly resistant strains. Specifically, we focussed on several of the most frequently

mutated genes *dacB*, *mexB* and *mexR*, the heavily upregulated *ampC* and its regulator *ampD* (Figure 5).

The mutations in *mexR* and *mexB* were introduced from strain 1_B, which exhibited high levels of

resistance to both CZA and MEM. The introduction of the L13P (T38C) mutation in *mexR*, which is

accompanied by a 20-fold increase in its expression upon MEM treatment of 1_B (Supplementary Table 6),

resulted in a 3-fold rise in MEM MIC and 2-fold rise in CZA MIC respectively. Introducing the R620C

(C1858T) mutation in *mexB* led to a 1.5-fold increase in CZA MIC without affecting MEM MIC (Table

1).

For the *ampC* deletion introduced based on the template from isolate 5_B, we also observed a moderate

increase of 1.5 fold in CZA resistance and no effect on MEM resistance (Table 1), likely explained by the

fact that this deletion was upstream of the avibactam binding site.

The strongest effect on CZA MIC was observed by introducing the *dacB* mutation acquired by strain 2_C

exposed to CZA (Table 1): the CZA MIC increased four fold, while no discernible impact on MEM

resistance was detected. In contrast, introducing another *dacB* mutation from CZA-resistant strain 5_B did

not result in any changes in the MIC of either CZA or MEM (Table 1).

Overall, our CRISPR-Cas9 assays could confirm the effects on CZA resistance of *dacB* and *mexR*,

although the effects were much less pronounced than the resistance patterns of the evolved strains. Along

with CRISPR-Cas9 assays, we also performed MIC assays with a subset of transposon insertion mutants

in the *P. aeruginosa* mPAO1 mutant library (n=15 mutants, Figure 5, Supplementary Table 16). From the

tested transposon mutants, we observed substantial differences to the wildtype CZA MIC in two out of

fifteen: 25% reduction in MIC in *mexB* mutant and 3-fold increase in MIC in *mexR* mutant

(Supplementary Figure 9, Supplementary Table 16). These results underline the composite nature of

antibiotic resistance mechanisms, as genetic perturbations of single genes cannot fully recapitulate the

resistance phenotypes of the evolved strains.

Table 1. Results of CZA and MEM resistance assays of the CRISPR/Cas9 edited mutants of P.

aeruginosa mPAO1.

Gene	Strain	Mutation	CZA MIC mPAO1	CZA MIC CRISPANT	CZA MIC fold change	MEM MIC mPAO1	MEM MIC CRISPANT	MEM MIC fold change
dacB	2_C	c.1282A>C p.Thr428Pro	0.50	2.00	4.00	0.50	0.50	1.00
dacB	5_B	c.1349T>A p.Val450Glu	0.50	0.50	1.00	0.50	0.50	1.00
mexR	1_B	c.38T>C p.Leu13Pro	0.50	1.00	2.00	0.50	1.50	3.00
mexB	1_B	c.1858C>T p.Arg620Cys	0.50	0.75	1.50	0.50	0.50	1.00
ampC	5_B	c.711_731del ACGGGTCG GTCCCGGC CCGCT p.Arg238_Le u244del	0.50	0.75	1.50	0.50	0.50	1.00

Introduction of the wild-type ampD gene reversed CZA resistance in the mutant strain

As CRISPR-Cas9 genome editing was unsuccessful for the *ampD* mutation despite several attempts, we

opted for an alternative approach. We cloned the wild-type *ampD* allele from the sensitive 2_C parent strain

(CZA MIC=2 mg/L) into the pUCP24 vector and introduced it into the evolved 2_C strain (CZA MIC=128

mg/L) via transformation assays using *Escherichia coli* TOP10. Complementation with the parent *ampD*

gene led to a 64-fold decrease of resistance in the mutant strain (CZA MIC=2 mg/L, Table 2), confirming

that *ampD* mutation was the leading cause of acquired CZA resistance in this strain.

Table 2. CZA MIC of clinical isolates expressing ampD gene on a pUCP24 vector. Parent 2$_C$ is the

unevolved clinical isolate, Mutant 2$_C$ is the 2$_C$ isolate evolved in MEM.

Strain	Parent (2$_C$)	Mutant (2$_C$)	Parent/ pUCP24	Mutant/ pUCP24	Parent/ pUCP*ampD*	Mutant/ pUCP*ampD*
CZA MIC, mg/L	2	128	2	128	2	2

Discussion

Treatment of *P. aeruginosa* infections is challenging due to the versatility of strains and their potential to

develop antimicrobial resistance. In this multi-omics, multi-assay investigation of the *in vitro* evolution of

resistance to meropenem (MEM) and ceftazidime-avibactam (CZA) in eight clinical *P. aeruginosa*

isolates, we addressed these challenges both from the clinical and from the systems microbiology

perspectives.

From the clinical perspective, our observations revealed that strains subjected to sub-inhibitory

concentrations of MEM exhibited a considerably higher rate of and faster resistance evolution compared

to those exposed to CZA. This suggests prolonged/inappropriate MEM use may accelerate resistance,

potentially diminishing its therapeutic efficacy. In contrast, CZA demonstrated a lower propensity for

resistance development.. These findings are in line with a recent study where *P. aeruginosa* isolates

displayed greater increases of the mean MIC for MEM compared to CZA when subjected to increasing

concentrations of the respective antibiotics (52).

Although cross-resistance to CZA appeared in fewer cases, continuous surveillance is important to

inform and guide individualized treatment decisions for *P. aeruginosa* infections (6,7,9).

While six genes were found to be mutated independently by at least three evolved strains, these mutations

were not clearly linked to resistance patterns (Figure 5). This reflects prior reports of overall lower

mutation frequency in *P. aeruginosa*, underpinning that phenotypic variability may play a stronger role in

the development of antibiotic resistance than adaptive mutations (53). Consistent with previous reports on

high transcriptional heterogeneity in *P. aeruginosa* clinical isolates (53,54), we observed that the parental

strain was the main driver of the transcriptional and proteomic profiles regardless of the antibiotic

exposure.

Despite heterogeneous responses, machine learning approaches have previously been successful at

predicting antibiotic resistance or virulence of clinical isolates based on genome or transcriptome features

(34,55,56). Gene expression was more informative than mutation information for predicting resistance to

meropenem and ceftazidime in *P. aeruginosa* (34). While our study's small sample size limits building

generalizable machine learning models, PLS-DA's feature importance analysis helped us to identify

candidate genes and proteins associated with resistance. Though potentially reflecting general drug stress

response rather than direct resistance mechanisms, the identified genes and proteins could still serve as

resistance and virulence biomarkers and inform combination treatments to suppress bacterial adaptation

mechanisms and reduce antibiotic resistance.

The main categories of genes and proteins associated with antibiotic exposure include expected antibiotic

resistance mechanisms such as membrane proteins, efflux pumps, and penicillin binding proteins, as well

as a large group of metabolic genes involved in energy production and lipopolysaccharide metabolism.

Energy metabolism was mostly downregulated, likely reflecting stress-induced reduced bacterial

metabolism Conversely, lipopolysaccharide metabolism genes were generally upregulated, including

those involved in alginate (PA0764 (*mucB*), PA0460) and lipid A biosynthesis (PA3552 (*arnB*), PA3554

(*arnA*)). Increased expression of arnA/B genes, along with mutations in their regulator *phoQ* are linked to

resistance against polymyxin antibiotics and antimicrobial peptides (53–55), and overexpression of *arnA*

and *arnB* were observed to be more common among *P. aeruginosa* isolates resistant to CZA, when

compared to other anti-pseudomonal antibiotics (60). Similarly increased expression of alginate

exopolysaccharide biosynthesis was previously observed in antibiotic resistant strains isolated from a

cystic fibrosis patient (54), and also appeared predictive of ceftazidime or ciprofloxacin resistance in

machine learning models (34,61). Since alginate production affects biofilm growth and decreases

clearance by the host immune response (54), targeting alginate biosynthesis may be a strategy to

complement antibiotic therapy.

The *mexAB-oprM* efflux system was preferentially upregulated in MEM-exposed strains. This system is

regulated by the transcriptional repressors *mexR* and *nalD*, with additional indirect influence from *nalC*

(62). Loss-of-function mutations in *mexR* and *nalD* are linked to increased resistance or elevated MICs to

CZA (24,60). In our assays *nalD* mutations appeared in two MEM-exposed strains with significant

upregulation of *mexA*, *mexB*, and *oprM*; however, these genes were also upregulated in three other strains

with *nalD* downregulation, indicating *nalD* mutations are not the sole mechanism modulating the

mexAB-oprM efflux system. Additionally, mutations in *oprD*, encoding the outer membrane porin *oprD*,

resulted in substantial downregulation of this gene and were observed exclusively in MEM-exposed

strains, further highlighting its role in carbapenem resistance (63).

In contrast, significant upregulation of *ampC* was primarily observed in strains exposed to CZA,

suggesting that this antibiotic induces the cephalosporinase. Mutations in *dacB*, encoding penicillin-

binding protein PBP4 and implicated in ampC derepression via ampR (30,59), were also found

exclusively in CZA-selected strains.

A recent study detected AmpC overexpression in many of the isolates resistant to CZA (64). Mushtaq et

al. (65) evaluated the role of *ampC* derepression in CZA resistance using 26 *Pseudomonas aeruginosa*

strains with ampC mutations. Their findings showed that avibactam effectively reversed high-level

ceftazidime resistance mediated by ampC, significantly reducing MICs in fully derepressed mutants.

However, one strain with both high-level efflux activity and *ampC* derepression exhibited persistent

resistance, highlighting the combined impact of efflux and *ampC* mutations on CZA resistance.

In our follow-up experiments with transposon insertion mutants in *P. aeruginosa* mPAO1 strain, only two

out of fifteen mutants showed significant resistance changes, with CZA MIC decreasing by 25% or

increasing up to three fold, though five other mutations were previously linked to altered resistance

against various antibiotics (38) (Supplementary Table 16). Since transposon insertions may not replicate

changes seen in evolved clinical isolates, this could explain the assay's inability to reproduce resistance

phenotypes.

To precisely address mutations found in genes associated with β-lactam resistance, we performed

CRISPR/Cas9 genome editing of four of the mutated genes: *dacB*, *ampC*, *mexB* and *mexR*.

Mutations in *ampC* have been shown to enhance the ability of *ampC* to break down ceftazidime and to

evade inhibition by avibactam more effectively (64). The introduction of a deletion within the *ampC,*

identified in one of our strains that exhibited high resistance to CZA, resulted in only a slight increase in

the CZA MIC level.

Previous studies identified a G115S mutation in *dacB* linked to *ampC* derepression (30), but its role in

CZA resistance was unclear. While mutations slightly increased resistance overall, the most notable

change was induced by a mutation in *dacB*, resulting in a four-fold increase in CZA MIC-levels, though

allMICs remained below the clinically relevant threshold of MIC>8 mg/L. These results underline that

antibiotic resistance mechanisms involve complex interactions beyond single-gene effects, emphasizing

the need for a systems-level approach to understand their multifaceted nature.

Cloning experiments revealed that introducing a wild-type copy of the *ampD* gene had the most

pronounced effect on restoring CZA sensitivity to mutant strains—highlighting its key role in acquiring

CZA resistance. However, attempts to introduce mutations via CRISPR/Cas9 were unsuccessful despite

several efforts; cloning served as an alternative approach but exaggerated *ampD*'s effect due to

overexpression from high-copy vectors like pUCP24 (62) compared to   chromosomally encoded *ampD*.

Editing through CRISPR/Cas9, by contrast, would better reflect bacterial physiology because this

approach enables modification of the endogenous *ampD* gene to be done in a precise manner that

maintains native expression levels and regulatory control, thus more closely mimicking natural genetic

variations in bacterial populations.

Together with previous reports, this study emphasizes the importance of comprehensive sampling and

multi-omics analyses for identifying biomarkers associated with antibiotic resistance and inform

combination therapies designed to suppress its evolution in clinicals isolates of *P. aeruginosa.*

Materials and Methods

Bacterial isolates and culture conditions

Eight clinical isolates of *Pseudomonas aeruginosa* were obtained from Kantonsspital St. Gallen hospital in Switzerland. The isolates were sourced from three different clinical sites, namely urine (n=4), sputum (n=3), and blood culture (n=1). For the multistep resistance selection process, bacterial cells were cultured in Luria Bertani (LB) Broth using Nunc™ CELLSTAR® 6-well plates (Greiner Bio-One GmbH, Frickenhausen, Germany). The cultures were maintained with shaking at 120 rpm on a microplate shaker (Edmund Bühler GmbH, Germany) at a temperature of 37°C.

Ceftazidime-avibactam (Zavicefta, Pfizer®) and meropenem (Meronem, Pfizer®) were procured from our hospital pharmacy at Kantonsspital St. Gallen. Antibiotic stock solutions were prepared by dissolving powder stocks in accordance with the manufacturer's instructions. These solutions were then filter-sterilized through a 0.2 µm filter and stored at -20°C until used.

MIC determination / antimicrobial susceptibility testing

Minimal inhibitory concentrations (MICs) for several antibiotics were determined by broth microdilution in Mueller–Hinton (MH) broth using the Sensititre GNX2F plates (Thermo Fisher scientific, East Grinstead, United Kingdom). MICs for meropenem (MEM) and ceftazidime-avibactam (CZA) were further determined by Etest-strips (bioMérieux, Chemin de l'Orme, France) and broth microdilution method (Supplementary Table 4).

MICs for ceftolozane-tazobactam (TOL-TAZ) were determined by broth microdilution in Mueller Hinton (MH) broth using ComASP™ Ceftolozane-tazobactam (Liofilchem, Brunschwig, Basel, Switzerland) and by Etest (bioMérieux, Chemin de l'Orme, France). The MIC determination adhered to the latest European Committee on Antimicrobial Susceptibility Testing (EUCAST) recommendations (66), with values interpreted in accordance with the 2022 EUCAST criteria (version 12.0).

Evolution for antimicrobial resistance (multi-step resistance selection)

Eight representative strains were selected from each of the original eight *Pseudomonas aeruginosa*

isolates. This process involved isolating a single colony forming unit (CFU), which was subsequently

streaked onto a fresh Mueller Hinton (MH) agar plate. From each of these plates, three distinct colonies

were carefully chosen to serve as parental strains for subsequent passage experiments. This resulted in 24

strains being passaged in each of the antibiotics (24 strains in CZA, 24 strains in MEM). For each

individual strain, separate minimal inhibitory concentration (MIC) measurements were conducted for

ceftazidime-avibactam (CZA) and meropenem (MEM). To initiate the experiments, a single colony of

each strain was selected after overnight growth at 37°C on an MH agar plate. Subsequently, each chosen

colony was grown in 5 ml of Luria Bertani (LB) broth at 37°C for 18 hours without the presence of

antibiotics (overnight culture). The strains were then concurrently exposed to sub-inhibitory

concentrations, equivalent to one-fourth of the MIC values for each respective strain, of either CZA (n=3)

or MEM (n=3). In brief, 5 ml of LB broth containing the corresponding antibiotic at a concentration of

one-fourth of the MIC was inoculated with 25 µl of the overnight broth. The cultures were then agitated

on a shaker under the previously described conditions, maintained at 37°C for a duration of 24 hours.

Subsequently, a daily transfer of 25 µL from the overnight culture was made to new LB broth containing

CZA or MEM concentrations corresponding to one-fourth of the measured MIC. Following six

consecutive days under the consistent influence of the same antibiotic concentration, the minimal

inhibitory concentrations (MICs) were reassessed using the Etest. Based on the outcomes of these re-

evaluated MICs, modifications were implemented to the antibiotic concentration for the subsequent cycle

of passage.

Aliquots of 50 µl of the bacterial cultures were sampled after each six passages and kept in 1 ml 10%

Glycerol at −80°C for further molecular investigations. Furthermore, a subset of strains was chosen at

random and underwent Matrix-Assisted Laser Desorption/Ionization Time-of-Flight (MALDI-TOF)

analysis to ensure the absence of contaminants. To thoroughly assess the trajectory of resistance

development and cross-resistance, the aforementioned protocol was replicated over a total of 18

sequential days. To comprehensively investigate the molecular mechanisms underlying resistance, those

specific strains that had not exceeded the clinical breakpoint defined for each respective antibiotic, as

specified by the EUCAST guidelines (67), underwent additional exposure to sub-inhibitory

concentrations of the corresponding antibiotics. This exposure was continued until the strains developed

resistance or for a maximum of 42 days. Concurrently, an identical procedure was conducted in the

absence of antibiotic pressure, serving as negative controls for comparison.

Whole Genome sequencing

DNA extractions were conducted following the protocols outlined in the handbooks of two distinct kits:

the QIAamp DNA Mini Kit (QIAGEN, Hilden, Germany) was employed for Illumina sequencing, while

the QIAGEN Genomic-tip 100/G + Genomic DNA Buffer Set (QIAGEN, Hilden, Germany) kits were

used for the parental strains that underwent PacBio sequencing. Whole genome sequencing was carried

out by Novogene (Cambridge, United Kingdom) and the Genomics Facility of the Department of

Biosystems Science and Engineering of ETH Zürich (Basel, Switzerland).

For PacBio sequencing, a 10 kb SMRTbell library was prepared using the SMRTbell Template Prep Kit

1.0-SPv3 (Pacific Biosciences of California, USA). The qualified high-molecular-weight DNA was

fragmented to approximately 10 kb, followed by processes such as damage repair, end repair, and adapter

ligation. Subsequently, size selection was accomplished using a Size-Selection System.

The SMRTbell-Polymerase Complex was prepared utilizing the Sequel™ Binding Kit 2.0 (Pacific

Biosciences of California, USA) and subsequently subjected to sequencing on a Sequel SMRT Cell

(Pacific Biosciences of California, USA).

For Illumina sequencing, DNA samples were processed for library preparation following the

manufacturer's recommendations of the NEBNext® DNA Library Prep Kit (New England BioLabs, US).

Index codes were incorporated for each sample. In summary, genomic DNA was randomly fragmented to

a size of 350 bp. The DNA fragments underwent end polishing, A-tailing, ligation with adapters, size

selection, and further PCR enrichment. Subsequently, PCR products were purified using the AMPure XP

system, their size distribution was assessed using the Agilent 2100 Bioanalyzer (Agilent Technologies,

CA, USA), and quantification was performed using real-time PCR. The library was then sequenced on a

NovaSeq 6000 S4 flow cell employing the PE150 strategy.

Mutation identification

Fasta files from the whole genome sequencing were analyzed with the snippy perl package (version 4.6.0)

(https://github.com/tseemann/snippy) to call the single nucleotide polymorphisms (SNPs) and other types

of mutations using each parent strain genome sequenced before treatment as a reference. Other than

options specifying the input files and output folder, snippy was called with default parameters and force

and report flags and parallelised on 16 cores.

Sample preparation for transcriptomic profiling

The gene expression profiles of the resistant strains and the corresponding susceptible parent strains were

analyzed with RNA sequencing involving three technical replicates for each condition. For RNA

extraction, the RNAprotect Bacteria Reagent & RNeasy Mini kits (QIAGEN, Hilden, Germany) were

utilized, following the RNAprotect® Bacteria Reagent Handbook protocols 4 (Enzymatic Lysis and

Proteinase K Digestion of Bacteria) and 7 (Purification of Total RNA from Bacterial Lysate Using the

RNeasy Mini Kit). An optional on-column DNase digestion was performed using the RNase-Free DNase

set (QIAGEN, Hilden, Germany), and ribosomal RNA was depleted with the NEBNext® rRNA

Depletion Kit (New England BioLabs, Allschwil, Switzerland). The resulting RNA was fragmented,

reverse transcribed, and prepared for sequencing on an Illumina NextSeq 500 platform. Sequencing

comprised 150 bp single-end reads with a target of 5-10E6 reads per strain.

Protein extraction

3ml of OD 1 bacteria culture was pelleted by centrifugation (14'000 x g for 5 minutes) and was washed

4x with 1x PBS and lysed with 100 µl of buffer (4% (w/v) SDS, 100 mM Tris/HCL pH 8.2, 0.1M DTT +

phosphoSTOP (Roche Diagnostics GmbH, Mannheim, Germany). To each sample 100 µl of lysis buffer

(4% Sodium dodecyl sulfate (SDS) in 50 mM triethylammonium bicarbonate buffer (TEAB) were added.

The samples were frozen in liquid nitrogen for 1 min and subsequently sonicated (sonication bath) for 2

min. These freeze/thaw cycles were repeated for a total of three times. To each sample 12.5 Units of

benzonase (Sigma-Aldrich) were added and incubated for 15 min at RT, 700 rpm in a thermoshaker

(Eppendorf).

The samples were treated with High Intensity Focused Ultrasound (HIFU) for 1 minute at an ultrasonic

amplitude of 85% followed by further extraction using a tissue homogenizer (TissueLyser II, QIAGEN)

for 4 min at 30 Hz. After another 1 min of HIFU the samples were centrifuged at 20'000 x g for 10 min.

The protein concentration was determined using a Lunatic (Unchained Labs) instrument. For each sample

50 µg of protein were taken and reduced with 2 mM TCEP (tris(2-carboxyethyl)phosphine) and alkylated

with 15 mM iodoacetamide at 60°C for 30 min.

Protein digestion

The sp3 protein purification, digest and peptide clean-up was performed using a KingFisher Flex System

(Thermo Fisher Scientific) and Carboxylate-Modified Magnetic Particles (GE Life Sciences) (68,69).

Beads were conditioned following the manufacturer's instructions, consisting of 3 washes with water at a

concentration of 1 ug/ul. Samples were diluted with an equal volume of 100% ethanol (50% ethanol final

concentration). The beads, wash solutions and samples were loaded into 96 deep well- or micro-plates

and transferred to the KingFisher.

Following steps were carried out on the robot: collection of beads from the last wash, protein binding to

beads (14 min), washing of beads in wash solutions 1-3 (80% ethanol, 3 min each), protein digestion

(offline from the KingFisher, overnight at RT with a trypsin:protein ratio of 1:50 in 50 mM TEAB) and

peptide elution from the magnetic beads (water, 6 min). The digest solution and water elution were

combined and dried to completeness. Afterwards the peptides were re-solubilized with 40 µl of 3%

acetonitrile, 0.1% formic acid for MS analysis.

Proteomic analysis with liquid chromatography-mass spectrometry

Mass spectrometry (MS) analysis was performed on an Orbitrap Fusion Lumos (Thermo Scientific)

equipped with a Digital PicoView source (New Objective) and coupled to a M-Class UPLC

(Waters). Solvent composition of the two channels was 0.1% formic acid for channel A and 0.1%

formic acid, 99.9% acetonitrile for channel B. For each sample 2 μl of peptides were loaded on a

commercial MZ Symmetry C18 Trap Column (100Å, 5 μm, 180 μm x 20 mm, Waters) followed by

nanoEase MZ C18 HSS T3 Column (100Å, 1.8 μm, 75 μm x 250 mm, Waters). The peptides were

eluted at a flow rate of 300 nl/min. After an initial hold at 5% B for 3 min, a gradient from 5 to 24%

B in 80 min and 36% B in 10 min was applied. The column was washed with 95% B for 10 min and

afterwards the column was re-equilibrated to starting conditions for additional 10 min. Samples were

acquired in a randomized order. The mass spectrometer was operated in data-dependent mode

(DDA) acquiring a full-scan MS spectrum (300–1'500 m/z) at a resolution of 120'000 at 200 m/z after

accumulation to a target value of 500'000. Data-dependent MS/MS were recorded in the linear ion

trap using quadrupole isolation with a window of 0.8 Da and HCD fragmentation with 35%

fragmentation energy. The ion trap was operated in rapid scan mode with a target value of 10'000

and a maximum injection time of 50 ms. Only precursors with intensity above 5'000 were selected

for MS/MS and the maximum cycle time was set to 3 s. Charge state screening was enabled. Singly,

unassigned, and charge states higher than seven were rejected. Precursor masses previously selected

for MS/MS measurement were excluded from further selection for 20 s, and the exclusion window

was set at 10 ppm. The samples were acquired using internal lock mass calibration on m/z 371.1012

and 445.1200. The mass spectrometry proteomics data were handled using the local laboratory

information management system (LIMS) (70) and all relevant data have been deposited to the

ProteomeXchange Consortium via the PRIDE (http://www.ebi.ac.uk/pride) partner repository with the

data set identifier PXD044998.

Protein identification and label free quantification

The acquired raw MS data were processed by MaxQuant (version 1.6.2.3), followed by protein

identification using the integrated Andromeda search engine (71). Spectra were searched against a

Pseudomonas aeruginosa reference proteome (version from 2021-01-12), concatenated to its decoyed

fasta database and common protein contaminants. Carbamidomethylation of cysteine was set as fixed

modification, while methionine oxidation and N-terminal protein acetylation were set as variable.

Enzyme specificity was set to trypsin/P allowing a minimal peptide length of 7 amino acids and a

maximum of two missed-cleavages. MaxQuant Orbitrap default search settings were used. The maximum

false discovery rate (FDR) was set to 0.01 for peptides and 0.05 for proteins. Label free quantification

was enabled and a 2 minutes window for match between runs was applied. In the MaxQuant experimental

design template, each file is kept separate in the experimental design to obtain individual quantitative

values. Protein fold changes were computed based on Intensity values reported in the proteinGroups.txt

file. The proteinGroups.txt file generated by MaxQuant was used for downstream analysis to compute a

moderated t-test (72) for all proteins quantified with at least 2 peptides employing the R package limma

(73). Protein abundances were normalised using quantile normalisation. A set of functions implemented

in the R package SRMService (74) was used to filter for proteins with 2 or more peptides allowing for a

maximum of 4 missing values, and to normalize the data with a modified robust z-score transformation

and to compute p-values using the t-test with pooled variance. If all measurements of a protein are

missing in one of the conditions, a pseudo fold change was computed replacing the missing group average

by the mean of 10% smallest protein intensities in that condition.

Transcriptomics data analysis

Quantification of transcriptomic data was performed by using Trimmomatic (version 0.39) (75) followed

by Salmon (version 1.10.1) (76) packages. DESeq2 (version 1.38.0) (77) R package was used to perform

normalisation and differential expression analysis, first by importing the data generated by Salmon

package using tximport R package (version 1.26.0) (78), then by running DESeqDataSetFromTximport

and DESeq functions. The aforementioned tools were run on all samples using a Snakemake pipeline

(58). Differential analysis was performed for each strain comparing expression levels in each evolved

strain (on CZA and MEM) to the parent strain expression, as well as between CZA- and MEM-evolved

strains of the same parent. Furthermore, to filter the results from PLS-DA feature importance analysis, all

the strains were grouped based on their MEM- or CZA-resistance levels with the threshold of MIC=32

mg/L indicating high resistance. Differential analysis was performed between all sensitive versus resistant

strains for each antibiotic (for MEM: all parent strains and CZA-exposed 2_C, 3_C, 4_C, 5_B and 6_C strains

versus CZA-exposed 1_B and all MEM-exposed strains; for CZA: all parent strains, CZA-exposed 3_C and

6_C and MEM-exposed 2_C, 3_C, 4_C and 5_B versus CZA-exposed 1_B, 2_C, 4_C, 5_B and MEM-exposed 1_B and 6_C

strains), as well as all sensitive versus resistant exposed to the same antibiotic (for MEM: CZA-exposed

2_C, 3_C, 4_C, 5_B and 6_C strains versus CZA-exposed 1_B strain; for CZA: CZA-exposed 3_C and 6_C versus

CZA-exposed 1_B, 2_C, 4_C, 5_B strains, or MEM-exposed 2_C, 3_C, 4_C and 5_B strains versus MEM-exposed 1_B

and 6_C strains). DESeq2 utilises Wald test to compare expression levels between conditions with

Benjamini-Hochberg correction for multiple hypothesis testing. All the processing of the data was

executed on the EMBL Heidelberg HPC cluster (80).

Partial least squares discriminant analysis

PLS-DA was performed by using the function PLSRegression from scikit-learn python package (version

1.3.0) to predict whether CZA or MEM resistance was high (1) or low (0). Threshold of MIC=32 mg/L

was used to define high resistance. First set of models used all of the transcriptomic and proteomic data

for fitting (n=51 samples), to determine the separation of samples with high and low MIC values in the

lower dimensional space. The second set of models used a leave-one-strain-out strategy - data related to

one of the strains left out of the training set (n=6-9 samples), then this data was used as a test set to

determine the accuracy of the predictions for the model trained on all the other strains (n=42-45 samples).

To determine a set of the most important features (genes or proteins) contributing to the separation of the

samples with high and low MIC values, top one hundred features by absolute values of weights were

selected from each leave-one-strain-out model for the first and the second PLS factor (200 features per

model in total).

 Gene ontology enrichment analysis

Gene ontology terms were determined by querying PantherDB REST API

(https://pantherdb.org/services/oai/pantherdb/enrich/overrep) with a set of genes or proteins that resulted

from differential expression and differential abundance analyses and PLS-DA analysis. Significantly up-

or down-regulated (FDR < 0.05 and abs(log2FC) > 1) genes and proteins from differential expression

and differential abundance analyses were passed to the PantherDB to perform over-representation

analysis. Similarly, genes and proteins determined by PLS-DA as contributing to the top two hundred

weights of leave-one-strain-out models and appearing in at least five models were passed to PantherDB

API. Over-representation analysis done by PantherDB service assumes that all *P. aeruginosa* genes are

present in the reference list.

Sequence typing and phylogenetic analysis

To have the genetic background of the strains in this study compared with each other and with genomes

of other publicly available strains, sequence typing analysis and phylogenetic analysis were performed.

Sequence typing was performed using PubMLST web service (https://pubmlst.org) (81). Phylogenetic

analysis was performed using Mashtree (version 1.4.6) (78), first comparing just the strains in this study,

then comparing them to strains with complete genomes available from a database of pseudomonas

genomes (https://pseudomonas.com) (82–84) and matplotlib (version 3.8.3) (85).

Comparison to public datasets

Several publicly available datasets were used for comparison with the results of the differential

expression analysis conducted in this study. Dong *et al. (2024)* dataset (86) was processed from raw

reads with the same pipeline as for the dataset in this study, specifically by using Salmon (version 1.10.1)

(76) followed by DESeq2 (version 1.38.0) (77). Khaledi *et al.* (2020) dataset, consisting of a counts table,

was processed with DESeq2 as well, by running DESeqDataSetFromMatrix and then DESeq functions.

Additionally, two datasets, Alam 2023 (87) and Castanheira 2022 (35), had already undergone differential

expression analysis. Significantly up- or down-regulated genes from each dataset were compared with

those from this study on a per-strain basis. This comparison was performed by calculating the ratio of the

size of the intersection of two given sets to the size of each set and applying Fisher's exact test to the

intersection. P-values were subsequently adjusted with the Benjamini-Hochberg procedure.

Data selection and visualisation for the summary plot

A data-driven approach was employed to select the genes and proteins shown in Figure 5 from mutation

analysis, differential analysis and PLS-DA. From mutation analysis, genes with mutations present in at

least two strains were selected. From differential analysis, genes and proteins significantly up- or

downregulated (abs(log2(fold change))>=1 in either of the omics, FDR<0.05 in both omics) by at least

five strains from transcriptomics, four strains from proteomics and three strains from both were selected.

From PLS-DA, genes and proteins that appear in the top two hundred features in at least five leave-one-

strain-out models and additionally significantly differentially abundant between the groups of sensitive

and resistant strains (see section *Transcriptomics data analysis* for details) were selected. The grouping of

the final list of genes and proteins, obtained by combining these sets of genes and proteins, was

performed by ChatGPT by providing the list of locus tags and using the prompt: "can you group the

following list of genes based on their functions" followed by manual checks and corrections.

MIC measurement of the selected mutant strains of P. aeruginosa mPAO1

P. aeruginosa mPAO1 parent strain and its transposon insertion mutants were obtained from the *P.*

aeruginosa Two-Allele Library of the Manoil Lab (88). Bacteria were activated on Plate-Count-Agar

(Sigma Aldrich, Steinheim, Germany). Single colonies from the plates were picked up and inoculated in

25 mL Müller Hinton Broth (MHB, Sigma Aldrich, Steinheim, Germany) and cultivated overnight at

37°C and 160 rpm. 250 μl of the overnight culture was added to 25 mL MHB and further incubated for 3

hours to obtain the exponentially growing cells. The final culture was diluted to an optical density (OD)

of 0.1 measured at 600 nm, which corresponds to approximately 10^8 CFU/ml, of which 500 μl was

transferred to 50 ml MHB and mixed well by vortexing. Antibiotic stock solutions for MEM were

prepared as mentioned above. CZA for micro broth dilution was prepared with ceftazidime (Sigma

Aldrich, Steinheim, Germany) and avibactam (Sigma Aldrich, Steinheim, Germany), the latter with a

fixed concentration of 4 mg/l per well.

The MIC assays of the MEM and CZA were performed in 96 well plates (BRAND® microplate

BRANDplates®, Sigma Aldrich, Steinheim, Germany) covered with an air-permeable foil (Breathe-Easy

sealing membrane Sigma Aldrich, Steinheim, Germany) and incubated at 37°C without shaking. All

assays were done in MBH medium supplemented with increasing concentrations of antibiotics, and

inoculated with mPAO1 mutants of $5*10^5$ CFU/ml. For each condition, four replicates were included.

After 24 h at 37°C, the bacterial growth was measured at OD 600 nm.

Cloning of ampD gene in the evolved CZA-resistant strain

The wildtype *ampD* allele was amplified from the parental 2_C isolate using primers ampD-Fw (5′-
GATGATGAGCTCGCGCCGCTGGATTAGAGT-3')   and   ampD-Rev   (5′-
GATGATGGATCCCAGCAGCAACACCAGGAAC-3'), and cloned into vector pUCP24, before

transformation into *E. coli* Top10. Transformants were selected on plates supplemented with gentamicin

(25 mg/L). Successful transformants were confirmed by PCR amplification and sequencing of the alleles,

then transformed into the parent and the CZA-evolved mutant 2_C strains. Susceptibility testing was

performed by disc diffusion, and MICs were determined by broth microdilution according to EUCAST

guidelines.

Crispr/Cas9 genome editing

CRISPR/Cas9 genome editing was performed using the two plasmid (pCasPA/pACRISPR) method

developed by Chen et al. (89) using the λ-Red recombination system.

sgRNA oligos were designed with CRISPOR (90) and the overhangs added according to Supplementary

Table 16.

Oligos were phosphorylated and annealed by mixing 35 µl H2O, 5 µl 10x T4 Polynucleotide Kinase

Buffer (New England BioLabs, Ipswich, MA, USA), 5 µl fresh 10mM ATP, 1 µl T4 polynucleotide

kinase (New England BioLabs, Ipswich, MA, USA), 2 µl of each oligo (50 µM). After one hour

incubation at 37°C, 2.5 µl 1M NaCl were added, the mix heated to 95°C for 3 minutes and then cooled to

25°C by decreasing the temperature -0.1°C per second. Annealed oligos were diluted 1:20 and 1 µl mixed

with 1.5 µl H2O, 5 µl 2x Quick Ligase Buffer (NEB), 20nM pACRISPR plasmid, 0.5µl Quick Ligase

(400U/ µl), 0.5 µl BsaI-HFv2 (NEB) and 0.5 µl fresh 10mM ATP. Reaction was kept in a thermocycler

for 25 cycles (37°C for 3 min, 16°C for 4 min), then 80°C for 15 min, and finally cooled to 10°C. The

product was transformed into zymo mix and go (Takara, Mountain View, CA, USA) competent *E. coli*

and selected on 50 mg/l carbenicillin plates. Insertion of the sgRNA sequence was verified by sequencing

(sgSeq-F primer). Primer for the repair 500 bp arms were designed with the Takara In-Fusion Cloning

Primer Design Tool to contain the desired restriction sites and containing the desired nucleotide change

and a synonymous change of the PAM sequence. Repair arms were inserted in between the XhoI and

XbaI restriction site of the pACRISPR plasmid following the Takara In-Fusion PCR Cloning systems

protocol and insertion verified by PCR and sequencing. *P. aeruginosa* mPAO1 was made

electrocompetent by diluting an overnight culture 1:100 in fresh LBB and incubating them at 37°C until

OD600 of ~0.5. Cells were chilled on ice for 30 minutes, then were harvested by centrifugation at 3300

rpm for 5 min and pellet washed twice with 20 ml of ice-cold 10% glycerol and then resuspended in 1 ml

10% glycerol. 50 µl electrocompetent cells were mixed with up to 5 µl of pCas plasmid and

electroporated in a 1 mm cuvette (1.8kV, 200 Ω, 25 µF). Immediately after the pulse 1 ml LBB was

added and the mix was incubated for 90 minutes at 37° C, 200 rpm followed by selection on 30 mg/l

tetracycline plates.

Afterwards one colony harbouring the pCasPA plasmid was prepared as electrocompetent cells, same as

above except for the addition of 2 mg/L total concentration L-arabinose at OD 0.3. pACRISPR with the

sgRNA sequence and repair arms were mixed and electroporation followed with the same conditions. 30

mg/l tetracycline and 100 mg/l carbenicillin were used for selection.

Transformants were plasmid cured by growing them in the absence of antibiotics until evident growth and

then plating a 1:10^4 dilution on 5% sucrose LBA. The PCR amplified region of the CRISPR site of

Plasmid cured bacteria was sequenced to confirm the CRISPR event.

Data availability

The genomics and transcriptomics data have been deposited in the European Nucleotide Archive (ENA)

repository https://www.ebi.ac.uk/ena/browser/view/PRJEB76120. Proteomics data have been deposited to

the ProteomeXchange Consortium via the PRIDE (http://www.ebi.ac.uk/pride) repository with the data

set identifier PXD044998. Supplementary tables and supporting data are available in the following

Zenodo archive https://zenodo.org/uploads/15235346.

Code availability

Code to reproduce all the analysis in this work can be found in the following git repository

https://git.embl.de/grp-zimmermann-kogadeeva/PseudomonasAntibioticsResistance.

Acknowledgements

We thank the members of the Zimmermann-Kogadeeva lab for helpful discussions; the Functional

Genomics Center Zurich for technical assistance with proteomics measurements, Stefanie Altenried and

Flavia Zuber at Empa for their assistance with analysing the mPAO1 insertion mutants, Valentin Gisler

from Cantonal Hospital Aarau for providing clinical isolates, the Genomics Facility Basel (Department of

Biosystems Science and Engineering, ETH Zürich) for WGS and RNAseq analysis. We also thank the

EMBL IT Services staff for managing and provision of access to the HPC resources.

This work has been presented at the 34th Congress of the European Society of Clinical Microbiology and

Infectious Diseases in Barcelona, Spain (EO243) and the Joint Annual Meeting of the Swiss Society of

Infectious Diseases and Microbiology in Bern, Switzerland (O077).

Funding

The work was supported by the European Molecular Biology Laboratory (BJB and MZ-K) and through a

grant from the Research Committee of the Kantonsspital St.Gallen (BBF).

Author Contributions

Study and experiment design: BBF and MZ-K. Data analysis: BJB, SS, MZ-K, AB. *In vitro* experiments:

AB, JF, QR, NRI. Clinical isolate work: AB, BBF, AE. Manuscript writing: BJB, MZ-K, BBF, AB.

Manuscript editing: BJB, AB, SS, NRI, QR, JF, AE, MZ-K, BBF. Funding acquisition: BBF, MZ-K. All
authors read and approved the manuscript.

Competing interests: The authors declare no competing interests.

Statement on using AI tools

In the preparation of this research paper, we utilised generative AI tools (ChatGPT 4o and perplexity.ai)

to enhance the clarity and coherence of our writing, including this section. We carefully reviewed and

edited the AI-generated recommendations to maintain the integrity and authenticity of our work.

Generative AI-tools were not used to produce new content, but only to rephrase human-written text to

improve its clarity. ChatGPT 4o was also used to suggest the clustering of genes based on their functions

in Figure 5 (described in the Methods section), which was subsequently edited by the authors.

References

1.   Cabot G, Zamorano L, Moyà B, Juan C, Navas A, Blázquez J, et al. Evolution of Pseudomonas
aeruginosa Antimicrobial Resistance and Fitness under Low and High Mutation Rates. Antimicrob
Agents Chemother. 2016 Feb 26;60(3):1767–78.

2.   Gellatly SL, Hancock REW. Pseudomonas aeruginosa : new insights into pathogenesis and host
defenses. Pathog Dis. 2013 Apr 1;67(3):159–73.

3.   Pang Z, Raudonis R, Glick BR, Lin TJ, Cheng Z. Antibiotic resistance in Pseudomonas aeruginosa:
mechanisms and alternative therapeutic strategies. Biotechnol Adv. 2019 Jan 1;37(1):177–92.

4.   Carmeli Y, Troillet N, Eliopoulos GM, Samore MH. Emergence of Antibiotic-Resistant
Pseudomonas aeruginosa: Comparison of Risks Associated with Different Antipseudomonal Agents.
Antimicrob Agents Chemother. 1999 Jun;43(6):1379–82.

5.   Cohen NR, Lobritz MA, Collins JJ. Microbial persistence and the road to drug resistance. Cell Host
Microbe. 2013 Jun 12;13(6):632–42.

6.   Ambreetha S, Zincke D, Balachandar D, Mathee K. Genomic and metabolic versatility of
Pseudomonas aeruginosa contributes to its inter-kingdom transmission and survival. J Med
Microbiol. 2024;73(2):001791.

7.   Livermore DM. Multiple Mechanisms of Antimicrobial Resistance in Pseudomonas aeruginosa: Our
Worst Nightmare? Clin Infect Dis. 2002 Mar 1;34(5):634–40.

8.   Reynolds D, Kollef M. The Epidemiology and Pathogenesis and Treatment of Pseudomonas
aeruginosa Infections: An Update. Drugs. 2021;81(18):2117–31.

9.   Chamot E, Boffi El Amari E, Rohner P, Van Delden C. Effectiveness of Combination Antimicrobial
Therapy for Pseudomonas aeruginosa Bacteremia. Antimicrob Agents Chemother. 2003
Sep;47(9):2756–64.

10. O'Callaghan CH, Acred P, Harper PB, Ryan DM, Kirby SM, Harding SM. GR 20263, a new broad-
spectrum cephalosporin with anti-pseudomonal activity. Antimicrob Agents Chemother. 1980
May;17(5):876–83.

11. Farra A, Islam S, Strålfors A, Sörberg M, Wretlind B. Role of outer membrane protein OprD and
penicillin-binding proteins in resistance of Pseudomonas aeruginosa to imipenem and meropenem.
Int J Antimicrob Agents. 2008 May 1;31(5):427–33.

12. Riera E, Cabot G, Mulet X, García-Castillo M, del Campo R, Juan C, et al. Pseudomonas aeruginosa
carbapenem resistance mechanisms in Spain: impact on the activity of imipenem, meropenem and
doripenem. J Antimicrob Chemother. 2011 Sep 1;66(9):2022–7.

13. Shu JC, Kuo AJ, Su LH, Liu TP, Lee MH, Su IN, et al. Development of carbapenem resistance in
Pseudomonas aeruginosa is associated with OprD polymorphisms, particularly the amino acid
substitution at codon 170. J Antimicrob Chemother. 2017 Sep 1;72(9):2489–95.

14. Drawz SM, Bonomo RA. Three Decades of β-Lactamase Inhibitors. Clin Microbiol Rev. 2010
Jan;23(1):160–201.

15. Shirley M. Ceftazidime-Avibactam: A Review in the Treatment of Serious Gram-Negative Bacterial
Infections. Drugs. 2018 Apr 1;78(6):675–92.

16. Mushtaq S, Vickers A, Woodford N, Livermore DM. Activity of aztreonam/avibactam and
ceftazidime/avibactam against Enterobacterales with carbapenemase-independent carbapenem
resistance. Int J Antimicrob Agents. 2024 Mar 1;63(3):107081.

17. Humphries RM, Yang S, Hemarajata P, Ward KW, Hindler JA, Miller SA, et al. First Report of
Ceftazidime-Avibactam Resistance in a KPC-3-Expressing Klebsiella pneumoniae Isolate.
Antimicrob Agents Chemother. 2015 Sep 18;59(10):6605–7.

18. Sauerborn E, Corredor NC, Reska T, Perlas A, Vargas da Fonseca Atum S, Goldman N, et al.
Detection of hidden antibiotic resistance through real-time genomics. Nat Commun. 2024 Jun
28;15(1):5494.

19. Lahiri SD, Walkup GK, Whiteaker JD, Palmer T, McCormack K, Tanudra MA, et al. Selection and
molecular characterization of ceftazidime/avibactam-resistant mutants in Pseudomonas aeruginosa
strains containing derepressed AmpC. J Antimicrob Chemother. 2015 Jun 1;70(6):1650–8.

20. Fraile-Ribot PA, Cabot G, Mulet X, Periañez L, Martín-Pena ML, Juan C, et al. Mechanisms leading
to in vivo ceftolozane/tazobactam resistance development during the treatment of infections caused
by MDR Pseudomonas aeruginosa. J Antimicrob Chemother. 2018 Mar 1;73(3):658–63.

21. Díaz-Cañestro M, Periañez L, Mulet X, Martin-Pena ML, Fraile-Ribot PA, Ayestarán I, et al.
Ceftolozane/tazobactam for the treatment of multidrug resistant Pseudomonas aeruginosa: experience
from the Balearic Islands. Eur J Clin Microbiol Infect Dis. 2018 Nov 1;37(11):2191–200.

22. dos Santos BS, da Silva LCN, da Silva TD, Rodrigues JFS, Grisotto MAG, Correia MT dos S, et al.

Application of Omics Technologies for Evaluation of Antibacterial Mechanisms of Action of Plant-Derived Products. Front Microbiol. 2016 Sep 27;7:1466.

23. Maity R, Zhang X, Liberati FR, Rossi CS, Cutruzzolà F, Rinaldo S, et al. Merging Multi-OMICs with Proteome Integral Solubility Alteration Unveils Antibiotic Mode of Action. eLife [Internet]. 2024 Apr 18 [cited 2024 Jul 7];13. Available from: https://elifesciences.org/reviewed-preprints/96343

24. Chernov VM, Chernova OA, Mouzykantov AA, Lopukhov LL, Aminov RI. Omics of antimicrobials and antimicrobial resistance. Expert Opin Drug Discov. 2019 May 4;14(5):455–68.

25. Molina-Mora JA, García F. Molecular Determinants of Antibiotic Resistance in the Costa Rican Pseudomonas aeruginosa AG1 by a Multi-omics Approach: A Review of 10 Years of Study. Phenomics. 2021 Jun 1;1(3):129–42.

26. Jo SH, Song WS, Park HG, Lee JS, Jeon HJ, Lee YH, et al. Multi-omics based characterization of antibiotic response in clinical isogenic isolates of methicillin-susceptible/-resistant Staphylococcus aureus. RSC Adv. 2020;10(46):27864–73.

27. Khil PP, Dulanto Chiang A, Ho J, Youn JH, Lemon JK, Gea-Banacloche J, et al. Dynamic Emergence of Mismatch Repair Deficiency Facilitates Rapid Evolution of Ceftazidime-Avibactam Resistance in Pseudomonas aeruginosa Acute Infection. mBio. 2019 Sep 17;10(5):10.1128/mbio.01822-19.

28. Castanheira M, Doyle TB, Smith CJ, Mendes RE, Sader HS. Combination of MexAB-OprM overexpression and mutations in efflux regulators, PBPs and chaperone proteins is responsible for ceftazidime/avibactam resistance in Pseudomonas aeruginosa clinical isolates from US hospitals. J Antimicrob Chemother. 2019 Sep 1;74(9):2588–95.

29. Babouee Flury B, Bösch A, Gisler V, Egli A, Seiffert SN, Nolte O, et al. Multifactorial resistance mechanisms associated with resistance to ceftazidime-avibactam in clinical Pseudomonas aeruginosa isolates from Switzerland. Front Cell Infect Microbiol [Internet]. 2023 [cited 2023 Dec 7];13. Available from: https://www.frontiersin.org/articles/10.3389/fcimb.2023.1098944

30. Sanz-García F, Hernando-Amado S, Martínez JL. Mutation-Driven Evolution of Pseudomonas aeruginosa in the Presence of either Ceftazidime or Ceftazidime-Avibactam. Antimicrob Agents Chemother. 2018 Sep 24;62(10):10.1128/aac.01379-18.

31. EUCAST. Breakpoint Tables for Interpretation of MICs and Zone diameters, Version 12.0 [Internet]. [cited 2023 Dec 7]. Available from: https://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_12.0_Breakpoint_Tables.pdf

32. Dong L, Sun L, Yang Y, Yuan L, Gao W, Yu D, et al. Non-antibiotic pharmaceutical phenylbutazone binding to MexR reduces the antibiotic susceptibility of *Pseudomonas aeruginosa*. Microbiol Res. 2024 Nov 1;288:127872.

33. Alam F, Blair JMA, Hall RA. Transcriptional profiling of Pseudomonas aeruginosa mature single- and dual-species biofilms in response to meropenem. Microbiol Read Engl. 2023 Jan;169(1):001271.

34. Khaledi A, Weimann A, Schniederjans M, Asgari E, Kuo T, Oliver A, et al. Predicting antimicrobial resistance in Pseudomonas aeruginosa with machine learning□enabled molecular diagnostics. EMBO Mol Med. 2020 Mar 6;12(3):e10264.

35. Castanheira M, Doyle TB, Hubler CM, Collingsworth TD, DeVries S, Mendes RE. The plethora of
resistance mechanisms in *Pseudomonas aeruginosa*: transcriptome analysis reveals a potential role of
lipopolysaccharide pathway proteins to novel β-lactam/β-lactamase inhibitor combinations. J Glob
Antimicrob Resist. 2022 Dec 1;31:72–9.

36. Ruiz-Perez D, Guan H, Madhivanan P, Mathee K, Narasimhan G. So you think you can PLS-DA?
BMC Bioinformatics. 2020 Dec 9;21(1):2.

37. Barker M, Rayens W. Partial least squares for discrimination. J Chemom. 2003;17(3):166–73.

38. Dötsch A, Becker T, Pommerenke C, Magnowska Z, Jänsch L, Häussler S. Genomewide
Identification of Genetic Determinants of Antimicrobial Drug Resistance in Pseudomonas
aeruginosa. Antimicrob Agents Chemother. 2009 Jun;53(6):2522–31.

39. Maunders EA, Triniman RC, Western J, Rahman T, Welch M. Global reprogramming of virulence
and antibiotic resistance in Pseudomonas aeruginosa by a single nucleotide polymorphism in
elongation factor, fusA1. J Biol Chem. 2020 Nov 27;295(48):16411–26.

40. Sindeldecker D, Stoodley P. The many antibiotic resistance and tolerance strategies of *Pseudomonas
aeruginosa*. Biofilm. 2021 Dec 1;3:100056.

41. Huse HK, Kwon T, Zlosnik JEA, Speert DP, Marcotte EM, Whiteley M. Pseudomonas aeruginosa
Enhances Production of a Non-Alginate Exopolysaccharide during Long-Term Colonization of the
Cystic Fibrosis Lung. Roop Ii RM, editor. PLoS ONE. 2013 Dec 6;8(12):e82621.

42. Scott NE, Hare NJ, White MY, Manos J, Cordwell SJ. Secretome of Transmissible Pseudomonas
aeruginosa AES-1R Grown in a Cystic Fibrosis Lung-Like Environment. J Proteome Res. 2013 Dec
6;12(12):5357–69.

43. Woods EC, McBride SM. Regulation of antimicrobial resistance by extracytoplasmic function (ECF)
sigma factors. Microbes Infect. 2017;19(4–5):238–48.

44. Kong KF, Schneper L, Mathee K. Beta-lactam Antibiotics: From Antibiosis to Resistance and
Bacteriology. APMIS Acta Pathol Microbiol Immunol Scand. 2010 Jan;118(1):1–36.

45. Czub MP, Zhang B, Chiarelli MP, Majorek KA, Joe L, Porebski PJ, et al. A Gcn5-Related
N□Acetyltransferase (GNAT) Capable of Acetylating Polymyxin B and Colistin Antibiotics in Vitro.
Biochemistry. 2018 Dec 26;57(51):7011–20.

46. Sigurdsson G, Fleming RMT, Heinken A, Thiele I. A Systems Biology Approach to Drug Targets in
Pseudomonas aeruginosa Biofilm. PLoS ONE. 2012 Apr 16;7(4):e34337.

47. Cirz RT, O'Neill BM, Hammond JA, Head SR, Romesberg FE. Defining the Pseudomonas
aeruginosa SOS Response and Its Role in the Global Response to the Antibiotic Ciprofloxacin. J
Bacteriol. 2006 Oct;188(20):7101–10.

48. Duang-Nkern J, Nontaleerak B, Thongphet A, Asano K, Chujan S, Satayavivad J, et al. Regulation of
curcumin reductase *curA* (*PA2197*) through sodium hypochlorite and *N*-ethylmaleimide sensing by
TetR family repressor CurR (PA2196) in *Pseudomonas aeruginosa*. Gene. 2024 Nov 15;927:148754.

49. Ramos PIP, Fernández Do Porto D, Lanzarotti E, Sosa EJ, Burguener G, Pardo AM, et al. An
integrative, multi-omics approach towards the prioritization of Klebsiella pneumoniae drug targets.
Sci Rep. 2018 Jul 17;8:10755.

50. Sobel ML, Hocquet D, Cao L, Plesiat P, Poole K. Mutations in PA3574 (nalD) Lead to Increased
MexAB-OprM Expression and Multidrug Resistance in Laboratory and Clinical Isolates of
Pseudomonas aeruginosa. Antimicrob Agents Chemother. 2005 May;49(5):1782–6.

51. Adewoye L, Sutherland A, Srikumar R, Poole K. The MexR Repressor of the mexAB-oprM
Multidrug Efflux Operon in Pseudomonas aeruginosa: Characterization of Mutations Compromising
Activity. J Bacteriol. 2002 Aug;184(15):4308–12.

52. Castanheira M, Kimbrough JH, Lindley J, Doyle TB, Ewald JM, Sader HS. In vitro development of
resistance against antipseudomonal agents: comparison of novel β-lactam/β-lactamase inhibitor
combinations and other β-lactam agents. Antimicrob Agents Chemother. 2024 Mar 25;68(5):e01363-
23.

53. Dötsch A, Schniederjans M, Khaledi A, Hornischer K, Schulz S, Bielecka A, et al. The Pseudomonas
aeruginosa Transcriptional Landscape Is Shaped by Environmental Heterogeneity and Genetic
Variation. mBio. 2015 Jun 30;6(4):10.1128/mbio.00749-15.

54. Rojas LJ, Yasmin M, Benjamino J, Marshall SM, DeRonde KJ, Krishnan NP, et al. Genomic
heterogeneity underlies multidrug resistance in Pseudomonas aeruginosa: A population-level analysis
beyond susceptibility testing. PLOS ONE. 2022 Mar 31;17(3):e0265129.

55. Burgaya J, Marin J, Royer G, Condamine B, Gachet B, Clermont O, et al. The bacterial genetic
determinants of Escherichia coli capacity to cause bloodstream infections in humans. PLOS Genet.
2023 Aug 2;19(8):e1010842.

56. Hu K, Meyer F, Deng ZL, Asgari E, Kuo TH, Münch PC, et al. Assessing computational predictions
of antimicrobial resistance phenotypes from microbial genomes. Brief Bioinform. 2024 May
1;25(3):bbae206.

57. Ben Jeddou F, Falconnet L, Luscher A, Siriwardena T, Reymond JL, van Delden C, et al. Adaptive
and Mutational Responses to Peptide Dendrimer Antimicrobials in Pseudomonas aeruginosa.
Antimicrob Agents Chemother. 2020 Mar 24;64(4):10.1128/aac.02040-19.

58. Gatzeva-Topalova PZ, May AP, Sousa MC. Structure and Mechanism of ArnA: Conformational
Change Implies Ordered Dehydrogenase Mechanism in Key Enzyme for Polymyxin Resistance.
Struct Lond Engl 1993. 2005 Jun;13(6):929–42.

59. Segev-Zarko LA, Kapach G, Josten M, Klug YA, Sahl HG, Shai Y. Deficient Lipid A Remodeling
by the arnB Gene Promotes Biofilm Formation in Antimicrobial Peptide Susceptible Pseudomonas
aeruginosa. Biochemistry. 2018 Apr 3;57(13):2024–34.

60. Castanheira M, Doyle TB, Hubler CM, Collingsworth TD, DeVries S, Mendes RE. The plethora of
resistance mechanisms in Pseudomonas aeruginosa: transcriptome analysis reveals a potential role of
lipopolysaccharide pathway proteins to novel β-lactam/β-lactamase inhibitor combinations. J Glob
Antimicrob Resist. 2022 Dec;31:72–9.

61. Khaledi A, Schniederjans M, Pohl S, Rainer R, Bodenhofer U, Xia B, et al. Transcriptome Profiling
of Antimicrobial Resistance in Pseudomonas aeruginosa. Antimicrob Agents Chemother. 2016 Jul
22;60(8):4722–33.

62. Arnold W, Pühler A. A family of high-copy-number plasmid vectors with single end-label sites for
rapid nucleotide sequencing. Gene. 1988 Oct 15;70(1):171–9.

63. Biggel M, Johler S, Roloff T, Tschudin-Sutter S, Bassetti S, Siegemund M, et al. PorinPredict: In

Silico Identification of OprD Loss from WGS Data for Improved Genotype-Phenotype Predictions of
P. aeruginosa Carbapenem Resistance. Microbiol Spectr. 2023 Jan 30;11(2):e0358822.

64. Slater CL, Winogrodzki J, Fraile-Ribot PA, Oliver A, Khajehpour M, Mark BL. Adding Insult to
Injury: Mechanistic Basis for How AmpC Mutations Allow Pseudomonas aeruginosa To Accelerate
Cephalosporin Hydrolysis and Evade Avibactam. Antimicrob Agents Chemother. 2020 Aug
20;64(9):10.1128/aac.00894-20.

65. Mushtaq S, Warner M, Livermore DM. In vitro activity of ceftazidime+NXL104 against
Pseudomonas aeruginosa and other non-fermenters. J Antimicrob Chemother. 2010 Nov
1;65(11):2376–81.

66. EUCAST: AST of bacteria [Internet]. [cited 2023 Dec 7]. Available from:
https://www.eucast.org/ast_of_bacteria

67. EUCAST. Breakpoint Tables for Interpretation of MICs and Zone Diameters, Version 12.0.
[Internet]. Available from:
https://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_12.0_Breakp
oint_Tables.pdf

68. Hughes CS, Foehr S, Garfield DA, Furlong EE, Steinmetz LM, Krijgsveld J. Ultrasensitive proteome
analysis using paramagnetic bead technology. Mol Syst Biol. 2014 Oct;10(10):757.

69. Leutert M, Rodríguez-Mias RA, Fukuda NK, Villén J. R2-P2 rapid-robotic phosphoproteomics
enables multidimensional cell signaling studies. Mol Syst Biol. 2019 Dec;15(12):e9021.

70. Türker C, Akal F, Joho D, Panse C, Barkow-Oesterreicher S, Rehrauer H, et al. B-Fabric: the Swiss
Army Knife for life sciences. In: Proceedings of the 13th International Conference on Extending
Database Technology [Internet]. New York, NY, USA: Association for Computing Machinery; 2010
[cited 2023 Dec 7]. p. 717–20. (EDBT '10). Available from:
https://doi.org/10.1145/1739041.1739135

71. Cox J, Mann M. MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass
accuracies and proteome-wide protein quantification. Nat Biotechnol. 2008 Dec;26(12):1367–72.

72. Smyth GK. Linear models and empirical bayes methods for assessing differential expression in
microarray experiments. Stat Appl Genet Mol Biol. 2004;3:Article3.

73. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential expression
analyses for RNA-sequencing and microarray studies. Nucleic Acids Res. 2015 Apr 20;43(7):e47.

74. Wolski, W, Grossmann, J, Panse, C PC. SRMService [Internet]. protViz; 2021 [cited 2023 Dec 7].
Available from: https://github.com/protViz/SRMService

75. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. 2014 Aug 1;30(15):2114–20.

76. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-aware
quantification of transcript expression. Nat Methods. 2017 Apr;14(4):417–9.

77. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data
with DESeq2. Genome Biol. 2014 Dec 5;15(12):550.

78. Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq: transcript-level estimates

improve gene-level inferences [Internet]. F1000Research; 2016 [cited 2023 Dec 7]. Available from:
https://f1000research.com/articles/4-1521

79. Mölder F, Jablonski KP, Letcher B, Hall MB, Tomkins-Tinch CH, Sochat V, et al. Sustainable data
analysis with Snakemake [Internet]. F1000Research; 2021 [cited 2024 Dec 18]. Available from:
https://f1000research.com/articles/10-33

80. European Molecular Biology Laboratory, Pečar J, Lueck R, Wahlers M. EMBL Heidelberg HPC
Cluster. 2020 Jan 1 [cited 2025 Apr 4]; Available from: https://zenodo.org/records/12785830

81. Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics: BIGSdb software, the
PubMLST.org website and their applications. Wellcome Open Res. 2018;3:124.

82. Katz LS, Griswold T, Morrison SS, Caravas JA, Zhang S, Bakker HC den, et al. Mashtree: a rapid
comparison of whole genome sequence files. J Open Source Softw. 2019 Dec 10;4(44):1762.

83. Cock PJA, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, et al. Biopython: freely available
Python tools for computational molecular biology and bioinformatics. Bioinformatics. 2009 Jun
1;25(11):1422–3.

84. Winsor GL, Griffiths EJ, Lo R, Dhillon BK, Shay JA, Brinkman FSL. Enhanced annotations and
features for comparing thousands of Pseudomonas genomes in the Pseudomonas genome database.
Nucleic Acids Res. 2016 Jan 4;44(D1):D646-653.

85. Hunter JD. Matplotlib: A 2D Graphics Environment. Comput Sci Eng. 2007 May;9(3):90–5.

86. Dong L, Sun L, Yang Y, Yuan L, Gao W, Yu D, et al. Non-antibiotic pharmaceutical phenylbutazone
binding to MexR reduces the antibiotic susceptibility of *Pseudomonas aeruginosa*. Microbiol Res.
2024 Nov 1;288:127872.

87. Alam F, Blair JMA, Hall RA. Transcriptional profiling of Pseudomonas aeruginosa mature single-
and dual-species biofilms in response to meropenem. Microbiology. 2023;169(1):001271.

88. Jacobs MA, Alwood A, Thaipisuttikul I, Spencer D, Haugen E, Ernst S, et al. Comprehensive
transposon mutant library of Pseudomonas aeruginosa. Proc Natl Acad Sci. 2003 Nov
25;100(24):14339–44.

89. Chen W, Zhang Y, Zhang Y, Pi Y, Gu T, Song L, et al. CRISPR/Cas9-based Genome Editing in
Pseudomonas aeruginosa and Cytidine Deaminase-Mediated Base Editing in Pseudomonas Species.
iScience. 2018 Aug 31;6:222–31.

90. Concordet JP, Haeussler M. CRISPOR: intuitive guide selection for CRISPR/Cas9 genome editing
experiments and screens. Nucleic Acids Res. 2018 Jul 2;46(W1):W242–5.

1066

	1_B - CZA vs PAR	1_B - MEM vs PAR	2_C - CZA vs PAR	3_C - CZA vs PAR	3_C - MEM vs PAR	4_C - CZA vs PAR	4_C - MEM vs PAR	5_B - CZA vs PAR	5_B - MEM vs PAR	6_C - CZA vs PAR	6_C - MEM vs PAR	Alam 2022 - PAO1 - MEM	Dong 2024 - PAO1 - MEM	Khaledi 2020 - 414 strains	Castanheira 2022 - 213 strains - CZA	Castanheira 2022 - 213 strains - MEM
1_B - CZA vs PAR (n=693)	1	0.6	0.23	0.081	0.27	0.34	0.26	0.063	0.029	0.0072	0.26	0.055	0.3	0.012	0.0087	0.0014
1_B - MEM vs PAR (n=570)	0.72	1	0.26	0.056	0.24	0.29	0.22	0.061	0.016	0.007	0.29	0.077	0.26	0.019	0.007	0.014
2_C - CZA vs PAR (n=503)	0.31	0.26	1	0.068	0.29	0.33	0.35	0.1	0.012	0.022	0.47	0.093	0.23	0.028	0.004	0.002
3_C - CZA vs PAR (n=126)	0.44	0.25	0.27	1	0.68	0.68	0.53	0.19	0	0.063	0.33	0.11	0.47	0.032	0	0.0079
3_C - MEM vs PAR (n=930)	0.2	0.14	0.16	0.092	1	0.42	0.28	0.07	0.018	0.012	0.42	0.055	0.34	0.019	0.0097	0.0011
4_C - CZA vs PAR (n=1390)	0.17	0.12	0.12	0.06	0.28	1	0.076	0.0072	0.0079	0.16	0.28	0.012	0	0	0	0
4_C - MEM vs PAR (n=659)	0.28	0.19	0.27	0.1	0.4	0.54	1	0.12	0.029	0.021	0.34	0.058	0.28	0.042	0.015	0
5_B - CZA vs PAR (n=230)	0.19	0.15	0.22	0.1	0.28	0.46	0.35	1	0.026	0.035	0.22	0.061	0.33	0.026	0	0.0043
5_B - MEM vs PAR (n=34)	0.59	0.26	0.18	0	0.5	0.29	0.18	0.18	1	0	0.18	0.059	0.5	0.059	0.059	0.029
6_C - CZA vs PAR (n=20)	0.25	0.2	0.55	0.4	0.55	0.55	0.7	0.4	0	1	0.25	0.2	0.25	0.2	0	0
6_C - MEM vs PAR (n=582)	0.31	0.29	0.41	0.07	0.32	0.38	0.086	0.01	0.027	1	1	0.079	0.24	0.021	0.0034	0.0017
Alam 2022 - PAO1 - MEM (n=167)	0.23	0.26	0.28	0.084	0.31	0.39	0.23	0.084	0.012	0.24	0.12	1	0.39	0.03	0.006	0.006
Dong 2024 - PAO1 - MEM (n=1153)	0.18	0.13	0.1	0.051	0.28	0.33	0.16	0.065	0.015	0.0043	0.12	0.056	1	0.017	0.0078	0.017
Khaledi 2020 - 414 strains (n=60)	0.13	0.18	0.23	0.067	0.3	0.47	0.47	0.1	0.033	0.067	0.2	0.083	0.33	1	0.05	0.033
Castanheira 2022 - 213 strains - CZA (n=13)	0.46	0.31	0.15	0	0.69	0.5	0.77	0	0.15	0	0.15	0.077	0.69	0.23	1	0
Castanheira 2022 - 213 strains - MEM (n=2)	0.5	0.5	0.5	0	0.5	0.5	0	0.5	0.5	0	0.5	0.5	1	1	0	1

Ratio of overlap

0.0   0.5   1.0

Abundance or expression

CZA MIC level

Abundance or expression

MEM MIC level

Condition
- CZA
- MEM
- PAR