══ REVIEW COMMONS MANUSCRIPT ══

IMPORTANT:

- Manuscripts submitted to Review Commons are peer reviewed in a journal-agnostic way.
- Upon transfer of the peer reviewed preprint to a journal, the referee reports will be available in full to the handling editor.
- The identity of the referees will NOT be communicated to the authors unless the reviewers choose to sign their report.
- The identity of the referee will be confidentially disclosed to any affiliate journals to which the manuscript is transferred.

GUIDELINES:

- For reviewers: https://www.reviewcommons.org/reviewers
- For authors: https://www.reviewcommons.org/authors

CONTACT:

The Review Commons office can be contacted directly at: office@reviewcommons.org

The Lipocone Superfamily: A Unifying Theme In Metabolism Of Lipids, Peptidoglycan And Exopolysaccharides, Inter-Organismal Conflicts And Immunity

A. Maxwell Burroughs†, Gianlucca G. Nicastro†, L Aravind*

Division of Intramural Research, National Library of Medicine, National Institutes of Health, USA

*To whom correspondence should be addressed. Tel: +1 (301)594-2445; Fax: +1(301)480-9241; Email: aravind@ncbi.nlm.nih.gov

†These authors contributed equally to this work

ABSTRACT

Wnt proteins are critical signaling molecules in developmental processes across animals. Despite intense study, their evolutionary roots have remained enigmatic. Using sensitive sequence analysis and structure modeling, we establish that the Wnts are part of a vast assemblage of domains, the Lipocone superfamily, defined here for the first time. It includes previously studied enzymatic domains like the phosphatidylserine synthases (PTDSS1/2) and the TelC toxin domain from Mycobacterium tuberculosis, the enigmatic VanZ proteins, the animal Serum Amyloid A (SAA) and a further host of uncharacterized proteins in a total of 30 families. Though the metazoan Wnts are catalytically inactive, we present evidence for a conserved active site across this superfamily, versions of which are consistently predicted to operate on head groups of either phospholipids or polyisoprenoid lipids, catalyzing transesterification and phosphate-containing head group severance reactions. We argue that this superfamily originated as membrane proteins, with one branch (including Wnt and SAA) evolving into soluble versions. By comprehensively analyzing contextual information networks derived from comparative genomics, we establish that they act in varied functional contexts, including regulation of membrane lipid composition, extracellular polysaccharide biosynthesis, and biogenesis of bacterial outer-membrane components, like lipopolysaccharides. On multiple occasions, members of this superfamily, including the bacterial progenitors of Wnt and SAA, have been recruited as effectors in biological conflicts spanning inter-organismal interactions and anti-viral immunity in both prokaryotes and eukaryotes. These findings establish a unifying theme in lipid biochemistry, explain the origins of Wnt signaling and provide new leads regarding immunity across the tree of life.

INTRODUCTION

The canonical Wnt signaling network is central to developmental decisions across animals relating to axis patterning, cell fate, cell migration and proliferation and systems morphogenesis at many levels (1-7). Other crucial pathways, dubbed non-canonical Wnt signaling pathways, include those that regulate planar cell polarity and intracellular calcium levels (8-10). With these roles in development and homeostasis, dysfunction of Wnt signaling is causally associated with a range of diseases, including diverse cancer types and type II diabetes (11,12). These paracrine and autocrine signaling networks are centered on the secreted Wnt proteins that act as diffusible, extracellular messenger molecules (13). Wnt proteins are ligands for the N-terminal, cysteine-rich CBD/Fz domains of the Frizzled class of G-protein coupled receptors (GPCRs) (14,15). Modifications of the Wnt proteins via palmitoleoylation and glycosylation at internal sites are associated with their secretion (16). Palmitoleoylation of Wnt occurs at a conserved serine residue and is also required for recognition by the Frizzled receptors (17). Binding of the Frizzled receptor by Wnt recruits the Disheveled (Dsh) protein on its cytoplasm face, in turn triggering a bevy of downstream responses, resulting in β-catenin stabilization in canonical pathways (18,19). When β-catenin concentrations cross a threshold, it is translocated into the nucleus, where it acts as a transcriptional coactivator, usually with an HMG domain transcription factor to stimulate multiple transcriptional programs (20-22).

Despite its initial discovery over 40 years ago, the evolutionary origins of the Wnt protein have, until recently, been mysterious (23). In 2020, our group reported the discovery of the first prokaryotic versions of the Wnt domain (24). Using comparative genomics, we showed that these bacterial Wnt domains present contexts characteristic of toxins or effectors in biological conflict systems (24,25). Prompted by these initial observations, we set out to comprehensively identify and computationally characterize the evolutionary relationships of these newly identified Wnt homologs in an effort to understand their evolutionary history and predict their functions.

Consequently, we were able to unify the Wnt family with several other domains into a large

superfamily described for the first time herein. These include two biochemically characterized families

that were hitherto not known as Wnt homologs: the phosphatidylserine synthase (PTDSS1/2, EC:

2.7.8.29) (26-28) and the toxin domain of TelC from *Mycobacterium tuberculosis* (29). However, the

majority of the families we unify are either reported for the first time or are functionally poorly

understood, including the animal Serum Amyloid A (SAA) (30) and the vancomycin resistance protein

VanZ families (31,32). Our comparative genomics analyses, paired with existing experimental evidence,

suggest that the superfamily is broadly comprised of enzymes operating on lipid head groups (e.g.,

transesterification reactions) in a diversity of biochemical contexts, notably including the regulation of

membrane composition, extracellular biopolymer metabolism and as effectors in biological conflicts.

Thus, we identify a unifying theme across diverse aspects of lipid metabolism.

RESULTS

Identification of the structural core of the Wnt domain

While the structure of Wnt was described over a decade prior (33-35), its origins have been a mystery as

it is phyletically restricted to Metazoa. Much attention has been focused on the three extended β-

hairpins and a poorly structured loop extruding out of the core, their stabilizing cysteine residues, and

the absolutely conserved serine residue, the site of palmitoleoylation (34,36) (Figure 1A). Our discovery

of the first prokaryotic Wnt domains helped define its ancestral α-helical core, revealing the cysteine-

rich extensions as Metazoa-specific insertions. Comparison of the core of the metazoan Wnt with

structural models of the prokaryotic versions (24) revealed a shared globular domain composed of five

α-helices (Figure 1A). As the prokaryotic homologs retained just the conserved core of the Wnt

proteins, we named these the minimal Wnt (Min-Wnt) family. The other core helices of this family

contained further absolutely conserved sequence motifs (Figure 2), consistent with the enzymatic

function we had earlier proposed for the Min-Wnts (24) (see below).

Having defined this shared core, we initiated sequence-based homology searches in an effort to identify

remote homologs. Iterative position-specific sequence matrix (PSSM)-based searches (see Methods)

initially recovered animal and bacterial versions of the Serum Amyloid A (SAA) proteins, and further

rounds of searching initiated from this set of sequences further recovered a vast expansion of

additional homologous families. As an example, a search initiated with a bacterial SAA-like sequence

from *Bdellovibrio bacteriovorus* (Genbank acc: AHZ84906.1) retrieved a sequence overlapping with the

Pfam model of a "Domain of unknown function", DUF2279 (acc: WP_146898260.1, iteration: 5, e-value:

0.004), the Pfam DUF4056 domain (acc: MBW8016507.1, iteration: 5, e-value: 0.005), and sequences

automatically annotated as "YfiM" (acc: WP_019077413.1, iteration: 4, e-value: 0.004). Sequence profile-

profile searches with HHpred confirmed these relationships and captured more distant ones. A single

sequence query from the *Bacteriovorax stolpii* Min-Wnt domain (acc: WP_102242990.1, residues 1-109)

recovered the Pfam Wnt profile (PF: PF00110.23, p-value: 1.5e-6) and the Pfam SAA profile (PF:

PF00277.22, p-value: 3.7e-5); while a single sequence query initiated with a Gemmatimonadetes

sequence (acc: PYP94660.1, residues 75..170) recovered the DUF2279 Pfam profile (PF: PF10043.12, p-

value: 5.4E-21), the DUF2238 profile (PF: PF09997.12, p-value: 1.3E-07), and the DUF4056 Pfam profile

(PF: PF13265.9, p-value: 2.5E-05), among others.

Exhaustion of these searches, followed by clustering and manual inspection of the retrieved sequences

(see Methods), revealed a shared four-helix core across all of them, hereinafter referred to as H1

through H4 (Figure 1A). This 4-helix core of the domain was further confirmed by inspection of

structural models constructed for representatives of the individual families, along with the rare

instances of solved experimental structures. These comparisons established that the above-mentioned

fifth C-terminal helix in the Wnt core is a synapomorphy (shared derived character) restricted to the

Wnts and closely related families like SAA (Figure 1A). In all, the results of our clustering analysis

tallied 30 distinct families. Remarkably, of these, 17 had no pre-existing annotations. Phyletic analysis

of individual families revealed a range of distributions, from families with broad conservation in

multiple superkingdoms of Life to those restricted to a small number of lineages (see below, Figure S1).

A relationship network was constructed based on p-value and e-value scores using alignments of each

family as a query in HHalign profile-profile searches against the rest (see Methods, Figure 1B). The

Leiden community detection algorithm (37) was then applied to this network to identify higher-order

assemblages. These groupings were also supported by structural synapomorphies, such as a circular

permutation and versions with a two-stranded 'handle' (see below, Methods).

The four helices conserved across the superfamily constitute a cone-like structure (Figure 1A), with the

helices tending to coalesce on one end and opening out into a pocket on the other, lined by the

conserved sequence positions (Figures 1A, S2). The core is also marked by a linker between H1 and H2,

which adopts characteristic extended conformations in certain families and higher-order groups. While

the linkers joining H2 and H3 and H3 and H4 tend to be more constrained, there are some exceptions;

for example, the extended loop insert housing the palmitoleoylated serine residue between H2 and H3

in the metazoan Wnt family (Figure 1A).

Dramatic variability in hydrophobicity of the conserved core across the superfamily

We observed that the Wnt-related families dramatically varied in their hydrophobicity. Using an index

for transmembrane propensity (38) (see Methods) and comparing that to known transmembrane (TM)

segments, we predict that the α-helices in 18 of the 30 families are hydrophobic enough to qualify as

TM domains (Figure 1C). Thus, these are predicted to be integral membrane domains. Further, these

'hydrophobic families' often evince a deeper phyletic distribution pattern than the less-hydrophobic

families (Figure S1), implying that the ancestral version of the superfamily was likely a TM domain.

Thus, their association with the lipid membrane, combined with the cone-like shape of the conserved

core (Figure 1A), leads us to refer to the whole superfamily hereinafter as the *Lipocone* superfamily.

Alphafold3-assisted transmembrane topology prediction (39) revealed that 16 of the 17 integral

membrane families are consistently orientated with the aperture of the cone-like structure opening

toward the outer face of the membrane. One family did not yield a consistent prediction for

orientation. The predicted TM topology is also generally consistent with the domain fusions when

present: e.g., domains that are typically cytoplasmic and those that have extracellular or periplasmic

functions are respectively predicted as projecting either inside or outside the membrane (see below).

A unified biochemistry for the Lipocone superfamily

Of the 30 identified families, 26 display a striking conservation pattern of polar residues associated with

the pocket of the Lipocone domain (Figure 2, S2). Of these, a set of three positions, one mapping to each

of H2, H3, and H4, can be inferred as being ancestrally present and were likely occupied by a histidine

(H2), glutamate (H3), and aspartate (H4), though in some families their identities have secondarily

changed (Figures 2, S2). A fourth well-conserved polar position is observed at or near the end of H3;

while its ancestral identity is difficult to establish, it is frequently an aspartate or glutamate (Figure 2).

Two further well-conserved positions are often seen in H4: a polar position downstream of the broadly

conserved aspartate residue and a glycine residue near the C-terminus of H4 (Figure 2). While the

ancestral pattern is noticeably degraded in the metazoan Wnt (Met-Wnt) family, it is strongly

preserved in the prokaryotic Min-Wnt family. In solved and modeled structures, the above set of 4

conserved positions forms a predicted active site in the aperture of the Lipocone domain. This, in turn,

implies a shared biochemistry across the superfamily, with secondary inactivation in some families like

Met-Wnt (see below, Figure 2). At the same time, the differences in the specific residues in the

conserved positions between different families point to a range of distinct but related activities across

the superfamily (40-42).

Consistent with these observations, two of the families with intact active sites, the PTDSS1/2 (28,43)

and TelC (29), which we identified in this work as members of the Lipocone superfamily, have been

characterized as active enzymes operating on different lipid substrates (Figure 3A). The eukaryotic

PTDSS1/2 localizes to the endoplasmic reticulum (ER) membrane and catalyzes a reaction on the polar

head group of phosphatidylcholine or phosphatidylethanolamine (44-46). PTDSS1 and PTDSS2,

respectively, exchange the phosphate-linked choline or ethanolamine head groups with L-serine (28)

(Figure 3A). The toxin domain of TelC acts on lipid II (29), the final intermediate in peptidoglycan

biosynthesis, which couples an undecaprenyl diphosphate tail to a head group comprised of a N-

acetylmuramic acid-N-acetylglucosamine disaccharide, with a pentapeptide further linked to the

former sugar (47,48). TelC severs the link between the undecaprenol and the diphosphate coupled to

the head group (29) (Figure 3B). The reaction is comparable to that catalyzed by PTDSS1/2, as both

attack phosphate linkages in lipid head groups. However, TelC apparently directs a water molecule for

the attack in lieu of the hydroxyl group of serine directed by PTDSS1/2 (Figure 3B).

Combining the above observations, we infer the unified biochemistry for the catalytically active

families thus: 1) They act on the head groups of lipids either by removing or swapping phosphate-

linked head groups (Figure 3A-B). These would be comparable to the phospholipase D (PLD),

transphosphatidylation or polyisoprenol phosphoesterase reactions (49). 2) Given the cone-like cavity

and the hydrophobicity of the helices, the lipid tail is predicted to be housed within the lipocone with

the head group positioned in the active site. 3) In the case of the integral membrane versions, their

orientation would predict the targeting of the head groups of the outer leaf of the bilayer.

Major clades of the Lipocone superfamily

The extreme sequence divergence of the superfamily, coupled with the small size of the domain,

prevents the use of simple phylogenetic tree analyses to resolve its deep evolutionary history. Hence,

we combined community finding algorithms applied on profile-profile similarity networks, comparison

of structural features and motifs, and phyletic patterns (Figures 1B, 2, S1) to reconstruct the most

parsimonious evolutionary scenario for the diversification of the Lipocone superfamily (Figure 4, see

Methods). In the below sections, we survey the higher-order clades, highlighting their specific features.

SAW (SAA-Wnt) clade

This clade consists of four families, with the two prokaryotic families (Min-Wnt and prok-SAA) (24,50),

respectively, giving rise to their counterpart eukaryotic families (Met-Wnt and Met-SAA; Figure 4). This

clade is structurally unified by the presence of a fifth helix that stacks in the space between the H2 and

H4 helices (Figure 1A, S2). In the Wnt families, this helix is comparable in length to the core helices,

while in the SAA families, the helix is shorter, usually containing only a few turns. The clade is further

unified by the pronounced conservation of a sNxxGR motif (where 's' is a small residue) encompassing

the conserved active site position in H4 (Figure 2). SAW clade Lipocones show low overall

hydrophobicity and are known or predicted to be soluble domains. Outside of the clearly inactive

eukaryotic Wnt family, the remaining three families largely conserve the core active site residues

(Figure 2).

VanZ-Skillet clade

This clade unites seven families: the two VanZ families, VanZ-1 and VanZ-2, prototyped by the bacterial

VanZ protein originally identified in the context of vancomycin resistance and the five Skillet families,

which form a distinct subclade. These are unified by a "handle"-like structure (hence, "Skillet")

adopting a helical conformation in the H1-H2 linker (Figure 1A, S2). Strikingly, a symmetric helical

handle is present in the H3-H4 loop of the Skillet-DUF2809 and Skillet-3 families (Figure S2) of this

clade. VanZ-1 features a conserved asparagine residue in the H2 position and a DxDDxxxN motif in H4,

while VanZ-2 features RKxxH and DxxxD motifs in these respective positions (Figure 2). The Skillet

families are largely unifiable in their conservation of an ExxQ motif in H3, an aspartate three positions

upstream of the canonical H4 aspartate, and another aspartate in the H2 contributing to the active site.

These first two features specifically ally them with the VanZ-1 family (Figure 2).

While the VanZ domain was previously reported as including a fifth TM helix which is C-terminal to the

4-helix Lipocone core defined here (51,52), our survey instead reveals a striking diversity of

configurations around the core 4-helix Lipocone domain (Supplemental Material). These range from

standalone Lipocone configurations to one or more TM-helices adorning the domain at its N- and/or C-

terminus. This variation is consistent with a further tendency for the VanZ families to feature an

extensive diversity of domain fusions to both soluble globular domains and discrete TM modules (see

below).

The VanZ families are deep-branching, as suggested by their wide phyletic spread (Figure S1). VanZ-2 is

the most widespread individual Lipocone family in bacteria, with several genomes encoding multiple

paralogs (Supplemental Material) (51,53). It is also found in certain eukaryotes, including a pan-fungal

presence and in some representatives of the SAR clade. Both VanZ-1 and VanZ-2 are particularly well-

represented in Gram-positive bacterial lineages like Actinomycetota and Firmicutes, while VanZ-2 is

notably nearly universally conserved in the Bacteriodetes/Chlorobi lineage (Figure S1). In contrast,

only one of the Skillet families, Skillet-DUF2809, is widely but sporadically distributed, with the four

others being more restricted (Figures 4,S1).

YfiM clade

This clade includes three families that are consistently centrally located in the profile-profile similarity

network (Figure 1B). This is likely due to their being close in sequence conservation to the ancestral

state of the superfamily (Figure 2). Consistent with this, the YfiM-1 family also presents a structurally

minimal Lipocone domain, comprised of just the 4-helix configuration with no further elaborations.

Notably, this also extends to a lack of domain fusions in this family. In contrast, YfiM-DUF2279 and

YfiM-Griddle (DUF3943) are structurally distinguished by an unusual H1-H2 linker, which wraps around

the outside and stacks against the H3-H4 linker (Figure S2). The YfiM-Griddle family further features a

unique 'flattened' surface around the aperture of the Lipocone formed by protruding loops (hence,

"Griddle"; Figure S2). This leaves the active site pocket more accessible relative to families with more

elaborately structured inter-helix linkers. The Griddle family also features a C-terminal extension with

a two-helix hairpin (with a hhsP motif in the turn between the two helices, where 'h' is a hydrophobic

residue and 's' is a small residue) (Figure S2). The three YfiM families straddle the membrane-

propensity boundary in the plot (Figure 1C). Further, the YfiM-DUF2279 and Griddle families are

strikingly absent in Gram-positive bacterial lineages (Figure S1). Concurrent with these features, they

are often predicted as outer-membrane proteins, suggesting a role in this subcellular location (see

below).

ClaspCone-CapCone-TelC clade

Members of this clade are unified by an elaborated H1-H2 linker that often contains one or more helical

segments that are typically predicted to guard the aperture of the Lipocone domain (Figure S2). This

linker ends in a "clasp"-like element, which forms a range of structures in different families of the clade

before leading into H2 (Figure S2). The clade is also unified by a striking reduction of overall

hydrophobicity, predicting that the members of this clade are soluble domains (Figure 1C). Outside of

the divergent TelC subclade, most of the families in this clade conserve a serine residue three positions

upstream of the active site aspartate in H4, often preceded by an aromatic residue which is typically

phenylalanine. H4 also usually features a conserved asparagine four positions downstream of the

conserved aspartate active site position, immediately preceded by a small residue (Figure 2). The

second H3 active site position is generally poorly conserved, though when present, it is usually an

aspartate residue. Finally, H2 contains either a DK or xD motif four positions upstream of the canonical

H2 active site histidine residue (Figure 2).

The most rudimentary clasps are found in the ClaspCone-1, -2, and -3 families, where it is little more

than a rounded loop, though, in ClaspCone-1, a small β-hairpin emerges within it. The three ClaspCone

families are further unified by the presence of a two-helix insert leading into H2 that stacks against the

Lipocone core (Figure S2). The three CapCone families, CapCone-DUF4056, CapCone-1, and CapCone-2,

are named so for an encasing structure over the active site resembling a cap (Figure S2). They share a

conserved glycine residue six positions upstream of the active site H2 histidine and a striking S/GxxSxx

motif upstream of the conserved H4 aspartate (Figure 2). They are further unified by a pronounced β-

hairpin clasp augmented by an additional strand (Figure S2). They also display varying degrees of

degeneration of H1, along with family-specific structural elaborations.

The TelC group of this clade, prototyped by the mycobacterial TelC toxin (29), is divided into two

families featuring prokaryotic (prok-TelC) (29) and metazoan versions (Met-TelC) (54). Both TelC

families feature a "cap" with contributions from inserts in the H1-H2 and H3-H4 loops (Figure S2).

Unique to these families is the conservation of an aspartate residue located six positions downstream of

the canonical active site aspartate of H4 (Figure 2). This aspartate points away from the center of the

Lipocone and interacts with a conserved arginine from a synapomorphic C-terminal helical extension.

cpCone clade

A widespread yet sporadically distributed clade of seven families emerging as a stable community in

the profile-profile similarity network (Figure 1B) is defined by a unique structural synapomorphy: a

circular permutation (55) (hence, cpCone) placing the normally N-terminal H1 at the C-terminus of H4

(Figure S2-3). This clade is also united by unique sequence features, viz., a polar residue (typically

aspartate) six positions upstream of the conserved H2 histidine and a second glutamate three positions

downstream of the conserved H3 glutamate (Figure 2). While the circular permutation is shared across

the clade, several structural variations are seen, often within the same family (Figure S3). These

include: 1) versions containing a duplication of the Lipocone domain. While the second copy in these

versions is catalytically inactive, the H1' from the second duplicate displaces the H1 from the first copy,

suggestive of an intermediate to the circular permutation. 2) Versions retaining a candidate H1 that has

been displaced by H1' in a five-helix arrangement. 3) Those containing just the circularly permuted

core. 4) Versions showing a degradation of the H1 helix, preserving just a 3-helix core (Figure S3).

Despite this propensity for structural variation, the active site residues are strongly conserved, with

the exception of the cpCONE-i family, which we infer to be catalytically inactive (Figure 2). The core

helices of the cpCone clade are strongly hydrophobic, and they are predicted to all be integral

membrane domains (Figure 1C). Consistent with this, the eukaryotic PTSSD1/2 domains reside in the ER

membrane (44,45).

Wok family

The Wok family (partly covered by the Pfam DUF2238 model) shows a higher order grouping with the

above circularly permuted clade (Figures 1B,4) but has a phyletic distribution only rivaled by the VanZ-

2 family (Figure S1), suggesting a deep-branching origin. This family has the general shape of a wok

formed by two distinguishing structural synapomorphies: a 2-TM helix N-terminal extension and a

unique "handle" formed by the linker between the H3 and H4 (Figure S2). It additionally features a C-

terminal, rapidly diversifying cytoplasmic tail. Nevertheless, it retains the inferred ancestral active site

configuration (Figure 2). The strongly hydrophobic core helices of the Wok family predict it to be an

integral membrane enzyme (Figure 1C).

Functional themes in the Lipocone superfamily

Given our inference of shared general biochemistry across the Lipocone superfamily in targeting

phosphate-containing linkages in head groups of both classic phospholipids and polyisoprenoid lipids,

we next used contextual information from conserved gene-neighborhoods, domain architectures and

phyletic pattern vectors, a powerful means of deciphering gene function (56), to narrow down the

predictions for specific families (Figure 5, Table S1, Supplementary Data). To this end, we constructed a

graph (network) wherein the nodes are individual domains and edges indicate adjacency in domain

architectures or conserved gene-neighborhoods (Figure 6, see Methods). We then identified cliques in

these networks and merged the individual cliques containing a particular Lipocone domain to define its

dense subgraph (Figures S4-S6). We then analyzed these subgraphs to identify statistically significant

functional categories (Table S2; see Methods). This data was combined with existing experimental

results and the sequence and structure analysis outlined above to arrive at the functional themes

surveyed in the below sections.

Lipocone domains in membrane lipid, peptidoglycan and exopolysaccharide modifications

Across different Lipocone families, we found statistically significant connections to roles in modifying

lipid head groups in various membranes and in lipids involved in the synthesis of extracellular matrix

polymers such as peptidoglycan and lipopolysaccharides (Figure 6, Table S2, Supplementary Data).

Archetypal lipid head group exchange reactions catalyzed by the cpCone clade

One of the few experimentally characterized Lipocone families is the eukaryotic PTDSS1/2 family of the

cpCone clade, members of which exchange the head group of essential membrane phospholipids to

generate phosphatidylserine from phosphatidylethanolamine or phosphatidylcholine (Figure 3A)

(28,57). Given the pervasive presence of this clade in archaea (Figure S1), it is thus tempting to

speculate that these archaeal cpCones may play a role in the modification of Archaea-specific lipids (58-

60) through a comparable head group exchange reaction (see below).

In bacteria, the related cpCone-1 family shows operonic association with a LolA-like lipoprotein which

shuttles lipoproteins to the outer membrane (61) and a novel 4TM protein (Figure 5A). This raises the

possibility that cpCone-1 might mediate the formation of membrane domains featuring lipids with a

modified head group that act as foci for the trafficking of lipoproteins. Curiously, the cpCone-1 gene

might also be inserted between the bacterial chromosome segregation and condensation complex

subunits the Kleisin ScpA and the wHTH ScpB (62-65). The bacterial cpCONE-DUF2585 is operonically

coupled to a GNAT family NH_2-group-acetyltransferase and further linked to genes for the glycolate

oxidase GlcE and GlcF (66) and the bacterial proteasome subunits HslV and HslU (67) (Figure 5A,

Supplementary Data). These might point to the coupling of membrane lipid head group modifications

with disparate processes, such as chromosome segregation during cell division or different responses

to stress (68-70).

The Wok and YfiM-1 families in cardiolipin and modified isoprenoid lipid pathways

We observed a set of conserved gene neighborhoods displaying the mutually exclusive presence of a

synaptojanin-like phosphatase gene with one either encoding a member of the Wok family or a

cardiolipin synthase of the HKD superfamily (71) (Figure 5B, Supplementary Data). This suggested that

the latter two are analogous enzymes catalyzing equivalent reactions. The cardiolipin synthase utilizes

two phosphatidyl glycerol molecules as substrates to generate cardiolipin with the release of one of the

glycerol head groups (72). Effectively, this is comparable to the head group exchange reaction catalyzed

by the cpCone clade (Figure 3A). Hence, we propose that these members of the Wok clade are

cardiolipin synthases (Figure 3C). Distinct phosphoesterases from the synaptojanin-like enzyme, like

calcineurin (73) and HAD (74), are also observed in association with the Wok, suggesting that they

might regulate lipid composition by acting on the phospholipids or their precursors (Figure 5C). In a

distinct neighborhood, the Wok clade enzyme is coupled to carotenoid biosynthesis genes (75,76).

(Figures 5D,S4). This raises the possibility that these members might also catalyze a comparable

reaction to the above on isoprenoid lipids: they could synthesize a carotenoid from two

geranylgeranyl-diphosphate molecules (77,78). In both these contexts, the actinobacterial operons

often include genes for GT-A family glycosyltransferases, suggesting the further synthesis of

glycosylated derivatives of the lipids or carotenoids (79) (Supplementary Data). In several bacteria, a

YfiM-1 family Lipocone is operonically coupled to a UbiA-like prenyltransferase (80). This gene

neighborhood additionally codes for a slew of enzymes, such as an amidophosphoribosyltransferase

(81), a RidA-like deaminase (82), and a pair of structurally distinct phosphoesterases, respectively,

containing an HD and a PHP domain (73,83) (Figures 5E,S4 Supplementary Data). This suggests a role for

the Lipocone and the associated enzymes in generating a modified polyisoprenoid metabolite.

VanZ families in lipid head group modifications in peptidoglycan and exopolysaccharide metabolism

The widespread VanZ-1 and VanZ-2 families frequently show either gene neighborhood associations or

direct domain fusions, with diverse genes involved in both peptidoglycan and other extracellular

polysaccharide pathways. Chief among these are the lipid carrier flippase (Pfam: MviN_MATE clan) (84-

86), the UDP-GlcNAc/MurNAc lipid transferases, which generate the lipid-linked exopolysaccharide

precursors (lipid I) (48,87), and UDP-N-acetylglucosamine (UDP-GlcNAc) biosynthesis enzymes (88,89).

Despite multiple examples of crossover in functional themes, the gene-neighborhood contexts of VanZ-

1 and VanZ-2 suggest a metabolic partitioning, with VanZ-2 significantly associating specifically with

peptidoglycan-related genes and VanZ-1 significantly linking with biosynthesis genes for other

exopolysaccharides (e.g., the outer-membrane-associated lipopolysaccharide) (90) (Figures 5F,6,S5,

Supplementary Data). The latter include WaaL-like lipid A transferase (91), the polysaccharide chain-

length determination domain Wzz (92), the Wzc kinase and the "extracellular antigen"-regulating ElyC-

like domain (Pfam: DUF218) (93), and numerous nucleotide-diphosphate sugar biosynthesis and

modification enzymes (94) (Figures 5F,6,S5).

The precursors of both peptidoglycan and exopolysaccharides are synthesized in the cytosol, linked to

lipid carriers via a diphosphate linkage, e.g., the polyisoprenoid lipid undecaprenol (bactoprenol)

(90,94-97). A key step in their maturation is the flipping by the flippase of the lipid-linked intermediates

associated with the inner membrane to the outer membrane. These flipped units are then incorporated

into the maturing chain (98,99) by the peptidoglycan glycosyltransferase (GTase) (100) and the chain

length determination protein, WzzE/polymerase, WzyE (92,101), in peptidoglycan and other

exopolysaccharide maturation pathways, respectively. Based on the precedence of the PTDSS1/2

reaction (Figure 3A), we propose that VanZ-1 and VanZ-2 analogously act on the lipid carrier-linked

sugar head groups to catalyze either the extension of the polysaccharide chains through

transesterification or the terminal release of the mature chain through phosphoester cleavage (Figure

3E).

The above activities could be linked to distinct types of peptidoglycan synthesis during different phases

of the life cycle, e.g., sporulation versus vegetative growth in Bacillota. Consistent with this, VanZ-1 is

pervasive in sporulating Gram-positive bacteria (Figure S1). Terminal transfer from the lipid carrier of

the Gram-negative bacterial O-antigen (as well as other exopolysaccharides attached to the lipid A

carrier) has been attributed to the WaaL-like enzymes (91,102). However, bacteria generate further

lineage-specific polysaccharide decorations, capsule structures, and other exopolysaccharides (e.g.,

xanthan, enterobacterial common antigen (ECA), alginate, colonic acid), as well as teichoic acids (e.g.,

wall teichoic acids, WTA) (103). Notably, the analogs of WaaL, i.e., the terminal transferases for several

exopolysaccharides, including ECA and WTA, have to date escaped identification (93), making the

VanZ-2 enzymes attractive candidates. Such a function would be consistent with its widespread yet

sporadic phyletic pattern (Figure S1), keeping with the rapid evolution of these exopolysaccharides due

to strong selection for variation (104) (Figure 6).

Atypical VanZ domains in uncharacterized modifications of peptidoglycan and the outer membrane

Certain representatives of the two VanZ families also show operonic associations indicative of distinct

outer membrane-associated or peptidoglycan modification functions distinct from those described

above (Figures 5G,6 Supplementary Data): 1) An operon in FCB group bacteria couples a VanZ-2 gene

with those coding for a SprA secretin-like channel protein (105), a glycine cleavage H (GCVH)-like

lipoyl-group carrier protein (106), a 2TM protein fused via a proline-rich linker to a C-terminal TonB-C

domain (107), and a secreted, second TonB-C domain fused to a Wzi-like outer membrane protein

(OMP) superfamily β-barrel (108) (Figures 5G, 6). 2) In betaproteobacteria, certain VanZ-1 domains are

duplicated with the C-terminal copy being inactive (VanZ-i) and found in an unusual four-gene operon

with a thioredoxin-fold [2Fe-2S] ferredoxin (109), a possible lipase of the α/β-hydrolase superfamily

(110), and a metallo-β-lactamase (MβL) fold D-Ala-D-Ala cross-linking transpeptidase (111,112). 3) A

patescibacterial operon encodes a VanZ-2 domain with an ABC ATPase transporter system, either of

two structurally distinct peptidases, namely a Papain-like or glycine-glycine peptidase (113,114), fused

to the same membrane-anchored N-terminal coiled-coil region, and a further TM protein containing

one or more external Lamin-Tail domains (LTDs) predicted to bind extracellular DNA or

polysaccharides (115) (Figures 5G,6,S5). The associations in the first of the above neighborhoods point

to a distinct outer membrane-associated lipid modification, while the other two might be involved in

lineage-specific decorations/modifications of peptidoglycan associated with peptide-crosslinking or

cleavage activities.

Lipocone domains operating in the outer membrane

Contextual associations, phyletic patterns, and localization predictions support the action of two

Lipocone families directly in the outer membrane. Notably, the Griddle and YfiM-DUF2279 families are

found nearly obligately directly fused or operonically linked to several distinct OMP β-barrels (116,117)

(Figures 5H,6,S4). Up to three Griddle Lipocones, usually with a cognate OMP β-barrel, might be

encoded next to each other in the genome. Additionally, Griddle family genes are often encoded in

operons with several components of the outer membrane lipid and protein trafficking apparatus,

including the LolA-like chaperone (118), the POTRA domain (119,120), the channel-blocking Plug

domains (121), and the TolA-binding TolB-N domain (122). Further, these operons might encode a

Patatin-like lipase (123), GT-B family glycosyltransferases (79), and a range of phosphoesterases (e.g., an

integral membrane phosphatidic acid phosphatase PAP2 (124), a lipobox-containing synaptojanin

superfamily phosphoesterase (125) and a secreted R-P phosphatase (126) (see Figure 6, Supplementary

Data)). In addition to the fusion to the OMP β-barrel, the YfiM-DUF2279 family (Figure 5H) shows

operonic associations with a secreted MltG-like peptidoglycan lytic transglycosylase (127,128), a lipid-

anchored cytochrome c heme-binding domain (129), a phosphoglucomutase/phosphomannomutase

enzyme (130), a GNAT acyltransferase (131), a diaminopimelate (DAP) epimerase (132), and a lysozyme-

like enzyme (133). In a distinct operon, YfiM-DUF2279 is combined with a GT-A glycosyltransferase

domain (79), a further OMP β-barrel, and a secreted PDZ-like domain fused to a ClpP-like serine

protease (134,135) (Figure 5H).

The strong linkage to the OMP β-barrel, together with their localization prediction, suggests that these

Griddle and YfiM-DUF2279 Lipocone domains operate in the outer membrane. As with the inner

membrane lipids, they could potentially catalyze modifications of head groups through

transesterification and/or linkage/release of outer membrane-associated polysaccharide chains

through action on lipid-head group phosphoesters.

Lipocone domains acting on lipids in transit to the outer membrane

The ClaspCone-1 and ClaspCone-3 families lack the hydrophobicity indicative of direct residence in the

membrane (Figure 1C); instead, they are predicted to localize to the periplasmic space. In the

ClaspCone-1 family, the Lipocone domain is fused at the extreme N-terminus to either a single TM or a

5TM domain predicted to anchor it to the cell membrane. Between this TM element and the Lipocone

domain, we detected a previously uncharacterized version of the Tubular lipid binding protein (TULIP)

domain (136,137) or an Ig-like and a Zincin-like metallopeptidase (MPTase) domain (138) (Figures 5I, 6).

These ClaspCone-1 genes may also show operonic associations with genes encoding a lipase of the

SGNH family (139) and a membrane-bound O-acyltransferase (MBOAT; Figure 5I, Supplementary Data)

(140). The TULIP domain superfamily has recently been characterized as a lipid-binding domain

(136,137), which in proteobacteria functions in outer membrane lipid transport (141,142). Thus, we

propose that the ClaspCone-1 family is likely to act in the periplasmic space on the head groups of

outer-membrane targeted lipids bound to the TULIP or Ig-like domains.

A Lipocone domain catalyzing a predicted lipoprotein lipid linkage reaction

The Skillet-1 Lipocone is strongly coupled in an operon with a downstream gene coding for a protein

with an unusual lipobox-like sequence followed by one of several extracellular domains (e.g.,

concanavalin, β-jelly roll, OB-fold, Ig-like, β-propeller) predicted to bind carbohydrates or other ligands

(143-147) (Figures 5J,6,S5 Supplementary Data). The lipobox-like sequence features a conserved GS

motif at its C-terminus instead of the usual GC of the classic lipobox of bacterial lipoproteins (148)

(Figure S7). In the canonical lipoprotein processing pathway, a thioether linkage is formed between the

sulfhydryl of the cysteine and a diacylglycerol lipid embedded in the inner membrane by the

lipoprotein diacylglyceryl transferase (lgt) enzyme, followed by the cleavage of the signal peptide at

the GC motif junction by the signal peptidase (149,150). Given the serine in place of the cysteine in

these lipobox-like sequences, we propose that it undergoes non-canonical lipidation by the associated

Skillet-1 Lipocone protein in lieu of lgt. We propose that comparable to PTDSS1/2, which act on free

serine, the Skillet-1 family links the conserved serine from the lipobox-like sequence to a phospholipid

(Figure 3A,D).

Lipocone domains in predicted lipid-associated signaling systems

Systems defined by standalone proteins with Lipocone domains

Several representatives of the two VanZ and Skillet-3 families are fused to a diverse array of known or

predicted extracellular ligand-binding domains (Figure 5K), where the architecture takes the form of

SP+X+TM+Lipocone or Lipocone+TM+X, where 'X' is the extracellular ligand-binding domain and SP is a

signal peptide. The ligand binding domains include: (i) carbohydrate-binding lectin domains such as

jelly-roll, concanavalin-like, NPCBM-like, CBD9-like, and other β-sandwiches (143,144,151-153)); (ii) a

lipid-binding helix-grip superfamily domain (154)); (iii) those binding other potential ligands (e.g., Ig,

OB-fold, YycI-like, DUF498-like, PepSY-like, β-helix, TPR, MORN, and β-propeller repeats (145-

147,153,155-158)) (Figure 5K, Supplementary Data). We interpret these architectures as implying

signaling, wherein the binding of the cognate ligand by one of the above domains regulates the

catalytic activity of the associated Lipocone domain. Among these, the extracellular domains fused to

the Skillet-3 family are particularly notable for their extreme variability (Figure 5K). This suggests their

diversification under an arms race scenario (also see below) in a biological conflict. Further, the genes

coding for the above are sporadically associated with exopolysaccharide metabolism genes

(Supplementary Data). Hence, it is conceivable that this signaling is associated with exopolysaccharide

variation (e.g., O-antigen phase-variation (159,160)), which might play a role in evading bacteriophage

attachment.

Additionally, VanZ-1 Lipocone domains are also fused to several known signaling domains confidently

predicted to reside in the cytoplasm, including the cyclic nucleotide-binding domain (cNMPBD),

phosphopeptide-binding FHA, and DNA-binding RHH and HTH domains (65,161-163) (Figure 5K). These

associations suggest potential VanZ regulation via a cytoplasmic cyclic nucleotide (sensed by cNMPBD)

or, conversely, VanZ acting as an allosteric regulator of a transcriptional program via the HTH or RHH

domain. One of the most common yet enigmatic fusions to VanZ is with the integral membrane RDD

domain (53). The role of this domain is unknown; however, our analysis indicates a potential active site

oriented towards the cytoplasmic face of the membrane (Nicastro GN, Burroughs AM, Aravind L,

unpublished observations). The VanZ-RDD fusion is sometimes further fused to other domains (Figure

5K), the most notable being a highly derived but active novel Histidine Kinase domain (Figure 5K).

These associations again point to the coupling of lipid modification with a signaling event on the

cytoplasmic face of the membrane, which might relate to the dynamic regulation of lipid-carrier-bound

exopolysaccharide precursors.

Multi-component associations of the Lipocone proteins in signaling

These systems resemble the above-discussed versions but are encoded by conserved gene

neighborhoods that separate the Lipocone and the signaling elements (typically predicted transcription

regulators) into distinct components. Our analysis recovered at least three such systems: 1) A VanZ-1

Lipocone in the recently described HAAS/PadR-HTH two-component systems, which sometimes

replace classical Histidine kinase-Receiver two-component systems (164). In these systems, the

detection of an extracellular or intramembrane stimulus by a sensor domain releases the PadR-HTH

transcription regulator bound to the sensor-fused HAAS domain. Here, VanZ-1 occupies the sensor

position (Figure 5L). 2) A Skillet-2 Lipocone is coupled in a core two-gene system to a conserved

upstream gene (Figure 5L,S4). That gene encodes a single TM protein with either a zinc ribbon (ZnR)

fused to a conserved helix or an HTH domain fused to a ClpS-ribosomal L7/L12 domain in its

cytoplasmic region (165). These neighborhoods might also code for an HMG-CoA reductase and GHMP

kinase that catalyze successive reactions in the production of phosphomevalonate, a precursor of

isoprenoid lipids (166,167). 3) A Skillet-DUF2809 Lipocone protein is operonically coupled with a 6TM

protein and a further predicted transcription factor with a wHTH protein. These operons are further

elaborated via additional embedded and flanking genes either coding for components of isoprenoid

lipid (e.g., undecaprenol) (168,169) or exopolysaccharide (e.g., ECA and related polysaccharides)

metabolism (94,97) (Figures 5L,6,S5 Supplementary Data).

The Lipocone domains in these systems are predicted to be active enzymes, which, together with their

operonic associations, points to functions involving the modification or transesterification of

isoprenoid lipid head groups, sometimes in the context of exopolysaccharide biosynthesis. However,

their associations with the intracellular HTH domains suggest that the Lipocone enzymatic activity is

potentially coupled with the transcriptional regulation of the production of precursors of the lipids or

exopolysaccharides. Given the high variability in the associated genes related to

exopolysaccharide/lipopolysaccharide biosynthesis, we anticipate that the associated transcriptional

regulation potentially responds to environmental stress, phages, predatory bacteria attacks, or host

immune response.

Lipocone domains as effectors in biological conflicts

Lipocone domains in antiviral immunity

The Min-Wnt domains that we originally identified were predicted to play a role in biological conflicts

with invasive selfish elements such as viruses (24). In this work, we better explain their potential

mechanism of action. These versions show no fusions to extracellular domains or secretory signals,

suggesting that they act as soluble versions within cells – a proposal also backed by their overall

hydrophobicity (Figure 1C). These Min-Wnts are typically fused to the DUF3892, which displays a fold

characterized by a three-stranded meander followed by a helix also seen in the dsRNA-binding domain

and the ribosome hibernation factors (HPF) (170,171) (Figure 5M). Hence, we propose that these

versions might potentially act to sense virally induced RNAs or modified ribosomes (24) to trigger a

dormancy or suicide response to limit viral infection via the Min-Wnt effector. Specifically, the Min-

Wnt might attack peptidoglycan precursors, such as lipid II, prior to their 'flipping' to restrict cell wall

synthesis (85,172,173) or other carrier lipids.

One other Min-Wnt domain N-terminally fused to a three-stranded β-meander is pervasive in the

Bacteroidetes clade. This is operonically coupled with genes encoding a TM-linked run of pentapeptide

repeats and two structurally distinct, secreted glycosyl hydrolase enzymes, respectively, containing a

TIM barrel domain and a run of β-helix repeats (Figure 5M, Supplementary Data). Further,

cyanobacteria show a standalone prok-TelC domain without any secretory signals. These could again

act as cell wall or membrane-lipid targeting effectors in response to intracellular invaders or stress

(Figure 5M). Interestingly, some tailed bacteriophages also code for intracellular Min-Wnt domains,

suggesting that they might also be deployed on the virus side in biological conflicts (Supplementary

Data).

Lipocones as toxin domains in polymorphic and allied conflict systems

Polymorphic toxins and related systems, widespread across bacteria and certain archaea, are

characterized by a highly variable C-terminal toxin domain ("toxin tip") that is preceded by a range of

more conserved domains typically required for autoproteolytic processing of the toxin, its packaging

and trafficking (e.g., RHS repeats), adhesion and secretion via one of several secretory systems

(174,175). The toxin might be delivered via one of the secretory systems into a target cell or else via

direct contact between interacting cells. Classical polymorphic toxins are usually involved in kin

discrimination and are accompanied by genomically linked cognate immunity proteins that protect

against self-intoxication (174,176). Keeping with the principle of effector sharing between systems

involved in distinct types of biological conflicts, we had originally identified the Min-Wnt domain

closely related to those described in the above subsection as a toxin tip in polymorphic toxin systems

(24). In the current work, we extend these findings to show that several distinct Lipocone families have

been independently recruited as toxin tips of polymorphic toxins and related systems, namely Min-

Wnt, pro-SAA, prok-TelC, CapCone-1, CapCone-2, ClaspCone-2 and VanZ-1 (Figures 4,5N,6,S6).

Certain CapCone-2 and Min-Wnt toxins define some of the simplest of these toxin systems that are

common in Gram-positive bacteria. Here, a standalone Lipocone domain is coupled to a signal peptide

or lipobox via a poorly structured linker. These are usually encoded in a two-gene configuration with

their cognate immunity protein (Figure 5O). More complex versions present, in addition to adhesion,

peptidoglycan-binding, lipid-binding and proteolytic processing domains, multiple hallmarks of

specific secretion systems. These include T4SS (VirD4-binding domain), T6SS (PAAR domain), T7SS

(WXG/LXG domain), T9SS, and MuF domains indicating that these toxins are delivered via diverse

secretion systems (174,176) (Figure 5P, Supplementary Data). Additionally, we recovered standalone

CapCone-1 domains encoded in an operon with a PsbP/MOG1-like domain diagnostic of the T6SS

(174,177) (Figure 5P), which predicts that these are likely toxins secreted via this system. Further, we

also found Min-Wnt domains fused to the N-termini of RTX-like β-roll repeats, suggestive of a T1SS-

mediated toxin export (178) (Figure 5P).

Our analysis also uncovered multiple, previously uncharacterized trafficking/packaging systems

associated with different Lipocone polymorphic toxins. Several Min-Wnt and CapCone-1 domains with

lipoboxes are fused to an N-terminal Cystatin-like superfamily domain (179) (Figure 5Q). The same

domain is also comparably fused to several other C-terminal toxin domains in related organisms, some

of which are also predicted to target lipid head groups: (i) a novel toxin domain we unified with the

lipid-targeting Colicin M fold (180); (ii) a lipid-binding START-domain-like helix-grip fold domain (154);

(iii) a papain-like fold fatty acyltransferase (181); (iv) a domain related to the VanY-like D-Ala-D-Ala

carboxypeptidase (182) (Figure 5Q, Supplementary Data). In all these cases, the toxins are coupled to a

related immunity protein (see below), suggesting that they define a distinct polymorphic toxin system.

We propose that this Cystatin-like domain defines a novel packaging or deployment system upon

secretion for the C-terminal toxin domain analogous to Cystatin domains in functioning with

eukaryotic proteases (183). The prok-TelC family Lipocones are found in distinctive architectures in

two poorly characterized, predicted polymorphic toxin systems. In one of them, they are fused to N-

terminal glucan-binding GbpC β-sandwich domain (184) and repeats of MucBP-like Ig domains (185),

which might anchor them to exopolysaccharides (Figures 5Q,S6 Supplementary Data). The second

variant found in association with T9SS components (186) shows fusions to one or more copies of a

previously undetected TPM domain (Figure 5Q). While the domain has been claimed to be a

phosphatase (187), our recent analysis indicates that this is unlikely to be the case (164). Instead, we

propose that the TPM domain might assist in the assembly of membrane-linked protein complexes, a

role that might be relevant to the trafficking of these toxins (164).

To date, the only experimentally characterized Lipocone domain from polymorphic toxins is of the

prok-TelC family, which are secreted via T7SS (29,188) (Figures 5P,6). Notably, prok-TelC has been

shown to be active only outside the cell and not in the cytoplasm (29). As noted above, it attacks lipid II

to cleave off the peptide-linked disaccharide pyrophosphate head group from the undecaprenol tail

(Figure 3B). Prok-TelC has also been speculated to similarly attack WTA-lipid II linkages (29). These

findings provide a template for other Lipocone superfamily effectors in potentially targeting lipid

carrier linkages in peptidoglycan and exopolysaccharide intermediates. However, given the diversity

within the family, it is conceivable that they also target other lipids (Figure 3F).

Immunity proteins of Lipocone polymorphic toxins indicate periplasmic/intramembrane action

To date, only a single immunity protein has been reported for Lipocone toxins, viz., TipC, which

counters the prok-TelC toxin in the periplasm (29,189) (Figure 5P,6). Here, we uncovered a range of

immunity proteins belonging to structurally distinct folds that counter the remaining Lipocone toxins

(Figures 5N-Q,S8, Supplementary Data). The most widespread of these is a rapidly evolving, membrane-

anchored member of the BamE-like superfamily that associates with not only Min-Wnt and CapCone

toxins but also other above-mentioned lipid-head-group targeting toxins (e.g., the novel Colicin M-like

domain). The BamE-like fold features a core two-helix hairpin followed by a run of three β-strands

(Figure S8). The classical BamE operates in a pathway for the assembly of OMP β-barrels (190,191),

suggesting that these immunity proteins were recruited from an ancestral BamE and, like it, function in

the periplasm. Additional candidate immunity proteins with more restricted phyletic spreads include

(Figure S8): (i) a β-jelly-roll fold-containing protein (144); (ii) an integral membrane protein with a 4-

TM core. These two are observed with Min-Wnt toxins. (iii) A novel domain combining an α-helix with

a run of 4 β-strands stabilized by four absolutely conserved cysteine residues. This is coupled to both

Min-Wnt and prok-SAA toxins; (iv) a protein with an OB-fold domain (145) (v) a protein with a β-

sandwich related to the eukaryotic centriolar assembly SAS-6 N-terminal domain (192). The last two are

coupled to CapCone-2 toxins (Supplementary Data). Notably, despite their structural diversity, these

immunity proteins are all TM or lipoproteins and, like TipC (29,189), are predicted to operate at the

membrane or in the periplasm (Figure 5N-Q, Supplementary Data). This suggests that they intercept

their cognate Lipocone toxin domain outside cells or in the membrane rather than within the cell.

Lipocone toxins in predator-prey and other inter-specific conflicts

In contrast to polymorphic toxins, which are typically deployed in intra-specific conflict between

competing strains of the same species, other toxin systems are deployed against more distantly related

target cells, such as prey and eukaryotic hosts (193). While some of these closely parallel polymorphic

toxins in their domain architecture, they are usually distinguished by the lack of an accompanying

immunity protein. The simplest of these systems are secreted Min-Wnt proteins from bacteria and

fungi. These present just a standalone Min-Wnt domain or one fused to a novel domain with a half β-

barrel wrapping around a helix (Figure 5R, Supplementary Data). These are probably deployed as

diffusible toxins that target rival organisms in the environment.

Another architectural theme is defined by Min-Wnt and prok-SAA Lipocones fused to an enigmatic,

novel, short C-terminal domain, which is comprised of a long β-hairpin with a characteristic "break" in

its central region, causing it to acquire an arch-like appearance (Figures 5S, S9). Hence, we refer to this

domain as the broken-hairpin. We found the broken-hairpin domain to be fused to a wide array of

predicted toxin domains across the bacterial superkingdom. These include effector domains found in

polymorphic toxin and allied systems that target peptidoglycan, carrier lipids and the membrane, such

as members of the Colicin M (180), Zeta toxin-kinase (194), lysozyme (195), an α/β-hydrolase

superfamilies (110) and nuclease toxins such as members of the HNH, HipA, SNase, and BECR

superfamilies (174) (Figures 5S, S9C). Remarkably, these proteins with the broken-hairpin tend to lack a

signal peptide or association with any other secretion system or immunity proteins (Figure 5S,

Supplemental Material). Hence, we propose that the broken-hairpin domain itself serves as a trafficking

mechanism for the externalization of these toxins in conflicts with rival environmental organisms.

Some predicted secreted Lipocones are found predominantly in predatory bacteria. The first of these

are CapCone-2 domains from lineages like Bdellovibrionota, which are encoded in two-gene systems,

with the second gene coding for a further secreted effector such as an α/β-hydrolase, Patatin, or

acyltransferase or an OMP β-barrel domain (110,116,117,123) (Figures 5T,S6). Secreted prok-SAA

domains from myxobacteria and some other lineages are fused to a Zincin-like metallopeptidase

domain at the N-terminal, and the first bacterial example of the von Willebrand Factor D (vWD) and Ig

domains at the C-terminus (196) (Figure 5T, Supplemental Material). In the recently described

predatory Patescibacterial branch of Omnitrophota species, Skillet-clade Lipocone domains are found

in gigantic proteins combined with several other domains and TM segments. Domains found in these

proteins include polysaccharide biosynthesis enzymes (94,97), signaling proteins involved in Histidine

kinase-Receiver relays (197), peptidases of the MPTase and Papain-like superfamily (113,138), diverse

methylases, and extracellular ligand-binding domains like the peptidoglycan-binding LysM domain

(198) (Figures 5T,6). Given the concentration of the above systems in predatory bacteria

(Supplementary Data), we posit that the above Lipocones might function as toxins targeting prey

membranes alongside a battery of effectors targeting other cell surface components. In particular, the

CapCone-2 systems are likely to target outer membranes. In animals, the vWD domain is involved in

adhesion (199) and might play a similar role in adhering to prey cells in the bacterial Prok-SAA toxins,

while the MPTase in these proteins potentially releases the toxin through autoproteolysis. Finally, the

giant proteins from the Patescibacteria are likely to combine signaling of prey presence with the

overcoming of prey defenses and targeting of membranes.

Certain prok-TelC proteins are observed as part of several distinctive systems that could be involved in

as-yet-undiscovered predatory interactions or in targeting environmental competitors. One such,

defined by large proteins from spore-forming Bacillota, combines a diversifying set of extracellular

ligand-binding domains (e.g., Ig-like, Cell-wall-binding β-hairpins and β-propellers (146,147,200)) with a

two-enzyme core formed by a prok-TelC and a N-acetylglucosamine (GlcNAc)-1-phosphodiester alpha-

N-acetylglucosaminidase (NAGPA). NAGPA catalyzes phosphoric-diester hydrolysis to release

phosphodiester-linked sugars (Figures 5U,6,S6) (201). Some of these proteins feature an additional

NlpC/p60 superfamily peptidase domain predicted to target peptidoglycan (181). The recombinational

diversity of ligand-binding domains in this system, even among closely related Bacillota species,

supports a possible arms race and involvement in a biological conflict. Other TelC domains in some

Bacillota, Actinomycetota, and fungi are fused to peptidoglycan-binding domains (PGBD) (202) and an

Rv2525c-like TIM-barrel (203) (Figures 5U,S6). In Actinomycetota, this protein is further combined in

operons with either of two mutually exclusive genes coding for rapidly evolving proteins (Figure 5U):

(i) a secreted protein containing a pair of Ig domains (200); (ii) a 3-TM protein (3TM-CCDN) with two

conserved cysteines, an aspartate and asparagine residues predicted to be located between the TM

segments outside the cell. This version is further coupled to a gene for a secreted VanY superfamily

peptidase (182) (Figure 5U, Supplementary Data). Common to these contexts are rapidly evolving and

variable domains on the one hand and peptidoglycan/exopolysaccharide binding or degrading domains

on the other. Hence, we interpret these as potential conflict systems that engage the cell wall and

target it and associated membranes in rival bacteria.

Lipocone domains in resistance to antimicrobial agents

VanZ-1 proteins were initially identified as encoded by a gene linked to that coding for the VanY D-

alanyl-D-alanine carboxypeptidase involved in resistance to glycopeptide antibiotics like vancomycin

and teicoplanin (31,204-206) (Figure 6). These antibiotics bind the terminal D-Ala-D-Ala in the peptide

moiety of peptidoglycan, preventing the transpeptidase cross-linking reaction necessary for its

maturation. Upon detection of these antibiotics, enzymes encoded by the core vancomycin resistance

operon re-engineer the exported peptidoglycan by inserting a D-Ala-D-Lac in place of the D-Ala-D-Ala

linkage, precluding antibiotic binding (53). The VanY peptidase, while not strictly required for

antibiotic resistance, acts as an accessory to this system by cleaving any remaining D-Ala-D-Ala

linkages generated via the canonical pathway (53,205). However, the role of VanZ in this system has so

far remained unknown. While only a small fraction of the VanZ-1 genes are found in these antibiotic

resistance contexts (Supplementary Data), interestingly, other Lipocone genes, namely those of the

VanZ-2 and the Skillet families, might also be linked to VanY in lieu of VanZ-1. Further, VanY might be

replaced by a structurally unrelated secreted D-Ala-D-Ala carboxypeptidase of the metallo-beta-

lactamase fold (111) in operonic contexts with VanZ-1 (Figure 5V). Hence, given our above prediction

regarding VanZ acting in peptidoglycan and/or exopolysaccharide metabolism, VanZ-1 and the

Lipocones displacing it might indeed play an accessory role with VanY at the membrane (204,205) in

antibiotic resistance. We posit that, in these contexts, it likely acts on the head group of Lipid II to

recycle canonical peptidoglycan intermediates for their accelerated or more thorough replacement

with the resistant versions (Figure 3G).

We also identified a conserved five-gene operon featuring a YfiM-1 family Lipocone that might play a

role in resistance to antibacterial agents (Figures 5W,S4). Other than YfiM-1, this operon contains genes

for: (i) a thioredoxin domain protein (109); (ii) A DTW clade RNA modifying enzymes of the SPOUT

superfamily (207,208); (iii) a protein with acyl-CoA ligase, GNAT superfamily N-acetyltransferase and

ATP-grasp domains (131,209,210); (iv) a PssA-like phosphatidylserine synthetase of the HKD

superfamily (211) (Figure 5W, Supplementary Data). Of these enzymes, the phosphatidylserine

synthetase is predicted to act in its usual capacity to generate a lipid with a serine head group (211). We

propose that this would then function as a substrate for the YfiM-1 Lipocone domain, which might

exchange the serine for another moiety via a reaction paralleling PTDSS1/2 (Figure 3A). This moiety

could then be modified by aminoacylation, further acylation and a redox modification by the third

protein listed above, together with the thioredoxin. Indeed, such peptide modifications of lipid head

groups by lysine, alanine, or arginine aminoacylation catalyzed by derived tRNA synthetases fused to

GNATs have been shown to be a key resistance mechanism against breaching of the membrane by

antibacterial peptides (212,213). Hence, we predict the modifications catalyzed by this system might

play a comparable role. The presence of a tRNA-modifying DTW domain suggests that in parallel to the

tRNA synthetases, the GNAT in this system might use a tRNA-linked acyl group as a substrate, as seen in

peptidoglycan biosynthesis (214,215).

Eukaryotic recruitments of the Lipocone superfamily

Lipocone domains have been transferred on several occasions from bacteria to eukaryotes (Figures

4,S1). While there is predicted functional overlap with the above-described, predominantly bacterial

versions, we discuss these separately as the inferred biological contexts of their deployment are often

distinct from the above.

Plant YfiM-1 and eukaryotic VanZ-2 proteins

A conserved YfiM-1 family protein typified by the *Arabidopsis* AT1G15900 was acquired from the

bacteroidetes lineage of bacteria at the base of the plant lineage prior to the chlorophyte-streptophyte

(including land plants) split and is predicted to be catalytically active (Figure S2, Supplementary Data).

In *Arabidopsis*, this gene is widely expressed across different tissue types, developmental stages, and

other tested conditions (216,217). Given the above-predicted roles for bacterial YfiM-1 proteins, it is

conceivable that the plant version plays a comparable role in the metabolism of a conserved plant-

specific lipid. In a similar vein, a distinct clade of standalone VanZ-2 domains typified by the

Saccharomyces cerevisiae YJR112W-A was acquired early in the fungal lineage. A similar transfer is also

seen in the SAR clade of eukaryotes (Figure S1). Since these eukaryotes lack peptidoglycan and other

bacterial-type isoprenoid lipid-borne exopolysaccharide intermediates, we suggest that this version

was recruited for modifications of a fungus-specific lipid (e.g., highly oxygenated isoprenoid lipids)

(218).

The Met-TelC proteins

The Met-TelC clade is comprised of versions of the TelC family with a reconfigured active site

transferred from bacteria to Metazoa prior to the divergence of the cnidarians, and most members are

predicted to be catalytically inactive (Figure 2). In cnidarians and arthropods, the Met-TelC domain is

found in a secreted protein fused to C-terminal adhesion-related vWA (219) and Ig domains, followed by

a TM helix (Figure 5X, Supplementary Data). The chordate version, typified by human PGLYRP2 (220), is

also secreted and is fused to a C-terminal Amidase targeting the N-acetylmuramoyl-L-alanine linkage

(Figure 5X, Supplementary Data). The chordate version is a key innate immunity factor against

bacterial pathogens that degrades sugar-peptide linkages in peptidoglycan via the Amidase domain

(221-223). As most Met-TelC proteins lack the active site residues but are modeled to retain the

substrate-binding pocket, we propose that they participate in anti-bacterial immunity as a Pathogen-

Associated Molecular Pattern (PAMP) receptor (224). Specifically, they could recognize polyisoprenoid

pyrophosphate-linkage-containing lipid intermediates of bacterial cell-surface molecules like

peptidoglycan or exopolysaccharides.

Eukaryotic Wnt proteins

Wnt family Lipocones were transferred on multiple occasions to eukaryotes. The best known of these

are Met-Wnt proteins, which were acquired from bacteria at the base of Metazoa after they had

separated from their closest sister group, the choanoflagellates. These lost the ancestral active site

residues and function as well-studied secreted signaling molecules and will not be detailed further in

this work (for review, see (1,225)). Independently of the Met-Wnt proteins, catalytically active, secreted

versions closely related to the bacterial Min-Wnt proteins were transferred to fungi and, within

Metazoa, to the rotifers and the hemichordate acorn worm *Saccoglossus kowalevskii*, where they are

lineage-specifically expanded (Figure S1, Supplementary Data). These versions are primarily standalone

versions of the Min-Wnt domain lacking the large inserts typical of the Met-Wnt proteins. We predict

that these eukaryotic Min-Wnt proteins retain their ancestral role and might participate in anti-

bacterial immunity in the lineages coding for them.

Met-SAA proteins

Met-SAA proteins were acquired from bacteria prior to the divergence of the cnidarians from the rest

of Metazoa. However, unlike the Met-Wnt and Met-TelC proteins, they often conserve the ancestral

active site residues indicating that they are usually enzymatically active (Figure 3). Human SAA has

been recognized as a key immune marker that dramatically increases in blood during the Acute Phase

Response (226). It has been reported to bind the *E. coli* outer membrane protein OmpA (227) and claimed

to function as an opsonin in innate immunity (228). Like Met-TelC, but in contrast to Met-Wnts, Met-

SAAs appear to have been lost or pseudogenized in several animal lineages (229-231) (Figure 3). This is

consistent with an arms-race scenario in immunity and the development of pathogen resistance

against the Met-SAAs, leading to loss. Keeping with an immune role for the Met-SAAs, we propose a

catalytic function for the active versions in severing lipid head groups of outer-membrane lipids or of

isoprenoid lipid carrier intermediates. Such action could also generate PAMPs that could explain the

activation of neutrophil- and macrophage-based immunity by SAA (228). Pertinent to these

observations, diverse OMP β-barrels have been linked to the translocation of polymorphic toxin

domains across the outer membrane of target cells (232-234). Given this and the origin of Met-SAA from

bacterial polymorphic toxin-related systems (Figure 4), its interaction with OmpA might help it cross

over into the periplasmic space and act on maturing peptidoglycan or teichoic acid intermediates.

SAA was first reported as a component of secondary amyloid deposits (235), and its capacity to form

amyloid fibrils upon protease cleavage was theorized as a potential PAMP activating the immune

response (30). Indeed, bacteria produce their own secreted amyloids, such as Curli and Fap, believed to

contribute to biofilm formation (236,237), and might be PAMPs recognized by animal immune systems

(238). Further, other animal amyloids, such as the β-amyloid, have been proposed to play a role as

physical barriers in immunity against bacteria (239). Thus, amyloid formation by protease cleavage

(including potentially by bacterial proteases) may represent a second line of defense mediated by Met-

SAA proteins.

DISCUSSION

Early Evolution of the Lipocone superfamily

No single well-defined Lipocone clade is universally conserved across the three superkingdoms of Life

(Figures 4, S1). However, the VanZ and Wok clades are both found across all major bacterial phyla

(notwithstanding sporadic losses in certain lineages) and in some archaeal lineages (Figure S1). At the

same time, the cpCone clade is found across most major archaeal lineages and is nearly universally

conserved in the eukaryotes (absent in Ascomycota and some choanoflagellates) (Figure S1). Notably,

the cpCone and Wok clades tend to group together in the profile-profile similarity network (Figure 1B).

These observations suggest that at least a single version of the Lipocone superfamily was likely present

in the Last Universal Common Ancestor (LUCA). The phyletic patterns suggest that the LUCA Lipocone

gave rise to the VanZ/Wok precursor in the bacterial lineage on the one hand and the cpCONE clade via

a circular permutation event in the archaeo-eukaryotic lineage on the other (Figure 4). Based on the

features of these deep-branching clades, the LUCA version is inferred to feature a hydrophobic domain

with a 4TM helix core, with the active site facing the outer leaf of the lipid bilayer (Figure 1C). Given

that extant versions operate both on classic phospholipids and isoprenoid lipids, it is difficult to infer

which of these might have been substrates for the LUCA version. It is not impossible that this early

version had a generic specificity that became specialized in the descendant clades.

Subsequent diversification of the Lipocone domain

The early diversification of the Lipocone domain appears to have had different drivers in the two

prokaryotic superkingdoms. The presence of an extensive repertoire of exopolysaccharides in the cell

wall (peptidoglycan, teichoic acids), cell surface (e.g., ECA), and outer membrane (e.g.,

lipopolysaccharide), synthesized via isoprenoid lipid-linked intermediates, like lipid-II, was the primary

driver in the bacterial superkingdom (240). Here, this diversification yielded 4 monophyletic groups:

the VanZs, Wok, YfiM/Griddle-like and Skillet (Figure 4). The deeper VanZ and Wok branches, which

were recruited first for lipid-II-related functions, were likely the predecessors of the more restricted

bacterial families with specialized functions. For instance, the emergence of the outer membrane in

certain bacteria was potentially coupled with the origin of the YfiM and Griddle clades (Figure 4).

Similarly, our predictions suggest that within these clades, further diversification accompanied the

acquisition of specialized functional roles in antibiotic resistance (Skillet-DUF2809, YfiM-1), secondary

sensor roles in single and multicomponent signaling (VanZ and Skillet families) and lipoprotein

processing (Skillet-1). The interoperability of Lipocone domains on lipid carriers shared across different

biosynthetic pathways (see above, Figures 3,S4-S5) appears to have been a key factor leading to this

versatility.

In the ancestral archaeo-eukaryotic lineage, the absence of peptidoglycan and an apparently lower

diversity of structures with exopolysaccharides was reflected in the lesser diversification of the

Lipocone clades (Figure 4). There are open questions regarding the biochemical functions of the

primary archaeo-eukaryotic Lipocone clade, the cpCONE. Although the eukaryotic cpCone PTDSS1/2

family has been shown to swap serine for ethanolamine or choline in lipid head groups (28,57), their

archaeal counterparts remain uncharacterized. Archaea have their own lipid with a serine in the head

group (archaeophosphatidyserine), but to date, its synthesis has been shown to depend on a patchwork

of different CDP-alcohol phosphatidyltransferase enzymes (CaPs) in different archaeal species

(58,241,242). While the CaPs are also integral membrane enzymes with a 6TM helix core catalyzing

similar reactions as the Lipocones on lipid head groups in archaea and eukaryotes (242), they are

evolutionarily unrelated domains. Nevertheless, we suggest that the archaeal cpCones, like their

eukaryotic counterparts, could contribute to distinct, as yet uncharacterized, pathways for the

generation of cell membrane phospholipids like archaeophosphatidylserine or those with other head

groups under special conditions.

Emergence of soluble versions of the Lipocone domain and their repeated recruitment in

biological conflicts

One of the remarkable aspects of the Lipocone superfamily is the loss of ancestral hydrophobicity in

several families (Figure 1C), transforming them from integral membrane proteins to soluble domains.

While unexpected, such a transition in integral membrane enzymes acting on lipid substrates is not

unprecedented. The PAP2 superfamily of integral membrane enzymes (e.g., diacylglycerol diphosphate

phosphatase) (124,243) also contains several soluble versions (244) that appear to have emerged from

an integral membrane ancestor (AMB and LA, unpublished observations). Most of the soluble Lipocone

domains retain their active site conservation (Figure 2) and, at least in one experimentally

characterized case, catalyze a comparable reaction as the TM version (29) (Figure 3B). The weight of the

evidence presented here, including the profile-profile similarity network (Figure 1B), phyletic patterns

(Figure S1), functional contexts (Figure 5-6), and the broadly shared structural features (Figures 1A, S2),

suggests that the loss of hydrophobicity occurred on a single occasion in the Lipocone superfamily,

followed by diversification of these soluble versions.

Our analysis of the soluble Lipocone families reveals repeated recruitment as toxins/effectors in anti-

viral and polymorphic toxin and allied systems (174), suggesting that their diversification was driven

by the arms races arising from the biological conflicts where they are deployed. Recruitment of a

representative of the VanZ-1 family as a polymorphic toxin on rare occasions (Figures 5N,6) suggests a

possible evolutionary pathway by which this might have happened: the soluble effector version of

Lipocones attacking lipids in competing bacteria likely emerged from an ancestral version that

catalyzed endogenous lipid-head-group modifications on the same lipids in metabolic pathways. Once

soluble versions emerged through reduced hydrophobicity, they could be deployed as diffusible

effectors that were shared across extracellular and intracellular conflict systems, a trend previously

recognized in many other effector domains (25).

Repeated acquisition of Lipocones of bacterial origin by eukaryotes

Unlike bacteria, eukaryotes as a whole do not possess a rich repertoire of Lipocone domains. The

PTDSS1/2 family, vertically inherited from the archaeal progenitor, is the only version that can be

inferred as being present in the Last Eukaryotic Common Ancestor (Figure 4). However, as noted above,

particular eukaryotic lineages have acquired and fixed distinct Lipocone families of ultimately bacterial

provenance: (i) YfiM-1 in the plant lineage; (ii) the fungal VanZ-2 domains typified by the *Saccharomyces*

cerevisiae YJR112W-A; (iii) Met-Wnt (discussed further below) (Figure S1). The fixation of these versions

early in the history of the eukaryotic lineages possessing them suggests that they were recruited for

"housekeeping" or developmental roles definitive of the respective lineages.

Beyond these, the fungal and metazoan lineages show more sporadically distributed versions, which

have all been acquired from bacterial secreted-toxin or antiviral systems: (i) Min-Wnt independently in

fungi and certain Metazoa; (ii) SAA; (iii) TelC; the latter two are absent in the basal-most metazoans, the

sponges, but are present in Cnidaria, suggesting a relatively early acquisition (Figure 4). The weight of

the evidence suggests that they have retained certain aspects of the ancestral bacterial effector

function for anti-pathogen immunity in eukaryotes. This is consistent with both their episodic loss and

lineage-specific expansion, the tendency to show rapid sequence divergence and, in the latter family,

loss of catalytic activity (Figures 2,S1, Supplementary Data).

This independent acquisition of at least 3 distinct Lipocone families in metazoan immunity from

polymorphic and allied effector systems of prokaryotes points to a persistent evolutionary trend.

Notably, the Lipocone domains participating in animal immunity have been drawn from secreted

effectors rather than the intracellular versions (bacterial intracellular Min-Wnts) predicted to

participate in bacterial anti-selfish element immunity. More generally, this adds to a growing list of

components drawn from secreted effector systems of prokaryotes in eukaryotic immune systems

(174,193,245). For example, this closely parallels another structurally unrelated effector domain, the Zn-

dependent deaminase (e.g., metazoan AID/APOBEC deaminases) (246). Hence, these observations add

further support to our hypothesis that the extensive expansion of effectors in diverse prokaryotic

inter-organismal conflict systems served as a reservoir from which eukaryotic immune systems

repeatedly acquired components (193,245). We propose that symbiotic associations between the early

animals and bacteria resulted in potential interactions via secreted effectors of the latter that aided the

former against antagonistic bacteria. This probably led to their eventual acquisition by animals and

incorporation into their immune processes.

Origin of Wnt as a signaling molecule

Earlier considerations on the evolution of Wnt signaling indicated that it emerged at the base of the

metazoan lineage and incorporated a wide range of components of different origins (e.g., the HMG

domain transcription factor TCF/LEF, the HEAT repeat protein β-catenin and the 7TM receptor Frizzled)

(1). However, the provenance of Met-Wnt itself had been mysterious and was seen as a possible example

of a metazoan innovation (225). While the Met-Wnt domains show Metazoa-specific elaborations (34),

its conserved core is a Lipocone domain (Figure 1A). We establish that the progenitor of Met-Wnt

emerged as part of the radiation of Lipocone domains in bacteria as effectors deployed in both

intracellular and inter-organismal conflict – the Min-Wnt proteins.

Whereas the prokaryotic Min-Wnt proteins are predicted to be secreted toxins targeting phosphate-

linked moieties in lipid head groups, the Met-Wnts underwent an ancestral inactivation through loss of

the catalytic residues (Figure 2). However, they retained their ancient involvement in cell-cell

interactions as secreted agents. The Met-Wnt residues recognized as essential for the receptor

(Frizzled) binding, including the absolutely conserved palmitoleoylated serine residue, are found in the

aforementioned Metazoa-specific hairpins and loops (34,36). However, despite their inactivation, the

Met-Wnts retain the ancestral substrate-binding pocket (Figures 1A, S2). This raises the possibility that

they might be involved in as-yet unexplored interactions with ligands such as lipids.

Our tracing of the provenance of Wnt back to an effector in secreted bacterial toxin systems adds it to a

growing list of components in metazoan signaling networks that have been acquired from such

systems. For instance, this is also the case with components of the other key metazoan signaling

pathway, Hedgehog (247). Here the Hedgehog protein itself contains an autoproteolytic HINT peptidase

domain that was likely drawn from a structurally and functionally cognate domain observed in

polymorphic toxin systems (174,247). Further, an intracellular component of the same signaling

pathway, Supressor of Fused (SuFu) was derived from a common immunity protein found in

polymorphic toxin systems (247). Similarly, the Teneurin/Odd Oz proteins mediating signaling in cell

migration, neuronal pathfinding, and fasciculation in Metazoa descended from a polymorphic toxin

protein with a C-terminal HNH endonuclease toxin tip (174). In a similar vein, the immunity protein of

certain CapCone toxins identified in this study might have given rise to the β-sandwich domain in the

eukaryotic centriolar assembly factor SAS-6. These observations suggest that in addition to immune

system components, interactions with symbiotic bacteria also potentially provided the progenitors of

components of eukaryotic signaling and cytoskeletal networks that were central to the emergence of

Metazoa as a clade of multicellular eukaryotes (248,249).

CONCLUSIONS

Using sensitive sequence and structure analysis, we unify a large, hitherto unrecognized superfamily of

enzymatic domains, the Lipocone. By combining analysis of the active site and the structure of the

Lipocone domain with contextual information from conserved gene-neighborhoods and domain

architectures, we present evidence that members of this superfamily target phosphate linkages in head

groups of both classical phospholipids and polyisoprenoid lipids. Specifically, they catalyze reactions

such as head group exchange or severing of the head group-diphosphate linkage from the

polyisoprenol. We present evidence that these activities have been recruited in a wide range of

biochemical contexts, including cell membrane lipid modification, metabolism of peptidoglycan and

exopolysaccharide lipid-carrier linked intermediates, lipoprotein modifications, bacterial outer

membrane modification, sensing of membrane-associated signals, effector activity in antiviral and

inter-organismal conflicts and resistance to antimicrobials. Further, catalytically inactive versions (e.g.,

Met-Wnt) have been recruited for signaling roles in Metazoa. We predict the catalytic activity and

potential biochemical pathways of numerous representatives for the first time, including some proteins

that have remained enigmatic for over two decades, like VanZ.

We identify three notable trends in Lipocone evolution. First, although we reconstruct the ancestral

member of the superfamily as being a 4TM integral membrane domain, a large monophyletic subset

underwent a dramatic loss of hydrophobicity, transforming them into diffusible versions, including the

Wnts and the SAAs (Figure 1C). Second, the superfamily expanded in two major functional niches in

bacteria, namely peptidoglycan/exopolysaccharide metabolism and effector domains of both secreted

toxins and immune systems (Figure 4). Finally, members of the Lipocone superfamily were acquired on

multiple occasions from bacteria by Metazoa and were reused in new functional contexts as signaling

messengers and immune factors (Figure 4).

Importantly, our predictions in this regard underscore that much remains unexplored in terms of

lineage-specific cell wall and membrane metabolism in prokaryotes. We present several testable

biochemical, functional hypotheses for the many poorly understood branches of the superfamily,

several of which are being recognized as enzymatic for the first time here. We hope this will also open

new avenues of research to fill key gaps in our understanding of lipid metabolism.

1004

METHODS

Sequence analysis

Sequence similarity searches were performed using PSI-BLAST (250) and JackHMMER (251) against the

NCBI non-redundant protein database (nr) (252) or a version clustered at 50% sequence identity (nr50).

The searches were initiated using the previously identified prokaryotic Wnt (24), with multiple rounds

of searches conducted, each using seeds collected from the preceding searches. Clusters based on

sequence similarity (percentage identity or bit-score) were generated using MMseqs (253). The

clustering parameters were adjusted according to specific goals, enabling redundancy removal, the

definition of homologous groups, and the creation of new profiles. Multiple sequence alignments (MSA)

were generated using the MAFFT program (254) with the local-pair algorithm, combined with the

parameters –maxiterate 3000, –op 1.5, and –ep 0.2, and were manually refined based on structural

superimpositions and profile-profile comparisons.

1017

Sequence similarity network analysis

The HHalign program (255) was used to perform profile-profile comparisons, with the resulting p-value

and e-value scores serving as edges for constructing a superfamily relationship network. These

networks were further analyzed using the Leiden community finding algorithm (37) to detect sub-

networks. Network visualization was performed using the R igraph (256) or Python networkX libraries

(257).

1024

Comparative genomics, domain identification, and phylogenetic analysis

Genomic neighborhoods were obtained from genomes available in the NCBI Genome database (252)

using in-house scripts written in Perl and Python. Conservation analysis of these genomic

neighborhoods was performed by clustering the protein products of neighboring genes. Domain

identification was conducted using a collection of HMMs and PSSMs maintained by the Aravind lab,

along with HMMs from the Pfam database (258), utilizing the RPSBLAST (259) and HMMSCAN (260)

programs. To further refine detection, domain identification was extended through remote homology

analysis using the HHpred (261) program, against profiles built from the Pfam (258) and PDB70 (262)

databases. Phylogenetic analyses were performed using FastTree (263) and iqTREE2 (264). Structural

comparisons, along with shared genomic associations, were used to further refine the detailed

interrelationships within and between the groups of the superfamily. Families with broader phyletic

patterns and conservation across multiple major lineages ("phyla") were assumed to be more ancient in

their emergence, and those with a more limited distribution were assumed to be later derivations

(Figures 4,S1).

Contextual Network construction

Each domain architecture and conserved gene neighborhood was decomposed into their constituent

domains. These domains were then labeled for their biochemical function and stored as a YAML file

(Supplementary Data). The contextual connections were then rendered as a graph with the domains as

its nodes and the adjacency relationships as its edges. Cliques containing a given Lipocone domain were

detected in this graph and merged to constitute their respective dense subgraphs. These subgraphs

were then examined for the statistically significant prevalence of particular labeled functions using the

Fisher exact test. Network analysis was performed using the functions of the R igraph or Python

networkX libraries.

Structure analysis

1051   Protein structures were modeled using Alphafold3 (265), with visualization and manipulation

1052   performed using either MOL* (266) or PyMOL software. Structural similarity searches were conducted

1053   using the DALIlite (267) and FOLDSEEK (268) programs. DALIlite was also used to generate structural

1054   alignments.

1055

1056   **Hydrophobicity analysis**

1057   To create the membrane propensity plots, for each protein P_i in a given family, we compute the average

1058   TM-propensity of its amino acids using the TM tendency scale (38). This score H_i for P_i is calculated as:

1059

1060
$$H_i = \frac{1}{n}\sum_{j=1}^{n} h_j$$

1061

1062   where H_j is the TM tendency of the j-th amino acid in the protein P_i, and n is its total length in amino

1063   acids.

1064

1065

FIGURE LEGENDS
Figure 1. (A) The four individual helices forming the core of the Lipocone superfamily are colored
consistently across the structures from representative families. The inter-helix linkers are colored gray,
and lineage-specific synapomorphic insertions and extensions are colored light brown. Active site and
other residues of interest are rendered as ball-and-stick. Protein Data Bank (PDB) IDs or Genbank
accessions used to generate AF3 models are provided. (B) Relationship network of the Lipocone
families. The thickness of the edges are scaled by the negative-log HHalign p-values. Families are
colored according to the community identified by the Leiden algorithm (37) (see Methods). (C) Box
plots displaying core helix transmembrane propensity scores of individual sequences within different
Lipocone families. The horizontal divider represents the boundary between typical TM and soluble
sequences.

Figure 2. Sequence logo conserved core elements of different Lipocone families, generally
corresponding to the core H2, H3, and H4 helices. The three conserved core active site residue positions
are boxed in dotted lines with the inferred ancestral residue indicated at the top of the alignment.
Families are grouped and labeled on the left in their higher-order clades.

Figure 3. Known and predicted Lipocone reaction mechanisms. Experimentally supported reactions are
boxed in blue (A-B), while a predicted reaction based on genome displacement by a Lipocone domain of
an experimentally characterized enzyme is boxed in orange (C). The remaining reactions (D-G) are
suggested based on the contextual inferences in this work.

Figure 4. Reconstructed evolutionary scenario for the diversification of the Lipocone superfamily. The
relative temporal epochs are demarcated by vertical lines and labeled at the bottom. The clades are
represented by colored lines indicating the maximum depth to which the families listed to the right can
be traced. Colors track the superkingdom level phyletic distribution of the family. Broken lines indicate
uncertainty in the starting point of a lineage(s). Inferred or experimentally characterized functions for
families are depicted by colored boxes to the left of family names.

Figure 5. Representative contexts for Lipocones, grouped by shared functional themes. Genes are
depicted by box arrows, with the arrowhead indicating the 3' end of genes. Genes encoding proteins
with multiple domains are broken into labeled sections corresponding to them. Domain architectures
are depicted by the individual domains represented by geometric shapes. TMs, lipoboxes (LPs), and SPs
are depicted as unlabeled, narrow yellow, blue, and red rectangles, respectively. All Lipocone domains
are consistently colored in orange. Genes marked with asterisks are labeled by the Genbank accession
number below each context. Colored labels above genes denote well-known gene names or gene cluster
modules. Abbreviations: PTase, peptidase; TFase, transferase; GlycosylTFase, Glycosyltransferase;
MPTase, metallopeptidase; TGase, transglycosylase; SLP, serine-containing lipobox; cNMPBD, cNMP-
binding domain; NCPBM, novel putative carbohydrate binding module; (w)HTH, (winged) helix-turn-
helix; ZnR, zinc ribbon; PPTs, pentapeptide repeats; Imm, immunity protein; βPs, β-propeller repeats;

Cystatin-FD, Cystatin fold domain; MTase, methylase; PGBD, peptidoglycan-binding domain; MβL, metallo-β-lactamase; L12-ClpS, ClpS-ribosomal L7/L12 domain; TA, teichoic acid.

Figure 6. Lipocone contextual network. The network represents the conserved contextual associations of Lipocone domains (represented as hexagonal nodes). Nodes and edges are colored based on the known or inferred functional categories of the domains, with groups labeled in matching colors. The nodes are scaled by degree. Gray coloring indicates domains without specific functional assignments. Examples of conserved gene neighborhoods and domain architectures supplementing those in Figure 5 illustrate contexts that bridge functional themes. Here, individual domains are colored to match network coloring. Additional abbreviations to those in Figure 5: APH-Pkinase, aminoglycoside phosphotransferase-like protein kinase; HUP, HIGH-signature, UspA and PP-ATPase superfamily-like domain; Alk-phosphatase, Alkaline phosphatase; dehyd, dehydrogenase; TPRs, tetratricopeptide repeats; PMM/PGM, phosphomannomutase/phosphoglucomutase; ZnF, zinc finger; APC-transporter, amino acid-polyamine-organocation transporter; LPS, lipopolysaccharide.

ACKNOWLEDGEMENTS

This research was supported by the Division of Intramural Research at the National Library of Medicine (NLM), National Institutes of Health (NIH). This research was supported in part by an appointment to the NLM Research Participation Program administered by the Oak Ridge Institute for Science and Education (ORISE) through an interagency agreement between the U.S. Department of Energy (DOE) and the NLM.

CONFLICT OF INTEREST STATEMENT

The authors declare that they have no conflict of interest.

REFERENCES

1.   Richards, G.S. and Degnan, B.M. (2009) The dawn of developmental signaling in the metazoa. *Cold Spring Harb Symp Quant Biol*, **74**, 81-90.

2.   Jessen, S., Gu, B. and Dai, X. (2008) Pygopus and the Wnt signaling pathway: a diverse set of connections. *Bioessays*, **30**, 448-456.

3.   Morata, G. and Lawrence, P.A. (1977) The development of wingless, a homeotic mutation of Drosophila. *Dev Biol*, **56**, 227-240.

4.   Rijsewijk, F., Schuermann, M., Wagenaar, E., Parren, P., Weigel, D. and Nusse, R. (1987) The Drosophila homolog of the mouse mammary oncogene int-1 is identical to the segment polarity gene wingless. *Cell*, **50**, 649-657.

5.   Sharma, R.P. (1973) Wingless a new mutant in Drosophila melanogaster. *Drosophila information service*, 134.

6.   Nusse, R. and Varmus, H.E. (1982) Many tumors induced by the mouse mammary tumor virus contain a provirus integrated in the same region of the host genome. *Cell*, **31**, 99-109.

7.   Nusse, R. and Varmus, H.E. (1992) Wnt genes. *Cell*, **69**, 1073-1087.

8.   Komiya, Y. and Habas, R. (2008) Wnt signal transduction pathways. *Organogenesis*, **4**, 68-75.

9.   Segalen, M. and Bellaiche, Y. (2009) Cell division orientation and planar cell polarity
pathways. *Semin Cell Dev Biol*, **20**, 972-977.

10.   Slusarski, D.C. and Pelegri, F. (2007) Calcium signaling in vertebrate embryonic patterning and
morphogenesis. *Dev Biol*, **307**, 1-13.

11.   Logan, C.Y. and Nusse, R. (2004) The Wnt signaling pathway in development and disease. *Annu*
Rev Cell Dev Biol, **20**, 781-810.

12.   Welters, H.J. and Kulkarni, R.N. (2008) Wnt signaling: relevance to beta-cell biology and
diabetes. *Trends Endocrinol Metab*, **19**, 349-355.

13.   Christian, J.L. (2000) BMP, Wnt and Hedgehog signals: how far can they go? *Curr Opin Cell Biol*,
12, 244-249.

14.   Schulte, G. and Bryja, V. (2007) The Frizzled family of unconventional G-protein-coupled
receptors. *Trends Pharmacol Sci*, **28**, 518-525.

15.   Bhanot, P., Brink, M., Samos, C.H., Hsieh, J.C., Wang, Y., Macke, J.P., Andrew, D., Nathans, J. and
Nusse, R. (1996) A new member of the frizzled family from Drosophila functions as a Wingless
receptor. *Nature*, **382**, 225-230.

16.   Takada, R., Satomi, Y., Kurata, T., Ueno, N., Norioka, S., Kondoh, H., Takao, T. and Takada, S.
(2006) Monounsaturated fatty acid modification of Wnt protein: its role in Wnt secretion. *Dev*
Cell, **11**, 791-801.

17.   Kurayoshi, M., Yamamoto, H., Izumi, S. and Kikuchi, A. (2007) Post-translational
palmitoylation and glycosylation of Wnt-5a are necessary for its signalling. *Biochem J*, **402**,
515-523.

18.   Gao, C. and Chen, Y.G. (2010) Dishevelled: The hub of Wnt signaling. *Cell Signal*, **22**, 717-727.

19.   Klingensmith, J., Nusse, R. and Perrimon, N. (1994) The Drosophila segment polarity gene
dishevelled encodes a novel protein required for response to the wingless signal. *Genes Dev*, **8**,
118-130.

20.   Kramps, T., Peter, O., Brunner, E., Nellen, D., Froesch, B., Chatterjee, S., Murone, M., Zullig, S.
and Basler, K. (2002) Wnt/wingless signaling requires BCL9/legless-mediated recruitment of
pygopus to the nuclear beta-catenin-TCF complex. *Cell*, **109**, 47-60.

21.   van Tienen, L.M., Mieszczanek, J., Fiedler, M., Rutherford, T.J. and Bienz, M. (2017)
Constitutive scaffolding of multiple Wnt enhanceosome components by Legless/BCL9. *Elife*, **6**.

22.   Archbold, H.C., Yang, Y.X., Chen, L. and Cadigan, K.M. (2012) How do they do Wnt they do?:
regulation of transcription by the Wnt/beta-catenin pathway. *Acta Physiol (Oxf)*, **204**, 74-109.

23.   Holstein, T.W. (2012) The evolution of the Wnt pathway. *Cold Spring Harb Perspect Biol*, **4**,
a007922.

24.   Burroughs, A.M. and Aravind, L. (2020) Identification of Uncharacterized Components of
Prokaryotic Immune Systems and Their Diverse Eukaryotic Reformulations. *J Bacteriol*, **202**.

25.   Aravind, L., Iyer, L.M. and Burroughs, A.M. (2022) Discovering Biological Conflict Systems
1185   Through Genome Analysis: Evolutionary Principles and Biochemical Novelty. *Annu Rev Biomed*
1186   *Data Sci*, **5**, 367-391.
1187   26.   Kuge, O., Saito, K. and Nishijima, M. (1997) Cloning of a Chinese hamster ovary (CHO) cDNA
1188   encoding phosphatidylserine synthase (PSS) II, overexpression of which suppresses the
1189   phosphatidylserine biosynthetic defect of a PSS I-lacking mutant of CHO-K1 cells. *J Biol Chem*,
1190   **272**, 19133-19139.
1191   27.   Kuge, O., Nishijima, M. and Akamatsu, Y. (1991) A Chinese hamster cDNA encoding a protein
1192   essential for phosphatidylserine synthase I activity. *J Biol Chem*, **266**, 24184-24189.
1193   28.   Stone, S.J. and Vance, J.E. (1999) Cloning and expression of murine liver phosphatidylserine
1194   synthase (PSS)-2: differential regulation of phospholipid metabolism by PSS1 and PSS2.
1195   *Biochem J*, **342 (Pt 1)**, 57-64.
1196   29.   Whitney, J.C., Peterson, S.B., Kim, J., Pazos, M., Verster, A.J., Radey, M.C., Kulasekara, H.D.,
1197   Ching, M.Q., Bullen, N.P., Bryant, D., Goo, Y.A., Surette, M.G., Borenstein, E., Vollmer, W. and
1198   Mougous, J.D. (2017) A broadly distributed toxin family mediates contact-dependent
1199   antagonism between gram-positive bacteria. *Elife*, **6**.
1200   30.   Sack, G.H., Jr. (2018) Serum amyloid A - a review. *Mol Med*, **24**, 46.
1201   31.   Arthur, M., Depardieu, F., Molinas, C., Reynolds, P. and Courvalin, P. (1995) The vanZ gene of
1202   Tn1546 from Enterococcus faecium BM4147 confers resistance to teicoplanin. *Gene*, **154**, 87-
1203   92.
1204   32.   Arthur, M., Depardieu, F., Reynolds, P. and Courvalin, P. (1999) Moderate-level resistance to
1205   glycopeptide LY333328 mediated by genes of the vanA and vanB clusters in enterococci.
1206   *Antimicrob Agents Chemother*, **43**, 1875-1880.
1207   33.   Bazan, J.F., Janda, C.Y. and Garcia, K.C. (2012) Structural architecture and functional evolution
1208   of Wnts. *Dev Cell*, **23**, 227-232.
1209   34.   Janda, C.Y., Waghray, D., Levin, A.M., Thomas, C. and Garcia, K.C. (2012) Structural basis of
1210   Wnt recognition by Frizzled. *Science*, **337**, 59-64.
1211   35.   Chu, M.L., Ahn, V.E., Choi, H.J., Daniels, D.L., Nusse, R. and Weis, W.I. (2013) structural Studies
1212   of Wnts and identification of an LRP6 binding site. *Structure*, **21**, 1235-1242.
1213   36.   Zhong, Q., Zhao, Y., Ye, F., Xiao, Z., Huang, G., Xu, M., Zhang, Y., Zhan, X., Sun, K., Wang, Z.,
1214   Cheng, S., Feng, S., Zhao, X., Zhang, J., Lu, P., Xu, W., Zhou, Q. and Ma, D. (2021) Cryo-EM
1215   structure of human Wntless in complex with Wnt3a. *Nat Commun*, **12**, 4541.
1216   37.   Traag, V.A., Waltman, L. and van Eck, N.J. (2019) From Louvain to Leiden: guaranteeing well-
1217   connected communities. *Sci Rep*, **9**, 5233.
1218   38.   Zhao, G. and London, E. (2006) An amino acid "transmembrane tendency" scale that
1219   approaches the theoretical limit to accuracy for prediction of transmembrane helices:
1220   relationship to biological hydrophobicity. *Protein Sci*, **15**, 1987-2001.
1221   39.   Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K.,
1222   Bates, R., Zidek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S.A.A., Ballard, A.J., Cowie, A.,
1223   Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T., Petersen, S., Reiman, D., Clancy, E.,

Zielinski, M., Steinegger, M., Pacholska, M., Berghammer, T., Bodenstein, S., Silver, D., Vinyals,
O., Senior, A.W., Kavukcuoglu, K., Kohli, P. and Hassabis, D. (2021) Highly accurate protein
structure prediction with AlphaFold. *Nature*, **596**, 583-589.

40.   Bastard, K., Smith, A.A., Vergne-Vaxelaire, C., Perret, A., Zaparucha, A., De Melo-Minardi, R.,
Mariage, A., Boutard, M., Debard, A., Lechaplais, C., Pelle, C., Pellouin, V., Perchat, N., Petit,
J.L., Kreimeyer, A., Medigue, C., Weissenbach, J., Artiguenave, F., De Berardinis, V., Vallenet, D.
and Salanoubat, M. (2014) Revealing the hidden functional diversity of an enzyme family. *Nat*
Chem Biol, **10**, 42-49.

41.   Glasner, M.E., Gerlt, J.A. and Babbitt, P.C. (2006) Evolution of enzyme superfamilies. *Curr Opin*
Chem Biol, **10**, 492-497.

42.   Zhang, D., Iyer, L.M., Burroughs, A.M. and Aravind, L. (2014) Resilience of biochemical activity
in protein domains in the face of structural divergence. *Curr Opin Struct Biol*, **26**, 92-103.

43.   Tomohiro, S., Kawaguti, A., Kawabe, Y., Kitada, S. and Kuge, O. (2009) Purification and
characterization of human phosphatidylserine synthases 1 and 2. *Biochem J*, **418**, 421-429.

44.   Saito, K., Kuge, O., Akamatsu, Y. and Nishijima, M. (1996) Immunochemical identification of
the pssA gene product as phosphatidylserine synthase I of Chinese hamster ovary cells. *FEBS*
Lett, **395**, 262-266.

45.   Stone, S.J. and Vance, J.E. (2000) Phosphatidylserine synthase-1 and -2 are localized to
mitochondria-associated membranes. *J Biol Chem*, **275**, 34534-34540.

46.   Miyata, N. and Kuge, O. (2021) Topology of phosphatidylserine synthase 1 in the endoplasmic
reticulum membrane. *Protein Sci*, **30**, 2346-2353.

47.   Anderson, J.S., Matsuhashi, M., Haskin, M.A. and Strominger, J.L. (1967) Biosythesis of the
peptidoglycan of bacterial cell walls. II. Phospholipid carriers in the reaction sequence. *J Biol*
Chem, **242**, 3180-3190.

48.   Higashi, Y., Strominger, J.L. and Sweeley, C.C. (1967) Structure of a lipid intermediate in cell
wall peptidoglycan synthesis: a derivative of a C55 isoprenoid alcohol. *Proc Natl Acad Sci U S A*,
57, 1878-1884.

49.   English, D. (1996) Phosphatidic acid: a lipid messenger involved in intracellular and
extracellular signalling. *Cell Signal*, **8**, 341-347.

50.   Zamocky, M. and Ferianc, P. (2023) Discovering the deep evolutionary roots of serum amyloid
A protein family. *Int J Biol Macromol*, **252**, 126537.

51.   Woods, E.C., Wetzel, D., Mukerjee, M. and McBride, S.M. (2018) Examination of the
Clostridioides (Clostridium) difficile VanZ ortholog, CD1240. *Anaerobe*, **53**, 108-115.

52.   Sur, V.P., Mazumdar, A., Vimberg, V., Stefani, T., Androvic, L., Kracikova, L., Laga, R., Kamenik,
Z. and Komrskova, K. (2021) Specific Inhibition of VanZ-Mediated Resistance to
Lipoglycopeptide Antibiotics. *Int J Mol Sci*, **23**.

53.   Stogios, P.J. and Savchenko, A. (2020) Molecular mechanisms of vancomycin resistance. *Protein*
Sci, **29**, 654-669.

54.   Dziarski, R. and Gupta, D. (2006) The peptidoglycan recognition proteins (PGRPs). *Genome Biol*,
7, 232.

55.   Bliven, S. and Prlic, A. (2012) Circular permutation in proteins. *PLoS Comput Biol*, **8**, e1002445.

56.   Aravind, L. (2000) Guilt by association: contextual information in genome analysis. *Genome*
1266   *Res*, **10**, 1074-1077.

1267   57.   Vance, J.E. (2018) Historical perspective: phosphatidylserine and phosphatidylethanolamine
1268   from the 1800s to the present. *J Lipid Res*, **59**, 923-944.

1269   58.   Koga, Y. and Morii, H. (2005) Recent advances in structural research on ether lipids from
1270   archaea including comparative and physiological aspects. *Biosci Biotechnol Biochem*, **69**, 2019-
1271   2034.

1272   59.   Caforio, A. and Driessen, A.J.M. (2017) Archaeal phospholipids: Structural properties and
1273   biosynthesis. *Biochim Biophys Acta Mol Cell Biol Lipids*, **1862**, 1325-1339.

1274   60.   Rezanka, T., Kyselova, L. and Murphy, D.J. (2023) Archaeal lipids. *Prog Lipid Res*, **91**, 101237.

1275   61.   Narita, S. and Tokuda, H. (2010) Sorting of bacterial lipoproteins to the outer membrane by
1276   the Lol system. *Methods Mol Biol*, **619**, 117-129.

1277   62.   Kamada, K., Miyata, M. and Hirano, T. (2013) Molecular basis of SMC ATPase activation: role of
1278   internal structural changes of the regulatory subcomplex ScpAB. *Structure*, **21**, 581-594.

1279   63.   Schleiffer, A., Kaitna, S., Maurer-Stroh, S., Glotzer, M., Nasmyth, K. and Eisenhaber, F. (2003)
1280   Kleisins: a superfamily of bacterial and eukaryotic SMC protein partners. *Mol Cell*, **11**, 571-575.

1281   64.   Soppa, J., Kobayashi, K., Noirot-Gros, M.F., Oesterhelt, D., Ehrlich, S.D., Dervyn, E., Ogasawara,
1282   N. and Moriya, S. (2002) Discovery of two novel families of proteins that are proposed to
1283   interact with prokaryotic SMC proteins, and characterization of the Bacillus subtilis family
1284   members ScpA and ScpB. *Mol Microbiol*, **45**, 59-71.

1285   65.   Aravind, L., Anantharaman, V., Balaji, S., Babu, M.M. and Iyer, L.M. (2005) The many faces of
1286   the helix-turn-helix domain: transcription regulation and beyond. *FEMS Microbiol Rev*, **29**, 231-
1287   262.

1288   66.   Pellicer, M.T., Badia, J., Aguilar, J. and Baldoma, L. (1996) glc locus of Escherichia coli:
1289   characterization of genes encoding the subunits of glycolate oxidase and the glc regulator
1290   protein. *J Bacteriol*, **178**, 2051-2059.

1291   67.   Ramachandran, R., Hartmann, C., Song, H.K., Huber, R. and Bochtler, M. (2002) Functional
1292   interactions of HslV (ClpQ) with the ATPase HslU (ClpY). *Proc Natl Acad Sci U S A*, **99**, 7396-7401.

1293   68.   Storck, E.M., Ozbalci, C. and Eggert, U.S. (2018) Lipid Cell Biology: A Focus on Lipids in Cell
1294   Division. *Annu Rev Biochem*, **87**, 839-869.

1295   69.   Barak, I. and Muchova, K. (2013) The role of lipid domains in bacterial cell processes. *Int J Mol*
1296   *Sci*, **14**, 4050-4065.

1297   70.   Hauck, A.K. and Bernlohr, D.A. (2016) Oxidative stress and lipotoxicity. *J Lipid Res*, **57**, 1976-
1298   1986.

1299   71.   Guo, D. and Tropp, B.E. (2000) A second Escherichia coli protein with CL synthase activity.
1300   *Biochim Biophys Acta*, **1483**, 263-274.

1301   72.   Tan, B.K., Bogdanov, M., Zhao, J., Dowhan, W., Raetz, C.R. and Guan, Z. (2012) Discovery of a
1302   cardiolipin synthase utilizing phosphatidylethanolamine and phosphatidylglycerol as
1303   substrates. *Proc Natl Acad Sci U S A*, **109**, 16504-16509.

73. Aravind, L. and Koonin, E.V. (1998) Phosphoesterase domains associated with DNA
polymerases of diverse origins. *Nucleic Acids Res*, **26**, 3746-3752.
74. Burroughs, A.M., Allen, K.N., Dunaway-Mariano, D. and Aravind, L. (2006) Evolutionary
genomics of the HAD superfamily: understanding the structural adaptations and catalytic
diversity in a superfamily of phosphoesterases and allied enzymes. *J Mol Biol*, **361**, 1003-1034.
75. Vershinin, A. (1999) Biological functions of carotenoids--diversity and evolution. *Biofactors*,
10, 99-104.
76. Šesták, Z., Britton, G., Liaaen-Jensen, S. and Pfander, H. (2004) Carotenoids. Handbook.
Photosynthetica, **42**, 186.
77. Sandmann, G. and Misawa, N. (1992) New functional assignment of the carotenogenic genes
crtB and crtE with constructs of these genes from Erwinia species. *FEMS Microbiol Lett*, **69**, 253-
257.
78. Chamovitz, D., Misawa, N., Sandmann, G. and Hirschberg, J. (1992) Molecular cloning and
expression in Escherichia coli of a cyanobacterial gene coding for phytoene synthase, a
carotenoid biosynthesis enzyme. *FEBS Lett*, **296**, 305-310.
79. Liu, J. and Mushegian, A. (2003) Three monophyletic superfamilies account for the majority of
the known glycosyltransferases. *Protein Sci*, **12**, 1418-1431.
80. Tran, U.C. and Clarke, C.F. (2007) Endogenous synthesis of coenzyme Q in eukaryotes.
Mitochondrion, **7 Suppl**, S62-71.
81. Massiere, F. and Badet-Denisot, M.A. (1998) The mechanism of glutamine-dependent
amidotransferases. *Cell Mol Life Sci*, **54**, 205-222.
82. Liu, X., Zeng, J., Chen, X. and Xie, W. (2016) Crystal structures of RidA, an important enzyme
for the prevention of toxic side products. *Sci Rep*, **6**, 30494.
83. Aravind, L. and Koonin, E.V. (1998) The HD domain defines a new superfamily of metal-
dependent phosphohydrolases. *Trends Biochem Sci*, **23**, 469-472.
84. Becker, A., Kleickmann, A., Kuster, H., Keller, M., Arnold, W. and Puhler, A. (1993) Analysis of
the Rhizobium meliloti genes exoU, exoV, exoW, exoT, and exoI involved in exopolysaccharide
biosynthesis and nodule invasion: exoU and exoW probably encode glucosyltransferases. *Mol*
Plant Microbe Interact, **6**, 735-744.
85. Ruiz, N. (2015) Lipid Flippases for Bacterial Peptidoglycan Biosynthesis. *Lipid Insights*, **8**, 21-31.
86. Ruiz, N. (2008) Bioinformatics identification of MurJ (MviN) as the peptidoglycan lipid II
flippase in Escherichia coli. *Proc Natl Acad Sci U S A*, **105**, 15553-15557.
87. Heydanek, M.G., Jr. and Neuhaus, F.C. (1969) The initial stage in peptidoglycan synthesis. IV.
Solubilization of phospho-N-acetylmuramyl-pentapeptide translocase. *Biochemistry*, **8**, 1474-
1481.
88. Mengin-Lecreulx, D. and van Heijenoort, J. (1993) Identification of the glmU gene encoding N-
acetylglucosamine-1-phosphate uridyltransferase in Escherichia coli. *J Bacteriol*, **175**, 6150-
6157.
89. Mengin-Lecreulx, D. and van Heijenoort, J. (1996) Characterization of the essential gene glmM
encoding phosphoglucosamine mutase in Escherichia coli. *J Biol Chem*, **271**, 32-39.

90. van Heijenoort, J. (2007) Lipid intermediates in the biosynthesis of bacterial peptidoglycan.
Microbiol Mol Biol Rev, **71**, 620-635.

91. Ashraf, K.U., Nygaard, R., Vickery, O.N., Erramilli, S.K., Herrera, C.M., McConville, T.H., Petrou,
V.I., Giacometti, S.I., Dufrisne, M.B., Nosol, K., Zinkle, A.P., Graham, C.L.B., Loukeris, M., Kloss,
B., Skorupinska-Tudek, K., Swiezewska, E., Roper, D.I., Clarke, O.B., Uhlemann, A.C.,
Kossiakoff, A.A., Trent, M.S., Stansfeld, P.J. and Mancia, F. (2022) Structural basis of
lipopolysaccharide maturation by the O-antigen ligase. *Nature*, **604**, 371-376.

92. Franco, A.V., Liu, D. and Reeves, P.R. (1998) The wzz (cld) protein in Escherichia coli: amino
acid sequence variation determines O-antigen chain length specificity. *J Bacteriol*, **180**, 2670-
2675.

93. Rai, A.K., Carr, J.F., Bautista, D.E., Wang, W. and Mitchell, A.M. (2021) ElyC and Cyclic
Enterobacterial Common Antigen Regulate Synthesis of Phosphoglyceride-Linked
Enterobacterial Common Antigen. *mBio*, **12**, e0284621.

94. Kumar, A.S., Mody, K. and Jha, B. (2007) Bacterial exopolysaccharides--a perception. *J Basic
Microbiol*, **47**, 103-117.

95. van Heijenoort, J. (2001) Formation of the glycan chains in the synthesis of bacterial
peptidoglycan. *Glycobiology*, **11**, 25R-36R.

96. Hong, Y., Hu, D., Verderosa, A.D., Qin, J., Totsika, M. and Reeves, P.R. (2023) Repeat-Unit
Elongations To Produce Bacterial Complex Long Polysaccharide Chains, an O-Antigen
Perspective. *EcoSal Plus*, **11**, eesp00202022.

97. Rai, A.K. and Mitchell, A.M. (2020) Enterobacterial Common Antigen: Synthesis and Function
of an Enigmatic Molecule. *mBio*, **11**.

98. Sham, L.T., Butler, E.K., Lebar, M.D., Kahne, D., Bernhardt, T.G. and Ruiz, N. (2014) Bacterial
cell wall. MurJ is the flippase of lipid-linked precursors for peptidoglycan biogenesis. *Science*,
345, 220-222.

99. Kim, S., Pires, M.M. and Im, W. (2018) Insight into Elongation Stages of Peptidoglycan
Processing in Bacterial Cytoplasmic Membranes. *Sci Rep*, **8**, 17704.

100. Di Guilmi, A.M., Dessen, A., Dideberg, O. and Vernet, T. (2003) The glycosyltransferase domain
of penicillin-binding protein 2a from Streptococcus pneumoniae catalyzes the polymerization
of murein glycan chains. *J Bacteriol*, **185**, 4418-4423.

101. Weckener, M., Woodward, L.S., Clarke, B.R., Liu, H., Ward, P.N., Le Bas, A., Bhella, D., Whitfield,
C. and Naismith, J.H. (2023) The lipid linked oligosaccharide polymerase Wzy and its
regulating co-polymerase, Wzz, from enterobacterial common antigen biosynthesis form a
complex. *Open Biol*, **13**, 220373.

102. Han, W., Wu, B., Li, L., Zhao, G., Woodward, R., Pettit, N., Cai, L., Thon, V. and Wang, P.G. (2012)
Defining function of lipopolysaccharide O-antigen ligase WaaL using chemoenzymatically
synthesized substrates. *J Biol Chem*, **287**, 5357-5365.

103. Imperiali, B. (2019) Bacterial carbohydrate diversity - a Brave New World. *Curr Opin Chem Biol*,
53, 1-8.

104. Mostowy, R.J. and Holt, K.E. (2018) Diversity-Generating Machines: Genetics of Bacterial
Sugar-Coating. *Trends Microbiol*, **26**, 1008-1021.
105. Saiki, K. and Konishi, K. (2007) Identification of a Porphyromonas gingivalis novel protein sov
required for the secretion of gingipains. *Microbiol Immunol*, **51**, 483-491.
106. Pares, S., Cohen-Addad, C., Sieker, L., Neuburger, M. and Douce, R. (1994) X-ray structure
determination at 2.6-A resolution of a lipoate-containing protein: the H-protein of the glycine
decarboxylase complex from pea leaves. *Proc Natl Acad Sci U S A*, **91**, 4850-4853.
107. Shultis, D.D., Purdy, M.D., Banchs, C.N. and Wiener, M.C. (2006) Outer membrane active
transport: structure of the BtuB:TonB complex. *Science*, **312**, 1396-1399.
108. Rahn, A., Beis, K., Naismith, J.H. and Whitfield, C. (2003) A novel outer membrane protein,
Wzi, is involved in surface assembly of the Escherichia coli K30 group 1 capsule. *J Bacteriol*,
185, 5882-5890.
109. Qi, Y. and Grishin, N.V. (2005) Structural classification of thioredoxin-like fold proteins.
Proteins, **58**, 376-388.
110. Suplatov, D.A., Besenmatter, W., Svedas, V.K. and Svendsen, A. (2012) Bioinformatic analysis of
alpha/beta-hydrolase fold enzymes reveals subfamily-specific positions responsible for
discrimination of amidase and lipase activities. *Protein Eng Des Sel*, **25**, 689-697.
111. Palomeque-Messia, P., Englebert, S., Leyh-Bouille, M., Nguyen-Disteche, M., Duez, C., Houba,
S., Dideberg, O., Van Beeumen, J. and Ghuysen, J.M. (1991) Amino acid sequence of the
penicillin-binding protein/DD-peptidase of Streptomyces K15. Predicted secondary
structures of the low Mr penicillin-binding proteins of class A. *Biochem J*, **279 (Pt 1)**, 223-230.
112. Aravind, L. (1999) An evolutionary classification of the metallo-beta-lactamase fold proteins.
In Silico Biol, **1**, 69-91.
113. Novinec, M. and Lenarcic, B. (2013) Papain-like peptidases: structure, function, and evolution.
Biomol Concepts, **4**, 287-308.
114. Razew, A., Schwarz, J.N., Mitkowski, P., Sabala, I. and Kaus-Drobek, M. (2022) One fold, many
functions-M23 family of peptidoglycan hydrolases. *Front Microbiol*, **13**, 1036964.
115. Mans, B.J., Anantharaman, V., Aravind, L. and Koonin, E.V. (2004) Comparative genomics,
evolution and origins of the nuclear envelope and nuclear pore complex. *Cell Cycle*, **3**, 1612-
1637.
116. Wimley, W.C. (2003) The versatile beta-barrel membrane protein. *Curr Opin Struct Biol*, **13**, 404-
411.
117. Fairman, J.W., Noinaj, N. and Buchanan, S.K. (2011) The structural biology of beta-barrel
membrane proteins: a summary of recent reports. *Curr Opin Struct Biol*, **21**, 523-531.
118. Tokuda, H. and Matsuyama, S. (2004) Sorting of lipoproteins to the outer membrane in E. coli.
Biochim Biophys Acta, **1693**, 5-13.
119. Sanchez-Pulido, L., Devos, D., Genevrois, S., Vicente, M. and Valencia, A. (2003) POTRA: a
conserved domain in the FtsQ family and a class of beta-barrel outer membrane proteins.
Trends Biochem Sci, **28**, 523-526.

120. Kim, S., Malinverni, J.C., Sliz, P., Silhavy, T.J., Harrison, S.C. and Kahne, D. (2007) Structure and
function of an essential component of the outer membrane protein assembly machine.
Science, **317**, 961-964.
121. Oke, M., Sarra, R., Ghirlando, R., Farnaud, S., Gorringe, A.R., Evans, R.W. and Buchanan, S.K.
(2004) The plug domain of a neisserial TonB-dependent transporter retains structural
integrity in the absence of its transmembrane beta-barrel. *FEBS Lett*, **564**, 294-300.
122. Carr, S., Penfold, C.N., Bamford, V., James, R. and Hemmings, A.M. (2000) The structure of
TolB, an essential component of the tol-dependent translocation system, and its protein-
protein interaction with the translocation domain of colicin E9. *Structure*, **8**, 57-66.
123. Ghosh, M., Tucker, D.E., Burchett, S.A. and Leslie, C.C. (2006) Properties of the Group IV
phospholipase A2 family. *Prog Lipid Res*, **45**, 487-510.
124. Stukey, J. and Carman, G.M. (1997) Identification of a novel phosphatase sequence motif.
Protein Sci, **6**, 469-472.
125. Whisstock, J.C., Romero, S., Gurung, R., Nandurkar, H., Ooms, L.M., Bottomley, S.P. and
Mitchell, C.A. (2000) The inositol polyphosphate 5-phosphatases and the
apurinic/apyrimidinic base excision repair endonucleases share a common mechanism for
catalysis. *J Biol Chem*, **275**, 37055-37061.
126. Burroughs, A.M. and Aravind, L. (2023) New biochemistry in the Rhodanese-phosphatase
superfamily: emerging roles in diverse metabolic processes, nucleic acid modifications, and
biological conflicts. *NAR Genom Bioinform*, **5**, lqad029.
127. Holtje, J.V., Mirelman, D., Sharon, N. and Schwarz, U. (1975) Novel type of murein
transglycosylase in Escherichia coli. *J Bacteriol*, **124**, 1067-1076.
128. Yunck, R., Cho, H. and Bernhardt, T.G. (2016) Identification of MltG as a potential terminase
for peptidoglycan polymerization in bacteria. *Mol Microbiol*, **99**, 700-718.
129. Einsle, O., Messerschmidt, A., Stach, P., Bourenkov, G.P., Bartunik, H.D., Huber, R. and Kroneck,
P.M. (1999) Structure of cytochrome c nitrite reductase. *Nature*, **400**, 476-480.
130. Levin, S., Almo, S.C. and Satir, B.H. (1999) Functional diversity of the phosphoglucomutase
superfamily: structural implications. *Protein Eng*, **12**, 737-746.
131. Dong, X., Kato-Murayama, M., Muramatsu, T., Mori, H., Shirouzu, M., Bessho, Y. and
Yokoyama, S. (2007) The crystal structure of leucyl/phenylalanyl-tRNA-protein transferase
from Escherichia coli. *Protein Sci*, **16**, 528-534.
132. Antia, M., Hoare, D.S. and Work, E. (1957) The stereoisomers of alpha epsilon-diaminopimelic
acid. III. Properties and distribution of diaminopimelic acid racemase, an enzyme causing
interconversion of the LL and meso isomers. *Biochem J*, **65**, 448-459.
133. Koraimann, G. (2003) Lytic transglycosylases in macromolecular transport systems of Gram-
negative bacteria. *Cell Mol Life Sci*, **60**, 2371-2388.
134. Muley, V.Y., Akhter, Y. and Galande, S. (2019) PDZ Domains Across the Microbial World:
Molecular Link to the Proteases, Stress Response, and Protein Synthesis. *Genome Biol Evol*, **11**,
644-659.

135. Hara, H., Yamamoto, Y., Higashitani, A., Suzuki, H. and Nishimura, Y. (1991) Cloning, mapping, and characterization of the Escherichia coli prc gene, which is involved in C-terminal processing of penicillin-binding protein 3. *J Bacteriol*, **173**, 4799-4813.

136. Wong, L.H. and Levine, T.P. (2017) Tubular lipid binding proteins (TULIPs) growing everywhere. *Biochim Biophys Acta Mol Cell Res*, **1864**, 1439-1449.

137. Levine, T.P. (2019) Remote homology searches identify bacterial homologues of eukaryotic lipid transfer proteins, including Chorein-N domains in TamB and AsmA and Mdm31p. *BMC Mol Cell Biol*, **20**, 43.

138. Dhanaraj, V., Ye, Q.Z., Johnson, L.L., Hupe, D.J., Ortwine, D.F., Dunbar, J.B., Jr., Rubin, J.R., Pavlovsky, A., Humblet, C. and Blundell, T.L. (1996) X-ray structure of a hydroxamate inhibitor complex of stromelysin catalytic domain and its comparison with members of the zinc metalloproteinase superfamily. *Structure*, **4**, 375-386.

139. Lo, Y.C., Lin, S.C., Shaw, J.F. and Liaw, Y.C. (2003) Crystal structure of Escherichia coli thioesterase I/protease I/lysophospholipase L1: consensus sequence blocks constitute the catalytic center of SGNH-hydrolases through a conserved hydrogen bond network. *J Mol Biol*, **330**, 539-551.

140. Hofmann, K. (2000) A superfamily of membrane-bound O-acyltransferases with implications for wnt signaling. *Trends Biochem Sci*, **25**, 111-112.

141. Rahlwes, K.C., Ha, S.A., Motooka, D., Mayfield, J.A., Baumoel, L.R., Strickland, J.N., Torres-Ocampo, A.P., Nakamura, S. and Morita, Y.S. (2017) The cell envelope-associated phospholipid-binding protein LmeA is required for mannan polymerization in mycobacteria. *J Biol Chem*, **292**, 17407-17417.

142. Yeow, J. and Chng, S.S. (2022) Of zones, bridges and chaperones - phospholipid transport in bacterial outer membrane assembly and homeostasis. *Microbiology (Reading)*, **168**.

143. Kadirvelraj, R., Foley, B.L., Dyekjaer, J.D. and Woods, R.J. (2008) Involvement of water in carbohydrate-protein binding: concanavalin A revisited. *J Am Chem Soc*, **130**, 16933-16942.

144. Flint, J., Nurizzo, D., Harding, S.E., Longman, E., Davies, G.J., Gilbert, H.J. and Bolam, D.N. (2004) Ligand-mediated dimerization of a carbohydrate-binding molecule reveals a novel mechanism for protein-carbohydrate recognition. *J Mol Biol*, **337**, 417-426.

145. Murzin, A.G. (1993) OB(oligonucleotide/oligosaccharide binding)-fold: common structural and functional solution for non-homologous sequences. *EMBO J*, **12**, 861-867.

146. Williams, A.F. and Barclay, A.N. (1988) The immunoglobulin superfamily--domains for cell surface recognition. *Annu Rev Immunol*, **6**, 381-405.

147. Chen, C.K., Chan, N.L. and Wang, A.H. (2011) The many blades of the beta-propeller proteins: conserved but versatile. *Trends Biochem Sci*, **36**, 553-561.

148. Babu, M.M., Priya, M.L., Selvan, A.T., Madera, M., Gough, J., Aravind, L. and Sankaran, K. (2006) A database of bacterial lipoproteins (DOLOP) with functional assignments to predicted lipoproteins. *J Bacteriol*, **188**, 2761-2773.

149. Sankaran, K. and Wu, H.C. (1994) Lipid modification of bacterial prolipoprotein. Transfer of diacylglyceryl moiety from phosphatidylglycerol. *J Biol Chem*, **269**, 19701-19706.

150. Tjalsma, H., Zanen, G., Venema, G., Bron, S. and van Dijl, J.M. (1999) The potential active site of
the lipoprotein-specific (type II) signal peptidase of Bacillus subtilis. *J Biol Chem*, **274**, 28191-
28197.
151. Notenboom, V., Boraston, A.B., Kilburn, D.G. and Rose, D.R. (2001) Crystal structures of the
family 9 carbohydrate-binding module from Thermotoga maritima xylanase 10A in native
and ligand-bound forms. *Biochemistry*, **40**, 6248-6256.
152. Rigden, D.J. (2005) Analysis of glycoside hydrolase family 98: catalytic machinery, mechanism
and a novel putative carbohydrate binding module. *FEBS Lett*, **579**, 5466-5472.
153. Boraston, A.B., Bolam, D.N., Gilbert, H.J. and Davies, G.J. (2004) Carbohydrate-binding
modules: fine-tuning polysaccharide recognition. *Biochem J*, **382**, 769-781.
154. Iyer, L.M., Koonin, E.V. and Aravind, L. (2001) Adaptations of the helix-grip fold for ligand
binding and catalysis in the START domain superfamily. *Proteins*, **43**, 134-144.
155. Santelli, E., Liddington, R.C., Mohan, M.A., Hoch, J.A. and Szurmant, H. (2007) The crystal
structure of Bacillus subtilis YycI reveals a common fold for two members of an unusual class
of sensor histidine kinase regulatory proteins. *J Bacteriol*, **189**, 3290-3295.
156. Das, K., Xiao, R., Wahlberg, E., Hsu, F., Arrowsmith, C.H., Montelione, G.T. and Arnold, E. (2001)
X-ray crystal structure of MTH938 from Methanobacterium thermoautotrophicum at 2.2 A
resolution reveals a novel tertiary protein fold. *Proteins*, **45**, 486-488.
157. Bennett, H.J., Davenport, J.B., Collins, R.F., Trafford, A.W., Pinali, C. and Kitmitto, A. (2013)
Human junctophilin-2 undergoes a structural rearrangement upon binding PtdIns(3,4,5)P3
and the S101R mutation identified in hypertrophic cardiomyopathy obviates this response.
Biochem J, **456**, 205-217.
158. Cortajarena, A.L. and Regan, L. (2006) Ligand binding by TPR domains. *Protein Sci*, **15**, 1193-
1198.
159. Seed, K.D., Faruque, S.M., Mekalanos, J.J., Calderwood, S.B., Qadri, F. and Camilli, A. (2012)
Phase variable O antigen biosynthetic genes control expression of the major protective
antigen and bacteriophage receptor in Vibrio cholerae O1. *PLoS Pathog*, **8**, e1002917.
160. Cai, R., Wang, G., Le, S., Wu, M., Cheng, M., Guo, Z., Ji, Y., Xi, H., Zhao, C., Wang, X., Xue, Y.,
Wang, Z., Zhang, H., Fu, Y., Sun, C., Feng, X., Lei, L., Yang, Y., Ur Rahman, S., Liu, X., Han, W.
and Gu, J. (2019) Three Capsular Polysaccharide Synthesis-Related Glucosyltransferases, GT-1,
GT-2 and WcaJ, Are Associated With Virulence and Phage Sensitivity of Klebsiella
pneumoniae. *Front Microbiol*, **10**, 1189.
161. Yau, K.W. (1994) Cyclic nucleotide-gated channels: an expanding new family of ion channels.
Proc Natl Acad Sci U S A, **91**, 3481-3483.
162. Durocher, D., Henckel, J., Fersht, A.R. and Jackson, S.P. (1999) The FHA domain is a modular
phosphopeptide recognition motif. *Mol Cell*, **4**, 387-394.
163. Schreiter, E.R. and Drennan, C.L. (2007) Ribbon-helix-helix transcription factors: variations on
a theme. *Nat Rev Microbiol*, **5**, 710-720.

164.   Ravi, J., Anantharaman, V., Chen, S.Z., Brenner, E.P., Datta, P., Aravind, L. and Gennaro, M.L.
1540   (2024) The phage shock protein (PSP) envelope stress response: discovery of novel partners
1541   and evolutionary history. *mSystems*, **9**, e0084723.

1542   165.   AhYoung, A.P., Koehl, A., Vizcarra, C.L., Cascio, D. and Egea, P.F. (2016) Structure of a putative
1543   ClpS N-end rule adaptor protein from the malaria pathogen Plasmodium falciparum. *Protein*
1544   *Sci*, **25**, 689-701.

1545   166.   Durr, I.F. and Rudney, H. (1960) The reduction of beta-hydroxy-beta-methyl-glutaryl
1546   coenzyme A to mevalonic acid. *J Biol Chem*, **235**, 2572-2578.

1547   167.   Tchen, T.T. (1958) Mevalonic kinase: purification and properties. *J Biol Chem*, **233**, 1100-1103.

1548   168.   Wolff, M., Seemann, M., Tse Sum Bui, B., Frapart, Y., Tritsch, D., Garcia Estrabot, A., Rodriguez-
1549   Concepcion, M., Boronat, A., Marquet, A. and Rohmer, M. (2003) Isoprenoid biosynthesis via
1550   the methylerythritol phosphate pathway: the (E)-4-hydroxy-3-methylbut-2-enyl diphosphate
1551   reductase (LytB/IspH) from Escherichia coli is a [4Fe-4S] protein. *FEBS Lett*, **541**, 115-120.

1552   169.   Guo, R.T., Ko, T.P., Chen, A.P., Kuo, C.J., Wang, A.H. and Liang, P.H. (2005) Crystal structures of
1553   undecaprenyl pyrophosphate synthase in complex with magnesium, isopentenyl
1554   pyrophosphate, and farnesyl thiopyrophosphate: roles of the metal ion and conserved
1555   residues in catalysis. *J Biol Chem*, **280**, 20762-20774.

1556   170.   Burroughs, A.M. and Aravind, L. (2016) RNA damage in biological conflicts and the diversity of
1557   responding RNA repair systems. *Nucleic Acids Res*, **44**, 8525-8555.

1558   171.   Ueta, M., Ohniwa, R.L., Yoshida, H., Maki, Y., Wada, C. and Wada, A. (2008) Role of HPF
1559   (hibernation promoting factor) in translational activity in Escherichia coli. *J Biochem*, **143**,
1560   425-433.

1561   172.   Akanuma, G., Kazo, Y., Tagami, K., Hiraoka, H., Yano, K., Suzuki, S., Hanai, R., Nanamiya, H.,
1562   Kato-Yamada, Y. and Kawamura, F. (2016) Ribosome dimerization is essential for the efficient
1563   regrowth of Bacillus subtilis. *Microbiology (Reading)*, **162**, 448-458.

1564   173.   Kuk, A.C.Y., Hao, A. and Lee, S.Y. (2022) Structure and Mechanism of the Lipid Flippase MurJ.
1565   *Annu Rev Biochem*, **91**, 705-729.

1566   174.   Zhang, D., de Souza, R.F., Anantharaman, V., Iyer, L.M. and Aravind, L. (2012) Polymorphic
1567   toxin systems: Comprehensive characterization of trafficking modes, processing, mechanisms
1568   of action, immunity and ecology using comparative genomics. *Biol Direct*, **7**, 18.

1569   175.   Iyer, L.M., Zhang, D., Rogozin, I.B. and Aravind, L. (2011) Evolution of the deaminase fold and
1570   multiple origins of eukaryotic editing and mutagenic nucleic acid deaminases from bacterial
1571   toxin systems. *Nucleic Acids Res*, **39**, 9473-9497.

1572   176.   Ruhe, Z.C., Low, D.A. and Hayes, C.S. (2020) Polymorphic Toxins and Their Immunity Proteins:
1573   Diversity, Evolution, and Mechanisms of Delivery. *Annu Rev Microbiol*, **74**, 497-520.

1574   177.   Alcoforado Diniz, J. and Coulthurst, S.J. (2015) Intraspecies Competition in Serratia
1575   marcescens Is Mediated by Type VI-Secreted Rhs Effectors and a Conserved Effector-
1576   Associated Accessory Protein. *J Bacteriol*, **197**, 2350-2360.

1577   178.   Satchell, K.J. (2011) Structure and function of MARTX toxins and other large repetitive RTX
1578   proteins. *Annu Rev Microbiol*, **65**, 71-90.

179. Kordis, D. and Turk, V. (2009) Phylogenomic analysis of the cystatin superfamily in eukaryotes
and prokaryotes. *BMC Evol Biol*, **9**, 266.
180. Cherier, D., Patin, D., Blanot, D., Touze, T. and Barreteau, H. (2021) The Biology of Colicin M
and Its Orthologs. *Antibiotics (Basel)*, **10**.
181. Anantharaman, V. and Aravind, L. (2003) Evolutionary history, structural features and
biochemical diversity of the NlpC/P60 superfamily of enzymes. *Genome Biol*, **4**, R11.
182. Arthur, M., Molinas, C. and Courvalin, P. (1992) Sequence of the vanY gene required for
production of a vancomycin-inducible D,D-carboxypeptidase in Enterococcus faecium
BM4147. *Gene*, **120**, 111-114.
183. Turk, V. and Bode, W. (1991) The cystatins: protein inhibitors of cysteine proteinases. *FEBS*
Lett, **285**, 213-219.
184. Sato, Y., Yamamoto, Y. and Kizaki, H. (1997) Cloning and sequence analysis of the gbpC gene
encoding a novel glucan-binding protein of Streptococcus mutans. *Infect Immun*, **65**, 668-675.
185. MacKenzie, D.A., Tailford, L.E., Hemmings, A.M. and Juge, N. (2009) Crystal structure of a
mucus-binding protein repeat reveals an unexpected functional immunoglobulin binding
activity. *J Biol Chem*, **284**, 32444-32453.
186. Chagnot, C., Zorgani, M.A., Astruc, T. and Desvaux, M. (2013) Proteinaceous determinants of
surface colonization in bacteria: bacterial adhesion and biofilm formation from a protein
secretion perspective. *Front Microbiol*, **4**, 303.
187. Wu, H.Y., Liu, M.S., Lin, T.P. and Cheng, Y.S. (2011) Structural and functional assays of
AtTLP18.3 identify its novel acid phosphatase activity in thylakoid lumen. *Plant Physiol*, **157**,
1015-1025.
188. Cao, Z., Casabona, M.G., Kneuper, H., Chalmers, J.D. and Palmer, T. (2016) The type VII
secretion system of Staphylococcus aureus secretes a nuclease toxin that targets competitor
bacteria. *Nat Microbiol*, **2**, 16183.
189. Klein, T.A., Pazos, M., Surette, M.G., Vollmer, W. and Whitney, J.C. (2018) Molecular Basis for
Immunity Protein Recognition of a Type VII Secretion System Exported Antibacterial Toxin. *J*
Mol Biol, **430**, 4344-4358.
190. Knowles, T.J., Browning, D.F., Jeeves, M., Maderbocus, R., Rajesh, S., Sridhar, P., Manoli, E.,
Emery, D., Sommer, U., Spencer, A., Leyton, D.L., Squire, D., Chaudhuri, R.R., Viant, M.R.,
Cunningham, A.F., Henderson, I.R. and Overduin, M. (2011) Structure and function of BamE
within the outer membrane and the beta-barrel assembly machine. *EMBO Rep*, **12**, 123-128.
191. Hagan, C.L. and Kahne, D. (2011) The reconstituted Escherichia coli Bam complex catalyzes
multiple rounds of beta-barrel assembly. *Biochemistry*, **50**, 7444-7446.
192. Kitagawa, D., Vakonakis, I., Olieric, N., Hilbert, M., Keller, D., Olieric, V., Bortfeld, M., Erat,
M.C., Fluckiger, I., Gonczy, P. and Steinmetz, M.O. (2011) Structural basis of the 9-fold
symmetry of centrioles. *Cell*, **144**, 364-375.
193. Aravind, L., Anantharaman, V., Zhang, D., de Souza, R.F. and Iyer, L.M. (2012) Gene flow and
biological conflict systems in the origin and evolution of eukaryotes. *Front Cell Infect Microbiol*,
2, 89.

194. Mutschler, H., Gebhardt, M., Shoeman, R.L. and Meinhart, A. (2011) A novel mechanism of
programmed cell death in bacteria by toxin-antitoxin systems corrupts peptidoglycan
synthesis. *PLoS Biol*, **9**, e1001033.

195. Monzingo, A.F., Marcotte, E.M., Hart, P.J. and Robertus, J.D. (1996) Chitinases, chitosanases,
and lysozymes can be divided into procaryotic and eucaryotic families sharing a conserved
core. *Nat Struct Biol*, **3**, 133-140.

196. Dong, X., Leksa, N.C., Chhabra, E.S., Arndt, J.W., Lu, Q., Knockenhauer, K.E., Peters, R.T. and
Springer, T.A. (2019) The von Willebrand factor D'D3 assembly and structural principles for
factor VIII binding and concatemer biogenesis. *Blood*, **133**, 1523-1533.

197. West, A.H. and Stock, A.M. (2001) Histidine kinases and response regulator proteins in two-
component signaling systems. *Trends Biochem Sci*, **26**, 369-376.

198. Costa, T., Isidro, A.L., Moran, C.P., Jr. and Henriques, A.O. (2006) Interaction between coat
morphogenetic proteins SafA and SpoVID. *J Bacteriol*, **188**, 7731-7741.

199. Zhang, C., Kelkar, A., Nasirikenari, M., Lau, J.T.Y., Sveinsson, M., Sharma, U.C., Pokharel, S. and
Neelamegham, S. (2018) The physical spacing between the von Willebrand factor D'D3 and A1
domains regulates platelet adhesion in vitro and in vivo. *J Thromb Haemost*, **16**, 571-582.

200. Hermoso, J.A., Monterroso, B., Albert, A., Galan, B., Ahrazem, O., Garcia, P., Martinez-Ripoll,
M., Garcia, J.L. and Menendez, M. (2003) Structural basis for selective recognition of
pneumococcal cell wall by modular endolysin from phage Cp-1. *Structure*, **11**, 1239-1249.

201. Das, D., Lee, W.S., Grant, J.C., Chiu, H.J., Farr, C.L., Vance, J., Klock, H.E., Knuth, M.W., Miller,
M.D., Elsliger, M.A., Deacon, A.M., Godzik, A., Lesley, S.A., Kornfeld, S. and Wilson, I.A. (2013)
Structure and function of the DUF2233 domain in bacteria and in the human mannose 6-
phosphate uncovering enzyme. *J Biol Chem*, **288**, 16789-16799.

202. Dideberg, O., Charlier, P., Dive, G., Joris, B., Frere, J.M. and Ghuysen, J.M. (1982) Structure of a
Zn2+-containing D-alanyl-D-alanine-cleaving carboxypeptidase at 2.5 A resolution. *Nature*,
299, 469-470.

203. Bellinzoni, M., Haouz, A., Miras, I., Magnet, S., Andre-Leroux, G., Mukherjee, R., Shepard, W.,
Cole, S.T. and Alzari, P.M. (2014) Structural studies suggest a peptidoglycan hydrolase
function for the Mycobacterium tuberculosis Tat-secreted protein Rv2525c. *J Struct Biol*, **188**,
156-164.

204. Wright, G.D., Molinas, C., Arthur, M., Courvalin, P. and Walsh, C.T. (1992) Characterization of
vanY, a DD-carboxypeptidase from vancomycin-resistant Enterococcus faecium BM4147.
Antimicrob Agents Chemother, **36**, 1514-1518.

205. Arthur, M., Depardieu, F., Cabanie, L., Reynolds, P. and Courvalin, P. (1998) Requirement of the
VanY and VanX D,D-peptidases for glycopeptide resistance in enterococci. *Mol Microbiol*, **30**,
819-830.

206. Arthur, M., Depardieu, F., Snaith, H.A., Reynolds, P.E. and Courvalin, P. (1994) Contribution of
VanY D,D-carboxypeptidase to glycopeptide resistance in Enterococcus faecalis by hydrolysis
of peptidoglycan precursors. *Antimicrob Agents Chemother*, **38**, 1899-1903.

207. Meyer, B., Wurm, J.P., Sharma, S., Immer, C., Pogoryelov, D., Kotter, P., Lafontaine, D.L.,
Wohnert, J. and Entian, K.D. (2016) Ribosome biogenesis factor Tsr3 is the
aminocarboxypropyl transferase responsible for 18S rRNA hypermodification in yeast and
humans. *Nucleic Acids Res*, **44**, 4304-4316.

208. Burroughs, A.M. and Aravind, L. (2014) Analysis of two domains with novel RNA-processing
activities throws light on the complex evolution of ribosomal RNA biogenesis. *Front Genet*, **5**,
424.

209. Fraser, M.E., Joyce, M.A., Ryan, D.G. and Wolodko, W.T. (2002) Two glutamate residues, Glu 208
alpha and Glu 197 beta, are crucial for phosphorylation and dephosphorylation of the active-
site histidine residue in succinyl-CoA synthetase. *Biochemistry*, **41**, 537-546.

210. Iyer, L.M., Abhiman, S., Maxwell Burroughs, A. and Aravind, L. (2009) Amidoligases with ATP-
grasp, glutamine synthetase-like and acetyltransferase-like domains: synthesis of novel
metabolites and peptide modifications of proteins. *Mol Biosyst*, **5**, 1636-1660.

211. Raetz, C.R. and Kennedy, E.P. (1974) Partial purification and properties of phosphatidylserine
synthetase from Escherichia coli. *J Biol Chem*, **249**, 5083-5045.

212. Fields, R.N. and Roy, H. (2018) Deciphering the tRNA-dependent lipid aminoacylation systems
in bacteria: Novel components and structural advances. *RNA Biol*, **15**, 480-491.

213. Hancock, R.E. (1997) Peptide antibiotics. *Lancet*, **349**, 418-422.

214. Benson, T.E., Prince, D.B., Mutchler, V.T., Curry, K.A., Ho, A.M., Sarver, R.W., Hagadorn, J.C.,
Choi, G.H. and Garlick, R.L. (2002) X-ray crystal structure of Staphylococcus aureus FemA.
Structure, **10**, 1107-1115.

215. Schneider, T., Senn, M.M., Berger-Bachi, B., Tossi, A., Sahl, H.G. and Wiedemann, I. (2004) In
vitro assembly of a complete, pentaglycine interpeptide bridge containing cell wall precursor
(lipid II-Gly5) of Staphylococcus aureus. *Mol Microbiol*, **53**, 675-685.

216. Schmid, M., Uhlenhaut, N.H., Godard, F., Demar, M., Bressan, R., Weigel, D. and Lohmann, J.U.
(2003) Dissection of floral induction pathways using global expression analysis. *Development*,
130, 6001-6012.

217. Berardini, T.Z., Reiser, L., Li, D., Mezheritsky, Y., Muller, R., Strait, E. and Huala, E. (2015) The
Arabidopsis information resource: Making and mining the "gold standard" annotated
reference plant genome. *Genesis*, **53**, 474-485.

218. Savidov, N., Gloriozova, T.A., Poroikov, V.V. and Dembitsky, V.M. (2018) Highly oxygenated
isoprenoid lipids derived from fungi and fungal endophytes: Origin and biological activities.
Steroids, **140**, 114-124.

219. Lee, J.O., Rieu, P., Arnaout, M.A. and Liddington, R. (1995) Crystal structure of the A domain
from the alpha subunit of integrin CR3 (CD11b/CD18). *Cell*, **80**, 631-638.

220. Zhang, Y., van der Fits, L., Voerman, J.S., Melief, M.J., Laman, J.D., Wang, M., Wang, H., Wang,
M., Li, X., Walls, C.D., Gupta, D. and Dziarski, R. (2005) Identification of serum N-
acetylmuramoyl-l-alanine amidase as liver peptidoglycan recognition protein 2. *Biochim
Biophys Acta*, **1752**, 34-46.

221.   Lee, J., Geddes, K., Streutker, C., Philpott, D.J. and Girardin, S.E. (2012) Role of mouse
peptidoglycan recognition protein PGLYRP2 in the innate immune response to Salmonella
enterica serovar Typhimurium infection in vivo. *Infect Immun*, **80**, 2645-2654.
222.   Dziarski, R. and Gupta, D. (2010) Review: Mammalian peptidoglycan recognition proteins
(PGRPs) in innate immunity. *Innate Immun*, **16**, 168-174.
223.   Dziarski, R. and Gupta, D. (2006) Mammalian PGRPs: novel antibacterial proteins. *Cell*
Microbiol, **8**, 1059-1069.
224.   Jones, J.D. and Dangl, J.L. (2006) The plant immune system. *Nature*, **444**, 323-329.
225.   Holzem, M., Boutros, M. and Holstein, T.W. (2024) The origin and evolution of Wnt signalling.
Nat Rev Genet, **25**, 500-512.
226.   Morrow, J.F., Stearman, R.S., Peltzman, C.G. and Potter, D.A. (1981) Induction of hepatic
synthesis of serum amyloid A protein and actin. *Proc Natl Acad Sci U S A*, **78**, 4718-4722.
227.   Hari-Dass, R., Shah, C., Meyer, D.J. and Raynes, J.G. (2005) Serum amyloid A protein binds to
outer membrane protein A of gram-negative bacteria. *J Biol Chem*, **280**, 18562-18567.
228.   Shah, C., Hari-Dass, R. and Raynes, J.G. (2006) Serum amyloid A is an innate immune opsonin
for Gram-negative bacteria. *Blood*, **108**, 1751-1757.
229.   Sack, G.H., Jr., Talbot, C.C., Jr., Seuanez, H. and O'Brien, S.J. (1989) Molecular analysis of the
human serum amyloid A (SAA) gene family. *Scand J Immunol*, **29**, 113-119.
230.   Uhlar, C.M., Burgess, C.J., Sharp, P.M. and Whitehead, A.S. (1994) Evolution of the serum
amyloid A (SAA) protein superfamily. *Genomics*, **19**, 228-235.
231.   Sun, L. and Ye, R.D. (2016) Serum amyloid A1: Structure, function and gene polymorphism.
Gene, **583**, 48-57.
232.   Aoki, S.K., Malinverni, J.C., Jacoby, K., Thomas, B., Pamma, R., Trinh, B.N., Remers, S., Webb, J.,
Braaten, B.A., Silhavy, T.J. and Low, D.A. (2008) Contact-dependent growth inhibition requires
the essential outer membrane protein BamA (YaeT) as the receptor and the inner membrane
transport protein AcrB. *Mol Microbiol*, **70**, 323-340.
233.   Virtanen, P., Waneskog, M. and Koskiniemi, S. (2019) Class II contact-dependent growth
inhibition (CDI) systems allow for broad-range cross-species toxin delivery within the
Enterobacteriaceae family. *Mol Microbiol*, **111**, 1109-1125.
234.   Ruhe, Z.C., Nguyen, J.Y., Xiong, J., Koskiniemi, S., Beck, C.M., Perkins, B.R., Low, D.A. and
Hayes, C.S. (2017) CdiA Effectors Use Modular Receptor-Binding Domains To Recognize Target
Bacteria. *mBio*, **8**.
235.   Levin, M., Franklin, E.C., Frangione, B. and Pras, M. (1972) The amino acid sequence of a major
nonimmunoglobulin component of some amyloid fibrils. *J Clin Invest*, **51**, 2773-2776.
236.   Blanco, L.P., Evans, M.L., Smith, D.R., Badtke, M.P. and Chapman, M.R. (2012) Diversity,
biogenesis and function of microbial amyloids. *Trends Microbiol*, **20**, 66-73.
237.   Rouse, S.L., Matthews, S.J. and Dueholm, M.S. (2018) Ecology and Biogenesis of Functional
Amyloids in Pseudomonas. *J Mol Biol*, **430**, 3685-3695.

238. Tukel, C., Wilson, R.P., Nishimori, J.H., Pezeshki, M., Chromy, B.A. and Baumler, A.J. (2009) Responses to amyloids of microbial and host origin are mediated through toll-like receptor 2. *Cell Host Microbe*, **6**, 45-53.

239. Prosswimmer, T., Heng, A. and Daggett, V. (2024) Mechanistic insights into the role of amyloid-beta in innate immunity. *Sci Rep*, **14**, 5376.

240. Egan, A.J.F., Errington, J. and Vollmer, W. (2020) Regulation of peptidoglycan synthesis and remodelling. *Nat Rev Microbiol*, **18**, 446-460.

241. Koga, Y. and Morii, H. (2007) Biosynthesis of ether-type polar lipids in archaea and evolutionary considerations. *Microbiol Mol Biol Rev*, **71**, 97-120.

242. Daiyasu, H., Kuma, K., Yokoi, T., Morii, H., Koga, Y. and Toh, H. (2005) A study of archaeal enzymes involved in polar lipid synthesis linking amino acid sequence information, genomic contexts and lipid composition. *Archaea*, **1**, 399-410.

243. Carman, G.M. and Han, G.S. (2006) Roles of phosphatidate phosphatase enzymes in lipid metabolism. *Trends Biochem Sci*, **31**, 694-699.

244. Neuwald, A.F. (1997) An unexpected structural relationship between integral membrane phosphatases and soluble haloperoxidases. *Protein Sci*, **6**, 1764-1767.

245. Aravind, L., Nicastro, G.G., Iyer, L.M. and Burroughs, A.M. (2024) The Prokaryotic Roots of Eukaryotic Immune Systems. *Annu Rev Genet*, **58**, 365-389.

246. Krishnan, A., Iyer, L.M., Holland, S.J., Boehm, T. and Aravind, L. (2018) Diversification of AID/APOBEC-like deaminases in metazoa: multiplicity of clades and widespread roles in immunity. *Proc Natl Acad Sci U S A*, **115**, E3201-E3210.

247. Zhang, D., Iyer, L.M. and Aravind, L. (2011) A novel immunity system for bacterial nucleic acid degrading toxins and its recruitment in various eukaryotic and DNA viral systems. *Nucleic Acids Res*, **39**, 4532-4552.

248. Kaur, G., Burroughs, A.M., Iyer, L.M. and Aravind, L. (2020) Highly regulated, diversifying NTP-dependent biological conflict systems with implications for the emergence of multicellularity. *Elife*, **9**.

249. Kaur, G., Iyer, L.M., Burroughs, A.M. and Aravind, L. (2021) Bacterial death and TRADD-N domains help define novel apoptosis and immunity mechanisms shared by prokaryotes and metazoans. *Elife*, **10**.

250. Altschul, S.F., Madden, T.L., Schaffer, A.A., Zhang, J., Zhang, Z., Miller, W. and Lipman, D.J. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res*, **25**, 3389-3402.

251. Johnson, L.S., Eddy, S.R. and Portugaly, E. (2010) Hidden Markov model speed heuristic and iterative HMM search procedure. *BMC Bioinformatics*, **11**, 431.

252. Sayers, E.W., Bolton, E.E., Brister, J.R., Canese, K., Chan, J., Comeau, D.C., Connor, R., Funk, K., Kelly, C., Kim, S., Madej, T., Marchler-Bauer, A., Lanczycki, C., Lathrop, S., Lu, Z., Thibaud-Nissen, F., Murphy, T., Phan, L., Skripchenko, Y., Tse, T., Wang, J., Williams, R., Trawick, B.W., Pruitt, K.D. and Sherry, S.T. (2022) Database resources of the national center for biotechnology information. *Nucleic Acids Res*, **50**, D20-D26.

253. Hauser, M., Steinegger, M. and Soding, J. (2016) MMseqs software suite for fast and deep
clustering and searching of large protein sequence sets. *Bioinformatics*, **32**, 1323-1330.
254. Katoh, K. and Standley, D.M. (2013) MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol*, **30**, 772-780.
255. Steinegger, M., Meier, M., Mirdita, M., Vohringer, H., Haunsberger, S.J. and Soding, J. (2019)
HH-suite3 for fast remote homology detection and deep protein annotation. *BMC*
Bioinformatics, **20**, 473.
256. Csardi, G. and Nepusz, T. (2006) The Igraph Software Package for Complex Network Research.
InterJournal, Complex Systems.
257. Hagberg, A.A., Schult, D.A., Swart, P. and Hagberg, J.M. (2008), *Proceedings of the Python in*
Science Conference.
258. Finn, R.D., Coggill, P., Eberhardt, R.Y., Eddy, S.R., Mistry, J., Mitchell, A.L., Potter, S.C., Punta,
M., Qureshi, M., Sangrador-Vegas, A., Salazar, G.A., Tate, J. and Bateman, A. (2016) The Pfam
protein families database: towards a more sustainable future. *Nucleic Acids Res*, **44**, D279-285.
259. Schaffer, A.A., Wolf, Y.I., Ponting, C.P., Koonin, E.V., Aravind, L. and Altschul, S.F. (1999)
IMPALA: matching a protein sequence against a collection of PSI-BLAST-constructed position-
specific score matrices. *Bioinformatics*, **15**, 1000-1011.
260. Eddy, S.R. (2011) Accelerated Profile HMM Searches. *PLoS Comput Biol*, **7**, e1002195.
261. Soding, J., Biegert, A. and Lupas, A.N. (2005) The HHpred interactive server for protein
homology detection and structure prediction. *Nucleic Acids Res*, **33**, W244-248.
262. Berman, H., Henrick, K., Nakamura, H. and Markley, J.L. (2007) The worldwide Protein Data
Bank (wwPDB): ensuring a single, uniform archive of PDB data. *Nucleic Acids Res*, **35**, D301-303.
263. Price, M.N., Dehal, P.S. and Arkin, A.P. (2010) FastTree 2--approximately maximum-likelihood
trees for large alignments. *PLoS One*, **5**, e9490.
264. Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A. and
Lanfear, R. (2020) IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in
the Genomic Era. *Mol Biol Evol*, **37**, 1530-1534.
265. Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O., Willmore,
L., Ballard, A.J., Bambrick, J., Bodenstein, S.W., Evans, D.A., Hung, C.C., O'Neill, M., Reiman, D.,
Tunyasuvunakool, K., Wu, Z., Zemgulyte, A., Arvaniti, E., Beattie, C., Bertolli, O., Bridgland, A.,
Cherepanov, A., Congreve, M., Cowen-Rivers, A.I., Cowie, A., Figurnov, M., Fuchs, F.B.,
Gladman, H., Jain, R., Khan, Y.A., Low, C.M.R., Perlin, K., Potapenko, A., Savy, P., Singh, S.,
Stecula, A., Thillaisundaram, A., Tong, C., Yakneen, S., Zhong, E.D., Zielinski, M., Zidek, A.,
Bapst, V., Kohli, P., Jaderberg, M., Hassabis, D. and Jumper, J.M. (2024) Accurate structure
prediction of biomolecular interactions with AlphaFold 3. *Nature*, **630**, 493-500.
266. Sehnal, D., Bittrich, S., Deshpande, M., Svobodova, R., Berka, K., Bazgier, V., Velankar, S.,
Burley, S.K., Koca, J. and Rose, A.S. (2021) Mol* Viewer: modern web app for 3D visualization
and analysis of large biomolecular structures. *Nucleic Acids Res*, **49**, W431-W437.
267. Holm, L. (2019) Benchmarking fold detection by DaliLite v.5. *Bioinformatics*, **35**, 5326-5327.

268.   van Kempen, M., Kim, S.S., Tumescheit, C., Mirdita, M., Lee, J., Gilchrist, C.L.M., Soding, J. and
Steinegger, M. (2024) Fast and accurate protein structure search with Foldseek. *Nat Biotechnol*,
42, 243-246.

1817

Figure 1

A Legend: blue - Helix-1; green - Helix-2; yellow - Helix-3; orange - Helix-4; pink - Inserts and N-, C-terminal extensions

Wnt — PDB:7DRT — Fifth helix; Cysteine-rich insertions/extensions; Palmitoleoylated serine

Min-Wnt — WP_095416411.1 — Fifth helix; N; C

Prok-SAA — WP_235649002.1 — Fifth helix; N

VanZ — WP_231507663.1 — Handle; N; C

B Network nodes: cpCone-Loki; cpCone-DUF2585; PTDSS1/2; pPTDSS; cpCone-2; cpCone-1; Wok-DUF2238; CapCone-DUF4056; CapCone-1; Met-SAA; Met-Wnt; SAA-prok; Min-Wnt; ClaspCone-3; ClaspCone-1; YfiM-1; VanZ-2; VanZ-1; YfiM-DUF2279; ClaspCone-2; CapCone-2; Skillet-4; Skillet-1; Skillet-DUF2809; YfiM-Griddle; TelC-prok; Met-TelC; Skillet-2; Skillet-3

C Transmembrane tendency plot

Y-axis: Normalized transmembrane tendency score

Higher hydrophobicity; Higher hydrophilicity

X-axis (Lipocone families): cpCone-DUF2585; cpCone-Loki; cpCone-1; cpCone-2; cpCone-i; Prok-PTDSS; PTDSS1/2; Wok-DUF2238; VanZ-1; VanZ-2; Skillet-DUF2809; Skillet-1; Skillet-4; Skillet-2; Skillet-3; YfiM-1; YfiM-DUF2279; YfiM-Griddle; Min-Wnt; Met-Wnt; Met-SAA; Prok-SAA; CapCone-1; CapCone-2; CapCone-DUF4056; ClaspCone-2; ClaspCone-1; ClaspCone-3; Prok-TelC; Met-TelC

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6