1

Decoding MASLD Progression: A Molecular Trajectory-Based Framework for Modelling Disease Dynamics

Ioannis Kamzolas[1,4*], Thodoris Koutsandreas[1,2*], Charlie George Barker[1,3], Anna Vathrakokoili Pournara[1], Harry Weston[1,4], Michele Vacca[4,5], Irene Papatheodorou[1,6], Antonio Vidal-Puig[4,7 +], Evangelia Petsalaki[1+]

[1]European Molecular Biology Laboratory - European Bioinformatics Institute, Wellcome Genome

Campus, Hinxton, CB10 1SD, UK

[2]Open Targets, Wellcome Genome Campus, Hinxton, CB10 1SD, UK

[3](current address) University College London Cancer Institute, London WC1E 6DD, UK

[4]MRC Institute of Metabolic Science, Metabolic Research Laboratories, University of Cambridge,

Cambridge, UK

[5]Interdisciplinary Department of Medicine, Università degli Studi di Bari "Aldo Moro", Bari, Italy;

Roger Williams Institute of Liver Studies, Foundation for Liver Research, London, SE5 9NT, UK

[6] (current address) Earlham Institute, Norwich Research Park, Colney Ln, Norwich NR4 7UZ

[7]Centro de Investigacion Principe Felipe, Valencia, Spain

+ correspondence should be addressed to Antonio Vidal-Puig ajv22@medschl.cam.ac.uk and

Evangelia Petsalaki: petsalaki@ebi.ac.uk

* These authors contributed equally to the work presented in this manuscript

Abstract

Metabolic Dysfunction-Associated Steatotic Liver Disease (MASLD) has emerged as a silent pandemic, affecting nearly one-third of the global population. This condition encompasses a spectrum of liver disorders, ranging from simple steatosis to Metabolic Dysfunction-Associated Steatohepatitis (MASH), which is characterised by liver steatosis, lipotoxicity, hepatocellular damage, inflammation, and fibrosis. Left unchecked, MASLD/MASH can progress to cirrhosis and hepatocellular carcinoma. Despite the progressive nature of MASLD/MASH, current research primarily relies on static, histopathologically defined stages, which fail to capture the dynamic disease continuum.

In this study, we present an integrative stratification approach that combines patient pseudo-temporal ordering, network analysis, and cell-type deconvolution to map the continuous disease trajectory. By analyzing transcriptomic profiles, we predict patients' positions along this trajectory, moving beyond conventional stage-based classifications. This approach reveals the sequence of critical molecular events driving MASLD/MASH progression, providing new insights into the disease's pathophysiology. Furthermore, we identify novel trajectory-specific biomarkers that support a more refined, personalised strategy for managing MASLD. This work highlights the potential of trajectory-based frameworks in advancing our understanding and treatment of complex metabolic diseases.

Introduction

Metabolic Dysfunction-Associated Steatotic Liver Disease (MASLD) encompasses a spectrum of conditions, ranging from simple fat accumulation (steatosis) to Metabolic Dysfunction-Associated Steatohepatitis (MASH), which can progress to fibrosis, cirrhosis, and hepatocellular carcinoma[1]. This global epidemic affects nearly one-third of Western populations and is primarily driven by the rising prevalence of metabolic syndrome, type 2 diabetes, and obesity[2,3]. Despite its significant burden, most MASLD cases remain asymptomatic and undiagnosed until advanced stages.
The diagnosis of MASLD typically relies on imaging techniques such as ultrasound and MRI spectroscopy alongside non-invasive fibrosis markers like the Fibrosis-4 index (FIB-4), with liver biopsy serving as the gold standard[4]. Although histological scoring reproducibility among

pathologists has improved[5], variability in sampling[6,7] and the invasiveness of biopsies[6,7]

underscore the urgent need for complementary non-invasive diagnostic methods.

Elastography techniques, including transient elastography (FibroScan) and magnetic resonance

elastography (MRE), have emerged as valuable tools for non-invasive fibrosis assessment. While

MRE offers high accuracy, its cost and limited accessibility are significant barriers. Conversely,

transient elastography is more affordable but less precise. Despite the availability of various

blood-based imaging and genetic biomarkers for detecting steatosis and fibrosis, their accuracy

remains suboptimal, and reliable options for diagnosing MASLD specifically are still lacking[4].

Advances in systems biology and omics technologies have propelled MASLD/MASH research

forward[4,8–11], linking the disease to metabolic syndrome, insulin resistance, and mechanisms such

as lipid accumulation, endoplasmic reticulum stress, and oxidative stress[12]. Genome-Wide

Association Studies (GWAS) have further highlighted genetic factors, such as the PNPLA3 SNP

rs738409 variant, which is associated with increased liver fat and heightened disease risk [13–15].

Despite these breakthroughs, the complexity of MASLD demands further research to elucidate

progression mechanisms and distinguish benign cases from those with higher mortality risks.

Histology has long been a cornerstone for understanding MASLD pathophysiology, but its

limitations as a sole proxy for disease mechanisms are increasingly evident. Pseudo-temporal

ordering, initially developed for microarray data[16] and now widely applied in single-cell genomics,

provides an alternative framework by arranging cells or patients along trajectories based on their

molecular states. This approach has recently been used to study Alzheimer's disease progression

through brain expression data[17,18], and offers a promising, histology-independent method for

investigating MASLD progression via transcriptomic profiles. By complementing traditional

assessments, pseudo-temporal ordering provides a data-driven perspective on the molecular

changes underpinning disease progression.

In this study, we applied pseudo-temporal ordering to MASLD/MASH patient data (Fig. 1A). We

integrated it with network analysis (Fig. 1B) and cell-type deconvolution (Fig. 1C). This

comprehensive approach allowed us to identify key molecular events specific to disease stages

and cell types. Additionally, by integrating our molecular map of MASLD progression with plasma

proteomics data, we identified liver-specific circulating biomarkers that can position patients along

the disease trajectory independently of histology. These findings not only provide new insights

into MASLD pathophysiology but also prioritize potential therapeutic targets associated with

disease progression (Figs. 1D–E).

Figure 1. Overview of approach used in this study to investigate MASLD progression. A. *Integration of public datasets and MASLD trajectory inference by pseudo-temporal ordering of patient biopsy transcriptomic data* ***B.*** *Combining phenotype-specific gene co-expression analysis with upstream pathway inference to define a MASLD network.* ***C.*** *Cell type deconvolution to study the dynamics of cell type composition changes.* ***D.*** *Random Forest regression model for the prediction of patients' position on the inferred MASLD trajectory based on a set of biomarkers. Training has been done on the initial dataset and testing on independent datasets. The model deployed different gene signatures, while random gene sets were used as controls to validate the high model's performance.* ***E.*** *Identification of deregulated cell type-specific processes and potential biomarker genes that could be targeted in specific disease stages.*

Results

Transcriptomics-based disease trajectory analysis captures MASLD progression.

We hypothesised that MASLD progression could be modelled as a continuum of patient states. To test this, we analysed RNA-Seq data from 136 patients[9,19] (Methods; Supplementary Table 1 & 2). After excluding one outlier, we applied pseudo-temporal ordering[20] to construct a transcriptomics-driven trajectory (Methods; Fig. 2A; Supplementary Fig. 1). This trajectory showed strong correlations with histopathology scores, including steatosis, ballooning, inflammation, fibrosis, and the NAS score, with Pearson R coefficients ranging from 0.96 to 1.0. These findings confirmed alignment between the transcriptomics-derived trajectory and established disease stages (ANOVA followed by Tukey's pairwise comparisons; Supplementary Fig. 2). (Methods; Supplementary Tables 1 & 2).

To validate the trajectory model, we tested it on two independent RNA-Seq datasets: the EPoS dataset, which includes healthy normal-weight and obese individuals, and the Gubra dataset, comprising MASLD and MASH patients (Supplementary Figs. 3A-B)[8,21]. In the EPoS dataset, the trajectory successfully captured disease stage progression. In the Gubra dataset it effectively distinguished healthy and obese samples from MASLD and MASH patients. However, additional variance in both datasets was observed, which did not fully correlate with disease progression.

To address this variability, we identified genes contributing to variance along the MASLD/MASH progression axis. Using a random forest approach, we identified 145 genes that significantly predicted MASLD stage and histopathological features (AUC: 62–73%; Supplementary Figs. 3C-D, Supplementary Table 3). These genes also recapitulated the disease trajectory in the independent datasets[8,21], demonstrating improved linearity and better differentiation of MASLD/MASH stages compared to the full transcriptome (Figs. 2B-C, Supplementary Fig. 3E). This underscores the value of targeted gene sets in refining trajectory models and improving stage prediction.

The identified genes were enriched in pathways central to MASLD/MASH progression, including ECM-receptor interaction and focal adhesion (associated with fibrosis), AGE-RAGE and IL-17 signalling (inflammatory responses), and TGF-beta and PI3K-Akt signalling (wound healing and

115 metabolic dysregulation). Other pathways, such as glycerolipid metabolism and cytokine-receptor
116 interaction, reflect the metabolic and immune dysregulation characteristic of MASLD/MASH (Fig.
117 2D). To further elucidate MASLD/MASH progression mechanisms, we developed a network-
118 based strategy, integrating molecular interactions and pathway analyses to identify critical drivers
119 of disease states.
120

*Figure 2. Pseudo-temporal ordering of patients captures disease progression and identifies MASLD gene signature. A. Pseudo-temporal ordering of patients based on transcriptomic data recapitulates disease progression based on individual phenotypes, i.e. steatosis, ballooning, inflammation and fibrosis, and based on the NAS and MASLD scores. B-C. Pseudotemporal ordering of two independent orthogonal datasets based on the 145 genes that are most predictive of the trajectory provides a linear and clear separation of the disease stages (EPoS dataset (B) and Gubra dataset (C)). D. Functional enrichment analysis results of these 145 genes using EnrichR, where the Combined Score (c) has been calculated as c = log(p_value) * z_score, with p_value representing the Fisher exact test outcome, and z-score the deviation from the expected rank.*

Data-Driven Global MASLD/MASH Network Recapitulates Key Molecular Mechanisms of Disease

To construct a MASLD reference network, we adapted our previously published method for generating phenotype-specific networks by integrating paired transcriptomics and phenotype data[22] (Methods). Using Weighted Gene Co-expression Network Analysis (WGCNA)[23], we identified gene co-expression modules and linked them to key phenotypic features, including the NAS score, steatosis, ballooning, and fibrosis (Fig. 3A; Supplementary Table 4). Linear regression analysis revealed modules that were either uniquely associated with specific phenotypes or shared across multiple phenotypes, offering valuable insights into the molecular underpinnings of MASLD progression.

Reactome pathway enrichment analysis[24] of the co-expression modules identified both shared and distinct mechanisms associated with MASLD progression. Several modules were enriched in pathways known to play central roles in MASLD-related metabolic processes (Supplementary Table 5). The MEbrown module was primarily enriched in protein translation pathways, while the MEblue module, characterised by its abundance of zinc-finger proteins, was linked to gene expression regulation. The MEred and MEsalmon modules were strongly associated with extracellular matrix organization and signal transduction, pathways critical for fibrosis and cellular communication. The MEtan module was tied to muscle contraction processes, suggesting potential roles in metabolic or structural regulation. Notably, the MEyellow module was exclusively enriched in pathways related to the immune system and cytokine signalling, including processes such as "antigen processing and cross-presentation", "interferon signalling", "interleukin signalling", and "immunoregulatory interactions between lymphoid and non-lymphoid cells". These findings highlight the specialized roles of these co-expression modules in MASLD progression, reflecting the complexity of the disease's molecular mechanisms.

Transcription factor (TF) enrichment analysis of significant gene modules identified 188 TF regulons across eight gene modules (adjusted p-value < 0.05) (Supplementary Table 6). Among these, 48 TFs were enriched in multiple modules, underscoring their pivotal roles in MASLD progression (Fig. 3B, Supplementary Table 6). Many of these TFs were linked to TGFβ1 signalling, a key driver of fibrosis through both SMAD-dependent and SMAD-independent pathways. For instance, the regulatory networks of SP1, SRF, and ETS1 interact with TGFβ1

152 signalling to promote the expression of cytoskeletal and ECM proteins, including actins and
153 integrins, which are critical to tissue remodelling and fibrosis[25–27].

Figure 3. Phenotype-specific gene modules associated with the different MASLD variables.
A. *Network of the different gene co-expression modules associated with NAS score, Steatosis,*
Ballooning, and Fibrosis. Each module is depicted by a different colour and can be linked with one
or multiple variables. ***B.*** *TFs whose regulons are enriched in the modules. The figure shows TFs*
per module coloured according to the adjusted p-value of the enrichment test. The presented TFs
have been identified as enriched in multiple modules and deregulated at least in one stage of the
disease trajectory (see sliding window analysis results). The full list of TFs derived from the
enrichment analysis is given in Supplementary Table 6. ***C.*** *Enrichment of MASLD network and its*
components in known MASLD genes curated from the literature (top) or from MSigDB (bottom).

EGR1, another TF regulated by TGFβ1 signalling, has been implicated in both steatosis and
fibrosis [28,29,30]. Similarly, SMAD3 and SMAD4, intracellular effectors of canonical TGFβ1/SMAD
signalling, were found to be deregulated and enriched in the MEred module, indicating a potential
disruption of the TGFβ1 feedback loop. Beyond TGFβ1 signalling, other deregulated TFs relevant

to MASLD include CREB1[31], NFKB1[32], and HIF1A[33,34], which are known to contribute to metabolic
and inflammatory pathways characteristic of MASLD progression (Supplementary Table 6).

Reactome pathway enrichment analysis[24] of TFs revealed the molecular pathways in which they
are known to participate (Supplementary Table 7). TF sets associated with steatosis and the NAS
score were enriched in pathways related to developmental biology, myogenesis, gene expression
and processes that are particularly relevant to the activation of hepatic stellate cells, a key driver
of fibrosis[35]. In contrast, TF sets linked to fibrosis, ballooning, and the NAS score exhibited
dysregulation in stress response and signalling pathways, including PUMA activation, FOXO-
mediated transcription, and Toll-like receptor cascades, all of which play critical roles in cellular
stress, apoptosis, and inflammatory responses during MASLD progression reflecting the liver's
response to injury.

Fibrosis-specific pathways, such as "Adherens junction interactions" and "Regulation of TP53
activity", underscore changes in cellular architecture and transcription that drive tissue
remodelling during fibrosis. These findings highlight the importance of targeted transcriptional
regulation and subtle alterations within specific cellular pathways in shaping the complex
dynamics of MASLD/MASH progression.

Leveraging this information, we constructed a comprehensive MASLD/MASH disease network to
examine dynamic changes in cellular processes along the disease trajectory (Methods;
Supplementary Table 8). The resulting network comprised 4,247 nodes and was significantly
enriched with genes previously associated with MASLD and integrated data from signalling and
metabolic pathways. Notably, the highest enrichment ratios were observed for the subnetworks
of ballooning and fibrosis (Fig. 3C; Supplementary Table 9). The inclusion of relevant transcription
factors, enriched pathways, and the statistically significant overlap of the network with known
MASLD/MASH-associated genes supports its robustness and suitability for downstream
analyses. This integrative framework provides a valuable tool for further investigation into the
molecular mechanisms driving MASLD/MASH progression.

Sliding Window Analysis Highlights Molecular Dysregulation in MASLD
Progression

To gain deeper insights into the molecular changes driving MASLD progression, we employed a
sliding window (SW) approach, dividing patients into groups along the disease trajectory. To

optimise the sequence of sliding windows, we developed a graph-based method that maximised

the information content within each window (Methods). Using this approach, patients were divided

into 13 subgroups along the trajectory for functional analysis (Methods; Supplementary Fig. 4;

Supplementary Tables 1 & 10). Analysis of genes within phenotype-associated modules revealed

patterns of collective dysregulation as the disease advanced into later stages (Supplementary

Fig. 5A). This approach enabled the identification of stage-specific molecular signatures and

provided a dynamic perspective on the progression of MASLD.

Initially, we identified transcription factors (TFs) that were differentially regulated along the

disease trajectory, demonstrating strong concordance with analyses based on traditional discrete

disease stages while providing greater granularity (Fig. 4A; Supplementary Table 11). Key

examples include HNF1A, a critical regulator of hepatic differentiation and metabolism[36], which

exhibited marked downregulation in sliding windows 5 and 13. This downregulation reflects

processes such as inflammation, epithelial-to-mesenchymal transition (EMT), and wound healing,

changes that were not captured by discrete stage-based approaches. Similarly, PPARα, a key

regulator of lipid metabolism[37], showed significant downregulation in the later stages of the

disease trajectory, highlighting its role in the progression of metabolic dysfunction. Other TFs,

including MYC, JUN, STAT1, and NFKB1/2, exhibited dynamic regulatory patterns, further

illustrating the complexity of gene expression changes during MASLD progression. These findings

underscore the value of trajectory-based analysis in capturing detailed molecular dysregulation

and identifying stage-specific regulatory mechanisms.

An intriguing cluster of TFs, including MEF2A, MEF2C, MEF2D, MYOG, NKX2-5, and ZBTB4,

exhibited unique activity patterns in our trajectory analysis that were not detected using traditional

discrete patient staging. These TFs, known primarily for their roles in myogenesis[38], have also

been implicated in hepatic stellate cell activation and their transition to a myofibroblast-like

phenotype, a critical driver of fibrosis[35]. Their activity peaked during SW5–SW6, corresponding to

heightened stellate cell activation, and then sharply declined at SW7, suggesting either a

resolution of activation or a shift in regulatory dynamics. Notably, no significant changes in their

activity were observed in later stages, indicating a potential stabilisation of their regulatory

influence as the disease progressed.

To explore the underlying molecular processes, we employed a network propagation-based

strategy to extract active network signatures for each sliding window from the MASLD reference

network (Methods). Reactome pathway enrichment analysis of these signatures revealed

Figure 4. Sliding window analysis to study the landscape of molecular changes along the MASLD trajectory. A. Results from TF activity analysis in the discrete patients stratification, and the pseudotemporal trajectory. The colour indicates change in TF activity compared to the previous disease stage. **B.** Mean phenotype and NAS pathologist scores along the SW trajectory.

dynamic, progressive changes along the disease trajectory. Signal transduction and immune
processes became increasingly active as the disease advanced, reflecting the intense
inflammatory and signalling dysregulation characteristic of later stages. Pathways related to
extracellular matrix organisation, strongly associated with fibrosis, exhibited the most significant
upregulation in advanced stages. In contrast, metabolism-related processes were consistently
downregulated throughout the trajectory, highlighting the persistent metabolic dysfunction driving
MASLD progression (Supplementary Fig. 5B).

We compared the dysregulation of molecular processes in each sliding window to traditional
MASLD histopathological scores, such as steatosis, ballooning, inflammation, fibrosis, and NAS
(Fig. 4B). While these scores generally increased with disease progression, some pathways
exhibited early dysregulation before corresponding changes in histopathological scores. For
example, extracellular matrix pathways were upregulated at SW8, preceding the observed
increase in the fibrosis score at SW10, suggesting that molecular markers may provide earlier
insights into MASLD progression than traditional assessments. Overall, the enhanced resolution
of our sliding window-based approach offers a comprehensive molecular landscape of MASLD
progression, shedding light on the interplay between immune activation, fibrotic remodelling, and
metabolic dysregulation over time.

Cell-Type Deconvolution Along the MASLD Trajectory Reveals Network
239

Changes Associated with Tissue Composition

Changes in cell type composition and cellular states within the liver are critical indicators of
MASLD progression, often driven by increased inflammation resulting from hepatic injury caused
by fat accumulation[1]. Understanding the molecular triggers underlying these cellular changes in
the hepatic microenvironment offers a valuable opportunity to identify stage-specific therapeutic
targets for reversing disease progression. Additionally, these insights could facilitate the discovery
of prognostic biomarkers, enabling earlier detection and more personalized MASLD treatment
strategies.

To investigate this, we performed cell type deconvolution on bulk patient data along the disease
trajectory to identify stage-specific shifts in cell composition and their association with the MASLD
disease network (Methods; Fig. 5A). The analysis revealed a decrease in hepatocyte and resident
NK cell proportions, highlighting the progressive infiltration or expansion of other cell types within
the liver. In contrast, increasing proportions of T cells, neutrophils, migratory dendritic cells,

cholangiocytes, and fibroblasts were observed, indicating heightened immune responses,
including inflammation and fibrogenesis. Interestingly, the stability of endothelial cell proportions
beyond the midpoint of the trajectory suggests their role in maintaining blood vessel integrity and
functionality, ensuring consistent liver blood flow and nutrient exchange even in advanced MASLD
stages. Notably, macrophage proportions remained relatively stable throughout the trajectory,
with only minor fluctuations, underscoring their consistent presence and potential involvement in
liver homeostasis and immune regulation across disease stages.

We next investigated the association between cell types and the deregulated processes identified
along the trajectory (Supplementary Fig. 5B). We identified genes whose differential expression
could be solely attributed to changes in cell type proportions along the trajectory, using pseudo-
bulk RNA-seq samples derived from a healthy liver single-cell RNA-seq dataset. These genes
were then analyzed to determine whether the enrichment of Reactome pathways was driven by
shifts in cell type proportions or by the cell type-specific deregulation of biological processes
(Methods). As expected, hepatocytes were primarily associated with deregulated processes
related to lipid metabolism, while other metabolic processes were also linked to cholangiocytes
and endothelial cells (Fig. 5B; Supplementary Table 12). Immune-related processes were
predominantly associated with macrophages, with the exception of 'signalling by interleukins,'
which was primarily linked to neutrophils, T cells, and other immune cell types. These findings
underscore the interplay between specific cell types and biological processes driving MASLD
progression. Deregulated processes related to extracellular matrix (ECM) organisation and
fibrosis were predominantly associated with fibroblasts, highlighting their central role in ECM
remodelling and fibrotic progression. Non-immune-related signalling pathways, such as Wnt
signalling, Akt signalling, and Rho signalling, were primarily linked to endothelial cells, with
contributions from fibroblasts and cholangiocytes in some cases. Interestingly, specific
processes, such as Wnt signalling, were also associated with specific immune cells, suggesting
a potential role in immune modulation or interactions with the stromal microenvironment. These
findings further underscore the complex crosstalk between cellular signalling, immune responses,
and tissue remodelling in MASLD progression.

Figure 5. Results of cell type deconvolution analysis. A. *Cell type deconvolution of patient transcriptomic data along our MASLD trajectory* ***B.*** *Prediction of cell types with which the deregulated processes are associated after excluding gene sets of these processes that could have been identified as differentiated by the mere change in abundance of cell types.*

MASLD Trajectory-Predictive Biomarkers Through Integrated Plasma-Liver Expression Analysis

To address the critical need for non-invasive diagnostic methods for MASLD/MASH staging, we sought to identify genes capable of predicting a patient's MASLD stage through blood proteomics. To establish a baseline for the predictive performance of selected gene sets, we employed a Random Forest regression model using a list of 145 gene signatures derived from the first principal component of our initial analysis (Figs. 2 & 6A). This gene list demonstrated high predictive performance on the UCAM/VCU dataset (Supplementary Fig. 3C), achieving an R^2 of 87.1% and a Mean Squared Error (MSE) of 0.005. Here, R^2 represents the proportion of variance in a patient's trajectory position explained by the model, underscoring the robustness of this gene set in trajectory prediction.

Govaere et al.[39] recently identified 194 genes with correlated expression in both blood plasma and liver across MASLD progression stages. Incorporating these plasma-liver correlated genes into a Random Forest model yielded strong but slightly lower predictive performance compared to the baseline on the UCAM/VCU dataset (R^2: 79.9%, MSE: 0.008). However, performance improved when we retained only the 153 genes with a positive variable importance score in the Random Forest model (R^2: 81.2%, MSE: 0.007; Fig. 6A).

To refine our focus on genes directly relevant to MASLD progression, we narrowed the list to 24 genes that overlapped with our MASLD network. This selection was based on the rationale that these genes are more likely to be directly involved in disease progression and, therefore, better suited for monitoring. The inclusion of our network provided functional context for their plasma protein modulation, enhancing both biological relevance and interpretability. This curated gene set achieved an R^2 of 69.4% and an MSE of 0.01 on the UCAM/VCU dataset, offering a balance between predictive accuracy, simplicity, and interpretability (Fig. 6A; Supplementary Fig. 6A, B).

To validate the ability of our 24-biomarker model to place patients along the trajectory using an independent dataset, we applied it to the EPoS dataset[8] (see first section of our Results & Methods). This dataset included patients for whom plasma protein data was extracted in the Govaere study[39]. Our model successfully recapitulated increasing trends in MASLD stage progression, NAS, and fibrosis scores corresponding to the predicted trajectory (R^2: 81%, MSE: 0.105; Supplementary Figs. 6C–F). Notably, this gene set explained 87.6% of the trajectory variance in this dataset. In contrast, when using random gene signatures of 1,000 and 24 genes,

the model failed to accurately predict patient positions along the trajectory accurately, yielding R^2 values of approximately 11% for both random sets (Supplementary Fig. 6G, H). This highlights the robustness and specificity of our curated 24-gene biomarker set in predicting MASLD progression.

To further validate our model on an entirely independent dataset, we applied it to the Gubra study dataset[21] (described in the first section of Results & Methods; Figs. 6B, C). This dataset effectively distinguished healthy/obese individuals from MASLD/MASH patients, accurately placing them along the trajectory with minimal errors (MSE for predicted vs. actual trajectory positions; Supplementary Fig. 7). Among the tested gene sets, the refined 153-gene set achieved the best performance (R^2: 87.8%, MSE: 0.22), while the 24-gene biomarker set also performed highly (R^2: 87.6%, MSE: 0.214) and explained 81% of trajectory variance. The model accurately classified 25 out of 30 healthy/obese individuals (83.3%) and 26 out of 27 MASLD/MASH patients (96.3%), achieving an overall accuracy of 89.5% (Supplementary Fig. 7). These results highlight the potential of this model for non-invasive MASLD/MASH staging and tracking disease progression.

The genes with the highest importance values in the model's predictions included those involved in extracellular matrix remodelling (COL3A1, MFAP4, THBS2), immune modulation (CXCL8, C7, CCL23, SAA1), metabolic regulation (AKR1B10, HSD17B8, APOF, MTHFS), and cellular stress response (PRDX6, HSPA6). By mapping their "activation" (significant changes compared to the previous trajectory window) and linking them to MASLD network subnetworks (Methods), we charted the timeline of pathway deregulation in blood samples (Fig. 6D). These results were consistent with findings from the liver-based trajectory pathway analysis (Supplementary Fig. 5).

To identify therapeutic opportunities, we hypothesised that targets capable of modulating these predictive biomarkers and present within the MASLD network should be enriched for candidates with a causal association to disease progression. Using the ChEMBL database[39], we searched for approved drugs targeting these biomarkers—either as agonists/activators when the biomarker was downregulated in the disease sliding window or as inhibitors when the biomarker was upregulated. This analysis identified 359 drugs targeting the biomarkers or their associated networks, including 136 drugs that had received regulatory approval (passed Phase 3), regardless of indication (Supplementary Table 13).

Figure 6. Random Forest regression model for the prediction of MASLD patients' position on the inferred disease trajectory. A. Model performance using different gene sets in the primary UCAM/VCU dataset (5-fold cross validation), and validation in two independent datasets used solely for testing. Randomisation was done on 1000 times. **B.** Compared predicted and actual patient positions on the trajectory using an independent dataset (GUBRA), based solely on a 24-gene signature. **C.** Violin plot showing the distribution of the predicted patients' position on the trajectory (GUBRA dataset). **D.** Network-based deregulation analysis of the 24-gene signature to identify potential targets. The color scale represents the -log10(adjusted p-value) of differential expression. Genes are deemed deregulated in a SW if part of the respective deregulated subnetworks. Drugs in max phase 4 were retrieved for targets in the relative deregulated pathways.

Examining the indications annotated in ChEMBL for these drugs, which include all indications

evaluated during clinical trials regardless of approval, revealed that several drugs target

conditions related to MASLD. For instance, fenofibrate and bezafibrate, both PPARα agonists,

are approved for hypertriglyceridemia but failed in MASLD clinical trials. Similarly, belinostat

(targeting HDAC1) and dasatinib (targeting PDGFRB) have shown potential for treating

hepatocellular carcinoma (HCC), although they are not yet approved for this indication. These

examples, along with others in our target list (Supplementary Table 13), underscore the potential

of leveraging our analysis for drug repurposing or the development of targeted single or

combination therapies for MASLD. By focusing on existing drugs with mechanistic alignment to

the disease's molecular drivers, this approach could accelerate therapeutic development and

improve outcomes for MASLD patients.

354 Discussion

MASLD has emerged as the leading cause of chronic liver disease and the second most common

reason for liver transplantation in the Western world[40]. Despite its growing prevalence, the

molecular mechanisms driving MASLD progression remain poorly understood. Traditional staging

methods, which rely on histological scores, face significant limitations, including variability due to

manual annotation and their inability to capture the continuous nature of disease progression. In

this study, we addressed these challenges by adopting a data-driven approach that leverages

pseudo-temporal ordering[20] of MASLD patients, coupled with network analysis, to create an

unbiased map of molecular changes across the MASLD progression continuum[20]. This

methodology offers a more dynamic and granular understanding of disease progression

compared to traditional staging techniques.

In addition to identifying a 145-gene signature predictive of histology scores at the liver biopsy

level, we developed a 24-protein biomarker set derived from our MASLD network. This set can

be measured non-invasively in plasma and used to predict a patient's position along the disease

trajectory. These biomarkers have the potential to enhance existing tools, such as the NAFLD

Fibrosis Score (NFS) and FIB-4, by improving stage precision, enabling multidimensional

assessments, and supporting personalized, stage-specific treatments[41]. Moreover, as these

biomarkers reflect molecular processes central to MASLD progression, they offer valuable

opportunities for monitoring treatment responses and linking molecular mechanisms to clinical

outcomes, paving the way for more effective therapeutic strategies.

By employing sliding windows along the disease trajectory instead of discrete groupings, we
identified both progressive and "switch-like" changes in TF activity and associated molecular
processes.

For example, immune signalling TFs, such as NFKB1 and STATs, exhibited consistent
upregulation at short and regular intervals throughout the trajectory. This trend was mirrored at
the network level, where immune processes were active from the earliest stages and showed
progressive increases in activation over time.

In contrast, profibrotic and myogenesis-related TFs, including MEF2C, MEF2A, MEF2D, ZBTB4,
GATA5, NKX2-5, and MYOG, displayed a distinct activation pattern during MASLD progression.
Their activity peaked between sliding windows 4 and 6, then declined and stabilised by window
7, a dynamic not detected by traditional disease staging. Among these, MEF2 is particularly
noteworthy, as it regulates hepatic stellate cell (HSC) activation and their transformation into a
myofibroblast-like phenotype[42]. HSCs are crucial in the liver's wound-healing response to injury,
but their persistent activation in MASLD—due to impaired apoptosis—drives pathological liver
fibrosis and sustained inflammation, leading to advanced disease stages, including cirrhosis and
potentially hepatocellular carcinoma[42]. Strategies targeting HSC apoptosis have shown
therapeutic promise. For instance, hydronidone has been proposed to improve liver fibrosis
phenotypes by inducing HSC apoptosis, providing a potential pathway for intervention[43].

At the pathway level, our trajectory-based analysis revealed stage-specific modulation of key
biological processes[8] For instance, the unfolded protein response becomes activated toward the
end of the trajectory[44]. In contrast, processes such as the response to hypoxia[45,46], extracellular
matrix organization pathways [47], and downregulation of lipid metabolism[48] emerge from sliding
window 3 onward. Notably, the downregulation of PPARα and HNF1A characterizes the latest
disease stages.

Certain signalling pathways exhibited distinct activity patterns. WNT[49], PDGF[50] and TGF-β
signalling[51] remain consistently activated throughout the trajectory, while Notch signalling shows
increased activity beginning at SW4, with a marked boost in the latter half of the trajectory[52].
Targeting TGF-β signalling has shown potential to reduce disease progression, making it a
promising therapeutic avenue[53–55]. Similarly, Wnt signalling is implicated in MASLD progression[49],
with animal studies demonstrating that inhibiting this pathway can suppress hepatic stellate cell
(HSC) activation, improving disease phenotypes[56]. Collectively, our findings highlight the dynamic

reprogramming of these processes at different time points, impacting multiple levels of cellular
regulation—including proteostasis, the cell cycle, cell signalling, and metabolism. These insights
offer valuable opportunities for stage-specific therapeutic interventions and a deeper
understanding of MASLD progression.

A hallmark of MASLD progression is the gradual shift in liver cell composition, driven by changes
in cellular signalling and plasticity. For example, hepatic stellate cells differentiate into
myofibroblast-like cells, while hepatocytes and cholangiocytes undergo transdifferentiation[57].
Cell-type deconvolution revealed a decrease in hepatocytes, likely due to cell death or
differentiation into cholangiocytes, accompanied by increases in cholangiocytes, fibroblasts, and
immune cells such as T cells, neutrophils, and circulating NK/NKT cells. Metabolic dysregulation
was predominantly linked to hepatocytes, while fibroblasts and cholangiocytes were major
contributors to signalling deregulation and extracellular matrix remodeling. Unsurprisingly,
immune cells—especially macrophages—were implicated in immune processes[58]. Macrophages,
including Kupffer cells, play a key role in regulating lipid metabolism during steatosis and initiating
steatohepatitis through proinflammatory signaling.[59]

Our findings highlight the intricate relationship between hepatic and systemic immunity during
MASLD progression. Future research using single-cell[57] and spatial data will be crucial to uncover
how liver zonation, structural organization, and local niches drive or modulate immune and cellular
processes in MASLD[59].

One limitation of our analysis is that, while it successfully captures molecular events across the
disease trajectory, it cannot fully distinguish drivers of cell deregulation from adaptive or passive
processes. Additionally, our reliance on cross-sectional data introduces uncertainty, as not all
patients in the early stages progress to advanced ones. Despite these limitations, our data-driven
MASLD network provides a valuable foundation for understanding disease mechanisms and
identifying stage-specific therapeutic targets. The network's modules, which are linked to distinct
histological phenotypes, support the development of tailored interventions for specific sub-
phenotypes or their combinations. Incorporating longitudinal data and metadata, such as diabetic
status or prior treatments, could further enhance the predictive power of the network, improve
biomarker discovery, and advance personalized treatment strategies for MASLD.

In summary, this study provides a comprehensive and integrative perspective on MASLD
progression, advancing our understanding of its molecular underpinnings. The data-driven

trajectory framework addresses the limitations of traditional histology-based staging by offering a reproducible and scalable approach to disease annotation. Importantly, this methodology is not restricted to MASLD; by partitioning patient data—or any trajectory data—into informative sliding windows, it can be adapted to explore other progressive diseases, including those modelled at the single-cell level.

The identified gene signature and plasma biomarkers enhance non-invasive diagnostics and enable real-time monitoring of disease dynamics, while the trajectory analysis reveals key molecular processes and identifies novel therapeutic targets. Although longitudinal data will be essential for further refinement, this study lays a strong foundation for future investigations into MASLD and other progressive diseases. Ultimately, this work advances the development of precision medicine, with the goal of improving outcomes for patients affected by this increasingly prevalent condition.

Materials & Methods

Datasets

For the analyses conducted in this study, 2 different publically available human datasets have been used: the "UCAM" dataset[19] (consisting of 58 consecutive patients recruited at the MASH Service at the Cambridge University Hospital) and the VCU dataset[9] (consisting of 4 obese bariatric controls without MASLD and 74 patients covering the whole MASLD spectrum). The raw data and relevant metadata have been downloaded from Array's Express database (E-MTAB-9815) and NCBI's GEO repository (GSE130970), respectively. Details for both datasets are given in Supplementary Table 1. "Sample_5" was removed from the analysis as it was considered an outlier (Supplementary Fig. 8A-B).

For the validation of the human trajectory inference method, we used two publicly available datasets (GUBRA[21,60] and EPoS[8]). The GUBRA dataset consists of n=26 healthy (n=14 normal-weight, n=12 overweight) individuals recruited at the Center for Diabetes Research (Copenhagen University, Denmark) and n=31 MASLD/MASH patients recruited at the Department of Hepatology and Gastroenterology (Aarhus University Hospital, Denmark). The EPoS dataset consists of 168 MASLD/MASH patients, originates from a large cohort recruited in multiple European parts, and has been previously used by our team[61]. In additionally to the expression

465   data, the EPoS dataset also provided information on the MASLD disease stage, NAS, and
466   Fibrosis scores per patient.

RNA-Seq Analysis and data processing

For the RNA-Seq analysis, the *FASTQC* software was used for generating quality-control (QC)
reports of individual FASTQ files (http://www.bioinformatics.babraham.ac.uk/projects/fastqc, v.
0.11.9) and reads were aligned to the human GRCh38 reference genome using hisat2[62] (v 2.1.0).
Then, the genes were counted with *HTSeq*[63] (v 0.11.1), and the *biomaRt package*[64] was used to
map "Ensembl gene IDs" to "HGNC symbols".

We used quantile normalisation to normalise the 2 datasets and Bioconductor's function
COMBAT[65] from R package *sva* to remove the batch effect observed upon integrating them
(Supplementary Fig. 8A-B). Quality control checks validate the robustness of the analysis,
showing that the expression of the genes in each MASLD stage separately for each dataset
remains similar after batch effect correction (Supplementary Fig. 8C-D), with a high correlation of
the first principal component of the PCA plot with the main MASLD characteristics (Supplementary
Fig. 8E), and a similar expression of genes previously linked to MASLD before and after the
correction (Supplementary Fig. 8F).

Pseudo-temporal ordering of patients

The merged dataset was not expected to have multiple disconnected trajectories with a particular
cyclic or (necessarily) complex topology. Thus, we selected *Slingshot*[20,66] as the most suitable
tool to infer the disease trajectory. We constructed an MST (Minimum Spanning Tree) on ordered
sets of clusters to capture the disease trajectory based on the points on the PCA plot. The inferred
horizontal trajectory was in agreement with the high correlation scores between PC1 and the
MASLD characteristics. Thus, this dimension could be considered as a pseudo-temporal
trajectory of MASLD, in which small values represent healthy or initial stages of disease, while
large values belong to severe fibrosis conditions. Based on this hypothesis, each point on the
PC1 axis could be considered as a pseudo-time. For the independent dataset, this process was
repeated with the complete transcriptome and using only the 145 gene signatures identified using
the random forest classification (see section below).

Patient stratification for annotation of trajectory

Based on the histological scores defined by the CRN scoring system[67] and following the older definition of NAFLD/NASH, our dataset (136 patients) has been divided into 4 bariatric obese controls and 132 MASLD, which has further been sub-clustered based on the fibrosis stage into mild (MASH F0), moderate (MASH F1-2) and severe (MASH F3-4) groups. This stratification aims to increase the interpretability of the analysis, which has been described in detail in our previous studies[19,61] (Supplementary Table 1).

Extraction of 145 gene signatures using Random Forest classification

Calling the parameter *rotation* from *prcomp* function in R (default stats package), we produced a list of ranked genes with the highest absolute values that explain the variance along with the disease progression on the horizontal axis. The top 200 genes were selected as the training set of a random forest classification model (R package *randomForest*) to predict the disease stage (MASL, MASHF01, MASHF2, MASHF34) and the different MASLD characteristics (steatosis, inflammation, ballooning, fibrosis). The accuracy was estimated using 5-fold cross-validation 1000 times. From this first run of the model, only genes with positive "Mean Decrease Accuracy" were selected, representing the set of top genes that contributed to the disease stage prediction. Those genes were used as the training set of a new random forest classification model.

Pathway enrichment analysis

The Reactome Pathway Knowledgebase[24] (accessed May 2024) was used in different steps of this study to perform pathway enrichment analysis. Fisher's exact test[68] was used for the statistical significance test, and the Benjamini–Hochberg method[69] was applied for the multiple hypothesis testing correction.

Identification of gene expression modules associated with MASLD histological phenotypes

We applied the weighted gene co-expression network analysis (WGCNA)[70], as in Barker et al.[22], through the R package *WGCNA*[23] (v 1.72-1) in combination with an optimisation task to identify the optimal set of gene co-expression modules. Initially, we used the soft-thresholding approach to construct a signed weighted gene co-expression network, and the power (soft thresholding

parameter) was set to 4, as it was the minimum value for which the weighted network

approximated scale-free topology (fit index equal to 0.9). Then, we modified the derived adjacency

weights into a topological overlap matrix (TOM) to finally get the dissimilarity matrix of genes (1-

TOM). We used the *cutreeDynamic* function of the *dynamicTreeCut* R package (v 1.63-1) to

analyse the dissimilarity matrix and extract the set of co-expression modules. Instead of using the

default parameters, we performed an optimization process to find the optimal set of "deepSplit"

and "minClusterSize" parameters. The criterion of this optimization task was the accuracy of the

derived modules to model the four MASLD phenotypes (steatosis, ballooning, fibrosis and the

NAS score) using linear regression. To that end, we performed cross-validation (k=4), separating

the patients into four groups with the *crossv_kfold* function of the *modelr* R package (v 0.1.11). At

each round, we modelled the phenotypes as linear functions of the derived eigen-genes of

modules (*lm* and *step* functions of the *stats* R package) using the training dataset. In order to

capture mechanisms underlying the progression of MASLD rather than downstream effects, we

used the inflammation score as a covariate in this model. Each linear model had the following

format:

$MASLD\ phenotype \sim intercept + eigengenes\ of\ modules + inflammation + age$

We applied a backward stepwise search and the Akaike Information Criterion (AIC)[71] to optimise

the coefficients of eigen-genes in each linear model, while modules with statistically non-

significant coefficients were filtered out (adjusted p-value<0.05, Benjamini-Hochberg

correction[69]). The test dataset was used to estimate the accuracy of the model, calculating the

mean squared error (MSE). In order to find the optimal parameter values, we calculated the

average test MSEs for the trained models of the four MASLD variables for each parameterisation

set. Following this approach, we found that the test error was minimum for ["deepSplit"=4,

"minClusterSize"=180]. We created the final linear models using this set of parameters and

keeping only the statistically significant modules for each phenotype. For the downstream

analysis, we filtered the gene content of each significant-identified module to select those genes

with Pearson's correlation equal to or greater than 0.4 with the respective eigen-gene.

Transcription factor activities analysis

We used *VIPER*[72] (Virtual Inference of Protein-activity by Enriched Regulon analysis; v 1.37.0)

and the CollecTRI database[73] (accessed May 2024) to identify the upstream transcriptional

regulators that can explain the observed gene expression changes. The calculated Normalised
Enrichment Score (NES) was used to determine the direction of each regulation ("activation" for
TFs with NES > 0 and "inhibition" for those with NES < 0). We performed the hypothesis testing
using the student's t-test, and then, the Benjamini-Hochberg method[69] was used to correct the p-
values, ensuring decreased Type I errors.

Generation of global MASLD network

This task aimed to create a global interaction network for MASLD, including information about the
molecular events that take place in different stages of the disease, either in signalling cascades
(upstream component) or the transcriptional regulation (downstream component). For that
reason, we exploited the results of WGCNA and differential expression analysis in SWs,
generating upstream and downstream subnetworks for each MASLD phenotype, as well as for
the significantly deregulated TFs. We used the *Prize-collecting Steiner Forest (PCSF)* algorithm,
implemented through the R package *PCSF*[74] (v 0.99.1), to create these subnetworks. Finally, we
connected the pairs of upstream and downstream subnetworks and unified all of them into the
global MASLD network.

PCSF returns as output a subset of a given reference network relevant to an input-weighted list
of prize-carrying nodes. To define such a list for each MASLD phenotype, we applied the following
workflow: We performed TF enrichment analysis on the filtered gene content of each significant-
identified WGCNA module, using the CollecTRI database[73] and other TF-regulon databases,
retrieved from the repository of *EnrichR*[75] (ENCODE and ChEA Consensus TFs from ChIP-X[76],
DNA binding preferences from JASPAR[77,78], TF protein-protein interactions). Then, we performed
Reactome[79] pathway enrichment analysis on each module-specific set of enriched TFs, and we
selected only the most semantically specific pathways, which are also related to signal
transduction. Consequently, each *WGCNA* module was linked with a group of TFs and a set of
Reactome pathways. The filtered gene sets of WGCNA modules (downstream nodes), the
enriched TFs, and the gene content of the enriched Reactome pathways (upstream nodes)
constituted the prize-carrying nodes for each MASLD variable.

We defined a similar set of prize-carrying nodes related to the results of differential expression
analysis. To that end, we first selected the TFs that have been identified as deregulated in at least
one SW (adjusted p-value < 0.05). Then, we identified the most specific signalling pathways
related to this set of TFs, similar to the previous task. Additionally, we found the intersection of

the selected TFs regulons (using the CollecTRI annotation) with the gene content of WGCNA

modules. This set of TF regulons (downstream nodes), the group of deregulated TFs and the

gene content of the enriched Reactome pathways (upstream nodes) constituted the prize-carrying

nodes for the deregulated network. We performed all the aforementioned enrichment tests using

the Fisher's exact test and Benjamini-Hochberg correction (adjusted p-value < 0.05), adjusting

the background annotation of the databases to include only genes that were present in the

transcriptomic dataset.

We constructed the reference interaction network, integrating various interaction databases with

the *union* function of the *igraph* R package (v 2.03)[80], including both cell signalling and metabolic

pathways (Reactome[79], KEGG[81,82], HumanCyc[83], ReconX[84], provided by Pathway Commons[85],

and OmniPath[86], accessed using the *OmniPathR* library in Bioconductor[87,88]). Regarding the

metabolic networks of HumanCyc and ReconX, we removed all the ChEBI entries with zero out-

coming edges, as they are ending points on the network. Additionally, we filtered the OmniPath

network to avoid weakly verified interactions, keeping only those with a number of resources and

curation efforts more than the 0.5 and 0.75 quantiles of the respective distributions. We split the

complexes to form fully connected groups of protein nodes, and we deleted all the interactions

annotated as inhibitions, as their molecular effect cannot be modelled using the type of network

analysis that we used (network propagation). The final reference network included 12,549 nodes

and 205,706 edges.

In order to run *PCSF*, we define the cost of the edges in the reference network equal to 0.001 and

the prizes of seed nodes equal to 100. For each of the five variables, we created an upstream

subnetwork (combining the upstream nodes and TFs) and a downstream subnetwork (combining

the downstream nodes and TFs), and then we connected them to form a unique network. This

gradual generation of the MASLD network was selected in order to keep information about the

origin of each node and edge. We used a variant of the *PCSF* algorithm (*PCSF_rand*) to generate

the subnetworks. This function executes the primary process several times (10 iterations were

selected here), adding random noise to the edge costs and merges all the solutions to create a

unique one. Finally, we unified the five networks into a global one using the *union* function of the

igraph R package (v 2.0.3).

We filtered the derived MASLD network so as to keep only functionally annotated genes and

weight the edges according to the semantic similarity of interactors. We used the *GOSemSim* R

package[89] (v 2.26.0) to calculate the semantic similarity of all pairs, according to their annotation

in Gene Ontology Biological Process (GO-BP)[90], based on the measures of Resnik[91], Lin[92] and

Wang[93]. We used different measures to score the similarities, aiming to mitigate each one's

potential drawbacks and biases. We averaged the similarity values to define the weight of each

edge. The lack of annotation of some genes produced edges with undefined weights. These

genes and the respective edges were removed from the network. Finally, we applied the

Laplacian normalisation in order to reduce the topological bias of hub nodes[94]. The final MASLD

network included 4,247 nodes and 38,847 edges.

To evaluate the relevance of the derived network to MASLD, we calculated its enrichment in two

gene signatures. The first signature (164 genes) was created based on recent literature[95–100],

including genes which have been associated with different stages of MASLD progression. The

second signature (155 genes) was retrieved from the MSigDB repository[101] and corresponds to

the canonical MASLD pathway of WikiPathways[102]. The overlap between these two signatures

was 15 genes. Fisher's exact test was applied to estimate the statistical significance of the

enrichment scores.

Patients stratification into sliding windows

We divided the ordered samples on the trajectory into sliding windows (SWs), namely as a

sequence of overlapping groups [SW_1, SW_2,..., SW_X], to investigate how the MASLD molecular

profile changes on the pseudo-temporal space. In order to define the SW-based stratification of

samples, we developed a graph-based method, which examines all the possible sequences of

SWs for a given set of SW sizes $S=[S_1,…, S_m]$ and an overlapping percentage $α$ between two

adjacent SWs and finds the optimal one. The concept of this novel method is based on the

representation of disease stages as nodes connected on a directed path, where each node is

annotated with a group of samples aligned to their pseudo-temporal order. A partial overlap of

samples between each pair of connected nodes ensures the continuity of the disease trajectory.

The optimisation process aims to detect a path of nodes that can sufficiently describe the

molecular events that occur along the disease progression through the sequential comparison of

the groups of samples. Generally, the method is implemented using a three-step workflow: the

creation of all the potential directed paths of SWs, their evaluation and filtering, and finally, the

selection of the optimal one.

The method starts with the mapping of samples to a set of indices $I=[1, 2,..., N]$, based on their

ranking on PC1, where N is the size of their set. Then, an iterative process is performed to

construct a directed acyclic graph (DAG), in which nodes correspond to groups of indices, and

directed edges connect two overlapping groups. Initially, an empty node is created as the starting

point of all paths. In the first iteration, a new node is created and annotated with the first S_i indices

for each size S_i of the given set S. As a result, M nodes are created and aligned with indices from

1 to S_i, constituting candidate versions of SW_1. Directed edges from the starting node to SW_1

nodes are created. In the next iteration, the SW_1 nodes are used as starting points to create

candidate nodes for SW_2. For an SW_1 with indices set K, the starting index of its subsequent SW_2

nodes is defined as *max(K)+1-ceiling(size(K)*α)*, which signifies that approximately the last *100*α*

percent of the SW_1 indices will also be included in the SW_2 nodes. New directed edges connect

each SW_1 with its subsequent SW_2 nodes. This iterative growth of the graph terminates when all

the samples have been mapped to SW nodes. Many nodes and edges could be generated

multiple times through the expansion of different paths. Thus, filtering is performed at the end of

each iteration to remove this redundancy. Finally, an empty node is created and connected with

the final SW nodes. All the directed paths that connect the starting and ending nodes constitute

potential representations of the pseudo-temporal sequence of disease stages (Supplementary

Fig. 9A).

Differential expression analysis with *DESeq2*[103] is used to examine each pair of connected SW

nodes as part of the optimal path. In the beginning, a pair is assumed as appropriate if the derived

number of differentially expressed genes (DEGs with adjusted p-value < 0.1) is in the range

R=[100, 1000]. These values were selected empirically, but the bounding of DEG size is based

on the hypothesis that smaller values indicate very weak differentiation, whereas higher values

might imply a strong differentiation signal, which can be explained in more than two SWs. Similarly

to the construction of the DAG, this step is performed in an iterative process, examining the graph

edges in batches based on their minimum distance from the starting node. To reduce the

execution time, a pruning step is used at the end of each iteration to remove edges and the

corresponding paths that do not satisfy the requirements of DEGs size. While this filtering reduces

the amount of *DESeq2* runs that need to be executed in the following iterations, it does not ensure

the connectivity of the graph, which is required in order to find a solution. Therefore, the limits of

range R are adjusted at each round so that the lower limit is not higher than the 80th percentile

and the upper limit is not lower than the 20th percentile of DEGs size distribution.

The second step typically produces numerous candidate paths, due to the relaxation of DEGs

size thresholds. Additional filtering is performed to reduce their amount and select the optimal

path. Firstly, the incoming edges of a single SW node are clustered based on the initial index of

the respective adjacent nodes. For each cluster, only the edge that corresponds to the highest

number of DEGs or (in case of equal numbers) the one that originates from the node with the

most extensive set of indices remains in the graph. Secondly, the edges which produce DEGs

less than the lower bound of the R range are selected to calculate their median number of DEGs.

This value is used to remove all the edges with the lower number of DEGs, ensuring that the

connectivity of the graph is not interrupted (Supplementary Fig. 9B). Finally, the product of the

following three max-scaled scores is used to prioritise the remaining candidate paths: 1. the

median number of DEGs, 2. the number of edges with DEGs more than the lower bound of R and

3. the coefficient of variation of the number of DEGs (Supplementary Fig. 9C). The described

method was implemented for the MASLD dataset (135 samples), using the following pairs of SW

sizes and overlap scores: ([6,15], 0.2), ([8,17], 0.25), ([8,24], 0.3), ([10,20], 0.3), ([12,25], 0.35).

Therefore, five graphs were constructed and filtered, and the candidate paths from all of them

were prioritized to find the optimal one.

Statistical analysis

Differential gene expression analysis was done using *DESeq2*[103] (v 1.26.0), and the Benjamini-

Hochberg method was applied to adjust the raw p-values to control the False Discovery Rate

(FDR). In the analyses that used the discrete stratification of patients' disease stages,

comparisons were made against the immediately previous stage. For all comparisons related to

the trajectory analysis (see section "Patients stratification into sliding windows"), sliding windows

(SWs) were compared to the previous SW (SW_x vs SW_{x-1}).

Disease stage-specific network analysis

We applied a network propagation-based approach to detect the deregulated components of the

MASLD network for each SW using the *igraph* R package (v 2.0.3). We used the personalised

PageRank algorithm[104] to run network propagation. This method accepts as input a vector of

weights for a set of nodes. We defined these weights based on the results of differential

expression analysis (for all genes except transcription factors) and TF activity analysis (for the

transcription factors). We were interested in finding a network signature for up- and down-

regulated conditions separately, so we created two weight vectors for each SW. In particular, we

defined the weights of node i (W_+ and W_-) using the direction of its differentiation (based on the

logarithmic fold change or NES score) and the log-transformed adjusted p-value of the respective

statistical test, to be bounded by a maximum value of 10:

● $If\ sign(deregulation) = +1: Weights_i : \{W_+ = min[-log_{10}(adj.pvalue) + 1, 10],\ W_- = 1\}$

● $If\ sign(deregulation) = -1: Weights_i : \{W_+ = 1,\ W_- = min[-log_{10}(adj.pvalue) + 1, 10]\}$

Network propagation was performed for each SW and direction of deregulation. To assess the

statistical significance of the derived PageRank scores, we run the propagation task 1000 times

using random weight distributions. Then, we calculated an empirical p-value for each node based

on the ratio of its random scores that are lower than the real one:

$$pvalue = 1 - \frac{|random\ scores\ <\ PageRank\ score|}{1000}$$

728
729

We implemented this approach for three values of damping factor: [0.5, 0.7, 0.85], and we

selected the gene nodes with p-value<0.05 in all three runs, as significantly propagated. Then,

we extracted the respective subnetwork signature as a subgraph of the MASLD network, and we

removed the significantly propagated nodes which belonged to small components (less than 4

members) because they have been identified due to isolated nodes with high input weight, far

from the main affected component of the network. Finally, the activation score of a gene in a SW

was defined as equal to its initial weight used for the network propagation if the gene had been

included in the derived SW-specific deregulated component. Otherwise, the activation score was

defined as equal to zero.

Definition of disease-relevant pathways

We used pathway enrichment analysis to link the deregulated subnetworks in each SW with

molecular pathways using the Reactome database (as described in the section "Pathway

enrichment analysis"). In order to create a unified and semantically non-redundant list of MASLD-

relevant pathways and describe their activation in the pseudo-temporal trajectory, we clustered

the enriched pathways from the different SWs. The clustering was performed based on the

hierarchical structure of the Reactome database and the sets of deregulated genes that were

associated with the enriched pathways. Initially, the Reactome graph was constructed using the

igraph Python package (v 0.4.11), connecting the ontological terms with semantic parent-child

relations. Then, we performed enrichment analysis to associate the genes of the global MASLD

network with an extended set of pathways (adjusted p-value < 0.01), and we extracted the

respective Reactome subgraph. This MASLD network-related subgraph was used as a reference

to filter the enriched pathways (adjusted p-value < 0.05) from the analysis of SW subnetwork

signatures, removing those which are not included in it. Then, we performed an iterative

agglomerative clustering on the remaining terms.

All terms with a distance greater than 4 were clustered to their parental terms because their

semantic meaning was primarily precise. We applied the following clustering steps to minimize

the redundancy of the gene sets of parent and child terms. Terms with distance 4, whose parent

had multiple child terms, were clustered to their parent. Terms with distance 2 or 3, whose parent

had only one child term, were clustered to their parent if the respective parent-child genomic

annotations were not identical (Jaccard similarity < 0.8). Otherwise, the child term was used to

substitute for its parent position on the graph. Finally, terms with distance 3, whose parent had

multiple child terms, were clustered to their parent, along with their sibling terms, if the respective

average genomic (Jaccard) similarity was more significant than 0.3, meaning that these terms

correspond to relatively similar gene sets. Otherwise, the clustering was not performed. We

selected the leaf nodes of the remaining graph to represent the disease-related pathways.

However, nodes with relatively low number of associated genes (a number less than the median

value minus one mean absolute deviation) were filtered out, in order to exclude extremely specific

pathways from the final list. For the genes that had been detected as deregulated, but they were

not linked to any of the final disease-related pathways, we defined additional terms. We used their

reference annotation to concatenate them into the generic mechanisms of Reactome, defining

the respective terms (e.g. "Other genes related to Metabolism"). Using this approach, we selected

82 MASLD-relevant pathways from the initial MASLD reference subgraph of 465 terms. We

calculated the activation of a pathway in a specific SW as the summation of the respective

activation scores of its genes, weighted by their pagerank centralities on the underlying pathway-

based subnetwork (Supplementary Fig. 5).

Cell-type deconvolution analysis

For the deconvolution analysis, we selected a 10X single-cell RNA-seq dataset from a healthy

human liver[105] as a reference. We re-analysed the single-cell dataset based on standard single-

cell analysis protocols[106]. We filtered out genes expressed in less than 3 cells and cells with less

than 200 genes expressed in them. Additionally, we filtered out cells with more than 7000 features

(putative doublets) and with more than 5% mitochondrial genes. To explore the suitability of the

dataset further, we proceeded with standard normalisation and vs (variance stabilising

transformation) scaling, selection of highly variable genes (HGV), PCA and UMAP analysis. For

the deconvolution analysis, we applied the CATD pipeline[107], using the raw counts as input. CATD

performs normalisation and additional preprocessing internally. After setting the preprocessing

parameters, we run the deconvolution analysis, selecting a subset of methods (EpiDISH[108],

bseqsc[109], DWLS[110], CIBERSORT[111], FARDEEP[112] and BayesPrism[113]). The consensus result

was calculated as the mean of cell-type proportions per cell-type across methods.

Identification of cell-type specific deregulated pathways

We assumed that the differentiated network signatures had been identified either due to changes

in cell type proportions or the deregulation of specific processes. In order to deconvolve the

differentiation signal and discern the deregulated processes that are related to specific cell types,

we applied the following workflow: Initially, we used the stratification of patients in SWs and the

calculated cell type proportions (see section "Cell type deconvolution analysis") to model

Gaussian kernel-based density functions for the cell type proportions in each SW. We used these

kernels to generate a number of pseudo-bulk RNA-seq samples for each SW, with predefined

library sizes, sampling cells from the reference single-cell RNA-seq dataset of healthy liver. The

number of SW pseudo-samples and the size of count libraries were selected randomly from the

ranges [6,20] and [20e6, 50e6], respectively. Differential expression, TF activity, and network-

based analyses were performed for the pseudo-dataset, similar to the real one, assuming that the

results would represent the differentiation of the MASLD network, which was caused only by

changes in cell type proportions. We generated 30 random pseudo-datasets to average the

results from differential expression and TF activity analyses. The network analysis extracted up-

and down-regulated network components, which were compared with those of the real dataset. If

a gene was included in the real- and pseudo-signature of a specific SW and its activation score

in the pseudo-signature exceeded the 90% of that in the real-signature, it was considered to be

differentiated due to changes in cell composition within that SW. We removed these genes from

the MASLD-relevant pathways (see section "Definition of disease-relevant pathways"), assuming

that the differentiation signal of the remaining gene sets is associated with the deregulation of the

respective pathways.

In order to examine if the deregulation of these pathways originates from specific cell types, we
first create extended lists of cell type markers. We used the reference single dataset and the
Seurat R package (v. 5.0.3)[114] to identify the markers. We filtered the derived lists, excluding
genes which were not included in the global MASLD network. Then, we used the PCSF algorithm
(using the same parameters as in the section "Generation of global MASLD network") to amplify
each set of marker genes, integrating other genes from the MASLD network that are topologically
close to them. Finally, we assessed the relevance of the deregulated pathways to the liver cell
types using enrichment tests (Fisher's exact test for hypothesis testing and the Benjamini–
Hochberg method for the multiple hypothesis testing correction).

Prediction of patients' position on pseudo-temporal trajectory

For the prediction of the patients' position on the trajectory, we created a Random Forest
regression model. In each iteration, the dataset was split into training (80%) and testing (20%)
sets and run 1000 times to ensure optimal performance. The model's performance was evaluated
for 1000 iterations by calculating the mean MSE (Mean Squared Error) and mean R^2 (coefficient
of determination), explaining the percentage of trajectory variance explained by the selected gene
set. Additionally, the assessment of variable importance reduced the gene set, identifying the
most predictive genes and excluding those with negative variable importance scores.

We tested multiple gene sets in our model to determine the top-performing predictors. First, we
tested the 145 gene signature derived from our random forest classifier and a set of 194 integrated
liver-plasma targets identified by Govaere et al.[39]. Next, the 194 liver-plasma targets were refined
further according to their positive contribution (negative variable-importance-score genes were
removed), assessing whether restricting the gene set improves the overall model performance.
Finally, we evaluated the intersection of this refined gene set with the differentiated part of the
MASLD network derived from our analysis (24 genes).

A crucial part of this analysis was the transformation of the expression data into z-scores based
only on the gene subset being analysed in each run. For instance, when using the 194 gene set,
the z-score transformation was applied only for this set of genes. This approach ensures
predictions even for incomplete proteomic data (measurements of specific plasma proteins),
allowing potential real-time prediction of the patients' position on our trajectory based only on the
most informative plasma proteins.

The same gene sets were also tested on two independent datasets (Gubra and EPoS, described
above) to predict their patients' position on our trajectory. This time, the model training was based
on our internal UCAM/VCU dataset, with the external datasets used solely for testing and
validating our model's generalisability. Furthermore, random gene sets were used as controls,
run 1000 times for each gene set and averaging the predictive metrics, highlighting the predictive
value of our selected gene signature.

Identification of drug targets related to liver-plasma biomarkers

We used the intersection of liver-plasma biomarkers with the deregulated genes (24 genes) in the
MASLD network to identify potential drug targets for the different stages and molecular events of
the disease. Each biomarker had been identified as up- or down-regulated in specific SWs based
on our network analysis. We searched for the associated deregulated pathways in these SWs
and retrieved the gene sets of these pathways, extending the list of biomarkers with functionally
similar genes, which were also deregulated in the same disease stages. Then, we searched in
the ChEMBL database[115] for approved or clinical candidate drugs (max_phase : [2,3,4]) that target
these genes. Specifically, we search for drugs whose action potentially inverses the deregulation
of the respective targets (agonists/activators for the down-regulated and inhibitors/antagonists for
the up-regulated genes). Finally, we created a mapping of drugs and drug targets for each
molecular event related to the 24 biomarkers (Supplementary Table 13).

Data and code availability

861 All data used in this study is publicly available as described in the methods. The code used for
862 the analyses can be found here: https://github.com/kamzolas/MASLD---Continuous-trajectory-
863 approach

Acknowledgements

The work was supported by the European Molecular Biology Laboratory (IK, TK, CGB, AVP, IP,
HW, EP) and Open targets (TK, IP, EP); The EMBL GeneCore Facility is acknowledged for
support in transcriptomics data collection. EMBL IT Support is acknowledged for the provision of
computer and data storage servers. Funding from the Medical Research Council (MRC)
supported I. Kamzolas. MRC MDU, Caixa Foundation and Spanish Government ATRAE
programme support AVP research on MASLD.

Author contributions

IK, TK: Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Data curation, Writing - Original Draft, Writing - Review & Editing, Visualization. CGB, AVP: Formal analysis, Investigation, HW: Data Curation, MV: Writing - Review & Editing, Supervision, IP: Supervision, Funding acquisition, AVP: Conceptualization, Supervision, Resources, Funding acquisition, Writing - Review & Editing EP: Conceptualization, Methodology, Resources, Supervision, Project administration, Visualization, Funding acquisition, Writing - Original Draft, Writing - Review & Editing.

References

1. Chan, W. K. *et al.* Metabolic Dysfunction-Associated Steatotic Liver Disease (MASLD): A State-of-the-Art Review. *Journal of obesity & metabolic syndrome* **32**, (2023).

2. Loomba, R. & Sanyal, A. J. The global NAFLD epidemic. *Nat. Rev. Gastroenterol. Hepatol.* **10**, (2013).

3. Estes, C. *et al.* Modeling NAFLD disease burden in China, France, Germany, Italy, Japan, Spain, United Kingdom, and United States for the period 2016-2030. *J. Hepatol.* **69**, (2018).

4. Wong, V. W.-S., Adams, L. A., de Lédinghen, V., Wong, G. L.-H. & Sookoian, S. Noninvasive biomarkers in NAFLD and NASH - current progress and future promise. *Nat. Rev. Gastroenterol. Hepatol.* **15**, 461–478 (2018).

5. Bedossa, P. Utility and appropriateness of the fatty liver inhibition of progression (FLIP) algorithm and steatosis, activity, and fibrosis (SAF) score in the evaluation of biopsies of nonalcoholic fatty liver disease. *Hepatology (Baltimore, Md.)* **60**, (2014).

6. McGill, D. B., Rakela, J., Zinsmeister, A. R. & Ott, B. J. A 21-year experience with major hemorrhage after percutaneous liver biopsy. *Gastroenterology* **99**, (1990).

7. Ratziu, V. *et al.* Sampling variability of liver biopsy in nonalcoholic fatty liver disease. *Gastroenterology* **128**, (2005).

896 8. Govaere, O. *et al.* Transcriptomic profiling across the nonalcoholic fatty liver disease

897  spectrum reveals gene signatures for steatohepatitis and fibrosis. *Sci. Transl. Med.* **12**,

898  (2020).

899 9. Hoang, S. A. *et al.* Gene Expression Predicts Histological Severity and Reveals Distinct

900  Molecular Profiles of Nonalcoholic Fatty Liver Disease. *Sci. Rep.* **9**, 12541 (2019).

901 10. Atabaki-Pasdar, N. *et al.* Predicting and elucidating the etiology of fatty liver disease: A

902  machine learning modeling and validation study in the IMI DIRECT cohorts. *PLoS Med.* **17**,

903  e1003149 (2020).

904 11. Anstee, Q. M., Seth, D. & Day, C. P. Genetic Factors That Affect Risk of Alcoholic and

905  Nonalcoholic Fatty Liver Disease. *Gastroenterology* **150**, (2016).

906 12. Eslam, M. & George, J. Genetic contributions to NAFLD: leveraging shared genetics to

907  uncover systems biology. *Nat. Rev. Gastroenterol. Hepatol.* **17**, 40–52 (2019).

908 13. Chen, Y. *et al.* Genome-wide association meta-analysis identifies 17 loci associated with

909  nonalcoholic fatty liver disease. *Nat. Genet.* **55**, 1640–1650 (2023).

910 14. Fairfield, C. J. *et al.* Genome-Wide Association Study of NAFLD Using Electronic Health

911  Records. *Hepatology Communications* **6**, 297 (2022).

912 15. Genome-wide association study of non-alcoholic fatty liver and steatohepatitis in a

913  histologically characterised cohort☆. *J. Hepatol.* **73**, 505–515 (2020).

914 16. Magwene, P. M., Lizardi, P. & Kim, J. Reconstructing the temporal ordering of biological

915  samples using microarray data. *Bioinformatics* **19**, 842–850 (2003).

916 17. Mukherjee, S. *et al.* Molecular estimation of neurodegeneration pseudotime in older brains.

917  *Nat. Commun.* **11**, 5781 (2020).

918 18. Young, A. L. *et al.* Data-driven modelling of neurodegenerative disease progression:

919  thinking outside the black box. *Nat. Rev. Neurosci.* **25**, 111–130 (2024).

920 19. Azzu, V. *et al.* Suppression of insulin-induced gene 1 (INSIG1) function promotes hepatic

lipid remodelling and restrains NASH progression. *Molecular metabolism* **48**, (2021).

20. Street, K. *et al.* Slingshot: cell lineage and pseudotime inference for single-cell

transcriptomics. *BMC Genomics* **19**, 1–16 (2018).

21. Suppli, M. P. *et al.* Hepatic transcriptome signatures in patients with varying degrees of

nonalcoholic fatty liver disease compared with healthy normal-weight individuals. *Am. J.*

Physiol. Gastrointest. Liver Physiol. **316**, (2019).

22. Barker, C. G. *et al.* Identification of phenotype-specific networks from paired gene

expression–cell shape imaging data. *Genome Res.* **32**, 750 (2022).

23. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network

analysis. *BMC Bioinformatics* **9**, 1–13 (2008).

24. Milacic, M. *et al.* The Reactome Pathway Knowledgebase 2024. *Nucleic Acids Res.* **52**,

D672–D678 (2023).

25. Córdova, G. *et al.* SMAD3 and SP1/SP3 Transcription Factors Collaborate to Regulate

Connective Tissue Growth Factor Gene Expression in Myoblasts in Response to

Transforming Growth Factor β. *J. Cell. Biochem.* **116**, 1880–1887 (2015).

26. Younesi, F. S., Miller, A. E., Barker, T. H., Rossi, F. M. V. & Hinz, B. Fibroblast and

myofibroblast activation in normal tissue repair and fibrosis. *Nat. Rev. Mol. Cell Biol.* **25**,

617–638 (2024).

27. Liu, D. *et al.* Ets-1 deficiency alleviates nonalcoholic steatohepatitis via weakening TGF-β1

signaling-mediated hepatocyte apoptosis. *Cell Death Dis.* **10**, 1–14 (2019).

28. Varga, T., Czimmerer, Z. & Nagy, L. PPARs are a unique set of fatty acid regulated

transcription factors controlling both lipid metabolism and inflammation. *Biochim. Biophys.*

Acta **1812**, 1007 (2011).

29. Vachliotis, I. D. & Polyzos, S. A. The Role of Tumor Necrosis Factor-Alpha in the

Pathogenesis and Treatment of Nonalcoholic Fatty Liver Disease. *Curr. Obes. Rep.* **12**,

191–206 (2023).

30. Tumor necrosis factor-α signaling in nonalcoholic steatohepatitis and targeted therapies. *J. Genet. Genomics* **49**, 269–278 (2022).

31. CREB family: A significant role in liver fibrosis. *Biochimie* **163**, 94–100 (2019).

32. Luedde, T. & Schwabe, R. F. NF-κB in the liver—linking injury, fibrosis and hepatocellular carcinoma. *Nat. Rev. Gastroenterol. Hepatol.* **8**, 108 (2011).

33. Luo, M., Li, T. & Sang, H. The role of hypoxia-inducible factor 1α in hepatic lipid metabolism. *J. Mol. Med.* **101**, 487–500 (2023).

34. Mesarwi, O. A. *et al.* Hepatocyte Hypoxia Inducible Factor-1 Mediates the Development of Liver Fibrosis in a Mouse Model of Nonalcoholic Fatty Liver Disease. *PLoS One* **11**, e0168572 (2016).

35. Mann, D. A. & Smart, D. E. Transcriptional regulation of hepatic stellate cell activation. *Gut* **50**, 891 (2002).

36. Tan, J. *et al.* HNF1α Controls Liver Lipid Metabolism and Insulin Resistance via Negatively Regulating the SOCS-3-STAT3 Signaling Pathway. *Journal of Diabetes Research* **2019**, 5483946 (2019).

37. Montagner, A. *et al.* Liver PPARα is crucial for whole-body fatty acid homeostasis and is protective against NAFLD. *Gut* **65**, 1202–1214 (2016).

38. Lu, J., McKinsey, T. A., Zhang, C. L. & Olson, E. N. Regulation of skeletal myogenesis by association of the MEF2 transcription factor with class II histone deacetylases. *Mol. Cell* **6**, (2000).

39. Govaere, O. *et al.* A proteo-transcriptomic map of non-alcoholic fatty liver disease signatures. *Nature Metabolism* **5**, 572 (2023).

40. Chitturi, S. *et al.* The Asia-Pacific Working Party on Non-alcoholic Fatty Liver Disease guidelines 2017-Part 2: Management and special groups. *J. Gastroenterol. Hepatol.* **33**,

971 (2018).

972 41. Younes, R. *et al.* Long-term outcomes and predictive ability of non-invasive scoring

973 systems in patients with non-alcoholic fatty liver disease. *Journal of hepatology* **75**, (2021).

974 42. Transcriptional regulation of hepatic stellate cells. *Adv. Drug Deliv. Rev.* **61**, 497–512

975 (2009).

976 43. Sun, Z. *et al.* Hydronidone induces apoptosis in activated hepatic stellate cells through

977 endoplasmic reticulum stress-associated mitochondrial apoptotic pathway. *J. Gastroenterol.*

978 *Hepatol.* (2024) doi:10.1111/jgh.16635.

979 44. Henkel, A. & Green, R. M. The Unfolded Protein Response in Fatty Liver Disease. *Semin.*

980 *Liver Dis.* **33**, 321 (2013).

981 45. Holzner, L. M. W. & Murray, A. J. Hypoxia-Inducible Factors as Key Players in the

982 Pathogenesis of Non-alcoholic Fatty Liver Disease and Non-alcoholic Steatohepatitis.

983 *Front. Med.* **8**, 753268 (2021).

984 46. Suzuki, T., Shinjo, S., Arai, T., Kanai, M. & Goda, N. Hypoxia and fatty liver. *World J.*

985 *Gastroenterol.* **20**, 15087 (2014).

986 47. Munsterman, I. D. *et al.* Extracellular matrix components indicate remodelling activity in

987 different fibrosis stages of human non-alcoholic fatty liver disease. *Histopathology* **73**,

988 (2018).

989 48. Ipsen, D. H., Lykkesfeldt, J. & Tveden-Nyborg, P. Molecular mechanisms of hepatic lipid

990 accumulation in non-alcoholic fatty liver disease. *Cell. Mol. Life Sci.* **75**, 3313 (2018).

991 49. Wnt/beta-catenin signaling and its modulators in nonalcoholic fatty liver diseases.

992 *Hepatobiliary Pancreat. Dis. Int* **22**, 333–345 (2023).

993 50. Ying, H.-Z. *et al.* PDGF signaling pathway in hepatic fibrosis pathogenesis and

994 therapeutics. *Mol. Med. Rep.* **16**, 7879 (2017).

995 51. TGF-β1 signaling can worsen NAFLD with liver fibrosis backdrop. *Exp. Mol. Pathol.* **124**,

996  104733 (2022).

997 52. Inhibition of Notch1 signaling reduces hepatocyte injury in nonalcoholic fatty liver disease

998  via autophagy. *Biochem. Biophys. Res. Commun.* **547**, 131–138 (2021).

999 53. Xu, X. *et al.* Targeted therapeutics and novel signaling pathways in non-alcohol-associated

1000  fatty liver/steatohepatitis (NAFL/NASH). *Signal Transduction and Targeted Therapy* **7**, 1–39

1001  (2022).

1002 54. Yingling, J. M. *et al.* Preclinical assessment of galunisertib (LY2157299 monohydrate), a

1003  first-in-class transforming growth factor-β receptor type I inhibitor. *Oncotarget* **9**, 6659–6677

1004  (2017).

1005 55. Luangmonkong, T. *et al.* Evaluating the antifibrotic potency of galunisertib in a human ex

1006  vivo model of liver fibrosis. *Br. J. Pharmacol.* **174**, 3107–3117 (2017).

1007 56. Ge, W.-S. *et al.* β-catenin is overexpressed in hepatic fibrosis and blockage of Wnt/β-

1008  catenin signaling inhibits hepatic stellate cell activation. *Mol. Med. Rep.* **9**, 2145–2151

1009  (2014).

1010 57. Gribben, C. *et al.* Acquisition of epithelial plasticity in human chronic liver disease. *Nature*

1011  **630**, 166–173 (2024).

1012 58. Vonderlin, J., Chavakis, T., Sieweke, M. & Tacke, F. The Multifaceted Roles of

1013  Macrophages in NAFLD Pathogenesis. *Cell Mol Gastroenterol Hepatol* **15**, 1311–1324

1014  (2023).

1015 59. Matchett, K. P., Paris, J., Teichmann, S. A. & Henderson, N. C. Spatial genomics: mapping

1016  human steatotic liver disease. *Nat. Rev. Gastroenterol. Hepatol.* 1–15 (2024).

1017 60. Heebøll, S. *et al.* Placebo-controlled, randomised clinical trial: high-dose resveratrol

1018  treatment for non-alcoholic fatty liver disease. *Scand. J. Gastroenterol.* (2016)

1019  doi:10.3109/00365521.2015.1107620.

1020 61. Vacca, M. *et al.* An unbiased ranking of murine dietary models based on their proximity to

human metabolic dysfunction-associated steatotic liver disease (MASLD). *Nature*

Metabolism **6**, 1178–1196 (2024).

62. Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory

requirements. *Nat. Methods* **12**, 357–360 (2015).

63. Anders, S., Pyl, P. T. & Huber, W. HTSeq—a Python framework to work with high-

throughput sequencing data. *Bioinformatics* **31**, 166–169 (2014).

64. Durinck, S., Spellman, P. T., Birney, E. & Huber, W. Mapping identifiers for the integration

of genomic datasets with the R/Bioconductor package biomaRt. *Nat. Protoc.* **4**, (2009).

65. Zhang, Y., Jenkins, D. F., Manimaran, S. & Johnson, W. E. Alternative empirical Bayes

models for adjusting for batch effects in genomic studies. *BMC Bioinformatics* **19**, 1–15

(2018).

66. Saelens, W., Cannoodt, R., Todorov, H. & Saeys, Y. A comparison of single-cell trajectory

inference methods. *Nat. Biotechnol.* **37**, 547–554 (2019).

67. Kleiner, D. E. *et al.* Design and validation of a histological scoring system for nonalcoholic

fatty liver disease. *Hepatology* **41**, 1313–1321 (2005).

68. Fisher, R. A. Statistical Methods for Research Workers. *Breakthroughs in Statistics* 66–70

(1992).

69. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Series B Stat. Methodol.* **57**, 289–

300 (1995).

70. Zhang, B. & Horvath, S. A General Framework for Weighted Gene Co-Expression Network

Analysis. *Stat. Appl. Genet. Mol. Biol.* **4**, (2005).

71. Stoica, P. & Selen, Y. Model-order selection: a review of information criterion rules.

https://ieeexplore.ieee.org/document/1311138.

72. Alvarez, M. J. *et al.* Network-based inference of protein activity helps functionalize the

genetic landscape of cancer. *Nat. Genet.* **48**, 838 (2016).

73. Müller-Dott, S. *et al.* Expanding the coverage of regulons from high-confidence prior

knowledge for accurate estimation of transcription factor activities. *Nucleic Acids Res* **51**,

10934–10949 (2023).

74. Akhmedov, M. *et al.* PCSF: An R-package for network-based interpretation of high-

throughput data. *PLoS Comput. Biol.* **13**, e1005694 (2017).

75. Kuleshov, M. V. *et al.* Enrichr: a comprehensive gene set enrichment analysis web server

2016 update. *Nucleic Acids Res.* **44**, W90–W97 (2016).

76. Lachmann, A. *et al.* ChEA: transcription factor regulation inferred from integrating genome-

wide ChIP-X experiments. *Bioinformatics* **26**, 2438–2444 (2010).

77. Stormo, G. D. Modeling the specificity of protein-DNA interactions. *Quantitative Biology* **1**,

115–130 (2013).

78. Wasserman, W. W. & Sandelin, A. Applied bioinformatics for the identification of regulatory

elements. *Nat. Rev. Genet.* **5**, 276–287 (2004).

79. Gillespie, M. *et al.* The reactome pathway knowledgebase 2022. *Nucleic Acids Res.* **50**,

D687–D692 (2021).

80. Csárdi, G. *et al.* Igraph: Network analysis and visualization. *CRAN: Contributed Packages*

The R Foundation https://doi.org/10.32614/cran.package.igraph (2006).

81. Kanehisa, M. & Goto, S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids*

Res. **28**, (2000).

82. Wrzodek, C., Büchel, F., Ruff, M., Dräger, A. & Zell, A. Precise generation of systems

biology models from KEGG pathways. *BMC Syst. Biol.* **7**, (2013).

83. Romero, P. *et al.* Computational prediction of human metabolic pathways from the

complete human genome. *Genome Biol.* **6**, (2005).

84. Thiele, I. *et al.* A community-driven global reconstruction of human metabolism. *Nat.*

Biotechnol. **31**, (2013).

85. Rodchenkov, I. *et al.* Pathway Commons 2019 Update: integration, analysis and

exploration of pathway data. *Nucleic Acids Res.* **48**, D489–D497 (2019).

86. Türei, D., Korcsmáros, T. & Saez-Rodriguez, J. OmniPath: guidelines and gateway for

literature-curated signaling pathway resources. *Nat. Methods* **13**, 966–967 (2016).

87. Türei, D. *et al.* Integrated intra- and intercellular signaling knowledge for multicellular omics

analysis. *Mol. Syst. Biol.* **17**, e9923 (2021).

88. OmnipathR. *Bioconductor* http://bioconductor.org/packages/OmnipathR/.

89. Yu, G. *et al.* GOSemSim: an R package for measuring semantic similarity among GO terms

and gene products. *Bioinformatics* **26**, 976–978 (2010).

90. Aleksander, S. A. *et al.* The Gene Ontology knowledgebase in 2023. *Genetics* **224**, (2023).

91. Resnik, P. Semantic Similarity in a Taxonomy: An Information-Based Measure and its

Application to Problems of Ambiguity in Natural Language. *jair* **11**, 95–130 (1999).

92. An Information-Theoretic Definition of Similarity.

https://dl.acm.org/doi/10.5555/645527.657297 doi:10.5555/645527.657297.

93. Wang, J. Z., Du, Z., Payattakool, R., Yu, P. S. & Chen, C.-F. A new method to measure the

semantic similarity of GO terms. *Bioinformatics* **23**, 1274–1281 (2007).

94. Charmpi, K., Chokkalingam, M., Johnen, R. & Beyer, A. Optimizing network propagation for

multi-omics data integration. *PLoS Comput. Biol.* **17**, e1009161 (2021).

95. Friedman, S. L., Neuschwander-Tetri, B. A., Rinella, M. & Sanyal, A. J. Mechanisms of

NAFLD development and therapeutic strategies. *Nat. Med.* **24**, 908–922 (2018).

96. En masse organoid phenotyping informs metabolic-associated genetic susceptibility to

NASH. *Cell* **185**, 4216–4232.e16 (2022).

97. Graupera, I. *et al.* LiverScreen project: study protocol for screening for liver fibrosis in the

general population in European countries. *BMC Public Health* **22**, 1–10 (2022).

1096 98. Hendriks, D. *et al.* Engineered human hepatocyte organoids enable CRISPR-based target

1097   discovery and drug screening for steatosis. *Nat. Biotechnol.* **41**, 1567–1581 (2023).

1098 99. Lee, E., Korf, H. & Vidal-Puig, A. An adipocentric perspective on the development and

1099   progression of non-alcoholic fatty liver disease. *Journal of hepatology* **78**, (2023).

1100 100.A network-based computational and experimental framework for repurposing compounds

1101   toward the treatment of non-alcoholic fatty liver disease. *iScience* **25**, 103890 (2022).

1102 101.The Molecular Signatures Database Hallmark Gene Set Collection. *Cell Systems* **1**, 417–

1103   425 (2015).

1104 102.Martens, M. *et al.* WikiPathways: connecting communities. *Nucleic Acids Res.* **49**, D613–

1105   D621 (2020).

1106 103.Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

1107   RNA-seq data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).

1108 104.The anatomy of a large-scale hypertextual Web search engine. *Computer Networks and*

1109   *ISDN Systems* **30**, 107–117 (1998).

1110 105.Guilliams, M. *et al.* Spatial proteogenomics reveals distinct and evolutionarily conserved

1111   hepatic macrophage niches. *Cell* **185**, (2022).

1112 106.Hao, Y. *et al.* Dictionary learning for integrative, multimodal and scalable single-cell

1113   analysis. *Nature Biotechnology* **42**, 293–304 (2023).

1114 107.Vathrakokoili, P. A. *et al.* CATD: a reproducible pipeline for selecting cell-type

1115   deconvolution methods across tissues. *Bioinformatics advances* **4**, (2024).

1116 108.Zheng, S. C. *et al.* EpiDISH web server: Epigenetic Dissection of Intra-Sample-

1117   Heterogeneity with online GUI. *Bioinformatics (Oxford, England)* **36**, (2019).

1118 109.Baron, M. *et al.* A Single-Cell Transcriptomic Map of the Human and Mouse Pancreas

1119   Reveals Inter- and Intra-cell Population Structure. *Cell systems* **3**, (2016).

1120 110.Tsoucas, D. *et al.* Accurate estimation of cell-type composition from gene expression data.

Nature Communications **10**, 1–9 (2019).

111. Newman, A. M. *et al.* Robust enumeration of cell subsets from tissue expression profiles. *Nature Methods* **12**, 453–457 (2015).

112. Hao, Y., Yan, M., Heath, B. R., Lei, Y. L. & Xie, Y. Fast and robust deconvolution of tumor infiltrating lymphocyte from expression profiles using least trimmed squares. *PLoS computational biology* **15**, (2019).

113. Chu, T., Wang, Z., Pe'er, D. & Danko, C. G. Cell type and gene expression deconvolution with BayesPrism enables Bayesian integrative analysis across bulk and single-cell RNA sequencing in oncology. *Nature Cancer* **3**, 505–517 (2022).

114. Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction of single-cell gene expression data. *Nature Biotechnology* **33**, 495–502 (2015).

115. Mendez, D. *et al.* ChEMBL: towards direct deposition of bioassay data. *Nucleic acids research* **47**, (2019).

116. Yilmaz, Y. & Eren, F. Serum biomarkers of fibrosis and extracellular matrix remodeling in patients with nonalcoholic fatty liver disease: association with liver histology. *European journal of gastroenterology & hepatology* **31**, (2019).

117. Coilly, A., Desterke, C., Guettier, C., Samuel, D. & Chiappini, F. FABP4 and MMP9 levels identified as predictive factors for poor prognosis in patients with nonalcoholic fatty liver using data mining approaches and gene expression analysis. *Scientific reports* **9**, (2019).

118. Toyoda, H. *et al.* Higher hepatic gene expression and serum levels of matrix metalloproteinase-2 are associated with steatohepatitis in non-alcoholic fatty liver diseases. *Biomarkers : biochemical indicators of exposure, response, and susceptibility to chemicals* **18**, (2013).

119. Qi, S., Wang, C., Li, C., Wang, P. & Liu, M. Candidate genes investigation for severe nonalcoholic fatty liver disease based on bioinformatics analysis. *Medicine* **96**, (2017).

Figure Legends

Figure 1. Overview of the approach used in this study to investigate MASLD progression. **A.** Integration of public datasets and MASLD trajectory inference by the pseudo-temporal ordering of patient biopsy transcriptomic data **B.** Combining phenotype-specific gene co-expression analysis with upstream pathway inference to define a MASLD network. **C.** Cell type deconvolution to study the dynamics of cell type composition changes. **D.** Random Forest regression model for the prediction of patient's position on the inferred MASLD trajectory based on a set of biomarkers. Training was done on the initial dataset, and testing was done on independent datasets. The model deployed different gene signatures, while random gene sets were used as controls to validate the high model's performance. **E.** Identification of deregulated cell type-specific processes and potential biomarker genes that could be targeted in specific disease stages.

Figure 2. Pseudo-temporal ordering of patients captures disease progression and identifies MASLD gene signature. A. Pseudo-temporal ordering of patients based on transcriptomic data recapitulates disease progression based on individual phenotypes, i.e. steatosis, ballooning, inflammation and fibrosis, and based on the NAS and MASLD scores. **B-C.** Pseudotemporal ordering of two independent orthogonal datasets based on the 145 genes that are most predictive of the trajectory provides a linear and clear separation of the disease stages (EPoS dataset (**B**) and Gubra dataset (**C**)). **D.** Functional enrichment analysis results of these 145 genes using EnrichR, where the Combined Score (c) has been calculated as $c = \log(p\_value)$ * z_score, with p_value representing the Fisher exact test outcome, and z-score the deviation from the expected rank.

Figure 3. Phenotype-specific gene modules are associated with the different MASLD variables. A. Network of the different gene co-expression modules is associated with NAS score, Steatosis, Ballooning, and Fibrosis. Each module is depicted by a different colour and can be linked with one or multiple variables. **B.** TFs whose regulons are enriched in the modules. The figure shows TFs per module coloured according to the adjusted p-value of the enrichment test. The presented TFs have been identified as enriched in multiple modules and deregulated at least in one stage of the disease trajectory (see sliding window analysis results). The complete list of TFs derived from the enrichment analysis is given in Supplementary Table 6. **C.** Enrichment of MASLD network in known MASLD genes curated from the literature (top) or from MSigDB (bottom).

Figure 4. Sliding window analysis to study the landscape of molecular changes along the MASLD trajectory. A. Results from TF activity analysis in the discrete patients stratification and the pseudotemporal trajectory. The colour indicates the change in TF activity compared to the previous disease stage. **B.** Mean phenotype and NAS pathologist scores along the SW trajectory.

Figure 5. Results of cell type deconvolution analysis. A. Cell type deconvolution of patient transcriptomic data along our MASLD trajectory **B.** Prediction of cell types with which the deregulated processes are associated after excluding processes that could have occurred by the mere change in abundance of cell types.

Figure 6. Random Forest regression model for the prediction of MASLD patients' position on the inferred disease trajectory. A. Model performance using different gene sets in the primary UCAM/VCU dataset (5-fold cross validation), and validation in two independent datasets used solely for testing. Randomisation was done on 1000 times. **B.** Compared predicted and actual patient positions on the trajectory using an independent dataset (GUBRA), based solely on a 24-gene signature. **C.** Violin plot showing the distribution of the predicted patients' position on the trajectory (GUBRA dataset). **D.** Network-based deregulation analysis of the 24-gene signature to identify potential targets. The color scale represents the -log10(adjusted p-value) of differential expression. Genes are deemed deregulated in a SW if part of the respective deregulated subnetworks. Drugs in max phase 4 were retrieved for targets in the relative deregulated pathways.

Supplementary Figure legends

Supplementary Figure 1. MASLD patients along the continuous transcriptomics-driven trajectory. The colouring is based on the different scores given by the histopathologists. The triangles show the average horizontal position of each scoring group on the trajectory. **A.** NAS score (ranging from 0 to 7). **B.** Steatosis scoring (0-3). **C.** Inflammatory damage (0-3). **D.** Ballooning (0-2). **E.** Fibrosis (0-4).

Supplementary Figure 2. Violin plots illustrate the distribution of the patients in each histopathological scoring group. A. Steatosis. **B.** Inflammatory damage. **C.** Ballooning. **D.** Fibrosis. **E.** NAS score. Each panel shows the Pearson correlation coefficient (R score) for the mean trajectory positions of each scoring group. The p-values are the results of ANOVA followed by Tukey's pairwise comparisons for the distributions of patients across the histopathological

scores. With dashed lines are indicated the statistically significant groups (*p-value < .05, ** p-

value < .005, *** p-value < .0005).

Supplementary Figure 3. Trajectory inference on independent MASLD transcriptomic

datasets and selection of a subset of genes based on a Random Forest classification

model. A. Trajectory inference for the independent EPoS dataset using the whole transcriptome.

Color coding depicts MASLD stages as published with the original RNA-Seq data. **B.** Trajectory

inference for the independent GUBRA dataset. The population was characterised as Healthy

(non-obese individuals without MASLD/MASH), Obese (healthy but obese individuals without

MASLD/MASH), MASL, and MASH. **C.** Distribution of the "Mean Decrease Accuracy" scores

derived from our Random Forest classification model. **D.** Selection of the 145 genes with positive

scores, representing the final gene signature able to predict a patient's disease state, as well as

key histopathology scores for steatosis, ballooning, inflammation, and fibrosis. **E.** Stage prediction

of the GUBRA dataset based on a Random Forest classifier trained on the original dataset (Stage

Division 1: predicting Healthy/Obese vs MASL vs MASH stages; Stage Division 2: predicting

Healthy/Obese/MASL vs MASH stages).

Supplementary Figure 4. Mean MASH component score in each Sliding Window (SW). The x-

axis represents the 13 SWs, while the y-axis represents the mean score in each one of these

SWs according to **A.** Steatosis, **B.** Inflammation, **C.** Ballooning, **D.** Fibrosis. The plots additionally

show the Pearson's correlation coefficient (R score) and the corresponding p-value.

Supplementary Figure 5 Overview of modulation of modules and pathways along sliding

windows. A. Enrichment of histological phenotype-specific modules along our MASLD trajectory

B. Reactome pathway enrichment analysis comparing the enrichment (and direction) using the

Sliding Window (SW) approach. The results are derived from the differentially expressed genes

and the deregulated transcription factors (Log2FoldChange, NES scores and FDR values) when

comparing consecutive SWs (i.e., SW_x vs SW_{x-1}). The "red" and "blue" colours represent the up-

and down-regulated pathways, respectively, according to the average activation score.

Supplementary Figure 6. Random Forest regression model for the prediction of MASLD

patients' position on the inferred disease trajectory for the primary and validation

(independent) dataset. A. Predicted vs real patients' positions on the trajectory for the primary

dataset. The dataset has been split into 80% training and 20% testing. The plot shows the average

after running the model 1000 times. The different colours show the different MASLD stages

according to the histopathological assessment. Only the 24 gene signature has been deployed

for the prediction. **B.** Violin plot showing the distribution of patients on the inferred trajectory based

on the predictions from panel A. **C-D.** Similar to panels "A" and "B", but for the independent

validation dataset (EPoS). The model training was done entirely on the primary dataset. **E-F.**

Violin plots showing the distribution of patients on the EPoS trajectory, based on the 24 gene

intersection, according to the NAS and Fibrosis scores (as assessed by the histopathologists). **G-**

H 1000 (G) and 24 (H) random genes are not able to place patients along the trajectory.

Supplementary Figure 7. Ranked independent dataset (Gubra) individuals according to

their predicted trajectory positions. The first half of the trajectory predicts the healthy/obese,

while the second part predicts the MASL/MASH. The Mean Squared Error is based on the results

from panel C.

Supplementary Figure 8. Sanity checks and validation for dataset integration. **A.** PCA plot

of the merged UCAM/VCU quantile normalised counts. There is a clear batch effect separating

the patients into two clusters according to the dataset from which they are derived. **B.** COMBAT

batch effect correction solved the problem of the batch effect with patients altogether in a cluster

and a clear disease progression pattern from the right towards the later disease stages on the

left. The outlier (sample 5) was removed from all downstream analyses. **C-D.** Normalised counts

before and after batch effect correction, separately for each disease stage. The data suggest that

COMBAT has not significantly changed the expression of the genes, given the high correlation

(R > 0.97 in all MASLD stages in VCU (C) and UCAM (D) datasets). **E.** Correlation of the principal

components (PCs) with the variables of interest. The first 10 PCs have been correlated using

Pearson correlation with 6 continuous variables of interest: age, NAS score, Fibrosis, Fat

(Steatosis), Ballooning, and Inflammation. **F.** Heatmap of the quantile normalised counts before

and after the merging/batch effect correction for UCAM and VCU datasets. The rows show genes

previously linked to MASLD[116–119]. The data are shown in a "fold change vs the first disease stage"

of each dataset, respectively ("red" = increase; "blue" = decrease; "white" = unchanged; "green

datasets": UCAM and VCU before correction; "blue datasets": UCAM and VCU after correction;

the asterisks "*" in the column names indicate that the samples have been batch effect corrected).

Note that the control group (first columns in each block) in the VCU dataset is CTRL, while in

UCAM is MASL.

Supplementary Table legends

Supplementary Table 1. Information about all MASLD patients used in this study (UCAM and VCU datasets). Each patient is given the histopathology score (steatosis, ballooning, inflammation, fibrosis, NAS score) and the MASLD stage, as well as additional metadata (age, sex), the Sliding Window (SW) number assigned to the patient, the order in the trajectory (1 to 136), the position according to PC1, the dataset (UCAM or VCU), and the batch corrected normalised expression value per gene.

Supplementary Table 2. Patients' clinical characteristics when divided in CTRL, MASL, MASH F0-1, MASH F2, MASH F3-4. Statistical significance is assessed by ANOVA. The letters next to the values indicate post-hoc analysis significance based on Tukey's test; "[a]" means reference group; when groups show different letters, they should be considered statistically different at the post-hoc comparison; groups showing the same letter are statistically not-significant at the post-hoc comparison.

Supplementary Table 3. Random forest's accuracy and AUC were tested on UCAM/VCU and GUBRA datasets using the top 145 genes. The UCAM/VCU dataset has been divided into training (80%) and testing (20%) sets and the metrics given in the table correspond to the average for running on the test set 1000 times. The gene signature predicts the different MASLD characteristics (Steatosis, Inflammation, Ballooning, Fibrosis) and the MASLD stage. For the GUBRA dataset predictions, the whole UCAM/VCU dataset has been used as a training set. The only available information for this external dataset is the MASLD stage. Finally, for both datasets, a random set of 20, 145, and 5000 genes (average for 1000 runs) verify the robustness of the selected genes.

Supplementary Table 4. Phenotype-specific gene co-expression modules associated ("Y") or not associated ("N") with the MASLD variables. Each column includes the set of genes constituting each gene module.

Supplementary Table 5. Reactome pathway enrichment analysis on the filtered gene sets of each module.

Supplementary Table 6. Transcription Factor (TF) enrichment analysis for each significant-identified gene module.

Supplementary Table 7. Reactome pathway enrichment analysis on the TFs of Supplementary

Table 6.

Supplementary Table 8. Unified inferred global MASLD network according to our analysis.

Supplementary Table 9. Genes previously associated with MASLD.

Supplementary Table 10. Patient assignment to each of the 13 Sliding Windows (SWs).

Supplementary Table 11. VIPER analysis results comparing consecutive Sliding Windows

(SWs). The results are derived from the differentially expressed genes of the previous steps

(Log2FoldChange and p-values). The table is organised in 3-plets showing for each comparison

the adjusted p-value and the Normalised Enrichment Score (NES). NES determines whether a

TF has significantly more "activated" than "inhibited" predictions (NES > 0) or vice versa (NES <

0).

Supplementary Table 12. Complete table of processes (pre-pruning) and associations to cell

types.

Supplementary Table 13. Table of drugs known to target genes within the 24 gene biomarker-

associated subgraphs extracted from our MASLD network

A — Human MASLD Trajectory

MASLD score → CONTROLS, MASL, MASHF0, MASHF1, MASHF2, MASHF3, MASHF4

NAS score → 0, 1, 2 3, 4, 5 6, 7

STEATOSIS → 0, 1, 2, 3

BALLOONING → 0, 1, 2

INFLAMMATION → 0, 1, 2, 3

FIBROSIS → 0, 1, 2, 3, 4

B — EPoS (Top 145 genes)

MASL, MASHF01, MASHF2, MASHF3, MASHF4

C — GUBRA (Top 145 genes)

HEALTHY, OBESE, MASL, MASH

D

	Pathway
***	ECM−receptor interaction
***	AGE−RAGE signaling
***	Protein digestion and absorption
***	Rheumatoid arthritis
***	IL−17 signaling pathway
**	Complement and coagulation cascades
***	Focal adhesion
**	Glycerolipid metabolism
***	Cytokine−cytokine receptor interaction
**	Relaxin signaling pathway
*	Leukocyte transendothelial migration
**	Chemokine signaling pathway
**	Proteoglycans in cancer
*	TGF−beta signaling pathway
*	PI3K−Akt signaling pathway
*	Pathways in cancer

Combined Score: 1, 0.8, 0.6, 0.4, 0.2, 0

A

Discrete approach | MASLD pseudotemporal trajectory (SWx vs SWx-1)

TF Activity: -log(FDR) x sign(NES)

Mild vs Control | Moderate vs Mild | Severe vs moderate

B

(Min-Max scores)

NAS score

Fibrosis
Steatosis
Inflammation
Ballooning