# Mapping spatially-resolved transcriptomes in systemic sclerosis

Zhijian Li[1,2], Aleix Rius Rigau[3,4], Wenjie Xie[5], Xiaohang Shao[6], Yi-Nan Li[7,8], Alexandru-Emil Matei[7,8], Hejian Zou[9,10,11], Luca Pinello[1,2,12], Jörg H.W. Distler[7,8]*, Rui He[5,13]*, and Minrui Liang[9,10,11]*

[1]Broad Institute of Harvard and MIT, Cambridge, MA, USA

[2]Molecular Pathology Unit, Center for Cancer Research, Massachusetts General Hospital, Boston, MA, USA

[3]Department of Internal Medicine 3, Rheumatology and Clinical Immunology, Friedrich-Alexander-University Erlangen-Nürnberg (FAU) and University Hospital Erlangen, Erlangen, Germany.

[4]Deutsches Zentrum Immuntherapie (DZI), FAU Erlangen-Nürnberg and University Hospital Erlangen, Erlangen, Germany.

[5]Department of Immunology, School of Basic Medical Sciences, Fudan University, Shanghai, China.

[6]Department of Nephrology and Rheumatology Immunology, Ningbo Medical Center Lihuili Hospital, Ningbo, China.

[7]Clinic for Rheumatology, University Hospital Düsseldorf, Medical Faculty of Heinrich Heine University, Düsseldorf, Germany.

[8]Hiller Research Center, University Hospital Düsseldorf, Medical Faculty of Heinrich Heine University, Düsseldorf, Germany.

[9]Department of Rheumatology, Huashan Hospital, Fudan University, Shanghai, China.

[10]Institute of Rheumatology, Immunology and Allergy, Fudan University, Shanghai, China.

[11]Huashan Rare Disease Center, Huashan Hospital, Fudan University, Shanghai, China.

[12]Department of Pathology, Harvard Medical School, Boston, MA, USA.

[13]National Clinical Research Center for Aging and Medicine, Huashan Hospital, Fudan University, Shanghai, 200040, China.

* Correspondence to Prof. Minrui Liang: mliang10@fudan.edu.cn; Prof. Rui He: ruihe@fudan.edu.cn; Univ. Prof. Dr. med. Jörg H.W. Distler: Joerg.Distler@med.uni-duesseldorf.de;

**Abstract**

Systemic sclerosis (SSc) is a prototypical fibrotic disease with high mortality and limited treatment options. Despite advances in single-cell RNA sequencing (scRNA-seq), the comprehensive understanding of cellular heterogeneity and cell-cell interaction within the fibrogenesis microenvironment remains limited. We generated spatially resolved transcriptome maps from healthy and SSc skin and built a scRNA-seq atlas to map the single-cell data to spatial space. This enabled us to identify a fibrotic niche, enriched with fibroblasts and macrophages, which is significantly expanded in SSc and correlated with clinical outcome. We revealed disease-specific cell states of fibroblasts and macrophages, and evaluated their spatial dependency on other cell types. We identified selective expression of ACKR3 in fibroblast progenitors that diminishes with SSc progression, which may serve to regulate CXCL12/CXCR4-mediated macrophage recruitment and fibrotic remodeling. Together, we provided an in-depth description at cellular and spatial level of fine-tuned regulatory events occurring in SSc, offering spatiotemporal insights.

**Introduction**

Systemic Sclerosis (SSc), an orphan autoimmune disease, is characterized by the progressive fibrosis affecting skin and multiple organs, holding the highest mortality among all rheumatic diseases[1]. The development of effective therapeutic strategies for SSc is impeded by a lack of comprehensive understanding of its pathogenesis and the limited translatability of current preclinical models. Importantly, studying SSc as a prototypical systemic fibrotic disease could yield new insights into the pathogenesis of fibrotic diseases, and aid in the discovery of novel therapeutic approaches.

SSc is characterized by a complex interplay of features that include vasculopathy, inflammation, and fibrosis. Within this triad, fibroblasts have been recognized as pivotal effector cells in the production of extracellular matrix (ECM), largely contributing to developing fibrotic tissue remodeling in SSc. Fibroblasts sense and receive signaling from

the neighboring cells, which are conveyed and converge to the shared and distinct

profibrotic pathways[2]. The coordination of fibroblasts with macrophages can shape their

cellular response, therefore orchestrating the initiation or resolution of fibrosis across

tissues[3–7]. Beyond the conventional macrophage polarization paradigm, new populations

of macrophages have been linked with tissue repair and fibrotic remodeling[3–5,7,8]. For

instance, phenotypically distinct macrophages (Lyve1$^{lo}$MHCII$^{hi}$ and Lyve1$^{hi}$MHCII$^{lo}$)

manifested specific niche-dependent functional roles in pulmonary and cardiac fibrosis[8].

However, it remains largely unexplored whether and how the spatial neighbourhood in

the tissue architecture affects the crosstalk between fibroblast and macrophage subsets,

potentially driving fibrotic remodeling in SSc.

Recent studies using single-cell RNA sequencing (scRNA-seq) have provided an

unprecedented opportunity to study the molecular mechanisms of SSc at single-cell

resolution[9–13]. Pioneering studies have characterized fibroblast types and states in

SSc[9,11,13]. While these studies have provided valuable insights, they are significantly

constrained by the lack of spatial resolution due to the reliance on disaggregated tissue

analysis. We recently demonstrated cellular niches centered around endothelial cells or

fibroblasts in SSc using a panel of up to 40 markers through multiplex spatial

proteomics[14,15]. Thus, a spatially-resolved characterization of genome-wide

transcriptomics of SSc is needed.

In this work, we profiled human skin from SSc patients and healthy controls using spatial

transcriptomics (ST) technique based on the 10x Visium formalin-fixed paraffin-

embedded (FFPE) platform and spatial enhanced resolution omics sequencing (Stereo-

seq)[16]. To increase the cellular resolution of our ST data, we integrated two publicly

available scRNA-seq datasets, creating a human skin transcriptome atlas and mapping

cell type information onto spatial regions. We validated our results using additional spatial

profiling techniques with higher resolution, such as imaging mass cytometry (IMC), as

well as time-course experimental skin fibrosis. These approaches enabled us to reveal a

detailed characterization of pathological cellular communication between fibroblast and

macrophage subsets. This has provided insights into therapeutic target discovery by

disrupting disease-driving cellular niches and circuits in SSc.

**Fig. 1 Spatial profiling of human skin. a,** *Overview of experimental design and analysis*

*workflow.* **b,** *UMAP showing integrated scRNA-seq datasets as colored by cell types (left) and*

*conditions (right). SSc skin n=104, Healthy skin n=64. **c,** Dotplot showing gene expression for*
*marker genes. The colors refer to the mean expression of the genes and the dot sizes represent*
*the proportion of cells expressing the marker genes. **d,** Visualization maps based on 10× Visium.*
*SSc skin n=10, Healthy skin n=4. Left: vWF, COL1A1, and KRT1 gene expressions in spatial*
*transcriptomics data. Middle: representative cell types estimated by cell2location. Right:*
*visualization of pathway activity in spatial space. **e,** Integration of scRNA-seq and Stereo-seq data*
*colored by protocols (left) and cell types (right). **f,** Visualization of cell types in spatial space based*
*on Stereo-seq data. SSc skin n=3, Healthy skin n=1. The experiments were performed on*
*adjacent tissue sections to those used for 10× Visium.*
109

**Results**

**Profiling of SSc and healthy skin with spatially-resolved transcriptomics**
We collected a total of 14 human skin biopsies from 10 SSc patients enriched for early-
stage, diffuse cutaneous involvement, with clinically active disease and 4 sex- and age-
matched normal controls (**Fig. 1a**; **Supplementary Table 1**). For each biopsy, we
generated spatially resolved transcriptomics (ST) using the 10x Visium platform. After
removing low-quality spots across all samples, we retained 13,204 high quality spots with
a median detection of 2,265 genes per spot (**Supplementary Fig. 1; Supplementary**
**Table 2**). To enhance spatial resolution, we also performed high-resolution ST profiles on
four samples, one from a healthy donor and three from SSc patients, using Stereo-seq.
After controlling data quality, we retained 40,566 spots across these samples. We
additionally analyzed two scRNA-seq datasets generated from skin tissues of SSc
patients (n = 104) and healthy donors (n = 64)[9,10]. The cells were re-annotated for each
dataset based on clustering or the original labels (see **Methods**; **Supplementary Fig.**
**2a-d**). We integrated the scRNA-seq datasets and observed that the same cell types were
well-aligned in the UMAP space, which was further confirmed by the cell-type-specific
marker genes identified based on the integrated profiles (**Fig. 1b-c; Supplementary Fig.**
**2e-g**). Together, these data provide a comprehensive multi-modal map of human skin
tissues in health and SSc, capturing both spatial and single-cell transcriptomic
landscapes.
130

We next deconvoluted the cell-type compositions of each spot in 10x Visium data using

the scRNA-seq as reference. The estimated cell type abundances from spatial

transcriptomics were generally co-localized with their respective marker genes (**Fig. 1d**).

Moreover, we observed that different cell types were mapped to the expected locations.

For example, keratinocytes were found localized within the epidermis and surrounding

the hair follicle. Endothelial cells were closely associated with vasculature in the dermis.

Fibroblasts were distributed throughout the papillary layer and reticular layer of the

dermis. We inferred the pathway activity for each spot and observed the enriched

expression of genes associated with some well-known pathways, such as Janus kinase

signal transducer and activator of transcription protein (JAK-STAT) and transforming

growth factor (TGF)-β, reflecting the link between cellular location and the corresponding

function (**Fig. 1d**). We transferred the cell type annotation from scRNA-seq to Stereo-seq

through data integration (**Fig. 1e**). Spatial visualization of the predicted labels in tissues

showed a similar cellular distribution as 10x Visium data (**Fig. 1f**). Collectively, our

integrative analyses on spatial transcriptomics and scRNA-seq data enable us to map

cell abundance and signaling pathways across zonations in skin samples.

**Fig. 2 Characterization of cell-type niches in SSc and healthy skin. a,** *UMAP showing the*

*integrated spatial data as colored by cell-type niches. **b,** Visualization of cell type proportion in UMAP space. **c,** Heatmap showing the enrichment of cell types in niches. Each column represents a niche, and each row represents a cell type. Colors refer to the z-score transformed proportions of each cell type across different niches. **d,** Boxplot comparing the proportion of each niche between SSc and healthy skin samples. Each dot represents a sample. P-values are calculated using the Wilcoxon rank-sum test (unpaired; two sides). **e,** Visualization of cell-type niches in spatial space for the healthy and SSc samples using 10x Visium data. **f,** Heatmap showing pathway activity for each niche. **g,** Visualization of fibroblasts and macrophages in spatial space using 10x Visium data. **h,** Visualization of cell types in spatial space using Stereo-seq data. **i,** Correlations between the proportion of the indicated niches with clinic statistics across all SSc subjects. Statistical significance was determined by the Spearman test for a two-variable correlation. The correlation coefficient r is indicated in figures (\* p < 0.05). **Abbreviations**: CRP (C-reactive protein), mRSS (Modified Rodnan Skin Score), PLT (platelet count), C3 (complement component 3), ILD (interstitial lung disease), ALT (alanine aminotransferase), C4 (complement component 4), ESR (erythrocyte sedimentation rate), HGB (hemoglobin), LDH (lactate dehydrogenase), AST (aspartate aminotransferase), CK (creatine kinase), DU (digital ulcer), IgG (immunoglobulin G), sPAP (systolic pulmonary arterial pressure), SpO2 (oxygen saturation), Cr (creatinine), TnT (troponin T), EUSTAR activity index (European Scleroderma Trials and Research Group activity index), r-GT (gamma-glutamyl transferase), TB (total bilirubin), WBC (white blood cell count), and Non-RP Disease Course (non-raynaud's phenomenon disease course).*

## Characterization of tissue organization in SSc and healthy skin

To characterize the cellular colocalization events in human skin, we integrated the 10x Visium samples (n = 14) based on the inferred cell-type proportions within each spot (**Supplementary Fig. 3a**). Specifically, we conducted spot deconvolution to determine the proportions of different cell types within each spot, and subsequently clustered the spots based on these identified proportions (**Fig. 2a**). Clustering analysis identified 12 compositionally distinct populations, defined as cell-type niches that represent potential structural building blocks shared across different samples. Visualization of the cell-type abundances and marker genes of these niches revealed that they were strongly associated with the corresponding cell types (**Fig. 2b; Supplementary Fig. 3b-c**). For example, niches 1 and 3 (defined as vasculature niches) were enriched with endothelial

cells and pericytes; niches 0 and 10 (defined as fibrotic niches) were enriched with

fibroblasts; niches 6, 7 and 8 (defined as epidermis niches) were largely associated with

keratinocyte as the representative cell types and along with the infiltration of Langerhans

cells; niche 11 was related to vascular smooth muscle cells (VSMC) and therefore defined

as a VSMC niche; niche 2, 4, 5, and 9 (defined as immune niches) were related to immune

cells (**Fig. 2c**).

Furthermore, we observed a significant increase in the proportions of three niches (0, 1,

and 4) in the SSc skin compared to healthy skin (**Fig. 2d-e; Supplementary Fig. 3d**).

Notably, niche 0 was infiltrated with a high prevalence of fibroblasts, macrophages,

conventional dendritic cells (cDCs), and mast cells, indicating a potential immune

microenvironment within a fibrogenic niche. Niche 1 was characterized by an enrichment

of endothelial cells, pericytes, macrophages, cDCs, and mast cells, suggesting a

perivascular niche associated with immune cell infiltration in SSc skin. Additionally, niche

4 was distinguished by a significant presence of plasma cells, melanocytes, and neural

cells (**Fig. 2c**). Moreover, SSc skin exhibited a more reorganized structure, highlighting a

tendency towards spatially educated organization, with a dominance in niche distribution

(**Fig. 2e**).

To investigate the featured signaling pathway that characterizes each specific niche, we

computed pathway activity. Fibroblast-dominant niche 0 showed the highest activity of

the TGF-β signaling pathway. Niche 9, composed of various immune cells, displayed the

strongest activity of JAK-STAT signaling (**Fig. 2f**). These findings supported the essential

roles of TGF-β and JAK-STAT signaling in the pathogenesis of SSc as expected[2],

suggesting a convergence of multiple signaling activities within the structural organization

in SSc fibrosis. KEGG pathway enrichment analysis based on marker genes also

revealed high activity in the TGF-β signaling in niche 0 (**Supplementary Fig. 3e**).

Indeed, we observed colocalization of fibroblasts and macrophages in our 10x Visium

and Stereo-seq data (**Fig. 2g-h**). To further validate these results, we performed imaging

mass cytometry (IMC) and identified the cellular neighborhood (CN) community as

described previously[14]. We observed similar niches comprising stromal cells and

macrophages (CN8), vascular ECs, and immune cells, including macrophages in CN5

(**Supplementary Fig. 4**). To explore the clinical relevance of these niches, we correlated

their proportions with clinical data across the SSc samples. Interestingly, we observed a

positive Spearman correlation between the proportion of niche 0 and the current total

modified Rodnan's skin score (mRSS) ($r = 0.721$, $p = 0.023$), C-reactive protein (CRP)

levels ($r = 0.75$, $p = 0.066$) and platelet counts ($r = 0.697$, $p = 0.031$), which are interpreted

as clinical indicators for disease severity or activity of SSc  (**Fig. 2i**). Additionally, we

observed a significant inverse correlation between the proportion of niche 1 and the non-

RP disease duration ($r = -0.756$, $p = 0.014$). In summary, our spatial multi-omics revealed

a distinct structural organization associated with disease outcome of SSc, suggesting a

potential coordinated regulation of immune-stromal interactions that may drive the

disease progression.

***Fig. 3 Characterization of fibroblast sub-populations. a,*** *UMAP showing sub-populations of*
*fibroblasts based on scRNA-seq data.* ***b,*** *Dot plot showing the marker genes of each subset.* ***c,***
*Bar graph comparing the proportion of each fibroblast subset between healthy and SSc samples.*
*Each dot represents a sample. P-values are calculated using the Wilcoxon rank-sum test*
*(unpaired; two sides).* ***d,*** *Visualization of SCARA5, POSTN, and COMP expression in spatial*
*space.* ***e,*** *Visualization of module score of Fib1, Fib2, and Fib9 in spatial space.* ***f,*** *Visualization*
*of TF activity of JUN and SMAD3 in spatial space.* ***g,*** *Visualization of TGF-β pathway activity in*
*spatial space.* ***h,*** *Heatmap showing correlations between expressions of SCARA5 and POSTN*
*and different clinical outcomes of SSc. The colors in the upper left square represent Spearman's*
*r value, while those in the lower right square represent p values (\* p < 0.05).* ***i,*** *Expression of*
*SCARA5, POSTN, ACTA2, and COMP overlaying with UMAP.* ***j,*** *UMAP showing the*
*differentiation trajectory from Fib1 to Fib2 (left), and Fib1 to Fib9 (right). Colors refer to pseudotime*
*points.* ***k,*** *Heatmap showing the gene expression dynamics during the trajectory from Fib1 to Fib2*
*(left), and Fib1 to Fib9 (right).* ***l,*** *co-localization between major cell types and fibroblast sub-*
*clusters (Fib1, Fib2 and Fib9) as measured within the same spot (left) and neighboring spots*
*(right).* ***m,*** *Visualization of the abundance of macrophages and VSMCs and the state scores of*
*Fib1 and Fib2 in a representative SSc skin sample.*

**Characterization of fibroblast sub-clusters in healthy and SSc skin**

To investigate the molecular mechanisms driving fibrogenesis in human SSc, we next

clustered scRNA-seq data for fibroblasts and identified ten sub-clusters across the

datasets and conditions (**Fig. 3a; Supplementary Fig. 5a**). Consistent with previous

studies[9,10], we found the corresponding fibroblast populations, including

*SCARA5$^+$/PCOLCE2$^+$/LGR5$^+$* fibroblasts (Fib1), *MYOC$^+$/C3$^+$/APOD$^+$* fibroblasts (Fib3),

*CCL19$^+$/C7$^+$/APOE$^+$* fibroblasts (Fib4), *SFPR2$^+$/WIF1/APCDD1$^+$* fibroblasts (Fib6),

*COCH$^+$/SPARCL1$^+$* fibroblasts (Fib7), *COMP$^+$/SFPR2$^+$/PRSS23$^+$* fibroblasts (Fib9), and

*IGFBP2$^+$/ANGPTL7$^+$* fibroblasts (Fib10) (**Fig. 3b**). Moreover, Fib5 was characterized by

the expression of *FOS*, *JUNB*, *JUN* and *FOSB* which were components of activator

protein 1 (AP1) known as the core transcription factors in SSc fibrosis[17,18]. Fib1

demonstrated the upregulation of *SCARA5*, which has been recently identified as a

marker for progenitor fibroblast in human myocardial and kidney fibrosis[19–21]. Fib2 and

Fib9 were characterized by the expression of Periostin (*POSTN*), which encodes an

extracellular matrix protein that can amplify TGF-β signaling[22,23]. In addition, Fib2 also

exhibited a high expression of myofibroblast hallmark *ACTA2*. Together, these suggested

that Fib2 and Fib9 as terminally-differentiated ECM-producing myofibroblasts.

Next, we compared the proportion of each sub-cluster between healthy and SSc samples.

We found that Fib1 showed a significant reduction in SSs skin (average fraction = 0.137)

compared to healthy skin (average fraction = 0.319) ($p$ = 3.5e-10) (**Fig. 3c**). In contrast,

Fib2 exhibited an increase in SSc skin (average fraction = 0.224) compared with that in

healthy skin (average fraction = 0.114) ($p$ = 5.7e-05). We also estimated transcription

factor (TF) activities for each subset and observed that Fib2 was characterized by high

TF activity of the hedgehog transcription factor GLI2 (**Supplementary Fib. 5b**). This

convergent driver can mediate Hedgehog pathways in activating fibroblast and promoting

fibrosis in SSc[18,24]. Moreover, Fib9 was marked by *SMAD3*, *SMAD4*, and *JUN*, all of which

are TGF-β downstream pathways to promote fibrosis. To validate our results, we re-

analyzed a bulk RNA-seq dataset from *PRESS* cohort[25] and observed the significant

increase of *POSTN* and decrease of *SCARA5* in SSc skin compared with healthy skin[25].

An inverse correlation was also observed in the gene expressions between *SCARA5* and

*POSTN* (**Supplementary Fig. 5c**). Overall, these findings further supported the potential

involvement of *SCARA5*[+] and *POSTN*[+] fibroblasts in the pathogenesis of SSc.

We next extended our analysis to spatial transcriptomics data. Visualization and

quantification of gene expression showed that *SCARA5* was significantly downregulated

in SSc skin, whereas *POSTN* was upregulated in SSc skin (**Supplementary Fig. 5d**).

Additionally, *SCARA5* has exhibited distinct spatial enrichment patterns that were

mutually exclusive to regions enriched for *POSTN* and *COMP* across multiple samples

(**Fig. 3d; Supplementary Fig. 5e**). To characterize the spatial distribution of fibroblast

subsets, we calculated module scores of each subset, defined as the average expression

of the top 100 marker genes, across spatial spots based on their marker genes. This

analysis revealed that Fib2 and Fib9 were spatially colocalized (**Fig. 3e; Supplementary**

**Fig. 5f**). To probe the regulatory networks underlying these subsets, we estimated the

activity of TFs and signaling pathways, and observed increased activities of JUN, SMAD3,

and the TGF-β pathway, which were closely associated with the spatial localization of Fib2 and Fib9. These findings align with their known pro-fibrotic roles, further confirming the spatial mapping of these sub-clusters (**Fig. 3f-g; Supplementary Fig. 5g-h**). To explore the clinical relevance of these findings, we correlated *SCARA5* and *POSTN* expression with clinical outcomes. Interestingly, *SCARA5* expression negatively correlated with the mRSS, whereas *POSTN* expression positively correlated with mRSS (**Fig. 3h**). Additionally, *SCARA5* showed an inverse correlation with platelet counts, while *POSTN* negatively correlated with non-RP disease duration. These results collectively suggested divergent effects of *SCARA5*⁺ and *POSTN*⁺ fibroblasts in SSc fibrosis.

**Differentiation of *SCARA5*⁺ *to POSTN*⁺ fibroblasts in SSc skin**

While pioneering studies have identified *SCARA5*⁺ fibroblasts as myofibroblast progenitors in fibrosis[19–21], their differentiation trajectories have not been constructed yet. To firstly determine if this phenomenon is consistent across different tissues such as SSc skin fibrosis, we analyzed scRNA-seq data from mouse skin challenged by bleomycin (BLM) at various time points (days 3, 7, 14, and 28) and compared them to untreated skin (day 0, steady state)[26]. This analysis identified 14 distinct cell types and nine fibroblast sub-clusters (**Supplementary Fig. 6a-b**). Among these, we found a progressive decrease in *Scara5*⁺ fibroblasts following BLM treatment, while *Postn*⁺ fibroblasts substantially increased on days 14 and 28 (**Supplementary Fig. 6c-d**). These temporal changes suggested a potential differentiation process from *Scara5*⁺ to *Postn*⁺ fibroblasts.

Based on these observations and the fact that Fib2 and Fib9 were marked by *POSTN*, we next inferred two pseudo-time trajectories from Fib1 to Fib2 (trajectory 1) and Fib1 to Fib9 (trajectory 2) (**Fig3. i-j**). To characterize transcriptional dynamics during the differentiation, we estimated the expression variation of each gene along the pseudo time and identified 1524 and 1515 genes associated with trajectory 1 and 2, respectively. For trajectory 1, we found many genes were down-regulated during the differentiation, including precursor marker *CD34*[27] and *SFRP2*[9] (**Fig. 3k; Supplementary Fig. 6e**). Moreover, we identified a set of genes, including *C3*, *C7*, *CCL2*, and *CCL19*, that were

particularly enriched at the intermediate stage of the trajectory. This suggested a role for chemokine and cytokine signaling, as well as complement system activity, in the differentiation process occurring at this stage. Genes associated with the late stage of this trajectory include *FOS* and *JUNB,* which are involved in the fibrosis-modulatory pathway and have been observed to be activated during the myofibroblast transition process[2]. For trajectory 2, we additionally observed that *TIMP2, TGFBR3* and *APOD* were associated with the intermediate stage. These results revealed distinct gene regulatory networks that are involved in the divergent differentiation of fibroblasts.

Of note, Sarthak Sinha et al conducted comparative single-cell multi-omics in reindeer following full-thickness injury and found that the velvet skin regenerates, while the back skin heals by forming a scar[7]. Similarly, in reindeer skin, *SCARA5* gradually decreased but *ACTA2* increased post wound in their back, but not in velvet (**Supplementary Fig. 7a-e**). This pattern is reminiscent of fibrotic tissue remodeling in human, which indicates the fibroblast dynamic changes could be conserved across species and tissues.

**Spatial co-localization of fibroblast sub-clusters and other cell types**

Apart from highly discordant fibroblast states, the distinct stromal-immune linkages also determine the fibrotic progression. Next, we investigated spatial dependencies between fibroblast subsets and other major cell types to better understand their spatial communications and contributions to fibrotic microenvironments. We conducted spatial colocalization analyses in two distinct proximity areas: 1) co-occurrence within one single spot, determined by Spearman correlation between the module score of each fibroblast sub-cluster and the proportion of each major cell type, and 2) extended neighbourhood within an annular region encompassing up to three surrounding spots for each target spot analyzed with the MISTy pipeline[28]. The results from both approaches were averaged across all samples to identify consistent spatial patterns. We focused specifically on Fib1, Fib2, and Fib9, as they have been demonstrated to exhibit significant alterations in SSc (**Fig. 3c**). Notably, we observed a consistent spatial dependency in both models, characterized by an increased co-occurrence between fibroblast progenitor *SCARA5*+ Fib1 and macrophages, whereas *POSTN*+/*ACTA2*+ myofibroblasts (Fib2 and Fib9) were found in proximity to VSMCs (**Fig. 3l-m**). These findings point towards the potential critical

355   role of macrophage in establishing the spatial microenvironment surrounding fibroblast

356   from the early stage.

**Characterization of macrophage heterogeneity**

Beyond the conventional polarization program of macrophage, new subpopulations have

been implicated in tissue fibrosis[3–5,29]. They are not the direct source of ECM, but rather

a major source of immune dysregulation in fibrotic conditions through release of

proinflammatory and profibrotic factors[30]. Nevertheless, the heterogeneity of

macrophages and their crosstalk with mesenchymal cells in the diseased context of SSc

remain poorly defined. We thus clustered the scRNA-seq data of macrophages and

identified three transcriptionally distinct subsets (**Fig. 4a; Supplemental Fig.8a**). We

observed one cluster highly expressing proinflammatory cytokine *IL-1β*, as well as

*CXCL8*, *CCL3*, *CXCL2* and *PTGS2*, thus termed as proinflammatory macrophages (**Fig.**

**4b**). Recent studies have suggested a role of IL-1β⁺proinflammatory macrophages in

idiopathic pulmonary fibrosis (IPF), and in cardiac tissue remodeling[3,4]. This subset also

highly expresses members of the NR4A subfamily of the nuclear hormone receptor

superfamily including *NR4A3, NR4A2 and NR4A1, which* have been identified as

immediate early genes in response to various stimuli and considered as promising

therapeutic targets due to their significant roles in regulating monocyte differentiation,

macrophage activation and inflammatory response[31] and have been linked to the

pathogenesis of fibrotic tissue remodeling[32].

*Fig. 4 Characterization of macrophage sub-clusters. a, UMAP showing sub-clusters of macrophages based on scRNA-seq data. b, Dotplot showing the marker genes for each sub-cluster. c, Boxplot comparing the proportion of each sub-cluster between healthy and SSc samples. Each dot represents one sample. P-values are calculated using the Wilcoxon rank-sum test (unpaired; two sides). d-e, Number of significant L-R pairs between the fibroblasts and macrophages. f, Heatmap showing significant ligand-receptor pairs between fibroblast subsets and proinflammatory macrophages, with fibroblast serving as signaling senders and macrophage*

We also identified a sub-cluster characterized by phagocytic markers, including *MARCO*,

*LYVE1*, *CD36*, *FCGRT and C1QA*, therefore defined as phagocytic macrophages. The

scavenging receptor CD36, known to mediate uptake in macrophage, has been

implicated in the pathogenesis of tissue fibrosis[33,34]. *MARCO,* a type II glycoprotein

scavenger receptor that binds a broad range of anionic ligands, was found to be

upregulated in tissue-resident macrophages and circulating monocytes upon activation.

*MARCO*[+] myeloid cells have been recognized as therapeutic targets for multiorgan

fibrosis in SSc[35,36]. LYVE1-expressing macrophage has been demonstrated to form

coordinated multi-cellular nest structures that are distributed proximal to vasculature[37].

Another subset of macrophage that may be involved in the antigen presentation process

is characterized by their overexpression of *HLA-B* and *HLA-C,* hereafter referred to as

antigen-presenting macrophages. Indeed, the antigen-presenting macrophage and

phagocytic macrophage defined here largely resembled previously identified

Lyve1[lo]MHCII[hi] and Lyve1[hi]MHCII[lo] macrophages, which are conserved across tissues[8].

Notably, the Lyve1[hi]MHCII[lo] macrophages have been demonstrated to play a critical role

in maintaining homeostasis, as depletion of Lyve1[hi]MHCII[lo] macrophages exacerbated

experimental lung and heart fibrosis, along with exacerbated vessel permeability, immune

infiltration and collagen deposition[8].

The comparison of cell proportion between healthy and SSc samples showed that

proinflammatory macrophages were significantly enriched in SSc ($p$ = 0.0095), whereas

the abundance of phagocytic macrophages was significantly decreased in SSc samples (*p* = 0.027) (**Fig. 4c**). We estimated TF activity for each cluster and observed a high activity of REL, RELA, and RELB in proinflammatory macrophages (**Supplementary Fib. 8b**). These TFs are key members of the NF-κB pathway that regulates immune and inflammatory responses[38], further supporting the functions of proinflammatory macrophages.

**Crosstalk between fibroblast and macrophage sub-clusters**

Our analysis of ST data revealed a strong spatial association between macrophages and fibroblasts, underscoring their potential functional interplay within the tissue microenvironment (**Fig. 2c; Fig. 3l**). To further investigate the cellular crosstalk between specific macrophage and fibroblast subtypes, we performed a detailed cell-cell communication analysis using CellChat[39] in the scRNA-seq data based on ligand-receptor (L-R) interaction networks. We retained only statistically significant L-R pairs between fibroblast and macrophage sub-clusters, resulting in the identification of 393 distinct interactions. These included 246 secreted autocrine/paracrine signaling interactions, 103 ECM-receptor interactions, and 44 direct cell-cell contact-mediated interactions (**Fig. 4d**). This categorization provides a comprehensive view of the diverse modes of communication utilized by these cell types to orchestrate their roles in the tissue microenvironment. Among macrophage subsets, pro-inflammatory macrophages exhibited the highest number of interactions with fibroblast subsets (n = 179 interactions), suggesting that they are central players in fibroblast-mediated signaling (**Fig. 4e**). Particularly, Fib9 emerged as a key fibroblast subtype with the strongest connectivity to macrophages (n = 83 interactions), outpacing other fibroblast subsets in its interaction capacity.

Visualization of the specific L-R interactions revealed that macrophages receive signals from fibroblasts through multiple signaling pathways, including THBS, MIF, CXCL, LAMININ, and COLLAGEN (**Supplementary Fig. 8c**). Among these interactions, the receptor CXCR4 was observed to interact with fibroblast-derived ligands such as MIF and CXCL12. This interaction suggests a mechanism by which fibroblasts recruit and retain proinflammatory macrophages, thereby sustaining local inflammation and contributing to

the progression of fibrosis (**Fig. 4f**). On the other hand, fibroblasts also acted as signal
receivers, engaging the receptor ACKR3 to interact with MIF and CXCL12
(**Supplementary Fig. 8d; Fig. 4g**). Notably, *MIF* and *CXCL12* were expressed by nearly
all subtypes of fibroblasts and macrophages (**Fig. 4h-i**), underscoring their widespread
involvement in cellular crosstalk.

However, the expression of *CXCR4* was specific to proinflammatory macrophages, while
*ACKR3* was uniquely expressed in the Fib1 fibroblast progenitor subcluster (**Fig. 4h**). To
confirm at the protein levels, we applied Tyramide Signal Amplification (TSA) high-plex
immunofluorescence onto adjacent tissue sections. Our TSA staining confirmed the
overlapping expression of ACKR3 in SCARA5$^+$ fibroblast (**Fig. 4j**). Again, we observed
IL-1β$^+$ macrophages, which were also positive for CXCR4, closely distributed adjacent to
fibroblasts (**Fig. 4k**). Together, these results revealed a highly organized and reciprocal
signaling network between fibroblasts and macrophages, underscored by distinct L-R
interactions. Of particular interest, the chemokine CXCL12 can bind to its classic receptor
CXCR4, a G protein-coupled receptor, facilitating the recruitment of CXCR4$^+$ cells, such
as proinflammatory macrophages in our case. Alternatively, CXCL12 can interact with its
atypical chemokine receptor ACKR3 independent of GPCR mediated signaling. ACKR3
is capable of actively internalizing its chemokine ligands CXCL12, through scavenging.
This function helps to modulate the availability and gradient of CXCL12 in the extracellular
environment[40]. The selective expression of ACKR3 in fibroblast progenitors may indicate
an endogenous compensation at an early stage. A dysfunction in this mechanism could
lead to the subsequent CXCR4 mediated recruitment of macrophages, during the
transition from fibroblast progenitor to myofibroblast over the course of SSc progression.

**A data-driven model of SSc**

Taken together, we proposed a data-driven disease model of SSc based on the results
of multi-omics data obtained from SSc, a prototypical systemic fibrotic condition (**Figure
5**). An imbalance in fibroblast subsets in SSc was characterized by a decrease in
*SCARA5*$^+$ fibroblast progenitors and an increase in *POSTN*$^+$ and *ACTA2*$^+$ terminally
differentiated myofibroblasts, which promote the excessive ECM deposition. The

decreased phagocytic macrophages (Lyve1$^{hi}$MHCII$^{lo}$) may contribute to the impairment

of vessel integrity. The infiltration of proinflammatory macrophages (IL-1β$^+$) was observed

within the fibrotic niche, where they were spatially localized in close proximity to fibroblast

subsets. These proinflammatory macrophages were overexpressed with *CXCR4*, which

may mediate their recruitment by interacting with *CXCL12*. Additionally, the expression

of ACKR3 on fibroblasts, particularly in progenitors under homeostatic conditions, may

regulate the local levels CXCL12 by internalizing the ligand, thereby limiting excessive

leukocyte recruitment and preventing unwanted tissue inflammation. However, this

regulatory mechanism might diminish in effectiveness as progenitor fibroblasts decrease

during the progression of SSc, ultimately resulting in uncontrolled CXCL12/CXCR4-

mediated inflammatory cell attraction. This regulatory network fosters a shift from tissue

homeostasis to a fibrogenic state, driving the progression of SSc. Therefore, the

integration of spatial omics with scRNA-seq provides insight into the immune-stromal

crosstalk involved in the pathogenesis of SSc.

***Fig 5. A Data-Driven Model of SSc.*** *Schematic summary illustrates the proposed cellular*

*interplay in SSc skin, which leads to a shift from tissue homeostasis to a fibrogenic environment.*

*The illustration was created by using images from Servier Medical Art (http://smart.servier.com/),*

*licensed under CC BY 3.0 (https://creativecommons.org/licenses/by/3.0/).*

**Discussion**

SSc is a multifaceted autoimmune disease characterized by a well-known triad of

manifestations: vasculopathy, autoimmunity and fibrosis. In this study, we provided a

comprehensive map of the skin ecosystem in SSc using genome-wide spatial

transcriptomes based on 10x Visium and Stereo-seq platforms combined with additional

scRNA-seq datasets. We identified disease-associated fibroblasts and macrophages and

characterized their intricate interplay within niches that potentially drive the transition from

a homeostatic to a fibrogenic state. The molecular signatures of cell populations were

validated through spatial proteomics, including IMC and TSA. The differentiation

trajectory of fibroblasts was cross-validated using time-course bleomycin induced SSc

mouse and wound-healing reindeer models. Further investigation into cell-cell

communications offered potential for identifying targetable disease-driving molecules for

therapeutic intervention.

Our research aimed to explore the potential to halt the disease progression by interrupting

the crosstalk between pathological cells. Spatial transcriptomics provides a unique

advantage in examining the structured compartment and the cellular composition and

states within specific microenvironments. To overcome the limitations of the 55-um

resolution per spot offered by the 10x Visium platform, we employed a higher-resolution

spatial transcriptomics technology, FFPE-based Stereo-seq, achieving a resolution of 0.5

um per spot. To further validate the genome-wide spatial data, we applied additionally

spatial proteomics, including IMC and TSA, which offering the molecular expression and

distribution of the identified cells in SSc skin at single-cell level with the resolutions of 1

$um^2$ and 0.25 $um^2$ per pixel, respectively. Thus, our study aimed to serve as a

comprehensive description of the skin landscape in SSc, revealing potential therapeutic

targets.

We identified distinct niches enriched with fibroblasts and macrophages, showing active

TGF-β signaling pathway and correlating with clinical outcomes in SSc. We then homed

in on distinct cell phenotypes and spatial relationships of fibroblasts and macrophages

within niches. Of note, we discovered a proportional imbalance of fibroblast subsets in

SSc, with a clear reduction in SCARA5[+] progenitor fibroblast and an upregulation in

*POSTN*[+]/*ACTA2*[+] terminally differentiated myofibroblasts compared with healthy controls.

SCARA5 positive fibroblasts have been recently reported as myofibroblast progenitors in

heart and kidney[19,20,41]. A cross-tissue human fibroblast atlas, derived from 517 human

samples encompassing 11 tissue types and diverse pathological states, demonstrated

the substantial overlaps among *SCARA5*[+] fibroblast and progenitor-like fibroblast, as well

as between *POSTN*[+] fibroblast and myofibroblast[42]. Additionally, the dynamic changes of

fibroblasts were not only identified through trajectory analysis in human SSc skin, but also

validated in time-course models of SSc mouse and wound-healing reindeer using the

publicly  available  datasets[7,26].  These  findings  indicated  that  the  differentiation  from

*SCARA5*[+] fibroblast progenitors to myofibroblasts represents a conserved phenotype

across different tissues and species. Targeting the shared mechanism has emerged as

a  strategy  for  basket  trial  design  especially  for  rare  diseases  that  share  the  same

triggering  or  perpetuating  signaling  pathways,  such  as  INBUILD  study[43].  Of  note,  we

revealed  an  intense  interaction  between  IL-1β[+]  proinflammatory  macrophages  and

fibroblast subsets in SSc. Infiltrating macrophages activate fibroblasts by secreting IL-1β,

whilst  systemic  IL-1β  inhibition  or  targeted  IL-1β  deletion  in  macrophages  prevents

fibroblast activation[3,4]. Furthermore, we found this crosstalk was predicted to be mediated

by  the  interaction  of  CXCL12  with  CXCR4  in  macrophages  and  ACKR3  in  fibroblast

progenitors. Indeed, we also found the deregulation of Lyve1[hi]MHCII[lo] macrophages in

SSc,  which  are  known  to  play  a  protective  role  by  maintaining  vascular  integrity  and

preventing immune cell infiltration in fibrotic conditions[8]. These sequential events such as

tissue entry and immune infiltration occuring in SSc may shift the hemostatic condition

towards  a  profibrotic  environment.  Importantly,  the  inability  to  constrain  CXCL12

overexpression in the milieu through internalization by fibroblast progenitors via ACKR3

leads to CXCR4-mediated proinflammatory macrophage recruitment, which is reflective
of the fine-tuned regulation by local fibroblasts.

In summary, spatial multi-omics combined with scRNA-seq helps to elucidate the
entanglement of fibroblasts and macrophages, forming a connectivity network in the
pathogenesis of SSc. Within this network, fibroblasts and macrophages with distinct
phenotypes contribute to tissue fibrotic remodeling in a coordinated manner. We thus
used this strategy to facilitate target discovery, identifying potential intervention axes for
fibrosis.

**METHODS**

**Human skin biopsy collection and processing**
Skin biopsies from 14 SSc subjects and 4 age- and gender-matched healthy controls
were obtained and proceeded for spatial transcriptomics using the 10x Visium platform.
All skin biopsies of SSc patients were taken on the forearm 15 ± 2 cm proximal from the
styloid process of the ulna using disposable biopsy punches (pfm medical, #48301, Gifu,
Japan). The epidemiological and clinical features of the SSc patients are presented in
Supplemental Table 1. After excluding 2 low-quality SSc samples, as shown in
Supplemental Figure 1, we finally included 14 skin biopsies in total, from10 SSc patients
and 4 healthy controls. SSc was diagnosed according to the American College of
Rheumatology (ACR) /European Alliance of Associations for Rheumatology (EULAR)
2013 criteria[44]. In particular, we enrolled SSc patients who had diffused cutaneous
involvement and had experienced their initial non-Raynaud's phenomenon symptoms for
less than three years. Among the SSc patients, 40% were female and 60% were male.
The median age at biopsy of SSc patients was 56.0 years (range 27-73). The median
disease duration from the first non-RP symptoms was 8 months (4-48). The median
mRSS was 21 (8-41). 40% of SSc patients suffered from interstitial lung disease (ILD).
All human studies were approved by the ethical committees of Huashan Hospital of Fudan
University. All patients and controls signed a consent form approved by the local
institutional review board after informed consent.

**Spatially profiling human skin with 10x Visium**

Formalin-fixed paraffin-embedded (FFPE) skin blocks from a total of 18 subjects were
sectioned at 5 um thickness. These sections were transferred to match the 6.5mm$^2$ oligo-
barcoded capture areas on the 10× Visium platform (FFPE v2) according to the
manufacturer's protocol. Prior to this, archived tissues from paraffin-embedded blocks
were tested for RNA quality control (DV200 > 30%), before being mounted on the 10×
CytAssist (6.5 mm × 6.5 mm) for library preparation and sequencing (~50,000 reads/spot).
Hematoxylin and eosin (H&E) staining was conducted for morphological assessment. The
Visium Human Transcriptome Probe Set v2.0 was applied. The final library pool was
sequenced on the Illumina Novaseq 6000 instrument using 150-base-pair paired-end
reads.

**Spatially profiling human skin with Stereo-seq**

To achieve an enhanced resolution and whole transcriptome in-situ analysis of skin
samples, we employed Stereo-seq technology according to the previous protocols[16,45],
which enables the spatial detection of transcriptome-wide gene expression at a resolution
of 0.5 um. Briefly, 5-um FFPE skin sections from 3 SSc patients and 1 healthy control,
which are overlapping with the ones who underwent the examination of 10x Visium, were
subjected to adhere to the Stereo-seq chips which have DNA nanoball (DNB) bins with
220 nm diameter and a center-to-center distance of 500 nm. After methanol fixation, the
formal sections were preceded with ssDNA staining and the adjacent sections with
hematoxylin and eosin (H&E) staining for morphological assessment. Next,
permeabilization of the tissue was performed to release mRNA. cDNA products are
synthesized by reversed transcription. After amplification, DNBs were then generated via
rolling cycle amplification (RCA) and loaded onto the patterned chips (65mm×65mm).
Next, to determine the distinct DNB-CID sequences at each spatial location, paired-end
sequencing was performed on a DNBSEQ-T7 sequencer (MGI Research, Shenzhen,
China) with a PE75 sequencing strategy.

**Processing 10x Visium and Stereo-seq data**

For 10x Visium data, we used SpaceRanger software (v2.0.0) to pre-process the raw
sequencing data based on the reference genome hg38. For each sample, we filtered the
spots by number of detected genes (< 100). For Stereo-seq data, we leveraged nucleic
acid staining from the same section to segment cells by projecting the staining image to
the Stereo-seq chips. The watershed algorithm was applied to the staining image for
single cell segmentations. For each of the segmented cells, UMI from all DNB within the
corresponding segmentation were aggregated per-gene and then summed to generate a
cell by gene matrix for downstream analysis. To overcome the low RNA capture
efficiencies on single DNB spots at a resolution of 500nm, the raw spatial expression
matrix was convoluted into larger pseudo-spots with a 50×50 window size (bin50 for
short), or more precisely, as 25µm squares. We then filtered the spots by number of
detected genes (< 50) and mitochondrial genes (> 10%).

**Analysis of publicly available scRNA-seq data**

To build an atlas of human skin cells, we downloaded scRNA-seq data from Gene
Expression Omnibus (GEO) with accession numbers GSE195452 (denoted by
Gur2022)[10] and GSE138669 (denoted by Tabib2021)[9]. For the Gur2022 dataset, we only
kept the cells from the skin. We further filtered the cells by number of UMIs (< 500),
number of detected genes (< 300 or > 5000), and mitochondrial genes (> 30%). We also
filtered the genes by the number of cells (n = 100). We re-annotated the cells by merging
the 49 original labels to 16 major cell types as follows: Fibroblast (Fibro_ATAC2,
Fibro_Bad, Fibro_COCH, Fibro_COMP, Fibro_IGFBP2, Fibro_LGR5, Fibro_MYOC1,
Fibro_MYOC2, Fibro_POSTN, Fibro_PTGDS, Fibro_POSTN_PTGDS), T cells
(sT_Effector, T_Effector, T_GD, T_Naive, sTreg_CXCR4, sTreg, Treg, sT), NK cells (NK,
NK_XCL1, NK_XCL1_CXCR4), cDC (DC, DC_CCL22, DC_CXCL10, DC_XCR1),
Macrophages (Mf, Mf_TREM2), Monocytes (Mo_CD16, M_CD16_IL1B, M_IL1B, Mo),
Mast cells (bMast_CLC, Mast), B cells (B, B_CXCR4), Pericytes (Peri_RGS5,
Peri_TGFBI), Endothelial cells (Vascular_ACKR1, Vascular_RBP7), Keratinocytes
(KRT1_KRT10, KRT14_ACTA2, KRT14_S100A2_GJA1). For the Tabib2021 dataset, we
filtered the cells by UMIs (<500), number of detected genes (<300 and >5000), and
mitochondrial gene expression (> 5%). Since the original annotation was unavailable, we

clustered and re-annotated the cells. This recovered 13 major cell types, among which

10 cell types were also identified by the Gur2022 dataset. To integrate these two datasets,

we merged the count matrices and normalized the data using Scanpy[46]. Next, we used

the Harmony algorithm[47] to correct the batch effects between the datasets and visualize

the integrated cells by UMAP.

## Cell type deconvolution of 10x Visium data

We used cell2location[48] and the above scRNA-seq data as a reference to estimate cell

type composition for each spot in the spatial transcriptomics samples. Specifically, we

first identified the signatures of each cell type using the scRNA-seq data based on a

Negative binomial regression model after accounting for the batch effects from different

datasets. Next, we removed all mitochondrial genes, subsetting the reference data and

spatial data using the identified signatures. We ran the function Cell2location

(N_cells_per_location=5; detection_alpha=20) to estimate posterior distributions of cell

abundance. We normalized the estimated cell type composition for each spot to [0, 1],

which was used for visualization and downstream analysis.

## Inference of pathway activity for 10x Visium data

We used decoupeR[49] (v1.8.0) to infer the pathway and transcription factor activity.

Specifically, for each sample, we normalized the expression counts matrix and identified

highly variable genes using Scanpy[46]. Next, we obtained a collection of pathways and

their target genes using the function get_progeny (organism='human', top=500). The

pathway enrichment scores were inferred using the multivariate linear model by running

the function run_mlm.

## Integration of scRNA-seq and Stereo-seq data

We used Seurat (v5.1.0) to integrate the scRNA-seq and Stereo-seq data. Specifically,

we merged the Stereo-seq samples and then identified a set of anchors between these

two datasets using the function FindTransferAnchors (reduction = "cca") based on the

shared genes (n = 11110). Next, we transferred the cell type labels from scRNA-seq to

Stereo-seq data using the function TransferData (weight.reduction = "cca", dims = 1:30).

To integrate the data in a common embedding space, we also transferred the batch-

corrected PCA components (n = 30) from scRNA-seq to Stereo-seq data with the function

TransferData. Next, we run the function RunUMAP to generate a co-embedding space to

visualize integrated scRNA-seq and Stereo-seq data.

## Identification of cell-type composition niches in 10x Visium data

We combined the cell type proportions estimated by cell2location from all samples to.

Next, we ran the function neighbors (n_neighbors=30) to create a neighborhood graph of

all spatial spots based on the cell type proportions. An UMAP embedding was generated

using the function umap with default settings in Scanpy. We clustered the spots with the

function leiden (resolution = 0.5), identifying 12 niches. To annotate the niches, we

computed the average proportion for each cell type within each niche. We next

normalized the values by z-score transformation and clustered the niches using

hierarchical approach.

## Sub-clustering of fibroblasts

We subsetted the fibroblasts and removed the samples with less than 100 cells to allow

for robust data integration. Next, we clustered the cells with the Leiden algorithm[50]

(resolution =1) based on the harmonized principal components. We removed one

population with high mitochondrial genes, resulting in a total of ten clusters identified. The

marker genes for each cluster were identified using the function rank_genes_groups

(method='wilcoxon') from Scanpy. To map these sub-clusters to spatial space, we

selected the top 100 markers for each cluster and then computed module scores using

the function AddModuleScore from Seurat[51].

## Analysis of scRNA-seq data from mouse skin

We downloaded scRNA-seq data from the Zenodo data archive
(https://zenodo.org/records/7455989). Cells were filtered by the number of UMIs (< 2000
or > 40000), number of detected genes (< 200 or > 6000) and fraction of mitochondrial
genes (> 5%), retaining 18503 cells. Genes were filtered by the number of cells (< 50).
Next, we normalized the count matrix using the functions normalize_total and log1p from
Scanpy. Highly variable genes were selected by the function highly_variable_genes. We
clustered the cells with the Leiden algorithm (resolution = 0.5) based on the 30 principle
components and identified 25 populations which were further grouped into 14 cell types
based on their canonical markers. We next subsetted all fibroblasts and clustered the
cells using the Leiden algorithm (resolution=0.5), obtaining 10 sub-clusters. We noticed
one cluster showing a low number of UMIs and detected genes, presumably
corresponding to low-quality cells. We therefore removed this cluster and rename the
remaining clusters as Fib1-9.

**Analysis of scRNA-seq data from reindeer**
The scRNA-seq data of  full-thickness injuries in skin from reindeer was analyzed with the
interface website (www.biernaskielab.ca/reindeer_atlas)[7].

**Trajectory analysis of fibroblasts**

We used ArchR[52] to perform trajectory analysis. Specifically, we inferred two pseudo-time
trajectories using the function AddTrajectory to characterize the differentiation processed
from Fib1 to Fib2 (denoted by trajectory 1), and from Fib1 to Fib9 (denoted trajectory 2)
based on the principle components (n = 30). The trajectory 1 involved six subsets (Fib1,
Fib3, Fib5, Fib4, Fib10, and Fib2), and the trajectory 2 included only Fib1 and Fib9. The
trajectories were visualized by the function TrajectoryPlot. To identify associated genes
for each trajectory, we obtained the expression matrix along the trajectory using the
function GetTrajectory (smoothWindow = 11). Next, we visualized the dynamics of gene
expression with the function plotTrajectoryHeatmap and selected the top 10% highly
variable genes across the trajectory.

**Spatial relationship analysis between fibroblast subsets and other cell types**

We used two approaches to infer spatial relationships between the subsets of fibroblasts and other major cell types. The first approach was based on Spearman correlation between the module score of each subset and the proportion of each major cell type across all spatial spots. This was performed for each 10x Visium sample independently, and results were averaged across all samples. The second approach was based on Misty. Specifically, we usd Misty to build a predictive model using the cell type proportions as features and module scores as target. This allows us to learn the spatial interactions between spots.

**Sub-clustering of macrophages**

We subsetted all macrophages from the integrated scRNA-seq atlas and removed samples with less than 50 cells. Next, clustered the cells using the Leiden algorithm[50] (resolution = 0.3) based on the harmonized principal components, identifying three sub-clusters. The marker genes for each cluster were identified using the function rank_genes_groups (method='wilcoxon') from Scanpy. To map these sub-clusters to spatial space, we selected the top 100 markers for each cluster and then computed module scores using the function AddModuleScore from Seurat[51].

**Cell-cell communication analysis between fibroblasts and macrophages**

We merged the fibroblast and macrophage subclusters to create a Seurat object. Next, we normalized the count matrix using the function NormalizeData from Seurat and identified highly variable genes using the function FindVariableFeatures. We created a CellChat object and grouped the cells based on the sub-clustering results as described above. We selected the database CellChatDB.human and excluded ligand-receptor pairs from non-protein signaling. The cell-cell communication network was inferred using the function computeCommunProb (type = "triMean") and was filtered using the function filterCommunication (min.cells = 10).

**Imaging mass cytometry (IMC)**

We performed data mining using our published IMC dataset[15], which included FFPE skin

biopsy sections from 19 SSc patients and 14 matched controls. The IMC panel included

up to 38 lanthanide labeled antibodies as well as Iridium 193 and 191 isotopes, which

were designed to target epitopes specific for different subsets of endothelial cells,

endothelial precursor cells, immune cells and mesenchymal cells, as well as structural

markers used for tissue segmentation. Data acquisition was performed on a Helios time-

of-flight mass cytometer coupled to a Hyperion Imaging System (Standard Biotools). After

single-cell segmentation and data transformation and normalization, we obtained the

single-cell dataset with spatial information alongside. We then proceeded with

dimensionality reduction, unsupervised clustering, and subsequent cell type annotation.

To investigate the local microenvironment of individual fibroblast subpopulations, we

identified cellular neighborhoods (CNs) by k-means clustering on the composition of

neighboring cells as described[14].

**Tyramide signal amplification (TSA)**

FFPE skin tissue sections obtained from 4 SSc patients and 4 control donors were stained

using a TSA system as previously described[15,53]. Briefly, the sections were proceeded

with dewaxing, rehydration, and antigen retrieval with Tris-EDTA(pH 9.0) for 15 minutes.

Endogenous peroxidase activity was neutralized with 3% $H_2O_2$ for 10 minutes, followed

by a 30-minute blocking step with antibody diluent/block buffer (ARD1001EA, AKOYA).

Next, sections were incubated for 90 min at room temperature with the following primary

antibodies of the indicated dilutions: SCARA5 (ab118894, 1:200, Abcam), CD90

(ab226123, 1:200, Abcam), α-SMA (#19245, 1:200, Cell signaling), CD68 (ab192847,

1:200, Abcam), IL-1β (ab283818, 1:200, Abcam), CXCR4 (ab181020, 1:200, Abcam) and

ACKR3 (20423-1-AP, 1:200, Proteintech). After multiple rounds of signal amplification

with Opal fluorophores and subsequent antibody stripping, sections were stained with

DAPI. Slides were scanned using a PhenoImager Fusion scanner (Akoya Biosciences)

to distinguish the spectra of the different fluorescence channels. All images were captured

and presented using the same equipment setting. Merging channels was conducted using

ImageJ / Fiji software (Release 2.16.0, https://fiji.sc).

**References**

1. Elhai, M. *et al.* Mapping and predicting mortality from systemic sclerosis. *Ann.*

*Rheum. Dis.* **76**, 1897–1905 (2017).

2. Distler, J. H. W. *et al.* Shared and distinct mechanisms of fibrosis. *Nat. Rev.*

*Rheumatol.* **15**, 705–730 (2019).

3. Alexanian, M. *et al.* Chromatin remodelling drives immune cell-fibroblast

communication in heart failure. *Nature* **635**, 434–443 (2024).

4. Amrute, J. M. *et al.* Targeting immune-fibroblast cell communication in heart failure.

*Nature* **635**, 423–433 (2024).

5. Mayr, C. H. *et al.* Spatial transcriptomic characterization of pathologic niches in IPF.

*Sci. Adv.* **10**, eadl5473 (2024).

6. Morse, C. *et al.* Proliferating SPP1/MERTK-expressing macrophages in idiopathic

pulmonary fibrosis. *Eur. Respir. J.* **54**, 1802441 (2019).

7. Sinha, S. *et al.* Fibroblast inflammatory priming determines regenerative versus

fibrotic skin repair in reindeer. *Cell* **185**, 4717–4736.e25 (2022).

8. Chakarov, S. *et al.* Two distinct interstitial macrophage populations coexist across

tissues in specific subtissular niches. *Science* **363**, eaau0964 (2019).

9. Tabib, T. *et al.* Myofibroblast transcriptome indicates SFRP2hi fibroblast progenitors

in systemic sclerosis skin. *Nat. Commun.* **12**, 4384 (2021).

10. Gur, C. *et al.* LGR5 expressing skin fibroblasts define a major cellular hub perturbed

in scleroderma. *Cell* **185**, 1373–1388.e20 (2022).

11. Clark, K. E. N. *et al.* Single-cell analysis reveals key differences between early-stage

and late-stage systemic sclerosis skin across autoantibody subgroups. *Ann. Rheum.*

*Dis.* **82**, 1568–1579 (2023).

12. Yin, H. *et al.* Endothelial response to type I interferon contributes to vasculopathy

and fibrosis and predicts disease progression of systemic sclerosis. *Arthritis*

*Rheumatol.* **76**, 78–91 (2024).

13. Ma, F. *et al.* Systems-based identification of the Hippo pathway for promoting fibrotic
mesenchymal differentiation in systemic sclerosis. *Nat. Commun.* **15**, 210 (2024).

14. Rius Rigau, A. *et al.* Imaging mass cytometry-based characterisation of fibroblast
subsets and their cellular niches in systemic sclerosis. *Ann. Rheum. Dis.* (2024)
doi:10.1136/ard-2024-226336.

15. Rius Rigau, A. *et al.* Characterization of vascular niche in systemic sclerosis by
spatial proteomics. *Circ. Res.* **134**, 875–891 (2024).

16. Chen, A. *et al.* Spatiotemporal transcriptomic atlas of mouse organogenesis using
DNA nanoball-patterned arrays. *Cell* **185**, 1777–1792.e21 (2022).

17. Avouac, J. *et al.* Inhibition of activator protein 1 signaling abrogates transforming
growth factor β-mediated activation of fibroblasts and prevents experimental fibrosis.
*Arthritis Rheum.* **64**, 1642–1652 (2012).

18. Bergmann, C. *et al.* Mutual amplification of GLI2/Hedgehog and cJUN/AP1 signaling
in fibroblasts in Systemic Sclerosis (SSc) - potential implications for combined
therapies. *Arthritis Rheumatol.* (2024) doi:10.1002/art.42979.

19. Kuppe, C. *et al.* Decoding myofibroblast origins in human kidney fibrosis. *Nature* **589**,
281–286 (2021).

20. Kuppe, C. *et al.* Spatial multi-omic map of human myocardial infarction. *Nature* **608**,
766–777 (2022).

21. Li, Z. *et al.* Chromatin-accessibility estimation from single-cell ATAC-seq data with
scOpen. *Nat. Commun.* **12**, 6386 (2021).

22. Nanri, Y. *et al.* Cross-talk between transforming growth factor-β and periostin can be
targeted for pulmonary fibrosis. *Am. J. Respir. Cell Mol. Biol.* **62**, 204–216 (2020).

23. Nanri, Y. *et al.* A positive loop formed by SOX11 and periostin upregulates TGF-β
signals leading to skin fibrosis. *J. Invest. Dermatol.* **143**, 989–998.e7 (2023).

24. Liang, R. *et al.* The transcription factor GLI2 as a downstream mediator of
transforming growth factor-β-induced fibroblast activation in SSc. *Ann. Rheum. Dis.*
**76**, 756–764 (2017).

25. Skaug, B. *et al.* Global skin gene expression analysis of early diffuse cutaneous
systemic sclerosis shows a prominent innate and adaptive inflammatory profile. *Ann.*
*Rheum. Dis.* **79**, 379–386 (2020).

26. Liang, Y. *et al.* Dynamic pathological analysis reveals a protective role against skin
fibrosis for TREM2-dependent macrophages. *Theranostics* **14**, 2232–2245 (2024).

27. Nazari, B. *et al.* Altered dermal fibroblasts in systemic sclerosis display podoplanin
and CD90. *Am. J. Pathol.* **186**, 2650–2664 (2016).

28. Tanevski, J., Flores, R. O. R., Gabor, A., Schapiro, D. & Saez-Rodriguez, J.
Explainable multiview framework for dissecting spatial relationships from highly
multiplexed data. *Genome Biol.* **23**, 97 (2022).

29. Buechler, M. B., Fu, W. & Turley, S. J. Fibroblast-macrophage reciprocal interactions
in health, fibrosis, and cancer. *Immunity* **54**, 903–915 (2021).

30. Bhandari, R. *et al.* Profibrotic activation of human macrophages in systemic sclerosis.
*Arthritis Rheumatol.* **72**, 1160–1169 (2020).

31. Boulet, S., Le Corre, L., Odagiu, L. & Labrecque, N. Role of NR4A family members
in myeloid cells and leukemia. *Curr. Res. Immunol.* **3**, 23–36 (2022).

32. Palumbo-Zerr, K. *et al.* Orphan nuclear receptor NR4A1 regulates transforming
growth factor-β signaling and fibrosis. *Nat. Med.* **21**, 150–158 (2015).

33. Kwak, D. *et al.* CD36/Lyn kinase interactions within macrophages promotes
pulmonary fibrosis in response to oxidized phospholipid. *Respir. Res.* **24**, 314 (2023).

34. Pennathur, S. *et al.* The macrophage phagocytic receptor CD36 promotes fibrogenic
pathways on removal of apoptotic cells during chronic kidney injury. *Am. J. Pathol.*
**185**, 2232–2245 (2015).

35. Papazoglou, A. *et al.* Epigenetic regulation of profibrotic macrophages in systemic
sclerosis-associated interstitial lung disease. *Arthritis Rheumatol.* **74**, 2003–2014
(2022).

36. Xu, D. *et al.* PLG nanoparticles target fibroblasts and MARCO+ monocytes to reverse
multiorgan fibrosis. *JCI Insight* **7**, (2022).

37. Anstee, J. E. *et al.* LYVE-1+ macrophages form a collaborative CCR5-dependent
perivascular niche that influences chemotherapy responses in murine breast cancer.
*Dev. Cell* **58**, 1548–1561.e10 (2023).

38. Liu, T., Zhang, L., Joo, D. & Sun, S.-C. NF-κB signaling in inflammation. *Signal
Transduct. Target. Ther.* **2**, (2017).

39. Jin, S., Plikus, M. V. & Nie, Q. CellChat for systematic analysis of cell-cell

communication from single-cell transcriptomics. *Nat. Protoc.* (2024)
doi:10.1038/s41596-024-01045-4.

40. Cuesta-Margolles, G., Schlecht-Louf, G. & Bachelerie, F. ACKR3 in skin
homeostasis, an overlooked player in the CXCR4/CXCL12 axis. *J. Invest. Dermatol.*
(2024) doi:10.1016/j.jid.2024.08.022.

41. Schreibing, F., Anslinger, T. M. & Kramann, R. Fibrosis in pathology of heart and
kidney: From deep RNA-sequencing to novel molecular targets. *Circ. Res.* **132**,
1013–1033 (2023).

42. Gao, Y. *et al.* Cross-tissue human fibroblast atlas reveals myofibroblast subtypes
with distinct roles in immune modulation. *Cancer Cell* **42**, 1764–1783.e10 (2024).

43. Flaherty, K. R. *et al.* Nintedanib in progressive fibrosing interstitial lung diseases. *N.*
*Engl. J. Med.* **381**, 1718–1727 (2019).

44. van den Hoogen, F. *et al.* 2013 classification criteria for systemic sclerosis: an
American College of Rheumatology/European League against Rheumatism
collaborative initiative: ACR/EULAR classification criteria for SSc. *Arthritis Rheum.*
**65**, 2737–2747 (2013).

45. Chen, A. *et al.* Single-cell spatial transcriptome reveals cell-type organization in the
macaque cortex. *Cell* **186**, 3726–3743.e24 (2023).

46. Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene
expression data analysis. *Genome Biol.* **19**, 15 (2018).

47. Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with
Harmony. *Nat. Methods* **16**, 1289–1296 (2019).

48. Kleshchevnikov, V. *et al.* Cell2location maps fine-grained cell types in spatial
transcriptomics. *Nat. Biotechnol.* **40**, 661–671 (2022).

49. Badia-I-Mompel, P. *et al.* decoupleR: ensemble of computational methods to infer
biological activities from omics data. *Bioinform Adv* **2**, vbac016 (2022).

50. Traag, V. A., Waltman, L. & van Eck, N. J. From Louvain to Leiden: guaranteeing
well-connected communities. *Sci. Rep.* **9**, 5233 (2019).

51. Hao, Y. *et al.* Dictionary learning for integrative, multimodal and scalable single-cell
analysis. *Nat. Biotechnol.* (2023) doi:10.1038/s41587-023-01767-y.

52. Granja, J. M. *et al.* ArchR is a scalable software package for integrative single-cell

chromatin accessibility analysis. *Nat. Genet.* **53**, 403–411 (2021).

53. Liang, M. *et al.* Identification and validation of anti-protein arginine methyltransferase

5 (PRMT5) antibody as a novel biomarker for systemic sclerosis (SSc). *Ann. Rheum.*

*Dis.* **83**, 1144–1155 (2024).

**Data Availability**

The scRNA-seq data for human skin were downloaded from GEO with the accession number GSE195452 (Gur2022) and GSE138669 (Tabib2021). A cross-tissue human fibroblast atlas was analyzed with the interface website http://pan-fib.cancer-pku.cn/ (Gao2024). The scRNA-seq data of full-thickness injuries in skin from reindeer was analyzed with the interface website www.biernaskielab.ca/reindeer_atlas (Sinha2022). The scRNA-seq data for mouse skin were downloaded from the Zenodo data archive (https://zenodo.org/records/7455989).

**Code Availability**

The codes used in this study are available on Github: https://github.com/lzj1769/Spatial_SSc_project.

**Acknowledgment**

This study was supported by the Natural Science Foundation of China (NSFC) (82371818) to ML, Joint Sino-German research project from NSFC and Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) (82161138022) to JHWD, HZ and ML, Huashan Cultivation research (30302164001) to ML, Huashan Innovation project (2024CX01) to ML, Shanghai Oriental Youth Talent program ( DFYCQN07) to ML, Rappaport MGH Research Scholar Award to LP.

**Competing interests**

Although no disclosures directly relate to the present study, JHWD has consultancy

relationships with Active Biotech, Anamar, ARXX, AstraZeneca, Bayer Pharma,
Boehringer Ingelheim, Callidatas, Celgene, Galapagos, GSK, Inventiva, Janssen,
Kyverna, Novartis, Pfizer, Quell Therapeutics and UCB, has received research funding
from Anamar, ARXX, BMS, Bayer Pharma, Boehringer Ingelheim, Cantargia, Celgene,
CSL Behring, Exo Therapeutics, Galapagos, GSK, Incyte, Inventiva, Kiniksa, Kyverna,
Lassen Therapeutics, Mestag, Sanofi-Aventis, RedX, UCB and ZenasBio and is CEO of
4D Science and scientific lead of FibroCure. LP has financial interests in Edilytics, Inc.,
and SeQure Dx, Inc.

**Author contributions statement**

ML and ZL designed the study. ZL performed all data analysis presented in this work. ZL,
ML, ARR, WX, XS, YNL and AEM were involved in acquisition and analysis of data. ZL,
ML, RH, LP and JHWD were involved in interpretation of data. ML provided essential
samples and resources. All authors were involved in manuscript preparation and proof-
reading.

970 **Supplementary Figures**

**Supplementary Fig. 1 Quality check of spatial transcriptomics data. a,** *Left: number of detected spots per sample in 10x Visium data. Middle: number of genes per spot in each sample. Right: number of UMIs in each sample.* **b,** *Left: number of detected spots per sample in Stereo-seq. Middle: number of genes per sample. Right: number of UMIs in each sample.* **c,** *Visualization*

*of H&E image and number of UMIs in spatial space for 10x Visium data. **d,** Visualization of*
*number of UMIs in spatial space for Stereo-seq data.*

**Supplementary Fig. 2 Analysis of publicly available scRNA-seq data. *a,*** *UMAP showing*
*scRNA-seq from the Tabib2021 dataset as colored by cell types and conditions. **b,** Heatmap*

*showing cell-type-specific marker genes. **b,** UMAP showing scRNA-seq as colored by cell types*

*and conditions. **c,** Same as **a** for the Gur2022 dataset.  **d,** Same as **b** for the Gur2022 dataset.*

*e, UMAP showing the integrated scRNA-seq data colored by different datasets. **f,** Visualization of*

*cell-type-specific marker genes. **g,** Barplot showing number of cells for each cell type.*

***Supplementary Fig. 3 Characterization of cell-type niches in human skin. a-b,** UMAP*

*embedding of the integrated spatial data as colored by samples, conditions, and the fraction of*

*each cell type**. c,** Marker genes of each cell-type niche. **d,** Visualization of cell-type niches in*

*spatial space. **e,** KEGG pathway enrichment analysis for each cell-type niche.*

**Supplementary Fig. 4 Validation of cellular neighbourhood (CN) by imaging mass cytometry (IMC) in healthy and SSc skin. a**, *Study design of IMC, as a validation spatial approach. We collected skin biopsies from 19 SSc patients and 14 healthy controls as we previously described[15].* **b,** *Heatmap showing the enrichment of cell types in CNs. Each column represents a cell type, and each row represents a CN. Colors refer to the proportions of each cell type across different CNs.* **c,** *Left: cellular neighborhood (CN) community detected using IMC. Colors refer to cellular clusters. Right: CN5 and CN8 are highlighted and colored by cell types.*

1001

1002

***Supplementary Fig. 5 Sub-clustering of fibroblasts. a,*** *UMAP embedding of the integrated*

*scRNA-seq fibroblasts as colored by datasets and conditions.* ***b,*** *Heamap of top four marker TFs*

*for each sub-cluster-based TF activity score.* ***c,*** *Left: barplot comparing the expression of*

*SCARA5 and POSTN between healthy and SSc conditions using bulk RNA-seq. Right: scatter*

*plot showing the Spearman correlation between SCARA5 and POSTN across patients.* ***d,*** *Left:*

*barplot comparing the expression of SCARA5 and POSTN between healthy and SSc conditions*

*using Visium data. Right: scatter plot showing the Spearman correlation between SCARA5 and*

*POSTN across Visium samples.* ***e-h,*** *Visualization of marker genes, module scores, TF and*

*pathway activities in spatial space.*

1012

1013

1014

1015

***Supplementary Fig. 6 Trajectory analysis of fibroblast differentiation. a,*** *UMAP of scRNA-*

*seq data from mouse skin[26] colored by time points (left) after bleomycin (BLM) treatment and*

*annotated cell types (right).* ***b,*** *UMAP of fibroblasts colored by time points (left) and sub-clusters*

*(right) and time points (right).* ***c,*** *Dot plots of marker genes for each sub-cluster. The colors refer*

*to the mean expression of the gene, and the size represents the fraction of cells that expressed*

*the gene.* ***d,*** *Box plot showing the fraction of each sub-cluster within different time points.* ***e,*** *Line*

*plot showing expression dynamics of selected genes along the differentiation trajectory from Fib1*

*to Fib2.* ***f,*** *Line plot showing expression dynamics of selected genes along the differentiation*

*trajectory from Fib1 to Fib9.*

1025

1026

1027

1028

1029

Reindeer fibroblast scRNA-seq data from Sarthak Sinha 2022

1030

1031 ***Supplementary Fig. 7 Cross-Species Validation of Fibroblast Dynamics.*** *Sarthak Sinha et al.*

1032 *utilized single-cell multi-omics to investigate full-thickness skin injuries in reindeer, revealing that*

1033 *their antler skin (velvet) regenerates, while their back skin forms fibrotic scars[7]. **a**, UMAP of*

*scRNA-seq data from reindeer skin colored by cell type. **b,c,** Visualization of SCARA5 (b) and*

1035 *ACTA2 (c) expression in UMAP, showing the gene expression on the back (left) and on the velvet*

1036 *(right). **d,e,** Line plot showing expression dynamics of SCARA5 and ACTA2 at the indicated days*

1037 *post wound in reindeer.*

1038

1039

1040 **Supplementary Fig. 8 Sub-clustering of macrophages. a,** *UMAP showing scRNA-seq of*

1041 *macrophages colored different datasets (left) and conditions (right). **b,** Heamap of top four marker*

1042 *TFs for each sub-cluster-based TF activity score. **c,** Heatmap showing all significant L-R pairs*

1043 *from fibroblasts to macrophages. **d,** Heatmap showing all significant L-R pairs from macrophages*

1044 *to fibroblasts.*