Single-cell RNA-seq reveals trans-sialidase-like superfamily gene expression heterogeneity
in *Trypanosoma cruzi* populations.

Lucas Inchausti[1,2], Lucía Bilbao[1,2], Vanina A. Campo[3,4], Joaquín Garat[5], José Sotelo-Silveira[5,6],
Gabriel Rinaldi[7], Virginia M. Howick[8], María Ana Duhagon[2], Javier G. De Gaudenzi[3,4], Pablo
Smircich[1,2]

1 - Laboratorio de Bioinformática, Departamento de Genómica, Instituto de Investigaciones
Biológicas Clemente Estable, Montevideo, Uruguay

2 - Sección Genómica Funcional, Facultad de Ciencias, Universidad de la República,
Montevideo, Uruguay

3 - Instituto de Investigaciones Biotecnológicas, Universidad Nacional de San Martín—Consejo
Nacional de Investigaciones Científicas y Técnicas, General San Martín, Prov. de Buenos Aires,
Argentina

4 - Escuela de Bio y Nanotecnologías (EByN), Universidad Nacional de San Martín, General
San Martín, Prov. de Buenos Aires, Argentina

5 - Departamento de Genómica, Instituto de Investigaciones Biológicas Clemente Estable,
Montevideo, Uruguay

6 - Sección Biología Celular, Facultad de Ciencias, Universidad de la República, Montevideo,
Uruguay

7 - Department of Biology, University of Oxford, Oxford, United Kingdom

8 - Institute of Biodiversity, Animal Health, and Comparative Medicine, University of Glasgow,
Glasgow, United Kingdom

Abstract

Trypanosoma cruzi, the causative agent of Chagas disease, presents a major public health
challenge in Central and South America, affecting approximately 8 million people and placing
millions more at risk. The *T. cruzi* life cycle includes transitions between epimastigote,
metacyclic trypomastigote, amastigote, and blood trypomastigote stages, each marked by distinct
morphological and molecular adaptations to different hosts and environments. Unlike other
trypanosomatids, *T. cruzi* does not employ antigenic variation but instead relies on a diverse
array of cell-surface-associated proteins encoded by large multi-copy gene families (multigene
families), essential for infectivity and immune evasion.

This study analyzes cell-specific transcriptomes using single-cell RNA sequencing of amastigote
and trypomastigote cells to characterize stage-specific surface protein expression during

mammalian infection. Through clustering and identification of cell-specific markers, we assigned cells to distinct parasite developmental forms. Analysis of individual cells revealed that surface protein-coding genes, especially members of the trans-sialidase TcS superfamily (TcS), are expressed with greater heterogeneity than single-copy genes. Additionally, no recurrent combinations of TcS genes were observed between individual cells in the population. Our findings thus reveal transcriptomic heterogeneity within trypomastigote populations where each cell displays unique TcS expression profiles. Focusing on the diversity of surface protein expression, this research aims to deepen our understanding of *T. cruzi* cellular biology and infection strategies.

Introduction

The protozoan parasite *Trypanosoma cruzi* is the etiological agent of Chagas disease, a highly prevalent infectious disease in Central and South America that affects ~8 million people, with several million more at risk of infection [1].

The life cycle of the parasite involves both an invertebrate vector (triatomine bug) and a vertebrate host. The change in environmental conditions triggers differentiation processes in the parasite developing across four main stages. The epimastigote form replicates within the insect and differentiates into metacyclic trypomastigotes in the insect's rectal tract. This latter form does not replicate, specializes in host infection, and can invade various cell types. Once inside the cell, the parasite differentiates into an amastigote, the replicative form within the mammalian host. After several rounds of division, amastigotes differentiate into bloodstream trypomastigotes, which, after cell lysis, can either infect new cells or be ingested by the vector [2]. Parasite developmental stages exhibit significant morphological and molecular differences. *T. cruzi* infection relies on a heterogeneous set of membrane proteins, encoded mainly by large multigene families [3]. Among these are trans-sialidases and trans-sialidase-like proteins (TcS), Mucins, MASPs, GP63, DGFs, and RHS proteins, most of which are involved in infection, tropism, and immune evasion [4].

The trans-sialidase (TcS) superfamily is involved in processes underlying host-parasite interactions [5]. Members that retain enzymatic activity catalyze the transfer of sialyl groups from host glycoconjugates to galactopyranosyl units on the parasite's surface, an essential activity, as *T. cruzi* is unable to synthesize *de novo* sialic acid [6]. This is the largest superfamily, with over 1,400 members, and is subdivided into eight groups based on amino acid sequence [5]. Few members have been functionally characterized, most expressed primarily in mammalian stages [7]. The TcS group I include proteins that retain TS activity, though members from all groups are involved in host-parasite interactions [8].

In recent years, single-cell RNA sequencing (scRNA-seq) has been employed in protozoa, with reports including *Plasmodium falciparum*, *Trypanosoma brucei,* and *Leishmania major* [9-18].

These studies revealed key aspects of the infection process undetectable with conventional
methods, highlighting the relevance of this approach for understanding individual variation in
gene expression in single-celled organisms [19]. In *T. brucei,* antigenic variation driven by
variant surface glycoproteins (VSGs) has been studied at single-cell resolution to understand the
mechanisms that enable subpopulations of this parasite to evade the immune system, as a bet
hedging strategy, ensuring parasite survival [14]. In this parasite, scRNA-seq revealed that pre-
metacyclic cells express multiple VSG transcripts simultaneously, in contrast to metacyclic
forms, which display a protein coat composed of a single VSG type.

While population-level heterogeneity in surface protein expression has been suggested as critical
for *T. cruzi* infection, immune evasion, and persistence [20, 21], this has not been studied at the
intra-population level, and the underlying mechanisms remain poorly understood. Here, we
present a scRNA-seq analysis of *T. cruzi*, to understand the heterogeneity in surface protein
expression within trypomastigote populations.

Results and Discussion

Identification of cell populations

To assess the expression of cell-surface protein-coding genes in *Trypanosoma cruzi*, we
conducted a 10X Chromium Single Cell 3' assay from a mixed population of amastigotes and
cell-derived trypomastigotes, aiming at sequencing the transcriptome of 5000 cells. After low-
quality cell filtering and gene expression quantification (see Materials and Methods), we
obtained 3192 single-cell transcriptomes with 14321 total genes detected, with a mean of 1088
genes and 2461 UMIs detected per cell. These results were comparable to other scRNA-seq
studies done in *Trypanosoma brucei* and recently reported as a preprint in *T. cruzi* using 10X
Chromium technology [15, 22, 23]. Cell populations (**Figure 1a**) were defined by identifying
cluster-specific gene markers (**Figure 1b, Supplementary File 1**). Two cell clusters were
assigned to trypomastigotes and amastigotes (cluster 0 and 1, respectively). Markers gene
expression was consistent with previous bulk RNA-seq data from Dm28c [24] (**Figures 1b, 1c**
and 1d, Supplementary Table 1). In turn, we hypothesize that cluster 2 reflects amastigote-
trypomastigote transitioning parasites, as its gene markers are differentially expressed in bulk
RNA-seq data, but some are upregulated in amastigotes and others in trypomastigotes.

Figure 1. (a) UMAP colored by detected clusters based on gene expression profiles, (b) Heatmap
of the top 10 gene markers upregulated in each of the 3 cell populations identified (log2FC > 1
and adjusted p-value < 0.05), (c) Expression of a cluster 0 marker gene (C4B63-16g183) on the
UMAP, and (d) Expression of a cluster 1 marker gene (C4B63-16g155) on the UMAP.

Expression pattern of surface protein-coding genes

We analyzed differences in gene expression between single-copy genes and multigene families,
between and within the identified cell populations. As expected, single-copy gene expression is
high in both amastigotes and trypomastigotes and similar on average between both cell types
(**Supplementary Figure 1a**), while surface protein-coding genes are more expressed in
trypomastigotes [25] (**Figure 2a**).

Within trypomastigotes, heterogenous expression of surface gene families was observed, with
high variation in the number of cells in which each surface gene was detected, as well as the total
expression level in each cell (**Figure 2b**). Even though expression heterogeneity is also observed
for single-copy genes (**Supplementary Figure 1a and 1b**), probably due to the sampling biases
that cause gene dropout in 10X Genomics technology, we investigated whether this phenomenon
was more pronounced in surface proteins. Therefore, we analyzed differences among single-copy
and multigene family genes (together or grouped by multigene family) in trypomastigotes, in
terms of the number of cells expressing each of the individual genes of each group (**Figure 2c**
and Supplementary Figure 1c). We additionally calculated their Gini index (**Figure 2d and**
Supplementary Figure 1d) [26] to assess the extent to which the expression levels of a given
gene were evenly distributed across individual cells. Ribosomal protein-coding genes were
included as a control group for homogeneous expression.

Interestingly, compared to single-copy genes, especially ribosomal genes, multigene family
genes showed a greater dispersion regarding the number of cells in which each gene was
detected (**Figure 2c and Supplementary Figure 1c**), and a higher value of its Gini indexes with
a lower dispersion (**Figure 2d and Supplementary Figure 1d**). Both observations indicate a
higher expression heterogeneity for surface protein expression in the trypomastigote population.
Moreover, when different families were analyzed separately, TcS genes showed a particularly
high heterogeneity (**Supplementary Figure 1c and 1d**).

Figure 2. (a) Summatory of expression levels values from all genes belonging to the disruptive
compartment for each cell from amastigote (Cluster 1) and trypomastigote (Cluster 0) cell
populations, (b) UMAP visualization of the expression patterns of multigene family genes. (c)
Violin plots showing the number of cells expressing a specific gene belonging to each group of
genes: subsampled single-copy and multigene families, ribosomal genes and trans-sialidases. To
avoid biases against size differences between compartments, we generated a subsampled single-
copy genes list, randomly selecting an equal number of genes as those from the multigene
family's gene set. The expression distribution of the subsampled single-copy genes is similar to
the distribution of the entire dataset. (d) Boxplots depict the Gini index, across single-copy and
multigene families, ribosomal genes and trans-sialidases. * p < 0.05, ** p < 0.01, *** p < 0.001,
**** p < 0.0001. For (c) and (d), all comparisons of the multigene families against single-copy
and ribosomal genes are statistically significant.

Expression pattern of the TcS superfamily

When we re-clustered the trypomastigote population based solely on trans-sialidase gene
expression, we identified two sub-populations: trypomastigote cluster 0 ("trypo_0" composed of
1186 cells), which over-expressed these genes compared to trypomastigote cluster 1 ("trypo_1"
composed of 1015 cells) (**Figure 3**).

Figure 3. Trypomastigote sub-populations identified based on trans-sialidase expression profiles.
(a) violin plot displaying average expression levels of ribosomal protein-coding genes across
sub-populations. (b) violin plot showing combined trans-sialidase expression levels for each sub-
population. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

The two trypomastigote sub-populations segregated by TcS expression, show only slight
differences in the expression of other gene categories, such as ribosomal protein-coding genes
(FC = 1.03), transporters (FC = 1.09), polymerase-related genes (FC = 1.15), and phosphatases
(FC = 1.10) (Supplementary Figure 2). Although there is a tendency for surface protein genes to
be more expressed in cluster trypo_0 (FC = 1.38), trans-sialidases displayed the highest fold
change between the two trypomastigote subpopulations (FC = 1.53) (**Supplementary Figure 2**).
It is tempting to speculate that this may reflect different infectivity amongst trypomastigote
subpopulations, consistent with reports of "broad" and "slender" forms [27]. Recently, while this
manuscript was being prepared, a similar approach by Laidlaw et al. described the phenomenon
by clustering trypomastigote cells based on the expression of all genes [23]. When applying this
strategy to our data, their observation was reproduced, further demonstrating the consistent
reproducibility of the scRNA-seq results across studies.

When all cells are considered, most TcS genes are expressed, but in each individual cell only
approximately 40 TcS genes were detected (**Figure 4d upper pannel**). Interestingly, we
observed that both trypomastigote subpopulations contain a subgroup of TcS genes that are
detected in a large portion of cells (>40%) (**Figure 4a, Supplementary Table 2**), indicating
high-level expression at the population level. However, these genes did not account for most of
the family´s expression in each individual cell. Even family members detected in only a few cells
significantly contribute to the total TcS expression in those cells, being quantified at similar
levels as those from the frequently detected group (**Figure 4b, 4c and 4d**). Gene dropouts might
favor random patterns of gene family's detection in scRNA-seq experiments, particularly
affecting genes with low expression. Since genes detected in a high percentage of cells are not
more highly expressed than those detected in only a few cells (**Figure 4b**), we do not believe that
this characteristic of scRNA-seq data is responsible for our observation. Actually, at the single-
cell level, each detected TcS gene contributes similarly to the overall family expression in those
cells, as shown by a small average Gini index for the percentage of the expression of the family
in each cell (**Figure 4c and 4d**). This highlights a key limitation of bulk RNA-seq, as it may
wrongly indicate that a set of few genes are highly expressed in each cell, while in fact, TcS
expression is evenly distributed among all detected family members, regardless of the number of
cells expressing each gene. Indeed, most TcS members detected in more than 40% of cells in this
work were found in the top 100 expressed TcS in an independent bulk RNA-seq study
(**Supplementary Figure 3a**).

When analyzing each trypomastigote subpopulation, no coordinated expression among specific
TcS members was observed, as no subclusters of cells were identified based on TcS detection
profiles. Even when clustering was restricted to genes detected in more than 40% of cells, no
clear subclusters of cells were identified (**Supplementary Figure 3b**).

The loci for TcS genes and other gene families of surface proteins are mostly grouped in specific
genomic compartments [28] that are regulated epigenetically by the activation or silencing of
chromatin folding domains [24]. The TcS that are detected in a high percentage of cells are
mostly dispersed throughout the genome (**Supplementary Table 2**). This suggests that their
preferential expression is likely not due to colocalization in one or a few ubiquitous activated
chromatin-folding domains. These observations raise intriguing questions about the mechanisms
governing the selection of TcS members expressed in each cell, warranting further investigation.

Figure 4. (a) Heatmap displaying the expression of TcS genes in each cell that together account for 75% of total TcS gene expression within cluster trypo 0. Cells are clustered by TcS expression profiles, with colors representing each gene's percentage contribution to the cell's total TcS expression. (b) Average expression of TcS genes grouped by the percentage of cells expressing each gene, (c) Gini index distribution for trypomastigotes cluster 0 cells considering only TcS detected in each cell. (d) Cumulative plot showing the percentage of TcS expression in each cell from cluster trypo_0. Genes are ordered by descending expression level and sequentially added to obtain each value on the Y-axis. The observed linear increase suggests a relatively equal contribution of each gene to the overall expression.

Final remarks

The expression of surface protein-coding genes varied between cell stages. Genes that belong to
the trans-sialidase-like superfamily, involved in host-parasite interactions, displayed significant
heterogeneity in expression levels within trypomastigotes. This variation in TcS expression
suggests that not all trypomastigotes express these genes uniformly. The lack of coordinated
expression among TcS members, observed in single-cell and bulk RNA-seq studies [3], suggests
a complex regulatory mechanism that enables adaptability in different *T. cruzi* subpopulations.
These findings reinforce the importance of TcS variability in driving phenotypic diversity and
pathogenicity in *T. cruzi*. Taken together, these findings not only demonstrate the sensitivity of
scRNA-seq in delineating parasite life stages and their associated gene expression profiles, but
also reveal a complex regulation of surface proteins and TcS family members, with tentative
implications for immune evasion and pathogenicity.

Materials and Methods

Trypanosoma cruzi and mammalian cell culture

Epimastigote forms of *Trypanosoma cruzi* strain Dm28c were derived from axenic cultures
cultivated in Brain-Heart Infusion medium (BHI, Oxoid) supplemented with 10% heat-
inactivated fetal bovine serum (FBS, Capricorn), penicillin (100 units/mL) and streptomycin
(100 µg/mL) as described [29]. Cultures were diluted 1/10 with fresh BHI medium every 3 days
and maintained at 28 °C.

Myoblast rat cell line H9c2 (ATCC CRL-1446) was maintained in hgDMEM medium (Gibco)
supplemented with 10% heat-inactivated fetal bovine serum, penicillin (100 units/mL) and
streptomycin (100 µg/mL) at 37 C in a humidified 5% CO2 incubator. Confluent cells were
washed with 1X phosphate-buffered saline (1X PBS), incubated for 5 min with trypsin-EDTA
(Gibco), diluted with culture medium and re-plated for maintenance.

Mycoplasma contamination in cell lines was regularly monitored using MycoAlert®
Mycoplasma Detection Kit (Lonza), following the manufacturer's protocol.

Isolation and purification of cellular trypomastigotes and intracellular amastigotes

Late stationary phase epimastigotes of Dm28c strain were used to infect H9c2 cells for a primary
infection. Six days post-infection, cellular trypomastigotes were obtained from the supernatant
and were used to infect 50% confluent H9c2 cells at a 10:1 rate. Twenty-four hours post-
infection the cell culture was washed twice with PBS 1X and maintained with fresh hgDMEM at
37 C in a humidified 5% CO2 incubator.

For amastigote purification, H9c2 cells grown in monolayers and infected for 48 hours were
washed with 1X PBS and incubated with trypsin-EDTA for 5 minutes at 37°C. The
trypsinization was stopped by adding an equal volume of hgDMEM with 10% FBS. The cell

suspension was repeatedly passed through a 27-gauge needle attached to a 30-mL syringe until
complete cell disruption was confirmed under the microscope. The supernatant, containing free
amastigotes, was collected and centrifuged at 500xg for 10 minutes at 4°C to remove large host-
cell debris. The resulting supernatant was then centrifuged at 4,000xg for 10 minutes at 4°C, and
the amastigote-containing pellet was washed twice in chilled 1X DPBS (Dulbecco's Phosphate-
Buffered Saline, No Calcium, No Magnesium) and resuspended in 1X DPBS at 200 μL per
$1x10^6$ cells, ready for the fixation step.

Cellular trypomastigotes derived from infected H9c2 cells and present in the cell supernatant
fraction were collected and centrifuged at 500xg for 10 minutes at 4°C to remove large host-cell
debris. The washing and resuspension steps in DPBS were performed for amastigotes as
described above.

scRNA-seq library preparation and sequencing

Cell fixation was performed using the Methanol Fixation Protocol for Single-Cell RNA
Sequencing [30] as recommended by 10X Genomics technical support. Briefly, chilled 100%
Methanol (for HPLC, ≥99.9%, Millipore) was added drop by drop ($1x10^6$ cells in 800 μl) and
incubated at -20°C for 30 min. For rehydration, fixed cells were first equilibrated at 4°C and then
centrifuged at 4,000xg for 5 min at 4°C, the supernatant was discarded and Wash-Resuspension
Buffer (3X SSC in Nuclease-free Water, 0.04% UltraPure Bovine Serum Albumin, 1mM DTT,
and 0.2 U/ml RNase Inhibitor) was added to the pellet. Cell debris and large clumps were
eliminated by passing the sample through a 40 μm Flowmi Cell Strainer.

10X Genomics library preparation was performed at the service provided by the Instituto de
Biología y Medicina Experimental (IBYME, Argentina). The library was sequenced by a service
provider (Macrogen, Korea) in a HiSeq2500 equipment (two lanes), generating approx. 880
million reads. Raw data is available in the SRA (https://www.ncbi.nlm.nih.gov/sra/) under
BioProject PRJNA1200704.

Transcript quantification

T. cruzi 2018 Dm28c genome [28] (release 62, TriTrypDB) and *T. cruzi* Dm28c maxi circle
kDNA sequence [31] were combined and used as reference. To improve the proportion of reads
assigned to genes, 11,362 3′UTR regions of the coding sequences (CDS) were annotated using
peaks2UTR [32]. Gene expression quantification was performed by pseudoalignment using
kallisto bustools [33], with the options --filter to remove potential noise from environmental
RNA and --em to apply the Expectation-Maximization (EM) algorithm. This algorithm
outperforms other mapping software in handling multimapping reads, enabling more accurate
quantification of multigene families [33].

scRNA-seq data processing and analysis

Count matrices from two technical replicates obtained by sequencing the same library were
merged: the common barcodes across both datasets were retained, and the count matrices were

combined to generate a unified dataset. Subsequently, the following metrics were calculated:
nUMI, nGene, and mitoRatio, as well as the log10 of genes per UMI (log10GenesperUMI). A
filtering criterion was applied, retaining cells with nUMI > 1200, nGene > 100,
log10GenesperUMI > 0.8, and mitoRatio < 0.1. Additionally, ribosomal genes were excluded
from subsequent analyses. Data normalization and scaling were performed using the Seurat R
version 5 package [34], employing NormalizeData and ScaleData functions.

FindNeighbors function was used to construct a k-nearest neighbors graph of cells using 10
principal components (PCs) and Louvain algorithm was employed for cell clustering using
FindClusters function.

To define marker genes, the FindAllMarkers function from Seurat package was used, selecting
those with an adjusted p-value < 0.05 and a log2 fold change (log2FC) > 1. To validate these
markers, bulk RNA-seq data of Dm28c was incorporated (NCBI BioProject ID PRJNA850400
[24]). Transcripts were quantified using Kallisto [35] (with -b 100 option to perform 100
bootstraps), followed by differential expression analysis conducted with Sleuth [36]. Genes with
an adjusted p-value < 0.05 and |log2FC| > 0.25 were filtered for further analysis. Finally, the
gene IDs of the Seurat-defined markers were cross-referenced with the IDs of the differentially
expressed genes obtained from the bulk RNA-seq analysis to corroborate stage-specific gene
expression. For 2D visualization of cell clusters and gene markers expression profiles across
cells, UMAP projection was employed [37].

Gene IDs corresponding to multigene families (trans-sialidase and trans-sialidase-like, MASP,
Mucins, GP63, RHS and DGF) were obtained by text searches using the current genome
annotation on the "description" field. Single and low copy number genes were defined as those
that did not belong to the latter gene list. For simplicity "single-copy" will be used to refer to
these genes throughout the manuscript.

Data processing was conducted using R version 4.2.0. Statistical analyses were performed using
the Wilcoxon rank-sum test and p-values < 0.05 were considered statistically significant.

To study expression inequality, we applied the Gini index, a metric originally developed in the
field of economics [26]. The Gini index serves as a measure of expression heterogeneity, and we
employed it in two distinct contexts: first, to evaluate the degree of inequality in the distribution
of a gene's expression levels across individual cells (**Figure 2d and Supplementary Figure 1d**);
and second, to assess the extent to which a given cell expresses individual genes at varying rates
(**Figure 4c**).

Author Contributions

320 LI: Data Curation, Formal Analysis, Investigation, Methodology, Visualization, Writing-Original
321 Draft Preparation, Writing-Review & Editing; LB: Methodology, Investigation, Writing-Review

& Editing; VAC: Methodology, Investigation, Writing-Review & Editing; JG: Methodology, Formal Analysis, Writing-Review & Editing; GR: Methodology, Writing-Review & Editing; VMH: Methodology, Writing-Review & Editing; MAD: Methodology, Resources, Writing-Review & Editing; JSS: Methodology, Resources, Writing-Review & Editing; JDG: Conceptualization, Methodology, Visualization, Writing-Original Draft Preparation, Writing-Review & Editing; PS: Conceptualization, Funding Acquisition, Methodology, Formal Analysis, Project Administration, Resources, Supervision, Validation, Writing-Original Draft Preparation, Writing-Review & Editing

Conflict of interest

The authors declare that they have no conflicts of interest

Financial Support

This project was supported by: CSIC, Universidad de la República, grant number: I+D-2020-505 awarded to PS; LI, LB, JG, MD, JSS and PS received financial support from PEDECIBA. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Tarleton, R.L., *Chagas Disease: A Solvable Problem, Ignored.* Trends Mol Med, 2016. **22**(10): p. 835-838.
2. Martin-Escolano, J., et al., *An Updated View of the Trypanosoma cruzi Life Cycle: Intervention Points for an Effective Treatment.* ACS Infect Dis, 2022. **8**(6): p. 1107-1115.
3. Belew, A.T., et al., *Comparative transcriptome profiling of virulent and non-virulent Trypanosoma cruzi underlines the role of surface proteins during infection.* PLoS Pathog, 2017. **13**(12): p. e1006767.
4. Pech-Canul, A.C., V. Monteon, and R.L. Solis-Oviedo, *A Brief View of the Surface Membrane Proteins from Trypanosoma cruzi.* J Parasitol Res, 2017. **2017**: p. 3751403.
5. Freitas, L.M., et al., *Genomic analyses, gene expression and antigenic profile of the trans-sialidase superfamily of Trypanosoma cruzi reveal an undetected level of complexity.* PLoS One, 2011. **6**(10): p. e25914.
6. Burle-Caldas, G.A., et al., *Disruption of Active Trans-Sialidase Genes Impairs Egress from Mammalian Host Cells and Generates Highly Attenuated Trypanosoma cruzi Parasites.* mBio, 2022: p. e0347821.
7. Berna, L., et al., *Transcriptomic analysis reveals metabolic switches and surface remodeling as key processes for stage transition in Trypanosoma cruzi.* PeerJ, 2017. **5**: p. e3017.

359   8.   Tonelli, R.R., et al., *Role of the gp85/trans-sialidases in Trypanosoma cruzi tissue*
360   *tropism: preferential binding of a conserved peptide motif to the vasculature in vivo.*
361   PLoS Negl Trop Dis, 2010. **4**(11): p. e864.

362   9.   Walzer, K.A., et al., *Single-Cell Analysis Reveals Distinct Gene Expression and*
363   *Heterogeneity in Male and Female Plasmodium falciparum Gametocytes.* mSphere, 2018.
364   **3**(2).

365   10.   Ngara, M., et al., *Exploring parasite heterogeneity using single-cell RNA-seq reveals a*
366   *gene signature among sexual stage Plasmodium falciparum parasites.* Exp Cell Res,
367   2018. **371**(1): p. 130-138.

368   11.   Reid, A.J., et al., *Single-cell RNA-seq reveals hidden transcriptional variation in malaria*
369   *parasites.* Elife, 2018. **7**.

370   12.   Howick, V.M., et al., *The Malaria Cell Atlas: Single parasite transcriptomes across the*
371   *complete Plasmodium life cycle.* Science, 2019. **365**(6455).

372   13.   Vigneron, A., et al., *Single-cell RNA sequencing of Trypanosoma brucei from tsetse*
373   *salivary glands unveils metacyclogenesis and identifies potential transmission blocking*
374   *antigens.* Proc Natl Acad Sci U S A, 2020. **117**(5): p. 2613-2621.

375   14.   Hutchinson, S., et al., *The establishment of variant surface glycoprotein monoallelic*
376   *expression revealed by single-cell RNA-seq of Trypanosoma brucei in the tsetse fly*
377   *salivary glands.* PLoS Pathog, 2021. **17**(9): p. e1009904.

378   15.   Briggs, E.M., et al., *Single-cell transcriptomic analysis of bloodstream Trypanosoma*
379   *brucei reconstructs cell cycle progression and developmental quorum sensing.* Nat
380   Commun, 2021. **12**(1): p. 5268.

381   16.   Howick, V.M., et al., *Single-cell transcriptomics reveals expression profiles of*
382   *Trypanosoma brucei sexual stages.* PLoS Pathog, 2022. **18**(3): p. e1010346.

383   17.   Quintana, J.F., et al., *Single cell and spatial transcriptomic analyses reveal microglia-*
384   *plasma cell crosstalk in the brain during Trypanosoma brucei infection.* Nat Commun,
385   2022. **13**(1): p. 5752.

386   18.   Catta-Preta, C.M.C., et al., *Single-cell atlas of Leishmania development in sandflies*
387   *reveals the heterogeneity of transmitted parasites and their role in infection.* Proc Natl
388   Acad Sci U S A, 2024. **121**(52): p. e2406776121.

389   19.   Abuchery, B.E., J.A. Black, and M.S. da Silva, *Single-cell transcriptomics reveals hidden*
390   *information in trypanosomatids.* Trends Parasitol, 2022. **38**(1): p. 4-6.

391   20.   Seco-Hidalgo, V., L.M. De Pablos, and A. Osuna, *Transcriptional and phenotypical*
392   *heterogeneity of Trypanosoma cruzi cell populations.* Open Biol, 2015. **5**(12): p. 150190.

393   21.   Luzak, V., et al., *Cell-to-Cell Heterogeneity in Trypanosomes.* Annu Rev Microbiol,
394   2021. **75**: p. 107-128.

395   22.   Bard, J.E., et al., *Life stage-specific poly(A) site selection regulated by Trypanosoma*
396   *brucei DRBD18.* Proc Natl Acad Sci U S A, 2024. **121**(29): p. e2403188121.

397   23.   Laidlaw, R., et al., *The Trypanosoma cruzi cell atlas; a single-cell resource for*
398   *understanding parasite population heterogeneity and differentiation.* bioRxiv, 2024: p.
399   2024.10.01.616042.

400   24.   Diaz-Viraque, F., et al., *Genome-wide chromatin interaction map for Trypanosoma cruzi.*
401   Nat Microbiol, 2023. **8**(11): p. 2103-2114.

402   25.   Li, Y., et al., *Transcriptome Remodeling in Trypanosoma cruzi and Human Cells during*
403   *Intracellular Infection.* PLoS Pathog, 2016. **12**(4): p. e1005511.

404   26.   Atkinson, A., *On the measurement of inequality.* Journal of Economic Theory, 1970.
405   **2**(3): p. 244-263.
406   27.   Schmatz, D.M., R.C. Boltz, and P.K. Murray, *Trypanosoma cruzi: separation of broad*
407   *and slender trypomastigotes using a continuous hypaque gradient.* Parasitology, 1983. **87**
408   **(Pt 2)**: p. 219-27.
409   28.   Berna, L., et al., *Expanding an expanded genome: long-read sequencing of Trypanosoma*
410   *cruzi.* Microb Genom, 2018.
411   29.   Smircich, P., et al., *Transcriptomic analysis of the adaptation to prolonged starvation of*
412   *the insect-dwelling Trypanosoma cruzi epimastigotes.* Front Cell Infect Microbiol, 2023.
413   **13**: p. 1138456.
414   30.   Gutierrez-Franco, A., et al., *Methanol fixation is the method of choice for droplet-based*
415   *single-cell transcriptomics of neural cells.* Commun Biol, 2023. **6**(1): p. 522.
416   31.   Berna, L., et al., *Maxicircle architecture and evolutionary insights into Trypanosoma*
417   *cruzi complex.* PLoS Negl Trop Dis, 2021. **15**(8): p. e0009719.
418   32.   Haese-Hill, W., K. Crouch, and T.D. Otto, *peaks2utr: a robust Python tool for the*
419   *annotation of 3' UTRs.* Bioinformatics, 2023. **39**(3).
420   33.   Sullivan, D.K., et al., *kallisto, bustools and kb-python for quantifying bulk, single-cell*
421   *and single-nucleus RNA-seq.* Nat Protoc, 2024.
422   34.   Hao, Y., et al., *Dictionary learning for integrative, multimodal and scalable single-cell*
423   *analysis.* Nat Biotechnol, 2024. **42**(2): p. 293-304.
424   35.   Bray, N.L., et al., *Near-optimal probabilistic RNA-seq quantification.* Nat Biotechnol,
425   2016. **34**(5): p. 525-7.
426   36.   Pimentel, H., et al., *Differential analysis of RNA-seq incorporating quantification*
427   *uncertainty.* Nat Methods, 2017. **14**(7): p. 687-690.
428   37.   McInnes, L., et al., *UMAP: Uniform Manifold Approximation and Projection.* Journal of
429   Open Source Software, 2018. **3**(29).

Supplementary Figures

Supplementary Figure 1. Gene expression in amastigote and trypomastigote cell populations. To minimize biases related to size differences between compartments, we generated a subsampled single-copy genes list, by randomly selecting an equal number of genes to match those from the multigene family's gene set. (a) For each gene set, we calculated the summatory expression of those detected genes for each cell (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$), (b) UMAP projection for 2D visualization of core gene expression among cells, (c) Violin plots showing the number of cells expressing a specific gene belonging to each group of genes: subsampled single-copy and multigene families, ribosomal genes and different multigene families, and (d) Gini-indexes for subsampled single-copy genes, multigene family genes (together or grouped by multigene families) and ribosomal protein-coding genes as a constitutive expression control group, in trypomastigotes. All comparisons among multigene families and single-copy and ribosomal genes yield statistically significant differences.

Supplementary Figure 2.

Trypomastigote sub-clusters identified based on trans-sialidase expression profiles. Violin plots
displaying average expression levels across sub-clusters and associated fold changes
($FC_{trypo\_0/trypo\_1}$) of (a) transporters coding genes, (b) DNA and RNA polymerase-associated
protein coding genes, (c) phosphatases coding genes and (d) multigene family genes. * $p < 0.05$,
** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Supplementary Figure 3. (a) Venn diagram showing the overlap between the top 100 most
expressed TcS from bulk RNA-seq data and TcS expressed in more than 40% of cells from
cluster trypo_0, and (b) Heatmap displaying the expression of TcS genes in each cell that
together account for 75% of total TcS gene expression and are expressed in more than 40% of
cells within cluster trypo_0. Cells are clustered by TcS expression profiles, with colors
representing each gene's percentage contribution to the cell's expression.

463