Single-cell RNA-seq reveals trans-sialidase-like superfamily gene expression heterogeneity in
***Trypanosoma cruzi* populations.**

Lucas Inchausti[1,2], Lucía Bilbao[1,2], Vanina A. Campo[3,4], Joaquín Garat[5], José Sotelo-Silveira[5,6],
Gabriel Rinaldi[7], Virginia M. Howick[8], María Ana Duhagon[2], Javier G. De Gaudenzi[3,4], Pablo
Smircich[1,2]

1 - Laboratorio de Bioinformática, Departamento de Genómica, Instituto de Investigaciones
Biológicas Clemente Estable, Montevideo, Uruguay

2 - Sección Genómica Funcional, Facultad de Ciencias, Universidad de la República, Montevideo,
Uruguay

3 - Instituto de Investigaciones Biotecnológicas, Universidad Nacional de San Martín-Consejo
Nacional de Investigaciones Científicas y Técnicas, General San Martín, Prov. de Buenos Aires,
Argentina

4 - Escuela de Bio y Nanotecnologías (EByN), Universidad Nacional de San Martín, General San
Martín, Prov. de Buenos Aires, Argentina

5 - Departamento de Genómica, Instituto de Investigaciones Biológicas Clemente Estable,
Montevideo, Uruguay

6 - Sección Biología Celular, Facultad de Ciencias, Universidad de la República, Montevideo,
Uruguay

7 - Department of Biology, University of Oxford, Oxford, United Kingdom

8 - Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow,
Glasgow, United Kingdom

Abstract

Trypanosoma cruzi, the causative agent of Chagas disease, presents a major public health
challenge in Central and South America, affecting approximately 8 million people and placing
millions more at risk. The *T. cruzi* life cycle includes transitions between epimastigote,
metacyclic trypomastigote, amastigote, and blood trypomastigote stages, each marked by
distinct morphological and molecular adaptations to different hosts and environments. Unlike
other trypanosomatids such as *Trypanosoma brucei*, *T. cruzi* does not employ a monoallelic
model of antigenic variation; instead, it relies on a diverse repertoire of cell-surface associated
proteins encoded by large multigene families, which are essential for infectivity and immune
evasion.

This study analyzes cell-specific transcriptomes using single-cell RNA sequencing of amastigote
and trypomastigote cells to characterize stage-specific surface protein expression during
mammalian infection. Through clustering and identification of cell-specific markers, we
assigned cells to distinct parasite developmental forms. Analysis of individual cells revealed that
surface protein-coding genes, especially members of the trans-sialidase like superfamily (TcS),
are expressed with greater heterogeneity than single-copy genes. Moreover, no recurrent
combinations of TcS genes were observed between individual cells in the population.
Remarkably, a small subset of TcS mRNAs, encoded by genes preferentially located in the core
genomic compartment, are frequently detected across the cell population, whereas the vast
majority of TcS mRNAs show low detection frequencies and are mainly encoded in the
disruptive compartment. Our findings thus reveal transcriptomic heterogeneity within
trypomastigote populations where each cell displays unique TcS expression profiles. Focusing
on the diversity of surface protein expression, this research aims to deepen our understanding
of *T. cruzi* cellular biology and infection strategies.

Introduction

The protozoan parasite *Trypanosoma cruzi* is the etiological agent of Chagas disease, a highly
prevalent infectious disease in Central and South America that affects approximately 8 million
people, with several million more at risk of infection (1).

The parasite life cycle involves both an invertebrate vector (triatomine bug) and a vertebrate
host. The change in environmental conditions triggers differentiation processes in the parasite
developing across four main stages. The epimastigote form replicates within the insect and
differentiates into metacyclic trypomastigotes in the insect's rectal tract. This latter form does
not replicate, specializes in host infection, and can invade various cell types. Once inside the cell,
the parasite differentiates into an amastigote, the replicative form within the mammalian host.
After several rounds of division, amastigotes differentiate into bloodstream trypomastigotes,
which, after cell lysis, can either infect new cells or be ingested by the vector (2). Parasite
developmental stages exhibit significant morphological and molecular differences, associated
with distinct gene expression profiles. In trypanosomatids, protein-coding genes are organized
into large polycistronic RNA polymerase II transcription units, with limited evidence for tight,
gene-specific transcriptional control (3). As a result, gene expression is largely regulated at the
post-transcriptional level, through mechanisms controlling mRNA steady-state abundance,
translation, and related processes (3). More recently, accumulating evidence indicates that
genome organization and chromatin state also shape transcriptional outputs (4–8). These
studies have identified two distinct chromatin compartments in *T. cruzi*: a core compartment, in
which genes tend to show more uniform expression consistent with permissive transcription

and predominant post-transcriptional regulation (9), and a disruptive compartment, enriched in
multigene families, whose genes exhibit lower average transcription and stronger chromatin-
associated regulation (4).

T. cruzi infection relies on a heterogeneous set of membrane proteins, encoded mainly by large
multigene families (10). Among these are trans-sialidases and trans-sialidase-like proteins (TcS),
mucins, MASPs, GP63, DGFs, and RHS proteins, most of which are involved in infection, tropism,
and immune evasion (11–16).

The trans-sialidase-like superfamily is involved in processes underlying host-parasite
interactions (12). Members that retain enzymatic activity catalyze the transfer of sialyl groups
from host glycoconjugates to galactopyranosyl units on the parasite's surface, an essential
activity, as *T. cruzi* is unable to synthesize *de novo* sialic acid (17). This is the largest superfamily,
with over 1,400 members, and is subdivided into eight groups based on amino acid sequence
(12). Few members have been functionally characterized, most expressed primarily in
mammalian stages (18). The TcS group I include proteins that retain trans-sialidase activity,
though members from all groups are involved in host-parasite interactions (19).

In recent years, single-cell RNA sequencing (scRNA-seq) has been employed in protozoa, with
reports including *Plasmodium falciparum*, *Trypanosoma brucei,* and *Leishmania major* (20–29).
These studies revealed key aspects of the infection process undetectable with conventional
methods, highlighting the relevance of this approach for understanding individual variation in
gene expression in single-celled organisms (30). In *T. brucei,* antigenic variation driven by
variant surface glycoproteins (VSGs) has been studied at single-cell resolution to understand
the mechanisms that enable subpopulations of this parasite to evade the immune system, as a
bet hedging strategy, ensuring parasite survival (25). In this parasite, scRNA-seq revealed that
pre-metacyclic cells express multiple VSG transcripts simultaneously, in contrast to metacyclic
forms, which display a protein coat composed of a single VSG type.

While population-level heterogeneity in surface protein expression has been suggested as
critical for *T. cruzi* infection, immune evasion, and persistence (31,32), this has not been studied
at the intra-population level, and the underlying mechanisms remain poorly understood. Here,
we present a scRNA-seq analysis of *T. cruzi*, to understand the heterogeneity in surface protein
expression within trypomastigote populations.

Results and Discussion

Identification of cell populations

To assess the expression of cell-surface protein-coding genes in *Trypanosoma cruzi*, we conducted a 10X Chromium Single Cell 3' assay from a mixed population of amastigotes and cell-derived trypomastigotes, aiming at sequencing the transcriptome of 5000 cells. After low-quality cell filtering and gene expression quantification (see Materials and Methods), we obtained 3192 single-cell transcriptomes with 14321 total genes detected, with a mean of 1088 genes and 2461 UMIs detected per cell. These results were comparable to other scRNA-seq studies done in *Trypanosoma brucei* and more recently addressed in *T. cruzi* (currently reported as a preprint) using 10X Chromium technology (26,33,34). Cell populations (**Figure 1a**) were defined by identifying cluster-specific gene markers (**Figure 1b and Supplementary File 1**). Two cell clusters were assigned to trypomastigotes and amastigotes: cluster 0 (2201 cells) and cluster 1 (824 cells), respectively. Markers gene expression was consistent with previous bulk RNA-seq data from Dm28c (4) (**Figures 1b, 1c and 1d, Supplementary Table 1**). In turn, we hypothesize that cluster 2, that comprised only 167 cells, reflects amastigote-trypomastigote transitioning parasites, as its gene markers are differentially expressed in bulk RNA-seq data, but some are upregulated in amastigotes and others in trypomastigotes (Supplementary Table 1).

Figure 1. Identification of amastigote and trypomastigote cell populations. (a) UMAP colored by detected clusters based on gene expression profiles. (b) Heatmap of the top 10 gene markers upregulated in each of the 3 cell populations identified (log2FC > 1 and adjusted p-value < 0.05). (c) Expression of a cluster 0 marker gene (C4B63-16g183) on the UMAP, and (d) Expression of a cluster 1 marker gene (C4B63-16g155) on the UMAP.

Expression pattern of surface protein-coding genes

We analyzed differences in gene expression between single-copy genes and multigene families, between and within the identified cell populations. As previously reported (4,18), multi gene family expression is increased in trypomastigotes (**Figure 2a**), consistent with the involvement of these genes in stage-specific functions. As control, we examined the expression of single-copy genes, for which this pattern was not observed as these genes are primarily associated with basic cellular functions and show more similar average expression levels between the two cell types (**Supplementary Figure 1a**)(33).

Within trypomastigotes, heterogenous expression of surface protein-coding gene families was observed, with high variation (compared to single-copy genes) in the number of cells in which each surface protein-coding gene was detected, as well as the total expression level in each cell

(**Figure 2b and 2c**). Even though expression heterogeneity is also observed for single-copy
genes (**Supplementary Figure 1a and Figure 1b**), probably due to the sampling biases that
cause gene dropout in 10X Genomics technology, we investigated whether this phenomenon
was more pronounced in surface multigene families. Therefore, we analyzed differences among
single-copy and multigene family genes (together or grouped by multigene family) in
trypomastigotes, in terms of the number of cells expressing each of the individual genes of each
group (**Figure 2c, Supplementary Figure 1c and Supplementary Table 2**). In addition, we
assessed expression inequality using Lorenz curves (**Figure 2d, Supplementary Figure 1d and**
Supplementary Table 2) to evaluate how unevenly gene expression is distributed within each
gene group. Ribosomal protein–coding genes were included as a control group.

Interestingly, compared to single-copy genes, especially ribosomal genes, multigene family
genes showed a greater dispersion regarding the number of cells in which each gene was
detected (**Figure 2c, Supplementary Figure 1c and Supplementary Table 2**), as well as a more
pronounced deviation from the diagonal in the Lorenz curves (**Figure 2d, Supplementary Figure**
1d), which represents perfect equality (reflected by different Gini indexes, see **Supplementary**
Table 2). Both observations indicate a higher expression heterogeneity for surface protein
expression in the trypomastigote population. When multigene families were analyzed
separately, several showed significant differences when compared to single-copy genes.
Heterogeneity of the MASP family has already been suggested by studying clonal populations
supporting our single cell results (31). Also, TcS genes exhibited a pronounced expression
heterogeneity (**Supplementary Figures 1c and 1d**). TcS constitute a well-established and
biologically central gene family in *T. cruzi*, playing key roles in host-parasite interactions
(11,12,17). In this context, we next focused on a more detailed characterization of TcS
expression heterogeneity in trypomastigotes.

Figure 2. Overview of expression patterns across amastigote and trypomastigote cells. (a)
Summatory of expression levels values from all multigene family genes for each cell from
amastigote (Cluster 1) and trypomastigote (Cluster 0) cell populations (**** p < 0.0001,
$mean_{Ama}$ = 137.2, SD_{Ama} = 48.7, $mean_{Trypo}$ = 201.1, SD_{Trypo} = 48.6, $FC_{Trypo/Ama}$ = 1.5). (b) UMAP
visualization of the expression patterns of multigene family genes; num_multigene indicates
the number of multigene family genes detected per cell (genes with >0 UMI counts).
Z_multigene reflects the relative expression level of multigene family genes per cell, calculated
as the z-score-standardized sum of their UMI counts, such that positive values reflect above-
average multigene family expression and negative values reflect below-average levels. (c)
Violin plots showing the number of cells expressing a specific gene belonging to each group of
genes: subsampled single-copy and multigene families, ribosomal genes and trans-sialidases. To
avoid biases against size differences between single-copy and multigene family genes we
generated a subsampled single-copy genes list, randomly selecting an equal number of genes as
those from the multigene family's gene set. The expression distribution of the subsampled
single-copy genes is similar to the distribution of the entire dataset (* p < 0.05, **** p < 0.0001.
See **Supplementary Table 2**). (d) Lorenz curves showing the cumulative proportion of gene
expression relative to the cumulative proportion of genes for subsampled single-copy,
multigene family genes, ribosomal protein coding genes and trans-sialidase-like genes. Genes
were ordered by total expression, and the dashed line indicates perfect equality. Curves that
deviate further from the diagonal reflect greater inequality, meaning that fewer genes account

for most of the expression within each category. Statistically significant differences between
groups for c) and d) are shown in **Supplementary Table 2**.

Expression pattern of the TcS superfamily

When we re-clustered the trypomastigote population based solely on TcS gene expression, we
identified two sub-populations: trypomastigote cluster 0 ("Trypo_0" composed of 1186 cells),
which over-expressed these genes compared to trypomastigote cluster 1 ("Trypo_1" composed
of 1015 cells) (**Figure 3a**).

Figure 3. Trypomastigote sub-populations identified based on trans-sialidase expression profiles.
(a) UMAP visualization colored by detected clusters based on gene expression profiles, with
trypomastigote subpopulations identified. (b) Violin plot displaying average expression levels of
ribosomal protein-coding genes across sub-populations. (c) Violin plot showing combined trans-
sialidase expression levels for each sub-population. * p < 0.05, **** p < 0.0001.

The two trypomastigote sub-populations segregated by TcS expression show only slight
differences in the expression of other gene categories, such as ribosomal protein-coding genes
(FC = 1.03), transporters (FC = 1.09), polymerase-related genes (FC = 1.15), and phosphatases
(FC = 1.10) (**Figure 3b and Supplementary Figure 2**, foldchange values correspond to
Trypo_0/Trypo_1 expression ratios). Although there is a tendency for surface protein genes to
be more expressed in cluster Trypo_0 (FC = 1.38), trans-sialidases displayed the highest fold
change between the two trypomastigote subpopulations (FC = 1.53) (**Figure 3c and
Supplementary Figure 2**). It is tempting to speculate that this may reflect different infectivity
amongst trypomastigote subpopulations, consistent with reports of "broad" and "slender"
forms (35). Recently, while this manuscript was being prepared, a similar approach by Laidlaw
et al. described the phenomenon by clustering trypomastigote cells based on the expression of
all genes (34). When applying this strategy to our data, their observation was reproduced.

When all cells are considered, most TcS genes are expressed, but in each individual cell only
approximately 40 TcS genes were detected. Interestingly, we observed that both
trypomastigote subpopulations contain a subgroup of TcS genes that are detected in a large
portion of cells (>40%) (**Figure 4a, and Supplementary Table 3**), indicating high-level expression
at the population level. This subset comprises 31 TcS genes belonging to subfamilies II-VI, and
VIII (**Supplementary Table 3**). Consistent with our findings, Laidlaw et al. (34) reported a similar
phenomenon, detecting on average approximately 40 TcS genes per cell, with a subset that are
frequently detected across the trypomastigote population, further supporting the
reproducibility of scRNA-seq results across studies. Gene dropouts in scRNA-seq experiments
can generate apparently stochastic detection patterns for multigene family members,
particularly affecting low-abundance transcripts, and thus represent an important potential
confounder in the analysis of multigene families. To evaluate whether this technical effect
could account for the observed TcS detection patterns, we examined the relationship between
detection frequency and expression level. If random sampling were the primary driver, genes
with similar average expression levels would be expected to exhibit comparable detection
frequencies. However, we observed that many TcS genes with similar average expression are
detected in markedly different proportions of cells (**Figure 4b**), arguing against a purely
stochastic dropout model. Notably, among the 50 TcS genes with the highest average
expression across cells, 62% are detected in fewer than 5% of cells (**Figure 4b**).

At the single-cell level, we found that detected TcS genes contribute relatively evenly to the
total family expression within each cell, as indicated by the low average Gini index of TcS
expression levels across each cell (**Figures 4c and 4d**). This suggests that, within each cell, total
TcS expression is shared across the detected TcS genes rather than being dominated by a few
very highly expressed transcripts in that cell. Consistent with this interpretation, most TcS genes
detected in a high fraction of cells in our scRNA-seq dataset also rank among the most highly

expressed TcS genes in an independent bulk RNA-seq study (**Supplementary Figure 3a**). This
highlights a key limitation of bulk RNA-seq, as it may wrongly indicate that a set of few genes
are highly expressed in each cell, while in fact, TcS expression is evenly distributed among all
detected family members, regardless of the number of cells expressing each gene. Taken
together with the results in **Figure 4b**, this observation further supports the conclusion that the
observed expression pattern is unlikely to arise from a purely stochastic detection of TcS
mRNAs.

While our data indicates transcriptional heterogeneity, it is important to note that future
studies will be required to determine the extent to which this diversity translates to protein
expression on the parasite surface. Consistent with our observations, recent proteomic
analyses (36) report pronounced variability in the expression of surface proteins, including TcS,
suggesting that the heterogeneous transcriptional patterns observed here likely reflect
biologically relevant differences.

When analyzing each trypomastigote subpopulation, no coordinated expression among specific
TcS members was observed, as no subclusters of cells were identified based on TcS detection
profiles. Even when clustering was restricted to genes detected in more than 40% of cells, no
clear subclusters of cells were identified (**Supplementary Figure 3b**).

Figure 4. Overview of TcS gene expression patterns in Trypo_0 cells. (a) Heatmap displaying the expression of TcS genes in each cell that together account for 75% of total TcS gene expression within cluster Trypo_0. Cells are clustered by TcS expression profiles, with colors representing each gene's percentage contribution to the cell's total TcS expression. (b) Average expression of TcS genes grouped by the percentage of cells expressing each gene. In red are highlighted the top 50 TcS with highest average expression. (c) Gini index distribution for trypomastigotes cluster 0 (Trypo_0) cells considering only TcS detected in each cell. (d) Lorenz curves showing, for each cell in cluster Trypo_0, the cumulative proportion of total TcS expression as a function of the cumulative proportion of detected TcS genes. Genes were ordered by total expression, and the dashed line indicates perfect equality (i.e., all detected TcS genes contribute equally to the total TcS expression of a given cell). Green and orange curves correspond to cells with higher and lower expression equality, respectively.

As discussed in the Introduction, although post-transcriptional control remains a central mechanism of gene expression regulation in *T. cruzi*, increasing evidence in recent years has revealed an important contribution of epigenetic mechanisms that modulate gene expression at the transcriptional level. (4–8). Specifically, the loci for TcS genes and other gene families of surface proteins are mostly grouped in specific genomic compartments (9) that are regulated epigenetically by the activation or silencing of chromatin folding domains (4). The TcS that are detected in a high percentage of cells are mostly dispersed throughout the genome (**Supplementary Table 3**). This suggests that their preferential expression is likely not due to colocalization in one or a few ubiquitous activated chromatin-folding domains. Nevertheless, mapping the genomic locations of TcS genes detected in a high proportion of cells revealed that most are flanked by core genes (**Figure 5**). The core compartment is enriched in conserved, single-copy genes that typically show more constitutive expression (9) and, as observed in this study, lower cell-to-cell variability. In contrast, TcS genes that are detected less frequently are preferentially located in the disruptive compartment (**Figure 5**), which is enriched in lineage-specific multigene families and associated with more variable, stage-specific, and potentially stochastic expression under tighter epigenetic control (4,9,18,36). Together, these findings suggest that the higher cellular prevalence of certain TcS transcripts is unlikely to be driven by colocalization within a small number of ubiquitously active chromatin domains but may instead reflect distinct regulatory regimes between the core and disruptive compartments. Future studies integrating single-cell chromatin profiling with scRNA-seq will be required to directly test this model.

Figure 5. Genomic context and neighborhood composition of frequently detected versus lowly detected TcS loci. (a) Representative genomic loci of frequently detected (top, high abundance) and lowly detected (bottom, low abundance) TcS genes. Genes are shown as arrows, colored according to genomic compartment: core (dark green), disruptive (salmon), and TcS genes under analysis (yellow). Chromosomal coordinates are indicated below each locus. (b) Comparison of the percentage of multigene-family neighbours within polycistronic transcription units containing frequently detected and lowly detected TcS genes. Lowly detected TcS genes were subsampled to $n = 30$. Wilcoxon rank-sum test: $p = 6.6 \times 10^{-7}$. (c) Mean percentage of multigene-family neighbours in polycistrons calculated from 50 random subsets ($n = 30$) of lowly detected TcS genes (mean = 37.76%). The orange dot indicates the corresponding mean for frequently detected TcS genes (9.27%).

Final remarks

The expression of surface protein–coding genes varied across parasite developmental stages. In particular, genes belonging to the TcS superfamily, which play key roles in host–parasite interactions, exhibited marked heterogeneity in expression among trypomastigotes. Notably, while most TcS genes were detected only in a small fraction of cells, a limited subset was frequently detected across the population. This pattern indicates that TcS expression is not uniform, suggesting the existence of distinct regulatory regimes within the family. Consistent with previous single-cell and bulk RNA-seq studies (34,36), the lack of coordinated expression among TcS members points to a complex regulatory framework that may enable functional diversification among *T. cruzi* subpopulations. Importantly, our analyses indicate that the frequent detection of this TcS subset may be partially explained by their preferential localization within the core genomic compartment, which is associated with more permissive transcriptional environments.

Taken together, our results demonstrate the sensitivity of scRNA-seq for resolving parasite life stages and their associated transcriptional programs, while also pointing to complex regulation of surface protein expression, particularly within the TcS family. Although these results should be considered with caution, as technical limitations inherent to scRNA-seq may influence the observed expression patterns, our interpretation is consistent with all observations presented here and aligns with emerging evidence from independent studies. Collectively, this supports a working hypothesis in which heterogeneous TcS expression may contribute to immune evasion and pathogenicity through a bet-hedging strategy.

Materials and Methods

Trypanosoma cruzi and mammalian cell culture

Epimastigote forms of *Trypanosoma cruzi* strain Dm28c were derived from axenic cultures cultivated in Brain-Heart Infusion medium (BHI, Oxoid) supplemented with 10% heat-inactivated fetal bovine serum (FBS, Capricorn), penicillin (100 units/mL) and streptomycin (100 μg/mL) as described (37). Cultures were diluted 1/10 with fresh BHI medium every 3 days and maintained at 28°C.

Myoblast rat cell line H9c2 (ATCC CRL-1446) was maintained in hgDMEM medium (Gibco) supplemented with 10% heat-inactivated fetal bovine serum, penicillin (100 units/mL) and streptomycin (100 μg/mL) at 37°C in a humidified 5% CO_2 incubator. Confluent cells were

332   washed with 1X phosphate-buffered saline (1X PBS), incubated for 5 min with trypsin-EDTA
333   (Gibco), diluted with culture medium and re-plated for maintenance.

334   Mycoplasma contamination in cell lines was regularly monitored using MycoAlert® Mycoplasma
335   Detection Kit (Lonza), following the manufacturer's protocol.

Isolation and purification of cellular trypomastigotes and intracellular amastigotes

Late stationary phase epimastigotes of Dm28c strain were used to infect H9c2 cells for a
primary infection. Six days post-infection, cellular trypomastigotes were obtained from the
supernatant and were used to infect 50% confluent H9c2 cells at a 10:1 rate. Twenty-four hours
post-infection the cell culture was washed twice with PBS 1X to remove any remaining
extracellular parasites and maintained with fresh hgDMEM at 37°C in a humidified 5% CO_2
incubator.

For amastigote purification, infected H9c2 cells incubated for 48 hours post infection were
washed with 1X PBS and incubated with trypsin-EDTA for 5 minutes at 37°C. The trypsinization
was stopped by adding an equal volume of hgDMEM with 10% FBS. The cell suspension was
repeatedly passed through a 27-gauge needle attached to a 30-mL syringe until complete cell
disruption was confirmed under the microscope. The supernatant, containing free amastigotes,
was collected and centrifuged at 500xg for 10 minutes at 4°C to remove large host-cell debris.
The resulting supernatant was then centrifuged at 4,000xg for 10 minutes at 4°C, and the
amastigote-containing pellet was washed twice in chilled 1X DPBS (Dulbecco's Phosphate-
Buffered Saline, No Calcium, No Magnesium) and resuspended in 1X DPBS at 200 µL per $1x10^6$
cells, ready for the fixation step.

Cellular trypomastigotes derived from infected H9c2 cells and present in the cell supernatant
fraction were collected and centrifuged at 500xg for 10 minutes at 4°C to remove large host-cell
debris. The washing and resuspension steps in DPBS were performed as previously described
for amastigotes.

scRNA-seq library preparation and sequencing

Cell fixation was performed using the Methanol Fixation Protocol for Single-Cell RNA
Sequencing (38), after resuspension in DPBS as described above, as recommended by 10X
Genomics technical support. Briefly, chilled 100% Methanol (for HPLC, ≥99.9%, Millipore) was
added drop by drop ($1x10^6$ cells in 800 µl) and incubated at -20°C for 30 min. For rehydration,
fixed cells were first equilibrated at 4°C and then centrifuged at 4,000xg for 5 min at 4°C, the
supernatant was discarded and Wash-Resuspension Buffer (3X SSC in Nuclease-free Water,
0.04% UltraPure Bovine Serum Albumin, 1mM DTT, and 0.2 U/ml RNase Inhibitor) was added to
the pellet. Cell debris and large clumps were eliminated by passing the sample through a 40 µm
Flowmi Cell Strainer.

10X Genomics library preparation was performed at the service provided by the Instituto de
Biología y Medicina Experimental (IBYME, Argentina). The library was sequenced by a service
provider (Macrogen, Korea) in a HiSeq2500 equipment (two lanes), generating
approximately880 million reads of 91 bp. Raw data is available in the SRA
(https://www.ncbi.nlm.nih.gov/sra/) under BioProject PRJNA1200704.

Transcript quantification

T. cruzi 2018 Dm28c genome (9) (release 62, TriTrypDB (39)) and *T. cruzi* Dm28c maxi circle
kDNA sequence (40) were combined and used as reference. To improve the proportion of reads
assigned to genes, 11,362 3'UTR regions of the coding sequences (CDS) were annotated using
peaks2UTR (41). Gene expression quantification was performed by pseudoalignment using
kallisto bustools (42), with the options --filter to remove potential noise from environmental
RNA and --em to apply the Expectation-Maximization (EM) algorithm. Default values were used
for the rest of the parameters. This algorithm outperforms other mapping software in handling
multimapping reads, enabling more accurate quantification of multigene families (42). This
resulted in 321 million reads mapped to the transcriptome.

scRNA-seq data processing and analysis

Count matrices from two technical replicates obtained by sequencing the same library were
merged: the common barcodes across both datasets were retained, and the count matrices
were combined to generate a unified dataset. Subsequently, the following metrics were
calculated: nUMI, nGene, and mitoRatio, as well as the log10 of genes per UMI
(log10GenesperUMI). A filtering criterion was applied, retaining cells with nUMI > 1200, nGene
> 100, log10GenesperUMI > 0.8, and mitoRatio < 0.1. Ribosomal rRNA genes were excluded
from subsequent analyses. Data normalization and scaling were performed using the Seurat R
version 5 package (43), employing NormalizeData and ScaleData functions.

After quality filtering, 3,192 single-cell transcriptomes were retained with an average of 8004
reads mapped per cell. In total, 14,321 genes were detected across all cells (93.5% of the
15,319 annotated protein-coding genes in the *T. cruzi* 2018 Dm28c genome). Per cell, we
observed a mean of 1,088 detected genes and 2,461 UMIs, which corresponds to ~7.1% of the
annotated protein-coding gene.

FindNeighbors function was used to construct a k-nearest neighbours graph of cells using 10
principal components (PCs) and Louvain algorithm was employed for cell clustering using
FindClusters function. Additionally, doublets were identified and removed using the
DoubletFinder R package (44); only 41 doublets were detected, all belonging to cluster 2.

To define marker genes, the FindAllMarkers function from Seurat package was used, selecting
those with an adjusted p-value < 0.05 and a log2 fold change (log2FC) > 1. To validate these

markers, bulk RNA-seq data of Dm28c was incorporated (NCBI BioProject ID PRJNA850400 [24]).
Transcripts were quantified using Kallisto (45) (with -b 100 option to perform 100 bootstraps),
followed by differential expression analysis conducted with Sleuth (46). Genes with an adjusted
p-value < 0.05 and |log2FC| > 0.25 were filtered for further analysis. Finally, the gene IDs of the
Seurat-defined markers were cross-referenced with the IDs of the differentially expressed
genes obtained from the bulk RNA-seq analysis to corroborate stage-specific gene expression.
For 2D visualization of cell clusters and gene markers expression profiles across cells, UMAP
projection was employed (47).

Gene IDs corresponding to multigene families (TcS, MASP, Mucins, GP63, RHS and DGF) were
obtained by text searches using the current genome annotation on the "description" field.
Throughout the manuscript, these genes are referred to as either surface protein genes or
multigene family genes. Single and low copy number genes were defined as those that did not
belong to the latter gene list. For simplicity "single-copy" will be used to refer to these genes
throughout the manuscript. To avoid biases arising from differences in list size between single-
copy and multigene family gene sets, we generated a subsampled single-copy gene list by
randomly selecting the same number of genes as in the multigene family set. The expression
distribution of this subsampled single-copy gene set is similar to that of the full dataset.

Data processing was conducted using R version 4.2.0. Statistical analyses were performed using
the Wilcoxon rank-sum test and p-values < 0.05 were considered statistically significant.

To study expression inequality, we plotted Lorenz curves (48) and applied the Gini index, a
metric originally developed in the field of economics (49). Both metrics serve as a measure of
expression heterogeneity, and we employed it in two distinct contexts: first, to evaluate the
degree of inequality in the distribution of a gene's expression levels across individual cells
(**Figure 2d, Supplementary Figure 1d and Supplementary Table 2**); and second, to assess the
extent to which a given cell expresses individual genes at varying rates (**Figure 4c**).

TcS genomic localization analysis was performed by defining directional gene clusters from the
GFF annotation: protein-coding genes were ordered by coordinate per contig and consecutive
genes on the same strand were grouped into a DGC, with a strand-switch (change from "+" to "-"
or vice versa) marking DGC boundaries. For each gene, we calculated the percentage of its
polycistronic neighbours (genes within the same DGC) that belong to the multigene family gene
set previously defined and compared high-abundance TcS vs low-abundance TcS (sampling n =
30). For statistical comparisons, we assessed the robustness of the analysis by sampling 50
iterations of n = 30 low-abudance TcS and distributions were compared to the high-abundance
set using two-sided Wilcoxon rank-sum tests with Benjamini-Hochberg correction.

Author Contributions

LI: Data Curation, Formal Analysis, Investigation, Methodology, Visualization, Writing-Original Draft Preparation, Writing-Review & Editing; LB: Methodology, Investigation, Writing-Review & Editing; VAC: Methodology, Investigation, Writing-Review & Editing; JG: Methodology, Formal Analysis, Writing-Review & Editing; GR: Methodology, Writing-Review & Editing; VMH: Methodology, Writing-Review & Editing; MAD: Methodology, Resources, Writing-Review & Editing; JSS: Methodology, Resources, Writing-Review & Editing; JDG: Conceptualization, Methodology, Visualization, Writing-Original Draft Preparation, Writing-Review & Editing; PS: Conceptualization, Funding Acquisition, Methodology, Formal Analysis, Project Administration, Resources, Supervision, Validation, Writing-Original Draft Preparation, Writing-Review & Editing

Conflict of interest

The authors declare that they have no conflicts of interest

Financial Support

This project was supported by: CSIC, Universidad de la República, grant number: I+D-2020-505 awarded to PS; LI, LB, JG, MD, JSS and PS received financial support from PEDECIBA. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Tarleton RL. Chagas Disease: A Solvable Problem, Ignored. Trends Mol Med. 2016 Oct 1;22(10):835–8.

2. Martín-Escolano J, Marín C, Rosales MJ, Tsaousis AD, Medina-Carmona E, Martín-Escolano R. An Updated View of the Trypanosoma cruzi Life Cycle: Intervention Points for an Effective Treatment. ACS Infect Dis. 2022 June 10;8(6):1107–15.

3. Clayton C. Regulation of gene expression in trypanosomatids: living with polycistronic transcription. Open Biol. 2019 June 5;9(6):190072.

4. Díaz-Viraqué F, Chiribao ML, Libisch MG, Robello C. Genome-wide chromatin interaction map for Trypanosoma cruzi. Nat Microbiol. 2023 Nov;8(11):2103–14.

5. Lima ARJ, Araujo CB de, Bispo S, Patané J, Silber AM, Elias MC, et al. Nucleosome landscape
reflects phenotypic differences in Trypanosoma cruzi life forms. PLOS Pathog. 2021 Jan
26;17(1):e1009272.

6. Lima ARJ, Silva HG de S, Poubel S, Rosón JN, de Lima LPO, Costa-Silva HM, et al. Open
chromatin analysis in Trypanosoma cruzi life forms highlights critical differences in genomic
compartments and developmental regulation at tDNA loci. Epigenetics Chromatin. 2022 June
1;15(1):22.

7. Beati P, Stepñicka MM, Larrea SCV, Smircich P, Alonso GD, Ocampo J. Improving genome-
wide mapping of nucleosomes in Trypanosome cruzi. PLOS ONE. 2023 Nov
21;18(11):e0293809.

8. Ocampo J, Carena S, López M del R, Vela VS, Zambrano Siri RT, Balestra SA, et al.
Trypanosomatid histones: the building blocks of the epigenetic code of highly divergent
eukaryotes. Biochem J. 2025 Mar 14;482(06):325–40.

9. Berná L, Rodriguez M, Chiribao ML, Parodi-Talice A, Pita S, Rijo G, et al. Expanding an
expanded genome: long-read sequencing of Trypanosoma cruzi. Microb Genomics [Internet].
2018 May 1 [cited 2020 Oct 27];4(5). Available from:
/pmc/articles/PMC5994713/?report=abstract

10.   Belew AT, Junqueira C, Rodrigues-Luiz GF, Valente BM, Oliveira AER, Polidoro RB, et al.
Comparative transcriptome profiling of virulent and non-virulent Trypanosoma cruzi
underlines the role of surface proteins during infection. PLOS Pathog. 2017 Dec
14;13(12):e1006767.

11.   Pech-Canul ÁDLC, Monteón V, Solís-Oviedo RL. A Brief View of the Surface Membrane
Proteins from Trypanosoma cruzi. J Parasitol Res [Internet]. 2017 [cited 2020 Nov 6];2017.
Available from: /pmc/articles/PMC5474541/?report=abstract

12.   Freitas LM, dos Santos SL, Rodrigues-Luiz GF, Mendes TAO, Rodrigues TS, Gazzinelli RT,
et al. Genomic Analyses, Gene Expression and Antigenic Profile of the Trans-Sialidase
Superfamily of Trypanosoma cruzi Reveal an Undetected Level of Complexity. PLOS ONE.
2011;6(10):e25914.

13.   Buscaglia CA, Campo VA, Frasch ACC, Di Noia JM. Trypanosoma cruzi surface mucins:
host-dependent coat diversity. Nat Rev Microbiol. 2006 Mar;4(3):229–36.

14.   Pereira-Chioccola VL, Acosta-Serrano A, de Almeida IC, Ferguson MAJ, Souto-Padron T,
Rodrigues MM, et al. Mucin-like molecules form a negatively charged coat that protects
Trypanosoma cruzitrypomastigotes from killing by human anti-α-galactosyl antibodies. J Cell
Sci. 2000 Apr 1;113(7):1299–307.

15.   De Pablos LM, González GG, Solano Parada J, Seco Hidalgo V, Díaz Lozano IM, Gómez
Samblás MM, et al. Differential Expression and Characterization of a Member of the Mucin-

Associated Surface Protein Family Secreted by Trypanosoma cruzi. Infect Immun. 2011 Sept
16;79(10):3993–4001.

16.   Bartholomeu DC, Cerqueira GC, Leão ACA, daRocha WD, Pais FS, Macedo C, et al.
Genomic organization and expression profile of the mucin-associated surface protein (masp)
family of the human pathogen Trypanosoma cruzi. Nucleic Acids Res. 2009 June
1;37(10):3407–17.

17.   Burle-Caldas G de A, dos Santos NSA, de Castro JT, Mugge FLB, Grazielle-Silva V, Oliveira
AER, et al. Disruption of Active Trans-Sialidase Genes Impairs Egress from Mammalian Host
Cells and Generates Highly Attenuated Trypanosoma cruzi Parasites. mBio. 2022 Jan
25;13(1):e03478-21.

18.   Berná L, Chiribao ML, Greif G, Rodriguez M, Alvarez-Valin F, Robello C. Transcriptomic
analysis reveals metabolic switches and surface remodeling as key processes for stage
transition in trypanosoma cruzi. PeerJ. 2017 Mar 8;2017(3):e3017.

19.   Tonelli RR, Giordano RJ, Barbu EM, Torrecilhas AC, Kobayashi GS, Langley RR, et al. Role
of the gp85/Trans-Sialidases in Trypanosoma cruzi Tissue Tropism: Preferential Binding of a
Conserved Peptide Motif to the Vasculature In Vivo. PLoS Negl Trop Dis. 2010 Nov
2;4(11):e864.

20.   Walzer KA, Kubicki DM, Tang X, Chi JTA. Single-Cell Analysis Reveals Distinct Gene
Expression and Heterogeneity in Male and Female Plasmodium falciparum Gametocytes.
mSphere. 2018 Apr 11;3(2):10.1128/msphere.00130-18.

21.   Ngara M, Palmkvist M, Sagasser S, Hjelmqvist D, Björklund ÅK, Wahlgren M, et al.
Exploring parasite heterogeneity using single-cell RNA-seq reveals a gene signature among
sexual stage *Plasmodium falciparum* parasites. Exp Cell Res. 2018 Oct 1;371(1):130–8.

22.   Reid AJ, Talman AM, Bennett HM, Gomes AR, Sanders MJ, Illingworth CJR, et al. Single-
cell RNA-seq reveals hidden transcriptional variation in malaria parasites. Krzych U, editor.
eLife. 2018 Mar 27;7:e33105.

23.   Howick VM, Russell AJC, Andrews T, Heaton H, Reid AJ, Natarajan K, et al. The Malaria
Cell Atlas: Single parasite transcriptomes across the complete Plasmodium life cycle. Science.
2019 Aug 23;365(6455):eaaw2619.

24.   Vigneron A, O'Neill MB, Weiss BL, Savage AF, Campbell OC, Kamhawi S, et al. Single-cell
RNA sequencing of Trypanosoma brucei from tsetse salivary glands unveils metacyclogenesis
and identifies potential transmission blocking antigens. Proc Natl Acad Sci. 2020 Feb
4;117(5):2613–21.

25.   Hutchinson S, Foulon S, Crouzols A, Menafra R, Rotureau B, Griffiths AD, et al. The
establishment of variant surface glycoprotein monoallelic expression revealed by single-cell

RNA-seq of Trypanosoma brucei in the tsetse fly salivary glands. PLOS Pathog. 2021 Sept
20;17(9):e1009904.

26.   Briggs EM, Rojas F, McCulloch R, Matthews KR, Otto TD. Single-cell transcriptomic
analysis of bloodstream Trypanosoma brucei reconstructs cell cycle progression and
developmental quorum sensing. Nat Commun. 2021 Sept 6;12(1):5268.

27.   Howick VM, Peacock L, Kay C, Collett C, Gibson W, Lawniczak MKN. Single-cell
transcriptomics reveals expression profiles of Trypanosoma brucei sexual stages. PLOS
Pathog. 2022 Mar 7;18(3):e1010346.

28.   Quintana JF, Chandrasegaran P, Sinton MC, Briggs EM, Otto TD, Heslop R, et al. Single
cell and spatial transcriptomic analyses reveal microglia-plasma cell crosstalk in the brain
during Trypanosoma brucei infection. Nat Commun. 2022 Sept 30;13(1):5752.

29.   Catta-Preta CMC, Ghosh K, Sacks DL, Ferreira TR. Single-cell atlas of Leishmania
development in sandflies reveals the heterogeneity of transmitted parasites and their role in
infection. Proc Natl Acad Sci. 2024 Dec 24;121(52):e2406776121.

30.   Abuchery BE, Black JA, Silva MS da. Single-cell transcriptomics reveals hidden
information in trypanosomatids. Trends Parasitol. 2022 Jan 1;38(1):4–6.

31.   Seco-Hidalgo V, De Pablos LM, Osuna A. Transcriptional and phenotypical heterogeneity
of Trypanosoma cruzi cell populations. Open Biol. 2015 Dec 1;5(12):150190.

32.   Luzak V, López-Escobar L, Siegel TN, Figueiredo LM. Cell-to-Cell Heterogeneity in
Trypanosomes. Annu Rev Microbiol. 2021 Oct 8;75(Volume 75, 2021):107–28.

33.   Li Y, Shah-Simpson S, Okrah K, Belew AT, Choi J, Caradonna KL, et al. Transcriptome
Remodeling in Trypanosoma cruzi and Human Cells during Intracellular Infection. PLoS
Pathog. 2016;12(4):1–30.

34.   Laidlaw R*, Garcia Sanchez M*, Pacheco JDS, Paradela LDS, Otto TD, Rycker MD. The
Trypanosoma cruzi cell atlas; a single-cell resource for understanding parasite population
heterogeneity and differentiation [Internet]. bioRxiv; 2025 [cited 2025 Dec 7]. p.
2024.10.01.616042. Available from:
https://www.biorxiv.org/content/10.1101/2024.10.01.616042v2

35.   Schmatz DM, Boltz RC, Murray PK. Trypanosoma cruzi: separation of broad and slender
trypomastigotes using a continuous hypaque gradient. Parasitology. 1983 Oct;87(2):219–27.

36.   Cruz-Saavedra L, Loock M, Antunes LB, Cestari I. Variation in surface protein expression
leads to heterogeneous Trypanosoma cruzi populations during host cell infection. Nat
Commun. 2025 Nov 12;16(1):9949.

37.   Smircich P, Pérez-Díaz L, Hernández F, Duhagon MA, Garat B. Transcriptomic analysis of
572   the adaptation to prolonged starvation of the insect-dwelling Trypanosoma cruzi
573   epimastigotes. Front Cell Infect Microbiol. 2023 Apr 6;13:374.

574   38.   Gutiérrez-Franco A, Ake F, Hassan MN, Cayuela NC, Mularoni L, Plass M. Methanol
575   fixation is the method of choice for droplet-based single-cell transcriptomics of neural cells.
576   Commun Biol. 2023 May 15;6(1):522.

577   39.   Aslett M, Aurrecoechea C, Berriman M, Brestelli J, Brunk BP, Carrington M, et al.
578   TriTrypDB: A functional genomic resource for the Trypanosomatidae. Nucleic Acids Res. 2009
579   Oct 20;38(SUPPL.1):D457.

580   40.   Berná L, Greif G, Pita S, Faral-Tello P, Díaz-Viraqué F, Souza RDCMD, et al. Maxicircle
581   architecture and evolutionary insights into Trypanosoma cruzi complex. PLoS Negl Trop Dis.
582   2021 Aug 26;15(8):e0009719.

583   41.   Haese-Hill W, Crouch K, Otto TD. peaks2utr: a robust Python tool for the annotation of
584   3' UTRs. Bioinformatics. 2023 Mar 1;39(3):btad112.

585   42.   Sullivan DK, Min KH (Joseph), Hjörleifsson KE, Luebbert L, Holley G, Moses L, et al.
586   kallisto, bustools and kb-python for quantifying bulk, single-cell and single-nucleus RNA-seq.
587   Nat Protoc. 2025 Mar;20(3):587–607.

588   43.   Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, et al. Dictionary
589   learning for integrative, multimodal and scalable single-cell analysis. Nat Biotechnol. 2024
590   Feb;42(2):293–304.

591   44.   McGinnis CS, Murrow LM, Gartner ZJ. DoubletFinder: Doublet Detection in Single-Cell
592   RNA Sequencing Data Using Artificial Nearest Neighbors. Cell Syst. 2019 Apr 24;8(4):329-
593   337.e4.

594   45.   Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq
595   quantification. Nat Biotechnol. 2016 May;34(5):525–7.

596   46.   Pimentel H, Bray NL, Puente S, Melsted P, Pachter L. Differential analysis of RNA-seq
597   incorporating quantification uncertainty. Nat Methods. 2017 July;14(7):687–90.

598   47.   McInnes L, Healy J, Saul N, Großberger L. UMAP: Uniform Manifold Approximation and
599   Projection. J Open Source Softw. 2018 Sept 2;3(29):861.

600   48.   Lorenz MO. Methods of Measuring the Concentration of Wealth. Publ Am Stat Assoc.
601   1905;9(70):209–19.

602   49.   Atkinson AB. On the measurement of inequality. J Econ Theory. 1970 Sept 1;2(3):244–63.

Supplementary Figures

Supplementary Figure 1. (a) Summatory of expression levels values from subsampled single-copy genes for each cell from amastigote (Cluster 1) and trypomastigote (Cluster 0) cell populations (**** p < 0.0001, meanAma = 394.9, SDAma = 101.7, meanTrypo = 353.1, SDTrypo = 62.1, FCAma/Trypo = 1.1). (b) UMAP projection for 2D visualization of core gene expression among cells, ; num_multigene indicates the number of multigene family genes detected in each cell, whereas z_multigene indicates the expression levels calculated by summing the UMI counts of all multigene family genes in each cell and then standardizing this value using a z-score transformation, such that positive values reflect above-average multigene family expression and negative values reflect below-average levels. (c) Violin plots showing the number of cells expressing a specific gene belonging to each group of genes: subsampled single-copy and multigene families, ribosomal genes and different multigene families. (d) Lorenz curves showing the cumulative proportion of gene expression relative to the cumulative proportion of genes for subsampled single-copy, multigene family genes (together or grouped by multigene family) and ribosomal protein coding genes. Genes were ordered by total expression, and the dashed line indicates perfect equality. Curves that deviate further from the diagonal reflect greater inequality, meaning that fewer genes account for most of the

expression within each category. Statistically significant differences between groups for c) and
d) are shown in **Supplementary Table 2**.

Supplementary Figure 2.

Trypomastigote sub-clusters identified based on trans-sialidase expression profiles. Violin plots
displaying average expression levels across sub-clusters and associated fold changes
($FC_{trypo\_0/trypo\_1}$) of (a) transporters coding genes, (b) DNA and RNA polymerase-associated
protein coding genes, (c) phosphatases coding genes and (d) multigene family genes. **** p <
0.0001.

Supplementary Figure 3. (a) Venn diagram showing the overlap between the top 100 most
expressed TcS from bulk RNA-seq data and TcS expressed in more than 40% of cells from cluster
trypo_0, and (b) Heatmap displaying the expression of TcS genes in each cell that together
account for 75% of total TcS gene expression and are expressed in more than 40% of cells
within cluster trypo_0. Cells are clustered by TcS expression profiles, with colors representing
each gene's percentage contribution to the cell's expression.

641