GCN5a is a telomeric lysine acetyltransferase whose loss primes

***Toxoplasma gondii* for latency**

Vishakha Dey[1], Michael J. Holmes[1], Sandeep Srivastava[2], Emma H. Wilson[2], William J.

Sullivan Jr.[1#]

[1]Department of Microbiology & Immunology, Indiana University School of Medicine,

Indianapolis, IN

[2]Division of Biomedical Sciences, School of Medicine, University of California,

Riverside, Riverside, CA

#Corresponding author: wjsulliv@iu.edu

Running title: Loss of telomeric GCN5a primes *Toxoplasma* for latency

Word count (abstract): 207

Word count (text): 4971

ABSTRACT

Toxoplasma gondii is a protozoan parasite that causes persistent infection in

warm-blooded vertebrates by undergoing differentiation from a replicative stage

(tachyzoites) to a latent encysted stage (bradyzoites). Stage differentiation is critical for

transmission and pathogenesis and relies on gene regulation driven by a network of

transcription and epigenetic factors. We previously found in non-cystogenic type I RH

strain parasites that the lysine acetyltransferase (KAT), GCN5a, is dispensable in

tachyzoites but required to upregulate stress-response genes, suggesting a link with

bradyzoite conversion. To address this possibility, we generated endogenously tagged

GCN5a parasites and a genetic knockout in cystogenic type II Pru strain. We show that

GCN5a protein, but not mRNA, increases during differentiation and complexes with

unique protein partners, most of which contain AP2 domains. Pru strain tachyzoites

lacking GCN5a augment bradyzoite-specific gene expression in the absence of stress.

Loss of *GCN5a* slowed tachyzoite replication and heightened sensitivity to bradyzoite

conversion but resulted in smaller cyst sizes compared to wildtype. Using CUT&Tag, we

delineated the chromosomal occupancy of GCN5a relative to the essential KAT,

GCN5b. While GCN5b localizes to coding regions, GCN5a surprisingly localizes

exclusively to telomeres. These findings suggest that the loss of GCN5a leads to

telomere dysfunction, which slows replication and promotes the transition to latency.

IMPORTANCE

Toxoplasma gondii is a single-celled parasite that persists in warm-blooded

hosts, including humans, because it converts into latent tissue cysts. Switching from its

replicating form into dormant cysts is a tightly regulated process that involves epigenetic

factors such as lysine acetyltransferases GCN5a and GCN5b. This study is the first to

examine the role of GCN5a in a cyst-forming *Toxoplasma* strain. We found that GCN5a

protein, but not mRNA, increases during cyst development. Additionally, parasites

lacking GCN5a replicate more slowly and are quicker to form cysts when stressed. We

show that GCN5a and GCN5b work in different multi-protein complexes and localize to

different areas of the genome; while GCN5b targets promoters of gene coding regions,

GCN5a is exclusively found at telomeric regions. Our findings suggest a novel role for

GCN5a in telomere biology that, when depleted, produces a fitness defect that favors

development of latent stages.

INTRODUCTION

Toxoplasma gondii is an obligate intracellular parasite that causes opportunistic infection. The life cycle is comprised of an asexual stage that occurs in the nucleated cells of warm-blooded animals and a sexual stage that takes place exclusively in felines (1). The sexual stage culminates in the release of oocysts that remain infectious for up to a year or more in the environment (2). Upon infection, tachyzoites reproduce asexually and disseminate throughout the host's body. Tachyzoites can cross the placental barrier and infect the fetus, causing miscarriage or congenital birth defects (3). Toxoplasma evades host immune clearance by converting into latent bradyzoites that reside inside infected cells in the form of tissue cysts, establishing chronic toxoplasmosis (4). Tissue cysts are resistant to frontline drug treatments, allowing the parasite to persist within an infected host for life. If the host becomes immunocompromised, bradyzoite cysts can reactivate into proliferating tachyzoites that can cause life-threatening tissue destruction. Tissue cysts are another crucial means of parasite spread as naïve hosts can become infected after consuming contaminated meat (5). Given the importance of bradyzoites in parasite pathogenesis and transmission, understanding the molecular events coordinating stage switching remains an intense area of investigation.

The conversion of tachyzoites into bradyzoites requires alterations in gene expression, and several transcription factors mediate bradyzoite formation to varying degree. Toxoplasma contains 67 genes possessing a plant-like DNA-binding domain called Apetela-2 (AP2), several of which impact tachyzoite-bradyzoite conversion (4, 6). The master regulator of bradyzoite differentiation harbors a myb-like DNA-binding

domain and is designated "bradyzoite-formation deficient 1" (*BFD1*). BFD1 binds to the

promoters of stage-specific genes that are upregulated in bradyzoites, and parasites fail

to initiate bradyzoite differentiation when *BFD1* is ablated (7).

Epigenetic machinery and histone modifications are also important for gene

expression changes required for stage conversion, including histone acetylation

mediated by general control non-repressed 5 (GCN5) proteins (4). Most early-branching

eukaryotes possess a single GCN5 lysine acetyltransferase (KAT) that modulates

stress responses (8, 9). In contrast, *Toxoplasma* contains two GCN5 KATs, designated

GCN5a and GCN5b. To date, these GCN5 KATs have only been studied in type I RH

parasites, which have largely lost their ability to readily convert to bradyzoites (10).

While *GCN5b* is essential for parasite viability and regulates housekeeping genes,

GCN5a is dispensable in tachyzoites (11, 12). Using a microarray for transcriptional

profiling, RH parasites lacking GCN5a failed to upregulate ~75% of the genes normally

activated during alkaline stress (12).

As type I RH strain has largely lost its developmental capacity, it is not ideal for

the study of factors involved in bradyzoite conversion. To address the potential role of

GCN5a in stress-induced bradyzoite differentiation, we generated a tagged line and a

genetic knockout in the cystogenic type II Prugniaud (Pru) strain. In contrast to RH

strain, type II parasites lacking GCN5a displayed a reduction in replication and

heightened sensitivity to stress-induced bradyzoite differentiation, although cyst sizes

remained smaller than those made by wildtype. Transcriptional profiling of unstressed

parasites revealed abnormal expression of bradyzoite genes in tachyzoites. We

performed CUT&Tag (Cleavage Under Targets and Tagmentation) on both GCN5a and

GCN5b, showing that GCN5a is exclusively associated with telomeric regions.

Together, these findings suggest that loss of *GCN5a* likely leads to telomere

dysfunction, thereby producing replicative delays and cellular stress that sensitizes the

parasite towards bradyzoite conversion.

RESULTS AND DISCUSSION

GCN5a protein, but not mRNA, increases during bradyzoite differentiation

We previously generated a *GCN5a* knockout in type I RH *Toxoplasma*, finding

that this lysine acetyltransferase (KAT) is dispensable for tachyzoite viability; however,

parasites lacking GCN5a were impaired in their ability to upregulate 74% of the genes

normally activated after 4 days in alkaline stress media, which is commonly used to

induce tissue cyst formation in vitro (12). These results suggested that GCN5a is a

stress-responsive KAT that may contribute to bradyzoite differentiation (13). Since type I

parasites do not develop into mature tissue cysts at high frequency, we sought to

characterize the role of GCN5a in a cystogenic type II strain. Using a CRIPSR/Cas9

approach, we first tagged endogenous *GCN5a* at its C-terminus with a 3xHA epitope in

type II Prugniaud (Pru) parasites lacking KU80 and HXGPRT (Fig. S1A). GCN5a^{HA} was

not detectable in unstressed tachyzoites but could be visualized by western blot and

immunofluorescence assay (IFA) following alkaline stress for 24 h (Fig. S1B, C). IFA of

stressed parasites showed that GCN5a^{HA} localizes exclusively to the parasite nucleus in

the Pru strain (Fig. S1B). Western blotting of GCN5a^{HA} parasites revealed a band at

130 kDa, the expected size for the epitope-tagged protein (Fig. S1C). To ensure fidelity

of the integration site, GCN5a^HA clones were further confirmed by PCR of genomic DNA

(Fig. S1D).

Previous work on GCN5a in RH strain employed an ectopically expressed

recombinant form of the protein, so endogenous expression patterns have yet to be

evaluated (12). Therefore, we monitored mRNA and protein levels before and after a

bradyzoite-inducing stress in our GCN5a^HA parasites. While *GCN5a^HA* mRNA levels

remain unchanged (Fig. 1A), GCN5a^HA protein expression is enhanced in response to

alkaline stress (Fig. 1B). These results suggest that GCN5a protein is stabilized during

alkaline stress, or its mRNA is subject to stress-induced preferential translation during

bradyzoite conversion like *BFD1* and *BFD2/ROCY1* transcripts (7, 14-16). Consistent

with this latter idea, the *GCN5a* mRNA contains a long and complex 5'-leader sequence

that contains 26 upstream open reading frames (uORFs), features that have been

linked to translational control during phosphorylation of eIF2α (17). In addition, a

genome-wide RNA CLIPseq analysis identified BFD2-binding sites within the 5'-leader

of *GCN5a* mRNA (16), which we've shown to be associated with cap-independent

mechanisms of translational control for *BFD1* (15).

Cystogenic parasites lacking GCN5a display reduced replication

We next generated a *GCN5a* knockout (ΔGCN5a) in the GCN5a^HA background.

We replaced the entire GCN5a^HA coding sequence with an HXGPRT minigene cassette

using a dual guide CRISPR/Cas9 strategy (Fig. S2A). The selected ΔGCN5a clone

displayed no HA signal by IFA or western blot under stress conditions (Fig. S2B, C).

The ΔGCN5a parasites were further validated by PCR of genomic DNA (Fig. S2D).

Previously, loss of GCN5a had no discernible effect on replication in type I RH

tachyzoites (10). To determine whether the loss of GCN5a affected the viability of type II

parasites, we conducted a plaque assay comparing GCN5a^{HA} and ΔGCN5a parasites

under tachyzoite culture conditions. After 14 days, we found that ΔGCN5a parasites

formed smaller plaques relative to GCN5a^{HA} parasites (Fig. 2A-B), although there was

no difference in the number of plaques (Fig. 2C). These data suggest that the loss of

GCN5a does not impact parasite invasion into host cells but does affect replication rate.

To examine this potential growth defect in ΔGCN5a parasites more closely, we

performed a parasite replication assay. Parasites were allowed to grow for 16 hours and

the number of parasites per vacuole was quantified. In contrast to the parental

GCN5a^{HA} parasites, ΔGCN5a parasites contained more vacuoles with 2 parasites and

fewer vacuoles with 4 parasites, confirming that the replication rate is slower for

ΔGCN5a parasites (Fig. 2D). We further ensured that tagging of GCN5a did not affect

the growth rate by comparing the PruΔKu80ΔHX parental strain with GCN5a^{HA} parasites

using plaque and replication assays (Fig. S3). These results indicate that GCN5a

contributes to parasite replication in type II cystogenic strains, unlike RH strain,

suggesting strain-specific variations in the function of GCN5a.

Increased expression of bradyzoite-associated genes in unstressed ΔGCN5a

parasites

To further investigate the impaired replication phenotype observed for ∆GCN5a

parasites, we performed differential expression transcriptional profiling of the parental

GCN5a^{HA} (WT) and ∆GCN5a parasites (Table S2). Both upregulated a similar set of

bradyzoite-specific genes in response to alkaline stress, including canonical markers

BAG1, *ENO1*, and *LDH2* (Table S2; Fig 3A-B). However, the presence of GCN5a

impacts the degree of the transcriptional response to stress. Although most genes were

regulated in the same direction following bradyzoite induction, the degree to which

these bradyzoite-associated genes are induced is blunted in the ∆GCN5a parasites

compared to WT (Fig. 3B). For example, the log$_2$ fold-change that occurs under stress

for *ENO1* is >9 in WT but only ~6 for ∆GCN5a parasites. The change in *BAG1* and

LDH2 expression is also much more pronounced in stressed GCN5a^{HA} parasites

compared to the knockout, suggesting that the loss of GCN5a produces a muted stress

response.

Since a blunting of the stress response could either indicate a decrease in the

induction of bradyzoite-associated genes or a change in basal gene expression in

tachyzoites, we compared gene expression in GCN5a^{HA} and ∆GCN5a tachyzoites

(Table S2; Fig 3C). Supporting the latter possibility, genes associated with early

bradyzoite formation (18), such as *ENO1* and *BAG1*, were upregulated in unstressed

∆GCN5a tachyzoites. In contrast, genes that are turned on later during bradyzoite

formation (18), such as *LDH2*, are unchanged in unstressed ∆GCN5a parasites. In fact,

most (66 of 77) genes that displayed elevated basal expression in ∆GCN5a compared

to WT tachyzoites were upregulated in both strains upon treatment with alkaline stress.

Furthermore, elevated expression of early bradyzoite genes in the absence of stress is

consistent with the slowed replication of ΔGCN5a parasites and supports the idea that

ΔGCN5a tachyzoites may be poised for bradyzoite formation.

Counterintuitively, these findings are consistent with our previous observations in

type I parasites lacking GCN5a. At the time, we interpreted a muted induction of

bradyzoite-associated gene expression in response to alkaline treatment as a failure to

initiate bradyzoite formation (12). Given our findings here, it is more likely that GCN5a

deficiency in type I parasites also leads to increased expression of bradyzoite-

associated genes in tachyzoites, blunting the response to alkaline stress, all while

maintaining the rapid growth that characterizes of type I strains.

Loss of GCN5a sensitizes tachyzoites towards bradyzoite conversion and results

in reduced tissue cyst size

Bradyzoite conversion can be triggered through slowed parasite replication (19).

In addition, our transcriptomic analysis revealed enhanced expression of several

bradyzoite genes in ΔGCN5a tachyzoites, suggesting they may be predisposed towards

differentiation. We infected HFF host cells with ΔGCN5a or WT (GCN5a[HA]) parasites

and initiated bradyzoite differentiation using alkaline stress. The frequency of tissue cyst

formation was monitored every 24 hours by staining for the cyst wall with *Dolichos*

biflorus agglutinin (DBA). We observed that ΔGCN5a parasites differentiated more

rapidly into cysts at 24 and 48 hours relative to WT (Fig. 4A left panel). By 72 hours, the

differentiation rate of WT parasites had caught up to the ΔGCN5a parasites.

Quantification of average tissue cyst sizes revealed that those made by ΔGCN5a

parasites were significantly smaller than those made by WT (Fig. 4A right panel). A

representative IFA showing the smaller cyst sizes of ΔGCN5a parasites is shown in

Figure 4B. These results confirm that the loss of GCN5a in a cystogenic strain makes

tachyzoites more prone to differentiation and shows that the slowed replication

produces smaller cyst sizes.

Differences between GCN5a and GCN5b complexes

To shed further light on the function of GCN5a, we sought to elucidate its

interacting proteins using co-immunoprecipitation (co-IP). As we could not obtain

sufficient material using Pru GCN5a^{HA} parasites, we recreated this line in

RHΔKu80ΔHX, which yields a higher biomass than Pru (Fig. S4A). We previously

employed a similar strategy to characterize the GCN5b interactome (20). IFA confirmed

that GCN5a^{HA} was localized to the nucleus as expected (Fig. S4B) and immunoblotting

confirmed the presence of the endogenously tagged GCN5a^{HA} protein at the expected

molecular weight of 130 kDa, increasing after stress as seen in Pru (Fig. S4C). Proper

insertion of the tag at the *GCN5a* locus was further confirmed by PCR analysis of

genomic DNA (Fig. S4D).

Results of three independent pull-downs of RH GCN5a^{HA} parasites under

unstressed or alkaline stress conditions are shown in Table 1. Speaking to the fidelity of

the pull-down data, we identified the coactivator protein ADA2-B (AP2VIIb-1), which we

previously determined to interact with GCN5a and not GCN5b using a yeast two-hybrid

assay (10). In contrast, GCN5b associates with exclusively with ADA2-A (10, 20).

We also found that GCN5a[HA] pulls down with a PHD-finger domain-containing

protein (TGGT1_234900), the putative cell cycle-associated SRPK (serine-arginine

protein kinase), and several AP2 factors: AP2XII-4, AP2X-5, and AP2V-1. (The

nanopore sequencing data available on ToxoDB (21) and nanopore sequencing data

(22) indicate that TGGT1_220520 and TGGT1_220530 are misannotated as two genes

and both should be attributed to AP2V-1). The PHD-finger protein and AP2XII-4 are

also components of the GCN5b complex, but the others are unique to the GCN5a

complex. Both GCN5 complexes contain a protein kinase, but they differ: GCN5a pulls

down with SRPK whereas GCN5b associates with GSK (20, 23).

The associated components of the *Toxoplasma* GCN5a complex are novel and

not conserved with other eukaryotic GCN5 complexes. The majority of the complex is

composed of AP2 factors of unknown function. *Toxoplasma* possesses 67 AP2 domain-

containing proteins, some of which have been linked to gene regulation while others to

mitoribosome formation via RNA-binding activities (24). While ADA2-B is a homologue

of the ADA2 coactivator known to enhance binding of acetyl-CoA to GCN5 for histone

acetylation (25), the unique presence of an AP2 domain speaks to another possible

function. The other AP2 factors contain no additional domains that provide clues to

function with exception of AP2V-5. AP2V-5 harbors a WHIM1 domain, which has been

implicated in chromatin remodeling as it pertains to nucleosome spacing (26). Notably

absent from the GCN5a complex are proteins unequivocally associated with

transcriptional activity, such as FACT (facilitates chromatin transcription) and TAF

proteins that are found in the GCN5b complex (11, 20).

As one might expect, peptide counts increase following alkaline stress and the

ensuing increase in GCN5a protein abundance. This represents a notable difference

from the significant reorganization of the GCN5b complex during alkaline stress, which

loses interactions with most of its AP2 factors, including AP2XII-4 (20). These findings

are consistent with the model that the GCN5a complex is mobilized during stress (12)

and support that GCN5a and GCN5b evolved to carry out distinct functions in the

parasite.

Chromosomal occupancy of GCN5a and GCN5b

Given the differences in complex composition and phenotype between the two

GCN5s, we performed a comparative analysis of genome occupancy for each under

stressed and unstressed conditions. To make a line comparable to Pru GCN5a[HA], we

engineered PruΔKu80ΔHX parasites to express endogenous GCN5b tagged with HA at

its C-terminus (Fig. S5A). Immunoblotting confirmed the presence of the tagged

GCN5b[HA] protein at the expected molecular weight of 115 kDa (Fig. S5B). IFA

demonstrated the expression of GCN5b[HA] in the nucleus as expected (Fig. S5C). The

accurate tagging of the *GCN5b* gene was further validated through PCR analysis (Fig.

S5D). In contrast to GCN5a, we noted that protein levels for GCN5b do not change in

response to alkaline stress (Fig. S5E).

CUT&Tag profiling revealed that GCN5b associates with the transcriptional start

sites of protein coding genes but GCN5a does not (Fig. 5A). Surprisingly, GCN5a is

robustly enriched at chromosomal ends over the repetitive sequence characteristic of

telomeres (Fig. 5B). Moreover, the association of GCN5a with telomeres becomes more

pronounced in the 48 h stress samples when GCN5a protein levels increase relative to

the unstressed conditions. We did not detect GCN5b at telomeric regions in either

culture condition.

To confirm the association between GCN5a and telomeres, we performed

fluorescent in situ hybridization (FISH) using a telomeric probe in GCN5a^{HA} parasites

(Fig. 6A). Co-localization analysis revealed partial overlap between GCN5a^{HA} and

telomere signals, consistent with the CUT&Tag data (Fig. 6B). Five additional nuclei

were examined, yielding similar results (Fig. S6). To further confirm this interaction, we

conducted an independent ChIP of GCN5a^{HA} (and parental parasites as a negative

control) followed by PCR using primers that amplify the genomic segment spanning the

subtelomeric region of chromosome ChrIb into its telomere. In agreement with the

CUT&Tag data, the ChrIb telomeric region was detected by PCR following ChIP of

GCN5a^{HA} (Fig. 6C). Also consistent with CUT&Tag data, we could not amplify *tubulin* or

B1 gene sequences following ChIP of GCN5a^{HA}, supporting that GCN5a localizes

specifically to telomeric regions and not coding sequences. These experiments

collectively demonstrate that GCN5a localizes to telomeric regions and affirms GCN5b

operates at coding sequences.

Concluding remarks

Toxoplasma is unusual among early-branching eukaryotes and fellow

apicomplexan parasites in possessing two GCN5 KATs. We previously postulated that

the non-essential KAT GCN5a resulted from a gene duplication of the essential KAT

GCN5b (10). The present study shows that each GCN5 family member operates in a

separate multi-protein complex to carry out distinct cellular functions. GCN5b fits the

profile of a standard eukaryotic KAT, partnering with proteins directly linked to

transcriptional activity (11, 20) and localizing to gene coding regions. Aside from

AP2XII-4 and the PHD-finger protein TGGT1_234900, GCN5a partners with unique

proteins and resides at telomeric regions. In addition, GCN5b protein levels do not

increase during stress whereas GCN5a protein levels do.

A model fitting our findings as well as those from previous studies suggests that

GCN5a evolved to promote telomere integrity, hence its loss generates a stress that

slows replication and primes the parasite for latency as a survival strategy. Evidence

from other species lends support to this model. In mammals, GCN5 interacts with

proteins at telomeric regions and its ablation in mouse or human cells leads to telomere

dysfunction (27, 28). Lysine acetylation is a means by which human telomeric protein

TRF2 is stabilized (29). The GCN5-family KAT PCAF facilitates the resolution of R-

loops that can accumulate at telomeres (30). Our findings open a new avenue for

continued interrogation of *Toxoplasma* GCN5a and how it may participate in telomere

biology.

MATERIALS AND METHODS

Host cell and parasite culture

Toxoplasma gondii PruΔKu80ΔHX (31) and RHΔKu80ΔHX (32) strains parasites

were cultured in confluent human foreskin fibroblast (HFF) cells (ATCC, SCRC-1041)

containing Dulbecco's modified Eagle medium (DMEM) (GIBCO, 10-017-CM)

supplemented with 5% inactivated fetal bovine serum (FBS) (R&D systems, S11150).

Parasite cultures were maintained in a humidified 5% CO_2 incubator at 37°C. HFF cells

were grown in DMEM supplemented with 10% inactivated FBS. Alkaline medium (pH

8.2) for bradyzoite culture conditions contained RPMI 1640 (GIBCO, 31800-022)

supplemented with 5% FBS and 50 mM HEPES. Parasites in alkaline medium were

grown in ambient CO_2 and the medium was changed daily.

Transgenic parasite generation

Primers for this study were ordered from Integrated DNA Technologies (Table

S1). GCN5a (TGME49_254555) and GCN5b (TGME49_243440) were endogenously

tagged with a triple hemagglutinin (3xHA) epitope tag at their C-termini using

CRISPR/Cas9 and double homologous recombination. A single guide RNA (sgRNA)

sequence near the stop codon was chosen using the EuPaGDT tool (33) and the

sequence was cloned into pCas9-GFP plasmid (34), designated as pCas9-GCN5a/b. A

donor amplicon with homologous regions to upstream of the stop codon and 3'-UTR of

the gene was PCR amplified to incorporate 3xHA sequence and the DHFR*-TS

selectable marker (35).The Cas9 plasmid and the donor amplicon were co-transfected

using a Lonza Biosciences Nucleofector and cultured for three passages in medium

containing 1.0 µM pyrimethamine (Sigma, SML3579) and cloned by limiting dilution. The

modified locus of the transgenic clone was validated by PCR analyses. The resulting

parasite lines generated in Pru are labeled GCN5a^{HA} and GCN5b^{HA}, while the line

generated in RH is denoted as RH-GCN5a^{HA}.

To generate the knockout GCN5a, two gRNAs were designed to replace the

entire *GCN5a* coding sequence and DHFR*-TS cassette in GCN5a^{HA} parasites using a

strategy that has been outlined previously (36). Both gRNAs were cloned into pCas9-

GFP plasmid (34) to target immediately upstream of the *GCN5a* coding sequence

(pCas9-gRNA1) and downstream of the *DHFR*-TS* 3'-UTR (pCas9-gRNA2). The

resulting plasmid was denoted pCas9-dual-Guide. A double-stranded donor containing

an HXGPRT selectable marker (37) flanked by homologous regions to the 5'- and 3'-

UTRs of *GCN5a* was PCR amplified. Parasites were co-transfected with pCas9-dual-

Guide and the repair template in GCN5a^{HA} parasites and selected for three passages in

medium containing 25 µg/ml mycophenolic acid (Sigma, 475913) and 50 µg/ml xanthine

(Sigma, X7375) and cloned by limiting dilution. The resulting parasite line was validated

with genomic PCRs and is termed ΔGCN5a.

Parasite growth assays

Standard growth assays were conducted as previously described (38). Briefly, for

plaque assays, 500 parasites were used to infect confluent HFF cells grown on 12-well

plates in tachyzoite medium. Plates were incubated undisturbed for 14 days until

plaques were formed, which were visualized by crystal violet staining. Plaques were

counted and the area was measured by ImageJ.

For replication assays, parasites were allowed to invade confluent HFFs for 4 h

in tachyzoite medium. After invasion, the medium was replaced to wash away

extracellular parasites. The number of parasites per vacuole was counted under light

microscopy at the indicated time points.

Immunofluorescence assay (IFA)

Parasites were allowed to infect confluent HFF cell monolayers on coverslips,

cultured under tachyzoite or bradyzoite conditions with media changes every 24 h.

Coverslips were fixed with 4% paraformaldehyde (Sigma, P6148), washed with PBS,

and treated with blocking buffer (3% BSA, Sigma, A9418) and 0.2% Triton X-100

(Sigma, 93443) for permeabilization. Primary antibody incubation (1:1,000 rat anti-HA,

Roche, 11867423001) was performed at 4°C for 16 h. After washing, cells were

incubated with secondary antibody (1:5,000 Alexa Fluor 598 anti-rat, Thermo Fisher, A-

11007) and 1 μg/ml DAPI (Invitrogen, D1306) for 1 h in the dark. For cyst walls, 1:500

FITC-conjugated *Dolichos biflorus* lectin (Vector Laboratories, FL1031) was added.

After washing, coverslips were mounted with ProLong Gold Antifade (Invitrogen,

P36930) and left to set overnight before imaging with a Nikon Eclipse E100080i

microscope.

Western blotting

Parasites were lysed with RIPA buffer supplemented with protease and

phosphatase inhibitor cocktail (Thermo Fisher, 78425). The lysate was sonicated,

centrifuged at 13,000×g for 2 min, and the supernatant was boiled for 10 min with 1X

SDS-PAGE loading buffer and β-mercaptoethanol (Sigma, M3148). Samples were

separated on NUPAGE 4-12% Bis-Tris gels (Invitrogen, NP0326BOX) and transferred

to nitrocellulose membrane (Cytiva, 10600001). Blots were blocked in 5% non-fat

milk/TBST for 1 h at room temperature, incubated with primary antibody at $4^{o}C$ for 16 h

using 1:1,000 dilution rat anti-HA (Roche, 11867423001) for HA-tagged proteins and

1:1,000 rabbit anti-BAG1 polyclonal antibodies (39) (provided by Vern Carruthers,

University of Michigan) or 1:5,000 rabbit anti-tubulin polyclonal antibodies (provided by

David Sibley, Washington University). After washing the blot with TBST, secondary

antibody was applied using the appropriate HRP-conjugated antibodies (GE healthcare)

at a 1:5,000 dilution. Blots were imaged using SuperSignal West Femto (Thermo Fisher,

34095) and detected via Bio-Rad ChemiDoc Imaging System.

Real-time reverse transcription PCR (qRT-PCR)

Confluent HFF cells in T-25 flasks were infected with 10^5 parasites. After 4 h, the

medium was replaced with tachyzoite or bradyzoite medium. Total RNA was isolated

using TRIzol-chloroform (Invitrogen, 10296028). cDNA synthesis was performed with

SuperScript IV First-Strand cDNA synthesis system (Invitrogen, 18091300). cDNA was

diluted 1:100 and used for real-time PCR with SYBR Green PCR Master Mix (Applied

Biosystems, 4368708) on an Applied Biosystems QuantStudio 5 system. β-tubulin

(TGME49_266960) was used for normalization. We verified that β-tubulin mRNA

expression did not change under alkaline stress when either actin (TGME49_209030) or

GAPDH (TGME49_289690) was used for normalization. Primers are listed in Table S1.

Stage-specific RNAseq

An equal number of parasites (10^6) were used to infect a confluent HFF cell

monolayers in T-75 flasks for 4 h. The media were then changed to tachyzoite or

bradyzoite medium and cultured for 48 h. Parasites were harvested by syringe

passaging followed by filtration. Total RNA was isolated using TRIzol-chloroform and

polyA-enriched RNAseq libraries were generated by Azenta Life Sciences using

standard Illumina protocols. Libraries were sequenced 2x150 bp. Sequencing reads

were trimmed of adapter sequences with Cutadapt (40) and mapped to the *Toxoplasma*

genome (v55) (21) with HISAT2 (41). Differential gene expression was assessed with

DESeq2 (42) after obtaining read counts with HTseq-count (43). Data pertaining to this

experiment can be obtained at GSE286090 (reviewer token: mtgzmkkopxqtpyn).

Immunoprecipitation of HA-tagged proteins

RH-GCN5a^{HA} and untagged RH parasites were cultured under both no stress

and 48 hr alkaline stress conditions. Due to the low expression of GCN5a, we used four

T-125 flasks for each biological replicate, with three biological replicates per sample.

Whole cell lysates were prepared by incubating the parasites in lysis buffer (50 mM Tris-

HCl pH 7.4, 150 mM NaCl, 1 mM MgCl$_2$, 0.5% NP-40, 10% glycerol). After clarifying the

lysates via centrifugation at 21,000 xg at 4°C, the lysate was precleared by incubating

with mouse IgG magnetic beads (Cell Signaling, 5873S) for 4 h at 4°C with rotation. The

supernatant was transferred onto α-HA magnetic beads (Thermo Fisher, 88836), which

were pre-washed with lysis buffer, and incubated overnight with rotation at 4°C. Beads

were washed with wash buffer (50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1 mM $MgCl_2$,

0.05% NP-40, 10% glycerol) and with PBS and submitted for MS at the Proteomics

Core Facility at IUSM as described before (44). Data were filtered to include only

proteins that were not detected in the parental strains and were identified by at least 5

peptides in two out of three replicates.

CUT&Tag

CUT&Tag was performed as described previously (45). Briefly, GCN5a^{HA},

GCN5b^{HA}, and untagged PruΔKu80ΔHX parental parasites were grown in confluent

HFF monolayers for 48 h in tachyzoite or bradyzoite conditions. Parasites were isolated

by syringe passage followed by filtration. Samples were processed using the CUT&Tag-

IT assay kit (Active Motif, 53160) according to the manufacturer's instructions with a

primary antibody directed against the HA tag (Invitrogen MA5-27915). Sequencing

reads were aligned to the *Toxoplasma* genome (v67) using bowtie2 allowing for

dovetailing (46). Chromosome occupancy was visualized using the IGV genome

browser (47). Data pertaining to this experiment can be obtained at GSE286088

(reviewer token: cvizwioknfepbmv).

Chromatin immunoprecipitation (ChIP)

GCN5a[HA] and untagged parental Pru parasites were cultured under alkaline

stress for 48 h. Three independent biological samples were processed simultaneously

for ChIP analysis. Parasites were fixed with 4% paraformaldehyde (Sigma, P6148) for

15 min, followed by PBS washes. Lysates were prepared as above for

immunoprecipitation. Ten percent of the lysate was reserved for input. The remaining

lysate was incubated onto α-HA magnetic beads and washed as above for

immunoprecipitation. DNA was extracted from ChIP and input samples using phenol-

chloroform (Invitrogen, 15593031) and precipitated with ethanol. qPCR was performed

in duplicate, with Ct values normalized to input. Subtelomere-specific forward and

telomere-specific reverse primers were used for analysis (Table S1).

Fluorescence *in situ* hybridization of telomeres

Parasites were inoculated onto confluent HFFs and grown under bradyzoite-

inducing conditions for 3 days. Monolayers were fixed in 4% paraformaldehyde for 15

min and incubated overnight at 4^{o}C in 70% EtOH. Coverslips were processed as

described in (48). Briefly, samples were denatured in 0.1N HCl for 15 min, then

neutralized in 2X saline sodium citrate (SSC) (Corning, 46-020-CM) for 5 min before

being placed in equilibration buffer (2X SSC, 50% formamide) for 30 min. Coverslips

were incubated in hybridization buffer (2X SSC, 50% formamide, 15% dextran sulfate,

1% Tween-20) containing 2 μM of a telomeric probe (TTTAGGG$_{X4}$-Cy5) overnight at

37°C. Coverslips were sequentially rinsed in 2X SSC, 1X SSC, 0.1X SSC, and PBS for

5 minutes then processed for IFA using anti-HA as above.

ACKNOWLEDGEMENTS

This research was supported by grants from the National Institutes of Health

(AI152583 and AI167662 to WJS). The authors thank Drs. Vern Carruthers (University

of Michigan) and David Sibley (Washington University) for generously supplying

antibodies used in this study.

The mass spectrometry work performed in this work was done by the Indiana

University School of Medicine Proteomics Core. Acquisition of the IUSM Proteomics

core instrumentation used for this project was provided by the Indiana University

Precision Health Initiative. The proteomics work was supported, in part, by the Indiana

Clinical and Translational Sciences Institute (funded in part by Award Number

UL1TR002529 from the NIH, National Center for Advancing Translational Sciences,

Clinical and Translational Sciences Award) and, in part, by the IU Simon

Comprehensive Cancer Center Support Grant (Award Number P30CA082709 from the

NCI).

REFERENCES

1.   Dubey JP. 2009. History of the discovery of the life cycle of Toxoplasma gondii.

Int J Parasitol 39:877-82.

2.   Shapiro K, Bahia-Oliveira L, Dixon B, Dumetre A, de Wit LA, VanWormer E,

Villena I. 2019. Environmental transmission of Toxoplasma gondii: Oocysts in

water, soil and food. Food Waterborne Parasitol 15:e00049.

3.   Bollani L, Auriti C, Achille C, Garofoli F, De Rose DU, Meroni V, Salvatori G,

490   Tzialla C. 2022. Congenital Toxoplasmosis: The State of the Art. Front Pediatr

491   10:894573.

492   4.   Jeffers V, Tampaki Z, Kim K, Sullivan WJ, Jr. 2018. A latent ability to persist:

493   differentiation in Toxoplasma gondii. Cell Mol Life Sci 75:2355-2373.

494   5.   Hill D, Dubey JP. 2002. Toxoplasma gondii: transmission, diagnosis and

495   prevention. Clin Microbiol Infect 8:634-40.

496   6.   White MW, Radke JR, Radke JB. 2014. Toxoplasma development - turn the

497   switch on or off? Cell Microbiol 16:466-72.

498   7.   Waldman BS, Schwarz D, Wadsworth MH, 2nd, Saeij JP, Shalek AK, Lourido S.

499   2020. Identification of a Master Regulator of Differentiation in Toxoplasma. Cell

500   180:359-372 e16.

501   8.   Lucky AB, Wang C, Shakri AR, Kalamuddin M, Chim-Ong A, Li X, Miao J. 2023.

502   Plasmodium falciparum GCN5 plays a key role in regulating artemisinin

503   resistance-related stress responses. Antimicrob Agents Chemother 67:e0057723.

504   9.   Johnsson A, Xue-Franzen Y, Lundin M, Wright AP. 2006. Stress-specific role of

505   fission yeast Gcn5 histone acetyltransferase in programming a subset of stress

506   response genes. Eukaryot Cell 5:1337-46.

507   10.   Bhatti MM, Livingston M, Mullapudi N, Sullivan WJ, Jr. 2006. Pair of unusual

508   GCN5 histone acetyltransferases and ADA2 homologues in the protozoan

509   parasite Toxoplasma gondii. Eukaryot Cell 5:62-76.

510   11.   Wang J, Dixon SE, Ting LM, Liu TK, Jeffers V, Croken MM, Calloway M, Cannella

511   D, Hakimi MA, Kim K, Sullivan WJ, Jr. 2014. Lysine acetyltransferase GCN5b

512  interacts with AP2 factors and is required for Toxoplasma gondii proliferation.

513  PLoS Pathog 10:e1003830.

514 12. Naguleswaran A, Elias EV, McClintick J, Edenberg HJ, Sullivan WJ, Jr. 2010.

515  Toxoplasma gondii lysine acetyltransferase GCN5-A functions in the cellular

516  response to alkaline stress and expression of cyst genes. PLoS Pathog

517  6:e1001232.

518 13. Sullivan WJ, Jr., Jeffers V. 2012. Mechanisms of Toxoplasma gondii persistence

519  and latency. FEMS Microbiol Rev 36:717-33.

520 14. Licon MH, Giuliano CJ, Chan AW, Chakladar S, Eberhard JN, Shallberg LA,

521  Chandrasekaran S, Waldman BS, Koshy AA, Hunter CA, Lourido S. 2023. A

522  positive feedback loop controls Toxoplasma chronic differentiation. Nat Microbiol

523  8:889-904.

524 15. Dey V, Holmes MJ, Bastos MS, Wek RC, Sullivan WJ, Jr. 2024. Cap-independent

525  translation directs stress-induced differentiation of the protozoan parasite

526  Toxoplasma gondii. J Biol Chem 300:107979.

527 16. Sokol-Borrelli SL, Reilly SM, Holmes MJ, Orchanian SB, Massmann MD, Sharp

528  KG, Cabo LF, Alrubaye HS, Martorelli Di Genova B, Lodoen MB, Sullivan WJ, Jr.,

529  Boyle JP. 2023. A transcriptional network required for bradyzoite development in

530  Toxoplasma gondii is dispensable for recrudescent disease. Nat Commun

531  14:6078.

532 17. Dever TE, Ivanov IP, Hinnebusch AG. 2023. Translational regulation by uORFs

533  and start codon selection stringency. Genes Dev 37:474-489.

534   18.   Cleary MD, Singh U, Blader IJ, Brewer JL, Boothroyd JC. 2002. Toxoplasma

535   gondii asexual development: identification of developmentally regulated genes

536   and distinct patterns of gene expression. Eukaryot Cell 1:329-40.

537   19.   Radke JR, Guerini MN, Jerome M, White MW. 2003. A change in the premitotic

538   period of the cell cycle is associated with bradyzoite differentiation in Toxoplasma

539   gondii. Mol Biochem Parasitol 131:119-27.

540   20.   Harris MT, Jeffers V, Martynowicz J, True JD, Mosley AL, Sullivan WJ, Jr. 2019. A

541   novel GCN5b lysine acetyltransferase complex associates with distinct

542   transcription factors in the protozoan parasite Toxoplasma gondii. Mol Biochem

543   Parasitol 232:111203.

544   21.   Harb OS, Roos DS. 2020. ToxoDB: Functional Genomics Resource for

545   Toxoplasma and Related Organisms. Methods Mol Biol 2071:27-47.

546   22.   Lee VV, Judd LM, Jex AR, Holt KE, Tonkin CJ, Ralph SA. 2021. Direct Nanopore

547   Sequencing of mRNA Reveals Landscape of Transcript Isoforms in

548   Apicomplexan Parasites. mSystems 6.

549   23.   Krueger A, Horjales S, Yang C, Blakely WJ, Francia ME, Arrizabalaga G. 2024.

550   The essential kinase TgGSK regulates centrosome division and endodyogeny in

551   Toxoplasma gondii. bioRxiv doi:10.1101/2024.09.27.615374.

552   24.   Wang C, Kassem S, Rocha REO, Sun P, Nguyen TT, Kloehn J, Liu X, Brusini L,

553   Bonavoglia A, Barua S, Boissier F, Lucia Del Cistia M, Peng H, Tang X, Xie F,

554   Wang Z, Vadas O, Suo X, Hashem Y, Soldati-Favre D, Jia Y. 2024. Apicomplexan

555   mitoribosome from highly fragmented rRNAs to a functional machine. Nat

556   Commun 15:10689.

25. Sun J, Paduch M, Kim SA, Kramer RM, Barrios AF, Lu V, Luke J, Usatyuk S,

Kossiakoff AA, Tan S. 2018. Structural basis for activation of SAGA histone

acetyltransferase Gcn5 by partner subunit Ada2. Proc Natl Acad Sci U S A

115:10010-10015.

26. Aravind L, Iyer LM. 2012. The HARE-HTH and associated domains: novel

modules in the coordination of epigenetic DNA and protein modifications. Cell

Cycle 11:119-31.

27. Atanassov BS, Evrard YA, Multani AS, Zhang Z, Tora L, Devys D, Chang S, Dent

SY. 2009. Gcn5 and SAGA regulate shelterin protein turnover and telomere

maintenance. Mol Cell 35:352-64.

28. Jeitany M, Bakhos-Douaihy D, Silvestre DC, Pineda JR, Ugolin N, Moussa A,

Gauthier LR, Busso D, Junier MP, Chneiweiss H, Chevillard S, Desmaze C,

Boussin FD. 2017. Opposite effects of GCN5 and PCAF knockdowns on the

alternative mechanism of telomere maintenance. Oncotarget 8:26269-26280.

29. Her YR, Chung IK. 2013. p300-mediated acetylation of TRF2 is required for

maintaining functional telomeres. Nucleic Acids Res 41:2267-83.

30. Lee SY, Lee SH, Choi NH, Kim JY, Kweon JH, Miller KM, Kim JJ. 2024. PCAF

promotes R-loop resolution via histone acetylation. Nucleic Acids Res 52:8643-

8660.

31. Fox BA, Falla A, Rommereim LM, Tomita T, Gigley JP, Mercier C, Cesbron-

Delauw MF, Weiss LM, Bzik DJ. 2011. Type II Toxoplasma gondii KU80 knockout

strains enable functional analysis of genes required for cyst development and

latent infection. Eukaryot Cell 10:1193-206.

32.   Huynh MH, Carruthers VB. 2009. Tagging of endogenous genes in a Toxoplasma

gondii strain lacking Ku80. Eukaryot Cell 8:530-9.

33.   Peng D, Tarleton R. 2015. EuPaGDT: a web tool tailored to design CRISPR

guide RNAs for eukaryotic pathogens. Microb Genom 1:e000033.

34.   Shen B, Brown KM, Lee TD, Sibley LD. 2014. Efficient gene disruption in diverse

strains of Toxoplasma gondii using CRISPR/CAS9. mBio 5:e01114-14.

35.   Donald RG, Roos DS. 1993. Stable molecular transformation of Toxoplasma

gondii: a selectable dihydrofolate reductase-thymidylate synthase marker based

on drug-resistance mutations in malaria. Proc Natl Acad Sci U S A 90:11703-7.

36.   Shen B, Brown K, Long S, Sibley LD. 2017. Development of CRISPR/Cas9 for

Efficient Genome Editing in Toxoplasma gondii. Methods Mol Biol 1498:79-103.

37.   Donald RG, Carter D, Ullman B, Roos DS. 1996. Insertional tagging, cloning, and

expression of the Toxoplasma gondii hypoxanthine-xanthine-guanine

phosphoribosyltransferase gene. Use as a selectable marker for stable

transformation. J Biol Chem 271:14010-9.

38.   Roos DS, Crawford MJ, Donald RG, Fohl LM, Hager KM, Kissinger JC, Reynolds

MG, Striepen B, Sullivan WJ, Jr. 1999. Transport and trafficking: Toxoplasma as

a model for Plasmodium. Novartis Found Symp 226:176-95; discussion 195-8.

39.   Smith D, Kannan G, Coppens I, Wang F, Nguyen HM, Cerutti A, Olafsson EB,

Rimple PA, Schultz TL, Mercado Soto NM, Di Cristina M, Besteiro S, Carruthers

VB. 2021. Toxoplasma TgATG9 is critical for autophagy and long-term

persistence in tissue cysts. Elife 10.

40.   Kechin A, Boyarskikh U, Kel A, Filipenko M. 2017. cutPrimers: A New Tool for

Accurate Cutting of Primers from Reads of Targeted Next Generation

Sequencing. J Comput Biol 24:1138-1143.

41.   Kim D, Langmead B, Salzberg SL. 2015. HISAT: a fast spliced aligner with low

memory requirements. Nat Methods 12:357-60.

42.   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol 15:550.

43.   Anders S, Pyl PT, Huber W. 2015. HTSeq--a Python framework to work with

high-throughput sequencing data. Bioinformatics 31:166-9.

44.   Holmes MJ, Bastos MS, Dey V, Severo V, Wek RC, Sullivan WJ, Jr. 2024. mRNA

cap-binding protein eIF4E1 is a novel regulator of Toxoplasma gondii latency.

mBio 15:e0295423.

45.   Srivastava S, Holmes MJ, White MW, Sullivan WJ, Jr. 2023. Toxoplasma gondii

AP2XII-2 Contributes to Transcriptional Repression for Sexual Commitment.

mSphere 8:e0060622.

46.   Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat

Methods 9:357-9.

47.   Thorvaldsdottir H, Robinson JT, Mesirov JP. 2013. Integrative Genomics Viewer

(IGV): high-performance genomics data visualization and exploration. Brief

Bioinform 14:178-92.

48.   Finn EH, Misteli T. 2021. A high-throughput DNA FISH protocol to visualize

genome regions in human cells. STAR Protoc 2:100741.

Table 1. GCN5a interactome analysis. The number of peptides from three

independent pull-downs is shown, generated from RH GCN5a[HA] tachyzoites subjected

to 48 h alkaline stress or not. Results are filtered for proteins represented by at least 5

peptides in two of the three replicates per condition. Proteins localized to organelles

outside the nucleus were excluded. Grey highlight indicates an associating protein

shared with the GCN5b complex.

GeneID	Description (CRISPR fitness score)	No Stress	48 h Stress
TGGT1_254555	**GCN5a (-1.0)**	**51|26|48**	**69|50|57**
TGGT1_234900	PHD-finger domain-containing protein (-4.4)	26|15|27	71|65|63
TGGT1_262420	AP2VIIb-1/ADA2-B (0.0)	17|14|16	45|56|44
TGGT1_247700	AP2XII-4 (-5.0)	7|5|11	41|44|36
TGGT1_220520 TGGT1_220530	AP2V-1 (0.8)	1|0|5	21|22|15
TGGT1_237090	AP2X-5 (-3.5)	0|4|0	14|11|9
TGGT1_253440	Cell cycle-associated protein kinase SRPK, putative (-3.0)	5|0|3	9|0|11

FIGURE LEGENDS

Figure 1. GCN5a^{HA} protein levels increase during alkaline stress. A. GCN5a^{HA}

parasites were cultured in tachyzoite (No Stress) or alkaline stress conditions for 48 h

(Stress). The level of *GCN5a* mRNA expression was monitored by RT-qPCR. The data

are represented as the average of 3 biological replicates ± standard deviation.

Statistical significance was measured by unpaired Student's t test; ns = not significant;

**p ≤ 0.01. B. Western blot of GCN5a^{HA} lysates from parasites cultured under

designated conditions with stage-specific controls (BAG1 and SAG1) and a

constitutively expressed control (TUB).

Figure 2. Phenotypic analysis of ∆GCN5a parasites. A. Plaque assay of GCN5a^{HA}

and ∆GCN5a parasites grown under tachyzoite culture conditions for 14 days. B. The

area of 50 randomly selected plaques was measured from 3 biological replicates.

Statistical significance was measured by unpaired Student's t test; **** represents p

<0.0001. C. Number of plaques formed in HFF monolayers by GCN5a^{HA} and ∆GCN5a

parasites. Each data point represents an average of 3 biological replicates ± standard

deviation. Statistical significance was measured by unpaired Student's t test; ns = not

significant. D. Replication assay for GCN5a^{HA} and ∆GCN5a parasites. Number of

parasites per vacuole was counted from 100 randomly selected vacuoles. Statistical

significance was measured by one-way ANOVA with Tukey's method for multiple

comparisons; **** represents p <0.0001.

Figure 3. ΔGCN5a tachyzoites show increased expression of bradyzoite-

associated genes. A. Venn diagram showing a high degree of gene co-regulation for

GCN5a[HA] and ΔGCN5a lines when they are subjected to bradyzoite-inducing culture

conditions (alkaline media for 48 h). B. Comparison of differential gene expression

between GCN5a[HA] and ΔGCN5a parasites during stress-induced bradyzoite formation.

Canonical stage-specific genes *BAG1*, *ENO1*, *LDH2*, and *SAG1* genes are highlighted

in orange. C) MA plot depicting differential gene expression between GCN5a[HA] and

ΔGCN5a tachyzoites. Up- and downregulated genes are colored in red and blue,

respectively. Canonical stage-specific genes *BAG1*, *ENO1*, *LDH2*, and *SAG1* genes are

highlighted in orange.

Figure 4. Cyst formation in ΔGCN5a parasites. A. (Left) Quantification of the number

of DBA positive vacuoles following 24, 48, 72, and 96 h in alkaline stress. Each data

point represents an average of 3 biological replicates ± standard deviation. Statistical

significance was measured by one-way ANOVA with Tukey's multiple comparison test;

ns = p>0.05, * = p ≤0.05. (Right) The area of at least 100 randomly selected cysts was

measured from 3 biological replicates. Statistical significance was measured by one-

way ANOVA with Tukey's multiple comparison test; ns = p>0.05, **** = p <0.0001. B.

IFA of GCN5a[HA] and ΔGCN5a parasite vacuoles after 72 h of alkaline stress. FITC-DBL

was used to demarcate bradyzoite cyst wall (yellow). Anti-HA antibody was used to

detect GCN5a[HA] (magenta) and DAPI was used to stain nuclei (blue). Scale bar = 10

μm.

Figure 5. Chromosomal occupancy of GCN5a versus GCN5b. A. Meta-gene plot

demonstrating enrichment of GCN5b, but not GCN5a, near the transcriptional start sites

(TSS) of protein coding genes as determined by CUT&Tag. Legend depicting sample

identification is shown in panel B. B. Meta-chromosomal plot demonstrating enrichment

of GCN5a, but not GCN5b, at telomeres.

Figure 6. Independent validation of GCN5a localization to telomeres. A. GCN5a^{HA}

parasites were cultured under alkaline stress conditions for 72 hours before IFA

analyses. A representative vacuole of telomere FISH is shown. DNA is stained with

DAPI (blue), GCN5a^{HA} is stained with anti-HA (yellow), telomere is stained with Cy5-

telomere probe (magenta). Scale bar = 10 μm. The lower right panel indicates the

profile regions that were assessed for signal intensity. B. Densitometry profile indicated

in lower right panel of A, showing overlap between GCN5a and telomeric fluorescent

signals. Profiles 2-5 are in Supplemental Figure S6. C. GCN5a^{HA} and PruΔKu80ΔHX

(Pru) parasites were cultured alkaline stress conditions for 48 h prior to harvesting for

ChIP-qPCR using anti-HA antibody. Immunoprecipitated DNA was examined by PCR

using primers spanning the telomeric region of ChrIb (ChrIb-tel) or designated controls.

Results are represented as a ratio of ChIP/Input DNA. Each data point represents an

average of 3 biological replicates ± standard deviation. Statistical significance was

measured by one-way ANOVA with Tukey's multiple comparison test; ns = $p > 0.05$, ****

= $p < 0.0001$.