Interactions between non-prion and prion domains of Rnq1

direct formation of amyloid vs liquid-like aggregates and

create transmission barriers

Short title:

Interactions between prion and non-prion domains of Rnq1

Sangeun Park[1], Dania M. Maldonado[1], Michele L. Kadnar[2], Monica Andrade[2, #a],

Anna P. Fomitchova[2, #b], Susan W. Liebman[1], Irina L. Derkatch[1,2,*]

[1] Department of Pharmacology, University of Nevada Reno, Reno, Nevada, United States of

America

[2] Department of Microbiology, New York University School of Medicine, New York, New York,

United States of America

[#a] Current address: Department of Biological and Vision Sciences, College of Optometry, State

University of New York, New York, New York, United States of America

[#b] Current address: New York Medical College / New York Metropolitan Hospital, New York, New

York, United States of America

* Corresponding Author:

E-mail: iderkatch@unr.edu

Abstract

Prions are self-propagating protein conformations usually existing as amyloid aggregates. [*PIN⁺*],

a prion form of the Rnq1 protein occasionally found in wild and laboratory yeast strains,

facilitates both the *de novo* formation and destabilization of other yeast prions, and affects

aggregation and toxicity of human misfolding disease proteins expressed in yeast. Rnq1 contains

a short N-terminus with no confirmed function (the non-prion domain, NPD) and a C-terminus

that carries four QN-rich regions and is sufficient for [*PIN⁺*] formation and maintenance (prion

domain, PD). In the current study, a genetic screen identified the NPD T27P mutation that blocks

transmission of the [*PIN⁺*] prion state from wild type Rnq1 ($Rnq1_{WT}$) to mutant $Rnq1_{T27P}$. The

mutation doesn't prevent $Rnq1_{T27P}$ from switching to a prion state when overexpressed *in vivo*, or

from forming amyloid fibers *in vitro*. Furthermore, like [$PIN⁺_{WT}$], the newly formed [$PIN⁺_{T27P}$]s

promote the *de novo* appearance of the Sup35-based prion [*PSI⁺*]. We conclude that the NPD

mutation creates a barrier for prion transmission from [$PIN⁺_{WT}$] to $Rnq1_{T27P}$. Because

fluorescence microscopy shows that $Rnq1_{T27P}$ efficiently joins [$PIN⁺_{WT}$] aggregates, the barrier is

likely due to the inability of $Rnq1_{T27P}$ to propagate the specific [$PIN⁺_{WT}$] conformational variant.

Indeed, the analysis of [$PIN⁺_{T27P}$]s resulting from rare transmission events from [$PIN⁺_{WT}$] indicates

that these [$PIN⁺_{T27P}$]s must undergo conformational adaptation to yield more stable prion

variants. Deletion analysis revealed that T27P constrains prion conformations through the first

two QN-rich regions within the PD. The finding that $Rnq1_{T27P}$-YFP readily forms non-amyloid

liquid-like droplets, which $Rnq1_{WT}$-YFP does not form, supports the idea that the NPD affects

aggregation properties of the PD. We propose that these aggregation properties are essential for

Rnq1's functions, such as controlling aggregation of other proteins. This provides new insight

into the role of heterologous proteins and transmission barriers in the origins of protein misfolding

diseases.

Keywords: yeast, prion, prion variant, prion strain, amyloid, liquid-like droplets, transmission

barrier, Rnq1, [*PIN⁺*], [*RNQ⁺*], Sup35, [*PSI⁺*].

Abbreviations: NPD – non-prion domain, PD – prion domain, WT – wild type, GuHCl- guanidine

hydrochloride, 5-FOA - 5-fluoroorotic acid, aa – amino acid, YFP – Yellow Fluorescent Protein,

CFP – Cyan Fluorescent Protein, EV – empty vector, TEM – transmission electron microscopy,

ThT - thioflavin T, Hex - 1,6 hexanediol –, RT - room temperature, HFIP - 1,1,1,3,3,3

hexafluoroisopropanol.

Author Summary

Proteins must fold into the right shapes to work properly. Sometimes they fold incorrectly and stick together, forming long fiber aggregates that damage cells. This kind of "protein misfolding" causes human diseases such as Alzheimer's. Certain yeast proteins behave similarly, making them useful to study this process. We investigate a yeast protein called Rnq1, which has a region that helps it misfold into fibers. These fibers can also cause other, unrelated proteins to misfold. We found that a mutation in a different part of Rnq1— outside the aggregation region — reduces the ability of non-mutant Rnq1 fibers to convert mutant Rnq1 into growing fiber aggregates. We also identified which section of the aggregation region is affected by this mutation. Interestingly, although the rarely converted mutant aggregates grow poorly at first, they can eventually "adapt" into a shape that grows better. The same mutation also pushes Rnq1 to form liquid-like droplets instead of fibers. Our findings show that the non-aggregating part of Rnq1 controls how Rnq1 aggregates, and, consequently, the appearance and elimination of aggregates formed by other proteins. Our work also helps explain how barriers to misfolded protein growth can be overcome, which is relevant to understanding human protein misfolding diseases.

Introduction

The prion hypothesis postulates that some proteins can acquire stable unconventional

conformations and pass these conformations to newly synthesized molecules with the same

primary sequence. Amyloid, the β-sheet-rich fibers with protein molecules stacked on top of one

another, is the structural basis for most prions and prion-like self-propagating protein

conformations (Chuang et al., 2018; Dobson, 2017; Ke et al., 2020). Prions and amyloids

attracted attention due to their connection to the so-called protein misfolding diseases that are

associated with the formation of amyloid aggregates. The list of such diseases is expanding

quickly from *bona fide* prion diseases, such as scrapie in sheep, mad cow disease in cattle, and

Creutzfeldt-Jakob disease (CJD) in humans, which prompted the development of the paradigm

of protein-based inheritance, to much more common neurodegenerative diseases, such as

Alzheimer's and Parkinson's, as well as type 2 diabetes and, possibly, cancer (Ayers et al., 2020;

Joshi and Ahuja, 2023).

Many protein misfolding diseases can be both sporadic, usually with a later age of onset,

and hereditary, e.g. due to mutations in amyloid-forming proteins. This underscores the

importance of uncovering the cellular mechanisms that drive and inhibit protein misfolding, as

well as the control of conformational changes within specific aggregation-prone proteins that lead

to disease-related aggregation. Prion diseases can also be infectious. Luckily, protective

measures were stepped up in recent decades to prevent iatrogenic human to human

transmission. Animal to human transmission is usually prevented by transmission barriers when

differences in primary sequences of protein homologs from different species make interspecies

transmission of the prion state inefficient (Beringue et al., 2008). However, as we learned from

the appearance of a novel human disease resulting from a cross of the transmission barrier

between cattle and humans, variant Creutzfeldt-Jakob disease (vCJD), such barriers may not be

strong enough (Houston and Andreoletti, 2019). An important aspect of the problem is possible

rapid adaptation of prion conformations in a new host after rare across-the-barrier transmission

events (Weissman, 2012). Also, across-the-barrier transmission leads to forms of disease with

distinct clinical manifestations compared to sporadic and hereditary diseases, like in the case of

sporadic CJD and vCJD transmitted from cattle (Igel-Egalon et al., 2018).

Prion research has been greatly facilitated by the discovery of naturally occurring prions

in the yeast *Saccharomyces cerevisiae* (Wickner, 1994; Liebman and Chernoff, 2012;

Zhouravleva et al., 2023). The same underlying amyloid structure of yeast prions and human

disease-related aggregation-prone proteins, as well as evolutionary conservation of genes

makes yeast an excellent model for studying protein aggregation and mechanisms that control it.

Many yeast prions carry long terminally located QN-rich domains that are unfolded in soluble

proteins but are required and sufficient for taking on prion conformations and maintaining them.

Hence their name – prion domains (PDs). Similar QN-rich or expanded polyQ domains also drive

aggregation of some human disease proteins (Fuenteabla et al., 2010; Lieberman et al., 2019).

[*PSI*+], the prion form of a translation termination factor Sup35 (eRF3), is the most extensively

studied yeast prion, and many important findings come from the [*PSI*+] prion research (Liebman

and Chernoff, 2012). One such finding is prion-prion interactions (Derkatch and Liebman, 2007).

Initially, it was postulated that overexpression of a prion-forming protein should induce the *de*

novo appearance of its respective prion conformation because increased protein concentration

makes switching to a prion state more likely (Wickner, 1994). It was further postulated that if a

prion is lost or cured from the cell, it should re-appear in the cell's progeny either rarely

spontaneously or frequently upon overexpression of the prion-forming protein (*ibid.*). Initial data

for [*PSI*+] appeared to support these postulates (Derkatch et al., 1996). However, once [*PSI*+]

was eliminated by the addition to yeast culture media of low concentrations of guanidine

hydrochloride (GuHCl), a generic yeast prion curing agent, efficient induction of [*PSI*+] by Sup35

overexpression was no longer possible. This was because the treatment also cured the

previously unknown [*PIN⁺*] prion required for induction of [*PSI⁺*] (Derkatch et al., 1997).

[*PIN⁺*] is formed by the Rnq1 protein (Sondheimer and Lindquist, 2000; Derkatch et al.,

2001). Rnq1 consists of a short N-terminal domain with no confirmed function (the non-prion

domain, NPD, aa 1-132; Fig 1A) and a C-terminal QN-rich domain sufficient for [*PIN⁺*] formation

and maintenance (prion domain, PD; aa 132-405). The prion domain of Rnq1 is complex. It

carries four discrete QN-rich regions. None of these alone is essential for [*PIN⁺*] prion

maintenance, and any two are sufficient (Kadnar et al., 2010). Also, bacterially expressed

purified Rnq1 fragments carrying NPD with only QN2, QN3 or QN4 form amyloid fibers *in vitro*

(*ibid.*). The QN1 region is very short and does not form amyloid when attached to the Rnq1 NPD,

but synthetic QN1 peptide does form amyloid fibers (*ibid.*). The [*mini-PIN⁺*]s formed by Rnq1

fragments with different combinations of QN-rich domains are all able to promote the *de novo*

formation of [*PSI⁺*] upon Sup35 overproduction, *i.e.* no specific QN-rich region of Rnq1's PD is

essential for this prion-prion interaction (*ibid.*).

Beyond promoting the *de novo* formation of [*PSI⁺*], [*PIN⁺*] is involved in other prion-prion

interactions, both positive and negative. For example, [*PIN⁺*] also facilitates the *de novo*

appearance of the [*URE3*] and [*SWI⁺*] prions with QN-rich PDs, even though the presence of

[*PIN⁺*] is not a requirement (Bradley et al., 2002; Du et al., 2017), as well as some artificial prions

(Borkhsenius et al., 2002) and the [HET-s]$_Y$ prion formed in yeast by the *Podospora anserine*

prion-forming protein Het-s with a non-QN-rich PD (Taneija et al., 2007). [*PIN⁺*] also interacts

with non-prion aggregation-prone proteins, such as human Huntington's and Machado-Joseph

disease-associated proteins with expanded polyQ stretches (Osherovich and Weissman, 2001;

Meriin et al., 2002) and TDP-43 associated with amyotrophic lateral sclerosis, frontotemporal

dementia and limbic-predominant age-related encephalopathy (Park et al., 2017). An example of

negative interactions is the mutual inhibition of stable inheritance by certain [*PSI⁺*] and [*PIN⁺*]

prion variants: some weak [*PSI+*] variants are very unstable in the presence of some "single dot"

[*PIN+*] variants, while some strong [*PSI+*] variants destabilize some "single dot" [*PIN+*]s (Bradley

and Liebman, 2003; Mathur et al., 2009; Villali et al., 2020). Yet another type of interaction is a

dramatic enhancement by [*PIN+*] of the [*SWI+*]-associated nonsense suppression in the strains

lacking Sup35 PD (aka [*NSI+*], Nizhnikov et al., 2016).

Like other prions, [*PIN+*] depends on chaperone proteins for its maintenance. Some of

these interactions are common with most other prions. For example, lack of the Hsp104

disaggregase cures [*PIN+*], as well as [*URE3*], [*PSI+*], and [*SWI+*] (Chernoff et al., 1995; Derkatch

et al., 1997; Moriama et al., 2000; Du et al., 2008), likely because Hsp104 promotes

conformational changes in prion aggregates and is needed to break growing prion aggregates to

create smaller seeds that can be transmitted to daughter cells (Chernoff et al., 1995; Wegrzyn et

al., 2001; Sapute-Krishnan et al., 2007). However, some prion-chaperone interactions are

specific for a particular prion: the same chaperone can affect the formation of one prion but not

others, and some chaperones have opposite effects on different prions (Barbitoff et al., 2022).

[*PIN+*] has a highly specific interaction with Sis1, an Hsp40 co-chaperone of Hsp70 chaperones:

not only does [*PIN+*] critically depend on Sis1 for maintenance (Sondheimer et al., 2001), but

Sis1 is present in [*PIN+*] aggregates in equimolar amounts with Rnq1 (Lopez et al., 2003). In

Sis1, the region critical for [*PIN+*] maintenance is within the G/F region that is not needed for the

maintenance of other prions (Lopez et al., 2003). In Rnq1, the major site for interaction with Sis1

is located not in the PD, but in the NPD, in the LGKLALl sequence (Douglas et al., 2008).

Strikingly, most prion-forming proteins, including PrP in mammals and Rnq1, Sup35, and

Ure2 in yeast, can form not just one but multiple prion variants (or prion strains) that differ in

numerous prion-associated phenotypes (Bessen and Marsh, 1992; Bessen et al., 1995; Derkatch

et al., 1996; Bradley et al., 2002; Schlumpberger et al., 2001). These prion variants manifest

different conformations of prion-forming proteins in prion aggregates thus leading to differences

in aggregate size, sensitivity to cellular chaperones and interactions with other prions *in vivo*, as

well as in stability of corresponding amyloids *in vitro* (see above). Extensive studies of mutations

within the Sup35 PD demonstrate that its primary sequence affects its conformational flexibility in

terms of what [*PSI⁺*] strains the mutant protein can maintain (DePace, 1998; Chang et al., 2008;

Bateman and Wickner, 2013).

Here we show that in Rnq1 the NPD is involved in determining the conformational

flexibility of its PD and its ability to form particular amyloid variants and non-amyloid aggregates.

Our findings are consistent with the hypothesis that the cellular function of the Rnq1 NPD is to

regulate the aggregation of its PD and, through this aggregation, to control aggregation and

disaggregation of other prions and prion-like aggregates.

Results

Genetic screen for *RNQ1* mutations leading to a Pin⁻ phenotype

Our goal was to investigate the structure-function organization of the Rnq1 protein in terms of its

ability to maintain the [*PIN⁺*] prion and to facilitate the induction of the heterologous [*PSI⁺*] prion.

Our approach was to screen for *RNQ1* mutant alleles (*RNQ1_{MUTANT}*) that, when substituted for

wild type *RNQ1 (RNQ1_{WT})* in a [*PIN⁺*] strain (Pin⁺ phenotype), lead to a Pin⁻ phenotype. The Pin

phenotype was scored by examining the *de novo* appearance of the [*PSI⁺*] prion upon transient

high-level expression of the PD of the [*PSI⁺*] prion forming protein Sup35 fused to YFP,

Sup35NM-YFP. The Pin⁺ *vs* Pin⁻ phenotype was assigned, respectively, when [*PSI⁺*] appearance

was detected *vs* not detected.

The screen is schematically shown in Fig 1B. **First**, *CEN LEU2* plasmids carrying PCR-

mutagenized *RNQ1* driven by its native non-mutagenized promoter and followed by its native

non-mutagenized terminator sequence were transformed into a [*PIN⁺*][*psi⁻*] strain where the

chromosomal copy of *RNQ1* was deleted and [*PIN⁺*] was maintained by a similar *RNQ1*$_{WT}$

plasmid (*CEN URA3*). The strain also carried the *CEN HIS3* plasmid with the *SUP35NM-YFP*

[*PSI⁺*] inducing construct controlled by the *GAL* promoter, and the *ade1-14* mutation allowing for

[*PSI⁺*] detection. Transformants were selected on and then patched onto synthetic media

selective for all plasmids, thus allowing for ~20 generations of co-expression of Rnq1$_{WT}$ and

Rnq1$_{MUTANT}$. **Second**, after two passages on media selective for only the mutagenized *RNQ1*

and [*PSI⁺*] inducing plasmids, the *RNQ1*$_{WT}$ maintainer plasmid was shuffled out on media

supplemented with 5-FOA and selective for the other two plasmids. **Third**, [*PSI⁺*] was induced by

high-level expression of Sup35NM-YFP on SGal-LeuHis media. **Fourth**, cells carrying [*PSI⁺*]

were selected on non-inducing media lacking adenine and not selective for plasmids. Selection

of [*PSI⁺*] cells on adenineless media is based on nonsense suppression of the *ade1*-14 (UGA)

mutation due to the recruitment of the Sup35 protein, a translation termination factor, into the

[*PSI⁺*] prion aggregates. For Pin⁺ cultures, thick, even if heterogeneous, growth was observed

over the entire patch. For the presumptive Pin⁻ cultures, either no growth or rare Ade⁺ colonies

were observed on media lacking adenine. In such cultures, the Pin⁻ phenotype was confirmed by

colony purifying yeast from the inducing SGal-LeuHis media onto YPD, where [*PSI⁺*] Ade⁺ cells

form pink or white colonies that are clearly distinct from red Ade⁻ colonies formed by [*psi⁻*] cells.

Finally, the Pin⁻ phenotype was confirmed by fluorescence microscopy of cells grown on the

[*PSI⁺*] inducing galactose media using the YFP-tagged Sup35NM as a reporter for the lack of

Sup35 aggregation. In Pin⁺ cultures the appearance of [*PSI⁺*] is visualized as ring- or dot-like foci,

while in [*psi⁻*] cells Sup35NM fluorescence is evenly distributed (Patino and Lindquist, 1996;

Zhou et al., 2001). To confirm that the Pin⁻ phenotype was due to a mutation in the shuffled-in

RNQ1 gene, *RNQ1*$_{MUTANT}$ plasmids were isolated from each candidate mutant, individually re-

transformed back into the yeast strain used for the original screen, and tested according to the

same scheme (see Materials and Methods for additional approaches used to improve the

candidate pool).

Approximately 12,000 yeast transformants were screened and 54 were either firmly Pin[-],

or their Pin phenotype was dramatically reduced. 53 of these carried plasmids with either

nonsense or frameshift *RNQ1* mutations. All nonsense mutations and most frameshifts were

located in the N-terminal part of *RNQ1*, either within its NPD, or before the end of the second

QN-rich domain within the PD (QN2, see Fig 1A). All five unique frameshift mutations that were

located downstream of QN2 led to long C-terminal extensions due to out-of-frame translation.

The only missense mutant that passed the screen carried the T27P (ACA ⇨ CCA) mutation in

the beginning of the Rnq1 NPD (Fig 1A; also see Fig 2B).

Identification in the screen of constructs synthesizing only the NPD or part of it was

expected because of the requirement of the PD of Rnq1 for the maintenance of the [*PIN+*] prion

(Vitrenko et al., 2007a; Kadnar et al., 2010). The fact that no nonsense mutations were detected

beyond the second QN-rich region can be explained based on our earlier deletion analysis of

Rnq1 (Kadnar et al., 2010): the QN-rich regions within Rnq1 PD are redundant in their ability to

acquire the [*PIN+*] prion state from Rnq1$_{WT}$ and maintain the Pin+ phenotype (i.e. allow for [*PSI+*]

induction). Specifically, the N-terminal Rnq1 fragments including at least QN1 and entire QN2

were sufficient to maintain Pin+, so mutants retaining these regions were not expected to pass

the screen. However, deletion analysis also indicated that the C-terminal location of the QN2

region in the truncated construct carrying only QN1 and QN2 was important for the transfer of

[*PIN+*] from Rnq1$_{WT}$ (*ibid*), so the identification of frameshifts leading to long extensions

downstream of QN2 is also not surprising. Finally, the fact that we have not identified PD-located

missense mutations can also be explained by the structural redundancy of the Rnq1 PD: a single

mutation that would just inactivate one of the four QN-rich regions is unlikely to block [*PIN+*]

propagation or the ability of the mutant to promote [*PSI+*] induction.

The inability to induce nonsense suppression in cultures where Rnq1$_{T27P}$ was

substituted for Rnq1$_{WT}$ is due to the dramatic reduction in the ability to induce

[*PSI⁺*]

For all experiments described here and below, we used plasmids with the T27P mutation

introduced by site-directed mutagenesis, to exclude the possibility that additional mutations in the

regions we did not sequence outside of the *RNQ1* gene may have contributed to the Pin⁻

phenotype during the screen. We also reasonably excluded the possibility that the absence of

[*PIN⁺$_{T27P}$*] in most post-shuffle Rnq1$_{T27P}$ cells was due to insufficient Rnq1$_{T27P}$. Although there is

an ~2-fold difference in protein abundance between Rnq1$_{T27P}$ and Rnq1$_{WT}$ (Fig 2A) this is unlikely

to be the main reason for the prion loss because (i) [*PIN⁺$_{WT}$*] is 100% stable in diploids

heterozygous for *RNQ1* deletions (Derkatch et al., 2001) and (ii) a similar reduction in

abundance compared to full-length Rnq1$_{WT}$ does not destabilize mini-[*PIN⁺*]s maintained by

Rnq1 fragments (Kadnar et al., 2010).

The goal for further experiments was to understand what leads to the loss of the Pin⁺

phenotype of the *rnq1-Δ* yeast cultures expressing Rnq1$_{T27P}$ after the elimination of Rnq1$_{WT}$. As

can be seen on Fig 2B, the cultures expressing Rnq1$_{T27P}$ were almost Pin⁻ compared to the

positive control expressing Rnq1$_{WT}$. However, compared to the negative control where Rnq1$_{WT}$

was substituted with the empty vector (EV), there were occasional Ade⁺ colonies on adenineless

media on the Rnq1$_{T27P}$ patches. Out of 24 such Ade⁺ microcolonies from six independent

Rnq1$_{T27P}$ transformants, 23 were [*PSI⁺*] according to the GuHCl test. The fluorescence analysis

of Ade⁺ cultures that retained the *GAL-SUP35NM-YFP* plasmid revealed dot-like foci in most

fluorescent cells in all eight cultures examined, and these foci were indistinguishable from [*PSI⁺*]

foci in the control Rnq1$_{WT}$ cultures. Thus, the Ade⁺ colonies in the cells expressing Rnq1$_{T27P}$ after

shuffle were due to [*PSI*+]s, so [*PSI*+] induction was not blocked completely due to the T27P

mutation.

We then asked whether the [*PSI*+] induction was indeed ~50-100-fold less frequent in the

Rnq1$_{T27P}$ *vs* the Rnq1$_{WT}$ cultures, as it appeared, or alternatively it was close to normal levels but

most newly forming [*PSI*+]s were not detected in the assay based on the ability of [*PSI*+] cells to

grow on adenineless media. This could happen for five reasons. (1) Most [*PSI*+] variants forming

in cells carrying [*PIN*+$_{T27P}$] are "very weak" and unable to ensure robust growth on adenineless

media. Indeed, some [*PIN*+] variants promote induction of predominantly weak [*PSI*+]s (Sharma

and Liebman, 2013). (2) Most [*PSI*+]s promoted by [*PIN*+$_{T27P}$] are highly unstable and are lost

during the first few cell divisions. Indeed, some [*PSI*+] variants are permanently unstable

(Sharma and Liebman, 2012), and some [*PIN*+] variants are known to destabilize weak [*PSI*+]s

they induce (Bradley and Liebman, 2003). (3) In [*PIN*+$_{T27P}$] cells, most newly forming Sup35

aggregates do not mature into [*PSI*+] or are lost in the absence of Sup35 overexpression. Such

[*PSI*+] variants have been described by Salnikova et al. (2005). (4) Most [*PSI*+]s in [*PIN*+$_{T27P}$]

cultures are extremely toxic and lead to cell death on the Sup35NM-YFP inducing media, prior to

transfer to adenineless media where Sup35NM-YFP is no longer expressed. Indeed, growth of

[*PSI*+] cells is inhibited by Sup35 overexpression in a [*PSI*+] variant-specific manner (Derkatch et

al., 1996). (5) Most [*PSI*+]s induced in [*PIN*+$_{T27P}$] are toxic even in the absence of Sup35NM-YFP

expression. Such "killer" [*PSI*+]s have also been described (McGlinchey et al., 2011).

Experiments discussed below essentially exclude all these possibilities.

To assess the strength of [*PSI*+] variants induced in cells expressing Rnq1$_{WT}$ and

Rnq1$_{T27P}$, yeast scraped up from patches incubated on SD-Ade media at 20^{0}C for ~25 days after

[*PSI*+] induction were colony purified on YPD, and individual colonies were randomly picked for

further analysis. Incubation at low temperature promotes nonsense suppression in weaker [*PSI*+]

variants (Derkatch et al., 1996) and prolonged incubation, in our experience, allows even the

weakest [*PSI*+]s to form microcolonies. We found that similar arrays of [*PSI*+] variants were

induced in the presence of Rnq1$_{WT}$ and Rnq1$_{T27P}$, with no prevalence for weak [*PSI*+]s in the

Rnq1$_{T27P}$ cultures. Specifically, moderate [*PSI*+]s constituted 85% and 91%, weak [*PSI*+]s – 6%

and 6%, and strong [*PSI*+]s – 10% and 3% for Rnq1$_{WT}$ and Rnq1$_{T27P}$, respectively (Table S1).

The slightly smaller percentage of strong [*PSI*+] variants induced in Rnq1$_{T27P}$ cells was not

statistically significant ($p > 0.1$) and could not explain the large difference in the Pin phenotype.

Furthermore, the dramatically reduced Pin+ phenotype in the Rnq1$_{T27P}$ after-shuffle

cultures was not due to extremely weak (essentially Ade⁻) or highly unstable [*PSI*+] variants. To

test this, yeast were colony purified directly from the [*PSI*+] inducing galactose media onto YPD.

Only ~0.7% of >5,000 colonies originating from 12 independent *RNQ1$_{T27P}$* transformants were

white or pink, and most of these colonies were [*PSI*+] based on the GuHCl test and, also, the

Sup35 aggregation test for those that retained the *SUP35NM-YFP* plasmid. The color of pink

colonies and their ratio to white colonies was similar to that in the positive control cultures with

RNQ1$_{WT}$ shuffled-in where white and pink (and [*PSI*+]) colonies constituted ~20%. In the negative

control cultures with EV shuffled-in, the rare (<0.1%) non-red colonies were usually white and not

[*PSI*+]. This is consistent with results shown in Fig 2B. Importantly, no dark pink colonies

indicative of very weak [*PSI*+] were detected in Rnq1$_{T27P}$ cultures, and analysis of all slightly

lighter shade of red colonies (~30) showed that they were not [*PSI*+]. In addition, fluorescence

microscopy of ~40 typical red colonies retaining *GAL-SUP35NM-YFP* did not reveal any [*PSI*+]s.

Finally, no increase of the proportion of sectored white/red or pink/red [*PSI*+] colonies was

detected for Rnq1$_{T27P}$ compared to Rnq1$_{WT}$. The fact that newly forming [*PSI*+]s are frequently

unstable is well-known, so the presence of some sectored colonies was expected (Sharma and

Liebman, 2012). All potentially sectored colonies in Rnq1$_{T27P}$ cultures yielded stable [*PSI*+]s upon

further analysis. Altogether, these experiments exclude possibilities (1) and (2) from the list

above.

Fluorescence microscopy of aggregate formation upon high expression of Sup35NM-YFP

was used to record the initial steps of [PSI^+] formation. In the after-shuffle Rnq1$_{T27P}$ cultures only

~2.5% cells had aggregates (Table 1; Fig 2C). This is significantly less compared to Rnq1$_{WT}$

(~57%) and significantly more compared to EV (<0.01%). Both ring-, line- and dot-shaped foci

were detected in the $RNQ1_{T27P}$ carrying cells, and these foci were not distinguishable from those

in cells with $RNQ1_{WT}$. The presence of dot-like foci indicative of mature [PSI^+] and occasional

mother-daughter pairs with aggregates in both cells indicated that newly forming Sup35

aggregates were heritable in the Rnq1$_{T27P}$ expressing cells. Also, there was no indication of

increased cell toxicity in cells co-expressing Rnq1$_{T27P}$ and Sup35NM-YFP: no abnormally shaped

elongated cells, no increase in the proportion of non-fluorescent cells, and no growth inhibition

compared to the Rnq1$_{WT}$ control (Fig S1). Finally, in the [PSI^+]s induced in the Rnq1$_{T27P}$

expressing cells, maintenance of the *LEU2 RNQ1$_{T27P}$* and *HIS3 GAL-SUP35NM-YFP* plasmids

was not reduced compared to the maintenance of *LEU2 RNQ1$_{WT}$* and *HIS3 GAL-SUP35NM-YFP*

in the [PSI^+]s induced in the Rnq1$_{WT}$ cells (see description of [PSI^+] isolation above and Table

S1). Together, these experiments exclude possibilities (3), (4), and (5) above.

In summary, the cultures expressing Rnq1$_{T27P}$ have a dramatically reduced but otherwise typical

Pin$^+$ phenotype.

Table 1. Analysis of Sup35NM-YFP aggregation in $RNQ1_{T27P}$ cells after shuffling out the

$RNQ1_{WT}$ plasmid indicates low-level [PSI^+] induction.

Construct after plasmid shuffle	% cells with Sup35NM-YFP aggregates		
	Average	SD	SEM
$RNQ1_{WT}$	56.8	± 14.5	± 3.0
$RNQ1_{T27P}$	2.6	± 3.1	± 0.8
EV	0	NA	NA

Table 1 Legend. Aggregation of Sup35NM-YFP was analyzed in YID146 cells after

substituting the $RNQ1_{T27P}$ *LEU CEN* or indicated control constructs for *URA3 CEN*

$RNQ1_{WT}$ (see Fig 1B and text). To induce the expression of Sup35NM-YFP, yeast

passaged on FOA media to eliminate the *URA3 CEN* $RNQ1_{WT}$ plasmid were replica

plated onto the SD-LeuHis and then SGal-LeuHis media and incubated for ~48 hours

at 30^0C. For $RNQ1_{WT}$ and $RNQ1_{T27P}$, at least 11 vision fields were scored for each

construct, ~500 cells total per construct. For EV, no aggregates were detected in

~1,000 cells scored. Based on T-test, differences are statistically significant between

$Rnq1_{T27P}$ *vs* $Rnq1_{WT}$ ($p<10^{-13}$), $Rnq1_{T27P}$ *vs* EV ($p<0.005$), and $Rnq1_{WT}$ *vs* EV ($p<10^{-14}$).

Qualitative assessment of aggregation in >10,000 cells in the cultures used for

quantification further confirmed the difference between the $RNQ1_{T27P}$ and EV: bright

ring- or dot-aggregates were readily detected in $Rnq1_{T27P}$ expressing cells, but no ring

aggregates were detected in EV cells, where there were only a few dot foci, mostly

atypically large. Also, similar observations were made upon qualitative assessments

after a shorter (overnight) incubation on SGal-LeuHis and after two passes on this

medium for over 4 days.

Overexpression of Rnq1$_{T27P}$ induces a heritable Pin$^+$ phenotype indicative of the

formation of [PIN^+_{T27P}]

In general, the *de novo* appearance of prions can be induced by their transient overexpression,

and the process of prion aggregate formation can be monitored by microscopy by attaching a

fluorescent tag to a prion forming protein (Zhou et al., 2001). However, $Rnq1_{WT}$ does not form *de*

novo [PIN^+] aggregates upon overexpression when attached to a C-terminal GFP family tag

(Derkatch et al., 2001), so a negative result for Rnq1$_{T27P}$ would not be informative. Thus, the *de*

novo induction of [*PIN+$_{T27P}$*] and [*PIN+$_{WT}$*] was attempted by overexpressing, respectively, the

untagged Rnq1$_{T27P}$ and Rnq1$_{WT}$ driven by the inducible *CUP* promoter and scoring for the Pin+

phenotype by [*PSI+*] induction upon high-level expression of Sup35NM-YFP.

For the first experiment (Fig 3A), the *CEN URA3* plasmids carrying either the *CUP-*

RNQ1$_{T27P}$ or *CUP-RNQ1$_{WT}$* inserts, or just the empty *CEN URA3* vector, were introduced into a

[*pin-*][*psi-*] *rnq1-Δ* strain already carrying the *HIS3 GAL-SUP3NM-YFP* [*PSI+*] inducing construct.

Transformants were patched on SD-UraHis media where plasmid-borne constructs were not

expressed. Then yeast cultures were replica plated onto the SGal-UraHis supplemented with

50μM CuSO$_4$ where Rnq1$_{T27P}$ or Rnq1$_{WT}$ were overexpressed together with Sup35NM-YFP, or

on control media where neither or only one of the two constructs was expressed (SD-UraHis,

SD-UraHis+50μM CuSO$_4$, and SGal-UraHis). Induction of [*PSI+*] was scored on adenineless

media to which yeast were replica plated from either SGal-UraHis+50μM CuSO$_4$ or control

media. Indeed, heterogeneous growth on adenineless media indicative of [*PSI+*] induction was

observed following co-overexpression of Sup35NM-YFP with either Rnq1$_{T27P}$ or Rnq1$_{WT}$, with no

obvious effect of the T27P mutation on the efficiency of [*PSI+*] induction (Fig 3A). The *de novo*

formation of [*PSI+*] was also confirmed by fluorescent microscopy: in cells from the SGal-

UraHis+50μM CuSO$_4$ media we observed ring-, line- and dot-shaped Sup35NM-YFP foci

indistinguishable in appearance between the Rnq1$_{T27P}$ and Rnq1$_{WT}$ expressing cultures (data not

shown). The limitation of this experiment was that it did not allow us to test if the Pin+ phenotypes

induced by the overexpression of Rnq1$_{T27P}$ and Rnq1$_{WT}$ were heritable, indicative of the

formation of, respectively, [*PIN$_{T27P}$+*] and [*PIN+$_{WT}$*], or only non-propagating aggregates were

formed by one or both constructs.

For the second experiment (Fig 3B), the [pin^-][psi^-] $rnq1$-Δ strain carrying the *HIS3 CEN*

GAL-SUP3NM-YFP plasmid was co-transformed with *URA3 CEN CUP-RNQ1$_{T27P}$* (to be the

inducer of the [PIN^+_{T27P}] prion state) and the *LEU2 CEN* plasmid carrying *RNQ1$_{T27P}$* under its

original promoter (to be the maintainer of this prion state after induction). Control samples were

co-transformed with corresponding *RNQ1$_{WT}$* plasmids (*URA3 CEN CUP-RNQ1$_{WT}$* and *LEU2*

CEN RNQ1$_{WT}$), or with empty vectors (*URA3 CEN* and *LEU2 CEN*). The Pin phenotype was

scored for by growth on adenineless media after overexpression of Rnq1$_{T27P\ (or\ WT)}$ and *GAL-*

SUP3NM-YFP induction but with 0 (co-overexpression), 1, 2, 3, or 4 passages on the non-

inducing glucose no-copper plasmid-selective media between the Rnq1$_{T27P\ (or\ WT)}$ overexpression

and *GAL-SUP3NM-YFP* induction. The efficiency of [PSI^+] induction was slightly but reproducibly

higher in the Rnq1$_{T27P}$ expressing cells compared to Rnq1$_{WT}$ in the nonsense suppression test for

most of ~50 co-transformants tested for each set of plasmids, but no statistically significant

difference was detected when aggregation was assessed by fluorescent microscopy (not

shown). As can be seen in Fig 3B, the Pin$^+$ phenotype of both Rnq1$_{T27P}$ and Rnq1$_{WT}$ expressing

cultures became slightly weaker upon subsequent passages on media not supplemented with

CuSO$_4$. This is expected considering that newly forming prions are frequently unstable (shown

before for spontaneous [PIN^+_{WT}]s by Derkatch et al. (2000) and newly induced [PSI^+]s by Zhou

et al. (2001)). Important is the retention of the robust Pin$^+$ phenotypes after ~30 generations (4

passages) on the no-copper media where Rnq1$_{T27P}$ and Rnq1$_{WT}$ were no longer overexpressed,

which is indicative of the formation of heritable [PIN_{T27P}^+] and [PIN^+_{WT}] prions (for comparison,

growth for a similar number of generations on 5mM GuHCl media that inhibits prion propagation

is sufficient to completely cure [PIN^+]). Also important is that Rnq1$_{T27P}$ is not deficient in acquiring

such a heritable [PSI^+]-inducing state compared to Rnq1$_{WT}$.

Rnq1$_{T27P}$ forms amyloid fibers *in vitro*

Another way to assess the aggregation properties of Rnq1$_{T27P}$ was to test the ability of the bacterially expressed and purified protein to form amyloid fibers *in vitro*. Previously, we found that the fragments of Rnq1$_{WT}$ encompassing the NPD and one or two QN-rich regions of the PD readily formed Thioflavin T (ThT)-binding amyloid fibers with reproducible kinetics and concentration dependence (Kadnar et al., 2010). However, the same assay was much less reproducible for the full-length Rnq1 even though full-length Rnq1 amyloid fibers could be detected by transmission electron microscopy (TEM) similar to fibers formed by shorter Rnq1$_{WT}$ fragments (Vitrenko et al., 2007b; Kadnar et al., 2010).

Since shorter Rnq1 fragments work better for the ThT-binding assay, the ΔB1C2D3$_{T27P}$ construct that carries the Rnq1 NPD, the QG dipeptide repeats, the E helical region and QN4 was made and tested alongside the ΔB1C2D3$_{WT}$ (see Fig 1A for the map of the Rnq1 regions). Both proteins ran according to their expected size on SDS-PAGE gels, although ΔB1C2D3$_{T27P}$ ran slightly higher than ΔB1C2D3$_{WT}$ (Fig 4A). The same change in mobility was observed for proteins carrying the T27P mutations when they were isolated from yeast, indicative of a conformational alteration due to the mutation even in the soluble state (see Fig 2A). Incubation of both proteins in ThT resulted in a shift of the ThT excitation spectrum and increase of ThT fluorescence at 483 nm, indicative of amyloid formation (Fig 4B). Sigmoidal fluorescence kinetics was consistent with the presence of a rate-limiting nucleation step followed by a fiber growth phase for both ΔB1C2D3$_{T27P}$ and ΔB1C2D3$_{WT}$. The length of the lag phase was concentration dependent. Although there was considerable variation between duplicate samples, all showed sigmoidal kinetics and, at equal protein concentrations, the lag phase was similar for the ΔB1C2D3$_{T27P}$ and ΔB1C2D3$_{wt}$ samples.

For TEM, full-length Rnq_{T27P} was used and compared to full-length $Rnq1_{WT}$. Fibers were

assembled in the presence of ThT for ~70 hours and an increase of ThT fluorescence was

confirmed. TEM revealed >1μm-long non-branched fibers consisting of laterally associated

thinner fibrils for both $Rnq1_{T27P}$ and $Rnq1_{WT}$ (Fig 4C). However, the Rnq_{T27P} fibers were less

twisted and wider than the $Rnq1_{WT}$ fibers (30 – 50nm $vs \sim$ 20nm) indicating a difference in

conformation due to the mutation in the amyloid state.

The T27P mutation creates a barrier for the transmission of the prion state from

[PIN^+_{WT}] to $Rnq1_{T27P}$

Next, we tested if the T27P mutation affects the transmission of the prion conformation from

$Rnq1_{WT}$ to $Rnq1_{T27P}$. If the transmission block were absolute, all post-shuffle cells expressing

$Rnq1_{T27P}$ would be [pin^-]. The incomplete Pin$^-$ phenotype observed in our experiments is

consistent with either the [PIN^+_{T27P}] prion being absent in most but not all post-shuffle $Rnq1_{T27P}$

cells, or with the prion state being transmitted and maintained in $Rnq1_{T27P}$ cells but the mutation

making [PIN^+_{T27P}] extremely weak in its ability to induce [PSI^+]. To distinguish between these

possibilities, we changed the protocol from the one where the Pin state of the entire post-shuffle

cultures was scored (e.g. in Fig 2B) to a protocol testing the Pin phenotype of individual cells in

these cultures (see scheme in Fig 5A). The key difference (circled in red) is that now post-shuffle

cultures were colony purified before being tested for the Pin phenotype.

We found that only 1.5 – 2% of the colonies from the $Rnq1_{T27P}$ post-shuffle cultures

remained Pin$^+$, whereas the remaining 98 – 98.5% were clearly Pin$^-$ (Table 2; also see a sample

plate in Fig 5B, but note that this plate was chosen to show the presence of Pin$^+$ patches while

many plates had no Pin$^+$ patches). We additionally confirmed that these Pin$^+$ isolates did not

retain the *URA3 RNQ1$_{WT}$* plasmid. Also, as expected, the Pin$^+$ phenotype depended on the

presence of the *LEU2 RNQ1$_{T27P}$* plasmid. In the positive control Rnq1$_{WT}$ post transmission

cultures, all but one colony were found to be Pin$^+$. This is consistent with the >99.5% [*PIN$^+$$_{WT}$*]

stability in the original yeast strain before transformation with any plasmids and plasmid shuffle.

In the negative control post-transmission cultures carrying just the EV, all but one colony were

found to be Pin$^-$ (see Table 2 note). The difference between the frequency of Pin$^+$ cells in the

Rnq1$_{T27P}$ and EV cultures was statistically significant in all experiments (p<0.05). Finally, the 1.5

– 2% Pin$^+$ cells detected in the Rnq1$_{T27P}$ post-transmission cultures were not due to *de novo*

[*PIN$^+$$_{T27P}$*] appearance. This is because [*PIN$^+$$_{T27P}$*] does not form in *rnq1-Δ* cells without Rnq1$_{T27P}$

overexpression (see Fig 3B bottom row). Also, when the *LEU2 CEN RNQ1$_{T27P}$* or EV plasmids

were introduced into the [*pin$^-$*][*psi$^-$*] *rnq1-Δ* strain carrying *HIS3 GAL-SUP3NM-YFP*, no Pin$^+$

transformants were obtained among ~1,000 individual transformants tested, and the frequency of

Ade$^+$ colonies following attempted [*PSI$^+$*] induction was no greater in the *RNQ1$_{T27P}$* transformants

compared to EV even after several passages on plasmid-selective media.

Table 2. Only a small proportion of cells remain Pin$^+$ in the *RNQ1$_{T27P}$* expressing cultures

after shuffling out the *RNQ1$_{WT}$* plasmid.

Experiment	Construct after plasmid shuffle	Total tested	Pin$^+$	Average Pin$^+$ (%)	SEM (%)	T-test
# 1	*RNQ1$_{WT}$*	435	435	100	± 0	
	RNQ1$_{T27P}$	286	7	2.23	± 0.79	
	EV	197	0	0	± 0	p=0.019
# 2	*RNQ1$_{WT}$*	420	419	99.76	± 0.23	
	RNQ1$_{T27P}$	796	15	1.49	± 0.53	
	EV	504	1*	0.19	± 0.19	p=0.038

Table 2 Legend. Shown are results of two independent experiments. For each experiment,

colonies from at least 11 post-shuffle Rnq1$_{T27P}$ cultures were tested. T-tests are only shown

for Rnq1$_{T27P}$ *vs* EV (for both experiments $p<0.05$); for all other combinations $p<1.0e^{-25}$.

* The only Pin$^+$ clone obtained in the EV control in Experiment 2 appears to be a non-Rnq1-

based prion.

Taken together, these findings support the idea of a transmission barrier between Rnq1$_{WT}$

and Rnq1$_{T27P}$. If this were true, the proportion of Pin$^+$ cells in the initial post-shuffle cultures

(Table 2) would be determined by both the frequency of the transmission of the prion state from

Rnq1$_{WT}$ to Rnq1$_{T27P}$ and the stability of the resulting [*PIN$^+$$_{T27P}$*] prion.

Because the phenotype of the Rnq1$_{T27P}$ Pin$^+$ isolates varied in strength (compare the two

Rnq1$_{T27P}$ Pin$^+$ patches on Fig 5B) and was moderately to significantly reduced relative to Rnq1$_{WT}$

after shorter incubation times or if grown on SD-Ade at 30^{0}C (not shown), we hypothesized that

the newly forming [*PIN$^+$$_{T27P}$*]s were unstable, so the Pin$^+$ Rnq1$_{T27P}$ patches contained both

[*PIN$^+$$_{T27P}$*] and [*pin$^-$$_{T27P}$*] cells and the strength of the Pin$^+$ phenotype reflected the proportion of the

[*PIN$^+$$_{T27P}$*] cells. To test this, four Pin$^+$ Rnq1$_{T27P}$ isolates were colony purified from the patches

made on the SD-LeuHis plasmid-selective media prior to testing for the Pin$^+$ phenotype (circled

in green in Fig 5A) onto SD-LeuHis, and then individual colonies were patched on the same

media and tested for the retention of [*PIN$^+$$_{T27P}$*] by [*PSI$^+$*] induction. Indeed, in all four Pin$^+$

Rnq1$_{T27P}$ isolates both Pin$^+$ and Pin$^-$ cells were present, indicative of prion instability. Also, as

hypothesized, the strength of the Pin$^+$ phenotype in the original patch correlated with the fraction

of the Pin$^+$ cells found (Table 3).

Table 3. In the post-shuffle Pin+ Rnq1$_{T27P}$ isolates the [*PIN+$_{T27P}$*] prion is mitotically

unstable and the strength of the Pin+ phenotype correlates with fraction of cells retaining

the prion.

Strength of Pin+ in the original isolate (Round 1)	% Pin+ cells (Round 2)	Average ± SD
Very weak	3.3	18.4 ± 12.3
Weak	14.3	
Moderate	27.3	
Moderate	28.6	

Table 3 Legend. See text and Fig 5A for experimental design (green circle indicates a step

at which Pin+ isolates were colony purified). The strength of the Pin+ phenotype in post-

shuffle Rnq1$_{T27P}$ isolates was determined by growth on SD-Ade and SEt-Ade media at

30^{0}C and SD-Ade media at 20^{0}C after [*PSI+*] induction. "Moderate", "weak" and "very weak"

corresponds to the Pin+ phenotypes compared to similarly obtained post-shuffle Rnq1$_{WT}$

isolates where the Pin+ phenotype is considered "strong" and 100% cells retained the

prion. Shown are percentages of Pin+ colonies obtained by colony purification of the

original Pin+ Rnq1$_{T27P}$ isolates, as described in Results.

This is reminiscent of what happens when PrP^{Sc} prions cross the inter-species prion

transmission barriers (reviewed in Baskakov, 2014) and when Rnq1 fragments encompassing

different QN regions of the PD cross intraspecies transmission barriers in yeast (Kadnar et al.,

2010). In these cases, crossing the barrier usually leads to poorly propagating prion variants, and

stable prion variants are only detected in later passages either after selection and/or

conformational adaptation. Thus, we asked if the unstable [*PIN+$_{T27P}$*] prions that formed during

co-expression of Rnq1$_{T27P}$ with Rnq1$_{WT}$ in [PIN^+_{WT}] cells would eventually mature into more stable

ones. The stability of [PIN^+_{T27P}] was first scored in 13 Pin$^+$ Rnq1$_{T27P}$ post-shuffle isolates as

described above (Fig 5A, green circle), but this time Pin$^+$ isolates were chosen randomly and ~84

colonies were scored for each isolate to estimate the stability more accurately (Round 2 in Fig

5C). Then, 21 [PIN^+_{T27P}] patches obtained in this round were again colony purified from pre-

induction SD-LeuHis and patched on SD-LeuHis for Round 3 of [PIN^+_{T27P}] stability analysis. This

was repeated with 42 Round 3 [PIN^+_{T27P}] patches for Round 4 analysis (Fig 5C). The number of

cell divisions between each round was equal and estimated to be ~19. As can be seen, while

[PIN^+_{T27P}] remained highly unstable in both Round 2 and Round 3 analyses, for some [PIN^+_{T27P}]

isolates in Round 3 stability increased significantly, with over 60% of cells retaining [PIN^+_{T27P}],

which was reflected by a moderate increase of the average stability in this round compared to

Round 2. In Round 4, however, no dramatic increase in stability was observed, suggesting that,

when [PIN^+_{T27P}] undergoes maturation without selection, prion variants characterized by different

stabilities can be formed and maintained.

To identify such prion variants, we repeated the analysis of post-shuffle [PIN^+_{T27P}]

stability, this time choosing only three independently obtained [PIN^+_{T27P}]s but keeping pedigree

data as separate sets and extending analysis beyond Round 4. In Pedigree 1 [PIN^+_{T27P}] was

extremely unstable early on but yielded an almost 100% stable [PIN^+_{T27P}] prion in Round 4, and

this prion remained stable in Round 5 (430/431 Pin$^+$ in 12 [PIN^+_{T27P}] patches tested; Fig 5D). This

[PIN^+_{T27P}] prion depended on the maintenance of the $RNQ1_{T27P}$ plasmid, was GuHCl-curable, and

promoted the induction of a typical array of [PSI^+] variants at a frequency of a "strong" [PIN^+_{WT}].

Such stable [PIN^+_{T27P}]s were occasionally detected in other experiments.

In Pedigree 2, [PIN^+_{T27P}] remained very unstable in Round 4 and this very low stability

was maintained in Rounds 5 and 6, indicative of the establishment of a prion variant not prone to

subsequent changes in heritability (Fig 5D). In Pedigree 3 [PIN^+_{T27P}] reached a moderate level of

stability, significantly higher than in Pedigree 2 (p<0.01; Fig 5D), and remained moderately stable

in most cultures in Rounds 5 and 6 but yielded occasional isolates with significantly lower or

higher heritable stability (red and green asterisks on Fig 5D), indicative of the establishment of a

prion variant that is capable of further conformational adaptation. Altogether, we interpret these

data as maturation of post-transmission [PIN^+_{T27P}]s to heritable prion variants with characteristic

stabilities that is mostly completed within ~60 cell divisions but may continue longer and go

through intermediate variants for some isolates.

Rnq1$_{T27P}$ can join the amyloid [PIN^+] aggregates formed by Rnq1$_{WT}$

The strength of transmission barriers is determined by two key factors: the efficiency of

transmission of the prion state from the donor prion, [PIN^+_{WT}], to the recipient protein, Rnq1$_{T27P}$,

and the ability of the recipient protein-based prion, [PIN^+_{T27P}], to propagate efficiently. The data

presented in the previous section indicate that the inefficient propagation of post-transmission

[PIN^+_{T27P}]s is definitely a contributing factor to the barrier. After the $RNQ1_{WT}$ plasmid was shuffled

out, [PIN^+_{T27P}] was very unstable in all isolates tested and further conformational adaptation was

required for [PIN^+_{T27P}] stability to increase (see Figs 5C and 5D). We now ask if extreme

instability of non-adapted [PIN^+_{T27P}] was likely the only reason for presence of only 1.5 - 2%

[PIN^+_{T27P}] cells in post-shuffle cultures or if other mechanisms were also contributing.

Here, we tested if Rnq1$_{T27P}$ could join the pre-existing [PIN^+_{WT}] prion aggregates because

a block at this step would inhibit the transmission of the prion state regardless of the resulting

stability of [PIN^+_{T27P}] prions. The Rnq1$_{WT}$-CFP and Rnq1$_{T27P}$-YFP were co-expressed in a

[PIN^+_{WT}] strain from CEN plasmids where constructs were controlled by the $CUP1$ promoter. To

make sure that no significant $de\ novo$ induction of [PIN^+_{T27P}] has occurred before cell analysis,

cultures of co-transformants were grown to mid-log in the plasmid-selective no-copper media,

and CuSO$_4$ was added to 5μM only 2 hours prior to microscopy (see Materials and Methods). At

this low concentration of CuSO$_4$, Rnq1 is expressed at levels approximately equal to those of

chromosomally encoded Rnq1 (Fig S3). As shown in Fig 6A, Rnq1$_{WT}$-CFP and Rnq1$_{T27P}$-YFP co-

localized in bright fluorescent foci in most cells, indicative of Rnq1$_{T27P}$ being able to join [*PIN$^+$$_{WT}$*]

aggregates. We confirmed that Rnq1$_{T27P}$-YFP joins [*PIN$^+$*] aggregates just as efficiently as

Rnq1$_{WT}$-CFP (Figs S2A and S2B). Co-localization was retained even after prolonged incubation

or a higher-level expression of the constructs (5μM CuSO$_4$ overnight, 20μMCuSO$_4$ for 2 hours or

overnight, or 50μMCuSO$_4$ for 4 hours; not shown). The appearance of Rnq1$_{T27P}$-YFP foci was

indistinguishable from that of Rnq1$_{WT}$-CFP foci and corresponded to a foci pattern characteristic

of the "multi-dot high" [*PIN$^+$*] variant (Bradley and Liebman, 2003) used for these experiments.

Also, there was no evidence suggesting that Rnq1$_{T27P}$-YFP aggregates were just seeded by

Rnq1$_{WT}$ but then formed next to pre-existing [*PIN$^+$*] aggregates not incorporating Rnq1$_{WT}$: the

overlap of Rnq1$_{T27P}$-YFP and Rnq1$_{WT}$-CFP foci was usually complete in the focal plane imaged

(Figs 6A, S2A, and S2B) and in the focal planes immediately above and below. The foci with co-

localized Rnq1$_{WT}$-CFP and Rnq1$_{T27P}$-YFP were not sensitive to 1,6 hexanediol (Hex; Fig 6A),

with Rnq1$_{T27P}$ retaininged after the treatment, suggesting that the fluorescently tagged proteins

were joining the typical amyloid [*PIN$^+$$_{WT}$*] aggregates (see also Fig 8A for ThT staining). Thus, the

transmission barrier is not due to the inability of Rnq1$_{T27P}$ to join pre-existing [*PIN$^+$$_{WT}$*] aggregates

formed by Rnq1$_{WT}$ or to convert into amyloid.

The co-expression of Rnq1$_{T27P}$ does not lead to loss of [*PIN$^+$$_{WT}$*]

We then asked if introduction of the Rnq1$_{T27P}$ expressing plasmid led to a rapid loss of [*PIN$^+$$_{WT}$*]

prior to the plasmid shuffle. One possibility is that incorporation of Rnq1$_{T27P}$ into the [*PIN$^+$$_{WT}$*]

aggregates "poisoned" these aggregates and blocked the prion propagation. This hypothesis

was used to explain the curing of the [*URE3*] prion upon the overexpression of Ure2-GFP or

some Ure2 fragments (Edskes et al., 1999) and curing of [*PSI⁺*] by dominant "[*PSI⁺*] no more"

PNM mutations within Sup35 PD (Kochneva-Pervukhova et al., 1998; Derkatch et al., 1999;

DiSalvo et al., 2011; Pei et al., 2017).

However, we have found that transforming the [PIN^+_{WT}] strain used in the genetic screen

([*PIN⁺*] [*psi⁻*] *rnq1-Δ* carrying the *CEN URA3 RNQ1$_{WT}$* [*PIN⁺*] maintainer and the *CEN HIS3 GAL-*

SUP35NM-YFP [*PSI⁺*] inducer; see Fig 1B) with the *CEN LEU2 RNQ1$_{T27P}$* plasmid did not lead to

loss of the [*PIN⁺*] prion. All 240 transformants tested in this "no shuffle" experiment remained Pin⁺

even after four more passages on media selective for all the plasmids. Furthermore, no reduction

of [*PSI⁺*] induction was observed in such cells compared to the control transformants with *CEN*

LEU2 RNQ1$_{WT}$ or *CEN LEU2* EV in the nonsense suppression test (Fig 6B) and when *de novo*

induction of [*PSI⁺*] was measured as a fraction of cells with Sup35NM-YFP aggregates during

overexpression of the Sup35NM-YFP construct (regardless of the plasmid introduced, ~50%

cells had line-, ring-, and dot-shaped aggregates after 48 hours of growth on SGal-UraLeuHis

media; data not shown).

Also, there was no indication of incompatibility of Rnq1$_{WT}$ and Rnq1$_{T27P}$: there was no

growth inhibition for the *RNQ1$_{T27P}$* carrying cells compared to the *RNQ1$_{WT}$* and EV transformants

(Fig 6B). The stability of the [PIN^+_{WT}] maintainer plasmid, *CEN URA3 RNQ1$_{WT}$*, was the same in

the *CEN LEU2 RNQ1$_{T27P}$*, *CEN LEU2 RNQ1$_{WT}$*, and *CEN LEU2* EV transformants (AVE ± SD are

38.4 ± 11.3%, 41.9 ± 10.3%, and 41.6 ± 5.8%, respectively). Also, there was no difference in the

stability of the *CEN LEU2 RNQ1$_{T27P}$*, *CEN LEU2 RNQ1$_{WT}$*, and *CEN LEU2* EV (AVE ± SD are

39.9 ± 7.8%, 34.8 ± 10.2%, and 38.0 ± 7.7%, respectively; see Materials and Methods for the

description of the test).

We also analyzed if similar or different arrays of [*PSI⁺*] variants were induced by

Sup35NM-YFP overexpression in cells co-expressing Rnq1$_{WT}$ and Rnq1$_{T27P}$ compared to cells

expressing two copies of Rnq1$_{WT}$. For either the cells co-expressing Rnq1$_{WT}$ and Rnq1$_{T27P}$ or

cells expressing two copies of Rnq1$_{WT}$ most [*PSI*$^+$] variants were moderate (76.1 and 89.7%,

respectively), and strong [*PSI*$^+$] variants were rare (~3%). However, the percentage of weak

[*PSI*$^+$] variants was significantly higher in cells co-expressing Rnq1$_{T27P}$ and Rnq1$_{WT}$ (21.0 *vs*

6.5%; p <0.01; Table 4).

Table 4. Co-expression of Rnq1$_{T27P}$ and Rnq1$_{WT}$ leads to a slight shift towards the

induction of weak [*PSI*$^+$] variants compared to expression of two copies of Rnq1$_{WT}$.

Plasmid introduced	[*PSI*$^+$] variants: number isolated (% ± SD)			
	Weak	Medium	Strong	Total
RNQ1$_{WT}$	9 (6.5 ± 5.5)	125 (89.7 ± 8.9)	5 (3.8 ± 6.6)	139 (100)
RNQ1$_{T27P}$	15 (21.1 ± 5.1)	54 (76.1 ± 4.9)	2 (2.8 ± 3.2)	71 (100)
EV	2 (3.1 ± 3.6)	64 (96.9 ± 3.6)	0 (0 ± 0)	66 (100)

Table 4 Legend. Following the induction of [*PSI*$^+$] by high level expression of Sup35NM-

YFP on SGal-UraLeuHis media selective for all plasmids, yeast were replica plated on

glucose SD-Ade media not selective for any plasmid and incubated at 20°C for 25 days.

Then Ade$^+$ cells were scraped from the SD-Ade plates and colony purified on YPD.

Random individual colonies were patched on YPD before color developed. [*PSI*$^+$]s isolated

from at least four independent transformants carrying the indicated plasmids were

analyzed by the color test on YPD, growth on adenineless media and curability by GuHCl.

Results of all three tests were consistent for ~99% [*PSI*$^+$] variants.

Thus, the barrier for the transmission of the [*PIN*$^+$$_{WT}$] prion state to Rnq1$_{T27P}$ is not due to

Rnq1$_{T27P}$ being incompatible with Rnq1$_{WT}$ co-expression or "poisoning" of the [*PIN*$^+$$_{WT}$] prion

aggregates, but upon joining [*PIN+_{WT}*] aggregates Rnq1$_{T27P}$ changes the ability of the Rnq1-

based prion to promote the induction of particular [*PSI+*] variants.

The transmission barrier created by the T27P mutation in the NPD is eliminated by

deletion of the first two QN-rich regions in Rnq1 PD

Data presented in the previous sections indicate that the major reason for the transmission

barrier is the inability of Rnq1$_{T27P}$ to attain an efficiently propagating prion conformation when

seeded by the [*PIN+_{WT}*] variant used in our studies. We now ask if other conformational variants

of Rnq1-based prions would be more acceptable donors of the prion state for Rnq1$_{T27P}$. In our

earlier studies we uncovered transmission barriers with subsequent conformational adaptation

for the transfer of prion conformations between the [*PIN+_{WT}*] variant used in the current study and

Rnq1 fragments lacking one or more QN-rich regions in the PD (Kadnar et al., 2010; see Fig 1A).

The stable prion variants that were formed by these Rnq1 fragments after crossing the

transmission barrier from full-length Rnq1 (Rnq1$_{FL}$) and consequent conformational adaptation

were collectively named [*mini-PIN+*]s and were confirmed to have prion conformations different

from those of [*PIN+_{FL}*]. Thus, we tested if the transmission barrier to Rnq1$_{T27P}$ was reduced when

these stable [*mini-PIN+*]s not carrying the T27P mutation were used as donors of the prion

conformation instead of full length [*PIN+_{WT}*]. From now on, the **nomenclature** will include both

the presence of full-length Rnq1 or of a Rnq1 fragment, as well as the WT or T27P allele (e.g.

[*PIN+_{FL-WT}*] or [*PIN+_{FL-T27P}*]). For the Rnq1 fragments, the nomenclature reflects the deleted part of

the PD, which includes the QN-rich region(s) numbered as in Fig 1A and an adjacent helical

stretch(es) indicated in Fig 1A by letters (e.g. [*PIN+_{ΔB1C2-WT}*] or [*PIN+_{ΔB1C2-T27P}*]).

The plasmid shuffle experiment described in Fig 1B was repeated using stable [*mini-*

PIN+_{WT}] stains instead of [*PIN+_{FL-WT}*]. According to the nonsense suppression-based [*PSI+*]

induction test, transmission of the prion state to Rnq1$_{FL-T27P}$ from [$PIN^+_{\Delta B1C2-WT}$] and [$PIN^+_{\Delta B1-WT}$]

but not from [$PIN^+_{\Delta 2D-WT}$] or [$PIN^+_{\Delta 3E-WT}$] was considerably more efficient compared to

transmission from [PIN^+_{FL-WT}] (Fig 7A). In the case of transmission from [$PIN^+_{\Delta B1C2-WT}$], a colony

purification experiment done as described in Fig 5A was used to confirm that growth on

adenineless media in Fig 7A was indeed due to a higher fraction of [$PIN^+_{FL-T27P}$] cells in the "post-

shuffle" cultures (Fig 7B; the difference is statistically significant, p<0.01). This supports the

hypothesis that the T27P mutation in the NPD changes the range of prion conformations

permissible for PD and that [$PIN^+_{\Delta B1C2-WT}$] and [$PIN^+_{\Delta B1-WT}$] fall into this range of acceptable

conformations while [PIN^+_{FL-WT}] does not.

Considering the indication that deletion of the B1C2 region of the Rnq1 PD reduced the

transmission barrier, we asked if this region fully determined the inability of Rnq1$_{FL-T27P}$ to take on

the prion conformation compatible with [PIN^+_{FL-WT}]. The T27P mutation was introduced into the

Rnq1 fragment lacking the B1C2 region (the Rnq1$_{\Delta B1C2-T27P}$ construct), and transmission of the

prion state to this construct was assessed using [$PIN^+_{\Delta B1C2-WT}$] as a donor of the prion state. We

found that a significantly higher fraction of Pin$^+$ cells was present in the "post-transmission"

cultures when both the donor and recipient of the prion state lacked the B1C2 region (Fig 7C;

p<0.01). However, no subsequent change of mitotic stability was observed following such

transmissions: after four rounds of consecutive colony purifications done as in Fig 5C, the

[$PIN^+_{\Delta B1C2-T27P}$] isolates retained the same stability as immediately after the transmission (Fig 7C,

no statistically significant difference between R1 and R2 through R4). Furthermore, one more

round of analysis of R4 [$PIN^+_{\Delta B1C2-T27P}$] isolates with the highest fraction of prion-retaining cells

did not indicate that these isolates underwent a heritable conformational shift leading to higher

mitotic stability (not shown). One explanation for these data is that the T27P mutation limits the

fragmentation of all or most [$PIN^+_{\Delta B1C2-T27P}$] prion variants thus preventing them from propagating

efficiently enough to achieve high mitotic stability. The alternative explanation is that there is no

transmission barrier between [$PIN^+_{\Delta B1C2-WT}$] and Rnq1$_{\Delta B1C2-T27P}$. In this case, transmission of the

prion state from [$PIN^+_{\Delta B1C2-WT}$] to Rnq1$_{\Delta B1C2-T27P}$ will result in all cells attaining the same heritable

[$PIN^+_{\Delta B1C2-WT}$] prion conformation with a particular mitotic stability (that happens to be ~45% after

~19 cell generations), and then this [$PIN^+_{\Delta B1C2-T27P}$] variant propagates faithfully without

undergoing any further conformational adaptation. Because there is no selection favoring the

maintenance of the [$PIN^+_{\Delta B1C2-T27P}$] prion in our experimental system, we do not expect to detect

rare switches to mitotically stable variants within the relatively small sample sizes tested. Also,

the proportion of Pin$^+$ cells in Round 1 analysis is expected to reflect the 100% efficient

transmission of the prion state between [$PIN^+_{\Delta B1C2-WT}$] and Rnq1$_{\Delta B1C2-T27P}$ and then the loss of

[$PIN^+_{\Delta B1C2-T27P}$] in cells divisions between the elimination of the $RNQ1_{\Delta B1C2-WT}$ plasmid and the

start of the analysis of "post-shuffle" cultures, which is ~18-20 cells divisions, i.e. similar to the

~19 number of divisions between subsequent rounds.

699   To distinguish between these two explanations, transmission of the prion state to

Rnq1$_{\Delta B1C2-T27P}$ was attempted using [PIN^+_{FL-WT}] as a donor. Data in Fig 7D indicate the existence

of a strong transmission barrier, similar to that between [PIN^+_{FL-WT}] and Rnq1$_{FL-T27P}$, and

subsequent conformational adaptation which allowed for the establishment of [$PIN^+_{\Delta B1C2-T27P}$]

variants with different levels of mitotic stability, including high mitotic stability that was maintained

in subsequent rounds of analysis (see asterisks in Fig 7D). This excludes the explanation that

the T27P mutation makes [$PIN^+_{\Delta B1C2-T27P}$] variants intrinsically unstable and supports the

alternative explanation that the T27P mutation in the NPD creates a transmission barrier through

the effect on the conformational flexibility of the B1C2 region of the PD. Comparison of data

presented in Figs 7C and 7D supports the idea that when both the donor and the recipient lack

this region, the mutant attains the prion conformation without a barrier but the mutation slightly

inhibits the stability of the resulting [$PIN^+_{\Delta B1C2-T27P}$] variant. The deletion of the B1C2 region only

in the T27P carrying recipient and not in the non-mutant donor is not sufficient to eliminate the

transmission barrier. However, in transmissions from [PIN^+_{FL-WT}], the subsequent increase of

mitotic stability was significantly faster for [$PIN^+_{\Delta B1C2-T27P}$] compared to [$PIN^+_{FL-T27P}$] (Fig 7D),

indicative of the effect of the B1C2 region of PD on the conformational adaptation properties of

T27P mutants.

Because the [PSI^+] induction test in Fig 7A indicated that deletion of only the B1 region

within the PD could also reduce the transmission barrier, we asked if the first QN-rich region of

the PD, QN1, was solely responsible for the effect on both the WT to T27P transmission barrier

and conformational adaptation of the T27P carrying prions observed for ΔB1C2 fragments. This

is particularly interesting because QN1 is very short, only 12aa (NSNNNNQQGQNQ). Thus, the

T27P mutation was introduced into the Rnq1 fragment lacking QN1 but retaining QN2 (the

Rnq1$_{\Delta B1-T27P}$ construct) and the [$PIN^+_{\Delta B1-WT}$] donor was used to transmit the prion state to this

construct. Like in the similar experiments with WT and T27P carrying ΔB1C2 Rnq1 fragments,

the fraction of [$PIN^+_{\Delta B1-T27P}$] cells in the "post-shuffle" cultures was significantly increased

compared to the transmissions between FL constructs (p<0.01; compare Fig 7E with 7D), even

though the fraction of Pin$^+$ cells was slightly lower in [$PIN^+_{\Delta B1-T27P}$] than in [$PIN^+_{\Delta B1C2-T27P}$] samples

(p<0.05; compare Figs 7C and 7E). However, in the [$PIN^+_{\Delta B1-T27P}$] isolates, subsequent

appearance of [$PIN^+_{\Delta B1-T27P}$]s with high mitotic stability, indicative of the ongoing conformational

adaptation, led us to conclude that removal of only the B1 region in both the donor and the

recipient did not eliminate the transmission barrier created by the T27P mutation. This finding

suggested that the QN2 region of the Rnq1 PD was also involved in the interaction of the PD

with the T27P mutation in the NPD. Yet, similar experiments attempting to transmit the prion

state from the [$PIN^+_{\Delta 2D-T27P}$] donor to the Rnq1$_{\Delta 2D-T27P}$ recipient did not suggest that deleting the

QN2 region reduced the transmission barrier or led to the establishment of the T27P carrying

prions not prone to further conformational adaptation (Fig 7F). Thus, the NPD exerts its effect on

the QN1 and QN2 regions of the PD acting in concert rather than individually. Finally, consistent

with data shown in Fig 7A, we obtained no evidence suggesting that the deletion of the QN3

region of the Rnq1 PD (Δ3E) could remove the transmission barrier created by the T27P

mutation or affect conformational adaptation of [$PIN^+_{\Delta 3E\text{-}T27P}$] prion variants after crossing the

barrier (Fig 7G).

Unlike Rnq1$_{WT}$-YFP, Rnq1$_{T27P}$-YFP readily forms liquid-like droplets when

expressed in [pin^-] cells

While Rnq1 aggregation in [PIN^+] strains can be visualized by expression of Rnq1$_{WT}$, its PD, or

PD fragments fused to C-terminal fluorescent tags from the GFP family (Sondheimer and

Lindquist, 2000; Derkatch et al., 2001; Osherovich and Weissman, 2001; Vitrenko et al., 2007b;

Kadnar et al., 2010), overexpression of Rnq1$_{WT}$ fused to such tags (e.g. Rnq1$_{WT}$-GFP) in [pin^-]

strains does not lead to the formation of visible aggregates and to the *de novo* appearance of

[PIN^+] (Derkatch et al., 2001; Kadnar et al., 2010). One possible explanation is that the extreme

C-terminus of the Rnq1 PD encompassing the last QN-rich region, QN4 (see Fig 1A), is

important during the step of the *de novo* [PIN^+] prion formation, and bulky C-terminal tags

interfere with this process (Galliamov et al., 2024). However, once [PIN^+] is formed, other QN-

rich regions within the RNQ1 PD get engaged in the maintenance of the prion structure and

blocking of QN4 by a tag is not critical for the prion maintenance (Kadnar et al., 2010).

Because our data indicate that the T27P mutation changes the conformational flexibility

of the Rnq1 PD through particular QN-rich regions, we asked if, in the presence of T27P, the C-

terminally tagged constructs retain aggregation properties despite the tag partially blocking the

QN4 region. As seen in Fig 8A, in contrast to Rnq1$_{WT}$, when Rnq1$_{T27P}$-YFP is expressed in [*pin$^-$*]

cells, it readily forms dot aggregates, one or several dots per cell. Such dot-like aggregates were

formed in ~40% of cells even when Rnq1$_{T27P}$-YFP was expressed briefly and at close to

physiological levels of expression (5μM CuSO$_4$ for 2 hours; Figs 8B and 8C; also see Fig S3 for

protein expression).

These dot aggregates are clearly distinct from ribbon-like aggregates of the *de novo*

forming [*PIN$^+_{WT}$*]s when GFP-tagged Rnq1 is overexpressed in [*URE3*] cells (Derkatch et al.,

2001). Therefore, we tested if Rnq1$_{T27P}$-YFP aggregates forming in [*pin$^-$*] cells are amyloid, and

found they are not because the aggregates are not stained by Thioflavin T (Fig 8A). Furthermore,

these aggregates are dissolved by a short-term Hexanediol (Hex) treatment, suggesting that they

are liquid-like droplets (Figs 8B and 8C). The presence of Rnq1$_{WT}$ is not required for Rnq1$_{T27P}$ to

form liquid-like droplets as they also form in *rnq1-Δ* cells (Fig 8D). Finally, Rnq1$_{WT}$-CFP does not

join the Rnq1$_{T27P}$-YFP when co-expressed in [*pin$^-$*] cells (Fig 8E). This further supports the idea

that the NPD mutation T27P modulates aggregation and conformational properties of QN-rich

regions within the PD of Rnq1.

Discussion

The T27P mutation in the NPD of Rnq1 creates a barrier for the transmission of the

[*PIN$^+$*] prion state from Rnq1$_{WT}$

Most known prion-forming proteins have chimeric organization containing: 1) a functional

non-prion domain, NPD, that is not essential for prion formation and is required for the protein to

perform its main cellular function; 2) a prion domain, PD, which is sufficient for prionization and

can form prions in the absence of NPD and even when attached to other proteins. At first glance,

Rnq1 shares such chimeric organization except that its NPD is very short, compared to its PD,

which constitutes up to 2/3 of the protein and has a very complex organization (see Introduction

and Fig 1A). The length of the NPD is ~132 – 153 aa but is not known exactly because *S.*

cerevisiae Rnq1, as well as its potential homologs from other species have no known function

and are poorly conserved (ILD unpublished observations).

We aimed at understanding the structure - function of Rnq1 related to its prion-forming

abilities. Two approaches were used. Previously, we performed extensive deletion analysis of all

potential structural features within the PD (Kadnar et al., 2010). Here, we used random

mutagenesis of *RNQ1*. We mutagenized both the PD and NPD because, first, the exact length of

the NPD is not known and, second, several lines of evidence indicated that the NPD is involved

in [*PIN+*] formation and maintenance (see below). The genetic screen was for the loss of the Pin+

phenotype, *i.e.* the ability of [*PIN+*] to facilitate the *de novo* induction of [*PSI+*]. This readout

allowed screening for mutations in the context of the complete Rnq1 not fused to reporters that

were previously shown to modify its prion forming ability (Sondheimer and Lindquist, 2000;

Derkatch et al., 2001; Bardill and True, 2009). Here we describe the $Rnq1_{T27P}$ mutant obtained in

this screen. We propose that, through interactions with specific QN-rich regions in the PD, this

NPD mutation alters the array of conformations the prion domain can attain and thus creates a

transmission barrier with a [*PIN+WT*] prion formed by non-mutant Rnq1. The mutation also alters

the aggregation properties of Rnq1 by promoting the formation of liquid-like droplets when

$Rnq1_{T27P}$ is fused to a C-terminal YFP tag.

The T27P mutation led to the loss of the [*PIN+*] prion in 98 – 98.5% of the cells when

$Rnq1_{T27P}$ was substituted for $Rnq1_{WT}$ in a [*PIN+WT*] strain (Table 2). We concluded that this occurs

because the mutation creates a barrier for the transmission of the prion state from $Rnq1_{WT}$ to

$Rnq1_{T27P}$ after not finding support for the alternative explanation that $Rnq1_{T27P}$ is intrinsically

defective in prion formation and propagation. Indeed, $Rnq1_{T27P}$ can form amyloid *in vitro* (Fig 4C).

Also, transient overexpression of $Rnq1_{T27P}$ in a *rnq1-Δ* strain leads to a heritable Pin+ phenotype

indicative of the *de novo* formation of [PIN^+_{T27P}] (Fig 3). Finally, a [PIN^+_{T27P}] variant with mitotic

stability of >99.9% was obtained upon prolonged mitotic propagation of an unstable [PIN^+_{T27P}]

retrieved immediately after transmission of the prion state from [PIN^+_{WT}] (Fig 5D).

The concept of transmission barriers dates to observations that experimental transmission of

scrapie between different mammalian species is frequently far less efficient than within the same

species (Pattison et al., 1965). These barriers were then attributed to differences in sequences of

the PrP protein in different species (Scott et al., 1989). Interspecies transmission barriers were

also uncovered for [PSI^+] and [*URE3*] (Santoso et al., 2000; Chen et al., 2007, 2010; Edskes et

al., 2009; Afanasieva et al., 2011; Sharma et al., 2016; Wickner et al., 2019). Importantly, the

analysis of these barriers focused on the conformational constraints resulting from sequence

differences **exclusively within the PDs** of these proteins. In closely related *Saccharomyces*

sensu stricto species, transmission barriers formed even though PDs are highly homologous and

carry only a few individual amino acid changes (Santoso et al., 2000; Chen et al., 2007, 2010).

Furthermore, mutagenesis of Sup35 and Rnq1 and analysis of Sup35 from wild isolates revealed

that transmission barriers can be created within *S. cerevisiae* by single mutations in the PDs

(DePace et al., 1998; Bardill and True, 2009; Bateman and Wickner 2012).

However, differences in amino acid sequences in the prion-forming regions are not the only

contributing factor to transmission barriers. Studies of mammalian prions eventually concluded

that the efficiency of interspecies transmissions is determined by both the PrP sequence and

particular PrP^{Sc} conformations in the donor species (Hill et al., 1997; Hill and Collinge 2001;

Collinge and Clarke, 2007; Collinge, 2012). Our analysis of the four QN-rich regions within the

PD of Rnq1 showed that while any two were generally sufficient for the maintenance of the [PIN^+]

prion, deletions of individual QN-rich regions created transmission barriers, and the possibility of

transmission across these barriers depended on the presence of at least one common QN-rich

region between the donor and recipient constructs. This underscored both the importance of

identical amino-acid stretches within the PD for prion transmission and the contribution of

conformational differences created by other parts of the PD (Kadnar et al., 2010). Our current

study contributes to understanding exclusively conformation-based transmission barriers

because the sequence of the entire PD is identical for $Rnq1_{WT}$ and $Rnq1_{T27P}$.

We found no evidence that the initial transmission of the prion state from [PIN^+_{WT}] to $Rnq1_{T27P}$

was blocked. $Rnq1_{T27P}$ joins [PIN^+_{WT}] amyloid aggregates as efficiently as $Rnq1_{WT}$ and does not

"poison" them (Figs 6A, 6B, 8A, 8B, and S2). But when we scored individual [PIN^+_{T27P}] isolates

from post-transmission cultures for prion maintenance, the striking observation is that all post-

transmission [PIN^+_{T27P}] isolates were extremely unstable. Scoring lineages of [PIN^+_{T27P}] isolates

for prion maintenance following a controlled number of mitotic divisions between rounds of

analysis revealed that, while on average mitotic stability of [PIN^+_{T27P}] increased compared to the

first round of analysis (Fig 5C), mitotic growth also led to the establishment of [PIN^+_{T27P}] variants

with distinct mitotic stabilities. Along with highly stable [PIN^+_{T27P}]s (>99.9% maintenance after 19

cell divisions) we obtained lineages with a heritable stability of ~20% (Fig 5D). Importantly, like

[PIN^+_{WT}] (Derkatch et al., 2000), [PIN^+_{T27P}] did not provide a detectable growth advantage relative

to [pin^-_{T27P}], so there was no selection for cells with more efficiently propagating [PIN^+_{T27P}]. Thus,

mitotic stability is just a quantitative characteristic that can be used to distinguish different

[PIN^+_{T27P}] variants established after crossing the transmission barrier.

Two models could explain the appearance of heritable [PIN^+_{T27P}] variants after multiple

mitotic divisions in the presence of [PIN^+_{WT}] (Fig 9). Both postulate that the T27P mutation puts

conformational constraints on the PD of $Rnq1_{T27P}$, so it cannot form a stable [PIN^+] that is

conformationally similar to [PIN^+_{WT}] even though $Rnq1_{T27P}$ can form stable prions *de novo*. The

first model proposes that multiple [PIN^+_{T27P}] variants are formed in the same cell after

transmission, and then daughter cells inherit them randomly, with a higher chance of inheriting

variants that produce more seeds (propagons) and are thus more stable in mitotic divisions. This

would lead to the rapid disappearance of the least stable variants in the population. However,

following individual [PIN^+_{T27P}] pedigrees allowed us to detect such unstable variants. In support of

this model, the co-existence of different prion variants in the same cell for several generations

has been confirmed for [PSI^+] (Norton et al., 2024). The **second model** predicts that the

transmission barrier leads to formation of a particular [PIN^+_{T27P}] variant, but this variant is not

conformationally stable and undergoes frequent conformational switches until more energetically

favorable states are attained. These stable conformations may have different, even low, mitotic

stabilities, but they are not changing frequently anymore. The process of such conformational

adaptation may involve multiple steps and some [PIN^+_{T27P}] variants may end up in efficiently

propagating but "undecided" states that have no preferred path towards conformational

stabilization. Our detection of such "undecided" [PIN^+_{T27P}] lineages that produce both more and

less mitotically stable variants even after 100 cell divisions (Fig 5D and data not shown) supports

the conformational adaptation model, but we do not see the models as strictly mutually exclusive.

Conformational adaptation in the absence of PD mutations has been proposed e.g. for *de novo*

induced [PIN^+] and [PSI^+], for permanent [PSI^+] variant switching in different chaperone

environments, and for conversion of *in vitro* made PrP[Res] into PrP[Sc] (Derkatch et al., 2001;

Sharma and Liebman, 2012; Baskakov, 2014; King, 2022).

The T27P mutation specifically interacts with the B1C2 part of the PD since its deletion

dramatically increased prion transmission to mutant constructs (Fig 7). Actually, data indicate

that there is no transmission barrier between [$PIN^+_{\Delta B1C2-WT}$] and Rnq1$_{\Delta B1C2-T27P}$. Even though the

mitotic stability of [$PIN^+_{\Delta B1C2-T27P}$] was not very high, this level of stability was clearly heritable

during subsequent mitotic growth, indicative of the establishment of a [$PIN^+_{\Delta B1C2-T27P}$] variant not

undergoing further conformational adaptation (Fig 7C). The alternative explanation that a prion

variant used as a [$PIN^+_{\Delta B1C2\text{-}WT}$] donor is more compatible with prion conformations that could be

attained by the T27P mutants is unlikely since deletion of B1C2 also affects the transmission

from [$PIN^+_{FL\text{-}WT}$] to the Rnq1$_{\Delta B1C2\text{-}T27P}$ mutant (Fig 7D). While initially the barrier was strong, mitotic

stability of the [$PIN^+_{\Delta B1C2\text{-}T27P}$] isolates increased much faster compared to [$PIN^+_{FL\text{-}T27P}$]s obtained

from the same donor. Thus, despite carrying the same mutation, Rnq1$_{FL\text{-}T27P}$ and Rnq1$_{\Delta B1C2\text{-}T27P}$

form different prion variants. Analysis of transmissions between Rnq1 fragments carrying

deletions of only QN1 (ΔB1) or QN2 (Δ2D) regions indicates that simultaneous deletion of both

regions is required for the lifting of the transmission barrier created by the T27P mutation, but

removing just B1 encompassing a 12 aa QN-rich region is sufficient to reduce the strength of the

barrier.

Another indication of a difference in amyloid conformations attained by Rnq1$_{WT}$ and Rnq1$_{T27P}$

is that amyloid fibers spontaneously formed by these proteins *in vitro* look different in TEM: the

bundles of Rnq1$_{WT}$ fibrils are tight and twisted whereas Rnq1$_{T27P}$ fibers are more relaxed and not

twisted (Fig 4C).

However, the T27P mutation affects more than the array of prion conformations that are

possible for the [PIN^+_{T27P}] prion. It changes the conformation of the protein even in the soluble

state, indicated by a slight but reproducible increase of mobility of SDS-PAGE gels when purified

from yeast or bacteria (Figs 2A and 4A). Indeed, T27P opens an alternative aggregation pathway

for the Rnq1 protein. The Rnq1$_{WT,}$ even when overexpressed, is not prone to aggregate when

fused to C-terminal GFP family tags unless a heterologous QN-rich amyloid aggregate is present

(Derkatch et al., 2001). This may be because a bulky tag blocks QN4, the last QN-rich

determinant in the PD, which has recently been shown to initiate [PIN^+] formation (Galliamov et

al., 2024, 2025), so heterologous amyloid seeds are needed to overcome this constraint. But

Rnq1$_{T27P}$ attached to YFP readily forms liquid-like droplets at physiological levels of expression

without the need for heterologous prions (Fig 8). This finding strengthens the evidence that the

T27P mutation affects aggregation properties of Rnq1.

Is the only function of the Rnq1 "NPD" to regulate the aggregation of Rnq1 PD?

Transmission barriers created by mutations **within the PDs** of yeast prions (see above) as well

as the existence of different prion variants can be explained by the parallel-in-register model of

protein molecules in amyloid fibrils (Wickner et al., 2008; 2011a). According to this model, prion

variants are the result of differences in the length of amyloid cores formed by PDs and relative

positioning of amyloid stretches due to turns within regions not involved in β-strand formation.

For Rnq1, amyloid formation was detected in the regions roughly corresponding to QN-rich

regions, but varies in different prion variants (Galliamov et al., 2024, 2025). So, mutations within

or deletions of these regions are expected to create conformational constrains leading to

different prion variants. But the T27P mutation is located **outside** the currently defined Rnq1 PD,

so its effect on variants of the Rnq1-based prions is not well explained by this model.

How could the T27P mutation in the NPD create a transmission barrier? Because NPDs

engage in interactions with cellular partners, the absence or inactivation of NPD can block these

interactions and thereby increase *de novo* prion formation (Derkatch et al., 1996; Maddelein and

Wickner, 1999). Increasing such interactions would, reciprocally, decrease prion formation

(Derkatch et al., 1998). However, if the prion-forming protein is indeed chimeric and the NPD and

PD domains fold independently, mutations that just change functionality of the NPD are not

expected to impose **specific** conformational constrains on the PD and should not lead to

transmission barriers. An alternative hypothesis is that in Rnq1 the N-terminal domain, presumed

NPD, regulates the conformation and aggregation properties of the PD.

Indeed, this hypothesis helps explain multiple previous findings indicating that the NPD of

Rnq1 is involved in essentially all prion-related phenotypes associated with [*PIN+*]. First, while

constructs where Rnq1 PD is separated from Rnq1 NPD and attached to Sup35MC can form

prions, these prions have distinct properties from [*PIN+*] formed by Rnq1$_{FL}$ (Sondheimer and

Lindquist, 2000; Bardill and True, 2009). Second, *in vitro* Rnq1$_{FL}$ forms amyloid considerably

slower than Rnq1 fragments lacking the NPD (Vitrenko et al., 2007b; Kadnar et al., 2010), which

is consistent with inhibition of aggregation by the NPD even in the absence of its cellular

interactors. Third, point mutations and deletions in Rnq1 NPD affect the propagation of [*PIN+*]

(Vitrenko et al., 2007a, Shibata et al., 2009; Kurahashi et al., 2011a). Curiously, among 13

missense mutations uncovered by a screen for loss of [*PIN+*], 12 were in the NPD and two

mutations, V23A and A29T, were located very close to our T27P mutation, within the same

predicted helical stretch (Shibata et al., 2009; Kurahashi et al., 2011a). The authors interpreted

their results as moderate destabilization of [*PIN+*] by the mutations and the possibility of

transmission barrier was not tested. However, our results suggest that data in these earlier

papers are also consistent with the creation of transmission barriers between [*PIN+$_{WT}$*] and at

least some of the mutants. The Shibata et al. study also supports the likelihood that our genetic

screen was close to saturating and that mutations creating strong transmission barriers, like

T27P, are extremely rare while most other changes have much milder effect (between 30 and

70% of cells carried [*PIN+$_{MUTANT}$*] prions in their experiments corresponding to Round 1 analysis

here). Fourth, a site for interaction with the Sis1 chaperone essential for [*PIN+*]

propagation is located in the NPD (aa 91-97) and mutations in this site lead to [*PIN+*]

destabilization (Douglas et al., 2008; Shibata et al., 2009). Overexpression of Rnq1$_{FL}$, but not

NPD or PD alone, is toxic in [*PIN+*] but not [*pin-*] cells due to oxidative stress, and this toxicity is

rescued by overexpression of the Sis1 chaperone (Douglas et al., 2008; Bharthi et al., 2016).

To explain this, it was hypothesized that Sis1 directs Rnq1 into non-toxic [*PIN+*] amyloid

aggregates, preventing it from forming off-pathway SDS-soluble aggregates that cause oxidative

stress (Douglas et al., 2008). Finally, the NPD of Rnq1 also affects interactions of [*PIN*+] with

other prions. Overexpression of a fragment with a specific deletion in NPD, Rnq1$_{\Delta 1\text{-}100}$, leads to

loss of [*PSI*+], [*URE3*] and huntingtin polyQ aggregates in [*PIN*+] but not in [*pin*-] cells (Kurahashi

et al., 2008). Later, similar [*PIN*+]-dependent curing of [*PSI*+] was reported for multiple missense

alleles in the Rnq1 NPD, including L26S, S28P and A29T located close to T27P (Kurahashi et

al., 2009, 2011a, 2011b). One possible explanation based on our study is that these mutant

proteins switch to variant(s) of [*PIN*+] that are incompatible with [*PSI*+], like those reported earlier

(Bradley and Liebman, 2003; Mathur et al., 2009; Villali et al., 2020).

Curiously, several studies mention that changes in the NPD of Rnq1 led it to form non-

amyloid aggregates when tagged with YFP, but these aggregates are different from Rnq1$_{T27P}$

liquid-like droplets detected in our study. The SDS-soluble aggregates formed by Rnq1 with a

mutated Sis1 binding site are highly toxic even in [*pin*-] cells (Douglas et al., 2008), while

Rnq1$_{T27P}$-YFP droplets do not inhibit growth. Also ThT-negative aggregates formed by Rnq1$_{\Delta 1\text{-}}$

$_{100}$-YFP (Kurahashi et al., 2008) actively recruit Rnq1$_{FL\text{-}WT}$-CFP which does not join liquid-like

droplets formed by T27P. Thus, the Rnq1 NPD can probably control formation of not one, but

multiple types of non-amyloid aggregates.

While the NPD of Rnq1 is associated with multiple aggregation-related phenotypes, there

is no evidence for its activity not related to Rnq1 aggregation. The *RNQ1* gene is not essential,

and its deletion or overexpression does not lead to any significant growth defects in [*pin*-] cells

(www.yeastgenome.org and our data).

The key rationale for [*PSI*+], [*URE3*], and [*PIN*+] being diseases rather than functional

prions is based on their lower-than-expected frequency in wild yeast populations (Wickner et al.,

2011b; Kelly and Wickner, 2013). But [*PIN*+] is detected in wild and commercial yeast isolates

much more frequently than [*PSI+*] and [*URE3*], >10% *vs* <1% (Resende et al., 2003; Nakayashiki

et al., 2005; Halfman et al.,2012; Westergard and True, 2014a; Kelly et al., 2014). So, the

evidence for its detrimental effect is based on careful assessment of the frequency of outcross

matings expected to spread prions and on polymorphism of the PD-encoding part of *RNQ1* locus

expected to create transmission barriers (Kelly et al., 2012; Kelly and Wickner, 2013). These

calculations assume stable inheritance of [*PIN+*]. However, our study reveals [*PIN+*] variants with

heritably low mitotic stability. In addition to conformational switches induced by a transmission

barrier, unstable [*PIN+*]s also form *de novo* and during stress or otherwise changing

environmental conditions (Derkatch et al., 2000; Newnam et al., 2011; Westergard and True

2014b), and their instability may be heritable. Also, meiotic stability of [*PIN+*] is not 100%

(Bradley et al., 2002), and at least some of >40 different [*PIN+*] variants (Huang et al., 2013;

Westergard and True, 2014a) may be even more meiotically unstable, like [*URE3*] and some

[*PSI+*]s (Liebman and All-Robyn 1984; Wickner, 1994; Bradley et al., 2002). Finally, redundancy

of the Rnq1 PD makes it resistant to transmission barriers created by mutations: both our genetic

screen here and screens mentioned above uncovered NPD rather than PD mutations affecting

[*PIN+*]. Thus, [*PIN+*] may not have a strong enough detrimental effect to be considered a disease.

In thinking about possible functions for the Rnq1 protein it is instructive to recognize that

PDs of both Ure2 and Sup35 do contribute to cellular functions of these proteins (Hoshino et al.,

1999; Hosoda et al., 2003; Shewmaker et al., 2007; Edskes et al., 2018), but they are not

required for their well-known major jobs. For Sup35, the PD is not required for an essential role

in translation termination, but it is critical for at least two non-essential "moonlighting" jobs. One

involves interactions with the Pab1 protein (Hosoda et al., 2003), a major component of stress

granules known to be enriched in aggregation-prone proteins contributing to their assembly

(Fomicheva and Ross, 2021). The other "moonlighting job for Sup35 uncovered in our work (Li et

al., 2014) involves formation of a two-protein amyloid composed of Sup35 PD and the QN-rich

domain of Pub1. The [*PUB1 / SUP35*] complex associates with microtubules and promotes

integrity of the cytoskeleton. Thus, while [*PSI*⁺] may indeed be an egoistic byproduct of Sup35

aggregation properties, Sup35 function requiring the PD depends on the ability to form prion-like

aggregates. We hypothesize that the same is true for Rnq1. Its cellular function may involve

aggregation through the PD, and the NPD's role is in regulating such aggregation. One such

possible function is positive and negative interactions with other aggregating proteins in order to

induce or eliminate heterologous protein aggregates. Whether this role is carried out by [*PIN*⁺] or

through other single- or multiprotein aggregates encompassing Rnq1 has yet to be established.

1009

Materials and Methods

1010

1011   Most of the genetic and biochemical methods used for analysis of the [*PIN*⁺] prion in yeast,

1012   especially those used in our laboratory, are described in a review by Liebman et al. (2006).

1013

Strains and Plasmids

Bacterial Strains. Unless specifically mentioned, strain DH5α was used (F⁻ φ80*lacZΔM15*

Δ(*lacZYA-argF*)*U169 recA1 endA1 hsdR17*(r$_K$⁻, m$_K$⁺) *phoA supE44* λ⁻*thi*-1 *gyrA96 relA1*). Strain

KC8 was used to select for the yeast *HIS3* and *LEU2* markers when transformed with DNA

isolated from cells harboring more than one plasmid (*pyrF::Tn5, hsdR, leuB600, trpC9830,*

lacX74, strA, galUK, hisB436). Strain BL21-AI was used for recombinant protein purification (F⁻

*omp*T *hsd*S$_B$ (r$_B$⁻, m$_B$⁻) *gal dcm ara*B*::T7RNAP-tetA*).

1021

Yeast Strains. The original [*PIN*⁺][*psi*⁻] isolate of the yeast 74-D694 yeast strain (1Y1, aka

L1749; aka "high multi-dot" [*PIN*⁺]; Derkatch et al., 1997; Bradley et al., 2002) and its derivatives

were used in all experiments. The genotype of 74-D694 is *MAT**a** ade1-14* (UGA) *leu2-3,112*

his3-Δ200 trp1-289 (UAG) *ura3-52* (Chernoff et al., 1995). The 74-D694-based [*PIN+*][*psi*] *rnq1-Δ*

strain, YID108, and independently obtained isolates of this strain, YID146.2 and 146.3, were

constructed as described in Kadnar et al. (2010). In these strains the entire *RNQ1* ORF and

~100 bp *RNQ1* terminator sequence is seamlessly disrupted, and [*PIN+*] is maintained by the

CEN URA3 plasmid carrying the $RNQ1_{WT}$ ORF driven by the native *RNQ1* promoter and followed

by the native *RNQ1* terminator sequence ($RNQ1_{WT;}$ pID127). We have previously confirmed

(Kadnar et al., 2010) that in these strains Rnq1 is expressed at a level similar to that of

endogenous Rnq1 and [*PIN+*] is stably maintained (>99.5% stability). Unless specifically

mentioned, the YID108, YID146.2, and YID146.3 strains also carry the *CEN HIS3 GAL-*

SUP35NM-YFP plasmid (pID106; Kadnar et al., 2010) allowing the presence of [*PIN+*] to be

scored through the *de novo* [*PSI+*] induction. When indicated, plasmids pID349 (ΔB1C2), pID341

(ΔB1), pID342 (Δ2D), and pID344 (Δ3E) carrying $RNQ1_{WT}$ fragments were substituted for the full-

length $RNQ1_{WT}$ as [*PIN+*] maintainers in the YID146.3 strain (see Plasmids). We have previously

demonstrated that these fragments are expressed at their expected sizes and can stably

maintain the [*mini-PIN+*] prions and promote [*PSI+*] induction (Kadnar et al., 2010). Stable [*mini-*

PIN+] isolates with these maintainers, YID412, YID405, YID407, and YID409, respectively, are

from the Kadnar et al. (2010) study.

A [*pin-*][*psi*] *rnq1-Δ* strain (YID193) used for analysis of the *de novo* [*PIN+*] induction by

overexpression of $Rnq1_{T27P}$ and analysis of the *de novo* aggregation of $Rnq1_{T27P}$-YFP in the

rnq1-Δ background was obtained by selectively eliminating the *URA3* [*PIN+*] maintainer from

YID146.3 on FOA media while maintaining the *GAL-SUP3NM-YFP* [*PSI+*] inducing plasmid. The

[*pin-*][*psi*] isolate of 74-D694 (1G4; Derkatch et al., 1997) obtained by curing the [*PIN+*] prion on

media containing 5mM GuHCl was used in experiments testing the *de novo* aggregation of

$Rnq1_{T27P}$-YFP and $Rnq1_{WT}$-YFP in the presence of chromosomally encoded $Rnq1_{WT}$.

1049

Plasmids. All yeast plasmids used in this study include the same modules: promoters are *Eco*RI

– *Bam*HI fragments, ORFs are *Bam*HI – *Sac*II fragments, and terminators / fluorescent tags are

*Sac*II – *Sac*I fragments inserted into the poly-linkers of the pRS series plasmids (Sikorski and

Hieter, 1989). Site-directed mutagenesis was done using the QuickChange kit (Stratagene)

according to manufacturer's instructions and primers #360 and #361 (see Table S2).

The pRS416-based *CEN URA3* (pID127) and the pRS415-based *LEU2 CEN* (pID129)

plasmids carrying the $RNQ1_{WT}$ ORF driven by the native *RNQ1* promoter and followed by the

native *RNQ1* terminator sequence ($RNQ1_{WT}$) are described in Kadnar et al. (2010). To make

these plasmids, the *RNQ1* ORF was PCR amplified from 74-D694 chromosomal DNA. An earlier

study (Kadnar et al., 2010) confirmed that from these constructs $Rnq1_{WT}$ is expressed at a level

similar to that of the chromosomally encoded Rnq1 which allows for stable maintenance of [*PIN*+]

and does not lead to the induction of the *de novo* appearance of [*PIN*+] in [*pin*-] cells.

The $RNQ1_{T27P}$ carrying plasmid and other plasmids in the library used for the genetic

screen are pRS415-based, i.e. *LEU2 CEN*, and are identical to pID129 except for the

mutagenized *RNQ1* ORFs. The $RNQ1_{T27P}$ carrying pID335 plasmid was obtained by site-directed

mutagenesis of the pID219 plasmid.

Plasmids carrying the T27P mutation within *RNQ1* fragments lacking the sequences

encoding the indicated QN regions and adjacent hydrophobic helices within Rnq1 PD were

constructed by site-directed mutagenesis introducing T27P into the constructs previously

described in Kadnar et al. (2010): pID331 encoding $Rnq1_{\Delta B1C2-T27P}$ is based on the $Rnq1_{\Delta B1C2-WT}$

(pID370; ΔB1C2) plasmid lacking QN1, QN2 and helices B and C; pID332 encoding $Rnq1_{\Delta B1-T27P}$

– on the $Rnq1_{\Delta B1-WT}$ (pID369; ΔB1) plasmid lacking QN1 and helix B); pID333 encoding $Rnq1_{\Delta 2D-}$

$_{T27P}$ – on the $Rnq1_{\Delta 2D-WT}$ (pID358; Δ2D) plasmid lacking QN2 and helix D; and pID334 encoding

Rnq1$_{\Delta3E-T27P}$ – on the Rnq1$_{\Delta3E-WT}$ (pID387; Δ3E) plasmid lacking QN3 and helix E (Fig 1A). The

pID370, pID369, pID358, and pID387 plasmids were used as controls for experiments with the

corresponding abovementioned *LEU2* constructs carrying *RNQ1*$_{T27P}$. Except for deletions in the

RNQ1 PD these plasmids are identical to pID335 (RNQ$_{FL-T27P}$) and pID129 (RNQ$_{FL-WT}$),

respectively: pRS415-based, *LEU2 CEN,* original *RNQ1* promoter and terminator. Plasmids

pID349 (Rnq1$_{\Delta B1C2-WT}$), pID341 (Rnq1$_{\Delta B1-WT}$), pID342 (Rnq1$_{\Delta 2D -WT}$), and pID344 (Rnq1$_{\Delta3E-WT}$) are

identical to pID370, pID369, pID358, and pID387, respectively, except that, like pID127, they

carry the *URA3* marker. These plasmids were used as [*PIN*$^+$] maintainers for experiments testing

transmission barriers to Rnq1$_{FL-T27P}$ or Rnq1 fragments carrying Rnq1$_{T27P}$.

To overexpress Rnq1$_{WT}$ and Rnq1$_{T27P}$, respectively, the pID336 and pID337 plasmids

were constructed. Both plasmids are pRS416-based, *URA3, CEN* and the full-length ORFs are

controlled by the *CUP1* promoter and followed by the original *RNQ1* terminator.

In the pRS416-based *CEN LEU2 CUP-RNQ$_{WT}$-CFP* plasmid (pID121; Kadnar et al.,

2010), a fusion of complete *RNQ1$_{WT}$* ORF to *CFP*, is driven by the inducible *CUP1* promoter. The

pRS415-based *URA3*-marked pID403 and pID404 plasmids are similar except they carry,

respectively, *CUP-RNQ$_{WT}$-YFP* and *CUP-RNQ$_{T27P}$-YFP*.

The pRS413-based *CEN HIS3 GAL-SUP35NM-YFP* plasmid (pID106, Kadnar et al.,

2010) carries *SUP35NM-YFP* encoding the PD of the [*PSI*$^+$]-forming Sup35 protein fused to the

YFP reporter and placed under control of the inducible *GAL1* promoter.

Bacterial expression plasmids for Rnq1$_{WT}$ and Rnq1$_{T27P}$ are pJC45-based and express N-

terminally 10xHIS tagged proteins. The ΔB1C2D3$_{WT}$ plasmid (pID257; aka QN4 only) and full-

length RNQ1$_{WT}$ (pID249/250) constructs are described in Kadnar et al. (2010) and Vitrenko et al.

(2007b), respectively. The ΔB1C2D3$_{WT}$ construct expresses the Rnq1NPD through the QG

repeat region (aa1-172) fused to the helical region E and QN4 in the PD (aa 320-405); the fusion

is seamless, *i.e* no additional amino acids are introduced in between. The ΔB1C2D3$_{T27P}$ and

RNQ1$_{T27P}$ differ only by the T27P mutation that was introduced by site-directed mutagenesis into

ΔB1C2D3$_{WT}$ and RNQ1$_{WT}$, respectively.

1100

Yeast Media and Cultivation

Standard yeast media and cultivation procedures were used (Sherman et al., 1987; Rose et al.,

1990; Amberg et al., 2005). Unless specifically mentioned yeast transformants were grown at

30^{0}C on solid synthetic glucose media selective for plasmid maintenance (e.g. SD-UraLeuHis). In

addition to synthetic glucose SD-Ade media, synthetic ethanol SEt-Ade media supplemented

with 2% ethanol as the only carbon source was used to score for [*PSI+*] (see below).

Untransformed strains and cultures for protein isolation were grown on complete glucose media,

YPD. The *GAL* promoter was induced on solid synthetic media selective for plasmid

maintenance and supplemented with 2% galactose as a single carbon source (e.g. SGal-

LeuHis). The *CUP* promoter was induced by supplementing SD media selective for plasmid

maintenance with the indicated concentration of CuSO$_4$ (5, 20 or 50µM; see Results and Figure

Legends for details of specific experiments). To selectively eliminate plasmids with the *URA3*

marker, yeast were passaged twice on solid SD supplemented with 5-fluoroorotic acid (5-FOA;

Boeke et al., 1984) selective for other plasmids. Solid YPD supplemented with 5mM guanidine

hydrochloride (GuHCl; Tuite et al., 1981) was used to cure the [*PSI+*] and [*PIN+*] prions.

YPGlycerol media supplemented with 2% glycerol as a single carbon source was used to detect

Pet$^-$ colonies that may be confused with [*PSI+*] by color.

1118

RNQ1 **Mutagenesis and Genetic Screen**

To mutagenize *RNQ1*, the complete 1.2 kb *RNQ1* ORF was PCR amplified from pID129 using

Taq polymerase (New England Biolabs), primers # 7 and # 8 (see Table S2), and manufacturer's

recommended reaction conditions, which was expected to yield ~1 mutation per 1,000 bps after

32 cycles. To eliminate mutagenic bias from any one PCR reaction, seven independent PCR

reactions were pooled. PCR products were digested with *Bam*HI and *Sac*II and re-introduced

into the *Bam*HI- and *Sac*II- digested pID129 vector between the *RNQ1* promoter and *RNQ1*

terminator sequences with a stop codon right after the *Sac*II site. Ligation reactions were

transformed into *E. coli*. Approximately 30,600 transformant colonies were washed off with ~1

ml LB from three transformant selection plates (banks pID142 – pID144) and ~200μl of

suspensions were added to 50ml LB+Amp (30 mg/l) and grown for five hours at 37^0C prior to

plasmid DNA isolation. Plasmid DNA was introduced into the YID108 yeast cells. Sequencing the

RNQ1 ORF from two randomly selected *E. coli* transformants with primers #119 and #8

confirmed the presence of 1 mutation per the Rnq1 PD. At this estimated rate of mutagenesis,

screening ~12,000 plasmids was expected to result in >95% chance of scoring all three possible

nucleotide changes at each base.

The genetic screen is described in Results and is illustrated in Fig 1B. Approximately 40

patches of yeast transformants were analyzed per plate. Each plate had negative (EV) and

positive (pID129) control transformants. Passages on all indicated media were by replica plating.

Two passages on SD-LeuHis not selective for the <u>*URA3 RNQ1$_{WT}$* plasmid preceded replica</u>

<u>plating to FOA to accumulate cells that lost this plasmid.</u> Two passages on 5-FOA were sufficient

for the selective elimination of all cells carrying the *URA3*-marked *RNQ1$_{WT}$* plasmid (confirmed

by replica plating to SD media lacking uracil). Induction of [*PSI⁺*] was scored, after one and two

passages on SGal-LeuHis, on SD-Ade at 30^0C and 20^0C and on SEt-Ade at 30^0C after up to 25

days of incubation, which is sufficient for the detection of even the weakest [*PSI⁺*] variants.

No [*PSI+*] induction was detected in ~500 transformants with mutated plasmids.

Immediately excluded were the candidates that: (i) were Pet⁻ (detected as lack of growth on

YPGlycerol to which all patches were also replica plated); (ii) carried *RNQ1_{MUTANT}* plasmids

lacking the full-length 1.2 kb *RNQ1* ORF insert (detected by PCR using primers #7 and #8 and

total yeast DNA from after-shuffle cultures). Also, to exclude plasmids that carried frameshift

mutations (usually -1) within the forward primer used to amplify the *RNQ1* ORF or stop codons in

the very beginning of the ORF, a blue-white ligation-based assay was developed. The first ~100

bps of the *RNQ1* ORF were PCR amplified using forward primers #127 or #124 and reverse

primer #129 and cloned into the pUC18 plasmid as *Bcl*I – *Hin*dIII, *Bam*HI – *Hin*dIII, or *Bgl*II –

*Hin*dIII fragments to create the in-frame, -1, and +1 fusions with the *lacZ* gene. The ligation

mixes were transformed into the DH5α *E. coli* strain to test for the complementation of the

φ80*lacZ*ΔM15 allele. White or light blue color of most bacterial transformants on the LB+Amp

media supplemented with 40μg/ml XGal was indicative of the disruption of LacZ synthesis in this

particular frame. The test was validated by sequencing ~30 plasmids with presumptive early

frameshift and nonsense mutations and >100 plasmids with no mutations in the beginning of the

RNQ1 ORF predicted by this assay.

For the remaining candidates, the *RNQ1_{MUTANT}* plasmids were isolated for further

analysis. To selectively isolate only the *LEU2*-marked *RNQ1_{MUTANT}* plasmids and not the *HIS3*-

marked *GAL-SUP35NM-YFP* plasmids, the *E. coli* KC8 strain (see *Strains* above) was

transformed with total yeast DNA extracts, and Leu⁺ transformants were selected on M9-based

media lacking leucine. Digestion of plasmids with restriction endonucleases was used to exclude

RNQ1_{MUTANT} plasmids with re-arrangements outside of the *RNQ1* ORF. Then, for each

candidate, DNA from at least two independent *E. coli* transformants was re-transformed into the

original yeast strain and re-tested according to the same scheme as in the original screen (see

Fig 1B); in total, at least 12 yeast transformants were tested for each candidate. The re-

transformation test provided proof that a mutation in the $RNQ1_{MUTANT}$ plasmid was the cause of

the Pin⁻ phenotype in the after-shuffle cultures. Alternatively, the Pin⁻ phenotype could be either

due to spontaneous loss of the [PIN^+] prion (0.2 – 0.5% in control experiments where the

$RNQ1_{WT}$ plasmid was shuffled-in) or due to chromosomal mutations e.g. either eliminating [PIN^+]

or inhibiting [PSI^+] induction. Primers #7, #8, #17, #18, #19, and #30 were used to sequence the

$RNQ1$ ORF and adjacent promoter and terminator regions (see Table S2).

1175

Scoring for the presence of the [PIN^+] prion and Pin phenotype

Induction of the *de novo* appearance of the [PSI^+] prion upon overexpression of the PD of the

Sup35 protein, Sup35NM, was used to score for the presence of [PIN^+] (see Liebman et al.,

2006). The assay is based on our finding that [PIN^+] is required for the efficient induction of [PSI^+]

(Derkatch et al., 1997). In [PIN^+] cells expressing Sup35NM-YFP at an ~5-fold increased level

compared to the expression of endogenous Sup35 for ~10 cell generations, [PSI^+] forms in ~10%

cells, whereas appearance of [PSI^+] in [pin^-] cultures is extremely rare, $<10^{-4}$. Transformants

carrying the *CEN HIS3 GAL-SUP35NM-YFP* [PSI^+] inducing construct and other indicated

plasmids were grown on SD media selective for all the plasmids and then replica plated onto the

inducing SGal media, also selective for all the plasmids. Growth on SGal was for 2 days (5-7

generations), a second passage on SGal was done to increase the number of generations and

thus the chance of [PSI^+] induction. As a negative control, the same cultures were passaged on

non-inducing SD media and then analyzed in parallel with induced cultures. Three methods were

used to score for the appearance of [PSI^+].

(1) For qualitative/semi-quantitative analysis, yeast were replica plated from SGal (or

control SD) media onto SD media lacking adenine and not selective for plasmids. An Ade⁺

phenotype after SGal and not after SD is indicative of [*PSI*+] induction. Growth of [*PSI*+] cells on

adenineless media is due to suppression of a premature UGA stop codon in the *ADE1* gene

present in our strains (*ade1-14*). The nonsense suppression is due to the sequestration of the

chromosomally encoded full-length Sup35 protein, a translation termination factor, into the [*PSI*+]

prion aggregates initiated by Sup35NM-YFP overexpression. For all such experiments, both SD-

Ade and SEt-Ade media were used. Cultures on SD-Ade were incubated at 30^{0}C and 20^{0}C;

cultures on SEt-Ade were incubated at 30^{0}C. Growth was recorded multiple times between 3 and

25 days of incubation. This approach allows detection of [*PSI*+] variants of different strengths

(some very weak [*PSI*+] variants grow only on SEt-Ade or at lower temperatures, and only after a

very long incubation).

(2) For quantitative analysis, yeast were colony purified on YPD from galactose (or

control glucose) media. On YPD, [*PSI*+] cells form white or pink colonies whereas [*psi*-] cells are

red due to the accumulation of a red pigment from the adenine precursor accumulating in the

ade1-14 cells. The appearance of white and pink Ade+ colonies (among red Ade- colonies)

following Sup35NM-YFP overexpression is indicative of [*PSI*+] induction.

However, nonsense suppression and white color in (1) and (2) can also be caused by

genomic mutations in *SUP35* and other genes. Such genomic mutations are rare compared to

the induced [*PSI*+] *de novo* formation in [*PIN*+] cultures. Yet, and especially in cultures where only

a few Ade+ or white / pink colonies were detected, such colonies were confirmed to be [*PSI*+]. For

the GuHCl test, white / pink colonies were patched onto YPD, then passaged three times on

YPD+5mM GuHCl and then returned to YPD. Because GuHCl cures [*PSI*+] (Tuite et al., 1981),

[*PSI*+] patches turn red on GuHCl media and remain red when returned to YPD. Chromosomal

mutants may turn red on GuHCl (Bradley et al., 2003) but return to their white or pink color once

back on YPD. Alternatively, for colonies that retained the *HIS3 GAL-SUP35NM-YFP* plasmid, the

presence of [PSI^+] was confirmed by fluorescence microscopy: after an overnight growth on

SGal-His media, dot-shaped foci are detected in most cells in [PSI^+] cultures.

(3) To directly visualize newly forming [PSI^+]s, yeast cells from SGal media were

analyzed by fluorescent microscopy utilizing the YFP tag attached to Sup35NM in the [PSI^+]

inducing construct. Newly forming [PSI^+]s can be seen as ring-, line- or dot-shaped foci; cells

with already established [PSI^+] have dot foci; in [psi^-] cells fluorescence is evenly distributed

(Patino et al., 1996; Zhou et al., 2001).

1223

Assessing the [PSI^+] prion variants

To assess the arrays of the [PSI^+] prion variants induced in cells expressing Rnq1$_{T27P}$ *vs* Rnq1$_{WT}$,

the Ade$^+$ cells from patches replica plated from SGal to SD-Ade [PSI^+] selective media and

grown for 25 days at 20^0C were thoroughly scraped and colony purified on YPD and individual

colonies were randomly picked up from YPD for further analysis before color had developed. At

least 12 transformants were analyzed for either Rnq1$_{T27P}$ or Rnq1$_{WT}$. The same approach was

used to obtain [PSI^+]s appearing in [pin^-] cultures carrying EV. Alternatively, yeast were colony

purified on YPD from the second pass on SGal-LeuHis media, where [PSI^+]s were induced.

Three assays were used to assess [PSI^+] strength. **(1)** Color on YPD (the lighter the color, the

stronger the [PSI^+] variant). **(2)** Growth on adenineless media: SD-Ade at 30^0C and at 20^0C, and

SEt-Ade at 30^0C (the more robust the Ade$^+$ phenotype, the stronger the [PSI^+]). **(3)** Efficiency of

curing by GuHCl: after three passages on YPD+GuHCl yeast were replica plated onto the same

set of adenineless media as in (**2**). While weak [PSI^+]s were usually cured completely, occasional

Ade$^+$ colonies were detected for stronger [PSI^+]s.

1238

1239   **Analysis of the *de novo* induction of [$PIN^+{}_{T27P}$]**

1240   Both variants of this experiment are succinctly explained in Results related to Figs 3A and B. See

1241   Supplementary Materials and Methods for plasmid numbers and details on controls.

1242

Plasmid stability analysis

To compare stability of the Rnq1 expressing plasmids in yeast transformants carrying the *CEN*

URA3 RNQ1$_{WT}$ and *CEN LEU2 RNQ1$_{T27P}$* plasmids *vs* transformants carrying two *RNQ1$_{WT}$*

plasmids (*CEN URA3* and *CEN LEU2*) or a *RNQ1$_{WT}$* (*CEN URA3*) plasmid and the EV (*CEN*

LEU2), fresh transformants were patched onto SD-UraLeu media selective for both plasmids,

then passaged twice on media selective for only one of the plasmids (SD-Ura or SD-Leu), then

colony purified on the same media and replica plated to both that media and to the reciprocal

media. Plasmid stability was determined as the fraction of colonies grown on the reciprocal

media where only colonies formed by cells retaining the plasmid despite of the lack of selection

were expected to grow.

1253

Western Blot Analysis of Rnq1$_{T27P}$ expression in yeast

Yeast cell lysates were prepared essentially as described in Liebman et al. (2006) and Kadnar et

al. (2010). See Supplementary Materials and Methods for a detailed protocol. Either 30 or 60μg

of total protein (concentration determined with BCA™ Protein Assay Kit, Pierce) were separated

on SDS-PAGE gels. Rnq1 was detected by Western blot with polyclonal antibodies raised in

rabbits against the first 185 aa of Rnq1 (type 1A, a kind gift from E. Craig, University of

Wisconsin-Madison). Sup35 was detected with a monoclonal antibody against the Sup35C

raised in mouse (BE4). HRP-conjugated anti-rabbit and anti-mouse secondary antibodies and

the Super Signal Western PICO ECL kit (Pierce) were used for detection.

1263

1264 **Recombinant protein purification from _E. coli_**

1265 Recombinant protein purification was essentially as described in Kadnar et al. (2010). See

1266 Supplementary Materials and Methods for a detailed protocol. Protein concentration was

1267 determined by the BCA™ Protein Assay Kit (Pierce). Purity of recombinant proteins was

1268 estimated by SDS-PAGE as >95% (see Fig 4A).

1269

Analysis of _in vitro_ fiber formation

Analysis of _in vitro_ fiber formation by $Rnq1_{WT}$ and $Rnq1_{T27P}$ was performed essentially as in

Kadnar et al. (2010). Specifically, 200 μl reactions were assembled in the fiber formation buffer

(1M urea, final concentration; 100mM NaH_2PO_4, pH 7.4; 300mM NaCl) in the presence of 5 μM

thioflavin T (ThT). Fiber formation was monitored by ThT fluorescence (LeVine, 1999) in a

Molecular Devices SpectraMax M-5 plate reader at RT (λ_{ex}=450 nm; λ_{em}=483 nm). Readings

were taken every 15 min, samples were shaken for 5 sec prior to each reading.

1277

Transmission Electron Microscopy (TEM)

TEM was performed at the Image Core Facility at the NYU School of Medicine. Following

incubation to allow for amyloid fiber formation, protein / fiber suspensions were diluted 2.5-fold in

water. Then 4 μl of diluted suspensions were applied to carbon-coated 400 mesh Cu/Rh grids

(Ted Pella Inc.). Grids were washed 3 times with water to get rid of urea present in the fiber

formation buffer. Then grids were negatively stained at 25^0C with 1% uranyl acetate (applied

twice briefly and then for 5 min). Images were captured using Philips CM12 TEM supplied with a

Gatan 1k61k digital camera and processed using the Gatan Digital Micrograph software.

1286

Fluorescence Microscopy, ThT staining, and 1,6 Hexanediol treatment

Cells were observed using a Nikon Eclipse E600 microscope with a 60x / 1.4 NA oil immersion

objective. YFP and CFP filters were used to observe YFP- and CFP-tagged constructs,

respectively. Fluorescent and brightfield images of the same field were captured with a Zeiss

AxioCam MRm digital camera and processed with AxioVision AxioVs40 V 4.8.2.0 software. In all

experiments, representative groups of cells were scored, including cells of different sizes, with

and without buds and with different levels of construct expression. For Rnq1 aggregation

analysis, cells from at least three transformants were counted, at least 250 cells from each. For

analysis of Sup35NM-YFP aggregation shown in Table 1 and Fig 2C, at least 11 vision fields

were scored for each construct, ~500 cells total per construct, while qualitative assessment of

aggregation was done on a much larger scale. ImageJ software was used for cell counting.

Photoshop was used for merging and adjusting images (only brightness and contrast were

adjusted, no other processing). For all images shown, differences in cell size, budding and

fluorescence intensity are within population variability range.

Staining with thioflavin T (ThT) was done essentially as described in Douglas et al.

(2008). See Supplementary Materials and Methods for a detailed protocol. The CFP filter was

used for ThT fluorescence analysis.

To dissolve liquid-like droplets with 1,6 Hexanediol (Hex), yeast transformants were

grown overnight in liquid SD media selective for all plasmids. Then OD_{595} was adjusted to 1.0,

$CuSO_4$ was added to 5μM, and cultures were grown for two more hrs. Cells were harvested,

resuspended in 100 – 500μl of the same SD media with 10% Hex (Fisher) and 0.2μg/μl digitonin

(Sigma; 200 mg/ml stock in water) and incubated on a shaker for 10 min at 30^0C. Then cells

were centrifuged, resuspended in the same SD, and imaged using the filter matching the Rnq1

fluorescent tag.

1311

Statistics

Standard methods of statistical analysis were used. Averages and Standard Deviations

(STDEV.S) were calculated in Excell (16.105.2 and earlier versions). Unless specifically

mentioned, SD is included. Standard Error of the Mean (SEM) was calculated by dividing

STDEV.S by square root of the number of datapoints. Excell and GraphPad Prism

(10.6.1(892))were used for graphs. Statistics Kingdom (statisticskingdom.com) was used for T-

tests (two sample Welch's T-test; unknown / unequal SD). Unless specifically mentioned, the

cutoff for the significance of variation was $p<0.01$.

Acknowledgements

We thank Elizabeth Craig for providing the anti-Rnq1 antibody. We are grateful to Gulnara

Articov for help with *in-vitro* experiments and to Alice Liang and Eric Roth at the NYU School of

Medicine Image Core Facility for performing TEM. We also thank Sei-Kyoung Park for helpful

discussions of the data.

Funding statement

This work was supported by grants from NIH - R01 GM070934-06 (ILD) and MIRA

1R35GM136229-05 (SWL). National Science Foundation - 0518482 (ILD), and US Army – ARO

W911NF-23-1-0122 (ILD). The funders had no role in study design, data collection and

analysis, decision to publish, or preparation of the manuscript.

Conflicts of Interest Statement: The authors declare no conflicts of interest.

Data availability

The authors confirm that all data underlying the findings are fully available without

restriction. All relevant data are within the paper and its Supporting Information files.

1336 **Author Contributions**

1337 Conceived and designed the experiments: ILD. Performed the experiments: SP, DMM, MLK, MA,

1338 APF, ILD. Analyzed the data: SP, MLK, SWL, ILD. Contributed reagents/materials/analysis tools:

1339 ILD, SWL. Wrote the manuscript: ILD, SWL.

1340

References

Afanasieva EG, Kushnirov VV, Tuite MF, Ter-Avanesyan MD. Molecular basis for transmission barrier and
interference between closely related prion proteins in yeast. J Biol Chem. 2011 May 6;286(18):15773-80.
doi: 10.1074/jbc.M110.183889. Epub 2011 Mar 15. PMID: 21454674; PMCID: PMC3091186.

Amberg, D.C., Burke, D., Strathern, J.N., and Cold Spring Harbor Laboratory. (2005). Methods in yeast
genetics: a Cold Spring Harbor Laboratory course manual, 2005 edn (Cold Spring Harbor, N.Y., Cold
Spring Harbor Laboratory Press).

Ayers JI, Paras NA, Prusiner SB. Expanding spectrum of prion diseases. Emerg Top Life Sci. 2020 Sep
8;4(2):155-167. Doi: 10.1042/ETLS20200037. PMID: 32803268.

Barbitoff YA, Matveenko AG, Zhouravleva GA. Differential Interactions of Molecular Chaperones and
Yeast Prions. J Fungi (Basel). 2022 Jan 27;8(2):122. Doi: 10.3390/jof8020122. PMID: 35205876; PMCID:
PMC8877571.

Bardill JP, True HL. Heterologous prion interactions are altered by mutations in the prion protein Rnq1p. J
Mol Biol. 2009 May 8;388(3):583-96. doi: 10.1016/j.jmb.2009.03.036. Epub 2009 Mar 24. PMID:
19324054; PMCID: PMC2706087.

Baskakov IV. The many shades of prion strain adaptation. Prion. 2014 Mar-Apr;8(2):169–72. doi:
10.4161/pri.27836. Epub 2014 Feb 11. PMID: 24518385; PMCID: PMC4189885.

Bateman DA, Wickner RB. [PSI+] Prion transmission barriers protect Saccharomyces cerevisiae from
infection: intraspecies 'species barriers'. Genetics. 2012 Feb;190(2):569-79. doi:
10.1534/genetics.111.136655. Epub 2011 Nov 17. PMID: 22095075; PMCID: PMC3276615.

Bateman DA, Wickner RB. The [PSI+] prion exists as a dynamic cloud of variants. PLoS Genet.
2013;9(1):e1003257. doi: 10.1371/journal.pgen.1003257. Epub 2013 Jan 31. Erratum in: PLoS Genet.
2017 Feb 3;13(2):e1006591. doi: 10.1371/journal.pgen.1006591. PMID: 23382698; PMCID: PMC3561065.

Béringue V, Vilotte JL, Laude H. Prion agent diversity and species barrier. Vet Res. 2008 Jul-Aug;39(4):47.
Doi: 10.1051/vetres:2008024. Epub 2008 Jun 3. PMID: 18519020.

Bessen RA, Marsh RF. Identification of two biologically distinct strains of transmissible mink
encephalopathy in hamsters. J Gen Virol. 1992 Feb;73 (Pt 2):329-34. Doi: 10.1099/0022-1317-73-2-329.
PMID: 1531675.

Bessen RA, Kocisko DA, Raymond GJ, Nandan S, Lansbury PT, Caughey B. Non-genetic propagation of

strain-specific properties of scrapie prion protein. Nature. 1995 Jun 22;375(6533):698-700. Doi:
10.1038/375698a0. PMID: 7791905.

Bharathi V, Girdhar A, Prasad A, Verma M, Taneja V, Patel BK. Use of ade1 and ade2 mutations for
development of a versatile red/white colour assay of amyloid-induced oxidative stress in saccharomyces
cerevisiae. Yeast. 2016 Dec;33(12):607-620. Doi: 10.1002/yea.3209. Epub 2016 Oct 13. PMID: 27654890.

Boeke JD, LaCroute F, Fink GR. A positive selection for mutants lacking orotidine-5'-phosphate
decarboxylase activity in yeast: 5-fluoro-orotic acid resistance. Mol Gen Genet. 1984;197(2):345-6. Doi:
10.1007/BF00330984. PMID: 6394957.

Borkhsenius AS, Sasnauskas K, Gedvilaĭte A, Inge-Vechtomov SG. Khimernye priony drozhzheĭ s
nestabil'nym nasledovaniem [Chimeric yeast prions with unstable inheritance]. Genetika. 2002
Mar;38(3):300-5. Russian. PMID: 11963556.

Bradley ME, Edskes HK, Hong JY, Wickner RB, Liebman SW. Interactions among prions and prion
"strains" in yeast. Proc Natl Acad Sci U S A. 2002 Dec 10;99 Suppl 4(Suppl 4):16392-9. Doi:
10.1073/pnas.152330699. Epub 2002 Jul 30. PMID: 12149514; PMCID: PMC139899.

1384

Bradley ME, Liebman SW. Destabilizing interactions among [PSI(+)] and [PIN(+)] yeast prion variants.
Genetics. 2003 Dec;165(4):1675-85. Doi: 10.1093/genetics/165.4.1675. Erratum in: Genetics. 2011
Aug;188(4):1023. PMID: 14704158; PMCID: PMC1462903.

Bradley ME, Bagriantsev S, Vishveshwara N, Liebman SW. Guanidine reduces stop codon read-through
caused by missense mutations in SUP35 or SUP45. Yeast. 2003 May;20(7):625-32. Doi: 10.1002/yea.985.
PMID: 12734800.

Chang HY, Lin JY, Lee HC, Wang HL, King CY. Strain-specific sequences required for yeast [PSI+] prion
propagation. Proc Natl Acad Sci U S A. 2008 Sep 9;105(36):13345-50. doi: 10.1073/pnas.0802215105.
Epub 2008 Aug 29. PMID: 18757753; PMCID: PMC2533192.

Chen B, Newnam GP, Chernoff YO. Prion species barrier between the closely related yeast proteins is
detected despite coaggregation. Proc Natl Acad Sci U S A. 2007 Feb 20;104(8):2791-6. doi:
10.1073/pnas.0611158104. Epub 2007 Feb 12. PMID: 17296932; PMCID: PMC1815260.

Chen B, Bruce KL, Newnam GP, Gyoneva S, Romanyuk AV, Chernoff YO. Genetic and epigenetic control
of the efficiency and fidelity of cross-species prion transmission. Mol Microbiol. 2010 Jun;76(6):1483-99.
doi: 10.1111/j.1365-2958.2010.07177.x. Epub 2010 Apr 23. PMID: 20444092; PMCID: PMC3025758.

Chernoff YO, Lindquist SL, Ono B, Inge-Vechtomov SG, Liebman SW. Role of the chaperone protein
Hsp104 in propagation of the yeast prion-like factor [psi+]. Science. 1995 May 12;268(5212):880-4. doi:
10.1126/science.7754373. PMID: 7754373.

Chuang E, Hori AM, Hesketh CD, Shorter J. Amyloid assembly and disassembly. J Cell Sci. 2018 Apr
13;131(8):jcs189928. doi: 10.1242/jcs.189928. PMID: 29654159; PMCID: PMC5963839.

Collinge J, Clarke AR. A general model of prion strains and their pathogenicity. Science. 2007 Nov
9;318(5852):930-6. doi: 10.1126/science.1138718. PMID: 17991853.

Collinge J. Cell biology. The risk of prion zoonoses. Science. 2012 Jan 27;335(6067):411-3. doi:
10.1126/science.1218167. PMID: 22282797.

DePace AH, Santoso A, Hillner P, Weissman JS. A critical role for amino-terminal glutamine/asparagine
repeats in the formation and propagation of a yeast prion. Cell. 1998 Jun 26;93(7):1241-52. doi:
10.1016/s0092-8674(00)81467-1. PMID: 9657156.

Derkatch IL, Chernoff YO, Kushnirov VV, Inge-Vechtomov SG, Liebman SW. Genesis and variability of
[PSI] prion factors in Saccharomyces cerevisiae. Genetics. 1996 Dec;144(4):1375-86. doi:
10.1093/genetics/144.4.1375. PMID: 8978027; PMCID: PMC1207691.

Derkatch IL, Bradley ME, Zhou P, Chernoff YO, Liebman SW. Genetic and environmental factors affecting
the de novo appearance of the [PSI+] prion in Saccharomyces cerevisiae. Genetics. 1997 Oct;147(2):507-
19. doi: 10.1093/genetics/147.2.507. PMID: 9335589; PMCID: PMC1208174.

Derkatch IL, Bradley ME, Liebman SW. Overexpression of the SUP45 gene encoding a Sup35p-binding
protein inhibits the induction of the de novo appearance of the [PSI+] prion. Proc Natl Acad Sci U S A.
1998 Mar 3;95(5):2400-5. doi: 10.1073/pnas.95.5.2400. PMID: 9482897; PMCID: PMC19355.

Derkatch IL, Bradley ME, Zhou P, Liebman SW. The PNM2 mutation in the prion protein domain of SUP35
has distinct effects on different variants of the [PSI+] prion in yeast. Curr Genet. 1999 Mar;35(2):59-67. doi:
10.1007/s002940050433. PMID: 10079323.

Derkatch IL, Bradley ME, Masse SV, Zadorsky SP, Polozkov GV, Inge-Vechtomov SG, Liebman SW.
Dependence and independence of [PSI+] and [PIN+]: a two-prion system in yeast? EMBO J. 2000 May
2;19(9):1942-52. doi: 10.1093/emboj/19.9.1942. PMID: 10790361; PMCID: PMC305693.

Derkatch IL, Bradley ME, Hong JY, Liebman SW. Prions affect the appearance of other prions: the story of
[PIN(+)]. Cell. 2001 Jul 27;106(2):171-82. doi: 10.1016/s0092-8674(01)00427-5. PMID: 11511345.

Derkatch IL, Liebman SW. Prion-prion interactions. Prion. 2007 Jul-Sep;1(3):161-9. doi:
10.4161/pri.1.3.4837. Epub 2007 Jul 6. PMID: 19164893; PMCID: PMC2634589.

DiSalvo S, Derdowski A, Pezza JA, Serio TR. Dominant prion mutants induce curing through pathways
that promote chaperone-mediated disaggregation. Nat Struct Mol Biol. 2011 Apr;18(4):486-92. doi:
10.1038/nsmb.2031. Epub 2011 Mar 20. PMID: 21423195; PMCID: PMC3082495.

Dobson CM. The Amyloid Phenomenon and Its Links with Human Disease. Cold Spring Harb Perspect
Biol. 2017 Jun 1;9(6):a023648. doi: 10.1101/cshperspect.a023648. PMID: 28062560; PMCID:
PMC5453392.

Douglas PM, Treusch S, Ren HY, Halfmann R, Duennwald ML, Lindquist S, Cyr DM. Chaperone-
dependent amyloid assembly protects cells from prion toxicity. Proc Natl Acad Sci U S A. 2008 May
20;105(20):7206-11. doi: 10.1073/pnas.0802593105. Epub 2008 May 14. PMID: 18480252; PMCID:
PMC2438228.

Du Z, Park KW, Yu H, Fan Q, Li L. Newly identified prion linked to the chromatin-remodeling factor Swi1 in
Saccharomyces cerevisiae. Nat Genet. 2008 Apr;40(4):460-5. doi: 10.1038/ng.112. Epub 2008 Mar 23.
PMID: 18362884; PMCID: PMC2633598.

Du Z, Goncharoff DK, Cheng X, Li L. Analysis of [SWI+] formation and propagation events. Mol Microbiol.

2017 Apr;104(1):105-124. doi: 10.1111/mmi.13616. Epub 2017 Jan 26. PMID: 28035761; PMCID: PMC5364053.

Edskes HK, Gray VT, Wickner RB. The [URE3] prion is an aggregated form of Ure2p that can be cured by overexpression of Ure2p fragments. Proc Natl Acad Sci U S A. 1999 Feb 16;96(4):1498-503. doi: 10.1073/pnas.96.4.1498. PMID: 9990052; PMCID: PMC15494.

Edskes HK, McCann LM, Hebert AM, Wickner RB. Prion variants and species barriers among Saccharomyces Ure2 proteins. Genetics. 2009 Mar;181(3):1159-67. doi: 10.1534/genetics.108.099929. Epub 2009 Jan 5. PMID: 19124570; PMCID: PMC2651050.

Edskes HK, Mukhamedova M, Edskes BK, Wickner RB. *Hermes* Transposon Mutagenesis Shows [URE3] Prion Pathology Prevented by a Ubiquitin-Targeting Protein: Evidence for Carbon/Nitrogen Assimilation Cross Talk and a Second Function for Ure2p in *Saccharomyces cerevisiae*. Genetics. 2018 Jul;209(3):789-800. doi: 10.1534/genetics.118.300981. Epub 2018 May 16. PMID: 29769283; PMCID: PMC6028256.

Fomicheva A, Ross ED. From Prions to Stress Granules: Defining the Compositional Features of Prion-Like Domains That Promote Different Types of Assemblies. Int J Mol Sci. 2021 Jan 27;22(3):1251. doi: 10.3390/ijms22031251. PMID: 33513942; PMCID: PMC7865556.

Fuentealba RA, Udan M, Bell S, Wegorzewska I, Shao J, Diamond MI, Weihl CC, Baloh RH. Interaction with polyglutamine aggregates reveals a Q/N-rich domain in TDP-43. J Biol Chem. 2010 Aug 20;285(34):26304-14. doi: 10.1074/jbc.M110.125039. Epub 2010 Jun 16. PMID: 20554523; PMCID: PMC2924052.

Galliamov AA, Malukhina AD, Kushnirov VV. Mapping of Prion Structures in the Yeast Rnq1. Int J Mol Sci. 2024 Mar 17;25(6):3397. doi: 10.3390/ijms25063397. PMID: 38542372; PMCID: PMC10970677.

Galliamov AA, Urakov VN, Dergalev AA, Kushnirov VV. On the Significance of the Terminal Location of Prion-Forming Regions of Yeast Proteins. Int J Mol Sci. 2025 Feb 14;26(4):1637. doi: 10.3390/ijms26041637. PMID: 40004101; PMCID: PMC11855515.

Halfmann R, Jarosz DF, Jones SK, Chang A, Lancaster AK, Lindquist S. Prions are a common mechanism for phenotypic inheritance in wild yeasts. Nature. 2012 Feb 15;482(7385):363-8. doi: 10.1038/nature10875. PMID: 22337056; PMCID: PMC3319070.

Hill AF, Desbruslais M, Joiner S, Sidle KC, Gowland I, Collinge J, Doey LJ, Lantos P. The same prion strain causes vCJD and BSE. Nature. 1997 Oct 2;389(6650):448-50, 526. doi: 10.1038/38925. PMID: 9333232.

Hill AF, Collinge J. Strain variations and species barriers. Contrib Microbiol. 2001;7:48-57. doi: 10.1159/000060375. PMID: 11923935.

Hoshino S, Imai M, Kobayashi T, Uchida N, Katada T. The eukaryotic polypeptide chain releasing factor (eRF3/GSPT) carrying the translation termination signal to the 3 -poly(A) tail of mRNA. J Biol Chem 1999;274:16677–80.

Hosoda N, Kobayashii T, Uchida N, Funakoshi Y, Kikuchi Y, Hoshino S, et al. Translation termination factor eRF3 mediates mRNA decay through the regulation of deadenylation. J Biol Chem 2003;278:38287–91.

Houston F, Andréoletti O. Animal prion diseases: the risks to human health. Brain Pathol. 2019

Mar;29(2):248-262. doi: 10.1111/bpa.12696. Epub 2019 Jan 22. PMID: 30588682; PMCID: PMC8028554.

Huang VJ, Stein KC, True HL. Spontaneous variants of the [RNQ+] prion in yeast demonstrate the
extensive conformational diversity possible with prion proteins. PLoS One. 2013 Oct 25;8(10):e79582. doi:
10.1371/journal.pone.0079582. PMID: 24205387; PMCID: PMC3808357.

Igel-Egalon A, Béringue V, Rezaei H, Sibille P. Prion Strains and Transmission Barrier Phenomena.
Pathogens. 2018 Jan 1;7(1):5. doi: 10.3390/pathogens7010005. PMID: 29301257; PMCID: PMC5874731.

Joshi T, Ahuja N. The Prion Basis of Progressive Neurodegenerative Disorders. Interdiscip Perspect Infect
Dis. 2023 Feb 14;2023:6687264. doi: 10.1155/2023/6687264. PMID: 36825209; PMCID: PMC9943612.

Kadnar ML, Articov G, Derkatch IL. Distinct type of transmission barrier revealed by study of multiple prion
determinants of Rnq1. PLoS Genet. 2010 Jan 22;6(1):e1000824. doi: 10.1371/journal.pgen.1000824.
PMID: 20107602; PMCID: PMC2809767.

Ke PC, Zhou R, Serpell LC, Riek R, Knowles TPJ, Lashuel HA, Gazit E, Hamley IW, Davis TP, Fändrich
M, Otzen DE, Chapman MR, Dobson CM, Eisenberg DS, Mezzenga R. Half a century of amyloids: past,
present and future. Chem Soc Rev. 2020 Aug 7;49(15):5473-5509. doi: 10.1039/c9cs00199a. Epub 2020
Jul 7. PMID: 32632432; PMCID: PMC7445747.

Kelly AC, Shewmaker FP, Kryndushkin D, Wickner RB. Sex, prions, and plasmids in yeast. Proc Natl Acad
Sci U S A. 2012 Oct 2;109(40):E2683-90. doi: 10.1073/pnas.1213449109. Epub 2012 Sep 4. PMID:
22949655; PMCID: PMC3479589.

Kelly AC, Wickner RB. Saccharomyces cerevisiae: a sexy yeast with a prion problem. Prion. 2013 May-
Jun;7(3):215-20. doi: 10.4161/pri.24845. PMID: 23764836; PMCID: PMC3783106.

Kelly AC, Busby B, Wickner RB. Effect of domestication on the spread of the [PIN+] prion in
Saccharomyces cerevisiae. Genetics. 2014 Jul;197(3):1007-24. doi: 10.1534/genetics.114.165670. Epub
2014 May 8. PMID: 24812307; PMCID: PMC4096356.

King CY. The Mutability of Yeast Prions. Viruses. 2022 Oct 25;14(11):2337. doi: 10.3390/v14112337.
PMID: 36366434; PMCID: PMC9696419.

Kochneva-Pervukhova NV, Paushkin SV, Kushnirov VV, Cox BS, Tuite MF, Ter-Avanesyan MD.
Mechanism of inhibition of Psi+ prion determinant propagation by a mutation of the N-terminus of the yeast
Sup35 protein. EMBO J. 1998 Oct 1;17(19):5805-10. doi: 10.1093/emboj/17.19.5805. PMID: 9755180;
PMCID: PMC1170908.

Kurahashi H, Ishiwata M, Shibata S, Nakamura Y. A regulatory role of the Rnq1 nonprion domain for prion
propagation and polyglutamine aggregates. Mol Cell Biol. 2008 May;28(10):3313-23. doi:
10.1128/MCB.01900-07. Epub 2008 Mar 10. PMID: 18332119; PMCID: PMC2423166.

Kurahashi H, Shibata S, Ishiwata M, Nakamura Y. Selfish prion of Rnq1 mutant in yeast. Genes Cells.
2009 May;14(5):659-68. doi: 10.1111/j.1365-2443.2009.01297.x. Epub 2009 Apr 15. PMID: 19371377.

Kurahashi H, Pack CG, Shibata S, Oishi K, Sako Y, Nakamura Y. [PSI(+)] aggregate enlargement in rnq1
nonprion domain mutants, leading to a loss of prion in yeast. Genes Cells. 2011(a) May;16(5):576-89. doi:
10.1111/j.1365-2443.2011.01511.x. Epub 2011 Apr 1. PMID: 21453425.

Kurahashi H, Oishi K, Nakamura Y. A bipolar personality of yeast prion proteins. Prion. 2011(b) Oct-
Dec;5(4):305-10. doi: 10.4161/pri.18307. Epub 2011 Oct 1. PMID: 22156730; PMCID: PMC4012401.

LeVine H 3rd. Quantification of beta-sheet amyloid fibril structures with thioflavin T. Methods Enzymol.
1999;309:274-84. doi: 10.1016/s0076-6879(99)09020-5. PMID: 10507030.

Li X, Rayman JB, Kandel ER, Derkatch IL. Functional role of Tia1/Pub1 and Sup35 prion domains:
directing protein synthesis machinery to the tubulin cytoskeleton. Mol Cell. 2014 Jul 17;55(2):305-18. doi:
10.1016/j.molcel.2014.05.027. Epub 2014 Jun 26. PMID: 24981173; PMCID: PMC4425694.

Lieberman AP, Shakkottai VG, Albin RL. Polyglutamine Repeats in Neurodegenerative Diseases. Annu

Rev Pathol. 2019 Jan 24;14:1-27. doi: 10.1146/annurev-pathmechdis-012418-012857. Epub 2018 Aug 8.
PMID: 30089230; PMCID: PMC6387631.

Liebman SW, All-Robyn JA. A non-Mendelian factor, [eta(+)], causes lethality of yeast omnipotent-
suppressor strains. Curr Genet. 1984 Oct;8(8):567-73. doi: 10.1007/BF00395701. PMID: 24177995.

Liebman SW, Bagriantsev SN, Derkatch IL. Biochemical and genetic methods for characterization of
[PIN+] prions in yeast. Methods. 2006 May;39(1):23-34. doi: 10.1016/j.ymeth.2006.04.010. PMID:
16793281.

Liebman SW, Chernoff YO. Prions in yeast. Genetics. 2012 Aug;191(4):1041-72. doi:
10.1534/genetics.111.137760. PMID: 22879407; PMCID: PMC3415993.

Lopez N, Aron R, Craig EA. Specificity of class II Hsp40 Sis1 in maintenance of yeast prion [RNQ+]. Mol
Biol Cell. 2003 Mar;14(3):1172-81. doi: 10.1091/mbc.e02-09-0593. PMID: 12631732; PMCID:
PMC151588.

Maddelein ML, Wickner RB. Two prion-inducing regions of Ure2p are nonoverlapping. Mol Cell Biol. 1999
Jun;19(6):4516-24. doi: 10.1128/MCB.19.6.4516. PMID: 10330190; PMCID: PMC104409.

Mathur V, Hong JY, Liebman SW. Ssa1 overexpression and [PIN(+)] variants cure [PSI(+)] by dilution of
aggregates. J Mol Biol. 2009 Jul 10;390(2):155-67. doi: 10.1016/j.jmb.2009.04.063. Epub 2009 May 5.
PMID: 19422835; PMCID: PMC2738641.

McGlinchey RP, Kryndushkin D, Wickner RB. Suicidal [PSI+] is a lethal yeast prion. Proc Natl Acad Sci U
S A. 2011 Mar 29;108(13):5337-41. doi: 10.1073/pnas.1102762108. Epub 2011 Mar 14. PMID: 21402947;
PMCID: PMC3069153.

Meriin AB, Zhang X, He X, Newnam GP, Chernoff YO, Sherman MY. Huntington toxicity in yeast model
depends on polyglutamine aggregation mediated by a prion-like protein Rnq1. J Cell Biol. 2002 Jun
10;157(6):997-1004. doi: 10.1083/jcb.200112104. Epub 2002 Jun 10. Erratum in: J Cell Biol 2002 Aug
5;158(3):591. PMID: 12058016; PMCID: PMC2174031.

Moriyama H, Edskes HK, Wickner RB. [URE3] prion propagation in Saccharomyces cerevisiae:
requirement for chaperone Hsp104 and curing by overexpressed chaperone Ydj1p. Mol Cell Biol. 2000
Dec;20(23):8916-22. doi: 10.1128/MCB.20.23.8916-8922.2000. PMID: 11073991; PMCID: PMC86546.

Nakayashiki T, Kurtzman CP, Edskes HK, Wickner RB. Yeast prions [URE3] and [PSI+] are diseases.
Proc Natl Acad Sci U S A. 2005 Jul 26;102(30):10575-80. doi: 10.1073/pnas.0504882102. Epub 2005 Jul
15. PMID: 16024723; PMCID: PMC1180808.

Newnam GP, Birchmore JL, Chernoff YO. Destabilization and recovery of a yeast prion after mild heat
shock. J Mol Biol. 2011 May 6;408(3):432-48. doi: 10.1016/j.jmb.2011.02.034. Epub 2011 Mar 15. PMID:
21392508; PMCID: PMC3095851.

Nizhnikov AA, Ryzhova TA, Volkov KV, Zadorsky SP, Sopova JV, Inge-Vechtomov SG, Galkin AP.
Interaction of Prions Causes Heritable Traits in Saccharomyces cerevisiae. PLoS Genet. 2016 Dec
27;12(12):e1006504. doi: 10.1371/journal.pgen.1006504. PMID: 28027291; PMCID: PMC5189945.

Norton J, Seah N, Santiago F, Sindi SS, Serio TR. Multiple aspects of amyloid dynamics *in vivo* integrate
to establish prion variant dominance in yeast. Front Mol Neurosci. 2024 Jul 30;17:1439442. doi:
10.3389/fnmol.2024.1439442. PMID: 39139213; PMCID: PMC11319303.

Osherovich LZ, Weissman JS. Multiple Gln/Asn-rich prion domains confer susceptibility to induction of the
yeast [PSI(+)] prion. Cell. 2001 Jul 27;106(2):183-94. doi: 10.1016/s0092-8674(01)00440-8. PMID:
11511346.

Park SK, Hong JY, Arslan F, Kanneganti V, Patel B, Tietsort A, Tank EMH, Li X, Barmada SJ, Liebman
SW. Overexpression of the essential Sis1 chaperone reduces TDP-43 effects on toxicity and proteolysis.
PLoS Genet. 2017 May 22;13(5):e1006805. doi: 10.1371/journal.pgen.1006805. PMID: 28531192;
PMCID: PMC5460882.

Patino MM, Liu JJ, Glover JR, Lindquist S. Support for the prion hypothesis for inheritance of a phenotypic
trait in yeast. Science. 1996 Aug 2;273(5275):622-6. doi: 10.1126/science.273.5275.622. PMID: 8662547.

Pattison, I. H. (1965) Experiments with scrapie with special reference to the nature of the agent and the
pathology of the disease: Slow, latent and temperate virus infections. NINDB Monograph 2, 249–257.

Pei F, DiSalvo S, Sindi SS, Serio TR. A dominant-negative mutant inhibits multiple prion variants through a
common mechanism. PLoS Genet. 2017 Oct 30;13(10):e1007085. doi: 10.1371/journal.pgen.1007085.
PMID: 29084237; PMCID: PMC5679637.

Resende CG, Outeiro TF, Sands L, Lindquist S, Tuite MF. Prion protein gene polymorphisms in
Saccharomyces cerevisiae. Mol Microbiol. 2003 Aug;49(4):1005-17. doi: 10.1046/j.1365-
2958.2003.03608.x. PMID: 12890024.

Rose M, Winston F, Hieter P. (1990) Methods in yeast genetics — A laboratory course manual by pp 198.
Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York. ISBN 0-87969-354-1.

Salnikova AB, Kryndushkin DS, Smirnov VN, Kushnirov VV, Ter-Avanesyan MD. Nonsense suppression in
yeast cells overproducing Sup35 (eRF3) is caused by its non-heritable amyloids. J Biol Chem. 2005 Mar
11;280(10):8808-12. doi: 10.1074/jbc.M410150200. Epub 2004 Dec 23. PMID: 15618222.

Santoso A, Chien P, Osherovich LZ, Weissman JS. Molecular basis of a yeast prion species barrier. Cell.
2000 Jan 21;100(2):277-88. doi: 10.1016/s0092-8674(00)81565-2. PMID: 10660050.

Satpute-Krishnan P, Langseth SX, Serio TR. Hsp104-dependent remodeling of prion complexes mediates
protein-only inheritance. PLoS Biol. 2007 Feb;5(2):e24. doi: 10.1371/journal.pbio.0050024. PMID:
17253904; PMCID: PMC1779812.

Schlumpberger M, Prusiner SB, Herskowitz I. Induction of distinct [URE3] yeast prion strains. Mol Cell Biol.
2001 Oct;21(20):7035-46. doi: 10.1128/MCB.21.20.7035-7046.2001. PMID: 11564886; PMCID:
PMC99879.

Scott M, Foster D, Mirenda C, Serban D, Coufal F, Wälchli M, Torchia M, Groth D, Carlson G, DeArmond
SJ, Westaway D, Prusiner SB. Transgenic mice expressing hamster prion protein produce species-specific
scrapie infectivity and amyloid plaques. Cell. 1989 Dec 1;59(5):847-57. doi: 10.1016/0092-8674(89)90608-
9. PMID: 2574076.

Sharma J, Liebman SW. [PSI(+)] prion variant establishment in yeast. Mol Microbiol. 2012 Nov;86(4):866-
81. doi: 10.1111/mmi.12024. Epub 2012 Sep 24. PMID: 22998111; PMCID: PMC3543502.

Sharma J, Liebman SW. Variant-specific prion interactions: Complicating factors. Cell Logist. 2013 Jan
1;3(1):e25698. doi: 10.4161/cl.25698. Epub 2013 Sep 12. PMID: 24475372; PMCID: PMC3891757.

Sherman, F., Fink, G., Hicks, J. (1987) *Methods in Yeast Genetics: A Laboratory Course Manual.* Cold
Spring Harbor Laboratory Press. ISBN 9780879691974.

Sharma A, Bruce KL, Chen B, Gyoneva S, Behrens SH, Bommarius AS, Chernoff YO. Contributions of the
Prion Protein Sequence, Strain, and Environment to the Species Barrier. J Biol Chem. 2016 Jan
15;291(3):1277-88. doi: 10.1074/jbc.M115.684100. Epub 2015 Nov 12. PMID: 26565023; PMCID:
PMC4714215.

Shewmaker F, Mull L, Nakayashiki T, Masison DC, Wickner RB. Ure2p function is enhanced by its prion
domain in Saccharomyces cerevisiae. Genetics 2007;176:1557–65.

Shibata S, Kurahashi H, Nakamura Y. Localization of prion-destabilizing mutations in the N-terminal non-
prion domain of Rnq1 in Saccharomyces cerevisiae. Prion. 2009 Oct-Dec;3(4):250-8. doi:
10.4161/pri.3.4.10388. Epub 2009 Oct 20. PMID: 20009538; PMCID: PMC2807699.

Sikorski RS, Hieter P. A system of shuttle vectors and yeast host strains designed for efficient
manipulation of DNA in Saccharomyces cerevisiae. Genetics. 1989 May;122(1):19-27. doi: 10.
1093/genetics/122.1.19. PMID: 2659436; PMCID: PMC1203683.

Sondheimer N, Lindquist S. Rnq1: an epigenetic modifier of protein function in yeast. Mol Cell. 2000
Jan;5(1):163-72. doi: 10.1016/s1097-2765(00)80412-8. PMID: 10678178.

Sondheimer N, Lopez N, Craig EA, Lindquist S. The role of Sis1 in the maintenance of the [RNQ+] prion.
EMBO J. 2001 May 15;20(10):2435-42. doi: 10.1093/emboj/20.10.2435. PMID: 11350932; PMCID:
PMC125465.

Taneja V, Maddelein ML, Talarek N, Saupe SJ, Liebman SW. A non-Q/N-rich prion domain of a foreign
prion, [Het-s], can propagate as a prion in yeast. Mol Cell. 2007 Jul 6;27(1):67-77. doi:
10.1016/j.molcel.2007.05.027. PMID: 17612491; PMCID: PMC1995001.

Tuite MF, Mundy CR, Cox BS. Agents that cause a high frequency of genetic change from [psi+] to [psi-] in
Saccharomyces cerevisiae. Genetics. 1981 Aug;98(4):691-711. doi: 10.1093/genetics/98.4.691. PMID:
7037537; PMCID: PMC1214469.

Villali J, Dark J, Brechtel TM, Pei F, Sindi SS, Serio TR. Nucleation seed size determines amyloid
clearance and establishes a barrier to prion appearance in yeast. Nat Struct Mol Biol. 2020 Jun;27(6):540-
549. doi: 10.1038/s41594-020-0416-6. Epub 2020 May 4. PMID: 32367069; PMCID: PMC7293557.

Vitrenko YA, Pavon ME, Stone SI, Liebman SW. Propagation of the [PIN+] prion by fragments of Rnq1
fused to GFP. Curr Genet. 2007(a) May;51(5):309-19. doi: 10.1007/s00294-007-0127-0. Epub 2007 Apr 6.
PMID: 17415568; PMCID: PMC2597802.

Vitrenko YA, Gracheva EO, Richmond JE, Liebman SW. Visualization of aggregation of the Rnq1 prion
domain and cross-seeding interactions with Sup35NM. J Biol Chem. 2007(b) Jan 19;282(3):1779-87. doi:
10.1074/jbc.M609269200. Epub 2006 Nov 22. PMID: 17121829.

Wegrzyn RD, Bapat K, Newnam GP, Zink AD, Chernoff YO. Mechanism of prion loss after Hsp104
inactivation in yeast. Mol Cell Biol. 2001 Jul;21(14):4656-69. doi: 10.1128/MCB.21.14.4656-4669.2001.
PMID: 11416143; PMCID: PMC87136.

Weissmann C. Mutation and selection of prions. PLoS Pathog. 2012;8(3):e1002582. doi:
10.1371/journal.ppat.1002582. Epub 2012 Mar 29. PMID: 22479179; PMCID: PMC3315487.

Westergard L, True HL. Wild yeast harbour a variety of distinct amyloid structures with strong prion-
inducing capabilities. Mol Microbiol. 2014a Apr;92(1):183-93. doi: 10.1111/mmi.12543. Epub 2014 Mar 7.
PMID: 24673812; PMCID: PMC4708258.

Westergard L, True HL. Extracellular environment modulates the formation and propagation of particular
amyloid structures. Mol Microbiol. 2014b May;92(4):698-715. doi: 10.1111/mmi.12579. Epub 2014 Apr 9.
PMID: 24628771; PMCID: PMC4739623.

Wickner RB. [URE3] as an altered URE2 protein: evidence for a prion analog in Saccharomyces
cerevisiae. Science. 1994 Apr 22;264(5158):566-9. doi: 10.1126/science.7909170. PMID: 7909170.

Wickner RB, Dyda F, Tycko R. Amyloid of Rnq1p, the basis of the [PIN+] prion, has a parallel in-register
beta-sheet structure. Proc Natl Acad Sci U S A. 2008 Feb 19;105(7):2403-8. doi:
10.1073/pnas.0712032105. Epub 2008 Feb 11. PMID: 18268327; PMCID: PMC2268149.

Wickner RB, Edskes HK, Kryndushkin D, McGlinchey R, Bateman D, Kelly A. Prion diseases of yeast:
amyloid structure and biology. Semin Cell Dev Biol. 2011(a) Jul;22(5):469-75. doi:
10.1016/j.semcdb.2011.02.021. Epub 2011 Feb 21. PMID: 21345375; PMCID: PMC3117979.

Wickner RB, Edskes HK, Bateman D, Kelly AC, Gorkovskiy A. The yeast prions [PSI+] and [URE3] are
molecular degenerative diseases. Prion. 2011(b) Oct-Dec;5(4):258-62. doi: 10.4161/pri.17748. Epub 2011
Oct 1. PMID: 22052353; PMCID: PMC4012404.

Wickner RB, Son M, Edskes HK. Prion Variants of Yeast are Numerous, Mutable, and Segregate on
Growth, Affecting Prion Pathogenesis, Transmission Barriers, and Sensitivity to Anti-Prion Systems.
Viruses. 2019 Mar 9;11(3):238. doi: 10.3390/v11030238. PMID: 30857327; PMCID: PMC6466074.

Zhou P, Derkatch IL, Liebman SW. The relationship between visible intracellular aggregates that appear
after overexpression of Sup35 and the yeast prion-like elements [PSI(+)] and [PIN(+)]. Mol Microbiol. 2001
Jan;39(1):37-46. doi: 10.1046/j.1365-2958.2001.02224.x. PMID: 11123686.

Zhouravleva GA, Bondarev SA, Trubitsina NP. How Big Is the Yeast Prion Universe? Int J Mol Sci. 2023
Jul 19;24(14):11651. doi: 10.3390/ijms241411651. PMID: 37511408; PMCID: PMC10380529.

1671

Figures

Fig 1. Genetic screen for mutations in Rnq1 that, when substituted for Rnq1$_{WT}$ in a [*PIN$^+$*] strain (Pin$^+$ phenotype), lead to a Pin$^-$ phenotype. (**A**) Structural organization of the Rnq1 protein. Shown are NPD (aa 1-132, yellow) with the position of the T27P mutation, and PD (aa 133-405) with a stretch of ten QG repeats (aa 153-172, blue), four distinct QN-regions (brown; QN1 through QN4), and five hydrophobic repeats with the LA^{S}/$_A$ ^{L}/$_M$ A core (light grey; A through E). The QN1 region is aa 185-198, QN2 - aa 218-263, QN3 - aa 279-319, QN4 - aa 337-405; patterned blocs indicate QN-rich repeats within QN2, QN3 and QN4). (**B)** Scheme of the genetic screen (see text).

Fig 2. The T27P mutation in Rnq1 NPD leads to a dramatic reduction of the Pin$^+$ phenotype when Rnq1$_{T27P}$ is substituted for Rnq1$_{WT}$ in a [*PIN$^+$*] strain. (**A**) Western blot analysis of the expression of Rnq1$_{T27P}$ and Rnq1$_{T27P}$ in *the rnq1-Δ* strain. The *rnq1-Δ* YID193 strain was transformed with the indicated *CEN LEU2* plasmids. For *RNQ1$_{WT}$* the pID129 plasmid was used. For*RNQ1$_{T27P}$*, shown is expression from the original plasmid from the mutagenized library and from two independent isolates of the pID335 plasmid where the T27P mutation was introduced by site-directed mutagenesis. Expression of Sup35 is used as a loading control. Both panels are from the same Western blot: anti-Rnq1 and anti-Sup35 antibodies were added simultaneously. Mobilities of Rnq1 and Sup35 are as expected based on protein molecular weights. Expression of Rnq1$_{T27P}$ is ~2-fold lower compared to Rnq1$_{WT}$. (**B)** Rnq1$_{T27P}$ leads to a dramatic reduction of the Pin$^+$ phenotype when substituted for Rnq1$_{WT}$ in a [*PIN$^+$*] strain. Shown is the image of the YID146.3 cells with *RNQ1$_{T27P}$* (or indicated control plasmids) substituted for *RNQ1$_{WT}$* incubated on SD-Ade media for 14 days at 30^{0}C after previously being grown either on the [*PSI$^+$*] inducing SGal-LeuHis media, or on the control SD-LeuHis (two passages). Each row includes patches of

5 independent transformants. See Fig S1 for images of the same cultures on SD-LeuHis and and

SGal-LeuHis from which cells were transferred onto SD-Ade. (**C**) Upon high-level expression of

Sup35NM-YFP in the cultures where $RNQ1_{T27P}$ is substituted for $RNQ1_{WT}$, the *de novo* forming

Sup35 aggregates are rare, but similar in appearance to aggregates forming in the control

$RNQ1_{WT}$ cultures. Shown are fluorescent and bright-field images of cells carrying *GAL-*

SUP35NM-YFP and the indicated *RNQ1* construct grown on SGal-LH for ~48hrs at 30^0C after

the plasmid shuffle described in Fig 1B. See Table 1 for quantification. A typical group of cells is

shown for the $RNQ1_{WT}$ culture. For $RNQ1_{T27P}$, all cells have the evenly distributed fluorescence

in a typical group of cells representative of the 97.5% of the culture. Additionally shown for

$RNQ1_{T27P}$ are rare cells with dot-, line-, and ring- aggregates, as well as aggregate-containing

mother-daughter pairs (a line and a dot, or a ring shared by a mother and a daughter).

1707

Fig 3. The T27P mutation does not interfere with the ability of the Rnq1 protein to acquire

a heritable Pin⁺ state upon overexpression, indicative of the *de novo* formation of the

[PIN^+_{T27P}] prion. For the description of Experiment 1 (**A**) and Experiment 2 (**B**) see Results and

Materials and Methods. The (++), (+), and ($\pm$) indicate the relative level of [*PSI⁺*] induction based

on growth on adenineless media compared to lack of [*PSI⁺*] induction (-), with (++) being the

most robust. In (B) some negative control experiments are grouped together.

* Occasional Ade⁺ colonies were detected after induction of the *GAL* promoter-driven Sup35NM-

YFP in cultures where *CUP1*-driven $RNQ1_{T27P\ (or\ WT)}$ constructs were not overexpressed. This is

indicative of a low-level of induction of Rnq1-based prions, probably due to residual expression

of *CUP1-RNQ1* constructs in all or some cells on regular SD media which contains $0.16\mu M$

$CuSO_4$. The number of such Ade⁺ colonies gradually increased after multiple passages prior to

the induction of Sup35NM-YFP but never reached the level of Pin+ phenotypes observed in

cultures that went through transient Rnq1$_{T27P (or WT)}$ overexpression on 50µM CuSO4.

1721

Fig 4. Full-length Rnq1$_{T27P}$ and ∆B1C2D3$_{T27P}$ can form amyloid fibers *in vitro*. (A) Western

blot analysis of bacterially expressed and purified ∆B1C2D3$_{T27P}$ (T27P; two independent

constructs) and ∆B1C2D3$_{WT}$ (WT). The expected size for both constructs is 27kDa. Note slightly

slower mobility for ∆B1C2D3$_{T27P}$. Anti-His tag antibody was used for detection. **(B)** Kinetics of *in*

vitro aggregation of ∆B1C2D3$_{T27P}$ and ∆B1C2D3$_{WT}$ monitored by ThT fluorescence. Samples and

concentrations are indicated on the graph. **(C)** TEM micrographs of negatively stained fibers of

Rnq1$_{T27P}$ and Rnq1$_{WT}$. Scale bars are for 100nm. Note lateral association of fibrils in the fibers for

Rnq1$_{T27P}$ and twisted appearance of Rnq1$_{WT}$ fibers.

1730

Fig 5. The T27P mutation creates a barrier for the transmission of the prion state from

[*PIN+$_{WT}$*] to Rnq1$_{T27P}$. (**A**) Scheme of experiments testing the frequency of [*PIN+*] loss after

substituting Rnq1$_{T27P}$ for Rnq1$_{WT}$ and mitotic stability of [*PIN+$_{T27P}$*] isolates. See text. Red circle

indicates a step at which yeast were colony purified for the analysis of the fraction of [*PIN+$_{T27P}$*]

cells in the cultures right after substituting *RNQ1$_{T27P}$* for *RNQ1$_{WT}$* (data shown in Table 2 and

illustrated in (B); also corresponds to Round 1 in (C) and (D), and, for other donor and recipient

construct combinations, to data in Fig 7B and Round 1 in Figs 7C through 7G). Green circle

indicates the step at which [*PIN+$_{T27P}$*] isolates were colony purified for the analysis of their mitotic

stability (corresponds to data in Table 3 and Round 2 in (C) and (D), and, for other donor and

recipient combinations, to Round 2 in Figs 7C through 7G). (**B**) After-shuffle cultures expressing

Rnq1$_{T27P}$ contain Pin+ and Pin- cells. Sample SD-Ade plates after [*PSI+*] induction on SGal-

LeuHis with patches expressing the indicated Rnq1 allele or carrying the empty vector. Shown is

growth on SD-Ade after 13 days of incubation at 20^{0}C. All patches were Ade$^-$ after incubation on

the control SD-LeuHis media (not shown). **(C)** Increase of mitotic stability of [PIN^+_{T27P}] indicative

of gradual conformational adaptation of the prion. See text and (A) for the description of the

experiment. Data for Round 1 come from colony purification of 12 Rnq1$_{T27P}$ post-shuffle cultures

transferred from FOA to SD-LeuHis. Data for Round 2 come from colony purification of the Pin$^+$

Rnq1$_{T27P}$ isolates obtained in Round 1; 13 out of 15 obtained isolates were analyzed (the

remaining 2 were lost due to contamination). For Round 3, seven Pin$^+$ cultures used in Round 2

analysis were randomly chosen, and for each three randomly chosen Pin$^+$ isolates were colony

purified and scored for Pin$^+$ (a total of 21 data points). In Round 4, the analysis was continued for

all twelve cultures originating from four out of seven Round 2 cultures (three randomly chosen

Pin$^+$ isolates for each culture, a total of 36 data points) and for partial lineages of two more

Round 2 isolates (a total of 6 more data points). Data for positive (Rnq1$_{WT}$ shuffled to Rnq1$_{WT}$)

and negative (Rnq1$_{WT}$ to EV) controls is not included for simplicity, but these experiments were

carried through Round 2 on a similar scale and through further rounds on a limited scale.

[PIN^+_{WT}] maintained ~100% stability in all isolates tested, with no switching to less stable

variants, and yeast with shuffled-in EV did not accumulate Pin$^+$ cells. (**D**) Maturation of post-

transmission [PIN^+_{T27P}] to heritable prion variants with characteristic stabilities. See Results for

the description of this experiment and explanation of pedigrees. The R4, R5 and R6 labels

correspond to the round of the analysis of mitotic stability of post-shuffle [PIN^+_{T27P}]. Asterisks

mark directly related parts of the pedigree of the meta-stable [PIN^+_{T27P}] prion variant that is prone

to yield more stable (red) and less stable (green) [PIN^+_{T27P}] variants . Differences between

pedigrees 1, 2, and 3 are statistically significant for all rounds of analysis (p<0.01). There are no

significant differences within pedigrees 1 and 2 between shown rounds of analysis. For pedigree

3, there is no significant difference between R4 and R5, but for R6 there is a significant

difference (p<0.01) due to gradual stabilization of [PIN^+_{T27P}] in some isolates (green asterisks).

1768

Fig 6. The transmission barrier between [*PIN⁺WT*] and Rnq1T27P is not caused by the

inability of Rnq1WT to co-aggregate with Rnq1WT or by loss of the [*PIN⁺*] prion during co-

expression of Rnq1WT and Rnq1T27P. (**A**) Rnq1$_{T27P}$ can join the [*PIN⁺WT*] prion aggregates. The

[*PIN⁺*] 74-D694 strain was co-transformed with the *CEN LEU2 CUP1-RNQ1WT-CFP* and *CEN*

URA3 CUP-RNQ1T27P-YFP plasmids. Cultures of co-transformants were grown overnight (~to

mid-log) in no-copper SD-LeuHis media, diluted to OD$_{595}$=1.0 into SD-LeuHis+5μM CuSO$_4$ and

grown for 2 more hours prior to observing the cells with or without Hex treatment. A

representative group of cells is shown. See Fig S2A for an image of a larger field and analysis of

Rnq1$_{WT}$ and Rnq1$_{T27P}$ co-localization individual cells. (**B**) Rnq1$_{T27P}$ does not affect the Pin$^+$

phenotype when co-expressed with Rnq1$_{WT}$ in a [*PIN⁺*] stain. Shown are the images of the

transformants of the YID146.3 cells carrying the *GAL-SUP35NM-YFP* [*PSI⁺*] inducing construct

along with the indicated plasmid combinations. "Galactose" and "Glucose" indicate SGal-

UraLeuHis and SD-UraLeuHis media, respectively. Images on these media were taken after 48

hours of growth at 30^{0}C. From these media yeast were replica plated to SD-Ade. Images were

taken after 14 days at 30^{0}C. Note that these are patches from the plates shown on Fig 2B and

Fig S1 but before plasmid shuffle.

1785

Fig 7. The transmission barrier created by the T27P mutation in the NPD is imposed

through the first two QN-rich regions in Rnq1 PD. (A) Deletion of B1 and B1C2 but not 2D

and 3E regions in the PD of Rnq1$_{WT}$ reduces the transmission barrier from [*mini-PIN⁺WT*]s

towards Rnq1$_{FL-T27P}$. The [*PSI⁺*] induction test was done in cultures where the Rnq1$_{FL-WT}$, Rnq1$_{FL-}$

$_{T27P}$ recipient constructs (or EV) were substituted for the indicated deletion constructs in the [*mini-*

PIN⁺WT] or [*PIN⁺FL-WT*] prion donor strains. Shown are images of growth on SD-Ade plates replica

plated from either the [*PSI⁺*] inducing SGal-LeuHis or the control glucose media and grown for 10

days at 30^{0}C. **(B – G)** Analysis of the role of QN-rich regions in the NPD of Rnq1 on the

transmission barriers and subsequent conformational adaptation of Rnq1-based prions in the

presence of the T27P mutation in the NPD. Experiments were performed as described in Fig 5.

Donor and recipient constructs for prion transmission are indicated; nomenclature marks the

deleted regions within the PD (ΔB1C2, ΔB1, Δ2D, or Δ3E; see Fig 1A) compared to the full-length

Rnq1 protein (FL) and the presence or absence of the T27P mutation in the construct (T27P or

WT, respectively). Shown is the fraction of prion-carrying cells in the cultures after plasmid

shuffle (B and R1 in C – G) and mitotic stability of the resulting prions during further propagation

of randomly chosen isolates (R2 through R4; ~19 cell generations between each round of

analysis). In (D) asteriscs indicate related progeny for cultures in R3 and R4 highlighting the

establishment of [$PIN^+_{\Delta B1C2}$] variants with high (red and blue) and low (green) mitotic stability.

Experiments in (B – G) were conducted in parallel, so all data can be compared between

the graphs. Some control experiments (e.g. analysis of transmissions between non-mutant

constructs and lack of transmissions in substitutions for EV) are omitted on graphs for simplicity:

the results were as expected and in full agreement with Kadnar et al. (2010). All conclusions in

Results are based on statistically significant differences or lack of thereof (0.05 or 0.01

probability cutoff).

1810

Fig 8. The T27P mutation in the NPD of Rnq1 promotes the formation of liquid-like

droplets by Rnq1$_{FL-T27P}$-YFP expressed in [pin^-] cells even at levels close to the expression

of endogenous Rnq1. (A) Unlike Rnq1$_{FL-WT}$-YFP, Rnq1$_{FL-T27P}$-YFP forms aggregates in [pin^-]

cells and these aggregates are not amyloid. Cultures of isogenic [PIN^+] and [pin^-] strains

transformed with plasmids carrying the indicated constructs under the control of *CUP1* promoter

were grown overninght in SD plasmid selective media to OD$_{595}$ ~ 0.8. Then expression of the

constructs was induced by addition of $CuSO_4$ to 50μM and growth was continued for 4 more

hours prior to ThT staining and imaging. **(B - C)** Aggregates formed by Rnq1$_{FL-T27P}$-YFP in [*pin*$^-$]

cells are liquid-like droplets. Yeast transformants were grown overnight in liquid SD media

selective for plasmids. Then OD$_{595}$ was adjusted to 1.0, $CuSO_4$ was added to 5μM, and cultures

were grown for two more hrs before Hex treatment. (C) is quantification of data in experiment

illustrated in (B). Asterisks indicate statistically significant differences (p<0.01). **(D)** Rnq1$_{FL-T27P}$-

YFP forms liquid-like droplets in *rnq1-Δ* cells. Experiment was performed as described for (B -

C). **(E)** Rnq1$_{FL-WT}$-CFP does not join liquid-like droplets formed by Rnq1$_{FL-T27P}$-YFP in [*pin*$^-$] cells.

The [*pin*$^-$] strain was co-transformed with the *CEN LEU2 CUP1-RNQ1$_{WT}$-CFP* and *CEN URA3*

CUP-RNQ1$_{T27P}$-YFP plasmids. Experiment was performed exactly as in Fig 6A where Rnq1$_{FL-}$

$_{T27P}$-YFP is shown co-aggregate with Rnq1$_{FL-WT}$-CFP in [*PIN*$^+$$_{WT}$] cells.

1828

1829   **Fig 9. Models explaining the establishment of heritable [*PIN*$^+$$_{T27P}$] variants after across-the-**

1830   **barrier transmissions from [*PIN*$^+$$_{WT}$].** See text.

1831

Supporting information

Fig S1. The T27P mutation in Rnq1 does not lead to inhibition of growth in the presence of

the highly expressed Sup35NM-YFP construct. *RNQ1$_{WT}$* was substituted with *RNQ1$_{T27P}$* or

control plasmids (indicated) in a [*PIN*$^+$] [*psi*$^-$] *rnq1-Δ* strain as described in Fig. 1B. Shown are the

images of patches of five transformants for each plasmid substitution after they were replica

plated from the *RNQ1$_{WT}$* (*URA3*) counter-selecting FOA media to Sup35NM-YFP inducing SGal-

LeuHis media or non-inducing SD-LeuHis. Before imaging, cells were incubated at 30^{0}C for

~48hrs. These are images of plates used for replica plating to adenineless media shown in

Figure 2B.

1841

Fig S2. Rnq1$_{T27P}$ joins [$PIN^+{}_{WT}$] prion aggregates as efficiently as Rnq1$_{WT}$. The [PIN^+] 74-

D694 strain was co-transformed, respectively, with the *CEN LEU2 CUP1-RNQ1$_{WT}$-CFP* and *CEN*

URA3 CUP-RNQ1$_{T27P}$-YFP plasmids (**A**), or with *CEN LEU2 CUP1-RNQ1$_{WT}$-CFP* and *CEN*

URA3 CUP-RNQ1$_{WT}$-YFP (**B**). Cultures of co-transformants were grown overnight (~to mid-log)

in no-copper SD-LeuHis media, diluted to OD$_{595}$=1.0 into SD-LeuHis+5μM CuSO$_4$ and grown for

2 more hours prior to observing the cells. In both images (A) and (B) CFP- and YFP-marked

constructs co-localized in bright fluorescent foci in most cells (squares 1 and 2 for both (A) and

(B); enlarged images are shown in panels below). However, there were cells with bright foci, for

which only YFP or only CFP fluorescence is easily detectable on merged images. To test if there

are cells in images from experiment (A) where only Rnq1$_{WT}$ but not Rnq1$_{T27P}$ joins [$PIN^+{}_{WT}$]

aggregates, at total of ~150 cells from three different transformants with only CFP detectable on

merge images were analyzed more carefully. For all of them, either no evenly distributed YFP

fluorescence was detected, indicative of the loss or rearrangement of the *CUP-RNQ1$_{T27P}$-YFP*

plasmid (see e.g. square 4 on Fig. S2A), or weak but definitely co-localizing YFP foci were

detected on YFP images, indicative of reduced amounts of Rnq1$_{T27P}$-YFP in the cell (e.g. square

5 on Fig. S2A). We also asked if there were Rnq1$_{T27P}$-YFP aggregates not incorporating Rnq1$_{WT}$.

However, after analyzing ~150 cells with only YFP detectable on the merge, we concluded that

the lack of the CFP signal within these foci was also either due to the *CUP-RNQ1$_{WT}$-CFP*

plasmid loss (e.g. square 3 on Fig. S2A) or reduced abundance of Rnq1$_{WT}$-CFP (e.g. square 6

on Fig. S6A). Analogous analysis in experiment (B) led to the same conclusion: squares 3 and 4

represent cells with reduced fluorescence for one of the constructs, and squares 5 and 6 – cells

where one of the constructs is lost or rearranged.

1864

Fig S3. Growth on media supplemented with 5μM CuSO$_4$ results in the level of Rnq1$_{T27P}$-YFP expression similar to that of chromosomally encoded Rnq1$_{WT}$. A transformant of a [*pin*$^-$] strain with the *CEN URA3 CUP-RNQ1$_{T27P}$-YFP* plasmid was grown overnight (~to mid-log) in the no-copper plasmid-selective SD-Ura media, then diluted to OD$_{595}$=1.0 into SD-Ura+5μM CuSO$_4$ and grown for 2 more hours prior to protein extraction (see Materials and Methods and Supplementary Materials and Methods). Approximately 30μg of total protein were separated on an SDS-PAGE gel. Rnq1 was detected by Western blot with polyclonal antibodies raised in rabbits against the first 185 aa of Rnq1 (type 1A Ab).

Table S1. Similar arrays of [*PSI*$^+$] variants are induced in cultures expressing Rnq1$_{T27P}$, Rnq1$_{WT}$, or not expressing Rnq1 after shuffling out the *CEN URA3 RNQ1$_{WT}$* [*PIN$^+_{WT}$*] maintainer plasmid. See Results and Materials and Methods for the description of the experiment. [*PSI*$^+$]s isolated from at least eight independent transformants carrying the indicated plasmids were analyzed by the color test on YPD, growth on adenineless media and curability by GuHCl. Results of all three assays were consistent for ~99% [*PSI*$^+$] variants.

Table S2. PCR primers used in this study. Sites for restriction endonucleases used for cloning are underlined; the ATG start codon is in bold.

File S1. Supplementary Materials and Methods.

A

B

A

B

C

Figure 3

A

[*PSI⁺*] induction
for indicated constructs

Rnq1 ↑ (CuSO₄)	Sup35NM-YFP ↑ (Gal)		Rnq1_{T27P}	Rnq1_{WT}	EV
YES	YES		+	+	-
NO	YES	Replica plate to -Ade	- *	- *	-
YES	NO		-	-	-
NO	NO		-	-	-

B

[*PSI⁺*] induction
for indicated constructs

Rnq1 ↑ (CuSO₄)	Passages on Glu w/o extra CuSO₄	Sup35NM-YFP ↑ (Gal)		Rnq1_{T27P}	Rnq1_{WT}	EV
YES	0	YES		++	+	-
YES	1	YES		++	+	-
YES	2	YES		+	±	-
YES	3	YES	Replica plate to -Ade	±	±	-
YES	4	YES		±	±	-
YES	0, 1, 2, 3 or 4	NO		-	-	-
NO	0, 1, 2, 3 or 4	NO		-	-	-
NO	0, 1, 2, 3 or 4	YES		- *	- *	-

Figure 4

A

B

C

Rnq1$_{T27P}$ Rnq1$_{T27P}$

Rnq1$_{WT}$ Rnq1$_{WT}$

Figure 5

A

[PIN⁺][psi⁻] rnq1-Δ
HIS3 GAL-SUP35NM-YFP
URA3 RNQ1_{WT} maintainer

LEU2 RNQ1_{T27P}

SD-UraLeuHis
co-express *RNQ1_{WT}* and *RNQ1_{T27P}*

URA3 RNQ1_{WT}

SD-LeuHis
express only *RNQ1_{T27P}*

Colony purify on SD-LeuHis

Patch colonies to SD-LeuHis

Score for Pin⁺

B

Rnq1_{T27P}

Rnq1_{WT}

EV

C   D

A

B

Figure 9

Across-the-barrier transmission

Subsequent mitotic propagation

Strain array

Variant 1 (retained)

Variant 3 (lost)

Variant 2 (retained)

Conformational adaptation

Variant 1 (occasionally lost)

Variant 2 (retained; adaptation of variant 1)

Donor prion [*PIN*+WT]

[*PIN*+T27P] prion (different strains)

Non-prion (soluble) conformation