IL-27 limits HSPC differentiation during infection and protects from stem cell exhaustion

Daniel L. Aldridge[1], Zachary Lanzar[1], Anthony T. Phan[1], David A. Christian[1], Ryan Pardy[1], Booki

Min[2], Ross M. Kedl[3], Christopher A. Hunter[1]

[1]University of Pennsylvania School of Veterinary Medicine, Philadelphia, PA, USA

[2]Department of Microbiology and Immunology, Feinber School of Medicine, Northwestern

University, Chicago, IL, USA

[3]University of Colorado, Anschuitz Medical Campus, Aurora, CO, USA

Abstract

Many inflammatory stimuli can induce progenitor cells in the bone marrow to produce

increased numbers of myeloid cells as part of the process of emergency myelopoiesis. These events

are associated with innate training and can have long-term impacts on hematopoietic stem and

progenitor cell (HSPC) development but can also compromise their function. While many

cytokines support emergency myelopoiesis, less is known about the mechanisms that temper these

events. When mice that lack the cytokine IL-27 were infected with *Toxoplasma gondii*, there was

enhanced generation of monocyte progenitors and increased numbers of inflammatory monocytes.

In the bone marrow of infected mice there was increased production of IL-27 that localized with

HSPCs and a survey of cytokine receptor expression highlighted that HSPCs were uniquely poised

to respond to IL-27. Furthermore, the use of *in vitro* differentiation assays and mixed bone marrow

chimeras revealed that HSPCs from IL-27 deficient mice are pre-disposed towards the monocyte

lineage. Additional studies highlighted that after infection loss of the IL-27R resulted in reduced

HSPC fitness that manifested as reduced proliferative responses and a decreased ability to

reconstitute the hematopoietic system. Thus, the ability of IL-27 to act on HSPC provides a

regulatory brake on differentiation to limit monocyte induction and preserve HSPC stemness.

Introduction

In response to inflammatory stimuli, cytokine production can directly affect progenitor populations in the bone marrow leading to alterations in hematopoiesis (Collins et al., 2021). For example, in response to microbial challenges, elevated levels of GM-CSF, IFN-γ, and IL-6 can modulate hematopoietic stem and progenitor cell (HSPC) differentiation and enhance myelopoiesis (Baldridge et al., 2010; Kaufmann et al., 2018; Khan et al., 2020; Link, 2000; MacNamara et al., 2011a; MacNamara et al., 2011b; Maeda et al., 2005; Matatall et al., 2014; Miner et al., 2021; Mirantes et al., 2014; Reynaud et al., 2011). These events can also lead to long-term epigenetic and metabolic changes in HSPCs that form the basis for the process of innate training (Ochando et al., 2023). Thus, many of the cytokines produced during different infections are associated with innate training (Kaufmann et al., 2018; Khan et al., 2020). For example, after challenge with *Mycobacterium avium* or SARS-COV-2, IFN-γ and IL-6, respectively, can skew HSPCs towards myeloid output and increase the functionality of their downstream progeny (Cheong et al., 2023; Matatall et al., 2014).

While innate training is associated with long-term changes to hematopoietic stem cells (HSC), inflammation can also reduce HSC functionality, which can manifest as decreased proliferative and differentiation potential. To mitigate these adverse effects, there are cell-intrinsic mechanisms such as autophagic processes to limit oxidative damage, epigenetic reprogramming, and metabolic changes that contribute to the maintenance of stemness (Ibneeva et al., 2024; Liu et al., 2020; Ma et al., 2020; Revuelta and Matheu, 2017; Singh et al., 2018; Zhao and Deininger, 2023). Without these protective mechanisms, HSCs accumulate cellular damage and lose their ability to self-renew and differentiate, a process termed stem cell exhaustion. While inflammation is linked to HSC exhaustion, less is known about the cell extrinsic factors that contribute to the

maintenance of HSC quiescence and integrity, but several cytokines have been implicated in this

process. For example, in response to repeated stimulation with poly I:C, IFN-α can remove actively

proliferating HSCs from the developmental pool as a mechanism to prevent the accumulation of

damaged HSCs (Pietras et al., 2014). In addition, during fetal development there is evidence that

IL-10 limits HSC responsiveness to inflammatory signaling and thereby limit emergency

myelopoiesis as a mechanism to preserve HSC functionality (Collins et al., 2024).

IL-27 is a member of the Type I family of cytokines that utilize JAK-STAT signaling to

mediate their biological effects. This family also includes IL-6, and both of these cytokines signal

through dimeric receptors that contain the gp130 receptor combined with unique IL-6Rα and IL-

27Rα components. Because T cells constitutively express high levels of the IL-27R (Pflanz et al.,

2004; Pflanz et al., 2002), the majority of studies on this cytokine have focused on its impact on

these populations. Thus, IL-27 can limit the development of pathological Th1, Th2, and Th17

CD4[+] T cell responses (Findlay et al., 2010; Hamano et al., 2003; Villarino et al., 2003), but several

studies have also indicated that the absence IL-27 results in an elevated population of inflammatory

monocytes that contribute to the development of immune pathology (Aldridge et al., 2024; Liu et

al., 2021). The observation that the loss of IL-27 also results in enhanced inflammatory monocyte

induction prior to a detectable T cell response suggests that IL-27 may antagonize the innate

events that promote these monocyte populations (Aldridge et al., 2024). Consistent with this

idea, there are additional reports that IL-27 can limit neutrophil and monocyte responses

and directly impact myelopoiesis (Furusawa et al., 2016; Liu et al., 2014; Liu et al., 2021; Seita

et al., 2008; Sun et al., 2017; Wirtz et al., 2006). However, the observation that monocytes do not

express the IL-27R suggests that the ability of IL-27 to modulate this population is indirect.

The studies presented here show that infection with *T. gondii* results in increased HPSC

activity associated with emergency myelopoiesis; however, in the absence of IL-27, HSPCs

display a skewed developmental program that resulted in enhanced monopoiesis. While HSPCs

showed preferential expression of the IL-27R, the loss of IL-27 during infection resulted in reduced

HSPC fitness. Thus, during infection, the ability of IL-27 to act on HSPC provides a regulatory

brake on differentiation to limit monocyte induction and preserve HSPC stemness.

Materials and Methods

Mice and infection

Female and male C57BL/6 mice were purchased (8-12 weeks old; strain # 000664) from

Jackson labs as age and sex matched controls for whole body knockout mice. CD45.1 donor mice

were either bred in house or purchased from Jackson labs (10-12 weeks old; strain # 033076).

CD45.2 recipient mice were purchased from Taconic labs (10-12 weeks old; model # B6-M). IL-

$27R^{-/-}$ (Jackson labs; strain # 018078) were bred in house, as were IL-27p28$^{-/-}$ mice, originally

generated by Lexicon Pharmaceuticals, Inc. and previously described (Hall et al., 2012). IL-27-

p28-GFP reporters (Kilgore et al., 2018) were similarly bred in house, and the Procr-Ai6 was

generated from crossing the Procr-ERT2-cre line (Procr: Jacson labs; strain # 033052) which had

been backcrossed 9 generations onto the C57BL/6 background and was provided by Dr. Nancy

Speck to the Ai6 mouse line (Jackson labs; strain# 007906). Mice that contain a floxed allele of

the IL-27Rα (IL-27R$^{fl/fl}$) were provided by Dr. Booki Min (Do et al., 2017), and maintained within

our mouse colony. This line was then crossed with a CD4-cre mouse (Jackson labs; strain# 022071)

to generate the CD4^{cre}-IL-27R$^{flox/flox}$ mouse line. M1Red mice, which express red fluorescent

protein (RFP, encoded by *dsRed2*) under control of the *Irgm1* promoter, were received from

Gregory Taylor (Duke University) and maintained in-house. All mice were housed and bred in specific pathogen-free (SPF) facilities in the Department of Pathobiology at the University of Pennsylvania in accordance with institutional guidelines (IACUC# 805045). Cysts of ME49 strain of *T. gondii* were collected from chronically infected CBA mice brain tissues. Experimental mice were then infected i.p. with 20 cysts. For experiments where uninfected bone marrow was sorted, a Prugniaud (PRU) strain of *T. gondii* expressing tdTomato was used (Pru-tdTomato). Both of these are type II strains and show similar virulence.

Functional Analysis of HSPC *in vivo*

For induction of the Procr-ERT2-cre, tamoxifen was given for three consecutive days (5 mg/mouse) prior to infection. For BRDU proliferation labeling, 5 mg of BRDU was given i.p. to each mouse the same day as infection. 1mg/ml of BRDU was then provided in the drinking water to mice throughout infection. To generate bone marrow chimeras, recipient mice were given a radiation dose of 900-1000 rads at least 6 hrs. prior to transferring donor marrow. Donor marrow was transferred (at least $2x10^6$ cells from each donor, unless otherwise stated; this resulted in a total of $4x10^6$ cells being transferred for mixed chimeras) intravenously (i.v.) via retro-orbital injection. Mice were then given 5 mL of Sulfamethoxazole/ Trimethoprim (200mg+40mg/5ml) in their drinking water and replaced every 2-3 days for two weeks.

Cell staining, flow cytometry, UMAP analysis, and detection of IL-27

Splenocytes were obtained by grinding spleens through a 70 μm filter and washing with RPMI supplemented with 5% fetal bovine serum (FBS). Red blood cells were then lysed by incubating samples with ACK lysis buffer (Thermo Fischer Scientific) for 5 minutes before

washing again with RPMI + 5% FBS. Both femurs and tibias were harvested from mice for bone

marrow isolation. Connective tissue was removed from the bones by gentle scraping with a scalpel

before one end of each bone was cut off to expose the marrow. Bones were then placed in a 0.5 ml

microcentrifuge tube, with a hole punctured in the bottom with a 31g needle. The 0.5ml tube was

then placed within a 1.5 ml microcentrifuge tube; the nested tubes were then spun down at 13, 000

rcf for 1.5 min. The pelleted cells were then ACK lysed and washed as above (centrifuging at 500

rcf) before being passed through a 70 μm strainer and resuspended in appropriate media (adapted

from (Au - Amend et al., 2016)). For peritoneal exudate cells (PECs), 8 mL of PBS was injected

into the peritoneum of mice (21G needle attached to a 10 mL syringe) and re-aspirated using the

same needle and syringe. Syringe contents were then expelled into a 15 mL conical tube, spun

down at 300 rcf, and resuspended in appropriate media.

After cell numbers were determined for each sample, equivalent numbers of cells were

plated in a 96-well plate and spun down at 300 rcf. Cells were then washed with FACS Buffer [1x

PBS, 0.2% bovine serum antigen, 1mM EDTA] before incubating with Fc block [99.5 % FACS

Buffer, 0.5% normal rat serum, 1 μg/ml 2.4G2 IgG antibody] prior to staining. If CD16/32 was

stained for, the staining antibody was substituted for unconjugated 2.4G2 antibody in the Fc block.

Cells were stained with the viability dye Ghost Dye Violet 510 (Tonbo biosciences; 12-0870), and

the following antibodies were used for subsequent staining: CD90.2 (BD; clone 30-H12), B220

(BD; clone RA3-6B2), Ly6G (BD; clone 1A8), CD11b (BD; clone M1/70), Ly6C (Biolegend;

clone HK1.4), Sca-1 (BD/Invitrogen; clone D7), CD117/cKit (Biolegend; clone ACK2), CD135

(Biolegend; clone A2F10), CD150 (Biolegend; clone TC15-12F12.2), CD48 (Biolegend/BD;

clone HM48-1), CD127 (Biolegend; clone A7R34), CD16/32 (BD; clone 2.4G2), CD34

(Invitrogen; clone RAM34), CD115 (Biolegend; clone AFS98), CCR2 (Biolegend; clone

SA203G11), CX3CR1 (Biolegend; clone SA011F11), CD11c (Biolegend; clone N418), XCR1

(Biolegend; clone ZET), CD172a (Biolegend; clone P84), CD64 (Biolegend; clone X54-5/7.1),

F4/80 (Invitrogen; clone BM8), MHCII (Biolegend; clone M5/114.15.2), gp130 (Biolegend; clone

4H1B35), IL-27Rα (BD; 2918), CD132 (BD; clone 4G3), CD131 (BD; clone JORO50), IL-6Rα

(Biolegend; clone D7715A7), GM-CSFRα (R&D systems; clone 698423), CD210/IL-10R

(Biolegend; clone 1B1.3a), IFN-γR β (Biolegend; clone MOB-47), IFNAR-1 (Biolegend; clone

MAR1-5A3), IL-21R (Biolegend; clone 4A9), CD120a/TNFRI (Biolegend; clone 55R-286),

CD120b/TNFRII (Biolegend; clone TR75-89), and CD217/IL-17RA (Biolegend; clone

S19010F).

Intracellular staining for Ki67 (BD; clone B56), and BCL2 (Biolegend; clone BCL/10C4)

was performed using the Foxp3/Transcription Factor Staining Buffer Set (eBioscience) according

to the manufacturer's instructions. BRDU staining was done following a previous published

protocol (Matatall, Methods Mol. Biol., 2018). Cells within the vasculature were labeled by

injecting 3 μg of florescent anti-CD45 antibody (Biolegend; clone 30-F11)/mouse for 3 min prior

to euthanasia. Samples were run on a FACSymphony A3 or A5 (BD) and analyzed using the

FlowJo Software analysis program (TreeStar). For UMAP analyses, either 500 or 10,000 live cells

(for 5 days of culture or 10, respectively) per sample were sampled using the downsampling plug-

in (TreeStar) and subsequently concatenated. The concatenated samples were then used for UMAP

analysis with the FlowJo plug-in (TreeStar), using Euclidean distances with the nearest neighbor

set to 15, minimum distance of 0.5, and number of components set to 2. All compensated

parameters were then used in the analysis. After the UMAP analysis was performed, the X-shift

plug-in (TreeStar) was used to further analyze each cluster for its defining features (Samusik et al.,

2016).

170   For IL-27p28 detection, mouse IL-27p28/IL-30 Quantikine ELISA kit (M2728; R&D

171 Systems) was used according to the manufacturer's protocol.

173 **Immunofluorescence of bone marrow and image analysis**

174   Procr^{tdTomato}IL-27p28^{GFP} mice were generated, infected, and long bones harvested at 5 dpi.

175 The bones were then fixed in 4 % PFA for 2.5-4 hrs, washed with PBS 3x, and then decalcified in

176 0.5mM EDTA for ≥72 hrs before being paraffin embedded. 5 μm sections were taken and placed

177 onto charged microscope slides. For staining, slides were de-paraffinized, rehydrated, and placed

178 in 1mM EDTA (pH 8.0) overnight in a 55°C water bath to recover epitopes prior to staining. Slides

179 were washed in diH$_2$O the next day before proceeding with staining. Staining was adapted from

180 (Im et al., 2019). Slides were blocked with 100μl of blocking buffer (1x PBS+0.3% triton X-

181 100+1:200 rat IgG) for 30 min at room temperature. Slides were then washed 3x, for 5 min each

182 with PBS. Endogenous streptavidin and biotin were then blocked using the ReadyProbes

183 Stretavidin/Biotin Blocking solution (Invitrogen) according to the manufacturer's directions. After

184 washing, GFP was stained for at 1:40 with anti-GFP AF488 (Biolegend; FM264G) overnight at

185 4°C. Slides were washed 3x with PBS, as above, and then mounted and coverslipped using 15μl

186 Fluoro-Gel (Electron Microsopy Sciences). Slides were then imaged on a Stellaris Falcon (Leica)

187 microscope.

188   Images were processed and analyzed using the Fiji software package (Schindelin et al.,

189 2012). Images were converted to binary images, manually thresholded, watersheding applied to

190 separate overlapping objects, and positive cells counted using the "analyze particles" tool.

192 **Statistics**

An unpaired, Welch's t-test or one-way ANOVA with Sidak's correction for multiple

comparisons were used to test for significant differences between groups. These were used as

indicated in each figure and unless otherwise stated. P values of less that 0.05 were considered

significant.

Results

IL-27 regulates monopoiesis during toxoplasmosis

Infection with *T. gondii* leads to increased production of monocytes that is enhanced in the

absence of IL-27 (Aldridge et al., 2024), but it is unclear if this is a result of differentiation from

HSPCs or from the more differentiated monocyte progenitors (MPs). Therefore, the Procr-ERT2-

cre mouse line, which expresses Cre in HSPCs (Gur-Cohen et al., 2015; Iwasaki et al., 2010), was

crossed with the Ai6 reporter line to provide a lineage tracing model. In these mice, tdTomato is

expressed in *procr*[+] cells and, upon tamoxifen (TAM) treatment, HSPCs will be tdTomato[+] and

GFP[+] but their differentiated progeny will only express zsGreen (Fig. 1A). For these experiments,

Procr-Ai6 mice were treated with TAM, infected, and then at 5 dpi the BM and peripheral cells

were analyzed. Consistent with other HSPC lineage tracing models (Chapple et al., 2018; Säwen

et al., 2018), in naïve bone marrow only a small proportion (~5%) of long-term hematopoietic

stem cells (LTHSCs) were labeled which corresponded to a low proportion of zsGreen[+] monocytes

in the peritoneum and spleen (Fig. 1B-C). After infection, the proportion of zsGreen[+] LTHSCs in

the marrow nearly doubled (Fig. 1B), and the proportion of monocyte dendritic cell progenitors

(MDP) also increased (Supp Fig. 1A). This corresponded to an increase in labeled monocytes and

neutrophils in the bone marrow of infected mice (Supp. Fig. 1A). In the periphery, there was also

an increased proportion of zsGreen$^+$ monocytes in the spleen, and the proportion of labeled

monocytes in the peritoneum, the site of initial infection, reached 100% (Fig. 1C). These

progenitors and their progeny were defined based on the expression of defining surface markers

as measured by flow cytometry (Supp. Fig. 2). Additional phenotyping revealed that both classical

(CCR2^{hi}, CX3CR1^{lo}) and non-classical (CCR2^{lo}, CX3CR1^{hi}) monocyte population were labeled to

equivalent degrees, indicating that infection impacts the recruitment of both subsets (Supp. Fig

1B). These data sets indicate that the increased myelopoiesis that occurs in response to infection

with *T. gondii* is due to expansion and enhanced differentiation of HSPCs, and not just MPs, into

downstream progeny, eventually resulting in increased monocyte production.

To determine the impact of IL-27 on this infection induced myelopoiesis, WT and IL-

27p28$^{-/-}$ mice were challenged with *T. gondii* and numbers of progenitors (from LTHSCs to MPs)

were analyzed (Fig. 1D). While infection led to an expansion of LTHSCs, the numbers of LTHSC,

granulocyte progenitors (GP), and other HSPCs (including common lymphocyte progenitors

(CLPs), common myeloid progenitors (CMPs), and granulocyte monocyte progenitors (GMPs))

(data not shown) were not impacted by the absence of IL-27 (Fig. 1E). Infection also resulted in

a transient increase in MPs in WT mice, but in the absence of IL-27 this was enhanced at 3 and 5

dpi before returning to baseline at 10 dpi (Fig. 1F). A similar pattern was observed for the MDPs

(Supp. Fig. 1C), and in the IL-27p28$^{-/-}$ mice this corresponded to an increase in mature monocytes

in the liver (Supp. Fig. 1D). The CCR2^{hi}CX3CR1^{lo} phenotype of these cells indicated that these

were inflammatory monocytes, that were predominantly within the tissue parenchyma (Supp. Fig.

1E), suggesting that these monocytes are poised to participate in the tissue pathology associated

with these mice. Thus, one function of IL-27 during acute toxoplasmosis is to limit monopoiesis

and the generation of inflammatory monocytes.

IL-27 regulates monocyte development in a cell intrinsic manner

Previous studies have shown that during toxoplasmosis or trypanosomiasis the absence of IL-27 results in enhanced CD4$^+$ T cell responses which contribute to increased accumulation of inflammatory monocytes (Aldridge et al., 2024; Liu et al., 2021). To test if the early (5 dpi) enhanced monopoiesis in the IL-27$^{-/-}$ mice observed here was a secondary consequence of the CD4$^+$ T cell response, a CD4^{cre} IL-27R$^{fl/fl}$ (CD4-IL-27R) mouse line was generated. In these mice IL-27R expression was efficiently deleted in T cells, but other IL-27R expressing cells, such as B cells, maintained expression (Supp. Fig. 3A). When infected with *T. gondii* these mice phenocopied whole body IL-27R$^{-/-}$ mice and developed lethal immune pathology > 10 days post-infection (data not shown). When control and CD4-IL-27R mice were infected and analyzed at 5 dpi there were equivalent numbers of monocyte progenitors (Fig. 2A). Thus, the enhanced monopoiesis observed in the whole-body IL-27p28$^{-/-}$ mice is independent of the effect of IL-27 on CD4$^+$ T cells.

Next, a mixed bone marrow chimera approach was used to determine if the enhanced monopoiesis observed in the absence of IL-27 was cell intrinsic. In these experiments, WT and IL-27R$^{-/-}$ marrow from naïve mice was used to generate a series of individual and mixed bone marrow chimeras (Fig 2B). In the single-transfer chimeras, at fourteen weeks after reconstitution WT and IL-27R$^{-/-}$ marrow reconstituted the hematopoietic system of irradiated recipients with 100% of hematopoietic cells derived from the donor (Supp. Fig. 3B-C, left bars). However, in the mixed chimeras, IL27R$^{-/-}$ cells had a reduced engraftment of LTHSCs and generation of downstream MPs (Supp. Fig. 3B, right bars, a 40:60 ratio KO: WT). Despite this reduction in the IL-27R$^{-/-}$ LTHSC the population of CCR2^{hi}CX3CR1^{lo} monocytes in the periphery were preferentially derived from

the IL-27R$^{-/-}$ lineage (Supp. Fig. 3C, right bars, 60:40 ratio KO:WT). To determine if infection

would impact this differentiation, these mixed chimeras were infected with *T. gondii* and analyzed

at 5 dpi (Fig. 2C). After infection, IL-27R$^{-/-}$ LTHSC remained deficient in the mixed chimeras, but

MPs from both lineages were equivalent (Fig. 2C, right bars). Additionally, the CCR2^{hi}CX3CR1^{lo}

monocytes were still predominately derived from the IL-27R$^{-/-}$ lineage (Fig. 2D, right bars).

Together, these data indicate that at homeostasis and during infection IL-27 can support LTHSC

populations but also temper the downstream processes that lead to monocyte production.

Additionally, this function is independent of IL-27 signals on CD4$^+$ T cells, but rather is due to a

cell intrinsic ability of IL-27 to limit the production of inflammatory monocytes.

IL-27p28 is expressed in the bone marrow at homeostasis and during infection

Given the impact of IL-27 on myelopoiesis described above, studies were performed to

characterize the cellular sources of IL-27p28 and their spatial location in the bone marrow of naïve

and infected mice. WT mice were infected, bone marrow harvested at 0, 3, 5, and 10 dpi, and local

production of the IL-27p28 subunit assayed (Fig. 3A). At 0 dpi, low constitutive levels of IL-27p28

were detected and increased throughout infection, with a peak at 5 dpi and elevated levels

maintained at 10 dpi (Fig. 3A). The use of an IL-27p28-GFP reporter mouse (Aldridge et al., 2024;

Kilgore et al., 2018) revealed a small population of p28-GFP$^+$ cells in the bone marrow of naïve

mice, but by 5dpi this was expanded (Fig. 3B-C). Additional analysis of the bone marrow and

peripheral tissues (spleen, blood, PEC) (Fig. 3B-C) revealed that although a small proportion of T

cells were GFP$^+$, consistent with other reports (Kimura et al., 2016; Lin et al., 2023), monocytes

and macrophages were the dominant source of IL-27 in these sites, by proportion of GFP$^+$ cells as

well as by cell number (Fig. 3B-C, Supp. Fig. 4A). Additionally, monocyte populations expressed

the highest levels of IL-27p28-GFP in comparison to other populations, such as dendritic cells

(Supp. Fig. 4B).

To visualize the spatial relationship between IL-27 expressing cells and HSPCs, femurs

from naive IL-27p28-GFP mice and imaged. In naïve mice, a stain for Sca-1 revealed low numbers

of HSPCs while p28^{GFP+} cells were present at a higher frequency; however, few of the cells in each

population colocalized (Supp. Fig. 4C, yellow arrows). During infection the expression of Sca-1

is upregulated by many cell populations (Baldridge et al., 2011; Morales-Mantilla et al., 2022),

and so the Procr$^{tdTomato+}$ cre reporter mouse (identifying HSPCs) was crossed with the IL-27p28^{GFP}

reporter. As HSCs are typically found within the hypoxic regions of the bone marrow (Eliasson

and Jönsson, 2010; Parmar et al., 2007), i.v. injection of anti-CD31 was used to label arterioles to

distinguish regions of the marrow that are highly vascularized or more hypoxic.

Immunofluorescence of the femur from 5 dpi highlighted that, similar to flow cytometric analysis

of the BM of the IL-27p28 reporters, infection resulted in increased numbers of p28^{GFP+} cells (Fig.

3D). Procr^{tdtomato} had a homogenous distribution in the femur (Supp. Fig. 4D, middle) whereas

p28^{GFP+} expression was more regional (Fig 3D; Supp. Fig. 4D, top). This was quantified by

sampling throughout the femur section (Fig 3D; Supp. Fig. 4D, regions 1-8). The area containing

almost exclusively Procr$^{tdTomato+}$ HSPCs, but few p28^{GFP+} cells, was an area of low vascularization,

consistent with the HSC niche (Fig. 3D; Supp. Fig. 4D, regions 3-5). However, in regions close to

the CD31$^+$ vasculature, HSPCs and IL-27p28$^+$ cells were found in proximity to one another (Fig.

3D; Supp. Fig. 4D, regions 1-2 and 6-8). When these regions were grouped based on this

vascularization, and GFP expressing cells normalized to tdTomato$^+$ cells, a marked decrease in

GFP$^+$ cells was apparent in hypovascularized regions (Fig. 3D). These data indicate that during

infection the p28-GFP⁺ BM monocytes associate with HSPCs, and this is most apparent near the

vasculature and away from the HSC niche.

Developmental expression of the IL-27R by hematopoietic cells

The hematopoietic tree begins with undifferentiated LT-HSCs that differentiate into distinct

hematopoietic lineages and mature progeny; the color scheme in Fig. 4A reflects these

developmentally related populations. To determine which progenitor stages in the marrow were

responsive to IL-27, the expression of the IL-27Rα was compared with other cytokine receptors

(gp130, the IL-2R common gamma chain (CD132), IL-6Rα, and GM-CSFRα) that modulate

hematopoiesis (Collins et al., 2021; Reynaud et al., 2011). Expression of these receptors was

measured on HSPCs and committed progeny in naïve and infected mice (Fig. 4B-C; Fig. 5A-D).

The gp130 receptor, one chain of the IL-27 receptor that is also utilized by other IL-6 family

members, was expressed by all developmental stages examined in both naïve and infected mice

(Fig. 4B). In contrast, the IL-27Rα was highly expressed by LTHSCs, but markedly decreased

with lineage commitment, with mature neutrophils and monocytes showing no expression (Fig.

4C). The use of the Haemosphere RNA-seq data set (Choi et al., 2018) and the BloodSpot online

tool (Bagger et al., 2015) for the analysis of normal mouse hematopoietic bone marrow transcripts

(Chambers et al., 2007; Di Tullio et al., 2011) confirmed that levels of *IL27RA* mRNA were high

in LSKs and LTHSCs but decreased with lineage commitment (Supp. Fig. 5A-B). To determine

if IL-27R signaling was functional on HSPCs, bulk bone marrow was collected from M1Red

reporter mice (Pardy et al., 2024; Stifter et al., 2019), which express RFP under the control of the

Irgm1 promoter, downstream of STAT1 signaling. When these cells were treated with IL-27 (an

activator of STAT1) the myeloid committed progenitors showed minimal RFP expression (Supp.

Fig. 5C), whereas HSPCS showed a time-dependent increase in reporter expression (Fig. 4D).

When expression of the common IL-2R gamma chain (utilized by IL-2, -4, -7, -9, -15, and

-21), was analyzed, like gp130, it was ubiquitously expressed by all developmental stages,

although infection did upregulate its expression (Fig. 5A). Similarly, CD131, which is shared by

GM-CSF, IL-3, and IL-5 signaling, was expressed in most developmental stages and was markedly

enhanced upon infection (Fig. 5B). In contrast to the IL-27Rα, the expression of IL-6Rα and GM-

CSFRα by HSPCs was negligible, and the highest expression of these was detected on fully

committed, mature neutrophils and monocytes (Fig. 5C-D). Additionally, both the IL-6Rα and

GM-CSFRα showed enhanced expression during infection, further differing from IL-27Rα. A

summary of the survey for IL-10R, IFNγRβ, IFNαR, IL-17Rα, TNFIR, and TNFIIR is presented

in Supp. Fig. 6, and highlights that while some of these were constitutively expressed across

the hematopoietic tree, none of these receptors showed preferential expression by HSPCs.

Thus, this survey of cytokine receptor expression highlights that HSPCs but not downstream

progenitors are uniquely positioned to respond to IL-27.

Infection derived IL-27 restrains a monocytic differentiation program in HSPCs

To determine the impact of infection-driven IL-27 on HSPC differentiation, LTHSCs were

sorted from naïve or infected WT and IL-27R$^{-/-}$ mice at 5 dpi (see Supp. Fig. 7 for gating strategy).

These LTHSCs were then cultured in MethoCult and their phenotype and ability to form colonies

was quantified by morphology after 5 or 10 days of culture (Supp. Fig. 8). As expected, cells from

naïve mice generated all colony types, but in cultures derived from infected mice there were

reduced granulocyte/monocyte (GM) and granulocyte (G) colonies (Supp. Fig. 8A). However, by

10 days post-culture (dpc), there was a slight enhancement in macrophage/monocyte colonies (M)

from infected mice. The significant loss of colonies from infected mice, though, made

quantification of these colonies difficult. Thus, to better quantify these differences these cultures

were analyzed by flow cytometry to measure expression of canonical developmental markers (see

materials and methods). Dimensional reduction was then performed via UMAP, and cluster

formation was analyzed with X-shift to determine the dominant markers associated with individual

clusters. When cells were concatenated from all collection timepoints, 14 clusters were delineated

(Supp. Fig. 8B). These clusters where driven, predominantly, by either collection time (Supp. Fig.

8B, top) or proliferative status (Supp. Fig. 8B, bottom). Clusters 1-9 contained only samples from

5 dpc, and clusters 1-6 corresponded to undifferentiated, quiescent colonies that were minimally

proliferative ($Ki67^{lo}$) while clusters 7-9 were starting to proliferate ($Ki67^{hi}$) and differentiate to

give rise to mature cell populations. By 10 dpc, there was a loss of the quiescent colonies and a

rise of populations (clusters 10-14) that are $Ki67^{hi}$ but have increased expression of mature myeloid

markers (CD11b, CD16/32, and Ly6C) (Supp. Fig. 8C). This reflects the increased differentiation

of colonies into mature myeloid cells. The UMAP analysis highlighted clusters 1, 6, 7, and 14 as

the dominant output of this assay (Supp. Fig. 8D): Cluster 1-undifferentiated progenitors ($Ki67^{lo}$,

$BCL2^{mid}$, no other expression), Cluster 6-myeloid biased HSPCs ($CD150^{hi}$), Cluster 7- monocyte

progenitors (intermediate expression of myeloid markers ($CD16/32^{lo}$) , but expression of

progenitor markers ($CD34^{int}$)), and cluster 14-mature monocyte ($CD11b^{hi}$, $Ly6C^{hi}$, $CD16/32^{int}$)

(Supp. Fig. 8C). Furthermore, this clustering analysis highlighted that infection resulted in a loss

of myeloid progenitors (cluster 7) that was further reduced in the absence of IL-27 (Supp. Fig. 8E).

However, by 10 dpc this corresponded to enhanced differentiation into mature monocytes (cluster

14).

To compare the ability of HSPCs from WT or IL-27R$^{-/-}$ mice to generate monocyte

populations, cells solely from 10 dpc were clustered (Fig. 6A), with the dominant clusters being

1, 2, 3, 10, and 11 (Fig. 6B). Of these, HSPCs were present in cluster 1 (Sca-1$^+$cKit/CD117$^+$),

while cluster 10 contained monocyte progenitors (CD11b^{hi}Ly6C^{low}CD34$^+$CD115$^+$), and cluster 11

expressed markers of mature monocytes (CD11b^{hi}Ly6C$^+$) (Fig. 6C-D). Further analysis (Fig. 6E),

revealed that HSPCs from infected mice produced increased numbers of monocytes and

macrophages (cluster 11), but only HSPCs from infected, IL-27R$^{-/-}$ mice were enhanced in their

differentiation into monocyte progenitors (cluster 10). Thus, during infection the loss of IL-27

signaling enhances the skewing of HSPCs *ex vivo* toward monocyte differentiation.

In a complimentary *in vivo* approach, HSPCs from naïve or infected WT or IL-27R$^{-/-}$ mice

were transferred into irradiated WT hosts and differentiation assessed (Fig. 6F). For these

experiments bulk marrow was utilized because, as discussed above, inflammation results in

upregulation of Sca-1 and other lineage defining markers (Baldridge et al., 2011; Morales-Mantilla

et al., 2022), making it difficult to define true HSPCs for transfer by cell-sorting. In addition, for

these studies mice were infected with a strain of *T. gondii* that expresses tdTomato (Pru-tdTomato)

(Christian et al., 2014; John et al., 2009) to allow parasite removal during sorting and prevent

infection of irradiated recipients. Analysis of spleens of recipient mice at 12 weeks, revealed that

donors from naive or infected IL-27R$^{-/-}$ mice produced more monocytes than those derived from

WT mice (Fig. 6F). These *ex vivo* and *in vivo* approaches indicate that exposure of LTHSCs to IL-

27 during infection results in a stable program that limits monocyte production.

IL-27 regulates the functionality and fitness of HSPCs during infection

While the data sets presented above indicate a role for IL-27 in the regulation of HSPC

differentiation, it was unclear if loss of IL-27 signals during infection would affect their

functionality and fitness. To test if there were alteration in HSPC proliferation during infection,

WT and IL-27R$^{-/-}$ mice were treated with BRDU throughout infection and HSPC proliferation

analyzed at 5 dpi. The combination of Ki67 expression and BRDU incorporation was used to

distinguish HSPCs entering cell cycle (Ki67^{+}BRDU^{-}), cells that have entered cell cycle and are

actively proliferating (Ki67^{+}BRDU^{+}), and those that have proliferated but are no longer in cell

cycle (Ki67^{-}BRDU^{+}). In WT mice, the majority of LSKs are quiescent (Ki67^{-}BRDU^{-}), a proportion

of cells have entered cell cycle (Ki67^{+} BRDU^{-}), and a distinct Ki67^{+}BRDU^{+} population, as well as

a smaller Ki67^{-}BRDU^{+} population, can be detected. In the absence of IL-27 signaling, the most

prominent difference was a reduced proportion of Ki67^{+}BRDU^{+} cells (Fig. 7A). To determine if

this decreased Ki67^{+}BRDU^{+} population might be due to increased cell death of HSPCs in the IL-

27R$^{-/-}$ mice, cells were stained for Annexin V and a free-amine reactive viability dye to

discriminate early- and late-apoptotic cells as well as necrotic cells. No difference in cell death

was observed between WT and IL-27R$^{-/-}$ HSPCs (Supp. Fig. 9A). These data need to be interpreted

with care, as total HSPC numbers are not different between WT and IL-27$^{-/-}$ mice (see Fig 1), but

they suggest that IL-27 may regulate either the proliferation or retention of HSPCs within the bone

marrow niche.

A key feature of healthy HSCs is their ability to reconstitute the hematopoietic system of

irradiated mice, even after serial transfer (Siminovitch et al., 1964; Weissman, 2000). However,

inflammation and over-proliferation of HSCs can impair their ability to rescue lethally irradiated

recipients (Feng et al., 2008; King et al., 2011; Pietras et al., 2014). To test the impact of IL-27 on

HSC fitness and functionality, serial transplantations of HSCs from infected WT and IL-27R$^{-/-}$

mice were performed. In these experiments, marrow from naïve or infected donors was used to

produce single chimeras or to provide a 1:1 mix of WT and IL-27R$^{-/-}$ populations that were

transferred to irradiated recipients (Fig. 7B). All transfers promoted host survival and at 9 weeks

post-transplantation, the WT or IL-27R$^{-/-}$ donors from naïve and infected mice reconstituted the

blood compartment with a similar efficiency (Supp. Fig. 9B). However, in the 1:1 chimeras from

infected mice, the peripheral compartment was dominated by cells derived from WT mice.

Analysis of the marrow of the single transfer chimeras at 12 weeks post-transplant highlighted that

when donor cells were derived from infected mice there was a significant reduction in the LTHSC

compartment (Fig. 7C). To test the functionality of these HSCs, they were then transplanted for a

second time into irradiated hosts (Fig. 7D). Mice that received a secondary transplant of marrow

from naïve WT and IL-27R$^{-/-}$ mice survived irradiation, consistent with the ability of healthy HSCs

to undergo at least three rounds of transplantation (Siminovitch et al., 1964; Weissman, 2000).

Additionally, secondary transplantation of WT HSCs from infected mice also resulted in survival

of these hosts. However, the serial transfer of marrow from infected IL-27R$^{-/-}$ mice did not save

irradiated recipients (Fig. 7D). Thus, while IL-27R$^{-/-}$ HSPCs from infected mice maintain a

differentiation bias towards the monocyte lineage, they have a decreased survival and proliferation

capacity and a reduced ability to rescue irradiated mice.

Discussion

During toxoplasmosis, the increased production of inflammatory monocytes is critical to

control infection (Dunay et al., 2008), but these populations also have regulatory properties that

help to limit immune-pathology. For example, as infection progresses, the presence of IFN-γ in the

bone marrow promotes the production of monocytes that produce IL-10 and PGE$_2$ (Askenase et

al., 2015) which contribute to the restoration of homeostasis. Similarly, during toxoplasmosis and

other infections in response to IFN-γ/STAT1 signals (Furusawa et al., 2016 and our own

unpublished observations) monocytes become a dominant source of IL-27 (Detavernier et al.,

2019; Hall et al., 2012). While IL-27 has a critical role in directly limiting pathological T cell

responses, the data presented here show that IL-27 is also part of a regulatory loop that acts on

HSPCs to limit emergency myelopoiesis. Infection-induced monopoiesis gives rise to a population

of IL-27$^+$ monocytes in the bone marrow that can then act on HSPCs to limit further monocyte

production. This observation is reminiscent of reports that IFN-γ activated

monocytes/macrophages can directly limit HSPCs (McCabe et al., 2015; Seyfried et al., 2020) or

the ability of IL-10 to limit emergency myelopoiesis in the fetus (Collins et al., 2024).

One challenge that faces HSCs during inflammation is the need to protect from stressors

associated with proliferation that adversely affect the functions of these cells (Caiado et al., 2021).

The concept of HSC "exhaustion" has been developed to describe HSCs that are no longer able to

fulfill their role in proliferation and reconstitution of steady-state hematopoiesis (Singh et al., 2020;

Zhao and Deininger, 2023). Unanticipated findings from our studies here were the reduced ability

of IL-27R$^{-/-}$ HSCs post-infection to compete with WT HSCs, as well as the inability of these HSCs

to reconstitute upon secondary transfer, hallmarks of functional exhaustion (Singh et al., 2020).

These results infer that the ability of IL-27 to limit emergency myelopoiesis is also associated with

the ability of IL-27 to limit HSC exhaustion. Aberrant HSPC responses are associated with

conditions such as clonal hematopoiesis and aging, and there has been interest in the development

of strategies to target these HSPCs. Thus, selective targeting and depletion of "aged" HSPCs via

anti-CD150 treatments has been reported to restore balanced hematopoietic output and "youthful"

blood phenotypes (Ross et al., 2024). Whether IL-27 treatment can be used to preserve HSPCs, or

be blocked to enhance monocyte production, remains to be tested.

A common feature of certain models of innate training that affect hematopoiesis is that

cytokines, such as IFN-γ and IL-6, promote long term changes that polarize HSPCs toward

myeloid differentiation (Cheong et al., 2023; Kaufmann et al., 2018; Khan et al., 2020). Inherent

in this process is the idea that HSCs or downstream progeny would be sensitive to different

cytokines. The ability to survey the expression of cytokine receptors that influence hematopoiesis

emphasized that the shared cytokine receptor chains gp130 and IL-2Rγ are uniformly

expressed throughout hematopoietic development. In contrast, the alpha chain receptors for

individual cytokines (IL-6Rα, GM-CSFRα, IL-10R, IFNγRβ, IFNAR, IL-21R, TNFRI, and

TNFRII) displayed a much more restricted pattern of expression that correlated with key

features of the biology of individual cytokines. For example, the limited expression of the IL-

6 receptor on downstream progeny is consistent with reports of its high expression on MPPs

but not on HSCs (Mirantes et al., 2014; Reynaud et al., 2011). Nevertheless, that HSCs express

high levels of the IL-27R, which decreases with differentiation, emphasizes that these stem

cells are especially sensitive to the effects of IL-27. Whether IL-27 tempers some of the

processes associated with innate training or is itself associated with long-term epigenetic

changes is the subject of ongoing studies.

The use of murine models of toxoplasmosis have revealed many of the inhibitory

activities of IL-27, and the datasets presented here highlight the ability of IL-27 to influence

HSPCs to restrain monocyte/macrophage induction. There are also reports during other

infections and models that show IL-27 can regulate neutrophil and monocyte development

(Furusawa et al., 2016; Liu et al., 2014; Liu et al., 2021; Seita et al., 2008; Sun et al., 2017; Wirtz

et al., 2006). For example, during infection with an attenuated strain of *Plasmodium berghei,* IL-

27 was required for the induction of a neutrophil response, and *in vitro* IL-27 blunted the ability

of HSPCs to differentiate into the macrophage lineage (Furusawa et al., 2016). However, in less

inflammatory settings, such as tumor growth, the overexpression of IL-27 resulted in LSK

expansion and the induction of an M1 macrophage population (Zhu et al., 2023). Likewise, in a

model of angiotensin-mediated atherosclerosis, IL-27 was needed to overcome HSC quiescence

and increase differentiation and output of mature myeloid cells (Peshkova et al., 2019). While it

can be difficult to ascribe some of these effects to IL-27 acting directly on HSC, the contrasting

nature of these data sets suggest that the impact of IL-27 on HSPCs is context dependent. Indeed,

reports that TNFα can upregulate HSPC expression of the IL-27R (He et al., 2020) suggests that

IL-27 responsiveness can be regulated by the local environment. A better appreciation of these

environmental signals at homeostasis and during different types of inflammation, then, should help

to define these context-dependent activities of IL-27 on hematopoiesis.

Acknowledgments and funding

We would like to thank Dr. Nancy Speck for generously providing us with the Procr-cre

mouse line. Additionally, we thank Dr. Gordon Ruthel for his assistance and instruction in

capturing images of the mouse bone marrow. Schematic were created using BioRender (agreement

license BU27RCNPBH).

Funding for this work was provided by the NIH equipment grant (S10 OD032305-01A1),

NIH fellowship support (F31 AI 161962-03), and the Emmerson Foundation.

Figure legends

Figure 1. IL-27 regulates monopoiesis during infection. A) Schematic of infection and TAM dosing strategy for Procr-Ai6 mice. B) Procr-Ai6 mice were infected and representative flow plots of LTHSCs (Lin (CD3, NK1.1, B220, Ly6G)$^-$, Sca-1$^+$, CD117 (cKit)$^+$, CD135$^-$, CD48$^-$, CD150$^+$) from naïve (left) and infected (right) mice are shown. Proportions of zsGreen$^+$ LTHSCs were then quantified in both groups. C) Proportions of zsGreen$^+$ monocytes (CD3$^-$, B220$^-$, CD11b$^+$, Ly6C$^+$, Ly6G$^-$) in the spleen and PECs of naïve and infected mice. D) Schematic of infection of WT and IL-27p28$^{-/-}$ mice for E-F. E) Numbers of LTHSCs and granulocyte progenitors (GPs) (CD3$^-$, NK1.1$^-$, B220$^-$, CD117$^+$, CD34$^+$, CD16/32^{hi}, Ly6C$^+$, CD135$^-$, CD115$^-$) in the bone marrow of WT and IL-27p28$^{-/-}$ infected mice throughout infection. F) Monocyte progenitors (MPs) (CD3$^-$, NK1.1$^-$, B220$^-$, CD117$^+$, CD34$^+$, CD16/32^{hi}, Ly6C$^+$, CD135$^-$, CD115$^+$; orange box) in the bone marrow of infected WT and IL-27p28$^{-/-}$ mice throughout infection. Representative flow plots are shown (left) and quantified (right). Statistical significance was tested by one-way ANOVA with Sidak's correction. *, **, and *** correspond to p-values $\leq$ 0.05, 0.01, and 0.001, respectively. N=3-5 mice/group and data shown are representative of 2-3 experiments.

Supplemental Figure 1. IL-27 regulates monopoiesis during infection. A) The proportion of zsGreen$^+$ monocyte dendritic cell progenitors (MDPs) (CD3$^-$, NK1.1$^-$, B220$^-$, CD117$^+$, CD34$^+$, CD16/32^{lo}, CD115$^+$), monocytes, and neutrophils (CD3$^-$, B220$^-$, CD11b$^+$, Ly6C$^+$, Ly6G$^+$) in the BM of naïve and infected Procr-Ai6 mice. B) The proportion of zsGreen$^+$ CCR2$^{hi/lo}$ CX3CR1$^{hi/lo}$ monocytes in the spleen and PECs in naïve and infected Procr-Ai6 mice. C) Number of MDPs in WT and IL-27p28$^{-/-}$ mice in the bone marrow during infection. D) Numbers of monocytes in the liver in WT and IL-27p28$^{-/-}$ throughout infection. E) WT and IL-27p28$^{-/-}$ mice were injected I.V.

with fluorescent anti-CD45 to label immune cells in the vasculature. Numbers of I.V. label⁻ and

CCR2^{hi}CX3CR1^{lo} monocytes in the liver were then quantified throughout infection. Statistical

significance was tested by one-way ANOVA with Sidak's correction. *, **, and *** correspond to

p-values $\leq$ 0.05, 0.01, and 0.001, respectively. N=3-5 mice/group and data shown are

representative of 2-3 experiments.

Supplemental Figure 2. Gating strategy for flow cytometric analysis of cells in the bone

marrow and periphery. Representative flow plots from the bone marrow (top) and spleen

(bottom) are shown indicating the gating strategy used to identify various cell populations. The

combination of surface markers used are listed in the tables (bottom) and are color coded to

correspond to both the developmental schematic pictured as well as the corresponding flow gate

on the representative plots.

Figure 2. IL-27 regulates monopoiesis during infection in a cell intrinsic manner. A) CD4-IL-

27R mice were infected and MPs (orange box) measured at 5 dpi in the BM. Representative flow

plots are shown (left) and quantified (right). B) Schematic of BM chimeras generated from WT

(CD45.1) and IL-27R$^{-/-}$ (CD45.1.2) bulk bone marrow that were then infected and used in C-D. C)

The proportion of LTHSCs (left) and MPs (right) from each respective donor lineage in the bone

marrow of both the single (left two bars) and mixed (right two bars) chimeras at 5 dpi. D)

Representative flow plots pre-gated on splenic monocytes (left plot) and then down-gated on

CCR2^{hi}CX3CR1^{lo} (right plot) monocytes at 5 dpi. WT derived monocytes are shown in the solid-

dark gray while IL-27R$^{-/-}$ derived cells are in the dashed-light gray. The proportion of each donor

lineage that contribute to mature CCR2^{hi}CX3CR1^{lo} splenic monocytes at 5 dpi were then

quantified (far right). Statistical significance was tested by either Welch's T-test (A) or one-way

ANOVA with Sidak's correction (C-D). *, **, and *** correspond to p-values $\leq$ 0.05, 0.01, and

0.001, respectively. N=3-5 mice/group and data shown are representative of 2-3 experiments.

Supplemental Figure 3. IL-27 regulates monopoiesis during infection and post-irradiation in

a cell intrinsic manner. A) IL-27Rα expression on CD4$^+$ (CD19$^-$CD3$^+$CD8α^-) T cells, CD8$^+$

(CD19$^-$CD3$^+$CD4$^-$) T cells, and B cells (CD19$^+$CD3$^-$) in CD4-27R mice at 5 dpi. B) The proportion

of LTHSCs (left) and MPs (right) from each respective donor lineage in the bone marrow post-

reconstitution and pre-infection. C) The proportion of CCR2^{hi}CX3CR1^{lo} monocytes from each

respective donor lineage in the spleen and liver post-reconstitution and pre-infection. Statistical

significance was tested by one-way ANOVA with Sidak's correction. *, **, and *** correspond to

p-values $\leq$ 0.05, 0.01, and 0.001, respectively. N=3-5 mice/group and data shown are

representative of 2 repeated experiments.

Figure 3. IL-27p28 is produced in the BM during infection. A) WT mice were infected, and the

bone marrow harvested throughout infection. BM was either washed with media or cultured for

24 hrs. and supernatant collected. Both the wash and supernatant were analyzed by ELISA for IL-

27p28. B) IL-27p28-GFP reporter mice were infected and their bone marrow analyzed by flow

cytometry throughout infection. A representative flow plot at 5 dpi of bulk marrow is shown (left)

and proportion of immune cell contribution to the GFP$^+$ population measured (right). C) Numbers

of IL-27p28-GFP$^+$ cells in the spleen, blood, and BM of infected mice were quantified throughout

infection (left). Proportions of GFP$^+$ CD45$^+$ cells were analyzed by cell type in the blood, spleen,

and PEC during infection (left). D) Femurs from infected Procr-tdTomato$^+$p28-GFP$^+$ mice were

harvested, sectioned, and imaged at 40x magnification. The total imaged femur is shown (top),

with zoomed in sections shown below for increased detail of cellular localization (bottom).

Numbers of Procr-tdTomato$^+$ and p28-GFP$^+$ cells were counted in eight randomly selected regions

throughout the marrow (see Supp. Fig. 4). GFP$^+$ cells were normalized to tdTomato$^+$ cells and the

eight counted regions classified as areas of hyper- or hypo-vascularization according to proximity

of the region to the labeled vasculature. This allowed quantification of the localization of cells

(bottom right).

Supplemental Figure 4. IL-27p28 is produced in the bone marrow during infection. A) IL-

27p28-GFP mice were infected, and the numbers of each CD45$^+$, immune cell type that expressed

GFP in the BM or periphery quantified throughout infection. B) Levels of IL-27p28-GFP

expression were measured in bulk monocytes (red) and dendritic cells (blue) in the bone marrow

(left) and spleen (right). C) Femurs from naïve IL-27p28-GFP mice were stained for Sca-1 and

imaged at 40x magnification. Zoomed in images are shown (right) of the indicated areas, with the

yellow arrows indicating positively stained, Sca-1$^+$ cells. D) The merged image shown in Fig. 3D

is separated by channel (GFP, tdTomato, and AF647, from top to bottom). Eight randomly selected

regions that were used for quantification of either GFP$^+$ or tdTomato$^+$ cells are shown (top), and

the quantification of cells in each region is shown next to the corresponding florescence channel

(right).

Figure 4. Expression of the IL-27R subunits gp130 and IL-27Rα during hematopoietic

development. A) Schematic ball-and-stick model of hierarchical hematopoietic development. Cell

type colors are maintained for reference in the following panels. B) Progenitors and progeny shown

in (A) were analyzed in WT and infected mice (at 5 dpi) by flow cytometry for the gp130 receptor.

10,000 live cells were concatenated from n=3 mice, and representative flow plots are shown. C)

Cells in (B) were analyzed for expression of the IL-27Rα. D) Bulk bone marrow was isolated from

IRGM1-dsRed reporter mice. 22×10^6 cells were plated per condition and stimulated with 20 ng/ml

of IL-27 for 0, 6, or 48 hours and RFP expression measured by flow cytometry.

Supplemental Figure 5. IL-27Rα transcript expression during hematopoiesis and differential

sensitivity of LTHSC and CMP to IL-27. A) Expression levels of *IL27RA* are shown from the

Haemosphere RNA-seq data set and online tool as well as the Blood Spot data set and visualizer

(B). C) Representative flow plots of IRGM1-RFP expression in LTHSCs and CMPs 48 hrs. post-

stimulation with IL-27.

Figure 5. Developmental expression of hematopoietic cytokine receptors. A) Expression of

CD132 (IL-2Rγ) and B) CD131 (GM-CSF/IL-3/IL-5Rβ) was analyzed by flow cytometry on

progenitors and progeny in the BM of naïve and 5 dpi mice as in Fig. 4. C) Expression of IL-6Rα

and D) GM-CSFRα were measured as in (A).

Supplemental Figure 6. Developmental expression of additional cytokine receptors. Indicated

cytokine receptors were analyzed in naive and infected mice as described in Fig. 4-5.

Figure 6. IL-27 limits infection induced HSPC polarization. WT and IL-27R⁻/⁻ mice were

infected for 5 days, LTHSCs sorted, and cultured in MethoCult for 10-12 days before being

analyzed by flow cytometry. A) 10,000 live cells from n=3-4 mice/ group (WT (naïve+5 dpi) and

IL-27R$^{-/-}$ (5 dpi)) were concatenated and dimensionally reduced via UMAP. B) Numbers of cells

in each cluster from (A) were measured. C) X-shift analysis was used to identify the expression

level of each marker used in the clustering performed in (A). 5 of the dominant clusters from (B)

are shown. D) Clusters 1, 10, and 11, all involved in monocyte development, are highlighted. E)

Contribution of Clusters 1, 10, and 11 to each genotype are shown (left) and quantified (right). F)

Bone marrow chimeras were generated from infected WT and IL-27R$^{-/-}$ mice (left). The

contribution of donor cells to mature monocytes was then measured at 14 weeks post-transplant

(right). Statistical significance was tested by one-way ANOVA with Sidak's correction. *, ***, and

**** correspond to p-values $\leq$ 0.05, 0.001, and 0.0001, respectively. N=3-5 mice/group and data

shown are representative of 2-3 repeated experiments.

Supplemental Figure 7. Gating strategy for sorting of LTHSCs used in monocyte

differentiation. LTHSCs were sorted from WT naïve, WT 5 dpi, and IL-27R$^{-/-}$ 5 dpi mice. The

gating strategy is shown as well as the purity of sorted samples (last panel).

Supplemental Figure 8. Development and validation of an unbiased flow cytometry approach

for the analysis of MethoCult colonies. WT and IL-27R$^{-/-}$ mice were infected, LTHSCs sorted

and cultured in MethoCult for either 5 or 10 days. A) Colony phenotypes were analyzed by

microscopy and counted by eye. B) Colonies were collected at either 5 or 10 days post-culture

(dpc) and analyzed by flow cytometry. 500 live cells from n=3-4 mice/ group were concatenated

and dimensionally reduced via UMAP. UMAP plots are shown, and Ki67 expression is overlayed.

C) X-shift analysis was used to identify the expression level of each marker used in the clustering

performed in (B). D) Numbers of cells in each cluster from (B) were measured. E) Contribution of

each genotype to each cluster was quantified. Statistical significance was tested by one-way

ANOVA with Sidak's correction. *, ***, and **** correspond to p-values $\leq$ 0.05, 0.001, and

0.0001, respectively. N=3-4 mice/group and data shown are from 1 of 3 experiments.

Figure 7. Testing the impact of IL-27 on functionality and fitness of HSPCs during infection.

A) WT and IL-27R$^{-/-}$ were infected, treated with 5 mg BRDU, and kept on 1mg/ml BRDU in the

drinking water throughout infection. Incorporation of BRDU was then measured in Lin$^-$Sca-

1$^+$cKit$^+$ (LSK) progenitor cells in comparison to Ki67 at 5 dpi. Representative flow plots are shown

(left) and the proportion of BRDU$^+$Ki67$^+$ cells quantified (right). B) Schematic for the generation

of BM chimeras from infected mice used in (C). C) Numbers of LTHSCs from recipients receiving

donor marrow from either uninfected or infected mice were measured based on genotype. D)

Secondary transplant chimeras from mice that originally received naïve or infected marrow where

generated as shown (left). Survival of recipients was then assessed (right).

Supplemental Figure 9. Testing HSPC survival and fitness. A) LTHSCs from WT and IL-27R$^-$

$^{/-}$ BM at 5 dpi was stained for Annexin V (AV) expression and free-amine staining viability dye

(LD). This allowed the measurement of apoptotic (AV$^+$LD$^-$), late apoptotic (AV$^+$LD$^+$), and necrotic

(AV$^-$LD$^+$) cells. The proportion of LTHSCs in each of these stages was quantified. B) Monocyte

genotype was analyzed at 9 weeks post-transplant (wpt) from either naïve WT (CD45.1) or IL-

27R$^{-/-}$ (CD45.1.2) donors (top) or from infected donors (bottom), including a 1:1 mix of infected

WT and IL-27R$^{-/-}$ donors. Statistical significance was tested by Welch's t-test.

References

674 Aldridge, D.L., D. Moodley, J. Park, A.T. Phan, M. Rausch, K.F. White, Y. Ren, K. Golin, E. Radaelli, R.
675   Kedl, P.M. Holland, J. Hill, and C.A. Hunter. 2024. Endogenous IL-27 during toxoplasmosis
676   limits early monocyte responses and their inflammatory activation by pathological T cells.
677   *mBio* 15:e0008324.
678 Askenase, M.H., S.J. Han, A.L. Byrd, D. Morais da Fonseca, N. Bouladoux, C. Wilhelm, J.E. Konkel,
679   T.W. Hand, N. Lacerda-Queiroz, X.Z. Su, G. Trinchieri, J.R. Grainger, and Y. Belkaid. 2015.
680   Bone-Marrow-Resident NK Cells Prime Monocytes for Regulatory Function during
681   Infection. *Immunity* 42:1130-1142.
682 Au - Amend, S.R., K.C. Au - Valkenburg, and K.J. Au - Pienta. 2016. Murine Hind Limb Long Bone
683   Dissection and Bone Marrow Isolation. *JoVE* e53936.
684 Bagger, F.O., D. Sasivarevic, S.H. Sohi, L.G. Laursen, S. Pundhir, C.K. Sønderby, O. Winther, N. Rapin,
685   and B.T. Porse. 2015. BloodSpot: a database of gene expression profiles and transcriptional
686   programs for healthy and malignant haematopoiesis. *Nucleic Acids Research* 44:D917-
687   D924.
688 Baldridge, M.T., K.Y. King, N.C. Boles, D.C. Weksberg, and M.A. Goodell. 2010. Quiescent
689   haematopoietic stem cells are activated by IFN-gamma in response to chronic infection.
690   *Nature* 465:793-797.
691 Baldridge, M.T., K.Y. King, and M.A. Goodell. 2011. Inflammatory signals regulate hematopoietic
692   stem cells. *Trends Immunol* 32:57-65.
693 Caiado, F., E.M. Pietras, and M.G. Manz. 2021. Inflammation as a regulator of hematopoietic stem
694   cell function in disease, aging, and clonal selection. *Journal of Experimental Medicine* 218:
695 Chambers, S.M., N.C. Boles, K.Y. Lin, M.P. Tierney, T.V. Bowman, S.B. Bradfute, A.J. Chen, A.A.
696   Merchant, O. Sirin, D.C. Weksberg, M.G. Merchant, C.J. Fisk, C.A. Shaw, and M.A. Goodell.
697   2007. Hematopoietic fingerprints: an expression database of stem cells and their progeny.
698   *Cell Stem Cell* 1:578-591.
699 Chapple, R.H., Y.J. Tseng, T. Hu, A. Kitano, M. Takeichi, K.A. Hoegenauer, and D. Nakada. 2018.
700   Lineage tracing of murine adult hematopoietic stem cells reveals active contribution to
701   steady-state hematopoiesis. *Blood Adv* 2:1220-1228.
702 Cheong, J.G., A. Ravishankar, S. Sharma, C.N. Parkhurst, S.A. Grassmann, C.K. Wingert, P. Laurent,
703   S. Ma, L. Paddock, I.C. Miranda, E.O. Karakaslar, D. Nehar-Belaid, A. Thibodeau, M.J. Bale,
704   V.K. Kartha, J.K. Yee, M.Y. Mays, C. Jiang, A.W. Daman, A. Martinez de Paz, D. Ahimovic, V.
705   Ramos, A. Lercher, E. Nielsen, S. Alvarez-Mulett, L. Zheng, A. Earl, A. Yallowitz, L. Robbins,
706   E. LaFond, K.L. Weidman, S. Racine-Brzostek, H.S. Yang, D.R. Price, L. Leyre, A.F. Rendeiro,
707   H. Ravichandran, J. Kim, A.C. Borczuk, C.M. Rice, R.B. Jones, E.J. Schenck, R.J. Kaner, A.
708   Chadburn, Z. Zhao, V. Pascual, O. Elemento, R.E. Schwartz, J.D. Buenrostro, R.E. Niec, F.J.
709   Barrat, L. Lief, J.C. Sun, D. Ucar, and S.Z. Josefowicz. 2023. Epigenetic memory of
710   coronavirus infection in innate immune cells and their progenitors. *Cell* 186:3882-
711   3902.e3824.
712 Choi, J., T.M. Baldwin, M. Wong, J.E. Bolden, K.A. Fairfax, E.C. Lucas, R. Cole, C. Biben, C. Morgan,
713   K.A. Ramsay, A.P. Ng, M. Kauppi, L.M. Corcoran, W. Shi, N. Wilson, M.J. Wilson, W.S.

Alexander, D.J. Hilton, and C.A. de Graaf. 2018. Haemopedia RNA-seq: a database of gene
expression during haematopoiesis in mice and humans. *Nucleic Acids Research* 47:D780-
D785.

Christian, D.A., A.A. Koshy, M.A. Reuter, M.R. Betts, J.C. Boothroyd, and C.A. Hunter. 2014. Use of
transgenic parasites and host reporters to dissect events that promote interleukin-12
production during toxoplasmosis. *Infect Immun* 82:4056-4067.

Collins, A., C.A. Mitchell, and E. Passegué. 2021. Inflammatory signaling regulates hematopoietic
stem and progenitor cell development and homeostasis. *Journal of Experimental Medicine*
218:

Collins, A., J.W. Swann, M.A. Proven, C.M. Patel, C.A. Mitchell, M. Kasbekar, P.V. Dellorusso, and E.
Passegué. 2024. Maternal inflammation regulates fetal emergency myelopoiesis. *Cell*
187:1402-1421.e1421.

Detavernier, A., A. Azouz, H. Shehade, M. Splittgerber, L. Van Maele, M. Nguyen, S. Thomas, Y.
Achouri, D. Svec, E. Calonne, F. Fuks, G. Oldenhove, and S. Goriely. 2019. Monocytes
undergo multi-step differentiation in mice during oral infection by Toxoplasma gondii.
Commun Biol 2:472.

Di Tullio, A., T.P. Vu Manh, A. Schubert, G. Castellano, R. Månsson, and T. Graf. 2011.
CCAAT/enhancer binding protein alpha (C/EBP(alpha))-induced transdifferentiation of
pre-B cells into macrophages involves no overt retrodifferentiation. *Proc Natl Acad Sci U S*
A 108:17016-17021.

Do, J., D. Kim, S. Kim, A. Valentin-Torres, N. Dvorina, E. Jang, V. Nagarajavel, T.M. DeSilva, X. Li,
A.H. Ting, D.A.A. Vignali, S.A. Stohlman, W.M. Baldwin, 3rd, and B. Min. 2017. Treg-specific
IL-27Rα deletion uncovers a key role for IL-27 in Treg function to control autoimmunity.
Proc Natl Acad Sci U S A 114:10190-10195.

Dunay, I.R., R.A. Damatta, B. Fux, R. Presti, S. Greco, M. Colonna, and L.D. Sibley. 2008. Gr1(+)
inflammatory monocytes are required for mucosal resistance to the pathogen Toxoplasma
gondii. *Immunity* 29:306-317.

Eliasson, P., and J.I. Jönsson. 2010. The hematopoietic stem cell niche: low in oxygen but a nice
place to be. *J Cell Physiol* 222:17-22.

Feng, C.G., D.C. Weksberg, G.A. Taylor, A. Sher, and M.A. Goodell. 2008. The p47 GTPase Lrg-47
(Irgm1) Links Host Defense and Hematopoietic Stem Cell Proliferation. *Cell Stem Cell* 2:83-
89.

Findlay, E.G., R. Greig, J.S. Stumhofer, J.C. Hafalla, J.B. de Souza, C.J. Saris, C.A. Hunter, E.M. Riley,
and K.N. Couper. 2010. Essential role for IL-27 receptor signaling in prevention of Th1-
mediated immunopathology during malaria infection. *J Immunol* 185:2482-2492.

Furusawa, J., I. Mizoguchi, Y. Chiba, M. Hisada, F. Kobayashi, H. Yoshida, S. Nakae, A. Tsuchida, T.
Matsumoto, H. Ema, J. Mizuguchi, and T. Yoshimoto. 2016. Promotion of Expansion and
Differentiation of Hematopoietic Stem Cells by Interleukin-27 into Myeloid Progenitors to
Control Infection in Emergency Myelopoiesis. *PLoS Pathog* 12:e1005507.

Gur-Cohen, S., T. Itkin, S. Chakrabarty, C. Graf, O. Kollet, A. Ludin, K. Golan, A. Kalinkovich, G.
Ledergor, E. Wong, E. Niemeyer, Z. Porat, A. Erez, I. Sagi, C.T. Esmon, W. Ruf, and T. Lapidot.
2015. PAR1 signaling regulates the retention and recruitment of EPCR-expressing bone
marrow hematopoietic stem cells. *Nat Med* 21:1307-1317.

Hall, A.O., D.P. Beiting, C. Tato, B. John, G. Oldenhove, C.G. Lombana, G.H. Pritchard, J.S. Silver, N.
Bouladoux, J.S. Stumhofer, T.H. Harris, J. Grainger, E.D. Wojno, S. Wagage, D.S. Roos, P.
Scott, L.A. Turka, S. Cherry, S.L. Reiner, D. Cua, Y. Belkaid, M.M. Elloso, and C.A. Hunter.
2012. The cytokines interleukin 27 and interferon-γ promote distinct Treg cell populations
required to limit infection-induced pathology. *Immunity* 37:511-523.
Hamano, S., K. Himeno, Y. Miyazaki, K. Ishii, A. Yamanaka, A. Takeda, M. Zhang, H. Hisaeda, T.W.
Mak, A. Yoshimura, and H. Yoshida. 2003. WSX-1 Is Required for Resistance to
Trypanosoma cruzi Infection by Regulation of Proinflammatory Cytokine Production.
Immunity 19:657-667.
He, H., P. Xu, X. Zhang, M. Liao, Q. Dong, T. Cong, B. Tang, X. Yang, M. Ye, Y. Chang, W. Liu, X. Wang,
Z. Ju, and J. Wang. 2020. Aging-induced IL27Ra signaling impairs hematopoietic stem cells.
Blood 136:183-198.
Ibneeva, L., S.P. Singh, A. Sinha, S.E. Eski, R. Wehner, L. Rupp, I. Kovtun, J.A. Pérez-Valencia, A.
Gerbaulet, S. Reinhardt, M. Wobus, M. von Bonin, J. Sancho, F. Lund, A. Dahl, M. Schmitz,
M. Bornhäuser, T. Chavakis, B. Wielockx, and T. Grinenko. 2024. CD38 promotes
hematopoietic stem cell dormancy. *PLoS Biol* 22:e3002517.
Im, K., S. Mareninov, M.F.P. Diaz, and W.H. Yong. 2019. An Introduction to Performing
Immunofluorescence Staining. *Methods Mol Biol* 1897:299-311.
Iwasaki, H., F. Arai, Y. Kubota, M. Dahl, and T. Suda. 2010. Endothelial protein C receptor-
expressing hematopoietic stem cells reside in the perisinusoidal niche in fetal liver. *Blood*
116:544-553.
John, B., T.H. Harris, E.D. Tait, E.H. Wilson, B. Gregg, L.G. Ng, P. Mrass, D.S. Roos, F. Dzierszinski,
W. Weninger, and C.A. Hunter. 2009. Dynamic Imaging of CD8(+) T cells and dendritic cells
during infection with Toxoplasma gondii. *PLoS Pathog* 5:e1000505.
Kaufmann, E., J. Sanz, J.L. Dunn, N. Khan, L.E. Mendonça, A. Pacis, F. Tzelepis, E. Pernet, A.
Dumaine, J.C. Grenier, F. Mailhot-Léonard, E. Ahmed, J. Belle, R. Besla, B. Mazer, I.L. King,
A. Nijnik, C.S. Robbins, L.B. Barreiro, and M. Divangahi. 2018. BCG Educates Hematopoietic
Stem Cells to Generate Protective Innate Immunity against Tuberculosis. *Cell* 172:176-
190.e119.
Khan, N., J. Downey, J. Sanz, E. Kaufmann, B. Blankenhaus, A. Pacis, E. Pernet, E. Ahmed, S.
Cardoso, A. Nijnik, B. Mazer, C. Sassetti, M.A. Behr, M.P. Soares, L.B. Barreiro, and M.
Divangahi. 2020. M. tuberculosis Reprograms Hematopoietic Stem Cells to Limit
Myelopoiesis and Impair Trained Immunity. *Cell* 183:752-770.e722.
Kilgore, A.M., S. Welsh, E.E. Cheney, A. Chitrakar, T.J. Blain, B.J. Kedl, C.A. Hunter, N.D. Pennock,
and R.M. Kedl. 2018. IL-27p28 Production by XCR1(+) Dendritic Cells and Monocytes
Effectively Predicts Adjuvant-Elicited CD8(+) T Cell Responses. *Immunohorizons* 2:1-11.
Kimura, D., M. Miyakoda, K. Kimura, K. Honma, H. Hara, H. Yoshida, and K. Yui. 2016. Interleukin-
27-Producing CD4(+) T Cells Regulate Protective Immunity during Malaria Parasite
Infection. *Immunity* 44:672-682.
King, K.Y., M.T. Baldridge, D.C. Weksberg, S.M. Chambers, G.L. Lukov, S. Wu, N.C. Boles, S.Y. Jung,
J. Qin, D. Liu, Z. Songyang, N.T. Eissa, G.A. Taylor, and M.A. Goodell. 2011. Irgm1 protects
hematopoietic stem cells by negative regulation of IFN signaling. *Blood* 118:1525-1533.
Lin, C.-H., C.-J. Wu, S. Cho, R. Patkar, W.J. Huth, L.-L. Lin, M.-C. Chen, E. Israelsson, J. Betts, M.
Niedzielska, S.A. Patel, H.G. Duong, R.R. Gerner, C.-Y. Hsu, M. Catley, R.A. Maciewicz, H.

Chu, M. Raffatellu, J.T. Chang, and L.-F. Lu. 2023. Selective IL-27 production by intestinal regulatory T cells permits gut-specific regulation of TH17 cell immunity. *Nature Immunology* 24:2108-2120.

Link, D.C. 2000. Mechanisms of granulocyte colony-stimulating factor-induced hematopoietic progenitor-cell mobilization. *Semin Hematol* 37:25-32.

Liu, F.D.M., E.E. Kenngott, M.F. Schröter, A. Kühl, S. Jennrich, R. Watzlawick, U. Hoffmann, T. Wolff, S. Norley, A. Scheffold, J.S. Stumhofer, C.J.M. Saris, J.M. Schwab, C.A. Hunter, G.F. Debes, and A. Hamann. 2014. Timed Action of IL-27 Protects from Immunopathology while Preserving Defense in Influenza. *PLOS Pathogens* 10:e1004110.

Liu, G., O. Abas, Y. Fu, Y. Chen, A.B. Strickland, D. Sun, and M. Shi. 2021. IL-27 Negatively Regulates Tip-DC Development during Infection. *mBio* 12:

Liu, Y., Y. Chen, X. Deng, and J. Zhou. 2020. ATF3 Prevents Stress-Induced Hematopoietic Stem Cell Exhaustion. *Front Cell Dev Biol* 8:585771.

Ma, Z., J. Xu, L. Wu, J. Wang, Q. Lin, F.A. Chowdhury, M.H.H. Mazumder, G. Hu, X. Li, and W. Du. 2020. Hes1 deficiency causes hematopoietic stem cell exhaustion. *Stem Cells* 38:756-768.

MacNamara, K.C., M. Jones, O. Martin, and G.M. Winslow. 2011a. Transient activation of hematopoietic stem and progenitor cells by IFNγ during acute bacterial infection. *PLoS One* 6:e28669.

MacNamara, K.C., K. Oduro, O. Martin, D.D. Jones, M. McLaughlin, K. Choi, D.L. Borjesson, and G.M. Winslow. 2011b. Infection-induced myelopoiesis during intracellular bacterial infection is critically dependent upon IFN-γ signaling. *J Immunol* 186:1032-1043.

Maeda, K., Y. Baba, Y. Nagai, K. Miyazaki, A. Malykhin, K. Nakamura, P.W. Kincade, N. Sakaguchi, and K.M. Coggeshall. 2005. IL-6 blocks a discrete early step in lymphopoiesis. *Blood* 106:879-885.

Matatall, K.A., C.C. Shen, G.A. Challen, and K.Y. King. 2014. Type II interferon promotes differentiation of myeloid-biased hematopoietic stem cells. *Stem Cells* 32:3023-3030.

McCabe, A., Y. Zhang, V. Thai, M. Jones, M.B. Jordan, and K.C. MacNamara. 2015. Macrophage-Lineage Cells Negatively Regulate the Hematopoietic Stem Cell Pool in Response to Interferon Gamma at Steady State and During Infection. *Stem Cells* 33:2294-2305.

Miner, X., Z. Shanshan, D. Fang, Z. Qingyun, W. Jinhong, W. Chenchen, Z. Caiying, Z. Sen, L. Bingqing, W. Peng, and E. Hideo. 2021. Granulocyte colony-stimulating factor directly acts on mouse lymphoid-biased but not myeloid-biased hematopoietic stem cells. *Haematologica* 106:1647-1658.

Mirantes, C., E. Passegué, and E.M. Pietras. 2014. Pro-inflammatory cytokines: emerging players regulating HSC function in normal and diseased hematopoiesis. *Exp Cell Res* 329:248-254.

Morales-Mantilla, D.E., B. Kain, D. Le, A.R. Flores, S. Paust, and K.Y. King. 2022. Hematopoietic stem and progenitor cells improve survival from sepsis by boosting immunomodulatory cells. *eLife* 11:e74561.

Ochando, J., W.J.M. Mulder, J.C. Madsen, M.G. Netea, and R. Duivenvoorden. 2023. Trained immunity — basic concepts and contributions to immunopathology. *Nature Reviews Nephrology* 19:23-37.

Pardy, R.D., K.A. Walzer, B.A. Wallbank, J.H. Byerly, K.M. O'Dea, I.S. Cohn, B.E. Haskins, J.L. Roncaioli, E.J. Smith, G.Y. Buenconsejo, B. Striepen, and C.A. Hunter. 2024. Analysis of

intestinal epithelial cell responses to Cryptosporidium highlights the temporal effects of
IFN-γ on parasite restriction. *PLOS Pathogens* 20:e1011820.

Parmar, K., P. Mauch, J.A. Vergilio, R. Sackstein, and J.D. Down. 2007. Distribution of
hematopoietic stem cells in the bone marrow according to regional hypoxia. *Proc Natl*
Acad Sci U S A 104:5431-5436.

Peshkova, I.O., T. Aghayev, A.R. Fatkhullina, P. Makhov, E.K. Titerina, S. Eguchi, Y.F. Tan, A.V.
Kossenkov, M.V. Khoreva, L.V. Gankovskaya, S.M. Sykes, and E.K. Koltsova. 2019. IL-27
receptor-regulated stress myelopoiesis drives abdominal aortic aneurysm development.
Nature Communications 10:5046.

Pflanz, S., L. Hibbert, J. Mattson, R. Rosales, E. Vaisberg, J.F. Bazan, J.H. Phillips, T.K. McClanahan,
R. de Waal Malefyt, and R.A. Kastelein. 2004. WSX-1 and Glycoprotein 130 Constitute a
Signal-Transducing Receptor for IL-27. *The Journal of Immunology* 172:2225-2231.

Pflanz, S., J.C. Timans, J. Cheung, R. Rosales, H. Kanzler, J. Gilbert, L. Hibbert, T. Churakova, M.
Travis, E. Vaisberg, W.M. Blumenschein, J.D. Mattson, J.L. Wagner, W. To, S. Zurawski, T.K.
McClanahan, D.M. Gorman, J.F. Bazan, R. de Waal Malefyt, D. Rennick, and R.A. Kastelein.
2002. IL-27, a Heterodimeric Cytokine Composed of EBI3 and p28 Protein, Induces
Proliferation of Naive CD4+ T Cells. *Immunity* 16:779-790.

Pietras, E.M., R. Lakshminarasimhan, J.M. Techner, S. Fong, J. Flach, M. Binnewies, and E.
Passegué. 2014. Re-entry into quiescence protects hematopoietic stem cells from the
killing effect of chronic exposure to type I interferons. *J Exp Med* 211:245-262.

Revuelta, M., and A. Matheu. 2017. Autophagy in stem cell aging. *Aging Cell* 16:912-915.

Reynaud, D., E. Pietras, K. Barry-Holson, A. Mir, M. Binnewies, M. Jeanne, O. Sala-Torra, J.P. Radich,
and E. Passegué. 2011. IL-6 controls leukemic multipotent progenitor cell fate and
contributes to chronic myelogenous leukemia development. *Cancer Cell* 20:661-673.

Ross, J.B., L.M. Myers, J.J. Noh, M.M. Collins, A.B. Carmody, R.J. Messer, E. Dhuey, K.J. Hasenkrug,
and I.L. Weissman. 2024. Depleting myeloid-biased haematopoietic stem cells rejuvenates
aged immunity. *Nature* 628:162-170.

Samusik, N., Z. Good, M.H. Spitzer, K.L. Davis, and G.P. Nolan. 2016. Automated mapping of
phenotype space with single-cell data. *Nat Methods* 13:493-496.

Säwen, P., M. Eldeeb, E. Erlandsson, T.A. Kristiansen, C. Laterza, Z. Kokaia, G. Karlsson, J. Yuan, S.
Soneji, P.K. Mandal, D.J. Rossi, and D. Bryder. 2018. Murine HSCs contribute actively to
native hematopoiesis but with reduced differentiation capacity upon aging. *eLife*
7:e41258.

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C.
Rueden, S. Saalfeld, B. Schmid, J.-Y. Tinevez, D.J. White, V. Hartenstein, K. Eliceiri, P.
Tomancak, and A. Cardona. 2012. Fiji: an open-source platform for biological-image
analysis. *Nature Methods* 9:676-682.

Seita, J., M. Asakawa, J. Ooehara, S. Takayanagi, Y. Morita, N. Watanabe, K. Fujita, M. Kudo, J.
Mizuguchi, H. Ema, H. Nakauchi, and T. Yoshimoto. 2008. Interleukin-27 directly induces
differentiation in hematopoietic stem cells. *Blood* 111:1903-1912.

Seyfried, A.N., J.M. Maloney, and K.C. MacNamara. 2020. Macrophages Orchestrate
Hematopoietic Programs and Regulate HSC Function During Inflammatory Stress. *Front*
Immunol 11:1499.

Siminovitch, L., J.E. Till, and E.A. McCulloch. 1964. Decline in colony-forming ability of marrow
cells subjected to serial transplantation into irradiated mice. *Journal of Cellular and*
Comparative Physiology 64:23-31.
Singh, S., B. Jakubison, and J.R. Keller. 2020. Protection of hematopoietic stem cells from stress-
induced exhaustion and aging. *Curr Opin Hematol* 27:225-231.
Singh, S.K., S. Singh, S. Gadomski, L. Sun, A. Pfannenstein, V. Magidson, X. Chen, S. Kozlov, L.
Tessarollo, K.D. Klarmann, and J.R. Keller. 2018. Id1 Ablation Protects
Hematopoietic Stem Cells from Stress-Induced Exhaustion and Aging. *Cell Stem Cell*
23:252-265.e258.
Stifter, S.A., N. Bhattacharyya, A.J. Sawyer, T.A. Cootes, J. Stambas, S.E. Doyle, L. Feigenbaum, W.E.
Paul, W.J. Britton, A. Sher, and C.G. Feng. 2019. Visualizing the Selectivity and Dynamics of
Interferon Signaling In Vivo. *Cell Rep* 29:3539-3550.e3534.
Sun, D., M. Zhang, and M. Shi. 2017. IL-27 limits neutrophil mediated pathology during pulmonary
infection with Cryptococcus neoformans. *The Journal of Immunology* 198:123.128-
123.128.
Villarino, A., L. Hibbert, L. Lieberman, E. Wilson, T. Mak, H. Yoshida, R.A. Kastelein, C. Saris, and
C.A. Hunter. 2003. The IL-27R (WSX-1) is required to suppress T cell hyperactivity during
infection. *Immunity* 19:645-655.
Weissman, I.L. 2000. Stem Cells: Units of Development, Units of Regeneration, and Units in
Evolution. *Cell* 100:157-168.
Wirtz , S., I. Tubbe , P.R. Galle , H.J. Schild , M. Birkenbach , R.S. Blumberg , and M.F. Neurath 2006.
Protection from lethal septic peritonitis by neutralizing the biological function of
interleukin 27. *Journal of Experimental Medicine* 203:1875-1881.
Zhao, H.G., and M. Deininger. 2023. Always stressed but never exhausted: how stem cells in
myeloid neoplasms avoid extinction in inflammatory conditions. *Blood* 141:2797-2812.
Zhu, J., J. Yu, A. Hu, J.Q. Liu, X. Pan, G. Xin, W.E. Carson, Z. Li, Y. Yang, and X.F. Bai. 2023. IL-27 Gene
Therapy Induces Stat3-Mediated Expansion of CD11b+Gr1+ Myeloid Cells and Promotes
Accumulation of M1 Macrophages in the Tumor Microenvironment. *J Immunol* 211:895-
902.
916

Figure 1

Supplemental Figure 1

Supplemental Figure 2

Bone marrow gating strategy

Periphery gating strategy

Bone marrow surface markers

Cell Type	Defining Surface Markers
Long-Term Hematopoietic Stem Cell (LTHSC)	Lin (CD3, B220, NK1.1, Ly6G)$^-$, Sca-1$^+$, CD117$^+$, CD135$^-$, CD48$^-$, CD150$^+$
Short-Term Hematopoietic Stem Cell (STHSC)	Lin (CD3, B220, NK1.1, Ly6G)$^-$, Sca-1$^+$, CD117$^+$, CD135$^-$, CD48$^-$, CD150$^-$
Multi-Potent Progenitor 3 (MPP3)	Lin (CD3, B220, NK1.1, Ly6G)$^-$, Sca-1$^+$, CD117$^+$, CD135$^-$, CD48$^+$, CD150$^-$
Multi-Potent Progenitor 4 (MPP4)	Lin (CD3, B220, NK1.1, Ly6G)$^-$, Sca-1$^+$, CD117$^+$, CD135$^+$
Common Myeloid Progenitor (CMP)	(CD3, B220, NK1.1)$^-$, CD117$^+$, CD34$^+$, CD16/32^{lo}, CD115$^-$
Monocyte-Dendritic cell Progenitor (MDP)	(CD3, B220, NK1.1)$^-$, CD117$^+$, CD34$^+$, CD16/32^{lo}, CD115$^+$
Common Monocyte Progenitor (cMoP)/ Monocyte Progenitor (MP)	(CD3, B220, NK1.1)$^-$, CD117$^+$, CD34$^+$, CD16/32^{hi}, Ly6C$^+$, CD135$^-$, CD115$^+$
Granulocyte-Monocyte Progenitor (GMP)	(CD3, B220, NK1.1)$^-$, CD117$^+$, CD34$^+$, CD16/32^{hi}, Ly6C$^-$, CD135$^-$, CD115$^-$
Granulocyte Progenitor (GP)	(CD3, B220, NK1.1)$^-$, CD117$^+$, CD34$^+$, CD16/32^{hi}, Ly6C$^+$, CD135$^-$, CD115$^-$

Periphery surface markers

Cell Type	Defining Surface Markers
T cells	CD45$^+$, CD19$^-$, CD3$^+$, TCRβ$^+$
B cells	CD45$^+$, CD19$^+$, CD3$^-$, TCRβ$^-$
Neutrophils	CD45$^+$, CD19$^-$, CD3$^-$, TCRβ$^-$, CD11b$^+$, Ly6C$^+$, Ly6G$^+$
Monocytes	CD45$^+$, CD19$^-$, CD3$^-$, TCRβ$^-$, CD11b$^+$, Ly6C$^+$, Ly6G$^-$
Macrophages	CD45$^+$, CD19$^-$, CD3$^-$, TCRβ$^-$, Ly6C$^-$, Ly6G$^-$, CD64$^+$
Dendritic Cells	CD45$^+$, CD19$^-$, CD3$^-$, TCRβ$^-$, Ly6C$^-$, Ly6G$^-$, CD64$^-$, CD11c$^+$, MHCII$^+$
Classical Dendritic Cells 1	CD45$^+$, CD19$^-$, CD3$^-$, TCRβ$^-$, Ly6C$^-$, Ly6G$^-$, CD64$^-$, CD11c$^+$, MHCII$^+$, XCR1$^+$, CD172a$^-$
Classical Dendritic Cells 2	CD45$^+$, CD19$^-$, CD3$^-$, TCRβ$^-$, Ly6C$^-$, Ly6G$^-$, CD64$^-$, CD11c$^+$, MHCII$^+$, XCR1$^+$, CD172a$^-$

Figure 2

Supplemental Figure 3

Figure 3

Supplemental Figure 4

A.

BM # of GFP+ | Spleen # of GFP+ | Blood # of GFP+ | PEC # of GFP+

Days Post Infection

# of GFP+ cells

Legend:
- T cells
- B cells
- Neutrophils
- Monocytes
- Macrophages
- DC1
- DC2

B.

BM | Spleen

Normalized to Mode

Monocyte
- 0 dpi
- 5dpi

Dendritic Cell
- 0 dpi
- 5dpi

p28-GFP

C.

Sca-1
IL-27p28

100 µm | 20 µm | 20 µm

D.

IL-27p28-GFP

200 µm

Procr-tdTomato

200 µm

CD31-AF647

200 µm

GFP+ counts/ region

# of cells

tdTomato+ counts/ region

# of cells

Region

Figure 4

Supplemental Figure 5

A.

IL27RA expression-Haemosphere

IL27RA expression-BloodSpot

B.

C.

Figure 5

A.

CD132/IL-2Rγ

B.

CD131/GM-CSF/IL3/IL5Rβ

C.

IL-6Rα

D.

GM-CSFRα

Supplemental Figure 6

Figure 6

Supplemental Figure 7

Supplemental Figure 8

Figure 7

Supplemental Figure 9

A.

Apoptotic (AV⁺LD⁻)

Late Apoptotic (AV⁺LD⁺)

Necrotic (AV⁻LD⁺)

B.

Donor marrow: WT IL-27R⁻/⁻ 1:1