1 **Research Article**

2 **1 Title page**

Analysis of the interactions between physiological substances and endogenous hormones in *Bombax ceiba* bud differentiation

ShuyuWANG,Zeping Cai,Ying ZHAO, Deqiao CUI,   Meixing WANG,Yanting ZHANG and

XudongYU*

1. College of Tropical Agriculture and Forestry, Hainan University, Haikou, 51700, China;

E-mail:doeasete@163.com

Keywords *Bombax ceiba*; flower bud differentiation; physiological substances; endogenous hormones;

reciprocal networks

2 Abstract

In order to reveal the morpho-physiological changes of bud differentiation in *Bombax ceiba* , this

study took Haikou City, Hainan Province as the study area, and used paraffin sections, plant

physiological assays, and LC-MS, etc., and proposed to explore the changing rules during its bud

differentiation. The results showed that 5Ds, CK, Auxin mainly acted in phase I and phase II, and ABA,

SA, GA, ETH, JA mainly acted in phase II and phase III. A total of 20 DEMs were screened to be

enriched in the biosynthesis of secondary metabolites and other pathways, and there were synergistic

and antagonistic effects of physiological substances and hormones to regulate the flower bud

differentiation. This study provides a theoretical basis for investigating the mechanism of flower

formation in *Bombax ceiba* and production practice.

3 Summary statement

To investigate the morphological and physiological characteristics of the bud differentiation process

in *Bombax ceiba*, and to reveal the mechanisms of physiological substances and hormone interactions.

4 Introduction

Plant flowering mechanism is a complex and precise regulatory network, in which physiological

perception system and hormone regulation system play an indispensable core role (Wang Rong, 2007),

which accurately regulate the developmental process of floral organs by participating in the

spatiotemporal specific expression of flower development. Its quality directly affects plant growth and

development, and is of great significance in improving crop yield, breeding of good seeds, and regulation

of flowering time. Flower bud differentiation in plants is a process of transition from nutrient growth to

reproductive growth, which is the result of the synergistic action of internal and external factors. In recent

years, new hormone-regulated receptors have been discovered and identified, and the mechanisms of

various kinase cascades and secondary messenger systems have been gradually elucidated in the study

of signal transduction pathways, and the study of hormone signals regulating the expression of flowering genes has also been deepened (Yongqiang Yu. et al., 2023; Wang, J. et al., 2024).

Bombax ceiba is a large deciduous tree of the genus *Bombax* in the family Bombacaceae, which combines multiple uses such as ornamental, medicinal, spinning, timber, and landscaping, and it is a rare species of ecological forests, landscape forests, and economic forests (Wu, J.S. and Yu, X.D.,2017). Hainan Island is very rich in *Bombax ceiba* resources, according to Wang Jian et al. survey, Hainan *Bombax ceiba* in the whole island are distributed, and the most concentrated distribution for the Baisha, Changjiang two counties, including Changjiang known as China's 'Bombax ceiba hometown', the *Bombax ceiba* planting area of the province accounted for about 17.71% of the total.

As the natural flowering period of *Bombax ceiba* is mainly concentrated in winter and spring, the ornamental period is shorter, the cultivation time is longer, and less basic research, and few studies on the interrelationship between physiological substances and endogenous hormones on the regulation of bud differentiation, less research on the mechanism of *Bombax ceiba* flower greatly affects the promotion and application of *Bombax ceiba*, in order to reveal the characteristics of the *Bombax ceiba* flower bud differentiation, to understand the reproductive strategy and adapt to the environment, and to investigate the energy metabolism and substance accumulation of *Bombax ceiba*, this study was carried out in the field of *Bombax ceiba* planting. In order to reveal the characteristics of *Bombax ceiba* flower bud differentiation, to understand the reproduction strategy and adaptation to the environment, and to investigate the energy metabolism and material accumulation of *Bombax ceiba*, this study took Haikou City, Hainan Province as the study area, and relied on the *Bombax ceiba* germplasm resource nursery of Hainan University and the Ministry of Education Key Laboratory of Tropical Speciality Forest Trees and Flower Genetics and Germplasm Innovation, to investigate the energy metabolism and material accumulation of *Bombax ceiba* .We used paraffin sections, plant physiological assays, LC-MS and other methods to measure the morphological and physiological characteristics of *Bombax ceiba* during the bud differentiation process, and to explore 1) morphological and structural characteristics and physiological changes during the bud differentiation process, 2) identification of 88 endogenous hormones and hormone-related differential metabolites during the bud differentiation process of *Bombax ceiba*, and 3) investigation of hormone and physiological substance interactions in the network of the bud differentiation stage. To provide theoretical basis and reference materials for later research on the mechanism of flower formation in *Bombax ceiba* and production practice.

5 Materials and methods

The test site is located in Hainan University, Haikou City, Hainan Province (20°3′19.38″N, 110°19′30.1″E) in 2021, selected 30 with the same genetic background, good growth condition, the same growth, tree height of 5.8-7.5 m, crown spoke of 4.2-6.5 m cottonwood adult experimental standard 5 plants, collected annual branches or the current year on the new shoots as test materials, and every 3 d from the flower bud differentiation to the wilting process. The flower buds were collected from annual branches or new shoots of the year as test materials, and the trees were exposed to uniform light on all sides, and the whole process from flower bud differentiation to withering was observed and recorded every 3 d. The standard plants produced in the field were observed and recorded at the same time. One of the field production of standard plants in the cultivars used in the normal conditions should not reach the height and number of leaves on the beginning of the buds of the varieties known as 'early flowering' *Bombax ceiba*, generally earlier than normal flowering

81   *Bombax ceiba* 5 ~ 10d, and vice versa for the 'late-flowering' *Bombax ceiba*, normal conditions Normal

82   conditions for normal flowering *Bombax ceiba*.

1. Observation of morphological structure of bud differentiation

From 15 October 2020 to 30 March 2021, 7~10 fresh flower buds with full buds and good development

were taken every 3 d and put into self-sealing bags and marked for morphological observation.

Paraffin sectioning method Flower buds were quickly cut into 3 mm×3 mm slices with a double-sided razor

blade and put into FAA fixative (38% formaldehyde, glacial acetic acid, and 70% anhydrous ethanol: 1:1:18

ratio), and three slices of the cut flower buds were completely immersed in FAA fixative for at least 24 h. Flower

buds were dehydrated with different gradients of ethanol (30%, 50%, 70%, 80%, 90%, and 100%) →

xyleneoxide. 100%) → xylene transparency → dipping wax → embedding → slicing → patching → dewaxing →

rehydration → 1% Senka red and 1% solid green staining → washing → dehydration of one slice → drying and

a series of other operations, and then placed the slices in a light microscope to observe and take pictures.

Observation by in-body microscope 20 flower buds were randomly collected from the 5th node of the new

shoots every 3 d from the beginning of bud differentiation of cottonwood, and on the day of sampling, the

morphology and structure of the flower buds were observed under in-body dissecting microscope by peeling off

the scales of the flower buds with bare hands and observing them with in-body microscope, counting and

recording the stage of differentiation of the flower buds, and the lengths of the transverse and longitudinal

stems of the flower buds were measured by using vernier calipers. When more than 75% of the flower buds

were in one of the above stages of differentiation, it was determined that the sampling period had entered that

stage of differentiation.

2 Measurement of physiological substances

From 15 October 2021 onwards, 0.5 g of fresh early-flowering cotton buds were sampled at the early stage

of bud differentiation (I), the calyx/petal primordial differentiation stage (II), and the stamen/gynoecium

primordial differentiation stage (III), for the determination of the contents of physiological substances (soluble

sugars, sucrose, soluble starch, soluble proteins, malondialdehyde (MDA), chlorophyll a, chlorophyll b, and

carotenoids) of the early-flowering and late-flowering cotton buds respectively. carotenoids) content. Three

replicates were used in each group, in which soluble sugar, sucrose and soluble starch were determined by

anthrone colourimetric method, soluble protein was determined by Caulophyllum blue G-250 method,

malondialdehyde (MDA) content was determined by thiobarbituric acid method, and chlorophyll was

determined by 95% ethanol milling and filtration method, and the methods were referred to the 'Experimental

Guidelines of Plant Physiology'.

3 Determination of endogenous hormones and differential metabolites

From 15 October 2021 onwards, 24 *Bombax ceiba* adult experimental standard strain buds were selected

for treatment and determination of hormone levels. 0.5 g of *Bombax ceiba* buds were collected at each stage

and stored at ultra-low temperature and set up three replicates, which were ground with a grinder (30 Hz, 1 min)

until powdered and added to 10 µL of the internal standard mixing solution with a concentration of 100 ng/mL, 1

mL of methanol/water/formic acid (15:4:1, v/v :1, v/v/v) extractant were mixed well; vortexed for 10 min at 4 ℃,

centrifuged at 12000 r/min for 5 min, and the supernatant was taken to a new centrifuge tube for concentration;

after that, 100 µL of 80% methanol/water solution was reconstituted over 0.22 µm filter membrane for

LC-MS/MS analysis.

4 Data analysis

122 Excel software was used to categorise and organise the data, and SPSS 27.0 software was used to
123 analyse the data by ANOVA and Duncan's new complex polarity method for multiple comparisons, and Power
124 point plotting.
125
6 Results

1 Morphological characteristics of bud differentiation
Bombax ceiba bud differentiation is divided into the early stage of bud differentiation (I), the
calyx/petal primordium differentiation stage (II), and the stamen/pistil primordium differentiation stage
(III).The new growth of *Bombax ceiba* new shoots stops in late October and gradually lignifies, with the
apical growth point raised upwards, the tip of the apical tip thin, the base broadly bulging and conical, and
the cytoplasm of the cells is thick and strong, with a length of the flower bud of 0.30 cm, and the width of
the flower bud of 0.21 cm (Fig.1 A, Fig.1 E). In early November, the calyx basal differentiation stage (Ⅱ)
was entered, in which the growth cone was sunken and widened, and the top of the growth cone was
gradually flattened from oblate to oblate, and both sides of the growth point produced protuberances that
grew and expanded, and the base of the differentiation elongated and gradually became wider and
curved inward to form the petal primordium to become the calyx primordium, and as the growth cone
continued to be widened, the base of the calyx primordium was 0.65 cm in length and 0.58 cm in width
(Figs.1 B and 1 F).
In early December, we entered the stage of androgynophore/gynophore protocorm differentiation
(III), as the flower buds continued to differentiate and grow, the top of the growth cone began to appear
concave, and multiple protruding stamen protocorms gradually formed on the inner side of the petal
protocorms (Xu, Y. & Liu, J., 2024; Malakshahi Kurdestani, A & Francioli, D., 2024), and by December By
mid- to late December, a distinct stamen cluster is formed and a growth cone emerges in the middle of
the stamen cluster, with the new gynobasal protuberances elongating and growing rapidly to eventually
cover the entire growth cone, and the flower buds are ca. 1.37 cm in length and ca. 1.01 cm in width (Figs.
1 C and 1 G) (Xiaoxiao Li & Wan, Chuan et al., 2021).
After the development of the flower primordium was completed, it entered the maturation peri
od of the floral organ under the influence of hormones in the plant and the regulation of externa
l conditions, which was divided into five stages: the primordial stage, the fast-growing stage (3-5
d), the large balloon stage (1 d), the beginning of the flowering stage (1 d), and the full bloom
stage (2-4 d). During the growing period of about 120 d, the pedicel elongates slowly, the bud
elongates and enlarges slowly, and the bud becomes conical (Fig.2 A, Fig.2 B); during the fast
growing period of 3-5 d, the pedicel elongates rapidly, the bud splits open at the tip, the petals
are exposed and elongate rapidly (Fig.2 C); during the blooming period, the buds grow into a lo
ng conic (Fig.2 D); during the initiation of flowering, most of the petals are highly revolute after
they open, and a few of them stretch straight out and are thick in texture, exposing the stamen
s and the pistil, and the anthers open up! Pollen dispersal, stigma secretes mucus, bees and bir
ds collect nectar for pollination (Fig.2 E); flower buds elongate to longest and largest at balloonin
g stage, most petals gradually recycle inward at unopened bloom stage to form straight or includ
ed floral pattern, a few maintain original revolute floral pattern and gradually become thinner, wit
h margins gradually wilting (Fig.2 F) (Chuan Wan & Dongyan Yang,2022).

2 Changes in physiological substances of bud differentiation

Physiological substances are involved in a variety of physiological processes such as energy metabolism, osmotic regulation, signalling, etc., and are an important source of energy storage in plants, as well as an important regulator of plant growth and development and gene expression (Nazir, F &. Péter, P,2024). In order to investigate the effects of flowering time and bud differentiation stage on the physiological substances of *Bombax ceiba,*two-factor ANOVA and multiple comparisons were conducted on different different differentiation stages of early-flowering and late-flowering *Bombax ceiba* , and the results showed that the soluble sugar of early-flowering *Bombax ceiba* showed a tendency to rise and then fall, and then gradually declined after rising to the peak value of 11.42 mg•g-1 in stage II, and the content of soluble sugar of late-flowering *Bombax ceiba* showed a tendency to gradually rise, and gradually rose to the peak value of 11.42 mg•g-1.The soluble sugar content of late-flowering *Bombax ceiba* showed a rising trend, gradually increasing to a peak value of 9.50 mg-g-1. in stage III, with higher significance of *0.06* and *0.43* in stages II and III, respectively (Fig. 3A). Early flowering and late flowering *Bombax ceiba*   sucrose showed a decreasing and then increasing trend, the content first decreased to the lowest value of 0.80 mg•g-1. and 0.75 mg•g-1 in phase II and then gradually increased in early flowering *Bombax ceiba*   more than late flowering *Bombax ceiba*   content increased by 1.47 mg•g-1. and 0.78 mg•g-1. from phase II to phase III, respectively (Fig. 3A). -1. and 0.78 mg•g-1 respectively, which was most significant at *0.01* in phase I (Fig. 3 B). The soluble starch of early and late flowering *Bombax ceiba*   showed an increasing and then decreasing trend, reaching a peak of 53.14 mg•g-1. and 47.47 mg•g-1. at stage Ⅱ, and consuming a large amount of starch after pollen mother cell development was the lowest at 12.26 mg•g-1. and 9.72 mg•g-1. with significance of *0.01*, *0.01* and *0.03* at stages I to III, respectively (Fig. 3 C). The soluble protein content of early and late flowering *Bombax ceiba* showed a trend of decreasing and then increasing, early soluble proteins were involved in the construction of cell morphology shaping, division and differentiation to reach a minimum value of 125.08 mg•g-1,125.60 mg•g-1. at stage Ⅱ, and reached a peak value of 150.29 mg•g-1. and 174.33 mg•g-1. with a significant *0.01* at stage I (Fig.3 D). Malondialdehyde is an important physiological index for osmotic regulation when plants are subjected to adversity stress (Yang, D & Chen, Y, 2024; Chi, et al. 2023) The content of early-flowering and late-flowering *Bombax ceiba* showed a decreasing and then an increasing trend, gradually decreasing to 1.63 mg•g-1. in phase II, 1.68 mg•g-1. and then increased to a peak of 3.15 mg•g-1. and 2.76 mg•g-1. Early-flowering *Bombax ceiba* responded more significantly to the external temperature, which was more significant at *0.01* in phase III (Fig. 3 E)

Chlorophylls and carotenoids in the chloroplasts during flower bud differentiation are involved in metabolism and transport and are reference indicators of plant senescence (Huang & Y.C.,2024). Early-flowering *Bombax ceiba* chlorophyll a and late-flowering *Bombax ceiba* chlorophyll b showed a tendency to gradually decrease first, increasing to peak values of 0.81 mg•g-1. and 0.26 mg•g-1. in stage I, and then decreasing to the lowest values of 0.52 mg•g-1. and 0.18 mg•g-1. Late-flowering *Bombax ceiba* chlorophyll a and early-flowering *Bombax ceiba* chlorophyll b showed a trend of gradual increase with bud differentiation, and the content of late-flowering *Bombax ceiba* chlorophyll a peaked at 0.78 mg•g-1 at stage III, with an increase of about 16%. Early flowering *Bombax ceiba* chlorophyll b content to the Ⅰ period reached a peak of 0.26 mg•g-1 after a gradual decline to 0.18 mg•g-1., chlorophyll a value of 0.26 mg•g-1. after a significant both *0.25* and *0.04* (Fig.3 F,G). Carotenoid anthocyanin deposition, petal and fruit colouration, the content of early flowering *Bombax ceiba* gradually increased from phase I to phase III, respectively, by 0.04 mg•g-1. and 0.01 mg•g-1 to a peak value of 0.27 mg•g-1. in phase III, and the

content of late-flowering *Bombax ceiba* carotenoids gradually decreased to a minimum value of 0.19
mg•g-1. (Fig.3 H). Early flowering *Bombax ceiba* chlorophyll (a+b)/carotenoid and late flowering *Bombax*
ceiba chlorophyll a/b content gradually decreased to peak at 4.62   mg•g-1. and after 3.82 mg•g-1. in
stage I and then gradually decreased to to stage III to a minimum value of 3.02 mg•g-1. and 1.59 mg•g-1.
after. Chlorophyll (a+b)/carotenoid content of late flowering *Bombax ceiba* and early flowering *Bombax*
ceiba chlorophyll a/b content gradually increased and reached a peak value of 5.27 mg•g-1. and 5.21
mg•g-1. at stage III, and chlorophyll (a+b)/carotenoid was significant at all three stages (Fig. 3 I J).

3 Flower bud differentiation hormone levels and differential metabolites

Flowering consists of a complex flowering control network formed by multiple exogenous and
endogenous signals, and phytohormones, as the most important endogenous signalling, play an
important role in the process of flower formation (Zhu, Y. & Zhang, J, 2024; Huang, P.K. & Schmit, J,
2024). By determining the content of phytohormones at the stage of flower bud differentiation, 88
phytohormones existed in 8 major phytohormone categories were classified to be changed, and in
general GA (GA24), 5Ds, CK (2MeSiP, BAP, iP7G, iP9G, cZ9G, oT9G, mT9G, K9G), IAA (IAA-Ala, ICA,
ICAld, TRA, MEIAA, IAA-Asp) and predominantly act in phase I and II for floral primordium differentiation
(Fig. 4 C E F G H). ABA (ABA, ABA-GE), SA (SA, SAG), GA (GA3, GA19), JA, IAA (MEIAA, IAA-Asp,
IAA-Trp, IAN, IAA-Val-Me, IAA, TRP), CK (cZROG, 2MeScZR, DHZ7G, DHZROG, IP, IPR, DHZR, tZR,
2MeSiP), and JA (JA-ILE, MEJA, JA, JA-Val, H2JA) acted mainly in the senescence of floral organs in
phase II and phase III varied significantly in phases I and II (Fig. 4 A B C F G H). ETH (ACC) content
peaked at stage III (Fig. 4 D).
To further explore the hormone regulation-related differential metabolites during flower bud
differentiation in *Bombax ceiba* , a total of 20 differential metabolites (DEMs) were screened, of which 14
were up-regulated and 6 were down-regulated, with up-regulated expression focusing on Auxin, CK, GA,
and SA, and down-regulated expression focusing on Auxin, CK, JA, and 5Ds (Fig.5 A).KEGG is enriched
for environmental information processing and genetic information processing, biosynthesis of secondary
metabolites, metabolic pathways, phytohormone signalling on hormones and nutrients during bud
differentiation are closely related to the involvement of early biological processes as response to stimuli,
transmembrane translocation, cleavage enzyme activity and structural molecule activity. Zeatin
biosynthesis, synthesis of various alkaloids, and metabolic pathways are associated with delayed organ
senescence and storage and preservation of freshness (Fig. 5 B, C, and D). The process of bud
differentiation contained a total of 24 hormone-related DEMs, with a gradual decrease in overall content,
a decrease and then an increase in the content of 5Ds from stage I to stage II, and the consumption of a
large amount of SA for senescence retardation by stage III (Fig.5 E). The 20 hormone-related DEMs
screened were correlated, and Auxin (IAA) and CK (cZ) were strongly correlated, synergising with each
other to promote cell division and floral organ growth, and growth hormone polar transport. Whereas
Auxin (IAA-Val-Me) and CK (KR) antagonise each other, cytokinin promotes apical dominance and
facilitates apical bud growth, and cytokinin inhibits apical dominance and promotes flower bud
differentiation. (Fig. 5 F).

4 Physiological substances and endogenous hormone interaction networks

Physiological substances and endogenous hormones have mutual synergistic and resistant effects in
epigenetic regulation and flowering promotion pathways. They were divided into two groups according to the

degree of association, physiological substances MDA, sucrose, soluble protein and chlorophyll a were closely

related with an average radius of 2.737, and the rest of them were weakly related with an average radius of

15.596. Hormones IAA, CK, and 5Ds were closely related with an average radius of 2.854, and had the highest

degree of association with physiological substances Sucrose, MDA, soluble protein, and chlorophyll a,

whereas JA, SA had lower correlation (Fig 6-B).Mantel test analysis and clustering heat map showed

significant positive correlation between soluble sugars and G19 *(0.01<p<0.05)*, soluble proteins and IAA (IAA,

IAA-Ala, IAA-Trp, IAA-Val-Me) and CK (tZOG, K), GA (GA19) *(0.01<p<0.05)*, the intermediates produced by

sucrose in photosynthesis can be used as raw materials for the synthesis of phytohormones, which are

involved in the synthesis of Auxin (IAA, IAA-Ala, TRP, IAA-Trp), CK (KR, tZR, K, mT9G), GA19, and JA

(JA-ILE), and the synthesis of soluble starch, IAA (IAA, IAA-Val-Me), and GA (GA19). Ala, IAA-Val-Me,

IAA-Trp), CK (tZOG, KR, BAP9G, DHZR, mT9G, K), and GA (GA19) were significantly positively correlated

(0.01<p<0.05), and the hormones provided carbon skeleton and energy by regulating the physiological state of

the plant, which induced the plant to convert more photosynthesis products into sugars and transported them

to the bud site Accelerating the initiation of flower bud differentiation. Cytokinin showed significant positive

correlation *(0.01<p<0.05)* with sucrose, soluble starch and malondialdehyde, which is conducive to the

promotion of flower differentiation, and it is closely related to the accumulation of carbohydrates, cytokinin

affects the activity of key enzymes such as sucrose synthase by regulating the expression of related genes, so

that sucrose and soluble starch accumulate in the buds, and malondialdehyde content is increased to promote

resistance to adversity, and provide the provide energy and material basis for flower bud differentiation (Fig

6-A, C).

7 Discussion

The transition from nutrient growth to reproductive growth in plants is regulated through a complex

network of interactions between genetic factors and external environmental conditions. After sensing changes

in the external environment, plants adjust the bud differentiation process to adapt to the changing

environmental conditions through the transmission and integration of physiological substances and hormone

signalling networks. Exploring the morphological and physiological characteristics of flower bud differentiation

in Bombax Bombax ceiba is of great significance in revealing the growth and development of Bombax Bombax

ceiba, precisely regulating the flowering period in agricultural production, improving the yield and quality of the

crop, and cultivating rare varieties in the field of horticulture.

Bombax ceiba flower buds are mixed buds of monoecious and dioecious plants, and according to the time

of germination, they can be divided into winter and spring differentiation type, which is formed by the

differentiation of terminal buds through the induction of flower formation. The transition from nutrient growth to

reproductive growth in *Bombax ceiba* flower bud differentiation is divided into two stages, one is the

transformation from the growing point to the flower bud known as floral primordium development (from late

October to January of the following year, which lasts 60 d), and the other is the maturation of the floral organ

(from early January to March of the following year, which lasts 75 d). This is similar to the results of Hu

Shuangling and Zhu (Hu Shuangling and Lin Gongtian, 2024; Zhu and Zhang Fangqiu, 2021). Physiological

substances are essential energy sources for maintaining normal life activities of flowers (Lin, B., & Wang, D

,2023). The results show that the same plant is affected by the photoperiod and temperature, the plant

physiological rhythm of its own growth process there are differences, which in turn affects the accumulation of

physiological substances, the early soluble sugar and soluble starch content increased hydrolysis into directly

usable energy substances for the initiation of flower bud differentiation, the large amount of sugar in the leaves,

phloem, and buds to transport nutrients for the development of the pollen mother cells in the late stage, and

then with the warming of the temperature gradually slowed down the stress. Soluble proteins are involved in

the construction of cell membrane morphology, early flowering *Bombax ceiba* soluble protein content increases
through a large number of enzymes to catalyse a variety of biosynthetic reactions, such as nucleic acid
synthesis, hormone synthesis, etc., to ensure that the cells can carry out the metabolic activities smoothly,
laying the groundwork for the initiation of the buds and the development of the foundation. Malondialdehyde
can play a certain osmotic regulation of water in the plant body under adverse conditions. When the
temperature gradually close to 24 ℃ to reach the appropriate growth temperature of *Bombax ceiba*, with the
temperature rising osmotic stress continues to increase, the buds appeared in varying degrees of physiological
water shortage, prompting the buds continue to produce and accumulate malondialdehyde in the body. When
the temperature exceeded 28°C, the increase in external temperature slowed down, and the change in
membrane peroxides slowed down accordingly (Hiroyuki Tsuji, & Moeko Sato, 2024).

Chlorophyll provides energy and material basis for plants through photosynthesis. Previous studies have
rarely measured the trends of chlorophyll and carotenoids during bud differentiation, and further studies are
needed to investigate the regulatory mechanisms of chlorophyll and carotenoids in the bud differentiation
process of *Bombax ceiba*. Chlorophyll a absorbs and converts light energy to convert carbon dioxide and water
into organic matter, and chlorophyll b assists chlorophyll a in photosynthesis. It is worth noting that in this study,
the trends of changes in chlorophyll a and chlorophyll b in *Bombax ceiba* were found to be opposite at the
stage of bud differentiation, and it is speculated that the photosynthetic efficiency is improved by adjusting its
own metabolic pathway to reduce the input of chlorophyll synthesis during the physiological differentiation
stage. Carotenoids, as precursors of abscisic acid and solanum lactone, act synergistically with light signalling
to regulate bud differentiation. In past studies, epigenetic processes have been associated with carotenoid
biosynthesis, accumulation and degradation in plant growth and development, and a gradual decrease in
carotenoid content at the stage of bud differentiation has been speculated to be associated with a feedback
effect with other genes in the carotenoid biosynthesis pathway as a way of regulate carotenoid biosynthesis in
plants.

Plant hormones are the most important endogenous signalling players. Studies have shown that there are
synergistic and interacting resistance among hormones. By determining the changes in hormone content
during flower bud differentiation in cottonwood, it was found that ABA (ABA, ABA-GE), as a hormone that
inhibits growth and promotes dormancy and senescence, changed significantly during phase I and phase II.
Differential metabolites related to hormone regulation of bud differentiation were mainly concentrated in
metabolism and environmental information processing. *Bombax ceiba* regulated hormone concentration by
changing the expression of different kinds of endogenous hormone synthesis and metabolism genes during
bud differentiation, and phytohormones in turn regulated *Bombax ceiba* bud differentiation by causing the
expression of downstream genes through the signal transduction pathway. The up-regulated expression was
mainly concentrated in Auxin, CK, GA and SA, and the down-regulated expression was mainly concentrated in
Auxin, CK, JA and 5Ds. The roles of CK and Auxin were the most significant in the process of *Bombax ceiba*
bud differentiation, and the metabolic pathways involved in the early biological process as a response to the
stimulus, and transmembrane transporter, cleaving enzyme activity, and structural molecule activity were
related to them. The co-expression network reveals that physiological substances and endogenous hormones
interact with each other. Hormones regulate carbohydrate metabolism and partitioning, and physiological
substances in turn regulate hormone synthesis and signalling in a feedback manner. Under low-temperature
stress at the early stage of bud differentiation, the physiological substances MDA, sucrose, soluble protein,
and chlorophyll a were positively correlated, and at this time, a large number of reactive oxygen radicals were
generated in the plant, which attacked unsaturated fatty acids in the cell membrane and triggered lipid
peroxidation to produce malondialdehyde.The Mantel test analysis and the clustering heatmap indicated that
sucrose Sucrose was involved in the synthesis of growth hormone and cytokinins, and it induced the transport

of growth hormone. carrier genes (CmPIN1 and CmAUX1) and inhibits the expression of growth hormone signalling genes (CmIAA3, CmIAA14 and CmIAA16), and down-regulates the expression of growth hormone response factor gene CmARF2 in the terminal buds, which enhances growth hormone transport and reduces the level of endogenous growth hormone in the terminal buds, thus promoting the growth of flower buds. Soluble proteins showed a significant positive correlation with Auxin and GA. Soluble proteins and Auxin showed a positive correlation, and soluble proteins such as Klotho proteins affected the expression of genes related to renal tubular epithelial cell fibrosis by inhibiting the STAT3 phosphorylation pathway in a high glucose environment. Meanwhile, PIN proteins, the growth hormone transport carrier, interacted with other soluble proteins, and played a polar role in the polar transport of growth hormone The growth hormone transport carrier PIN protein interacts with other soluble proteins and plays a key role in polar transport of growth hormone.

In summary, the network of phytohormones and physiological substances regulating bud differentiation is a highly complex and ordered system, in which various phytohormones act in time and space in a balanced manner, controlling nutrient metabolism, regulating physiological processes at the cellular level to the regulation of gene expression, and comprehensively and finely regulating the whole process of bud differentiation from its initiation to its completion, in order to adapt to the various environmental changes in the process of plant growth and development. The following is a list of some of the most important aspects of the plant's development.

8 Acknowledgements

We are also grateful to the Key Laboratory of Genetics and Germplasm Innovation of Tropical Speciality Forestry and Floriculture of the Ministry of Education for its strong support of plant physiological experiments.

9 Competing interests

Shuyu Wang was the main author of the review, completing the collection and division of relevant literature and the writing of the first draft of the paper; Deqiao Cui, Meixing Wang, and Yanting Zhang were involved in the analysis of the literature, collation, and experimental personnel; Zeping Cai and Ying Zhao were involved in the experimental design and experimental study, and Xudong Yu was the conceptualiser and the person in charge of the project, guiding the whole experimental design, data analysis, and the paper writing and revision. All authors read and agreed to the final text.

10 Funding

This study was funded by the sub-project of Major Science and Technology Special Project of Hainan Province (No.323RC535).

11 Data and resource availability

All relevant data and resource can be found within the article and its supplementary information.

12 Diversity and inclusion statement

No Diversity and Inclusion Statement.

13 References

[1] **WANG R**.(2007). Study on the characteristics of flower bud differentiation and secondary flowering in peony,Beijing Forestry University,China.

[2] **YONGQIANG YU.et al.**(2023), ABLs and TMKs are co-receptors for extracellular auxin,Cell,Volume186, Issue 25,Pages 5457-5471.e17,ISSN 0092-8674 https://doi.org/10.1016/j.cell.2023.10.017

[3] **JIAHUI WANG, YUANYUAN LI, YANRU HU, SIRUI ZHU.**(2024). Jasmonate induces translation of the Ar-abidopsis transfer RNA-binding protein YUELAO1, which activates MYC2 in jasmonate signaling, *The Plant cell,* 37(1), koae294*.* https://doi.org/10.1093/plcell/koae294

[4] **WU JINSHAN,YU XUDONG,XU ZEKANG,et al.**(2017). Study on photosynthetic characteristics and carbon sink capacity of Haina-n Bombax ceiba in different months[J]. *China Tropical Agriculture*.(03):67-71.

[5] **WANG JIAN,SHUI QINGYAN,SHI JING, et al.**(2010). A preliminary study on the investigation and classification of the plant resources of *Bombax ceiba* in Hainan[J].*Subtropical Plant Science*,39(01):53-56.

[6] **HUANG JINRONG,XIE JINLAN,ZHANG QINYUAN.**(2009). Biological characteristics and seedling technology of *Bombax ceiba* [J].*Agricultural Technology Service.*26(05):149.

[7] **QIN RENQUAN,LIN JIANG,WANG HONGXING, et al.**(2017). Identification and biological chara-cterization of the causal agent of brown spot disease of *Bombax ceiba*[J].*Jiangsu Agricultural Science*. 45 (09):101-104

[8] **PAN HONGBING,DU BANG.**(2008). Grafting and breeding technology of *Bombax ceiba*[J].*Sichuan Fore-stry Science and Technology*,(01):86-87

[9] **LUO XIAOTOU,HU SHIJUN,MA HUANCHENG.**(2020). Analysis of the relationship between sta m-en charact-ers and their reproductive strategies in *Bombax ceiba*[J].*Forestry Survey and Planning*,45(02):143-148.

[9] **WANG GUOKAI,LIN BINBIN,QIN MINJIAN.**(2014). Studies on the chemical constituents of the leaves of *Bombax ceiba*(Ⅱ)[J].*Chinese Materia Medica*,37(02):240-242(王国凯,林彬彬,秦民坚.2014木棉叶化学成分研究(Ⅱ)[J].中药材,37(02):240-242.

[10] **LI XUSEN,XU GUANGKAI,WANG GUOKAI,et al.**(**2015).** Pr-ogress in the study of chemical c om-position and pharmacological effects of *Bombax ceiba*[J].*China Wild Plant Resources*,34(03):42-45+4 8.

[11] **TANG AICUN,YU YUAN, HUANG MIN, et al.**(2020)**.** Progress of chemical composition and pharmaco-logical effects of *Bombax ceiba*[J].*Chinese Et-hnic and Folk Medicine*,29(23):74-79.

[12] **SHAO L, JIN S,CHEN J, et al.**(2024), High-quality genomes of *Bombax ceiba* and Ceiba pen ta-ndra provide insights into the evolution of Malvace-aespecies and differences in their natural fiber d ev-elopment.*Plant Commun*.;5(5):100832 https://doi.org/10.1016/j.xplc.2024.100832

[13] **XU, Y.,LIU,J.,REN,C.,NIU,X.,ZHANG, T.& HUO, K.**(2024). Mulched Drip Fertigation with Growth Inhibitors Reduces Bundle-Sheath Cell Leakage and Improves Photosynthesis Capacity and Barley Production in Semi-Arid Regions. *Plants* (Basel, Switzerland), 13(2), 239. https://doi.org/10.3390/plants13020239

[14] **MALAKSHAHI KURDESTANI A, FRANCIOLI D,RUSER R, et al.**(2024).Optimizing nitrogen fer tili-zation in maize: the impact of nitrification inhibitors, phosphorus application,and microbial interaction s on enhancing nutrient efficiency and crop performance.*Frontiers in plant science*,15,1451573 https://doi.org/10.3389/fpls.2024.1451573

[15] **XIAOXIAO LI,CHUYU LIN,CHENGHAO LAN,ZENG TAO**,(2024),Genetic and epigenetic basis of phytohormonal control of floral transition in plants, *Journal of Experimental Botany*, Volume 75,Issue 14,23,Pages 4180–4194. https://doi.org/10.1093/jxb/erae105

[16] **WAN, CHUAN. et al.**(2022), "1′-OH of ABA and its analogs is a crucial functional group corr es-pondence to seed germination and development of plants." *Journal of Molecular Structure* 1249(1):1 31650. https://doi.org/10.1016/j.molstruc.2021.131650

[17] **YANG, D., CHEN, Y., WANG, R., HE, Y., MA, X., SHEN, J., He, Z., & Lai, H**. (2024). Effects of
Exogenous Abscisic Acid on the Physiological and Biochemical Responses of Camellia oleifera Seedlings
under Drought Stress. *Plants*, 13(2), 225. https://doi.org/10.3390/plants13020225

[18] **CHI, HY., OU, SL., WANG, MC. et al**.(2023) Physiological responses and Ethylene-Response
AP2/ERF Factor expression in Indica rice seedlings subjected to submergence and osmotic stress. *BMC Plant*
Biol 23, 372 . https://doi.org/10.1186/s12870-023-04380-y2

[19] **NAZIR, FAROZA. et al**. (2024) Plant hormone ethylene: A leading edge in conferring drought stress
tolerance.*Physiologia Plantar-umn*. pag. https://doi.org/10.1111/ppl.14151

[20] **HUANG, Y. C., YEH, T. H., & YANG, C. Y.**(2019). Ethylene signaling involves in seeds germination
upon submergence and antioxidant response elicited confers submergence tolerance to rice seedlings. *Rice*
(New York, N.Y.),12(1),23. https://doi.org/10.1186/s12284-019-0284-z

[21] **ZHU, Y., ZHANG, J., ZHAO, X., & XU, H.**(2024) Analysis of the Impact of Environmental Te-
mperature and Humidity on Cotton Fiber Quality During Initial Processing. *Journal of Natural Fibers,* 2
1(1). https://doi.org/10.1080/15440478.2024.2338538

[22] **HUANG, P. K., SCHMITT, J., & RUNCIE, D. E.** (2024). Exploring the molecular regulation of
v-ernalization-induced flowering sy-nchrony in Arabidopsis. *The New phytologist*, 242(3), 947–959. http
s://doi.org/10.1111/nph.19680

[23] **WANG, L., WANG, B., YU, H. et al.**(2020). Transcriptional regulation of strigolactone signalling in
Arabidopsis. *Nature* 583, 277–281 https://doi.org/10.1038/s41586-020-2382-x

[24] **HU SHUANGLING,LIN GONGTIAN,CHEN ZHEHONG, et al.**(2024),Morphological and structural
characteristics of flower bud differentiation in *Bombax ceiba* and changes in endogenous hormone content
during its differentiation[J/OL]. *Fujian Journal of A-griculture*,1-9.

[25] **ZHU ZHOU,ZHANG FANGQIU QIU, PAN WEN, et al**.(2016),Morphological and structural char
ac-teristics and division of ornamental types of *Bombax ceiba*[J].China Wild Plant Resources,35(06):42-
46+52.

[26] **LIN, B., WANG, D., LI, Z., LIU, J., XU, W., LI, X., & HUANG, J**.(2023). Morphological Development
and Nutritional Metabolism during Floral Bud Differentiation in *Acca Sellowiana (Feijoa)*. *International Journal*
of Fruit Science, 23(1), 102–115. https://doi.org/10.1080/15538362.2023.2216806

[27] **HIROYUKI TSUJI, MOEKO SATO**.(2024), The Function of Florigen in the Vegetative-to-Reproductive
Phase Transition in and around the Shoot Apical Meristem, *Plant and Cell Physiology*, Volume 65, Issue 3,
March 2024, Pages 322–337. https://doi.org/10.1093/pcp/pcae001

14 Figure legends

1. Morphological characteristics of flower bud differentiation in *Bombax ceiba*

Fig. 1 Morphological characteristics of flower bud differentiation in *Bombax ceiba*

A, E: early stage of flower bud differentiation (I); B, F: stage of calyx/petal primordium differentiation (II);

C, G: stage of stamen/pistil primordium differentiation (III)

Note: GC: growth cone; Cap: calyx primordium; PeP: petal primordium; Ca: calyx; Pe: petal; PiP: pistil

primordium; StP: stamen primordium

2.Characterization of external morphological structure of single flowers

Fig.2 Characterization of external morphological structure of single flowers

Note:A-B: primordial stage ; C: fastigiate stage; D: large balloon stage; E: initiation stage; B-F: full bloom

stage

3.Variation of physiological substance content in flower bud differentiation

465
Fig. 3 Variation of physiological substance content in flower bud differentiation

Note:A: Soluble sugar content (mg•g-1. FW); B: Sucrose content (mg•g-1. FW); C: Soluble starch content
(mg•g-1. FW); D: Soluble protein content (mg•g-1. FW); E: MDA content (mg•g-1. FW); F: Chlorophyll a
content (mg•g-1. FW); G: Chlorophyll b content (mg•g-1. FW); H: Carotenoids content (mg•g-1. FW); I:
Chlorophyll (a+b)/Carotenoid; J: Chlorophyll a/b

4.Dynamic changes of hormone levels in flower bud differentiation

472
473 Fig 4 Dynamic changes of hormone levels in flower bud differentiation

474 Note:A:Abscisic acid analogs (ABA); B:Salicylic acid analogs (SA); C:Gibberellins (GA); D:Ethylene
475 analogs (ETH); E:Solanum lactone (SL); F:Cytokinins (CK); G:Growth hormone (Auxin) H:Jasmonic acid
476 (JA)

477
478
479

5.Screening of differential metabolites of flower bud differentiation

 Fig 5 Screening of differential metabolites of flower bud differentiation

Note:A: Differential metabolite content map; B: Differential metabolite KEGG classification map (stage I and II); C: Differential metabolite KEGG classification map (stage I and III); D: Differential metabolite KEGG classification map (stage II and III); E: Differential metabolite Venn and Upset maps; F: Differential metabolite related network map

6.Correlation network of physiological and hormonal substances for flower bud differentiation

490
491   Fig 6 Correlation network of physiological and hormonal substances for flower bud differentiation
492   Note:A: Manteltest analysis; B: Correlation network diagram; C: Clustering heat map