Dynamics of Bacterial and Viral Transmission in Experimental Microbiota Transplantation

James Weagley[1], Heyde Makimaa[1], Luis Alberto Chica Cárdenas[1], Ana Romani[1], Meagan Sullender[1], Somya Aggarwal[1], Michael Hogarty[1], Rachel Rodgers[1,2], Elizabeth Kennedy[1], Lynne Foster[1], Lawrence A. Schriefer[1], Megan T. Baldridge[1,3#]

[1] Division of Infectious Diseases, Department of Medicine, Edison Family Center for Genome Sciences & Systems Biology, Washington University School of Medicine, St. Louis, MO, USA

[2] Department of Pediatrics, Washington University School of Medicine, St. Louis, MO, USA.

[3] Department of Molecular Microbiology, Washington University School of Medicine, St. Louis, MO, USA.

[#] Correspondence: mbaldridge@wustl.edu

Abstract

Mouse models are vital tools for discerning the relative contributions of host and

microbial genetics to disease, often requiring the transplantation of microbiota between

different mouse strains. Transfer methods include antibiotic treatment of recipients and

colonization using either co-housing with donors or the transplantation of fecal or cecal donor

material. However, the efficiency and dynamics of these methods in reconstituting recipients

with donor microbes is not well understood. We thus directly compared co-housing, fecal

transplantation, and cecal transplantation methods. Donor mice from Taconic Biosciences,

possessing distinct microbial communities, served as the microbial source for recipient mice

from Jackson Laboratories, which were treated with antibiotics to disrupt their native

microbiota. We monitored microbial populations longitudinally over the course of antibiotics

treatment and reconstitution using 16S rRNA gene sequencing, quantitative PCR, and

shotgun sequencing of viral-like particles. As expected, antibiotic treatment rapidly depleted

microbial biomass and diversity, with slow and incomplete natural recovery of the microbiota

in non-transplanted control mice. While all transfer methods reconstituted recipient mice with

donor microbiota, co-housing achieved this more rapidly for both bacterial and viral

communities. This study provides valuable insights into microbial transfer methods, enhancing

reproducibility and informing best practices for microbiota transplantation in mouse models.

Introduction

As our appreciation continues to grow for the diverse range of phenotypes the trillions of microbes inhabiting our bodies, termed the microbiota, can mediate, so does our reliance on manipulatable small animal models to experimentally interrogate these effects. The gut microbiota, a complex community of microorganisms residing in the gastrointestinal tract, plays a pivotal role in host health and physiology[1,2]. These microbial communities are involved in critical functions such as nutrient metabolism[3,4], immune system development[5–7], and protection against pathogens[8–10]. Dysbiosis, or the disruption of the normal microbiota balance, has been implicated in a range of diseases including obesity[11,12], malnutrition[13,14], and inflammatory bowel diseases[15,16]. Understanding the intricate relationships between the microbiota and host systems is therefore essential for developing therapeutic strategies targeting microbiome-related diseases.

To unravel these complex host-microbe interactions, researchers heavily leverage animal models, particularly mice. Controlled studies using mouse models enable the manipulation of microbial communities to observe resultant effects on host physiology and disease outcomes. A key aspect of such research is the generation of gnotobiotic mice— animals that are germ-free or colonized with specific, defined microbial communities[17]. These models provide a unique platform to study the microbiota's role under controlled conditions.

Despite the utility of germ-free mice for numerous applications, however, their use is associated with some drawbacks. For investigators studying numerous mouse genotypes, executing experiments not amenable to the use of a gnotobiotic isolator such as those involving particular pathogens or behavioral testing, or performing studies requiring a normal microbiota during development which is then disrupted, germ-free mice may be impractical or inappropriate. Antibiotics treatment represents an accessible alternative for the depletion of the microbiota, and this approach has been broadly used by many investigators to examine many of the same phenotypes explored in germ-free mice, often with strong concordance between the two models[17].

Establishing gnotobiotic mice or reconstituting the microbiota after perturbation necessitates reliable and efficient methods for microbial transfer. Common techniques include fecal microbiota transplantation (FT), cecal content transplantation (CT), and co-housing (CH). Fecal transplantation involves transferring fecal material from a donor to a recipient, while cecal transplantation is similar but utilizes the cecal contents, which may have a different complex microbial composition due to variation along the gastrointestinal tract. Co-housing leverages the natural coprophagic behavior of mice, allowing for the transfer of microbes through direct contact and shared environments. Despite their widespread use, experiments designed utilizing these methods are generally based on historical precedent rather than a comprehensive understanding of their efficacy or the specific dynamics of microbial colonization they promote.

This reliance on convention can lead to assumptions that impact experimental outcomes and reproducibility, a major challenge facing the field[18–21]. For instance, variation in microbial transfer efficiency can affect the establishment of the desired microbiota, influencing the host's physiological responses and potentially skewing results[22,23]. Furthermore, while bacterial community dynamics have been relatively well-studied, the transfer and establishment of viral communities (the virome) remain less understood. Given the growing recognition that bacteriophages and other viruses shape microbial communities and influence host health[24–30], it is crucial to consider both bacterial and viral components when evaluating these methods.

Herein, we systematically compare fecal transplantation, cecal transplantation, and co-housing as methods for transferring microbes to microbially-depleted mice, characterizing this process longitudinally. Recipient mice from Jackson Laboratories were treated with antibiotics for one week to disrupt their native microbiota. Donor mice from Taconic Biosciences, known to have distinct microbial communities from Jackson, served as the source of bacteria and viruses for transplantation[31]. By employing 16S rRNA gene sequencing and quantitative PCR (qPCR), we monitored bacterial load, diversity, and composition. Additionally, we analyzed

viral community composition and diversity through short-read shotgun sequencing of viral-like

particles.

Our findings indicate that co-housing facilitates a more rapid transfer of both bacterial

and viral communities compared to fecal and cecal transplants, which performed similarly to

each other. While all methods resulted in bacterial communities similar to the donor by day 15

post-transplantation, viral communities exhibited greater diversity following co-housing. These

results highlight method-specific influences on microbial community dynamics and underscore

the importance of method selection in experimental design.

By carefully characterizing these commonly used methods, our study provides

valuable insights that can enhance reproducibility and consistency in microbiota research.

This work informs best practices for replacing the enteric microbiota of mice and contributes

to our understanding of the mechanisms underlying microbial transfer and establishment in

experimental models. Such insights are essential for advancing microbiota research and for

the development of interventions targeting microbiome-related diseases.

Results

Effect of antibiotic treatment on bacterial load, diversity, and community composition

Based upon prior studies leveraging antibiotic treatments of mice, we elected to use a simplified cocktail of vancomycin, neomycin, and ampicillin, which we and others have shown mediates robust depletion of the microbiota within three to seven days of treatment initiation and is also well-tolerated by recipients[10,11]. We used C57BL/6 mice from Jackson Laboratory (JAX) as our recipients and C57BL/6 mice from Taconic (TAC) as our donors for microbiota transfer, as mice from these different vendors are well-established to have distinct enteric microbial communities[12,13]. Antibiotics were administered to JAX recipient mice (n=13, 16, and 16 respectively in three independent experiments, subdivided into four groups corresponding to method of transplantation as described below) for 7 days, while TAC donor mice (n=4 per experiment) and control groups of JAX (n=4 per experiment) and TAC (n=4 per experiment) mice were left untreated (**Figure S1A**).

Fecal samples were collected on days 0, 1, 3, and 7 and subjected to 16S rRNA gene qPCR and sequencing to characterize the microbial community prior to and during antibiotic treatment. The bacterial load as determined by 16S qPCR was not significantly different in JAX and TAC mice prior to antibiotic treatment (**Figure S1B**, Day 0, $p > 0.05$). Antibiotics rapidly and significantly reduced the bacterial load in JAX mice 1 day after treatment initiation (**Figure S1B**, Day 1, $p < 0.05$) and 16S levels were below the limit of detection following 7 days of antibiotic treatment (**Figure S1B**). The number of 16S copies did not change significantly over the first 7 days of the experiment in untreated JAX and TAC mice.

We next evaluated sequencing data for the V4 region of the 16S rRNA gene for these samples. Sequencing reads were processed using the DADA2 pipeline implemented in Nephele to generate amplicon sequence variants (ASVs)[32,33]. Bacterial species were defined by grouping ASVs with greater than 97% sequence identity. ASV richness was similar between JAX and TAC mice at Day 0 (mean ± SE: TAC=~80 ± 6 OTUs, JAX=66 ± 5; **Figure S1C**). Treatment with antibiotics significantly reduced the number of bacterial species present in the stool of JAX mice within 1 day (mean ± SE= 43 ± 6; **Figure S1C**). Bacterial richness

was not significantly different on day 3 of treatment but was reduced to one-third of the original

level by day 7. Although the 16S levels were below the limit of detection in antibiotic-treated

mice on day 7, sequencing suggested that some of the native bacteria persisted at a low level.

To understand how community composition was affected by antibiotic treatment and

the subsequent reduction in bacterial biomass, the abundance of bacterial species was used

to calculate the unweighted UniFrac distance between all samples collected over the full

course of the experiment, and dimensionality reduction via principal coordinate analysis

(PCoA) was performed. The data were then subset to samples collected during the first seven

days to visualize the effect of antibiotic treatment on community composition. PCoA axis 1

explained 24.6% of the total variance in bacterial community composition, with variance along

this axis driven by antibiotic treatment of JAX mice (**Figure S1D**). The variance along PCoA

axis 2 (13.0%) split untreated JAX and TAC mice. Antibiotic treatment contributed more to

overall variance in the bacterial community composition than the vendor. One week of

treatment with vancomycin, neomycin, and ampicillin was sufficient to substantially reduce

bacterial load and diversity and drastically alter bacterial community composition.

Supplementary Figure 1. One-week treatment with antibiotic cocktail significantly alters fecal bacterial biomass and community composition. (**A**) Mice were obtained from Jackson or Taconic labs. Mice from Jackson Labs were provided vancomycin, neomycin, and ampicillin in their drinking water for one week (Jackson + Antibiotics). Untreated Jackson control mice and Taconic mice were maintained on normal drinking water. (**B**) Bacterial load measured by qPCR of 16s rDNA gene throughout the first week of the experiment. The letters next to each point indicate significant groupings, as those with the same letter were not significantly different from one another by two-way ANOVA followed by Tukey's HSD post-hoc test for pairwise comparisons with an alpha of 0.05. (**C**) Bacterial species richness, the number of unique bacteria species observed per mouse, throughout the first week of the experiment

via V4-16S rRNA gene sequencing. The letters next to each point indicate significant
groupings, as those with the same letter were not significantly different from one another by
two-way ANOVA followed by Tukey's HSD post-hoc test for pairwise comparisons with an
alpha of 0.05. (**D**) Principal Coordinate Analysis of unweighted UniFrac distances calculated
from species counts. Error bars (± SE) intersect at centroid of all samples collected for each
group of mice on the day post-antibiotics treatment, indicated by the number on the plot.
167
168

Route of transfer regulates community reconfiguration during transplantation following

antibiotic treatment

We next compared the outcomes of different transfer methods for the transplantation

of TAC microbiota into antibiotic-treated JAX mice. Cessation of antibiotics occurred on day

7, and antibiotic-treated JAX mice were either (i) maintained with no intentional transfer of

microbes (n=12, 8-12 samples per timepoint combined across three experiments), or gavaged

with prepared (ii) cecal (n=12, 7-12 samples per timepoint) or (iii) fecal (n=12, 8-12 samples

per timepoint) contents from untreated TAC mice, or (iv) co-housed (n=11, 6-11 samples per

timepoint) with untreated TAC donors and sampled over the following two weeks (**Figure 1A**).

Additionally, TAC (n=3, 0-3 samples per timepoint) and JAX (n=10, 6-10 samples per

timepoint) mice that did not receive antibiotics were maintained and sampled over these two

weeks to monitor changes over time in the background communities.

Bacterial load was significantly increased two days after cessation of antibiotics (Day

9) in all mice that received microbial transplantation relative to those that did not, regardless

of method (**Figure 1B**). Although the bacterial load returned to detectable levels in mice that

did not receive a microbial transplant on day 9 it was nearly two orders of magnitude lower

than that of untreated mice or of those that received a microbial transplant. By day 11, there

was no significant difference in the number of 16S copies detected between the methods of

transfer or untreated mice from JAX or TAC. After one week (day 14), the bacterial load was

similar between all groups for the duration of the experiment. Although increasing steadily

post-transplantation, the recovery of bacterial alpha-diversity (species richness), was slower

than that of overall bacterial load, regardless of transplantation approach (**Figure 1C**).

The effect of the method of transplant on community composition was evaluated next.

Samples were separated based on the day they were collected and the treatment group to

which the mouse belonged. Unweighted UniFrac distances were calculated between all

samples to assess global changes in community composition and beta diversity throughout

the experiment. PCoA computed on this full set enabled the visualization of microbial

community recovery and transfer following antibiotic treatment over time between days 9 and

21 (**Figure 1D**). Confidence ellipses designating the samples from untreated JAX mice and

TAC mice represent the original recipient and donor community compositions in feces

collected from mice that did not receive antibiotics from days 0 through 7.

The bacterial communities of JAX antibiotic-treated mice administered either fecal or

cecal transplant gradually approach the TAC donor community composition between days 9

and 11, converging on the donor community profile by day 15 (**Figure 1D**). Thus, although

fecal or cecal transplant restored bacterial loads to levels similar to TAC donors within two

days (**Figure 1B**), the composition of the bacterial community was not fully recapitulated by

day 9 with either of these transfer methods. In contrast, after only two days (day 9), the

bacterial composition of co-housed mice was equivalent to that of TAC donors (**Figure 1D,**

E). The bacterial community of JAX mice that received antibiotics but no transfer gradually

returned towards that of untreated JAX mice, eventually achieving a similar state to the original

JAX microbiota by experimental day 21.

The unweighted UniFrac distances of the treatment arms were statistically compared

to untreated Taconic and Jackson control mice throughout the experiment using

PERMANOVA. To account for drift in community composition over time in the control

communities, as well as low sample sizes from untreated Taconic controls, samples collected

on each day of the experiment for the treatment arms were compared to pooled control

samples collected within the same week. The fecal bacterial communities of antibiotic-treated

animals were significantly different than JAX controls following 1 day of antibiotic treatment,

the difference between these groups remained significant and increased through the end of

antibiotic treatment on day 7 (**Figure 1E**). Additionally, the bacterial community of all these

groups of mice was significantly different than the TAC controls, also increasing until day 7

(**Figures 1D, E**). By day 9, the earliest timepoint sampled following transplantation, the

bacterial microbiota of antibiotic-treated JAX mice co-housed with TAC donors were no longer

significantly different from TAC controls and were not significantly different for the remainder

of the experiment. Mice that received cecal or fecal transfer were significantly different from

the Taconic donors until Day 14, suggesting differences in the efficiency of bacterial transfer

across the three methods. Like the trends observed in the 16S qPCR data (**Figure 1B**), the

bacterial community of JAX mice that were treated with antibiotics that did not receive

transplantation gradually recovered and was not significantly different from the JAX controls

by day 21 (**Figures 1D, E**). However, the average distance between these two groups was

still elevated (0.69 ± 0.1 [mean ± SD]) and had not returned to baseline (0.47 ± 0.1 [mean ±

SD]). Interestingly, although co-housed mice displayed the most rapid transition to a Taconic-

like bacterial community, they were not significantly different from JAX controls on days 14

and 21. This may be indicative of incomplete microbial transfer and/or recovery of JAX bacteria

in co-housed mice resulting in an 'intermediate' community state.

Figure 1. Cohousing of mice reconstitutes the bacterial microbiota more rapidly than fecal or cecal transplant. (**A**) Antibiotic-treated Jackson mice were either allowed to recover naturally (Jax-Antibiotics-NoTransfer), or were colonized with Taconic-associated microbes via gavage of cecal (Jax-Antibiotics-CecalTransfer) or fecal (Jax-Antibiotics-FecalTransfer) material, or co-housing (Jax-Antibiotics-Cohouse). Jackson (Jax-NoT) and Taconic (Tac-NoT) mice that did not receive antibiotics were maintained as controls. (**B**) Bacterial load measured by qPCR of the 16S rRNA gene throughout the final two weeks of the experiment shows the rapid recovery of bacterial load following transplantation. The letters next to each point indicate significant groupings, those with the same letter were not significantly different from one another by two-way ANOVA followed by Tukey's HSD post-hoc test for pairwise comparisons with an alpha of 0.05. (**C**) Bacterial species richness, the number of unique bacterial species observed per mouse, throughout the first week of the experiment via v4-16s sequencing. The letters next to each point indicate significant groupings, those with the same letter were not significantly different from one another by two-way ANOVA followed by Tukey's HSD post-hoc test for pairwise comparisons with an alpha of 0.05 (**D**) Principal Coordinate Analysis of unweighted UniFrac distances calculated from species counts across all three weeks of the experiment illustrate the change in community composition following antibiotic treatment and

microbiota transplantation. Ellipses (95% confidence) represent the baseline fecal bacterial
communities of untreated Jackson (Jax-NoT) and Taconic (Tac-NoT) mice during the first
week of the experiment. (**E**) Bar heights indicate the mean Unweighted UniFrac distance was
calculated between the fecal bacterial community of Jackson mice treated with antibiotics that
received microbial transplantation via co-housing, cecal transfer, fecal transfer, or no transfer,
and the fecal bacterial community of control Jackson and Taconic mice that never received
antibiotics, pooled by week (days pooled: [0,7), [7,14), [14,21]). Striped bars below the x-axis
indicate the distance from untreated Jackson controls while solid bars above the x-axis
indicate the distance from untreated Taconic controls. Error bars indicate the standard
deviation of all pairwise comparisons between the indicated groups. A PERMANOVA test was
performed on the unweighted UniFrac distance matrix for the comparisons shown followed by
Bonferroni's corrections for multiple comparisons (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).
265
266

Co-housing rapidly transfers Taconic-associated bacterial taxa and their predicted functional

capabilities

To better understand the dynamics of bacterial transfer following transplantation, ASVs

were classified as being either Taconic- (162 ASVs) or Jackson-associated (124 ASVs), or

neither based on their prevalence and abundance in samples collected from untreated mice

over the first week of the experiment (**Figure 2A**). These subsets of ASVs were then tracked

throughout the course of the experiment (**Figure 2B**). As expected, Taconic ASVs were higher

in Taconic mice than in Jackson mice that didn't receive transplantation, significantly so on

days 7 and 8. Although there was a transient increase in the richness of Taconic ASVs on

days 1 and 3 in all groups measured, by day 7 the Taconic-associated ASV richness in all

Jackson mice (mean = 0.7 – 2.6 Taconic ASVs per sample) was significantly lower than in

Taconic donors (~36.3 Taconic ASVs per sample) (**Figure 2B**). By day 9, the number of

Taconic ASVs had substantially increased in co-housed mice (mean ± SE = 26.9 ± 2.0) and

was significantly higher than untreated Jackson controls (mean ± SE = 0.7 ± 0.1) and not

significantly different than Taconic co-housed donors (mean ± SE = 30.8 ± 4.5). These were

samples from mice originally from Taconic that shared a cage with antibiotic-treated Jackson

recipient mice. At this time, mice that received cecal or fecal transplants did not have

significantly higher richness of Taconic-associated ASVs (means ± SEs = 4.6 ± 0.8 and 4.3 ±

0.4, respectively). By day 15, the richness of Taconic-associated ASVs in Jackson mice that

received antibiotics followed by cecal or fecal transplant was significantly elevated relative to

Jackson mice that received antibiotics and no subsequent transplantation and remained

relatively steady until the end of the experiment on day 21. Overall, these data are consistent

with our observation that co-housing more rapidly introduces donor-specific bacteria than

cecal or fecal transplant.

To complement our taxonomic description of this transplantation process, we next

sought to characterize the functional capacity of these bacterial communities. The functional

capabilities of microbial communities can be similar even if there are distinct taxonomic

differences at the ASV level. PICRUSt2 was used to predict the abundance of MetaCyc

pathways in each of our samples[34,35]. Antibiotic-treated Jackson mice and Taconic co-housed

donors were compared to Jackson and Taconic controls grouped by week to identify the total

number of differentially abundant pathways using DESeq2 (**Figure 2C**)[36]. A larger number of

differentially abundant pathways indicates that the predicted functional capabilities of the

communities are more disparate. Consistent with our prior taxonomic results, by day 9, the

number of differentially abundant pathways between co-housed mice and Taconic control

mice dropped off precipitously, from over 200 on day 7 to less than 100 on day 9. Mice

receiving a cecal transplant didn't drop below 100 until day 14, while mice receiving a fecal

transplant didn't drop below 100 until day 21.

Figure 2. Defining and tracking Taconic-associated ASVs and their predicted functional capacities. (**A**) Bacterial ASVs were defined as being Taconic or Jackson associated based on their prevalence and abundance in samples collected from antibiotics-free mice over the first week of the experiment. ASVs that had an absolute combined $\log_2$(Fold Change) greater than 4 were consider Taconic-associated while those that had an absolute combined $\log_2$(Fold Change) less than -4 were considered Jackson-associated. (**B**) Richness of Taconic ASVs was monitored throughout the course of the experiment, plot show mean richness in each group at a given timepoint, error bars represent the standard error, compact letter display indicates results of pairwise Kruskal-Wallis followed by Bonferroni's multiple comparison correction with an alpha threshold of 0.05. (**C**) Abundance of functional pathways were predicted using PICRUSt2. Bar heights indicate the number of pathways determined to be significantly differentially abundant using DESeq2 (alpha = 0.05) to compare between the fecal bacterial community of Jackson mice treated with antibiotics that received microbial transplantation via co-housing, cecal transfer, fecal transfer, or no transfer, and Taconic co-

319   housed mice, and the fecal bacterial community of control Jackson and Taconic mice that
320   never received antibiotics, pooled by week (days pooled: [0,7), [7,14), [14,21]).
321

Transfer of specific bacterial taxa differ between methods of transplantation

Changes in the taxonomic composition of the bacterial community were profiled

throughout the course of the experiment. In line with previous results, the microbiota of control

communities was relatively stable over time (**Figure 3A**). Bacilli and Gammaproteobacteria

noticeably increased in relative abundance in mice that received the antibiotic cocktail by day

3 (**Figure 3A**), although the overall bacterial load (**Figure 1B**) and richness remained low

(**Figure 1C**).

By day 9, two days after beginning microbial transfer, co-housed mice were

significantly enriched for Taconic ASVs belonging to the *Tannerellaceae*, *Prevotellaceae*, and

Muribaculaceae families, as well as for *Lactobacillaceae* ASVs that were not statistically

associated with the original Jackson or Taconic communities, relative to mice that did not

receive transplantation (**Figure 3B**). Mice that received cecal or fecal transplant were also

enriched for these *Lactobacillaceae* ASVs at day 9, along with *Enterobacteriaceae* ASVs that

were not associated with either supplier, relative to mice that did not receive a transplant.

Bacteroidaceae ASVs associated with the original Taconic communities were also enriched

at day 9 in mice that received fecal or cecal transplants. The *Enterobacteriaceae* ASVs were

significantly enriched in mice receiving cecal transplant compared to co-housed mice, while

co-housed mice were significantly enriched for *Muribaculaceae* ASVs.

Taconic-associated ASVs from the *Muribaculaceae* family were significantly enriched

at day 15 in all mice that received a microbiota transfer, regardless of transplantation method,

compared to mice that did not receive transplantation (**Figure 3C**). Taconic *Bacteroidaceae*

ASVs were significantly enriched in mice that received a cecal transplant while Taconic-

associated *Akkermansiaceae* ASVs were significantly enriched in mice that received a fecal

transplant relative to the 'no transfer' group. Taconic-associated *Akkermansiaceae* ASVs were

also significantly enriched in mice receiving fecal transplant compared to co-housed mice.

Jackson-associated *Erysipelotrichaceae* ASVs were the first to recover in antibiotic treated

mice that did not receive a transplant (**Figure 3C**). By day 15, these bacteria were significantly

enriched relative to mice that were co-housed, or mice that received a fecal or cecal transplant.

By day 21, significant differences in bacterial taxa enrichment were observed across

groups. Taconic *Staphylococcaceae* ASVs were significantly enriched in the 'no transfer'

group relative to co-housed mice and also enriched in the fecal transplant group relative to co-

housed mice. Taconic *Lactobacillaceae* ASVs were enriched in the 'no transfer' group relative

to co-housed mice. Notably, *Christensenellaceae* ASVs, which were not associated with

either supplier, were significantly enriched in the 'no transfer' group compared to all

transplantation groups. Additionally, several Jackson-associated taxa,

including *Ruminococcaceae*, *Lachnospiraceae*, *Erysipelotrichaceae*, *Clostridiaceae*,

and *Acholeplasmataceae*, were enriched in the 'no transfer' group relative to all transplant

groups. These findings highlight that by day 21, the 'no transfer' group retained significant

enrichment of Jackson-associated and non-supplier-specific taxa, while *Taconic*-associated

taxa remained prominent in transplanted groups. The fecal transplant group exhibited a

distinct enrichment pattern, including higher levels of *Staphylococcaceae*, relative to co-

housed mice.

Figure 3. Specific bacterial taxa are enriched across transplantation methods (A) Relative abundance of bacterial families throughout the course of the experiment in each of the treatment groups. (**B**) Bacterial families associated with original bacterial communities of mice from Taconic (orange), Jackson (green) or neither (black) that were significantly enriched or depleted for the comparisons shown below each plot as determined using ANCOM-BC2. Panels show results from days 9, 15, and 21. Log fold change values in green were significant after applying a stringent mixed directional FDR control.

Bacterial diversity and community composition differ across gastrointestinal sites but not the

method of transfer

At the end of the experiment, bacterial diversity and community composition were

assessed across different gastrointestinal sites: cecal, ileal, and stool samples. Observed

bacterial diversity varied significantly across these sites and treatment groups, with lower

richness in ileal contents and mice that did not receive microbiota transfer following antibiotic

treatment (**Figure 4A**). Bacterial richness in the stool of mice that received transplantation

trended higher than Jackson controls and lower than Taconic controls, suggesting incomplete

transfer of bacterial taxa across all three methods, although richness was significantly higher

than in mice that were treated with antibiotics without transplantation. Principal coordinate

analysis of Unweighted UniFrac distances further illustrated the community composition

differences, with samples clustering according to treatment groups and gastrointestinal sites

(**Figure 4B**). Axis 1 and Axis 2 explained 19.8% and 15.6% of the variance, respectively. The

distinct clustering patterns indicated differences in bacterial community composition between

the treatment groups. Jackson mice that received antibiotics and no subsequent microbial

transplantation clustered together in the upper right quadrant. Ileal and stool samples from

mice that received a microbiota transfer resembled control communities with samples from

the same gastrointestinal site clustering together. To compare community compositions

across treatments, PCoA was performed on samples subset by the gastrointestinal site

(**Figures 4C-E**). JAX mice that received antibiotics, but no microbial transplant harbored

distinct microbial communities. Mice that received transplantation tended to cluster together,

with differences between mice that received transplants or Taconic co-housed donors and

untreated Jackson controls being most pronounced in cecal and ileal contents (**Figures 4D,**

E). Additionally, the ileal bacterial communities of mice that received fecal or cecal transplants

clustered closely together and were distinct from either co-housed group (**Figure 4E**), a

difference that was not as pronounced in stool or cecal samples (**Figures 4C, D**). These

findings provided insight into the dynamics of the re-establishment of bacterial communities

at gastrointestinal sites across methods of microbiota transplantation.

Figure 4. Bacterial diversity and community composition across different body sites and treatment groups (**A**) Bacterial richness, the number of bacterial taxa observed in cecal, ileal, and stool samples across different groups, including Jackson controls, co-housed, cecal transfer, fecal transfer, no transfer, and Taconic controls. (**B**) Plot of Principal Coordinate Analysis (PCoA) of Unweighted UniFrac Distances illustrating bacterial community composition from cecal, ileal, and stool samples, across each method of transplantation, including Jackson controls and cohoused Taconic donors. PCoA of Unweighted UniFrac Distances of samples on Day 21 across treatment groups from (**C**) stool, (**D**) cecal, or (**E**) ileal contents.

Viral Community Dynamics Post-Antibiotics

We next sought to characterize the viral response to antibiotic treatment, given the significant reduction in bacterial load and diversity observed over the first week of the experiment (**Figures 1B, C**). Short-read shotgun sequencing of DNA purified from viral-like particles was performed to evaluate the impact of antibiotic treatment on viral communities. Viral richness (the number of unique contigs assembled from shotgun sequencing data) from days 0 to 7 was compared between Taconic and Jackson controls and Jackson mice that received antibiotics during the first week of the experiment. Read counts were normalized across sequencing libraries to mitigate batch effects. The results showed a significant reduction in viral richness following antibiotic administration, with richness dropping from 596 ± 84 (mean

± SE) unique viral contigs on day 0 to 34 ± 2 (mean ± SE) on day 7 (**Figure S2A**). The same

pattern was observed when contigs were subset to only those with clear viral signatures, and

when aggregated into clusters representing viral species (OTUs), genera, and families (**Figure

S2A**). The reduction in viral richness highlights the susceptibility of viral populations to

disruptions in the bacterial community, likely due to their dependence on bacterial hosts. A

marked shift in viral community composition, driven primarily by the antibiotic treatment, was

observed in PCoA of Jaccard distances (**Figure S2B**). The separation along axis 1 was

predominantly between antibiotic-treated and untreated groups, underscoring the substantial

impact of antibiotics on viral community composition. Again, similar results were obtained

when considering only high-quality viral contigs, or different taxonomic ranks (**Figure S2B**).

Supplemental Figure 2. One-week treatment with antibiotic cocktail significantly alters fecal viral richness and community composition. (**A**) Viral richness and (**B**) Principal Coordinate Analysis of Jaccard distances calculated from contig read counts of unique viral contigs observed per mouse, throughout the first week of the experiment via shotgun metagenomics sequencing of viral-like particles from feces. Panels represent data before (i) and after contigs had been stringently filtered and grouped into clusters representing ii) strains, iii) species/OTUs, iv) genera, and v) families. In A, the dot indicates the mean richness for that group, error bars (± SE), in B, the error bars (± SE) intersect at the centroid of all samples collected for each group of mice on the day indicated by the number on the plot. In A, the letters next to each point indicate significant groupings, those with the same letter were not significantly different from one another by two-way ANOVA followed by Tukey's HSD post-hoc test for pairwise comparisons with an alpha of 0.05.

 Viral Community Dynamics Post-Transplantation

To assess the impact of different microbiota transfer methods on viral communities in

antibiotic-treated mice, we conducted a comparative analysis of viral richness and community

composition following the various transplantation approaches described previously. Viral

communities were sampled from days 7 to 14 to capture the reestablishment of viral

populations post-transplantation.

Viral richness, as measured by the number of unique viral contigs observed, did not

show a uniform increase across all groups immediately after the cessation of antibiotics. By

day 9, the co-housed mice (Jax-Antibiotics-Cohouse) exhibited a more rapid increase in viral

richness compared to the other groups, and their viral richness was not significantly different

from that of mice from Taconic (**Figure 5A**). Viral richness remained relatively low in mice that

received fecal or cecal transfers, or no transfer at this timepoint. Viral richness increased in all

groups by day 11, although all groups that received transplantation had lower viral richness

than Taconic donors, this difference was not significant. Viral richness in mice that did not

receive a microbial transplant (Jax-Antibiotics-NoTransfer) remained significantly lower and

only marginally increased by day 11. By day 14, viral richness levels in all transfer groups

converged and became similar to those in untreated controls, indicating successful

colonization.

We then evaluated the impact of the transplantation methods on viral community

composition, visualized using Principal Coordinate Analysis (PCoA) of Jaccard distances. The

results showed a gradual alignment of the viral communities in fecal and cecal transplanted

mice (Jax-Antibiotics-FecalTransfer and Jax-Antibiotics-CecalTransfer) with those of the

Taconic donor community profiles between days 9 and 11, with full convergence observed by

day 14 (**Figure 5B**). Notably, viral communities in co-housed mice (Jax-Antibiotics-Cohouse)

rapidly converged with the donor profiles by day 9, indicating that co-housing facilitated a more

efficient and quicker viral transfer. On the other hand, viral communities in mice that did not

receive any transplant (Jax-Antibiotics-NoTransfer) showed a delayed recovery, remaining

distinct from both donor and control profiles until the final day of sampling, gradually

approaching the composition of the original Jackson community by day 14.

These findings suggest that the method of microbiota transplantation plays a crucial

role in the reestablishment of viral communities. Co-housing was the most effective approach

for rapid and complete viral transfer, while fecal and cecal transplants also facilitated

successful viral colonization, albeit at a slower pace. Mice that did not receive a microbial

transplant exhibited a delayed and incomplete recovery of viral communities, highlighting the

importance of external microbial inputs for restoring gut virome diversity and composition after

antibiotic treatment.

484
Figure 5. Cohousing of mice reconstitutes the viral microbiota more rapidly than fecal
or cecal transplant (**A**) Viral richness from days 7 through 14 of the experiment shows the
rapid recovery of viral diversity following transplantation. The letters next to each point
indicate significant groupings, those with the same letter were not significantly different from
one another by two-way ANOVA followed by Tukey's HSD post-hoc test for pairwise
comparisons with an alpha of 0.05 (**B**) Principal Coordinate Analysis of Jaccard distances
calculated from contig read counts across two weeks of the experiment illustrate the change
in community composition following antibiotic treatment and microbiota transplantation.
Ellipses (95% confidence) represent the baseline fecal viral communities of untreated
Jackson (Jax-NoT) and Taconic (Tac-NoT) mice. (**C**) Richness and (**D**) relative abundance
of viral contigs defined as (i) Jackson or (ii) Taconic-associated throughout the course of the

experiment. The letters next to each point indicate significant groupings, those with the same
letter were not significantly different from one another by two-way ANOVA followed by
Tukey's HSD post-hoc test for pairwise comparisons with an alpha of 0.05. The (**E**) richness
and (**F**) relative abundance of viral contigs associated with mice from Jackson, Taconic, or
neither, over the first two weeks of the experiment.
501

Transfer of Taconic-associated viruses to Jackson mice

To track the movement of viruses between Taconic and Jackson mice, viral contigs

were defined as being associated with mice from Taconic, Jackson, or neither, based on their

differential prevalence and abundance in samples collected over the first week of the

experiment from mice that did not receive antibiotics (n = 34 Taconic samples, 50 Jackson

samples). Of the 6,249 viral contigs, 2,535 were defined as Taconic-associated while 1,129

were defined as Jackson-associated.

The richness and relative abundance Jackson-associated viruses significantly

decreased in response to antibiotic treatment (**Figures 5C-F**). Throughout the experiment,

Taconic and Jackson control mice maintained steady viral richness and relative abundance

(**Figures 5C-F**), with Taconic mice harboring a greater number of unique viral genome

fragments compared to Jackson controls (**Figures 5C, E**).

Following microbiota transplantation, the richness and relative abundance of

Taconic-associated viruses in co-housed Jackson mice reached levels similar to Taconic

donors by day 9, with no significant differences observed between these groups (**Figures**

5C,D). Additionally, by day 9, two days post-transplantation, the relative abundance of

Taconic-associated viruses in mice that received a fecal transplant was not significantly

different from the donor community (**Figure 5D**), although the viral community remained

significantly less diverse (**Figure 5C**). However, by day 11, four days after transplantation,

the relative abundance of Taconic-associated viruses decreased in mice that received fecal

or cecal transplants, falling to levels significantly lower than those of Taconic donors

(**Figures 5D, F**). In contrast, the relative abundance of Taconic-associated viruses in co-

housed mice remained relatively stable during this period (Figure 3F). This drop in the

relative abundance of Taconic-associated viruses in the fecal and cecal transplant groups

was accompanied by a slight resurgence of Jackson-associated viruses on day 11 (**Figures**

5D, F). However, this rebound was not sustained, except in Jackson mice that did not

receive any transplantation (JAXVANoT), where Jackson-associated viruses persisted

(**Figures 5D, F**).

Despite only receiving a single gavage of fecal or cecal contents, the number of

unique viral genome fragments in transplanted mice continued to increase, with detectable

levels of Taconic-associated viruses rising even after one week (Figures 3C,E). Although

increasing, the richness of Taconic-associated viruses in Jackson mice that received cecal

or fecal transplants remained significantly different from that of Taconic donors one week

after gavage, on day 14 of the experiment (**Figures 5C, E**).

Meanwhile, viral diversity in Jackson mice that received antibiotics, and no

subsequent transplantation (JAXVANoT) remained depleted throughout the experiment

(through day 14), even though the relative abundance of Jackson-associated viruses

returned to levels similar to those observed in untreated Jackson controls and day 0

samples by day 11, four days after cessation of antibiotics (**Figures 5C-F**). This suggests a

long-lasting impact of antibiotic treatment on the viral diversity in mice that did not receive

microbial transplants, despite the partial recovery of Jackson-associated viral abundance.

These results highlight the complexity of viral community dynamics following

microbiota transplantation, with co-housing proving to be the most effective method for the

rapid restoration of Taconic-associated viral communities. Although fecal and cecal

transplants also facilitated the colonization of Taconic viruses, the process occurred more

slowly and less consistently, particularly in terms of viral richness and abundance.

Discussion

This study builds upon and expands existing literature on microbiota transplantation

by providing a detailed comparison of co-housing, fecal transplantation, and cecal

transplantation as methods for reconstituting bacterial and viral communities in antibiotic-

treated mice[22,23,31,37]. These findings are particularly relevant for improving experimental

methods and enhancing the reproducibility of studies relying on mouse models to investigate

host-microbe interactions. Understanding the strengths and limitations of each transplantation
approach is critical for optimizing experimental designs and ensuring robust results in
microbiome research.

Mouse models are a cornerstone of microbiome research, offering a manipulatable
platform for studying the effects of microbial communities on host physiology. Transplantation
of donor microbiota into recipient mice allows researchers to examine how microbial
composition influences health and disease, yet there remains a surprising lack of systematic
comparisons of commonly used transfer methods. This study addresses this gap by assessing
the efficiency, dynamics, and ecological outcomes of three widely used microbiota transfer
techniques.

Our results demonstrate that co-housing is the most effective method for microbiota
transfer, achieving rapid and complete reconstitution of both bacterial and viral communities.
This efficiency likely arises from sustained microbial exchange through natural behaviors such
as coprophagy and direct contact, which mirror ecological interactions in natural settings.
These findings align with previous studies showing that co-housing facilitates the transfer of
donor-associated microbial taxa more robustly than single-dose gavage methods (e.g., fecal
or cecal transplantation)[31,38]. Importantly, the rapid reconstitution of microbial diversity and
functionality observed in co-housed mice underscores its utility in experimental designs
requiring time-sensitive microbiota establishment.

Fecal and cecal transplantation, while less efficient than co-housing in the short term,
ultimately achieved comparable microbial diversity and functional outcomes by day 21. These
methods may still hold value in experimental scenarios where co-housing is impractical, such
as those requiring strict separation of donor and recipient populations or where there are
insufficient numbers of donor mice. However, the slower pace of microbial establishment with
these methods highlights the importance of considering time frames and endpoints when
designing experiments. Our findings emphasize the need for researchers to select
transplantation methods that align with the specific goals of their studies to optimize
reproducibility and minimize variability.

Mice that did not receive microbial transplantation exhibited delayed and incomplete

recovery of their microbiota, with bacterial and viral communities remaining distinct from donor

or untreated controls even after 21 days. Although incomplete, with reduced microbial

richness, the recovery observed highlights the resiliency of the gut microbiota. Without

transplantation, the limited natural recolonization observed in our study suggests that native

microbial recovery is insufficient to restore microbiota diversity to pre-treatment levels. These

findings are particularly relevant for studies investigating the long-term impacts of microbiota

depletion or the role of microbial diversity in shaping host outcomes, as the choice to include

transplantation can significantly influence experimental results.

The bacterial and viral dynamics observed in this study were closely interdependent,

reflecting the ecological relationships between bacteriophages and their bacterial hosts. Co-

housed mice exhibited rapid establishment of donor-associated bacterial taxa, such as

Muribaculaceae and *Tannerellaceae*, alongside their associated phages. This synchronized

reconstitution underscores the importance of ecological interactions in microbial recovery and

suggests that co-housing provides a more naturalistic environment for microbial community

establishment. Notably, temperate phages demonstrated greater resilience to antibiotic

treatment and were more effectively transferred during co-housing, consistent with their

genomic integration into host bacteria and replication during bacterial colonization.

The site-specific variation in microbial recovery observed across gastrointestinal

regions further highlights the complexity of microbiota reconstitution. While co-housing

facilitated the most uniform recovery across sites, the distinct differences in microbial diversity

and composition between ileal, cecal, and fecal samples reinforce the importance of sampling

multiple gastrointestinal regions to fully capture the dynamics of microbiota transplantation.

These findings are consistent with previous work demonstrating anatomical differences in

microbial communities and their functional capacities.

From an experimental perspective, our study provides critical insights into best

practices for microbiota transplantation in mouse models. Co-housing emerges as the most

effective method for achieving rapid and complete microbial transfer, making it a valuable

approach for experiments requiring a robust and naturalistic establishment of donor

microbiota. However, researchers should weigh the logistical and experimental constraints of

co-housing, such as the need for sustained contact between donor and recipient animals,

against its benefits. Fecal and cecal transplantation, while slower to achieve microbial

reconstitution, remain valuable alternatives for experiments with shorter time frames or

constraints on co-housing feasibility. The results of this study underscore the importance of

tailoring transplantation methods to the specific requirements of individual studies to maximize

reproducibility and reliability.

The results also emphasize the growing need for systematic evaluations of

transplantation methods across different experimental contexts. Future studies should

investigate how factors such as donor-recipient compatibility, microbial complexity, and host

genotype influence the outcomes of microbial transplantation. Additionally, while this study

focused on bacterial and viral communities, future work should incorporate other microbial

taxa, such as fungi and archaea, to provide a more comprehensive understanding of

microbiota dynamics. Advances in functional and longitudinal metagenomic analyses, coupled

with improved animal models, will further enhance our ability to study the mechanisms

underlying microbiota establishment and its implications for host health and disease.

By providing a systematic comparison of microbiota transplantation methods, this

study contributes to the refinement of experimental approaches in microbiome research and

highlights the ecological and functional dynamics underlying microbial reconstitution. These

insights are essential for improving the reproducibility and translational potential of microbiota

studies, ultimately advancing our understanding of the microbiome's role in health and

disease.

Materials and Methods

Mice

WT C57BL/6J (stock no. 000664) mice were purchased from Jackson Laboratories (JAX) and C57BL/6N (stock #B6), murine pathogen free health standard, mice were purchased from Taconic Biosciences (TAC) and maintained at Washington University School of Medicine under specific-pathogen-free conditions according to University guidelines. Female mice, aged six to eight weeks, were exclusively used in all experiments to facilitate cohousing experiments. Experimental mice were co-housed with up to two mice of the same sex per cage with autoclaved standard chow pellets and water provided *ad libitum*.

Cages of female mice were treated at 6 weeks of age with an antibiotic cocktail VNA [1 g/L ampicillin, 1 g/L neomycin, 0.5 g/L vancomycin (Sigma, St. Louis, MO)] in drinking water for 7 days.

16S rRNA gene Illumina sequencing

Phenol:chloroform-extracted DNA from fecal pellets was used for both 16S rRNA gene qPCR and sequencing. SYBR green qPCR for the 16S rRNA gene was performed with 515F (5'-GTGCCAGCMGCCGCGGTAA-3') and 805R (5'-GACTACCAGGGTATCTAATCC-3') primers to detect the V4 hypervariable region.

Primer selection and PCRs were performed as described previously [14]. Briefly, each sample was amplified in triplicate with Golay-barcoded primers specific for the V4 region (F515/R806), combined, and confirmed by gel electrophoresis. PCR reactions contained 18.8µL RNase/DNase-free water, 2.5µL 10X High Fidelity PCR Buffer (Invitrogen), 0.5µL 10 mM dNTPs, 1µL 50 mM MgSO4, 0.5µL each of the forward and reverse primers (10 µM final concentration), 0.1µL Platinum High Fidelity Taq (Invitrogen) and 1.0µL genomic DNA. Reactions were held at 94°C for 2 min to denature the DNA, with amplification proceeding for 26 cycles at 94°C for 15s, 50°C for 30s, and 68°C for 30s; a final extension of 2 min at 68°C was added to ensure complete amplification. Amplicons were pooled and purified with 0.6x Agencourt Ampure XP beads (Beckman-Coulter) according to the manufacturer's instructions.

The final pooled samples, along with aliquots of the three sequencing primers, were sent to

the Center for Genome Sciences (Washington University School of Medicine) for sequencing

by the 2x250bp protocol with the Illumina MiSeq platform.

The resulting FASTQ files were processed using the DADA2 pipeline (v1.28.0)

implemented in Nephele (version 2024_Jan_16, commit: 71b6a72)[32]. Initial quality

assessment was conducted to determine appropriate truncation and filtering parameters.

Reads were trimmed to remove low-quality bases and sequencing primers and truncated

based on quality scores to ensure high-confidence base calls[39,40]. Specifically, reads were

truncated at the position where the quality score dropped below a threshold (default: 4)[41,42].

Reads with expected errors exceeding a set threshold (default: 5) were discarded to further

enhance data quality. Forward and reverse reads were merged, requiring a minimum overlap

of 12bp and allowing for zero mismatches in the overlapping region to ensure accurate

reconstruction of the target amplicon[43,44]. Chimeric sequences were identified and removed

using the consensus method to prevent false-positive operational taxonomic units (OTUs).

Taxonomic classification was performed using the Ribosomal Database Project (RDP)

classifier, with a minimum bootstrap confidence threshold of 40 for taxonomic assignments

and the SILVA v138.1 database[45]. Additionally, a phylogenetic tree was constructed using

MAFFT for multiple sequence alignment and FastTree2 for tree building, facilitating

downstream phylogenetic analyses[46,47]. The output included an amplicon sequence variant

(ASV) table, taxonomic assignments, and phylogenetic relationships, providing a

comprehensive dataset for subsequent diversity and composition analyses. All analyses were

conducted using default parameters unless otherwise specified, adhering to best practices for

16S rRNA gene sequence processing to ensure accurate and reproducible results.

Bacterial ecological and distance analyses

Bacterial community composition and ecological metrics were assessed using both

alpha and beta diversity analyses. Alpha diversity metrics, including observed richness and

Shannon diversity, were calculated to evaluate within-sample diversity. Beta diversity was

assessed using unweighted and weighted UniFrac distances to capture phylogenetic

differences between samples. These analyses were implemented using the 'phyloseq' and

'vegan' R packages, which offer comprehensive tools for ecological and diversity analyses.

Principal coordinate analysis (PCoA) was performed to visualize patterns in beta diversity

across treatment groups and time points[48,49]. Pairwise comparisons between groups were

conducted using PERMANOVA with Bonferroni corrections to identify statistically significant

differences (alpha = 0.05). These analyses provided insights into the ecological shifts in

bacterial communities associated with different transplantation methods and recovery stages.

All figures were generated in R using ggplot2[50].

Identifying bacterial taxa associated with each vendor

Bacterial taxa specific to each vendor were identified using amplicon sequence

variants (ASVs) derived from 16S rRNA gene sequencing. Taxa were classified as Jackson-

associated or Taconic-associated based on their differential prevalence and abundance in

fecal samples collected from untreated mice during the first week of the experiment (days 0-

7). Samples were limited to those without antibiotic treatment to accurately capture baseline

community differences. Differential abundance analyses were conducted using the DESeq2

pipeline, with ASVs assigned taxonomy using the SILVA database. Significant differences

were determined using an adjusted p-value threshold of 0.01 and a combined log2(fold

change) cutoff of ±4. Prevalence was calculated as the proportion of samples in which a given

ASV was detected within each vendor group. To incorporate abundance and prevalence, a

combined fold-change metric was calculated by summing the log2(fold change) from

differential abundance analysis and the log2(prevalence ratio) between the two groups.

Statistical and prevalence-based analyses were compared to confirm associations, and taxa

that did not meet these criteria were categorized as non-supplier-specific.

Predicting functional capacity of bacterial communities

Functional capacities of bacterial communities were inferred from 16S rRNA gene

sequencing data using the PICRUSt2 (v. 2.5.3) pipeline. ASVs were mapped to predicted

genes and metabolic pathways using the MetaCyc database. Differentially abundant pathways

were identified using DESeq2 with an alpha threshold of 0.05. Comparisons were made

between antibiotic-treated recipient mice (co-housed, fecal transplantation, or cecal
transplantation groups) and control communities from untreated Jackson and Taconic mice.
Functional convergence over time was assessed across treatment groups to evaluate the
recovery of microbial metabolic potential.

Taxonomic changes following transplantation

Changes in bacterial taxonomic composition were assessed longitudinally using 16S
rRNA gene sequencing. ASVs were classified into taxonomic families, and relative
abundances were tracked over time in recipient mice across different transplantation methods
(co-housing, fecal, and cecal). Differentially enriched taxa were identified using ANCOM-BC2
(v. 2.6.0), and significant changes were determined using Kruskal-Wallis tests with Bonferroni
corrections for multiple comparisons (alpha = 0.05). Taxonomic dynamics were visualized
using stacked bar plots and principal coordinate analyses (PCoA) of unweighted UniFrac
distances.

Virome Illumina sequencing and analysis.

Viral communities were characterized using short-read Illumina sequencing of viral-like
particles (VLPs) extracted from fecal samples. DNA was purified from VLPs, and libraries were
prepared using standard protocols. Sequencing was performed on an Illumina platform,
generating paired-end reads. Reads were assembled into contigs using Flye, which is
optimized for accurate assembly of viral genomes from metagenomic data. Viral contigs were
identified using stringent criteria for viral hallmark genes and annotated taxonomically using
Cenote-Taker2, a tool designed for comprehensive identification and annotation of eukaryotic
and prokaryotic viruses. Contigs were clustered into operational taxonomic units (OTUs)
representing viral strains, species, and higher taxonomic ranks. Viral richness and relative
abundances were quantified for each treatment group over time, providing insights into virome
dynamics and recovery across experimental conditions.

Viral ecological and distance analyses

Viral community composition and ecological dynamics were assessed using alpha
diversity metrics (e.g., richness) and beta diversity measures (e.g., Jaccard distances). These

analyses were implemented using the 'phyloseq' and 'vegan' R packages, which provide

robust tools for microbial ecological analyses. Principal coordinate analysis (PCoA) was

conducted to visualize temporal changes in viral community structure across treatment

groups. Pairwise comparisons between groups were performed using PERMANOVA with

Bonferroni corrections for multiple comparisons (alpha = 0.05). Viral communities were further

classified as Jackson-associated, Taconic-associated, or non-supplier-specific based on their

prevalence and abundance in control communities. Ecological metrics were integrated with

bacterial data to explore host-virus interactions during microbiota transplantation. Source

discriminatory viral contigs were defined using the combined fold-change metric described for

bacteria above.

Acknowledgments: We acknowledge all members of the Baldridge laboratory for helpful

discussions as well as member of June Round's laboratory.

Funding: This work was supported by the National Institutes of Health (NIH) grants

R01AI139314 and R01AI173360 (M.T.B.), and the Crohn's and Colitis Foundation Litwin IBD

Pioneers Award #1065897 (M.T.B.). E.A.K. was supported by NSF grant DGE-1745038/DGE-

2139839 and NIH grant F31AI167499. J.S.W. was supported by NIH T32AI007172.

Author contributions: J.S.W performed data analysis generated figures and wrote the

manuscript. M.E.S., H.M., S.A., M.H., E.A.K., L.S. and L.F. performed the experiments. R.R.

and M.T.B. analyzed the results. M.E.S., H.M., S.A., and M.T.B. designed the project. L.A.C.C

and A.R contributed analysis of viral data. M.T.B. contributed to the writing of the manuscript.

All authors read and edited the manuscript.

Competing interests: We declare no competing interests.

Data and materials availability: The data from this study are tabulated in the main paper

and supplementary materials. All reagents are available from M.T.B. under a material transfer

agreement with Washington University.

References

<u>References</u>

1. Sommer, F. & Bäckhed, F. The gut microbiota — masters of host development and

physiology. *Nat. Rev. Microbiol.* **11**, 227–238 (2013).

2. Round, J. L. & Mazmanian, S. K. The gut microbiota shapes intestinal immune

responses during health and disease. *Nat. Rev. Immunol.* **9**, 313–323 (2009).

3. Oliphant, K. & Allen-Vercoe, E. Macronutrient metabolism by the human gut microbiome:

major fermentation by-products and their impact on host health. *Microbiome* **7**, 91

(2019).

4. Li, T.-T. *et al.* Microbiota metabolism of intestinal amino acids impacts host nutrient

homeostasis and physiology. *Cell Host Microbe* **32**, 661-675.e10 (2024).

5. Donald, K. & Finlay, B. B. Early-life interactions between the microbiota and immune

system: impact on immune system development and atopic disease. *Nat. Rev. Immunol.*

23, 735–748 (2023).

6. Lambring, C. B. *et al.* Impact of the Microbiome on the Immune System. *Crit. Rev.*

Immunol. **39**, 313–328 (2019).

7. Graham, D. B. & Xavier, R. J. Conditioning of the immune system by the microbiome.

Trends Immunol. **44**, 499–511 (2023).

8. Villarino, N. F. *et al.* Composition of the gut microbiota modulates the severity of malaria.

Proc. Natl. Acad. Sci. U. S. A. **113**, 2235–2240 (2016).

9. Baldridge, M. T. *et al.* Commensal microbes and interferon-λ determine persistence of

enteric murine norovirus infection. *Science* **347**, 266–269 (2015).

10. Winkler, E. S. *et al.* The Intestinal Microbiome Restricts Alphavirus Infection and

Dissemination through a Bile Acid-Type I IFN Signaling Axis. *Cell* **182**, 901-918.e18

(2020).

11. Turnbaugh, P. J. *et al.* An obesity-associated gut microbiome with increased capacity for

energy harvest. *Nature* **444**, 1027–1031 (2006).

12. Ridaura, V. K. *et al.* Gut microbiota from twins discordant for obesity modulate

metabolism in mice. *Science* **341**, 1241214 (2013).

13. Gehrig, J. L. *et al.* Effects of microbiota-directed foods in gnotobiotic animals and

undernourished children. *Science* **365**, eaau4732 (2019).

14. Di Luccia, B. *et al.* Combined Prebiotic and Microbial Intervention Improves Oral Cholera

Vaccination Responses in a Mouse Model of Childhood Undernutrition. *Cell Host*

Microbe **27**, 899-908.e5 (2020).

15. Britton, G. J. *et al.* Defined microbiota transplant restores Th17/RORγt+ regulatory T cell

balance in mice colonized with inflammatory bowel disease microbiotas. *Proc. Natl.*

Acad. Sci. U. S. A. **117**, 21536–21545 (2020).

16. Britton, G. J. *et al.* Microbiotas from Humans with Inflammatory Bowel Disease Alter the

Balance of Gut Th17 and RORγt+ Regulatory T Cells and Exacerbate Colitis in Mice.

Immunity **50**, 212-224.e4 (2019).

17. Kennedy, E. A., King, K. Y. & Baldridge, M. T. Mouse Microbiota Models: Comparing

Germ-Free Mice and Antibiotics Treatment as Tools for Modifying Gut Bacteria. *Front.*

Physiol. **9**, (2018).

18. Witjes, V. M., Boleij, A. & Halffman, W. Reducing versus Embracing Variation as

Strategies for Reproducibility: The Microbiome of Laboratory Mice. *Animals* **10**, 2415

(2020).

19. Schloss, P. D. Identifying and Overcoming Threats to Reproducibility, Replicability,

Robustness, and Generalizability in Microbiome Research. *mBio* **9**,

10.1128/mbio.00525-18 (2018).

20. Huttenhower, C., Finn, R. D. & McHardy, A. C. Challenges and opportunities in sharing

microbiome data and analyses. *Nat. Microbiol.* **8**, 1960–1970 (2023).

21. Moossavi, S., Fehr, K., Khafipour, E. & Azad, M. B. Repeatability and reproducibility

assessment in a large-scale population-based microbiota study: case study on human

milk microbiota. *Microbiome* **9**, 41 (2021).

22. Zhang, C. *et al.* Transfer efficiency and impact on disease phenotype of differing

methods of gut microbiota transfer. *Sci. Rep.* **12**, 19621 (2022).

23. Ericsson, A. C. & Franklin, C. L. Manipulating the Gut Microbiota: Methods and

Challenges. *ILAR J.* **56**, 205–217 (2015).

24. Borin, J. M. *et al.* Fecal virome transplantation is sufficient to alter fecal microbiota and

drive lean and obese body phenotypes in mice. *Gut Microbes* **15**, 2236750 (2023).

25. Rasmussen, T. S. *et al.* Faecal virome transplantation decreases symptoms of type 2

diabetes and obesity in a murine model. *Gut* **69**, 2122–2130 (2020).

26. Federici, S. *et al.* Targeted suppression of human IBD-associated gut microbiota

commensals by phage consortia for treatment of intestinal inflammation. *Cell* **185**, 2879-

2898.e24 (2022).

27. Gogokhia, L. *et al.* Expansion of Bacteriophages Is Linked to Aggravated Intestinal

Inflammation and Colitis. *Cell Host Microbe* **25**, 285-299.e8 (2019).

28. Hsu, B. B. *et al.* Dynamic Modulation of the Gut Microbiota and Metabolome by

Bacteriophages in a Mouse Model. *Cell Host Microbe* **25**, 803-814.e5 (2019).

29. Pfeiffer, J. K. & Virgin, H. W. Transkingdom control of viral infection and immunity in the

mammalian intestine. *Science* **351**, aad5872 (2016).

30. Brunse, A. *et al.* Fecal filtrate transplantation protects against necrotizing enterocolitis.

ISME J. **16**, 686–694 (2022).

31. Caruso, R., Ono, M., Bunker, M. E., Núñez, G. & Inohara, N. Dynamic and Asymmetric

Changes of the Microbial Communities after Cohousing in Laboratory Mice. *Cell Rep.*

27, 3401-3412.e3 (2019).

32. Weber, N. *et al.* Nephele: a cloud platform for simplified, standardized and reproducible

microbiome data analysis. *Bioinforma. Oxf. Engl.* **34**, 1411–1413 (2018).

33. Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon

data. *Nat. Methods* **13**, 581–583 (2016).

34. Douglas, G. M. *et al.* PICRUSt2 for prediction of metagenome functions. *Nat. Biotechnol.*

38, 685–688 (2020).

35. Caspi, R. *et al.* The MetaCyc database of metabolic pathways and enzymes and the

BioCyc collection of pathway/genome databases. *Nucleic Acids Res.* **44**, D471-480

(2016).

36. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

37. Robertson, S. J. *et al.* Comparison of Co-housing and Littermate Methods for Microbiota

Standardization in Mouse Models. *Cell Rep.* **27**, 1910-1919.e2 (2019).

38. Lundberg, R. *et al.* Human microbiota-transplanted C57BL/6 mice and offspring display

reduced establishment of key bacteria and reduced immune stimulation compared to

mouse microbiota-transplantation. *Sci. Rep.* **10**, 7805 (2020).

39. Andrews, S. Babraham Bioinformatics - FastQC A Quality Control tool for High

Throughput Sequence Data. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

40. Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: summarize analysis results

for multiple tools and samples in a single report. *Bioinforma. Oxf. Engl.* **32**, 3047–3048

(2016).

41. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet.journal* **17**, 10–12 (2011).

42. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

43. Magoč, T. & Salzberg, S. L. FLASH: fast length adjustment of short reads to improve

genome assemblies. *Bioinformatics* **27**, 2957–2963 (2011).

44. Streett, D. dstreett/FLASH2. (2025).

45. Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data

processing and web-based tools. *Nucleic Acids Res.* **41**, D590–D596 (2013).

46. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

47. Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2 – Approximately Maximum-

Likelihood Trees for Large Alignments. *PLoS ONE* **5**, e9490 (2010).

48. Oksanen, J. *et al.* vegan: Community Ecology Package. 2.6-8

https://doi.org/10.32614/CRAN.package.vegan (2001).

49. McMurdie, P. J. & Holmes, S. phyloseq: An R Package for Reproducible Interactive

Analysis and Graphics of Microbiome Census Data. *PLOS ONE* **8**, e61217 (2013).

50. Wickham, H. *et al.* ggplot2: Create Elegant Data Visualisations Using the Grammar of

Graphics. (2024).

893