Detection of Iron Protein Supercomplexes in *Pseudomonas aeruginosa* by Native Metalloproteomics

Mak A. Saito[1]* and Matthew R. McIlvin[1]

[1]Marine Chemistry and Geochemistry Department, Woods Hole Oceanographic Institution
Woods Hole MA 02543 USA

*Corresponding author: msaito@whoi.edu
Watson Biogeochemistry Building
360 Woods Hole Rd
Woods Hole MA 02543
USA

Running head: The Iron Metalloproteome of *Pseudomonas aeruginosa*

Abstract

Pseudomonas aeruginosa is a major contributor to human infections and is widely distributed in the environment. Its ability for growth under aerobic and anaerobic conditions provides adaptability to environmental changes and to confront immune responses. We applied high-throughput native 2-dimensional metalloproteomics to *P. aeruginosa* to examine how use of iron within the metallome responds to oxic and anoxic conditions. Metalloproteomic analyses revealed four major iron peaks, each comprised of metalloproteins with synergistic functions, including: 1) respiratory and metabolic enzymes, 2) oxidative stress response enzymes, 3) DNA synthesis and nitrogen assimilation enzymes, and 4) denitrification enzymes and related copper enzymes. Three ferritins co-eluted with the first and third iron peaks, localizing iron storage with these functions. Several metalloenzymes were more abundant at low oxygen, including alkylhydroperoxide reductase C that deactivates organic radicals produced by denitrification, all three classes of ribonucleotide reductases, ferritin (increasing in ratio relative to bacterioferritin), and denitrification enzymes. Superoxide dismutase and homogentisate 1,2-dioxygenase were more abundant at high oxygen. The co-eluting Fe peaks contained multiple iron metalloproteins of varying size, implying the presence protein supercomplexes with related functionality and co-localized iron storage. This coordination with functionality implies cellular organization at the protein complex level optimized for metal trafficking that contributes to efficient iron use and prioritization, particularly for *Pseudomonas* with its large genome and flexible metabolism. This study provides insight into prokaryotic metallome dynamics in response to oxygen availability and demonstrates the capabilities of native metalloproteomic methods in understanding metal use and protein-protein interactions in biological systems.

Graphical Abstract

The iron metalloproteome of *Pseudomonas aeruginosa* was examined using native (non-denaturing) 2-dimensional chromatographic separation coupled to elemental and proteomic mass spectrometries. (A) Four major iron peaks were observed that corresponded to multi-protein complexes associated with respiratory, (B) antioxidant, DNA production, and denitrification functions, and associated iron storage and supply. The results demonstrate the presence of large protein complexes (super complexes) and their potential roles in iron homeostasis and trafficking.

Introduction

The pseudomonad bacterium are a group of widely distributed microbes found in both the environment and as pathogens[1,2]. The species *Pseudomonas aeruginosa* is a major contributor to infections in humans including in wounds and lungs of cystic fibrosis patients, as well as the cause of nosocomial infections such as pneumonia and septicemia[3]. Due to the prevalence of *P. aeruginosa* infections and increasing occurrence of antimicrobial resistance, the World Health Organization has classified it as a priority pathogen and recommended development of new antimicrobial strategies[4]. Furthermore, the role of metals is important in nutritional immunity, where immune cells target the availability of metals required by pathogens[5]. *P. aeruginosa* can also be in nature, within soils and aquatic environments, such as sediments and sinking particles, as part of the microbial community that is decomposing organic matter[6–8].

Pseudomonads are capable of rapid growth, consistent with a copiotrophic lifestyle that allows them to proliferate quickly. Moreover, Pseudomonads are capable of multiple respiratory pathways that allows them to continue growth under suboxic or anoxic conditions, such as by growth on the oxidant nitrate via denitrification[9,10]. When infecting host tissues, *Pseudomonas aeruginosa's* oxygen flexibility is advantageous, where it resists neutrophil cells' oxidative bursts through the use of multiple superoxide dismutases, as well as also actively consuming oxygen to reduce the neutrophils' effectiveness[11]. *Pseudomonas* has a large genome for a bacterium, with over 5000 genes in the core genome[12], and 5570 genes in the genome of PAO1[13], reflecting these dynamic capabilities. Adaptations to acquire metals for nutrition is an important trait of *Pseudomonas aeruginosa,* and *P. aeruginosa* use of iron is particularly complex with regards to acquisition and homeostasis capabilities. As a result, an active area of development for antimicrobials involves drugs that target iron acquisition or trafficking, such antibiotic conjugated to siderophores, or inhibitors for iron release from ferritin. Amongst *P. aeruginosa*'s sophisticated iron capabilities are the biosynthesis capability for two siderophores, pyoverdine and pyochelin, which are fluorescent and give cultures a characteristic visible coloration. *P. aeruginosa* also acquires iron through exogenous siderophores, through a ferrous iron transporter, and when complexed to heme, citrate, and catechols[14]. Metal insufficiencies caused through mutation of transporter systems often result in decreased pathogenicity, demonstrating the importance of metal nutrition in *P. aeruginosa* infections[15] and potential avenues for antibiotic treatment through inhibition of transporters and metal complexation by exogenous metallophores[14,16].

With this array of metal uses within *P. aeruginosa*, understanding the overall composition of this microbe's metallome is of broad interest. The study of metalloproteins often focuses on characterization of individual proteins to provide essential mechanistic insights. In parallel, metallome studies can focus on the overall organismal or tissue metal content by bulk metal analyses and intracellularly localization of metals. Metalloproteomic techniques can complement these approaches by reconstructing the contribution of the composite of metalloproteins to the overall metallome. This approach combines native extraction of metalloproteins and their detection by dual mass spectrometry analyses to determine both metal and protein content simultaneously to assess major metalloproteins inventories within an organism[17–20]. In this study we conducted a metalloproteomic analysis of *P. aeruginosa* grown under aerobic and anaerobic conditions to assess how the iron metalloprotein inventory changes in response to these environmental conditions. A follow-up study will present the results of metals beyond iron in the *P. aeruginosa* metalloproteome.

Methods

Isolation and Cultivation: Liquid media and plates for isolation were made with marine broth media base (Zobell's media), which was prepared from 500 mL of 0.2 μm-filtered coastal seawater (Vineyard Sound)

with the addition of 2.5 g peptone (Fisher Scientific) and 0.5 g yeast extract (BD Difco), with 7.5 g of agar
(for plates only), followed by autoclaving for 40 min. Plates were streaked from 51 micron filter size
fraction collected on a large volume submersible pump from 400m at Station 5 (0°S 202°E) on October
13, 2011 in the Central Pacific Ocean on the METZYME expedition (KM1128) aboard the R/V Kilo Moana
and allowed to incubate at room temperature. Picked colonies from plates grown in liquid media
displayed coloration characteristic of *Pseudomonas aeruginosa*. Once colonies had grown, a colony was
restreaked and frozen at -80 in glycerol and shipped to the laboratory. Experiments were conducted
with strain 2-54. While *P. aeruginosa* is known to be found in natural environments, including coastal
environments, it is not often reported from open ocean (oceanic) environments. The sampling location
was oceanic, although there were islands present in the broad geographic vicinity (Hawaii and Christmas
Island of Kiribati). While this was an attempted isolation from the natural ocean environment, given
genomic results and the lack of *Pseudomonas aeruginosa* observed in co-collected metagenomes and
metaproteomes datasets[21,22], we infer that this isolate was likely a contaminant from the ship or its
personnel during the isolation process.
15
16 *Genomic analyses:* A bacterial isolate was genomic sequenced was sequenced at the Johns Hopkins
17 Deep Sequencing and Microarray Core Facility using Oxford Nanopore and assembled using Canu,
18 Nanopolish, and circulator pipeline. BLAST analyses and resulting annotations determined the microbe
19 to be *Pseudomonas aeruginosa*.
20
21 *Genome Database for Proteomics Analysis:* Given minor differences between PAO1 and our isolate, and
22 the high-quality annotations associated with PAO1, the PAO1 genome was used for proteome
23 annotation. PAO1 was downloaded from NCBI in January of 2019, allowing use of the well-established
24 PA identification numbers.
25
Anoxic Media: Anoxic media (anaerobic) treatments were conducted by autoclaving 1L bottles of media
into followed by cooling, sterile-filtered sodium nitrate was added to a concentration of 0.88mM as an
alternate oxidant to oxygen and enabling denitrification. The media was equilibrated with a loose cap in
an anerobic glove box for several days to allow equilibration to the no oxygen environment. Anoxic
bottles were then inoculated in the glove box, closed tightly, and vacuum heat sealed within a plastic
bag. Oxic cultures were allowed to grow with a loose cap and moderate shaking to allow gas exchange.
Cultures were grown in an incubator at 37°C with continuous moderate orbital mixing. Culture aliquots
were harvested by centrifugation in 50mL centrifuge tubes at 3220 × *g* for 40 min at 4 °C using an
Eppendorf 5810R centrifuge, the media was decanted, and pellets were frozen until analysis. 50mL
aliquots of each treatment were extracted for both metalloproteomics and global proteomics (the latter
to verify metabolic responses to environmental oxygen).
37
Protein analyses
Protein extraction for global proteomics: Pellets were resuspended in 5mL of protein extraction buffer
(50mM HEPES pH 8.5 (Boston BioProducts #BB-2082), 1% SDS in HPLC grade water). All reagents in this
protocol are made with HPLC grade water. Samples were heated at 95°C for 10 minutes and shaken at
room temperature for 30 minutes. Samples were then spun for 30 minutes at 3220 x *g* in an Eppendorf
5810 centrifuge. Supernatant was removed from pellet and transferred to a Vivaspin 5K MWCO
ultrafiltration unit (Sartorius Stedim #VS0611). Protein extract was concentrated to approximately 350
µL, washed with 1mL of lysis buffer and transferred to a 2mL ETOH washed microtube (all tubes from
this point on are ETOH washed). Vivaspins were rinsed with small volumes of protein extraction buffer
to remove all concentrated protein, and all samples were brought up to 400µL.
48

Protein clean up and digestion. SpeedBead Magnetic Carboxylate Modified Particles (GE Healthcare #65152105050250 and #45152105050250) were prepared according to Hughes et al. (2014). 20 µL (20 µg/µL) of magnetic beads were added to 400 µL of extracted protein sample. Samples were heated at 37˚C periodically to avoid precipitation. Samples were acidified to a pH of 2-3 by adding 50 µl of 10% formic acid. 2X volume (1100 µL) of acetonitrile was immediately added. Samples were incubated at 37˚C for 15 minutes and then at room temperature for 30 minutes. Samples were placed on a magnetic rack, incubated for 2 minutes, supernatant was removed and discarded. Samples were washed 2 times removing and discarding supernatants with 1400 µL of 70% ETOH for 30 seconds on the magnetic rack. 1400 µL of acetonitrile was added to each sample for 30 seconds on the magnetic rack. Supernatant was removed and discarded. Samples air dried for approximately 4 minutes until acetonitrile had just evaporated. Samples were removed from the magnetic rack and beads were reconstituted in 90 µL of 50 mM HEPES pH 8.0.

Protein reduction and alkylation. 50 units (2 µL) of benzonase nuclease (Novagen #70746-3) was added to each sample and incubated at 37˚C for 30 minutes. Samples were reduced by adding 20 µL of 200 mM DTT (Fisher #BP172-5) in 50mM HEPES pH 8.5 at 45˚C for 30 minutes. Samples were alkylated by adding 40 µL of 400mM iodoacetamide (Acros #122270050) in HEPES pH 8.5 for 30 minutes at 24˚C, occasionally heating to 37˚C to prevent precipitation. The reaction was quenched by adding 40 µL of 200 mM DTT in 50mM HEPES pH 8.5. Total protein was then quantified according to the method described below (Total protein analysis).

Peptide recovery and preparation. Acetonitrile was added to digested peptides at a concentration of ≥ 95% and incubated for 20 minutes at room temperature. Samples were then placed on the magnetic rack for 2 minutes and the supernatant was removed and discarded. 1400 µL of acetonitrile was added to samples on the magnetic rack for 15 seconds. Supernatant was removed and discarded. Samples were air dried for approximately 4 minutes, until acetonitrile had just evaporated. Beads were reconstituted in 90 µL of 2% DMSO and incubated off the rack at room temperature for ≥ 15 minutes. Samples were centrifuged slowly and briefly at 900 rcf to remove liquid from the tube walls. Samples were incubated on the magnetic rack for 15 minutes and supernatant containing peptides was transferred to a new ETOH washed 1.5 mL microtube. This step was repeated to ensure removal of all magnetic beads. 1% trifluoroacetic acid was added to samples for a final concentration of 0.1%. Samples were zip tipped with Pierce C18 tips (Fisher #87784) according to manufacturer's protocol with a final resuspension in 25 µL of 70% acetonitrile, 0.1% formic acid. Samples were evaporated to approximately 10 µL in a DNA110 Speedvac (ThermoSavant). Samples were finally resuspended to a peptide concentration of 1 µg/µL in buffer B (2% acetonitrile, 01% formic acid).

Total protein analysis. Total protein was quantified protein using 2µL of sample in duplicate and the BCA method (Thermo Scientific Micro BCA Protein Assay Kit #23235). Absorbance was measured on a Nanodrop ND-1000 spectrophotometer (Thermo Scientific). Standard curves were generated using albumin standard (Thermo Scientific #23210). The samples were then digested with trypsin (Promega #V5280) dissolved in HEPES pH 8.0 at a concentration of 0.5 µg/µL was added to samples at a 1:25 trypsin to protein ratio and incubated at 37˚C overnight.

Peptide analysis. Protein extracts were analyzed by liquid chromatography-mass spectrometry (LC-MS) (Michrom Advance HPLC coupled to a Thermo Scientific Fusion Orbitrap mass spectrometer with a Thermo Flex source). 1 µg of each sample (measured before trypsin digestion) was concentrated onto a trap column (0.2 x 10 mm ID, 5 µm particle size, 120 Å pore size, C18 Reprosil-Gold, Dr. Maisch GmbH) and rinsed with 100 µL 0.1% formic acid, 2% acetonitrile (ACN), 97.9% water before gradient elution

through a reverse phase C18 column (0.1 x 250 mm ID, 3 µm particle size, 120 Å pore size, C18 Reprosil-Gold, Dr. Maisch GmbH) at a flow rate of 500 nL/min. The chromatography consisted of a nonlinear 160 min gradient from 5% to 95% buffer B, where A was 0.1% formic acid in water and B was 0.1% formic acid in ACN (all solvents were Fisher Optima grade). The mass spectrometer was set to perform MS scans on the orbitrap (240000 resolution at 200 m/z) with a scan range of 380 m/z to 1580 m/z. MS/MS was performed on the ion trap using data-dependent settings (top speed, dynamic exclusion of 15 seconds, excluding unassigned and singly charged ions, and precursor mass tolerance of ±3ppm, with a maximum injection time of 150 ms).

Proteomics informatics. Mass spectra were searched against *Pseudomonas aeruginosa* PA01 proteome downloaded from NCBI in January of 2019 using Proteome Discoverer 2.0 using the SEQUEST HT algorithm (Thermo) with a parent tolerance of 10 ppm and a fragment tolerance of 0.6 Da. Proteome Discoverer output files were then loaded in Scaffold (Proteome Software) with a protein threshold maximum of 1.0% false discovery rate.

Metalloproteomic Method: 2D Native Separation and Dual Mass Spectrometric Analyses
Protein Separation. Cell pellets from 50 mL centrifuge tubes were thawed in an anaerobic chamber (Coy Laboratory Products Inc.) and suspended in 5 mL of 50 mM TRIS buffer (pH 8.8, all buffers Chelex treated to remove background metals). The solution was transferred to a 15 mL PET plastic centrifuge tube (Fisher 055391) before sonication on ice for 2 min (1 second on/off pulses with a 5 min stop after 1 min, Fisherbrand Model 120 sonic dismembrator). The sonicated pellet was diluted to a total volume of 30 mL in two 15 mL tubes and centrifuged in a gas tight bucket rotor for 60 min at 3220 × g at 5 °C (Eppendorf 5810R). The supernatant was loaded onto a GE HiTrap Q HP anion exchange column (1st dimension, AE hereon) with a peristaltic pump at 0.25 mL/min. The column was then attached to an Agilent 1100 series HPLC pump and eluted with a gradient of sodium chloride at 0.5 mL/min, where buffer A was 50 mM TRIS buffer (pH 8.8), and buffer B was 1 M NaCl in 50 mM Tris buffer (pH 8.8). The gradient was 2 min at buffer A, followed by a linear increase to 60% buffer B over 12 min, then another linear increase to 100% buffer B over 6 min. Eluent was collected with a Bio-Rad 2110 fraction collector in 2 mL microcentrifuge tubes at a rate of 1 fraction every 2 min and stored on ice. Anion exchange column fractions were concentrated using 5000 Da molecular weight cutoff Vivaspin columns from 500 µL to 100 µL before size exclusion separation. Concentrated samples were then injected into a Tosoh Bioscience TSKgel G3000SWxl size exclusion column (2nd dimension, SE hereon) with guard column attached to a Michrom Paradigm MS4 LC system with a CTC Analytics HTC Pal autosampler, and samples were eluted with a 0.5 mL/min isocratic gradient of 50 mM Tris buffer (pH 7.5, 50 mM sodium chloride). Fractions were collected in 1 min intervals into 1.2 mL 96 well plates. 250 µL of each well was transferred to another 96 well plate for metals analysis by liquid handling robot.

Protein analysis of metalloproteins. In preparation for proteomic analyses, separated proteins were reduced, alkylated, and tryptic digested according to the following protocols. 200 µL of each sample was combined with 10 µL acetonitrile and 15 µL of 10 mM dithiothreitol in 100 mM ammonium bicarbonate. The samples were incubated for 30 min at 70 °C while shaking at 450 rpm. After cooling to room temperature, 30 µL of 20 mM iodoacetamide in 100 mM ammonium bicarbonate was added to each well and incubated for 30 min in the dark. 10 µL of 0.03 µg/µL trypsin (Promega Gold) was added to each well and incubated overnight at 37 °C while shaking at 450 rpm.

Trypsin digested size fractionated samples were analyzed using a Michrom Advance HPLC system with reverse phase chromatography coupled to a Thermo Scientific Q-Exactive Orbitrap mass spectrometer with a Michrom Advance CaptiveSpray source. Each sample was concentrated onto a trap column (0.2 x

10 mm ID, 5 µm particle size, 120 Å pore size, C18 Reprosil-Gold, Dr. Maisch GmbH) and rinsed with 100
µL 0.1% formic acid, 2% acetonitrile (ACN), 97.9% water before gradient elution through a reverse phase
C18 column (0.1 x 150 mm ID, 3 µm particle size, 120 Å pore size, C18 Reprosil-Gold, Dr. Maisch GmbH)
at a flow rate of 500 nL/min. The chromatography consisted of a nonlinear 50 min gradient from 5% to
95% buffer B, where A was 0.1% formic acid in water and B was 0.1% formic acid in ACN (all solvents
were Fisher Optima grade). The mass spectrometer monitored MS1 scans from 380 m/z to 1580 m/z at
70K resolution. MS2 scans were performed at 15K resolution on the top 10 ions with an isolation
window of 2.0 m/z and a 10 second exclusion time.

Metals analysis of metalloproteins. 250 µL of each size fractionated sample was combined with 50 µL of
30% hydrogen peroxide (MilliporeSigma Supelco, trace metal grade) and digested overnight. 500 µL of
5% nitric acid containing 1 ppb Indium was then added and digested overnight. Samples were analyzed
on an ICAP-Q inductively coupled plasma-mass spectrometer (Thermo) with an SC-4 DX FAST
autosampler (Elemental Scientific, Inc.). Samples were analyzed in KED mode with helium as the collision
gas. The following metal isotopes were analyzed: Fe 56, P 31, Co 59, Zn 66, Ni60, Cu63, Al27, Ti48, V51,
Cr52, Mn55, Fe57, As75, and Mo95. Metals were calibrated to standard curves prepared the same as
the samples (96 well plates, identical reagents), which were run between each set of anion exchange
fractions.

Comparability between Oxic and Anoxic Metalloproteomes: Efforts were made to make the results
between oxic and anoxic metalloproteomes consistent to allow comparisons. Total protein yields from
the 1st dimension were measured by total protein assay (see Total Protein Analysis above) for both oxic
and anoxic treatments (Figure S1). This was the protein material loaded into the 2nd SE dimension. The
total protein was relatively similar between oxic and anoxic treatments, ranging between 9% and 56%
more in the anoxic treatment in all cases, with the smallest difference occurring at the peak protein
fraction for both consistent with similar protein loadings between each treatment (300-400mM; Table
S1 and Figure S1).

Data analysis
The metals and protein datasets had slightly different ranges of data coverage, as summarized in Table
S1. The 2D data was visualized in 6-8 anion exchange (AE) fractions as the first dimension (100-200mM,
200-300mM, 300-400mM, 400-500mM, 500-600mM, 600-800mM, 800-1000mM, 1000-1000mM, where
the last two ranges were only available in the oxic metals dataset) and 38 size exclusion (SEC) fractions
(fractions 1-38) in the second dimension for a grid of 304 samples. Two initial AE fractions containing
largely dead volume (0 and 0-100mM) were not analyzed for metals or proteins. AE values were plotted
as the lower value of their AE ranges (100, 200, 300, 400, 500, 600, 800, 1000). Metals and proteins
were analyzed from a subset of SEC fractions, 7-38 for metals and 10-35 for proteins, resulting in ~312
proteomic analyses (156 per treatment) and ~448 metal analyses (256 and 192 for oxic and anoxic,
respectively). The four datasets were organized in a uniform metalloproteome data template with
consistent formatting of dataframes including empty fields where no data was present (since the extent
of metal and protein coverage varied) to allow import into the Python notebook.

Informatics Methods
Data analysis and visualization were conducted by importing CSV files (see Data Availability section) into
a customized Metalloproteomic Viewer written in Python (Version 3.10.9 Anaconda package) using
Matplotlib and Plotly packages to create visualizations, identify maxima, and conduct optimization
fittings and implemented as a Jupyter notebook (Version 7.2.2). Code for the Metalloproteomic-Viewer
is available on GitHub at (see Data Availability section). The notebook workflow consisted of identifying

major metal peaks, identifying the proteins with maxima on or adjacent to the metal peak, identification of known metalloproteins within the list by searching using Copilot AI. For Fe Peak 1, the contributions of proteins to the iron peak in the SE dimension was estimated using a local search optimization algorithm (L-BFGS-B; or Broyden, Fletcher, Goldfarb, and Shannon optimization) and minimize function in Python). Protein structures and putative protein-protein interactions were created using Alphafold3.[23]

Results and Discussion

Global Proteomics

Cultures grown under aerobic and anoxic conditions were screened using proteomics to verify acclimation to each treatment prior to metalloproteome analyses. These results are briefly summarized here, with a more detailed proteomic analysis of the influence of oxygen on *P. aeruginosa* to be presented in a future study. The proteomic results confirmed acclimation to the presence or absence of oxygen, with characteristic proteins showing changes in protein abundance in response to O_2, including the denitrification pathway, which contain numerous metalloenzymes involving Fe, Cu and other metals (the nitrate, nitrite, nitrous, nitric oxide reductases; Figure 1). A number of other metal-related proteins also were more abundant under anoxic conditions including molybdopterin biosynthesis proteins, cobalt chelatase CobN from cobalamin biosynthesis pathway[24], and the CopR copper regulator[25]. Anaerobic molybdopterin biosynthesis is consistent with Mo use within nitrate reductase.

The cobalt chelatase CobN, a component of the B_{12} biosynthetic pathway, increased in abundance under anoxic conditions. This contrasted a prior model that proposed aerobic B_{12} production, secretion and diffusion into anaerobic layers and uptake[26]. Because *P. aeruginosa* only has the oxygen-requiring enzymes for cobalt insertion in B_{12} biosynthesis (CobNST; as opposed to the oxygen independent CbiK/X found in other microbes), Crespo et al. proposed a model where B_{12} is produced aerobically in the upper biofilm layer, then must be secreted and diffuse to inner anaerobic layers where it is used for NrdJ (the B_{12}-requiring for of the enzyme); and once depleted the Fe RNR-III (NrdD) is used below[26]. Ribonucleotide reductase enzymes also showed large changes in protein abundance between aerobic and anaerobic conditions (Figure 1; also see Fe Peak 3 section)[26]. Under aerobic conditions Class I NrdAB proteins were most abundant, with less of the B_{12}-requiring class II NrdJab, and no detectable signal for the class III NrdD. Under anaerobic conditions, all three classes of RNR's were present and in higher abundance than in oxic treatments (Figure 1). Similar to CobN, these results also contrast with the prior model, where a cascade of expression from NrdA to NrdJ to NrdG was expected to coincide with decreasing B_{12} availability[26]. Our experiments were conducted with mild orbital shaking, which discouraged accumulation of biofilms and may contribute to the differences with the prior study.

Overall, the results of the global proteomic survey demonstrated clear differences between oxic and anoxic treatments, particularly with regards to the denitrification apparatus. As a result, metalloproteomic analyses were conducted to determine changes in the metallome under oxic and anoxic conditions.

Metalloproteomics Overview

Metalloproteomic analyses were conducted on sample splits from the oxic and anoxic treatments described above in the global proteomic results. This study focuses primarily on the iron metalloproteome, with the additional contextual Cu data. A future study will examine additional metals in the metalloproteome of *P. aeruginosa*.

The *P. aeruginosa* iron metalloproteome had four major peaks, labeled Fe Peaks 1-4 hereon, within the native 2-dimensional space created by the 1st dimension of anion exchange (AE) and 2nd dimension of size exclusion (SE) (Figures 2 and 3, Figure S2, Table 1). Notably the protein content of these Fe peaks was associated with major cellular functions, with Fe Peak 1 containing the oxic

respiratory system and the bulk of iron storage. Fe Peak 2 was close in proximity to Fe Peak 1 and contained antioxidant functions. Fe Peak 3 was located higher in the AE dimension and contained enzymes associated with DNA synthesis and nitrogen use. Finally, Fe Peak 4 was found at the low end of the AE dimension and contained proteins associated with the denitrification capabilities. Notably Fe Peaks 3 and 4 showed increased Fe and protein content under the anoxic condition. The following sections describe the contents of the four major iron peaks, followed by a section about iron storage capabilities. A list of metalloproteins within each Fe Peak is provided in Table 2.

While the metalloproteomic methodology was developed with the intention of maintaining the native metal-protein relationship, the results presented here also support notion that the native extraction methodology preserve some multi-protein complexes (MPCs). As a result, the co-elution of multiple iron (and other) proteins within the four iron peaks described below may be neither coincidental nor due to limited chromatographic resolution, but remnants of *in vivo* protein-protein interactions. The final section of this manuscript briefly discusses this potential useful aspect of the metalloproteomic method.

Fe Peak 1

The largest peak (Fe Peak 1) was located at AE-SE location 400-14 in oxic conditions and 400-15 in anoxic conditions (Figure 2, Table 1). Six iron proteins co-eluted within this region (Figure 4, labeled 1a-1f below), aconitase (annotated as aconitate hydratase 2 PA1787), homogentisate 1,2-dioxygenase (HmgA, PA2009): and hydrazone dehydrogenase (HdhA, PA4022), catalase (PA02360), and two ferritins (annotated as bacterial ferritin PA4235 and bacterioferritin PA3531).

Fe Peak 1a: The key respiratory protein aconitase was the most abundant iron protein with a large and very broad peak in both oxic and anoxic datasets that extended through multiple anion exchange fractions (AE 300-500 along SE 16 in oxic and AE 300-600 along SE 16-17; Figures 3C, 3D, 4A, 4B), reaching 173 and 228 spectral counts at each AE 400 maxima, respectively. However, both aconitase maxima were offset from the Fe Peak 1 maxima (along AE 400): under oxic conditions aconitase was at SE 16 versus Fe Peak 1 at SE 14, while under anoxic conditions aconitase was at SE 17 versus Fe Peak 1 at SE 15. There is a shoulder in Fe Peak 1 towards higher SE fractions, and the aconitase offset likely contributes to this. A similar offset of aconitase was observed in the metalloproteome of the marine bacterium *Pseudoalteromonas*, where a large second peak that eluted at a larger size fraction was offset from one that aligned with iron, and was interpreted as a potential apo form[19]. Given the breadth of aconitase elution in the *P. aeruginosa* metalloproteome, it also seems feasible that the metalated form occurs at the lower SE range, where the bacterioferritin and ferritin co-eluted (Figure 4C and 4D). The breadth of this peak could also reflect the extent of protein-protein interactions and resulting MPCs, where ferritin proteins can provide the source of iron to aconitase and other metalloproteins in Fe Peak 1 (see last section).

Fe Peak 1b: Another enzyme within Fe Peak 1, homogentisate 1,2-dioxygenase (PA2009, HmgA), was over 11-fold more abundant in the oxic treatment relative to the anoxic treatment (maximum of 67 and 6 spectral counts, respectively). This enzyme is involved in the degradation of aromatic rings including tyrosine and phenylalanine to produce the metabolite homogentisate (2,5-dihydroxyphenylacetate, HG), and requires oxygen for activity. It is a homo-6mer with an iron cupin site, where the cupin β-strands have a characteristic metal binding motif[27]. HmgA was observed to be one of the few genes downregulated in the hypervirulent Australian *P. aeruginosa* clinical strain AES-1 under chronic infection cystic fibrosis patients, relative to acute infection systems, and hence its regulation has been described as an important component of *P. aeruginosa's* adaptation to the cystic fibrosis lung environment[28]. HmgA is also important in humans: a mutation to HmgA causes the rare genetic disease alkaptonuria, which causes an accumulation of homogentisic acid, due to the inability to break down tyrosine and phenylalanine, which polymerizes inside the body and as a dark pigmentation and causing

severe arthropathy[29]. HmgA's abundance within the largest iron peak and responsiveness to oxygen in

this study is consistent with these prior studies and implies that if HmgA in the AES-1 strain responds

similarly to oxygen as observed here, then hypoxic conditions would be characteristic of the cystic

fibrosis lung microbiome environment, and the decreased expression of HmgA under those conditions

could enable a shift of iron to other, non-oxygen requiring functions (see below and Fe Peak 4).[30]

Fe Peak 1c: The iron enzyme hydrazone dehydrogenase HdhA (PA4022) also co-eluted with Peak

1, and transitions from being a minor peak in the oxic conditions to a major peak that also aligns with

the iron maxima in the anoxic conditions. HdhA transforms hydrazones, molecules with a C=N-N moiety,

to hydrazides and acids. HdhA uses NAD^+ as the oxidant to for the oxidative degradation of hydrazone

molecules which PAO can grow on as a carbon source[31]. Prior studies observed that PAO1 grown on the

hydrazone adipic acid bis(ethylidene hydrazide) as its carbon source upregulated *hdhA*, and mutants of

the gene had depressed growth[31]. It is unclear to us why HdhA would be more abundant under anoxic

conditions, since NAD^+, which is regenerated by oxygen, is used by both aconitase and HdhA, but

perhaps due to increased availability of hydrazones under anoxic and denitrifying conditions.

Fe Peak 1 was four times larger in anoxic conditions compared to oxic conditions, despite only

having 9% more total protein used in the metallome and proteome analyses (Table S1). The alignment of

HdhA with Fe peak 1 and HdhA's increased abundance under anoxic conditions implies it contributes to

the larger anoxic Fe Peak 1.

Comparing this with the prior HmgA, both proteins co-eluted at Fe Peak 1, but one protein was

more abundant under oxic conditions (HmgA) and the other under anoxic conditions (HdhA), implying a

shifting influence of proteins contributing to Fe Peak 1 with the change in oxygen availability.

Fe Peak 1d: The Fe metalloprotein catalase (PA0236) also eluted within Fe Peak 1 in both oxic

and anoxic conditions. This enzyme is responsible for the deactivation of hydrogen peroxide. With

aconitase being highly sensitive to reactive oxygen species that cause degradation of its iron-sulfur

clusters[32], the close proximity of these two enzymes within Fe Peak 1 would be beneficial.

Fe Peaks 1e and 1f: Two ferritin iron storage proteins contributed to Fe Peak 1, and a

comprehensive discussion of ferritins in the *P. aeruginosa* metalloproteome will be discussed in a later

section.

Optimization of contributions to Fe Peak 1: quantitative reconstruction of Fe Peak 1 was

attempted using an optimization approach (see Methods) along the AE 400 1st dimension for both oxic

and anoxic conditions (Figure 5, Table 3). Notably, the double peak of homogentisate 1,2-dioxygenase

(HmgA, PA2009), where the second peak may have been apo based on lack of iron detected there,

results in it being penalized in the optimization. When the second HmgA peak was manually adjusted to

have zero protein spectral counts prior to optimization to test this notion of it having an apo form, it

became a significant component to Fe Peak 1 (Table 3; Figure S3).

Given the multiple co-elution of iron metalloproteins within Fe Peak 1 (and other Fe Peaks, see

below) in *P. aeruginosa*, the application of optimization approaches was challenging. Indeed, instead of

independent metalloproteins that eluted uniquely based on their native charge and size properties as

assumed in the optimization, there may be protein-protein associations creating co-dependencies

between metalloproteins negating assumptions of their independent co-elution. While this makes

teasing apart the contribution of multiple iron proteins to each major iron peak difficult, there are

potentially important biological implications of these co-elutions (see last section on supercomplexes).

Summary of Fe Peak 1: Fe Peak 1 appeared to be dominated by metalloproteins associated use

of catabolism of carbon substrates and respiratory processes, such as via the Citric Acid cycle

(aconitase), or degradation of aromatic rings (homogentisate 1,2-dioxygenase) under oxic conditions or

hydrazones (hydrazone dehydrogenase) under anoxic conditions. In addition, catalase and iron storage

proteins were both present, protecting Fe-S clusters (in aconitase), and as a ready supplier of iron.

Fe Peak 2

The second largest iron peak within *Pseudomonas* was at 300-17 under oxic conditions and 300-18 (Figure 2, Table 1). Several iron proteins related to antioxidant activity had maxima associated with Fe Peak 2: the iron superoxide dismutase (SodB, PA4366), two copies of alkylhydroperoxide reductase C (PA3529 and PA0139) and quinone oxidoreductase (PA0023; Figure 6). Fe Peak 2 was also 'diagonally' adjacent to Fe Peak 1 described above and shared overlap of protein features: within oxic Fe peak 2 there was a shoulder towards the lower SE fractions, caused by a second peak at 300-15 that is likely associated with the aconitate hydratase 2 protein (PA1787; Fe Peak 1a) and its broad elution pattern mentioned above that continues at 300-16 (Figure 2, Figure 3A, 3E) and that contributes to both the AE 300 and AE 400 dimensions.

Fe Peak 2a: In *P. aeruginosa*, FeSOD (SodB, PA4366) has been reported to be produced constitutively, including under anaerobic conditions, while the MnSOD (SodA, PA 4468) is only produced under oxic stationary phase conditions[33,34]. These SODs, with catalase, contribute to defense against oxidative bursts from host neutrophil cells[11]. Our results are consistent with these prior findings, where the MnSOD was not identified in the oxic or anoxic metalloproteomes (see supplemental datasets within Data Availability section), which were both harvested in exponential growth, and the FeSOD was abundant in both treatments. Moreover, the Mn maximum was not aligned with the FeSOD implying no substitution in this SOD isoform (Figure 6G and 6H).

Fe Peak 2b and 2c: The two copies of alkylhydroperoxide reductase C (AhpC; PA3529 and PA0139) also co-elute in the 2nd largest iron peak (Figures 6A and 6B). These sequences were checked using METATRYP software and do not share any tryptic peptides, meaning their proteomic identifications in the metalloproteome datasets here were independent of each other[35]. This enzyme has an important role in the physiology of *P. aeruginosa*: AhpC contributes to the cell's antioxidant capability by removing organic peroxides such as peroxynitrite and is part of the 2-Cys peroxiredoxin family. As a result AhpC has been described as the "most widely distributed" reactive nitrogen intermediate (RNI) resistance gene[36]. AhpC is also the primary scavenger of hydrogen peroxide in *E. coli*, where catalase mutants rely on this enzyme for anti-peroxide activity[37]. Hare et al.[38] observed the PA3529 alkylhydroperoxide reductase C to have increased abundance under peroxide stress treatments in *P. aeruginosa* strain PAO1, in addition to positive responses from superoxide dismutase and catalase. PA3529 immediately precedes bacterioferritin-associated ferredoxin Bfd (PA3530) and bacterioferritin (PA353) in the genome, implying co-regulation of this gene neighborhood. AhpC also exhibits chaperone activity, preventing misfolding and thermal aggregation of proteins[39]. This dual function of AhpC is thought to convey its adaptability to oxidative and environmental stresses[39]. AhpC exists as either a dimer or a decamer in physiological solutions.

Both AhpC proteins were present in this metalloproteomic analysis contributing to Fe Peak 2, with PA3529 having higher spectral counts than PA0139 in both oxic and anoxic conditions (Figure 6A, 6B). One notable difference between conditions is that PA3529 goes from being the 3rd most abundant iron protein in Fe Peak 2 after superoxide dismutase and aconitase under oxic conditions (where it was tied with aconitase at the iron maximum with 96 spectral counts), to the *primary* iron protein under anoxic conditions at 319 spectral counts. The minor AhpC (PA0139) also increased under anoxic conditions from 33 to 52 spectral counts. Both AhpC proteins were quite minor constituents to the adjacent Fe Peak 1, with under 18 or fewer spectral counts for oxic conditions, and 9 or less spectral counts for anoxic conditions offset from the peak 1 maximum (centered between SE 11-13 in both cases). Given the switch to nitrate-based respiration in the anoxic treatment, the increased abundances of both AhpC's and their contribution to the overall metallome was likely due to the increased production of organic peroxides like peroxynitrite, and the need to deactivate them to prevent toxicity. Given the prevalence of hypoxic conditions within *P. aeruginosa* infections and its increased abundances, inhibitors to the AhpC could be a potential antimicrobial therapeutic target[36].

Fe Peak 2d: Quinone oxidoreductase is an antioxidant enzyme catalyzes the reduction of quinones preventing the formation of semiquinones that generate reactive oxygen species.[40] In bacteria the enzyme plays a critical role in the electron transport chain, catalyzing the transfer of electrons from NADH to quinone, and contains multiple iron-sulfur clusters[41]. Interestingly, NAD(P)H quinone oxidoreductase is often highly expressed in cancers, and hence is considered a drug target[42].

Summary of Fe Peak 2: Overall, Fe Peak 2 displayed key antioxidant capabilities (SodB and AhpC) being deployed both under oxic and anoxic conditions (AhpC for reactive nitrogen intermediates) in close proximity (in native chromatographic space) to respiratory systems of Fe Peak 1 and iron storage systems.

Fe Peak 3

The third largest iron peak in the *P. aeruginosa* metalloproteome occurred in the 1^{st} dimension AE 600 mM and AE 500mM fractions. Here we consider oxic peak AE 600-SE 18 and anoxic peak AE 600-SE 14 both as Fe Peak 3 given their relative proximity in the iron visualizations and some shared proteins (Table 1, Figures 2, 3, 7A-F; note Figure 2 has expanded AE range to AE 1000 which includes Fe Peak 3). Notably Fe peak 3 was 5-fold larger in the anoxic treatment than the oxic treatment (comparing across the AE 600 dimension). The total protein in the anoxic 600mM AE fraction was only 23.1% larger than the oxic (Table S1), not enough to account for the 5-fold Fe difference, hence we interpret this as a biological signal. After discussion of a technical issue, the identification of metalloproteins within oxic and anoxic treatments of Fe Peak 3 are briefly outlined (Table S2, also described as Fe Peak 3a-k), followed by discussion of selected proteins and their biological role across both treatments, and overall assessment of Fe Peak 3's metabolic roles.

There was a divot in overall signal intensity of protein abundances at anoxic Fe enzymes at AE600-SE14 (Figure 7B, D) where the iron maxima occurred. This divot was not consistent with protein data points on either side of its chromatographic elution, resulting in a 'missing peaks' for those proteins. The cause of the divot appears to be an instrument or sample processing issue rather than representing an actual biological signal, as evidenced by the overall mass spectrometry ion count signal in the sample being $1/10^{th}$ that of the surrounding samples in the raw data files (raw data available within supplemental raw datasets submitted to PRIDE). To provide redundancy of perspective for the missing peaks, we also examined for overlap of proteins and iron with the adjacent AE 500 dimension, where there was an Fe peak at SE 15 for both, relatively close to the anoxic AE600-SE14, and a bit more removed from the oxic AE600-SE18. As a result, AE 500mM is included in the Fe Peak 3 discussion here (Figure 7G-J, Table S1).

Proteins in Oxic Treatment: At oxic AE 600-18 there were no proteins with maxima that co-eluted at that precise location. Of all 33 proteins identified at that location, only three were known iron proteins (Figure 7A, C, E): glutamine synthase (PA5119), aconitate hydratase 2 (PA1787), and L-cysteine desulfurase (pyridoxal phosphate-dependent, PA3814). As mentioned above, the oxic Fe Peak 3 signal was much 5-fold smaller than the anoxic treatment.

Proteins in Anoxic Treatment: At anoxic 600-14, there were also no proteins with maxima at this precise location, although this was likely caused by the signal divot mentioned above. Of the 59 proteins detected at that location, the three iron proteins found in oxic AE 600-18 were present here, as well as additional the potential iron proteins (Figure 7B, E, F): aconitate hydratase 1 (PA1562), nitrite reductase precursor (PA0519), electron transfer flavoprotein alpha-subunit (PA2951), DNA-binding protein from starved cells, Dps (PA0962), and ferric uptake regulation protein (PA4764), NrdA, catalytic component of class Ia ribonucleotide reductase (PA1156) and NrdB, tyrosyl radical-harboring component of class Ia ribonucleotide reductase (PA1155), class III (anaerobic) ribonucleoside-triphosphate reductase subunit, NrdD (PA 1920). There were 8 unannotated proteins that eluted with maxima at 600-18 under anoxic conditions (PA3309, PA2765, PA0916, PA2817, PA3982, PA2151, PA3850, PA0457).

Fe Peak 3a: Glutamine synthetase (GS) was the most abundant enzyme in both oxic and anoxic treatments on the Fe Peak 3 (along the AE600 dimension, Figure 7A and B), yet the GS peak maxima was offset from the iron peak in the size exclusion spectrum at SE 13 in both treatments, implying that perhaps the apo form may have been associated with a larger protein complex. This enzyme is important in nitrogen metabolism that converts glutamate and ammonia into glutamine[43].

Fe Peak 3b and 3c: The ribonucleotide reductase subunits NrdAB were the next most abundant iron metalloproteins in the anoxic AE500 and AE 600 dimensions (Figure 7C, H), and the AE 500, but the AE 600 dimensions of the oxic treatment (Figure 7C, G). The ribonucleotide reductases are a key enzyme in DNA replication by providing the necessary dNTP substrates. *P. aeruginosa* is unusual in that it contains all three classes of ribonucleotide reductases within their genomes (RNR I-III)[44,45]. Most RNRs are metalloenzymes, with RNR-I (NrdAB) subclasses having Fe-Fe, Fe-Mn, Mn-Mn, or no metal active sites, RNR-II (NrdJab) requiring B_{12}, and RNR-III (NrdDE) requiring two Fe atoms[45–47]. Notably, mutants in the anaerobic RNRs (Class II and III) having deficient biofilm formation[15] and impaired anaerobic growth and virulence[16]. The factors necessitating this triple redundancy in *Pseudomonas* are thought to be related to hypoxia and are not well understood.

The observation that NrdB shared a maxima with iron in both oxic and anoxic treatments of AE 500 implied this ribonucleotide reductase isoform was a major contributor to the Fe signal at this site, which is known to have 3 Fe atoms per NrdB polypeptide[48]. Interestingly, the distribution of the NrdA subunit overlapped but with a much wider distribution (SE 12-30 in oxic, Figure 7G, and SE12-35 in anoxic, Figure 7H) and had different maxima than NrdB, implying the NrdAB complex may not have been stoichiometrically assembled throughout the dimension. The alignment of NrdB with the Fe Peak in the AE 500 dimension was consistent that the divot in anoxic AE600 SE14 being an artifact of sample processing. The three most abundant iron proteins in this dimension, NrdA, NrdB, and glutamine synthetase, all show adjacent points that imply higher protein abundance if extrapolated between them (e.g. between SE13 and SE15). When linearly extrapolated these three proteins would be the major proteins with maxima close to or potentially on the Fe Peak 3 site (SE 14), implying their contribution to the iron demand here.

The increased contribution of ribonucleotide reductase to Fe Peak 3 was also consistent with the global proteome results (Figures 1 and 8). In our global experiment, the NrdAB is the dominant ribonucleotide reductase under oxic conditions, as expected from prior studies[26]. Under aerobic conditions Class I NrdAB proteins were most abundant, with less of the class II NrdJab, and no observed class III NrdD. Under anaerobic conditions all three classes of RNR's were present and in higher abundance than in oxic treatments. Similar to CobN, these results also conflict with the model proposed by Crespo et al. (2018), where a cascade of expression from NrdA to NrdJ to NrdG was expected to coincide with decreasing B_{12} availability. Together these results demonstrate a greater role for RNR under anaerobic conditions in *P. aeruginosa* and suggests this enzyme or its cofactors (Fe or B_{12}), could be potential antimicrobial targets.

The major presence of RNR within the iron metalloproteome and the observation of the triple redundancy of RNR classes present in the proteome under a variety of redox conditions can be attributed to *P. aeruginosa*'s need to maintain high growth rates and corresponding DNA synthesis rates under a variety of oxygen and micronutritional conditions.

Fe Peak 3d: A third ferritin annotated as "probably bacterioferritin" (PA4880) was present in the anoxic treatment adjacent to this location with a maximum at SE 18 in the AE 600 dimension (Figure 7D). The ferritins are discussed in greater detail below.

Summary of Fe Peak 3: The iron within this peak appeared to be comprised from nitrogen metabolism and DNA synthesis metalloenzymes, as well as co-located iron storage bacterioferritin.

Fe Peak 4

Although a relatively minor peak compared to Fe Peaks 1 and 2, the fourth largest peak within the *P. aeruginosa* metalloproteome contained key components of the denitrification apparatus. This peak was found in the AE 200 dimension, at SE-20 and SE-22 in oxic and anoxic treatments, respectively (Table 1, Figures 2 and 3). Similar to Peak 3, the peak was much more pronounced in the anoxic treatment compared to the oxic peak with 7.0 times more Fe in the anoxic at their respective maximum locations (Figures 9G, H). This was large Fe difference occurred despite the anoxic treatment having only 9.2% more total protein (Table S1). Yet Fe Peak 4 was still the smallest of the four iron peaks, with Fe Peak 1 being 5.5 larger in the anoxic treatment. Fe Peak 4 in the oxic treatment was particularly minor, approaching background sounding baseline levels, consistent with minimal need for denitrification apparatus under oxic conditions. Notably, the Fe Peak 4 is quite broad and the known iron metalloproteins within it range from SE 18 to SE 32 (Figures 9B, D, F, and H). The denitrification system is known to be a membrane localized protein complex. Despite the high abundance of some of these proteins, the iron peak was smaller than the other three Fe peaks as mentioned above. This may be a result of underrecovery of the denitrification membrane proteins in the native, non-detergent, system.

Proteins in the Anoxic Treatment: A total of 148 proteins were detected at the anoxic maximum of Fe Peak 4, of those 18 proteins shared a maximum at that location (AE 200 SE-22, Table S2). Of those 18, NirN (PA0509) was the only known iron metalloprotein present and 11 of those were annotated as hypothetical proteins (Figures 9B, D, F). Among the remaining proteins detected at SE-22 (not just having a maxima), 9 were known iron proteins, including components of the denitrification system (Figures 9B, D, F).

Proteins in the Oxic Treatment: 162 proteins were detected at the oxic maximum of Fe Peak 4, with 27 sharing a maximum at that location (AE 200 SE-20, Table S2). Of those 27 proteins with a shared maxima none were known iron proteins and 14 were annotated as hypothetical proteins. Among the proteins detected at SE-20 (Figure 9A, C, E), four were known iron proteins, all of which were also on the list of iron proteins detected in the anoxic peak (Table S2), including nitrite reductase precursor (PA0519), cytochrome c4 precursor (PA5490), probable iron-binding protein IscU (PA3813), and cytochrome c551 peroxidase precursor (PA4587).

Fe Peak 4a-c : Under anoxic and added nitrate media conditions *Pseudomonas* utilizes denitrification-based respiration. A number of iron proteins are enlisted including NirS (4a: PA0519 nitrite reductase precursor), NapA (4b: periplasmic nitrate reductase protein NapA PA1174), and NirN (4c: PA0509). NirN is a NirS homologue, and forms a 8 blade β-propeller with a cytochrome c domain[49].

Nitrite reductase (NirS, PA0519) was highly abundant in the anoxic treatment. With reaching 475 spectral counts at its maximum, it was the 2nd most protein in the metalloproteomic dataset. While its peak is somewhat offset from the Fe/Cu peak by two SE fractions it still was at 250 spectral counts on the Fe maximum (Figures 9B, J, I).

Azurin Cu protein: The copper containing azurin enzyme (PA4922) is an electron donor to nitrite reductase (NirS)[50], and the residues involved in their protein-protein interaction have been identified characterized[51]. Here we observed the Cu peak in the AE 200 dimension to co-elute with Fe Peak 4 in both oxic and anoxic treatments (Figures 9I, J). This protein was also one of the most abundant proteins in the anoxic metalloproteome. While it was present with 30 spectral counts at the maximum in AE 200, it was highly abundant in AE 100 reaching 449 spectral counts at 100-22, where few other proteins elute.

Fe Peak 4d: IscU is an Fe-S cluster assembly scaffold, and part of the ISC (Iron Sulfur Cluster) biosynthetic system.[52] Its iron signal may be due to the presence of Fe-S within the protein scaffold. Unlike *E. coli* which has the SUF and ISC Fe-S biosynthesis systems, *P. aeruginosa* only has the ISC system and is essential for viable *Pseudomonas*, has been proposed as a target for antibiotic treatment, although the human mitochondria use a homolog of the system that could be sensitive to treatments.

Fe Peak 4e: Corresponding to the occurrence of five cytochromes in this area (see below), a
heme-transport protein, PhuT, (PA4708) also co-eluted, consistent with the need to assemble the
prosthetic groups into the cytochromes. PhuS in *P. aeruginosa* has been shown to contribute heme to
heme oxygenase and to in sensing and maintaining iron homeostasis.[53] PhuT may have a role in
contributing to denitrification cytochromes.
Fe Peak 4f-j: Five cytochrome proteins were observed in this range, including cytochrome c4
precursor, cytochrome c-551 precursor, cytochrome c551 peroxidase precursor, cytochrome c5, and
cytochrome c-type protein NapB precursor (PA5490, PA0518, PA4587, PA5300, PA1173). Cytochrome
proteins help to conduct electrons, they contain the prosthetic group heme, which coordinates an Fe
atom at its center. The co-elution of these five cytochromes at the Fe Peak 4 was consistent with their
involvement in the denitrification system, and a cytochrome from *Thermus thermophilus* has been
proposed to be part of a super complex with nitrate reductase[54].
Fe Peak 4 - Non-Fe metalloproteins: In addition to azurin two additional non-iron
metalloproteins were detected at Fe Peak 4, nitrous-oxide reductase precursor (PA3392, Cu, anoxic
only), probable glutathione peroxidase (PA0838, Se, oxic and anoxic).
Summary of Fe Peak 4: This peak was dominated by denitrification metabolism metalloenzymes
and associated Fe-S cluster biosynthesis and cytochrome electron transport capabilities.
18
Iron Storage Proteins – Bacterioferritin and Ferritin
Given the importance of iron for the variety of metalloenzymes and biochemical functions
described above, the ability to store iron when excess is available and to prevent intracellular toxicity is
important for bacterial cells. Ferritins are proteins whose 12 or 24 subunits combine to form a protein
sphere that stores as many as ~4000 iron atoms per complex as iron oxide. Bacterioferritins are bacterial
versions of ferritin that also differ Eukaryotic ferritins in containing heme between subunits[55]. In a prior
application of metalloproteomics on the bacterium *Pseudoalteromonas*, iron storage within two copies
of bacterioferritin represented the dominant reservoir of iron within that microbe, indicating a
substantial storage capability[19]. Within *P. aeruginosa's* genome (PAO1) there are three annotated
ferritin genes, a bacterioferritin (BfrB, PA3531), a "bacterial ferritin" (FtnA, PA4235), and a probable
bacterioferritin (PA4880). There is also a bacterioferritin comigratory protein in the PAO1 genome
(PA1008).
Prior structural analysis revealed that *P. aeruginosa* BfrB coordinates heme iron between
subunits, while FtnA does not[56], and that a knockout BfrB *P. aeruginosa* strain did not accumulate iron
within any other ferritin molecule, leading to their conclusion that BfrB (PA3531) was the main iron
storage molecule[57]. Moreover, the *P. aeruginosa* FtnA (PA4235) protein was described as a ferritin
distinct from bacterioferritin[58] that assembles as a heterooligomer 24mer from both BfrB and FtnA
subunits. A decreasing ratio of BfrB to FtnA was observed at low oxygen, with FtnA becoming the
dominant subunit under low oxygen conditions reaching ~30% to 70% of the subunits[56].
While these first two ferritins (PA3531 and PA4235) have been previously studied, less work has
been done on the 2nd bacterioferritin (and third ferritin overall), PA4880. Recent structural studies by
Rajapaksha et al[59] showed that PA4880 adopts a 12-mer structure similar to Dps proteins, but lacks the
iron coordinating ligands of Dps. Rajapaksha et al. named this gene DpsL (for Dps-like) and suggested
that this protein may contribute to avoiding iron toxicity and as well as "innate immune mechanisms
consisting of restriction endonucleases and cognate methyl transferases."
The current metalloproteome study provides a whole organism assessment of the iron storage
capabilities in *P. aeruginosa*. Since both the global and metalloproteomic techniques used here measure
tryptic peptides, we verified that these three ferritins do not share any tryptic peptides that could cause
misidentification from their peptidic constituents (see Figure S4-S6 for sequence alignments). In our
metalloproteomes, PA3531 (BfrB) and PA4235 (FtnA) both co-elute with in Fe Peak 1 (Figures 4 and 10)

and PA4880 (DpsL) elutes in Fe Peak 3 (Figures 7 and 10). The bacterioferritin-associated ferredoxin Bfd (PA3530) was not identified in either the anoxic or oxic metalloproteome.

The hypothesis that the contribution of ferritin (PA4235) increases under anoxic conditions by Yao et al.[56] was supported by our percentage contribution results: PA4235 was the only one of the three proteins to increase in percent contribution across the treatments from 8.2% to 22.2% between oxic and anoxic experiments (Table S1). There also appeared to be support for the accompanying hypothesis that BfrB and FtnA combine as heterogenous subunits to form a single ferritin 24mer protein complex[56]. We observed the ratio of BfrB:Ftn (PA4235:PA3531) to be highest in the AE 400 anoxic treatment with ratios of 1.1 and 7 between fractions 17 and 25 where many of these ferritins eluted (Figure 10F).

Notably the contribution to the native proteome in terms of total spectral counts showed PA4880 to be the most abundant iron storage protein, contrasting with the prior Eshelman et al. (2017) report of PA3531 as the dominant *P. aeruginosa* iron storage molecule as mentioned above. PA4880 had 76.5% of all of the ferritin molecule spectral counts in the oxic treatment and 60.9% in the anoxic treatment (Table S1). PA4880 was the dominant ferritin in AE 500 and AE 600 and coincided with the iron distributions, at SE 17 (oxic) and 18 (anoxic) for AE 500 (Figure 10G and 10H). Notably, PA4880 was not present in oxic AE 600, while it showed a large peak at SE 26 in anoxic AE 600, implying increased use of this bacterioferritin under anoxic conditions. This comparison assumes that the ionization efficiency of tryptic peptide constituents across these three proteins to be approximately averaged out, and that there are similar number of peptides based on their similar small size. More accurate assessments could be conducted in the future by use of targeted proteomics methods[60].

In terms of the amount of iron stored within each of the three ferritins, the metalloproteomic approach could assess this. Yet the high complexity of the *P. aeruginosa* iron metalloproteome makes this less straightforward compared to the straightforward approach used in our prior study on the simpler marine bacterium *Pseudoalteromonas*[19]. From the metalloproteomic data and media conditions of this study, it appears that the extent that the Fe metallome was stored varied considerably between oxic and anoxic treatments. In oxic treatment the two Fe Peak 1 ferritins BfrB and FtnA (PA3531 and PA4235) co-eluted with the major iron peak (Figures 4 and 10), whereas under anoxic conditions they were offset by one higher SE contributing to the shoulder. The extent of iron stored within these three ferritin proteins does not appear to dominate the iron metallome as it did in *Pseudoalteromonas*[19], being a relatively minor peaks compared to the iron metalloproteins, which are offset from the iron maxima, even with the high stoichiometry when fully loaded.

Detection of Prokaryotic Iron Supercomplexes Related to Key Cellular Metabolisms

The native metalloproteomic method used in this study purposefully does not use detergents in order to maintain metal-protein interactions. An additional advantage to omitting detergents is that protein complexes also have the potential to remain intact within the 2D chromatographic separations. As a result, the elution of proteins in native 2D space would not be expected to follow the size and charge state of the protein monomers but instead will reflect those of multimers and protein complexes. An indication of this is shown in Figure 11, where the distribution of molecular weights with size exclusion in each anion exchange fraction follows a downward trend, but also includes many proteins that do not follow this trend, eluting at sizes higher than their monomer molecular weight.

Here the four major Fe peaks identified in our study also appear to be related to protein complexes present in the cytosol of *P. aeruginosa*. In particular, the co-elution of proteins with related functions and co-located with dedicated iron storage systems implies a level of cellular organization at the protein-protein interaction level that is optimized for prioritization of metal distribution and trafficking. The presence of protein supercomplexes, assembles of multiple enzymes complexes (complexes of complexes) has been observed by native gel and chromatographic methods. For example the respiratory complexes, termed respirasomes, within mitochondria and bacteria contain assemblies

of respiratory chain complexes III and IV, each containing one or more enzymes[61,62]. The potential ability to observe and study supercomplexes from the entire soluble proteome and to observe their association with iron homeostasis is exciting. Application of recent AI-based structural modeling (Alphafold3) to examining protein-protein interactions[23] allowed exploration of putative protein complexes among co-eluting proteins. For example, a trimer protein complex was modeled with constituents of Fe Peak 1 (homogentisate 1,2-dioxygenase (PA2009), aconitate hydratase (PA1787), and hydrazone dehydrogenase (PA4022, HdhA, Figures 12 and 4). Similarly, protein-protein interactions for a putative trimer of proteins associated with the nearby Fe Peak 2 was modeled (Figures 12 and 6): PA4366 (superoxide dismutase), PA3529 (alkylhydroperoxide reductase C), and PA1787 again (aconitate hydratase 2). Bacterioferritins are 12mer protein complexes, and as mentioned above have been hypothesized to be interchangeable with ferritin subunits. While simple protein-protein interactions associations are beginning to be able to be modeled using artificial intelligence based tools (e.g. Alphafold3), the associations of large multimers like ferritins with other co-eluting proteins (e.g. aconitase) did not produce any predicted interactions, and hence the interactions that enable iron storage co-elution currently remain unknown to us.

It is likely that the many protein-protein interactions needed to create these supercomplexes not only involves a combination of protein binding as briefly explored here, but also the "crosstalk" of post-translational modifications (PTMs) between proteins. In this latter case, PTMs between proteins may contribute to the associations and functionality. For example, the eukaryotic aconitase enzyme has been shown to undergo reversible oxidation of its iron sulfur cluster and cysteine residues, as well as phosphorylation and transamidation PTMs, all of which contribute to cellular functions[63], and would create a variety of isoforms that elute and associate distinctly. While *in vitro* studies are more common in regulatory PTMs studies[64], *in vivo* datasets such as this native metalloproteomic approach offer an avenue to discover complex biological processes in use and their interaction with other cellular systems. Adding PTM analysis and interpretation of potential protein-protein interactions adds challenges to this already complex metalloproteomic analysis and presents a perhaps daunting level of biological complexity, yet it also presents great opportunities for discovery as this complexity is important to *in vivo* biology. Synergies are possible, such as further use of structural biology to help validate potential PTM sites[65] and protein-protein interactions. Future studies could take advantage of the large amount of raw spectra associated and their distributions within native separations space, such as from this study, to elucidate PTMs dynamics associated with protein-protein interactions particularly for those associated with protein supercomplexes[61,62].

From a systems biology perspective, these empirical observations of co-eluting and functionally related metalloproteins imply a cellular organization combining key metabolic functions with iron supply and biosynthesis of Fe-S cluster prosthetic groups within *Pseudomonas*. As bacteria tend to have only a few organelle structures (such as photosynthetic thylakoids, and protein-bound carboxysomes and vacuoles) compared to Eukaryotes[66], the strategies that allow targeting of enzymes and transporting of metals to organelles are less available to them. Yet, the challenges, including the potential for inefficiencies or outright failures, associated with directing metal resources towards specific reactions must be considerable, particularly in rapidly growing and metabolically flexible organisms such as *Pseudomonas*. Hence the co-locating of multiple metabolically-related iron enzymes and ferritins storage molecules would be advantageous, contributing to both the metalation and the prioritization of iron distribution to those metabolisms. The four supercomplexes observed in this study were not necessarily specific to iron metalloproteins: other metalloenzymes and proteins also co-eluted, such as copper enzymes in the Fe Peak 4 region and other non-metalloproteins, hence their subcellular organization may be related to both shared metabolic function and broader metal use. Future studies could further explore these iron supercomplexes and interrogate the regulation and structural biology behind them. While originally conceived as a methodology capable of investigating metalloproteins, this

native 2D approach also serves to contribute empirical information about protein-protein interactions that pertain to cellular metal homeostasis.

Data Availability

The processed metalloproteomic metal and protein data is available available as supplemental datasets and at Zenodo at 10.5281/zenodo.14950404. Four CSV data files are available: 2025_0214_pao_metals_oxic.csv, 2025_0214_pao_metals_anoxic.csv, 2025_0214_pao_proteins_oxic.csv, and 2025_0214_pao_proteins_oxic.csv. Raw proteomic spectra are available at ProteomeXchange and PRIDE (review access available upon request). The Jupyter notebook code for data analysis and visualization for this project is available at https://github.com/maksaito/Metalloproteomic-Viewer (Version 12).

Acknowledgements

We thank Dawn Moran for microbiological isolations and Viktoria Steck for assistance with genome sequencing samples. We thank Kevin Waldron for helpful discussions. The research was supported by NIH R01GM135709, the Simons Foundation, and the Center for Chemical Currencies on a Microbial Planet (NSF 2019589; contribution #069), and NSF projects 2123055, 2125063, 2048774.

References

N. J. Palleroni, *Microbiology*, 2003, **149**, 1–7.

N. Palleroni and I. Genius, *Bergey's Manual of Systematic Bacteriology*, 1984, **1**, 141–199.

G. P. Bodey, R. Bolivar, V. Fainstein and L. Jadeja, *Reviews of Infectious Diseases*, 1983, **5**, 279--313.

E. Tacconelli, E. Carrara, A. Savoldi, S. Harbarth, M. Mendelson, D. L. Monnet and et al., *The Lancet Infectious Diseases*, 2018, **18**, 318--327.

E. P. Skaar and M. Raffatellu, *Metallomics*, 2015, **7**, 926–928.

N. Kimata, T. Nishino, S. Suzuki and K. Kogure, *Microb Ecol*, 2004, **47**, 41–47.

S. Crone, M. Vives-Flórez, L. Kvich, A. M. Saunders, M. Malone, M. H. Nicolaisen, E. Martínez-García, C. Rojas-Acosta, M. Catalina Gomez-Puerto, H. Calum, M. Whiteley, R. Kolter and T. Bjarnsholt, *APMIS*, 2020, **128**, 220–231.

B. Borer, I. H. Zhang, A. E. Baker, G. A. O'Toole and A. R. Babbin, *PNAS Nexus*, 2023, **2**, pgac311.

C. A. Carlson and J. L. Ingraham, *Applied and Environmental Microbiology*, 1983, **45**, 1247-1253.

K. L. Thomas, D. Lloyd and L. Boddy, *FEMS Microbiology Letters*, 1994, **118**, 181-186.

A. J. Jesaitis, M. J. Franklin, D. Berglund, M. Sasaki, C. I. Lord, J. B. Bleazard, J. E. Duffy, H. Beyenal and Z. Lewandowski, *The Journal of Immunology*, 2003, **171**, 4329-4339.

B. E. Poulsen, R. Yang, A. E. Clatworthy, T. White, S. J. Osmulski, L. Li, C. Penaranda, E. S. Lander, N. Shoresh and D. T. Hung, *Proc Natl Acad Sci U S A*, 2019, **116**, 10072–10080.

C. Stover, X. Pham, A. Erwin, S. Mizoguchi, P. Warrener, M. Hickey, F. Brinkman, W. Hufnagle, D. Kowalik and M. Lagrou, *Nature*, 2000, **406**, 959–964.

I. J. Schalk and Q. Perraud, *Environmental Microbiology*, 2023, **25**, 811–831.

M. C. Mastropasqua, M. D'Orazio, M. Cerasi, F. Pacello, A. Gismondi, A. Canini, L. Canuti, A. Consalvo, D. Ciavardelli, B. Chirullo and P. Pasquali, *Molecular Microbiology*, 2017, **106**, 543-561.

M. M. Golden, A. C. Heppe, C. L. Zaremba and W. Wuest, *RSC Chemical Biology*, DOI:10.1039/D4CB00175C.

J. P. Barnett, D. J. Scanlan and C. A. Blindauer, *Anal Bioanal Chem*, 2012, **402**, 3371–3377.

A. Cvetkovic, A. L. Menon, M. P. Thorgersen, J. W. Scott, F. L. Poole II, F. E. Jenney Jr, W. A. Lancaster, J. L. Praissman, S. Shanmukh and B. J. Vaccaro, *Nature*, 2010, **466**, 779–782.

M. G. Mazzotta, M. R. McIlvin and M. A. Saito, *Metallomics*, 2020, **12**, 654–667.

M. G. Mazzotta, M. R. McIlvin, D. M. Moran, D. T. Wang, K. D. Bidle, C. H. Lamborg and M. A. Saito, *Metallomics*, 2021, **13**, mfab060.

J. K. Saunders, M. R. McIlvin, C. L. Dupont, D. Kaul, D. M. Moran, T. Horner, S. M. Laperriere, E. A. Webb, T. Bosak and A. E. Santoro, *Proceedings of the National Academy of Sciences*, 2022, **119**, e2200014119.

N. R. Cohen, M. R. McIlvin, D. M. Moran, N. A. Held, J. K. Saunders, N. J. Hawco, M. Brosnahan, G. R. DiTullio, C. Lamborg and J. P. McCrow, *Nature Microbiology*, 2021, **6**, 173–186.

J. Abramson, J. Adler, J. Dunger, R. Evans, T. Green, A. Pritzel, O. Ronneberger, L. Willmore, A. J. Ballard and J. Bambrick, *Nature*, 2024, 1–3.

D. A. Rodionov, A. G. Vitreschak, A. A. Mironov and M. S. Gelfand, *Journal of Biological Chemistry*, 2003, **278**, 41148–41159.

G. M. Teitzel, A. Geddie, S. K. De Long, M. J. Kirisits, M. Whiteley and M. R. Parsek, *Journal of Bacteriology*, 2006, **188**, 7242–7256.

A. Crespo, N. Blanco-Cabra and E. Torrents, *Frontiers in Microbiology*, 2018, **9**, 986.

J. M. Dunwell, A. Purvis and S. Khuri, *Phytochemistry*, 2004, **65**, 7–17.

C. J. Harmer, M. Wynn, R. Pinto, S. Cordwell, B. R. Rose, C. Harbour, J. A. Triccas and J. Manos, *PLoS One*, 2015, **10**, e0134229.

D. B. Ascher, O. Spiga, M. Sekelska, D. E. V. Pires, A. Bernini, M. Tiezzi, J. Kralovicova, I. Borovska, A. Soltysova, B. Olsson, S. Galderisi, V. Cicaloni, L. Ranganath, A. Santucci and A. Zatkova, *Eur J Hum Genet*, 2019, **27**, 888–902.

D. J. Hassett, M. D. Sutton, M. J. Schurr, A. B. Herr, C. C. Caldwell and J. O. Matu, *Trends in Microbiology*, 2009, **17**, 130–138.

K. Taniyama, H. Itoh, A. Takuwa, Y. Sasaki, S. Yajima, M. Toyofuku, N. Nomura and N. Takaya, *Journal of Bacteriology*, 2012, **194**, 1447–1456.

L. Castro, V. Tórtora, S. Mansilla and R. Radi, *Accounts of Chemical Research*, 2019, **52**, 2609–2619.

D. J. Hassett, W. A. Woodruff, D. J. Wozniak, M. L. Vasil, M. S. Cohen and D. E. Ohman, *J Bacteriol*, 1993, **175**, 7658–7665.

B. Polack, D. Dacheux, I. Delic-Attree, B. Toussaint and P. M. Vignais, *Infect Immun*, 1996, **64**, 2216–2219.

J. K. Saunders, D. A. Gaylord, N. A. Held, N. Symmonds, C. L. Dupont, A. Shepherd, D. B. Kinkade and M. A. Saito, *Journal of Proteome Research*, 2020, **19**, 4718–4729.

L. Chen, Q. W. Xie and C. Nathan, *Mol Cell*, 1998, **1**, 795–805.

L. C. Seaver and J. A. Imlay, *Journal of Bacteriology*, 2001, **183**, 7173–7181.

N. J. Hare, N. E. Scott, E. H. H. Shin, A. M. Connolly, M. R. Larsen, G. Palmisano and S. J. Cordwell, *Proteomics*, 2011, **11**, 3056–3069.

J. T. Lee, S. S. Lee, S. Mondal, B. N. Tripathi, S. Kim, K. W. Lee, S. H. Hong, H.-W. Bai, J.-Y. Cho and B. Y. Chung, *Mol Cells*, 2016, **39**, 594–602.

K. Nishida-Tamehiro, A. Kimura, T. Tsubata, S. Takahashi and H. Suzuki, *Plos One*, 2022, **17**, e0272090.

T. Yagi, *Journal of Bioenergetics and Biomembranes*, 1991, **23**, 211–225.

A. L. Pey, C. F. Megarity and D. J. Timson, *Bioscience Reports*, 2019, **39**, BSR20180459–BSR20180459.

K. Brown-Grant and J. Tata, *The Journal of Physiology*, 1961, **157**, 157.

I. Borovok, B. Gorovitz, M. Yanku, R. Schreiber, B. Gust, K. Chater, Y. Aharonowitz and G. Cohen, *Molecular Microbiology*, 2004, **54**, 1022-1035.

A. Jordan, E. Torrents, I. Sala, U. Hellman, I. Gibert and P. J. Reichard, *Journal of Bacteriology*, 1999, **181**, 3974-3980.

E. J. Blaesi, G. M. Palowitch, K. Hu, A. J. Kim, H. R. Rose, R. Alapati, M. G. Lougee, H. J. Kim, A. T. Taguchi and K. O. Tan, *Proceedings of the National Academy of Sciences*, 2018, **115**, 10022-10027.

J. Stubbe and M. R. Seyedsayamdost, *ACS Publications*.

E. Torrents, M. Westman, M. Sahlin and B.-M. Sjoberg, *Journal of Biological Chemistry*, 2006, **281**, 25287–25296.

T. Klünemann, A. Preuß, J. Adamczack, L. F. M. Rosa, F. Harnisch, G. Layer and W. Blankenfeldt, *J Mol Biol*, 2019, **431**, 3246–3260.

T. Kakutani, H. Watanabe, K. Arima and T. Beppu, *J Biochem*, 1981, **89**, 463–472.

M. Kukimoto, M. Nishiyama, M. Tanokura, E. T. Adman and S. Horinouchi, *J Biol Chem*, 1996, **271**, 13680–13683.

A. Romsang, J. M. Dubbs and S. Mongkolsuk, *Stress and Environmental Regulation of Gene Expression and Adaptation in Bacteria*, 2016, 1090–1102.

A. P. Kaur, I. B. Lansky and A. Wilks, *Journal of Biological Chemistry*, 2009, **284**, 56–66.

F. Cava, O. Zafra and J. Berenguer, *Molecular Microbiology*, 2008, **70**, 507–518.

M. J. Grossman, S. M. Hinton, V. Minak-Bernero, C. Slaughter and E. I. Stiefel, *Proc Natl Acad Sci USA*, 1992, **89**, 2419–2423.

H. Yao, A. Soldano, L. Fontenot, F. Donnarumma, S. Lovell, J. R. Chandler and M. Rivera, *Biomolecules*, 2022, **12**, 366.

K. Eshelman, H. Yao, A. N. D. Punchi Hewage, J. J. Deay, J. R. Chandler and M. Rivera, *Metallomics*, 2017, **9**, 646–659.

H. Yao, G. Jepkorir, S. Lovell, P. V. Nama, S. Weeratunga, K. P. Battaile and M. Rivera, *Biochemistry*, 2011, **50**, 5236–5248.

N. Rajapaksha, H. Yao, A. Cook, S. Seibold, L. Liu, K. P. Battaile, L. Fontenot, F. Donnarumma, S. Lovell and M. Rivera, *Front Mol Biosci*, 2024, **11**, 1390745.

M. A. Saito, M. R. McIlvin, D. M. Moran, A. E. Santoro, C. L. Dupont, P. A. Rafter, J. K. Saunders, D. Kaul, C. H. Lamborg and M. Westley, *Nature Geoscience*, 2020, **13**, 355–362.

R. Vartak, C. A.-M. Porras and Y. Bai, *Protein & cell*, 2013, **4**, 582–590.

A. Stroh, O. Anderka, K. Pfeiffer, T. Yagi, M. Finel, B. Ludwig and H. Schägger, *Journal of Biological Chemistry*, 2004, **279**, 5000–5007.

F. G. Oleh V. Lushchak Marta Piroddi and V. I. Lushchak, *Redox Report*, 2014, **19**, 8–15.

A. S. Venne, L. Kollipara and R. P. Zahedi, *Proteomics*, 2014, **14**, 513–524.

E. Vandermarliere and L. Martens, *Proteomics*, 2013, **13**, 1028–1035.

66. D. Murat, M. Byrne, and A. Komeili, Cold Spring Harbor Perspectives in Biology, 2010, **2**, a000422.

2 **Table 1.** Major Fe peaks in the oxic and anoxic metalloproteome of *Pseudomonas aeruginosa*.

Peak Number and Rank	Location Oxic (AE-SE)	Fe value Oxic (ppb)	Location Anoxic (AE-SE)	Fe value Anoxic (ppb)	Biochemical Function(s)
1	400-14	40.5	400-15	158.96	Respiration
2	300-17	25.7	300-18	46.2	Oxidative Stress
3	600-18	11.6	600-14	57.8	DNA production
4	200-20	4.1	200-22	28.7	Denitrification

3
4

1 **Table 2**. Iron proteins found within Fe Peaks 1-4 of the *Pseudomonas aeruginosa* metalloproteome.
2 Individual proteins are labelled with a letter suffix after the iron peak number.

Fe Peak Number	Metalloprotein (genome annotation)	PA ID	Comments
1a	aconitate hydratase 2	PA1787	
1b	homogentisate 1,2-dioxygenase (HmgA)	PA2009	6mer
1c	hydrazone dehydrogenase (HdhA)	PA4022	
1d	catalase	PA0236	
1e	bacterial ferritin	PA4235	
1f	bacterioferritin	PA3531	
2a	iron superoxide dismutase (SodB)	PA4366	
2b	alkylhydroperoxide reductase C	PA3529	dimer or decamer
2c	alkylhydroperoxide reductase C	PA0139	dimer or decamer
3a	glutamine synthase	PA5119	
3b	catalytic component of class Ia ribonucleotide reductase (NrdA)	PA1156	
3c	NrdB, tyrosyl radical-harboring component of class Ia ribonucleotide reductase	PA1155	
3d	probable bacterioferritin	PA4880	
3e	aconitate hydratase 1	PA1562	
3f	nitrite reductase precursor	PA0519	
3g	electron transfer flavoprotein alpha-subunit	PA2951	
3h	DNA-binding protein from starved cells, Dps	PA0962	
3i	ferric uptake regulation protein	PA4764	
3j	L-cysteine desulfurase (pyridoxal phosphate-dependent)	PA3814	
3k	class III (anaerobic) ribonucleoside-triphosphate reductase subunit, NrdD	PA1920	
4a	NirS nitrite reductase precursor	PA0519	anoxic/oxic
4b	NapA periplasmic nitrate reductase protein	PA1174	anoxic
4c	NirN	PA0509	anoxic
4d	probable iron-binding protein IscU	PA3813	anoxic/oxic
4e	Heme-transport protein, PhuT	PA4708	anoxic
4f	cytochrome c4 precursor	PA5490	anoxic/oxic
4g	cytochrome c-551 precursor	PA0518	anoxic
4h	cytochrome c551 peroxidase precursor	PA4587	anoxic/oxic
4i	cytochrome c5	PA5300	anoxic
4j	cytochrome c-type protein NapB precursor	PA1173	anoxic
	Selected Additional Metalloproteins of Interest		
4	nitrous-oxide reductase precursor	PA3392	anoxic
4	azurin precursor	PA4922	anoxic/oxic
4	probable glutathione peroxidase	PA0838	anoxic/oxic

Table 3. Optimization results for protein contributions to Fe Peak 1 (see Figure 5) as percent
contributions to each optimization (oxic and anoxic). PA2009 (homogentisate 1,2-dioxygenase) had a
second peak at higher SE fractions perhaps corresponding to an apo form, once removed PA2009 and
PA4022 (hydrazone dehydrogenase HdhA) reversed their contribution estimates, illustrating the
difficulty of determining the optimization when multiple proteins co-elute. Optimized coefficients are
shown in parentheses multiplied by 100 for readability.

Protein ID	Oxic: Original	Oxic: 2nd PA2009 peak removed	Anoxic
PA1787	32.9% (4.6)	34% (4.7)	0.0% (0)
PA2009	0.00% (0)	45.7% (41.9)	39.5% (2496)
PA4022	34.1% (57)	0.00% (0)	11.2% (17.6)
PA3531	24.8% (148)	12.8% (75.5)	7.8% (55.5)
PA4235	8.3% (33)	7.5% (29.7)	41.5% (130)

Figure Captions

Figure 1. Global proteomic abundance data for selected metabolic and metal related systems in *P.*
aeruginosa grown under aerobic and anaerobic conditions, where yellow is more abundant (values
normalized to the sum of spectral counts for the two treatments; extraction triplicates were averaged).
Strong responses were observed in nitrogen, iron, molybdenum and cobalt related systems. These
global proteomes are sample splits of those used for the metalloproteomic dataset in this study.

Figure 2. The Fe metalloproteome of *P. aeruginosa* under oxic (left) and anoxic (right) conditions in 3D
(top) and 2D (bottom) reveals four major iron peaks (numbered, Table 1). The four peaks were
associated with major cellular processes, 1) oxic respiration, 2) ROS detoxification, with 3) RNA
production, and 4) denitrification. AE refers to the first dimension of anion exchange (in units of mM;
note scale of oxic 2D is extended to 1000mM). SE refers to the second dimension of size exclusion and
represents fraction numbers. Iron (Fe 56) is in units of ppb.

Figure 3. The iron metallome in 2D native space under oxic (A) and anoxic (B) conditions (similar to
Figure 2 but with uniform scale for comparison to proteins), compared with the iron metalloproteins:
aconitate hydratase 2 (C, D; PA1787), homogentisate 1,2-dioxygenase (E, D; PA2009), hydrazone
dehydrogenase HdhA (G, H; PA4022), bacterial ferritin (I, J; PA4235), and bacterioferritin (K, L; PA3531).
Note some iron proteins have broad peaks (PA1787, PA4022) or double peaks (PA2009, PA4235) that
extend beyond the iron maximum region, which may be due to apo (unmetalled) or partially metalated
isoforms.

Figure 4. Iron proteins co-eluting with Peak 1 along the 400 AE 1^{st} dimension under oxic (A, E) and anoxic
(B, F) conditions. The iron maxima in this dimension aligns with the peak maxima of homogentisate 1,2-
dioxygenase (PA2009) in both oxic and anoxic conditions, while the much larger aconitate hydratase 2
(PA1787) peak is offset by slightly offset by two SE fractions under both conditions, however this protein
has a large broad distribution in native space (see Figure 2). The iron enzyme hydrazone dehydrogenase
HdhA (PA4022) also co-eluted with Peak 1, and transitions from being a minor peak in the oxic
conditions to a major peak in the anoxic conditions. (C-D) The ferritins (PA3531 and PA4235) align with
the iron maxima under oxic conditions but are also slightly offset by SE 1 fraction under anoxic
conditions. The iron peak has a shoulder in the 16-21 SE fraction range 9 (E-F), in both oxic and anoxic
conditions from the confluence of these metalloprotein peaks.

Figure 5. Optimization of protein contributions to Fe peak 1 under oxic (A) and anoxic (C) conditions.
Note that PA2009 has a second peak at SE 23 under oxic conditions (see Figure 4) that may be an apo
form. Because there is no iron peak at that location, the optimization results in a 0% contribution (A;
Table 3). When the second peak of PA2009 was removed (B), PA2009 contributes 45.7% of the Peak 1
area and PA4022 goes to 0% (Table 3). In the anoxic treatment (C) the optimization produces a 0.0%
(Table 3) contribution of PA1787, which similarly may be due to the broad and potentially apo forms of
this abundant enzyme.

Figure 6. Iron proteins that contribute to Fe Peak 2, examined in 1D along the size exclusion fractions
derived from the 300 AE dimension. The iron maxima in this dimension aligns with the peak maxima of
superoxide disumutase (PA4366) in both oxic and anoxic conditions. Aconitate hydratase 2 (PA1787) is
observed in Fe Peak 2, and it also observed in Fe Peak 1 (Figures 3, 4). As in Peak 1, it was also offset by
1-2 SE fractions under both conditions and has a large broad distribution in native space (see Figure 2).

Two copies of the iron enzyme alkylhydroperoxide (PA3529 and PA0139) co-eluted with Peak 2, with PA3529 going from the 3rd largest peak in oxic conditions to the largest peak in anoxic conditions, likely due to the need to address increased organic peroxides when growing on nitrate. and transitions from being a minor peak in the oxic conditions to a major peak in the anoxic conditions. Minor contributors to Fe Peak 2 included quinone oxidoreductase, a second copy of aconitate hydratase, and 3-isopropylmalate dehydratase large subunit (middle panels). There is a shoulder in Fe Peak 2 on the left side of the oxic peak at SE 15, likely contributed to by PA1787. The dotted grey line in all panels represents the local of the Fe maximum in AE 300, and the Fe anoxic peak was larger than under oxic conditions.

Figure 7. Proteins eluting at Fe Peak 3 in the AE 600 (A-F) and AE 500 (G-J) 1st dimension fractions. Dotted grey represents the iron maxima.

Figure 8. Abundance of proteins associated with the ribonucleotide reductase enzyme complexes within oxic and anoxic treatments from the global proteome data. NrdA/B, and NrdD/G require iron, while NrdJa/Jb requires vitamin B_{12}. Bars are the average and error bars represent standard deviation of the extraction triplicates in normalized spectral count units.

Figure 9. Metalloproteins associated with Fe Peak 4 20 oxic and at anoxic treatments (panels A-F), dashed line is the location of the iron maxima under each condition (AE 200 SE and AE 200 SE 22). Element traces along the AE 200 dimension for Fe, Cu, Ni, Zn, and P traces along the AE 200 dimension (panels G-P), dashed line is the maximum for each element. Left and right panels are oxic and anoxic treatments, respectively. Panels C and D replicate A and B, but without PA0519 to allow examination of lower abundance proteins. Proteins shown are known metalloproteins detected at the iron maxima positions, see Table S2 for list of proteins.

Figure 10. Abundance (oxic: A, D, G, J, anoxic: B, E, H, K; 300, 400, 500, 600) and ratios (C, F, I, L; oxic and anoxic) of ferritin iron storage in *P. aeruginosa*. Distributions of bacterioferritin (PA3531), ferritin (PA4235), and probable bacterioferritin/Dps-like protein (DpsL; PA4880) shown in the 300-600 anion exchange (AE) fractions of the first dimension and 10-35 size exclusion (SE) fractions of the second dimension. PA3531, PA4235, and PA4880 each had the highest spectral counts in AE 300, 400, and 500-600, respectively. Consistent with Yao et al., 2022, the ratio of BfrB:Ftn (PA4235:PA3531; left panels) was highest in the AE 400 anoxic treatment with ratios of 1.1 and 7 between fractions 17 and 25, where much of both ferritins eluted. The PA3531 and PA4235 maxima were slightly offset between treatments (panels D at SE14 in oxic and SE16 in anoxic), likely due to slightly different chromatographies between experiment. No ferritins were observed in the AE 600 oxic fraction.

Figure 11. Molecular weight distributions relative to size exclusion in each of the size anion exchange fractions (for oxic dataset). The decreasing trend in each fraction is consistent with larger proteins eluting earlier (lower fraction numbers). The presence of proteins in the bottom left corner of each plot is consistent with the presence of protein complexes, where smaller proteins form complexes and elute at lower SE fractions than expected based on their size.

Figure 12. Predicted structures and protein-protein interactions with example putative protein complexes comprised of co-eluting proteins in the native metalloproteome. Left: Trimer of proteins associated with Fe Peak 1: PA2009 (homogentisate 1,2-dioxygenase), PA1787 (aconitate hydratase 2) and PA4022 (hydrazone dehydrogenase, HdhA) (Figure 4). Right: Trimer of proteins associated with Fe

Peak 2: PA4366 (superoxide dismutase), PA3529 (alkylhydroperoxide reductase C), PA1787 (aconitate hydratase 2). Results generated using Alphafold3. The co-eluting supercomplexes (Figure 2) involve more proteins than the three in each case shown here, hence these Alphafold models are examples of putative protein-protein interactions that may exist within them.

Figure 1.

Figure with rows labeled top to bottom:
respiratory nitrate reductase alpha chain
respiratory nitrate reductase beta chain
respiratory nitrate reductase delta chain
respiratory nitrate reductase gamma chain
periplasmic nitrate reductase protein NapA
nitrate-inducible formate dehydrogenase
beta subunit nitrite reductase precursor
probable sulfite or nitrite reductase
regulatory protein NirQ
nitrite extrusion protein 2
nitric-oxide reductase subunit B
nitric-oxide reductase subunit C
NosD protein
NosL protein
nitrous-oxide reductase precursor
probable dinitrification protein NorD
probable binding protein component of ABC iron transporter
cytochrome c-551 precursor
ferripyoverdine receptor
probable non-ribosomal peptide synthetase
phenazine biosynthesis protein PhzE
probable phenazine biosynthesis protein
probable phenazine biosynthesis protein
ZnuA
Putative copper transport outer membrane porin OprC precursor
two-component response regulator, CopR
molybdenum cofactor biosynthetic protein A1
molybdopterin biosynthetic protein B1
molybdopterin biosynthetic protein C
cobalamin biosynthetic protein CobN

Columns: anoxic, oxic

1   **Figure 2.**

1   **Figure 3.**

1   **Figure 5.**

Figure 7.

1   **Figure 8.**

Figure 9.

1   **Figure 10.**

1   **Figure 11.**

1   **Figure 12.**
2
3