1   **A stress-activated neuronal ensemble in the supramammillary**

2   **nucleus produces anxiety-like behavior in male mice**

4   Jinming Zhang[1], Kexin Yu[1], Junmin Zhang[1], Yuan Chang[2], Xiao Sun[1], Zhaoqiang Qian[3],
5   Zongpeng Sun[4], Yanning Qiao[1], Zhiqiang Liu[1], Wei Ren[1,5], Jing Han[1,6,*]
6
7
1.   Key Laboratory of Modern Teaching Technology, Ministry of Education, Shaanxi
Normal University, Xi'an, 710062, China
2.   Department of Histology and Embryology, School of Basic Medical Science, Xi'an
Medical University, Xi'an, 710021, China
3.   Laboratory Animal Center, Shaanxi Normal University, Xi'an, 710119, China
4.   School of Psychology, Shaanxi Normal University, Xi'an, 710062, China
5.   Faculty of Education, Shaanxi Normal University, Xi'an, 710062, China
6.   Lead contact
16
*Correspondence:
Jing Han, Ph.D.
Key Laboratory of Modern Teaching Technology,
Ministry of Education,
Shaanxi Normal University,
9115 - 199 South Changan Road,
Xi'an, China, 710000
Tel/fax: +86-29-85308178
Email: jhan2012@snnu.edu.cn

Abstract

Anxiety is a prevalent negative emotional state induced by stress; however, the neural

mechanism underlying anxiety is still largely unknown. We used acute and chronic stress to

induce anxiety and test anxiety-like behavior; immunostaining, multichannel extracellular

electrophysiological recording and Ca^{2+} imaging to evaluate neuronal activity; and

virus-based neuronal tracing to label circuits and manipulate circuitry activity. Here, we

identified a hypothalamic region, the supramammillary nucleus (SuM), plays important role

in anxiety-like behavior. We then characterized a small ensemble of stress-activated neurons

(SANs) that are recruited by stress. These SANs respond specifically to stress, and their

activation robustly increases anxiety-like behavior in male mice. We also found that ventral

subiculum (vSub)-SuM projection but not dorsal subiculum (dSub)-SuM projections encode

anxiety-like behavior and that inhibition of these vSub-SuM projections has an antianxiety

effect. These results indicate that the reactivation of stress-activated supramammillary cells

and relevant neural circuits are important neural processes underlying anxiety-like behavior.

Keywords

Anxiety, Calcium activity, Supramammillary nucleus, Subiculum, Social stress

Introduction

Anxiety is a fundamental negative emotion observed in almost all mammal species. Long-lasting and uncontrollable anxiety often leads to several mental disorders, anxiety disorders, and even depression[5]. Recent studies have shown that the supramammillary nucleus (SuM), a part of the hypothalamus, regulates sleep[6], memory[7, 8], novelty exploration[7, 9], social memory[10, 11], neurogenesis[12], consciousness[13], locomotor activity[14], and theta oscillations in the hippocampus[15]. Projections from the SuM to the hippocampus have been largely studied and were found to modulate either episodic memory[7] or social memory[10, 11] in depending on the subregion of the hippocampus targeted. Although the SuM is located near the mammillary nucleus, a key region implicated in emotion regulation via the Papez circuit, its role in regulating emotion has been explored only superficially, without in-depth investigation. Despite some discussions of this role of the SuM, no consistent conclusion has been reached thus far[12, 16, 17].

Activity-dependent activation of cells have been studied in many brain areas[18, 19]. Tagged cells react to specific stimuli, such as conditional stimuli[20, 21], pain[22], food[23], and even peripheral inflammation[24], and mediate the storage and retrieval of relevant memories. The manipulation of those believed memory-associated cells can alleviate neurodegenerative diseases[25] or inflammation[24]. Naturally, this has led us to consider whether there is special neuronal ensemble that play roles in regulating anxiety or anxiety-like behaviors.Click or tap here to enter text. Recent studies have focused on the role of the hippocampus and related neuronal afferents and efferents[27–29]. The dorsal part of the hippocampus mainly contributes to cognition, whereas the ventral hippocampus is often associated with emotion[30, 31]. Although the ventral hippocampus□hypothalamus circuit was reported to modulate anxiety[27, 28], it is still unknown whether the SuM is part of this regulatory circuit. Although the SuM sends and receives dense neuronal projections, few studies have focused on its afferents or its ability to modulate behavior and emotion[32].

In this study, we hypothesize that stress can recruit a special neuronal ensemble that exclusively encodes anxiety. To test this hypothesis, we first used multiple methods to assess whether the SuM responds to acute or chronic stress. The activity of the SuM was chemogenetically manipulated, and anxiety-like behavior in rodents was tested. We subsequently employed the targeted recombination in active populations (TRAP) strategy to label and manipulate stress-activated neurons in the SuM in terms of anxiety-like behavior. After demonstrating how the SuM modulates anxiety, we sought to identify upstream brain areas that may contribute to supramammillary function. We also examined the functions of neuronal projections from the ventral subiculum (vSub) to the SuM using fiber photometry to measure calcium dynamics, as well as chemogenetic manipulation. These results allowed us to characterize a previously unreported role of SuM in regulating anxiety-like behavior. In addition, we showed that projections from the vSub but not the dorsal subiculum (dSub) to the SuM govern chronic stress-induced anxiety-like behaviors.

Results

1. Stress increases neuronal activity in the SuM

c-Fos protein expression was assessed after acute stress exposure to test whether the SuM was activated (Figure 1 A). The number of c-Fos$^+$ cells was significantly increased by foot shock exposure (Figure 1 B-C). To investigate if SuM would be responsive to diverse stressors, we next examined whether chronic stress, which different mechanism underlying, affects neuronal activity in the SuM (Figure 1 D-E). We chose in vivo electrophysiological extracellular recordings to reveal the neuronal activity before and after chronic social defeat stress, the data shows that the firing rate of regular-spiking neurons (RNs) (Figure 1 F) but not fast-spiking neurons (FNs) (Supplemental Figure 1 A-B) increased after CSDS. Regarding local field potentials, there were no noticeable difference between the naïve and CSDS groups according to power spectrum analysis (Supplemental Figure 1 C-D). These results indicate that acute and chronic stress can strongly activate the supramammillary nucleus.

2. Activation of SuM produces anxiety-like behavior

After confirming the activation of SuM caused different types of stress, we then further investigate whether the SuM regulates anxiety behavior in mice. Chemogenetic manipulations were conducted to activate SuM neurons. The experiments were performed as shown in the workflow (Figure 2 A-B). The mice were subjected to the open field (OF) and EZM tests at least 2 weeks after virus injection, followed by a reward-seeking test (Figure 2 E-H, Supplemental Figure 2 A). CNO was administered intraperitoneally 30 minutes before the test. Chemogenetic activation of the SuM did not affect the performance of mice in the OF test (Figure 2 E-F, Supplemental Figure 2 B). Compared with control mice, mice in which the SuM was activated explored the open arms of the EZM less (Figure 2 G), despite no change in distance traveled (Supplemental Figure 2 C). Moreover, mice in which the SuM was activated consumed less food than control mice did (Figure 2 F). These data suggest that there are neuronal ensembles that control the expression of anxiety behavior.

3. Identification of stress-activated neurons in the SuM

We next investigated whether an ensemble that encodes stress and controls the expression of anxiety exists. By crossbreeding Fos 2A-iCreERT2(TRAP2) and Rosa26-LSL-tdTomato (Ai14) mice, we generated TRAP2;Ai14 mice in which activated cells were genetically tagged for visualization (Figure 3 A). Foot shock exposure strongly activated neurons in the SuM but not adjacent areas (Figure 3 B-C). We hypothesized that these stress-activated neurons (SANs) respond exclusively to stress, but not to other stimuli such as reward. To validate the activity-dependent labeling in TRAP2;Ai14 mice, we labeled SANs in two cohorts of mice exposed either to the home cage condition or to foot shock. Several days after labeling, the mice were exposed to either sucrose pellets or social stress to induce reward-related or stress-related c-Fos expression, respectively. The reactivation of SANs under reward stimulation and stress was then compared (Figure 3 D-H). Foot shocks dramatically activated and labeled neurons in the SuM (Figure 3 F). Social stress but not reward (presentation of sucrose pellets) induced neuronal activation (Figure 3 G) and led to a much greater chance of reactivation of SANs (Figure 3 H). These data suggest the specific regulatory effect of the SuM on the effects of stress but not reward.

4. Reactivation of SuM^SANs promotes anxiety-like behavior

To make sure mice are on similar basal condition while applying chemo-genetic
manipulation, we subjected mice to an acute stress protocol involving foot shocks and then
performed the elevated plus maze (EPM) and elevated zero maze (EZM) tests to evaluate
anxiety on days 2 and 7 (Figure 4 A). The mice that experienced foot shocks showed
decreases in the exploration time in the open arms on day 2. However, acute stress-induced
anxiety was not detected on day 7 (Figure 4 B), which allow us to compare the reactivation of
SANs produced anxiety-like behavior between groups at the same baseline. Seven days after
SANs tagging, specific activation of SANs significantly increased the concentration of
corticosterone (a peripheral indicator of stress) in the mouse serum (Figure 4 C-D).
Experiments involving chemogenetic manipulation also revealed the sound-selective
activation of SANs in the SuM (Figure 4 E-F). We then tested whether manipulating SANs in
the SuM influences the anxiety-like behavior of the mice. The mice were subjected to the OF
and EZM tests at least 1 week after SANs were tagged, followed by reward-seeking tests
(Figure 4 G-H). CNO was administered intraperitoneally 30 minutes before the test.
Chemogenetic activation of SANs decreased the total distance traveled by the mice in the OF
and EZM tests (Supplemental Figure 2 F-G). The mice also presented decreases in the
distance traveled in the central area (Figure 4 I) and time spent in the central area in the OF
test (Figure 4 J), time spent in the open arms in the EZM test (Figure 4 K) and food
consumption (Figure 4 L). These data suggest that SANs in the SuM encode anxiety-like
behavior.

5. vSub-SuM projections encode anxiety-like behavior

The SuM receives afferents from various brain areas, and we identified projections to the
SuM by using a nonvirus- and virus-based retrograde tracing strategy (Figure 5 A,
Supplemental Figure 3 A-C). Afferents from the dSub and vSub were identified using
CTB-647 and AAV (Supplemental Figure 4 A-B). These projection neurons expressed Vglut1
RNA but not Vgat RNA (Figure 5 B), suggesting that Sub-SuM projections are excitatory
neuronal projections, as Vglut1 is a crucial marker of glutamatergic neurons. We then
performed an electrophysiological experiment. Optogenetics-evoked postsynaptic currents in
SuM neurons were blocked by perfusion with DNQX, which indicates the existence of
glutamatergic projections from the Sub to the SuM (Figure 5 C-E). To investigate how
Sub-SuM projections modulate stress and anxiety-like behavior, we then used fiber
photometry to measure the calcium concentration to assess the activity patterns of the
projection neurons (Figure 5 F-H). The projection neurons in the vSub but not those in the
dSub were more strongly activated when the mice moved into the open arms from the closed
arms of the EZM, (Figure 5 I-M). On the other hand, vSub but not dSub projection neurons
show lower calcium activity while mice backing to the closed arms (Supplemental Figure 5
A-D). Following exposure to acute stress, both dSub-SuM and vSub-SuM projection neurons
presented increased calcium activity (Figure 5 N-R). These data suggest that vSub-SuM
projections, but not dSub-SuM projections, may participate in regulating anxiety-liker
behavior.

6. Chronic inhibition of vSub-SuM projections alleviates anxiety-like behavior

After confirming the regulatory role of vSub-SuM projections in anxiety, we hypothesis
that inhibition of this projection would alleviate chronic stress-induced anxiety. In the
following experiments, the activity of these projections was chronically inhibited via a
chemogenetic strategy. The mice were exposed to CSDS after the expression of the Gi protein
was induced specifically on vSub-SuM projection neurons and their axons (Figure 6 A-B, D).
The body weights of the mice were monitored throughout the entire procedure to assess their
health (Figure 6 C). The mice showed no change in social interaction test scores after CSDS
exposure (Figure 6 E). In the EZM test, the mice in which vSub-SuM projections were
inhibited presented less anxiety-like behavior, as indicated by a longer time spent in the open
arms of the EZM but no significant change in the distance traveled (Figure 6 F-G). Taken
together, these data suggest that vSub-SuM projections are the essential neuronal projections
for regulating chronic stress induced anxiety-like behavior.
181
Discussion

In this study, we combined multiple methods to determine whether the SuM is a brain
region that involve in modulating anxiety. SuM neurons strongly respond to acute and chronic
stress, and their activation results in robust increases in anxiety-like behavior in mice. We
then defined a small ensemble of neurons that are activated by stress, called SANs. These
SANs specifically respond to stressful stimuli but not reward. Selective activation of SANs in
the SuM increases the serum concentration of corticosterone and anxiety-like behavior in
mice. The neuronal circuits that may be underlie the regulation of anxiety were also
determined in this study. The subiculum sends glutamatergic projections to the SuM and can
be activated by stress, whereas only the vSub has a potential effect on the transition to anxiety.
We finally determined that inhibition of SANs in the vSub project to the SuM is sufficient to
alleviate anxiety in mice after CSDS exposure.

Regulation of anxiety avoidance by the SuM

The SuM has been demonstrated to respond to novel environments, social stimulation[7,
33], and stress[34, 35]. Given these findings, we assume that the SuM may be activated by
foot shocks, a quantifiable acute stressor used in animal studies. Consistent with this
hypothesis, we found that the SuM robustly positively responds to both acute and chronic
stress through observable increases in c-Fos expression and increases in the neuronal firing
rate. The activation of the SuM was also demonstrated to be essential for maintaining
arousal[6, 13]. Sensitization to stressful events and high arousal are often associated with
anxiety[36]. Thus, our data strongly suggest that the SuM potentially modulates anxiety. To
further confirm whether the SuM participates in anxiety regulation, we recorded neuronal
action potentials via multichannel extracellular recording while the mice were moving in the
EPM, a traditional type of maze used to test anxiety in rodents. The change in the neuronal
firing frequency when mice transitioned from the open arms to the closed arms supports the
idea that the SuM may somewhat modulate anxiety. We then manipulated neuronal activity in
the SuM via a chemogenetic method and subjected the mice to the EZM test, an improved test
for assessing anxiety in rodents. The decrease in the exploration time in the open arms by
mice in which the SuM was activated by hM3Dq indicated increased anxiety. We noted that

these results are inconsistent with those of some previous reports. Some studies have reported
that lesions in the SuM and adjacent areas decrease anxiogenic behavior in rats[15, 37, 38].
López-Ferreras et al. performed the OF test, a complicated test, and reported that the
chemogenetic activation of neurons in the SuM results in increased anxiety-like behaviors in
rats[16, 17]. However, further experiments involving specific test (e.g., the EZM test) are
needed to confirm whether there is a potential difference across species.

The role of stress-activated neurons in regulating anxiety avoidance

Recent studies have highlighted the importance of activity-tagged neuronal ensembles in
regulating various behaviors, particularly memory[20, 39–41], food consumption[22], the
inflammatory response[24] and emotion[26, 42]. A negative experience-related neuronal
ensemble in the hippocampus was found to increase susceptibility to chronic stress[26]. A
recent study reported that the lateral habenula contains a small population of neurons that are
recruited in response to stress and mediate the development of depression in mice[42]. These
studies suggest that SANs may be important for emotional regulation. In this study, we found
that the SuM was more strongly activated by acute stress than was adjacent areas. SANs were
more likely to be reactivated by social stress than by sucrose reward, indicating their potential
to specifically encode anxiety. The serum corticosterone concentration can be used as a
marker of stress-induced change in the peripheral blood. Previous studies showed serum
corticosterone can be increased by various stress stimulation[39–42]Click or tap here to enter
text.; meanwhile, intentionally supplementing the diet with corticosterone can induce
anxiety-like behaviors in rodents[43]. Our data showed that the chemogenetic activation of
SANs in the SuM increased the serum corticosterone concentration, whereas the inactivation
of SANs had no effect, suggesting the chemogenetic manipulation of SANs may cause similar
anxiety effect like real stressors. These findings in combination with the results of the OF and
EZM suggest that SANs in the SuM are more likely involved in modulating anxiety-like
avoidance. However, the reactivation rate of SANs caused by different stressor was relatively
lower than the initial activation rate caused by foot shock (Figure 3). This suggests that
stress-activated neuronal clusters may have more flexible recruitment principles, with only a
small number of neurons potentially encoding emotional information, while most other
neurons remain involved in encoding other neural activities. Studies in other field,
particularly studies of memory engram, has shown that the sets of neurons activated during
learning are dynamic and exhibit high flexibility[44, 45]. While the activation of SANs
produced anxiety-like behavior, the future study will examine whether silencing SuM SANs,
either during stress exposure or during anxiety testing, can prevent or reduce stress-induced
anxiety. We also found that both the nonselective activation of SuM neurons and the selective
activation of SANs in the SuM significantly suppressed the consumption of sucrose pellets.
This result may be attributed to the anxiety-induced suppression of reward seeking[43].
However, further experiments are still needed to confirm whether this effect is anxiety
dependent and whether basal food consumption is affected.

A relevant neural circuit that regulates anxiety avoidance

The SuM recruits and is targeted by neuronal projections in the hippocampus, medial
septum, and cortex[32]. To further understand the circuitry through which the SuM regulates

anxiety, we identified projections from the dSub and the vSub to the SuM[46]. Fiber
photometry was used to measure the Ca^{2+} concentration in projection neurons in the dSub and
vSub, and the results revealed increased Ca^{2+} activity in vSub-SuM projection neurons but not
dSub-SuM projection neurons when the mice transited from the closed arms to the open arms,
indicating that vSub-SuM projections encode anxiety. To confirm the regulation of anxiety by
vSub-SuM projections, we exposed mice to CSDS and found that constant inhibition of
vSub-SuM activity significantly abolished CSDS-induced anxiety in mice. Unlike the dorsal
hippocampus, which is involved in the regulation of cognition, the ventral hippocampus is
often involved in regulating emotion[47]. The ventral CA1 area and its projections to the
lateral hypothalamic area were found to mediate innate anxiety, and its activation increases
anxiety-like behavior in mice[28]. Although very close spatially, neurons in the subiculum are
somewhat different from those in the CA1 region[48, 49]. The vSub and its downstream brain
areas were found to regulate anxiety[50, 51]. Jing-Jing et al. reported that the vSub and its
projections to the anterior hypothalamic nucleus are essential for anxiety because the
inhibition of these projections decreases anxiety-like behavior[27]. Our data are consistent
with these findings and suggest the mediating role of the vSub and its projections to different
subareas in the hypothalamus.

In summary, the activation of SuM increases anxiety-like behavior. Stressful event
recruits a neuronal ensemble in SuM. The activation of SANs also significantly increases
anxiety-like behavior and suppresses reward seeking. SuM receives glutamatergic projections
from the vSub, and inhibition of these projections can diminish CSDS-induced anxiety-like
behavior. These results suggest that SuM plays important role in regulating anxiety-like
behavior and furthermore studies are worthy performing.

Acknowledgment

We thank Dr. Wenting Wang for his generous gift of the transgenic mice. We thank all the
members of MTT for their valuable comments.

This research was funded by the National Natural Science Foundation of China (No.
82371518, 82071516, and 82441060), STI 2030—Major Projects 2021ZD0200500, the
Humanities and Social Science Fund of the Ministry of Education of China (No.
22XJC880005), the Innovation Capability Support Program of Shaanxi (Program No.
2021PT-055), the Natural Science Basic Research Plan in Shaanxi Province of China
(Program No. 2024JC-YBQN-0902 and 2023-JC-YB-189), and the Scientific and
Technological Innovation Team of Shaanxi Innovation Capability Support Plan (No.
2022TD-47).

Disclosures

The authors declare that they have no competing interests.

Methods and Materials

Animals

Male C57BL6/J mice aged 12–20 weeks were used. Fos$^{2A\text{-}iCreERT2}$ (TRAP2) mice were a
gift from Wenting Wang (JAX, cat. No. 030323). Rosa26-CAG-LSL-tdTomato (Ai14) mice
were purchased from the Shanghai Model Organisms Center (cat. No. NM-KI-225042). Male
CD-1 mice aged 8–10 months were purchased from Charles River (cat. No. 201). To construct
TRAP2;Ai14 mice, homozygous male TRAP2 mice and homozygous female Ai14 mice were
bred. Homozygous TRAP2;Ai14 mice were maintained and used in the experiments. We
performed genotyping for TRAP2 and Ai14 via PCR with the following primers: TRAP2
(wild-type: 357 bp, mutant: 232 bp): wild-type forward: GTCCGGTTCCTTCTATGCAG,
mutant forward: CCTTGCAAAAGTATTACATCACG, common:
GAACCTTCGAGGGAAGACG; Ai14 (wild-type: 297 bp, mutant: 196 bp): wild-type
forward: AAGGGAGCTGCAGTGGAGTA, wild-type reverse:
CCGAAAATCTGTGGGAAGTC, mutant forward: GGCATTAAAGCAGCGTATCC, mutant
reverse: CTGTTCCTGTACGGCATGG.

The mice were housed 4–5 per cage at a constant temperature and humidity (22 ± 1°C,
30–40% RH)on a day–night cycle (lights on from 08:00–20:00) with a fixed-intensity light
source. Each mouse was acclimated to the testing environment for 1□2 min. Acclimation was
performed for three days before the behavioral experiments. The mice were fed ad libitum
and euthanized with CO_2 after all the tests were finished. The experimental protocols
described here were approved by the Animal Ethics Committee of Shaanxi Normal
University.

Behavioral procedure

Acute stress exposure

316   The mice were exposed to acute stress according to a previously reported procedure[52];
317 specifically, they were exposed to 20 foot shocks with an intensity of 0.5 mA that were
318 randomly delivered across 10 minutes. The foot shocks were delivered in a fear conditioning
319 box (Med-associates).

Chronic stress exposure

321   We used a CSDS protocol to induce anxiety and depression in the mice[53]. Each
322 C57BL6/J mouse was housed with one CD-1 mouse, and the mice were separated by a
323 transparent plexiglass board with several small holes. The mice were allowed to contact each
324 other directly for 10 minutes every day for 10 days. Body weight was measured and recorded
325 every day before contact.

OF test

327   The OF test was carried out in a 50 × 50 × 35 cm arena made of white plexiglass. The
328 mice were allowed to move freely in the arena for 10 minutes, and the distance the mice
329 traveled and the time the mice spent in the central area were recorded and analyzed.

EPM test

331   The EPM consisted of two open arms (30 × 7 cm), two closed arms (30 × 7 × 14 cm) and
332 a central area (7 × 7 cm). The mice were allowed to move freely in the arena for 10 min, and
333 the time the mice spent in the open arms was recorded and analyzed.

EZM test

The EZM was used to test whether the mice were anxious. The EZM used in this study
was made of organic glass (height of 60 cm), with an inner diameter of 51.8 cm and an outer
diameter of 65 cm. The closed arms of the EZM were separated by two 15 cm-high pieces of
organic glass, the outer one of which was opaque. After a 15-minute habituation period, the
mice were placed into the EZM and allowed to move freely for 10 minutes. Videos were
recorded and analyzed using EthoVisionXT software. The time that the mice spent in the open
arms was compared between the groups to evaluate anxiety-like behavior.

Reward seeking

343 On the first day, sucrose pellets were provided for habituation. The mice were then
344 deprived of food on the second day. On the third day, the mice were placed in a new home
345 cage without bedding and allowed to eat freely for 2 hours. The pellets were weighed to
346 evaluate whether the experimental manipulation influenced reward seeking by the mice.

Social interaction test (SIT)

The SIT was conducted as previously described[53]. The SIT involved two 2.5-minute
phases. In the first phase (no-target phase), we placed each C57BL6/J mouse in the periphery
of the arena opposite the social interaction area (SIA). We allowed the animal to explore the
arena freely. In the second phase (with-target phase), each C57BL6/J mouse was placed in the
arena again, with a new CD-1 mouse in the SIA. The social interaction ratio (SIR) was
calculated using the following formula:

$$Social\ interaction\ ratio\ (SIR) = \frac{Time\ in\ SIA^{With-target} - Time\ in\ SIA^{No-target}}{Time\ in\ SIA^{With-target} + Time\ in\ SIA^{No-target}}$$

Click or tap here to enter text.**Neuronal tagging of stress-activated neurons**

To specifically label SANs, TRAP2 or TRAP2;Ai14 mice were intraperitoneally (i.p.)
injected with 4-hydroxytamoxifen (4-OHT, 50 mg/kg) immediately after acute stress exposure
or the learning phase of the CFC test. The mice were subjected to the next experiment or test
after 7 days to allow Cre-dependent recombination.

4-OHT (CAS No. 68392-35-8. Sigma, cat. No. H6278 or Bidepharm, cat. No.
BD00958757) was dissolved in DMSO at a concentration of 62.5 mg/mL and diluted with
vehicle (containing 10% TWEEN-80 and 80% saline) on the day of neuronal tagging. The
final concentration of DMSO was kept below 10% to avoid toxicity.

Observation of the reactivation of SANs

One week after neuronal tagging, whether previously tagged SANs were reactivated in
TRAP2;Ai14 mice when they were subjected to social stress was assessed. The mice in the
first group were underwent neuronal tagging in their home cages, and c-Fos expression was
induced by sucrose pellets. The mice in the second group were subjected to neuronal tagging
in response to foot shock exposure, and c-Fos expression was induced by sucrose pellets. The
mice in the third group were subjected to neuronal tagging in response to foot shock exposure,
and c-Fos expression was induced by social stress (one CD-1 mouse was placed in the home
cage). The mice were then sacrificed 90 minutes after sucrose pellet feeding or social stress
exposure, and c-Fos immunofluorescence staining was performed. The number of
c-Fos-positive neurons was counted to determine whether reward and cross-strain social stress

could activate neurons in the SuM. The likelihood of SAN reactivation was calculated using
the following formula:

$$Reactivation\ chance\ (\%) = (c - Fos^+/DAPI^+) \times (tdTomato^+/DAPI^+) \times 100$$

Viral vectors

An adeno-associated virus (AAV) vector was used to label and manipulate specific
neurons or determine the calcium concentration. To manipulate the neuronal activity in the
SuM, AAV2/9-hSyn-hM3Dq-EGFP (titer: 5.00E+12 GC/mL, BrainVTA, cat. No. PT-0891) or
its control vector AAV2/9-hSyn-EGFP (titer: 5.00E+12 GC/mL, BrainVTA, cat. No. PT-1990)
was injected into the SuM of mice.

To manipulate the activity of SANs in the SuM, AAV2/9-hSyn-DIO-hM3Dq -EGFP (titer:
5.00E+12 GC/mL, BrainVTA, cat. No. PT-0891) or its control vector
AAV2/9-hSyn-DIO-EGFP (titer: 5.00E+12 GC/mL, BrainVTA, cat. No. PT-1103) was
injected into the SuM of TRAP2 mice.

To chronically inhibit vSub-SuM circuitry activity, AAV2/Retro-hSyn-Cre (titer:
2.00E+12 GC/mL, Taitool, cat. No. S0278) was injected into the SuM, and
AAV2/9-hSyn-DIO-hM4Di-mCherry (titer: 2.00E+12 GC/mL, Taitool, cat. No. S0193) or its
control vector AAV2/9-hSyn-DIO-mCherry (titer: 2.00E+12 GC/mL, Braincase, cat. No.
BC-0025) was injected into the vSub of wild-type mice.

To determine the calcium concentration in dSub/vSub-SuM projection neurons,
AAV2/Retro-hSyn-Cre (titer: 2.00E+12 GC/mL, Taitool, cat. No. S0278) was injected into the
SuM, and AAV2/9-hSyn-DIO-GCaMP7b (titer: 5.00E+12 GC/mL, BrainVTA, cat. No.
PT-2892) was injected into the dSub/vSub of wild-type mice.

For the ex vivo electrophysiological experiment, AAV2/9-hSyn-ChR2-mCherry (titer:
5.00E+12 GC/mL, BrainVTA, cat. No. PT-0150) was injected into the vSub of wild-type
mice.

Neuronal tracing

Retrograde neuronal tracing was initially performed via injection of a serotype-2 AAV
vector (AAV2/Retro-hSyn-EGFP, titer: 5.00E+12 GC/mL, Taitool, cat. No. S0237) and
CTB-647 (1 µg/µL, Thermo Fisher, cat. No. C34778) into the SuM of wild-type mice. The
mice were then sacrificed after 2 weeks, and the brains were cut into coronal slices for
imaging.

To precisely trace neuronal afferents projecting to SuM^{SANs}, AAV2/9-Ef1α-DIO-RVG
(titer: 5.00E+12 GC/mL, BrainVTA, cat. No. PT-0061) and
AAV2/9-Ef1α-DIO-mCherry-F2A-TVA (titer: 5.00E+12 GC/mL, BrainVTA, cat. No.
PT-0207) were injected into the SuM of TRAP2 mice simultaneously. The rabies virus (RV)
vector RV-ENVA-ΔG-EGFP (titer: 2.00E+08 IFU/mL, BrainVTA, cat. No. R01001) was
injected into the SuM 2 weeks after neuronal tagging. The mice were then sacrificed after 2
weeks, and the brains were cut into coronal slices for imaging.

Stereotaxic surgery

The mice were anesthetized using isoflurane at a concentration of 1.5~2.0%. Virus was
injected into the SuM (AP: -2.8, ML: 0, DV: -4.5 mm), dSub (AP: -2.8, ML: ±0.7, DV: -1.7

mm) or vSub (AP: -3.5, ML: ±3.0, DV: -4.6 mm) according to the experimental design. If
only one type of virus needed to be injected into a single brain area, the final volume was
typically 150 nL. Otherwise, the final volume of the virus mixture was 200 nL. The viruses
were injected at a rate of 50 nL/min. The syringe was held in place for at least 5 minutes and
carefully removed from the brain. The mice were then returned to their home cages, and their
health was monitored on the following days. All the mice that underwent surgery were
subjected to the subsequent experiment after 2 weeks or more to allow virus expression.

For fiber photometry, ceramic ferrules (outer diameter: 2.5 mm, core diameter: 0.2 mm,
NA: 0.50) were inserted into the dSub (AP: -2.8, ML: ±0.7, DV: -1.5 mm) or vSub (AP: -3.5,
ML: ±3.0, DV: -4.4 mm) 2 weeks after virus injection under the guidance of a laser
(wavelength: 470 nm). Calcium imaging was conducted at least 1 week after ferrule
implementation.

Fiber photometry

Commercially available equipment (Thinker Tech) was used to determine the calcium
concentration. The fluorescence signal was activated by a laser at 470 nm, and the signal was
transmitted through a low-autofluorescence fiber□optic patch cord and rotary (doric lenses)
and collected. The final activation intensity was set to ~40 µW. The sampling rate was 50 Hz
for all the recordings. The mice were habituated to the fiber□optic patch cord for 3
consecutive days before recording. A TTL lasting 0.1 s was delivered by the software to mark
the timepoint when the mouse moved from a closed arm to an open arms in the EPM
(USB-IO box, Noldus). Continuous data were stored as *.tdms files and analyzed using
custom-made software in MATLAB.

Chemogenetic manipulation

To manipulate neuronal activity in the SuM in wild-type and TRAP2 mice, clozapine
N-oxide (CNO, 5 mg/kg; Cayman, cat. No. 25780) was injected i.p. 30 minutes before
behavioral tests were performed. For chronic inhibition of circuit activity, CNO was
administered orally (25 mg/L).

For acute and chronic experiments, CNO was dissolved in DMSO at a concentration of
10 mg/mL and stored at -20°C or in saline at a concentration of 1 mg/mL and stored at -80°C.
The storage solution was diluted with saline to a concentration of 0.75 mg/mL to prepare a
working solution for acute manipulation or to a concentration of 25 mg/L to prepare a
working solution for chronic inhibition on the day of the experiment.

Immunofluorescence

The mice were anesthetized with 20% urethane and perfused with PBS or saline. The
mouse brain was dissected and immersed in 4% paraformaldehyde (PFA) at 4°C overnight.
The PFA solution was then replaced with a 30% sucrose solution. After the brain sank to the
bottom, it was embedded in optimal cutting temperature (OCT) compound and frozen in a
cryostat (CM1950, Leica). Coronal slices (40 µm) were cut and collected in a 24-well plate.
After the residual OCT was removed with PBS, the slices were blocked with 0.3% Triton
X-100 and 10% normal donkey serum at room temperature (RT) for 2 hours. The slices were
then incubated with diluted primary antibody (rabbit anti-c-Fos, 1:500, Cell Signaling
Technology, cat. No. 2250) at 4°C overnight. The next day, the slices were washed and

incubated with secondary antibody dilutions (donkey anti-rabbit conjugated to AF647, 1:500,
Jackson ImmunoResearch, cat. No. 706-605-148) at RT for 2 hours. After washing, the slices
were transferred to slides and mounted with an antifade reagent (Thermo Fisher, cat. No.
P36981). Images of the slices were collected using a Zeiss M2 microscope and then analyzed.

RNA fluorescence in-situ hybridization

The samples were processed as described in the *Immunofluorescence* section. Slices (10
µm thick) were cut and dried at RT for ~15 minutes and then heated at 37°C for 30 minutes in
a hybridization oven. The baked slides were then moved to precooled 4% PFA solution for
fixation (~15 minutes). The slices were dehydrated in 100% ethanol at RT for 5 minutes. The
dehydration step was then repeated. The following steps were performed as recommended by
the manufacturer (ACDbio, cat. No. 323100). To label vglut1, vglut2 and vgat RNA, the
slices were hybridized with Mm-*Slc17a7* (ACDbio, cat. No. 416631-C1), Mm-*Slc17a6*
(ACDbio, cat. No. 319171-C1) and Mm-*Slc32a1* (ACDbio, cat. No. 319191-C3), respectively.
The samples were then stained with Opal dye.

Costaining of protein and RNA

After confirming RNA staining, the slices were blocked in 10% normal goat serum for 1
hour. The blocking solution was removed, and the slices were incubated with diluted primary
antibody (mouse anti-GFP, 1:500, Thermo Fisher, cat. No. MA5-16256; rabbit anti-tdTomato,
1:500, Oasis BioFarm, cat. No. OB-PRB013) at 4°C overnight. The slides were washed with
PBS and incubated with secondary antibody solution (goat anti-rabbit/mouse conjugated to
HRP, Proteintech, cat. No. PR30009) at RT for 1 hour (in the dark). After washing, the slices
were stained with Opal dye at RT for 30 minutes. The slides were then mounted and imaged.

Corticosterone assay

479 Mouse whole blood was collected 90 min after CNO injection (5 mg/kg, i.p.). The
480 samples were subsequently centrifuged at 2000×g for 10 minutes at 4°C after being left to
481 stand at RT for 30–60 minutes. The supernatant was then carefully collected as the serum.
482 Corticosterone levels were then measured using a commercial ELISA kit (Beyotime, cat. No.
483 PC100) according to the manufacturer's instructions.

Ex vivo electrophysiology

The mice were anesthetized with urethane and then decapitated. The brain was quickly
removed from the skull and immersed in precooled sucrose-based cutting solution (in mM,
225 sucrose, 2.5 KCl, 1.25 NaH_2PO_4, 26 $NaHCO_3$, 11 D-glucose, 5 L-ascorbic acid, 3 sodium
pyruvate, 7 $MgSO_4 \cdot 7H_2O$, 0.5 $CaCl_2$). After being fixed on a metal pallet, the brain was cut
into 300-µm slices. The slices were then collected and incubated in artificial cerebrospinal
fluid (ACSF) containing (in mM) 122 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 26 $NaHCO_3$, 11
D-glucose, 2 $MgSO_4 \cdot 7H_2O$, and 2 $CaCl_2$ equilibrated with 95% O_2-5% CO_2 at 28°C for at
least 1 hour before recording.

To evoke postsynaptic currents (PSCs) using light, whole-cell recording of global SuM
neurons near axons illuminated by ChR2-mCherry injected into the vSub was performed. The
final light intensity at the end of the optical fiber was set to ~5 mW/mm^2. Then, blue light
(470 nm, width: 10 ms, frequency: 0.05 Hz) was used to evoke optically induced PSCs
(oPSCs). DNQX (20 µM) was perfused into the ACSF to isolate AMPA-dependent currents

from the oPSCs after establishing a 5-minute baseline.

In vivo electrophysiology

The mice were anesthetized with 2% isoflurane and fixed to a stereotaxic device. A

sixteen-channel microwire electrode array (KD-MWA, KedouBC), a 4x4 array of 25 µm NiTi

wires spaced 200 µm apart, was slowly inserted into the mouse brain. Four small nails were

first inserted into the skull, with a ground wire presoldered onto one of them. The electrode

array was left in the SuM (AP: -2.8, ML: 0, DV: -4.55 mm), and then dental cement was used

to fix it onto the skull.

The mice were introduced to the recording area at least one week after surgery. During

the day, the electrode array attached to the mouse skull was connected to the OpenEphys

acquisition board through an Intan head stage. An OpenEphys GUI was used to visualize and

save electrical signals. The mice were allowed to move freely inside a home cage-like arena

for at least 20 minutes. Only data acquired during the last 5 min were saved and then analyzed

via Python-based software and a customized Python script.

Spikes were detected and divided into single units using SpikeInterface

(https://github.com/SpikeInterface/spikeinterface). Continuous binary raw data (sampling rate:

30 kHz) were imported and filtered using a bandpass butter filter at a cutoff value of 300 Hz.

Movement artifacts were removed by subtracting medians across all channels. The templates

were then extracted and fitted using spyking-circus2 inside the SpikeInterface frame. Neurons

meeting the following criteria were excluded from the subsequent analysis: (1) spikes with

refracting period violations smaller than 1 ms, accounting for more than 2% of total spikes,

and (2) a total frequency lower than 0.2 Hz. Neurons with spike frequencies $\geq$ 10 Hz were

considered RNs, whereas those with spike frequencies < 10 Hz were considered FNs, as

reported in a previous study[7]. The local field potential was extracted and analyzed using the

power spectrum analysis tool in MATLAB.

Statistical analysis

The data are presented as the means $\pm$ SEMs in all the figures in this manuscript. For

normally distributed data with equal standard deviations, independent t tests for unpaired data

and dependent t tests for paired data were performed in GraphPad software to compare mean

values between two groups. Otherwise, the Mann☐Whitney test for unpaired data and the

Wilcoxon test for paired data were performed instead. One-way ANOVA followed by Tukey's

post hoc test and two-way ANOVA followed by Sidak's post hoc test were performed to

compare mean values among more than three groups. A p value less than 0.05 was considered

to indicate a statistically significant difference between groups. "*" represents $p < 0.05$, "**"

represents $p < 0.01$, and "***" represents $p < 0.001$. Detailed statistical information referring

to specific figure can be found in Table 1.

References

1.   Nettle D, Bateson M. The evolutionary origins of mood and its disorders. Current
Biology. 2012;22.

2.   Liang M, Jian T, Tao J, Wang X, Wang R, Jin W, et al. Hypothalamic supramammillary
neurons that project to the medial septum modulate wakefulness in mice. Commun
Biol. 2023;6.

3.   Chen S, He L, Huang AJY, Boehringer R, Robert V, Wintzer ME, et al. A
hypothalamic novelty signal modulates hippocampal memory. Nature.
2020;586:270–274.

4.   Li Y, Bao H, Luo Y, Yoan C, Sullivan HA, Quintanilla L, et al. Supramammillary
nucleus synchronizes with dentate gyrus to regulate spatial memory retrieval through
glutamate release. Elife. 2020;9.

5.   Kesner AJ, Shin R, Calva CB, Don RF, Junn S, Potter CT, et al. Supramammillary
neurons projecting to the septum regulate dopamine and motivation for environmental
interaction in mice. Nat Commun. 2021;12.

6.   Qin H, Fu L, Jian T, Jin W, Liang M, Li J, et al. REM sleep-active hypothalamic
neurons may contribute to hippocampal social-memory consolidation. Neuron.
2022:1–15.

7.   Li J, Sun X, You Y, Li Q, Wei C, Zhao L, et al. Auts2 deletion involves in DG
hypoplasia and social recognition deficit: The developmental and neural circuit
mechanisms. Sci Adv. 2022;8:1238.

8.   Li YD, Luo YJ, Chen ZK, Quintanilla L, Cherasse Y, Zhang L, et al. Hypothalamic
modulation of adult hippocampal neurogenesis in mice confers activity-dependent
regulation of memory and anxiety-like behavior. Nat Neurosci. 2022;25:630–645.

9.   Pedersen NP, Ferrari L, Venner A, Wang JL, Abbott SBG, Vujovic N, et al.
Supramammillary glutamate neurons are a key node of the arousal system. Nat
Commun. 2017;8.

10.   Farrell JS, Lovett-Barron M, Klein PM, Sparks FT, Gschwind T, Ortiz AL, et al.
Supramammillary regulation of locomotion and hippocampal activity. Science (1979).

2021;374:1492–1496.

11.   Pan WX, McNaughton N. The role of the medial supramammillary nucleus in the
control of hippocampal theta activity and behaviour in rats. European Journal of
Neuroscience. 2002;16:1797–1809.

12.   López-Ferreras L, Eerola K, Shevchouk OT, Richard JE, Nilsson FH, Jansson LE, et al.
The supramammillary nucleus controls anxiety-like behavior; key role of GLP-1R.
Psychoneuroendocrinology. 2020;119.

13.   López-Ferreras L, Eerola K, Mishra D, Shevchouk OT, Richard JE, Nilsson FH, et al.
GLP-1 modulates the supramammillary nucleus-lateral hypothalamic neurocircuit to
control ingestive and motivated behavior in a sex divergent manner. Mol Metab.
2019;20:178–193.

14.   Tonegawa S, Liu X, Ramirez S, Redondo R. Memory Engram Cells Have Come of
Age. Neuron. 2015;87:918–931.

15.   Josselyn SA, Tonegawa S. Memory engrams: Recalling the past and imagining the
future. Science (1979). 2020;367.

16.   Liu X, Ramirez S, Pang PT, Puryear CB, Govindarajan A, Deisseroth K, et al.
Optogenetic stimulation of a hippocampal engram activates fear memory recall.
Nature. 2012;484:381–385.

17.   Guenthner CJ, Miyamichi K, Yang HH, Heller HC, Luo L. Permanent genetic access
to transiently active neurons via TRAP: Targeted recombination in active populations.
Neuron. 2013;78:773–784.

18.   Sun H, Li Z, Qiu Z, Shen Y, Guo Q, Hu SW, et al. A common neuronal ensemble in
nucleus accumbens regulates pain-like behaviour and sleep. Nat Commun. 2023;14.

19.   Azevedo EP, Pomeranz L, Cheng J, Schneeberger M, Vaughan R, Stern SA, et al. A
Role of Drd2 Hippocampal Neurons in Context-Dependent Food Intake. Neuron.
2019;102:873-886.e5.

20.   Koren T, Yifa R, Amer M, Krot M, Boshnak N, Ben-Shaanan TL, et al. Insular cortex
neurons encode and retrieve specific immune responses. Cell.
2021;184:5902-5915.e17.

21.   Ryan TJ, Roy DS, Pignatelli M, Arons A, Tonegawa S. Engram Cells Retain Memory
Under Retrograde Amnesia. Science (1979). 2015;348:1007–1013.

22.   Zhang TR, Larosa A, Di Raddo ME, Wong V, Wong AS, Wong TP. Negative memory

engrams in the hippocampus enhance the susceptibility to chronic social defeat stress.
Journal of Neuroscience. 2019;39:7576–7590.

23. Yan JJ, Ding XJ, He T, Chen AX, Zhang W, Yu ZX, et al. A circuit from the ventral
subiculum to anterior hypothalamic nucleus GABAergic neurons essential for
anxiety-like behavioral avoidance. Nat Commun. 2022;13.

24. Jimenez JC, Su K, Goldberg AR, Luna VM, Biane JS, Ordek G, et al. Anxiety Cells in
a Hippocampal-Hypothalamic Circuit. Neuron. 2018;97:670-683.e6.

25. Forro T, Volitaki E, Malagon-Vina H, Klausberger T, Nevian T, Ciocchi S.
Anxiety-related activity of ventral hippocampal interneurons. Prog Neurobiol.
2022;219.

26. Strange BA, Witter MP, Lein ES, Moser EI. Functional organization of the
hippocampal longitudinal axis. Nat Rev Neurosci. 2014;15:655–669.

27. Fanselow MS, Dong HW. Are the Dorsal and Ventral Hippocampus Functionally
Distinct Structures? Neuron. 2010;65:7–19.

28. Kesner AJ, Mozaffarilegha M, Rajamani KT, Arima Y, Harony-Nicolas H,
Hashimotodani Y, et al. Hypothalamic Supramammillary Control of Cognition and
Motivation. Journal of Neuroscience, vol. 43, Society for Neuroscience; 2023. p.
7538–7546.

29. Cumbers MR, Chung ST, Wakerley JB. A neuromodulatory role for oxytocin within
the supramammillary nucleus. Neuropeptides. 2007;41:217–226.

30. Cristina M, Silveira L, Sandner G, Graeff FG. Induction of Fos immunoreactivity in
the brain by exposure to the elevated plus-maze. Behavioural Brain Research.
1993;56:115–118.

31. Escobedo A, Holloway SA, Votoupal M, Cone AL, Skelton H, Legaria AA, et al.
Glutamatergic supramammillary nucleus neurons respond to threatening stressors and
promote active coping. Elife. 2023;12.

32. Leduke DO, Borio M, Miranda R, Tye KM. Anxiety and depression: A top-down,
bottom-up model of circuit function. Ann N Y Acad Sci. 2023;1525:70–87.

33. Aranda L, Santín LJ, Begega A, Aguirre JA, Arias JL. Supramammillary and adjacent
nuclei lesions impair spatial working memory and induce anxiolitic-like behavior.
Behavioural Brain Research. 2006;167:156–164.

34. Beck' CHM, Fibiger HC. Conditioned Fear-induced Changes in Behavior and in the

Expression of the Immediate Early Gene c-fos: With and Without Diazepam

Pretreatment. The Journal of Neuroscience. 1995;15:709–720.

35.   Tonegawa S, Morrissey MD, Kitamura T. The role of engram cells in the systems

consolidation of memory. Nat Rev Neurosci. 2018;19:485–498.

36.   Sun X, Bernstein MJ, Meng M, Rao S, Sørensen AT, Yao L, et al. Functionally Distinct

Neuronal Ensembles within the Memory Engram. Cell. 2020;181:410-423.e17.

37.   Lacagnina AF, Brockway ET, Crovetti CR, Shue F, McCarty MJ, Sattler KP, et al.

Distinct hippocampal engrams control extinction and relapse of fear memory. Nat

Neurosci. 2019;22:753–761.

38.   Zheng Z, Liu Y, Mu R, Guo X, Feng Y, Guo C, et al. A small population of

stress-responsive neurons in the hypothalamus-habenula circuit mediates development

of depression-like behavior in mice. Neuron. 2024. 2024.

https://doi.org/10.1016/j.neuron.2024.09.012.

39.   Baez M, Siriczman I, Volosin M. Corticosterone Is Involved in Foot Shock-Induced

Inactivity in Rats. Physiol Behav. 1996;60:795–801.

40.   Chen Y, Zhou X, Chu B, Xie Q, Liu Z, Luo D, et al. Restraint Stress, Foot Shock and

Corticosterone Differentially Alter Autophagy in the Rat Hippocampus, Basolateral

Amygdala and Prefrontal Cortex. Neurochem Res. 2024;49:492–506.

41.   Handa RJ, Nunley KM, Lorens SA, Louie JP, Mcgivern RF, Bollnow MR, et al.

Androgen Regulation of Adrenocorticotropin and Corticosterone Secretion in the Male

Rat Following Novelty and Foot Shock Stressors. Physiol Behav. 1994;55:117–124.

42.   dos Santos Corrêa M, Vaz B dos S, Grisanti GDV, de Paiva JPQ, Tiba PA, Fornari RV.

Relationship between footshock intensity, post-training corticosterone release and

contextual fear memory specificity over time. Psychoneuroendocrinology. 2019;110.

43.   Peng B, Xu Q, Liu J, Guo S, Borgland SL, Liu S. Corticosterone attenuates

reward-seeking behavior and increases anxiety via D2 receptor signaling in ventral

tegmental area dopamine neurons. Journal of Neuroscience. 2021;41:1566–1581.

44.   Zaki Y, Cai DJ. Memory engram stability and flexibility. Neuropsychopharmacology.

2024;50:285–293.

45.   Sweis BM, Mau W, Rabinowitz S, Cai DJ. Dynamic and heterogeneous neural

ensembles contribute to a memory engram. Curr Opin Neurobiol. 2021;67:199–206.

46.   Tang H, Wu GS, Xie J, He X, Deng K, Wang H, et al. Brain-wide map of projections

from mice ventral subiculum. Neurosci Lett. 2016;629:171–179.

47.   Shi HJ, Wang S, Wang XP, Zhang RX, Zhu LJ. Hippocampus: Molecular, Cellular, and

Circuit Features in Anxiety. Neurosci Bull. 2023;39:1009–1026.

48.   Aggleton JP, Christiansen K. The subiculum: The heart of the extended hippocampal

system. Prog Brain Res. 2015;219:65–82.

49.   Ding SL, Yao Z, Hirokawa KE, Nguyen TN, Graybuck LT, Fong O, et al. Distinct

Transcriptomic Cell Types and Neural Circuits of the Subiculum and Prosubiculum

along the Dorsal-Ventral Axis. Cell Rep. 2020;31.

50.   Mueller NK, Dolgas CM, Herman JP. Stressor-selective role of the ventral subiculum

in regulation of neuroendocrine stress responses. Endocrinology.

2004;145:3763–3768.

51.   Ghasemi M, Navidhamidi M, Rezaei F, Azizikia A, Mehranfard N. Anxiety and

hippocampal neuronal activity: Relationship and potential mechanisms. Cogn Affect

Behav Neurosci. 2022;22:431–449.

52.   Marcus DJ, Bedse G, Gaulden AD, Ryan JD, Kondev V, Winters ND, et al.

Endocannabinoid Signaling Collapse Mediates Stress-Induced Amygdalo-Cortical

Strengthening. Neuron. 2020;105:1062-1076.e6.

53.   Kim H-D, Call T, Carotenuto S, Johnson R, Ferguson D. Testing Depression in Mice: a

Chronic Social Defeat Stress Model. Bio Protoc. 2017;7.

54.   Zhang J, Zhang J, Yuan R, Han W, Chang Y, Kong L, et al. Inhibition of cannabinoid

degradation enhances hippocampal contextual fear memory and exhibits anxiolytic

effects. IScience. 2024;27:108919.

55.   Chang Y, Zhang J, Zhang J, Zhu W, Zheng Q, Qian Z, et al. N6-methyladenosine RNA

modification of glutamatergic neurons is associated with contextual fear

discrimination. Physiol Behav. 2022;248:113741.

Figure 1. Stress activates the SuM

(A) Workflow of the c-Fos staining. **(B)** Representative images of c-Fos staining (DAPI: blue; c-Fos: white;

scale bar: 100 μm). **(C)** Statistical analysis of the number of c-Fos-positive cells displayed in panel B. n =

10–13 per group; unpaired t test. **(D)** Workflow of CSDS exposure and in vivo recording. **(E)**

Representative spikes acquired by multichannel recording. **(F)** Statistical analysis of the firing rates of

RNs at baseline and after CSDS exposure. n = 16⌐23 per group; two-way ANOVA followed by Sidak's

post hoc test. The bars in C and F indicate the Min to Max of all data point, and the "+" indicates mean

value of all data point. "**", p <0.01; "***", p < 0.001. CSDS: chronic social stress; FS: foot shock. Also

see **Supplemental Figure 1.**

Figure 2. Activation of SuM produces anxiety-like behavior

(A-B) Virus injection information (A) and workflow for chemogenetic manipulation (B). **(C)** Representative

c-Fos images. **(D)** Statistical analysis of the number of c-Fos-positive cells displayed in panel C. n = 4 per

group; unpaired t test. **(E-F)** Statistical analysis of the distance traveled in the central area (E) and the

time that the mice spent in the central area (F) in the OF test. n = 10 per group; unpaired t test for the

data in (E) and the Mann⌐Whitney test for the data in (F). **(G)** Statistical analysis of the time that the mice

spent in the open arms of the EZM. n = 8 per group; unpaired t test. **(H)** Statistical analysis of sucrose

pellets consumed. n = 8–10 per group; Mann⌐Whitney test. The bars in D-H indicate the Min to Max of

all data point, and the "+" indicates mean value of all data point. "ns", p >0.05; "**", p <0.01. Also see

Supplemental Figure 2.

Figure 3. SANs in the SuM selectively respond to social stress but not reward

(A) Workflow of neuronal tagging. **(B)** Representative image of SANs in the SuM (DAPI: blue, tagged

cells: red). **(C)** Quantitative statistics of stress-tagged cells in several brain areas. n = 3 per area;

one-way ANOVA followed by Tukey'4s post hoc test. **(D)** Workflow of neuronal tagging and c-Fos staining.

(E) Representative images of stress-tagged cells and c-Fos expression induced by sucrose and social

stress (DAPI: blue, tdTomato: red, c-Fos: green). **(F)** Statistical analysis of the number of stress-tagged

cells in the SuM. n = 3 per group; one-way ANOVA followed by Tukey's post hoc test. **(G)** Statistical

analysis of the number of c-Fos^{+} cells after sucrose or social stress exposure. n = 3 per group; one-way

ANOVA followed by Tukey's post hoc test. **(H)** Statistical analysis of the reactivation of stress-tagged cells.

n = 3 per group; one-way ANOVA followed by Tukey's post hoc test. The data in C and F-H are presented

as the means ± SEMs. "ns", $p > 0.05$; "*", $p < 0.05$; "**", $p < 0.01$; "****", $p < 0.001$.

Figure 4. Selective chemogenetic activation of SANs elevates corticosterone level and produces

anxiety-like behavior

(A) Workflow of the acute stress and anxiety tests. **(B)** Statistical analysis of the time that the mice spent

in the open arms of the EZM. n = 10∟11 per group; two-way ANOVA followed by Sidak post hoc test. **(C)**

Workflow of the CORT assay and c-Fos staining. **(D)** Statistical analysis of the serum concentration of

corticosterone after the application of CNO. n = 4-5 per group; unpaired t test. **(E)** Representative images

of stress-tagged cells and c-Fos expression induced by chemogenetic manipulation (DAPI: blue, EGFP:

green, c-Fos: violet). **(F)** Statistical analysis of the percentage of costained cells relative to EGPF[+] cells in

the SuM. n = 4-5 per group; unpaired t test. **(G-H)** Virus injection information and workflow of

chemogenetic manipulation. **(I-J)** Statistical analysis of the distance traveled in the central area (I) and

the time that the mice spent in the central area (J) in the OF test. n = 14–15 per group; unpaired t test. **(K)**

Statistical analysis of the time that the mice spent in the open arms of the EZM. n = 14–15 per group;

Mann⊔Whitney test. **(L)** Statistical analysis of sucrose pellets consumed. n = 13–15 per group; unpaired

t test. The data in D & F are presented as the means ± SEMs. The bars in B and I-L indicate the Min to

Max of all data point, and the "+" indicates mean value of all data point. "*", p <0.05; "**", p <0.01; "***", p

< 0.001. Also see **Supplemental Figure 2.**

Figure 5. vSub-SuM projections encoding anxiety-like behavior

(A) Workflow of virus-based retrograde neuronal tracing. **(B)** Representative images of in situ RNA
staining (DAPI: blue, Slc32a1: green, Slc17a7: red, EGFP: white). **(C)** Workflow of ex vivo
electrophysiological recording. **(D)** Schematic of oPSCs in the SuM. **(E)** Representative traces of oPSCs.
(F) Workflow of Ca^{2+} imaging. **(G)** Schematic of Ca^{2+} imaging of dSub and vSub projection neurons. **(H)**
Representative images of GCaMP7b expression in the dSub, vSub and SuM (DAPI: blue, GCaMP7b:
green). **(I)** Heatmap of the Ca^{2+} fluorescence intensity during the transition from the closed to the open
arms. **(J)** Representative Ca^{2+} activity during the transition from the closed arms to the open arms. **(K)**
Average ΔF/F of Ca^{2+} recorded in the dSub and vSub. **(L)** Statistical analysis of the peak Ca^{2+} activity. n =
5 per group; unpaired t test. **(M)** Statistical analysis of the area under the curve of Ca^{2+} activity. n = 5 per
group; unpaired t test. **(N)** Heatmap of the Ca^{2+} fluorescence intensity during exposure to foot shocks. **(O)**
Representative Ca^{2+} activity during exposure to foot shocks. **(P)** Average ΔF/F of Ca^{2+} recorded in the
dSub and vSub. **(Q)** Statistical analysis of the peak Ca^{2+} activity. n = 5⌐6 per group; unpaired t test. **(R)**
Statistical analysis of the area under the curve of Ca^{2+} activity. n = 5⌐6 per group; unpaired t test. The
bars in L-M and Q-R indicate the Min to Max of all data point, and the "+" indicates mean value of all data
point. "ns", $p > 0.05$; "*", $p < 0.05$. Also see **Supplemental Figure 4** & **Supplemental Figure 5.**

766
Figure 6. Selective inhibition of vSub-SuM projections alleviated CSDS-induced anxiety
(A) Workflow of CSDS and chemogenetic manipulation. **(B)** Schematic of chemogenetic manipulation of
specific projections. **(C)** Body weight during CSDS exposure. **(D)** Representative images of virus
expression. **(E)** Statistical analysis of the social interaction ratio after CSDS exposure. n = 6-10 per group;
two-way ANOVA followed by Sidak's post hoc test. **(F)** Statistical analysis of the distance that the mice
traveled in the EZM. n = 6-10 per group; two-way ANOVA followed by Sidak's post hoc test. **(G)** Statistical
analysis of the time that the mice spent in the open arms of the EZM; two-way ANOVA followed by Sidak
post hoc test. The bars in E-G indicate the Min to Max of all data point, and the "+" indicates mean value
of all data point. "ns", $p > 0.05$; "*", $p < 0.05$; "**", $p < 0.01$; "****", $p < 0.001$.
776