The Brugada syndrome associated gene *WT1* impacts on *SCN5A* expression and cardiac

conduction

E. Madelief J. Marsman*[1], Joost A. Offerhaus*[1,2], Thomas Stervinou[3], Isabella Mengarelli[1],

Leander Beekman[1,2], Lisanne M.W. Wilde[1], Simona Casini[1], Tom Streef[4], Maurice J.B. van den

Hoff[5], Marie-José Goumans[4], Arie O. Verkerk[1,5], Sean J. Jurgens[1], Arthur A.M. Wilde[1,2],

Guillaume Lamirault[3], Nathalie Gaborit[3], Bastiaan J. Boukens[6], Carol Ann Remme[1], Connie R.

Bezzina*[1], Fernanda M. Bosada*[1].

[1] Department of Experimental Cardiology, Amsterdam Cardiovascular Sciences, Amsterdam

University Medical Centers, University of Amsterdam, Amsterdam, The Netherlands.

[2] Department of Clinical Cardiology, Heart Center, Amsterdam Cardiovascular Sciences,

Amsterdam University Medical Centers, University of Amsterdam, Amsterdam, The Netherlands.

[3] Nantes Université, CHU Nantes, CNRS, INSERM, l'institut du thorax, F-44000 Nantes, France.

[4] Department of Cell and Chemical Biology, Leiden University Medical Center, Leiden,

Netherlands.

[5] Department of Medical Biology, Amsterdam University Medical Centers, University of

Amsterdam, Amsterdam, The Netherlands.

[6] Department of Physiology, Cardiovascular Research Institute Maastricht University Maastricht

Maastricht, The Netherlands.

Abstract

Brugada syndrome (BrS) is an inherited cardiac arrhythmic disorder caused by conduction

slowing primarily affecting the right ventricular (RV) outflow tract (RVOT). A recent genome-wide

association study (GWAS) implicated a genomic region in chromosome 11, overlapping the

transcription factor *WT1*, in BrS susceptibility. Here, we investigated the role of *WT1* on cardiac

conduction using a heterozygous knockout mouse model ($Wt1^{+/-}$). Transcriptomic analysis

revealed increased *Scn5a* predominantly in $Wt1^{+/-}$ cardiomyocytes located subepicardially in the

RV and RVOT without any changes in electrical properties. To unmask an effect on cardiac

conduction, we performed optical mapping in a severely challenged setting offered by *Scn5a*

haploinsufficiency, ageing, and exposure to the sodium channel blocker ajmaline and found that

diminished *Wt1* improved the observed slowed conduction. Examination of human single-nuclei

cardiac datasets indicated a strong negative correlation between *WT1* and *SCN5A* expression.

In line with this observation, cardiac samples from patients carrying mutations in *SCN5A* showed

increased WT1 protein abundance in histological sections, suggesting that increased *WT1*, and

not loss, is associated with BrS pathophysiology. By deleting the mouse orthologue of a BrS-

associated noncoding region (RE) harboring a candidate regulatory element, we established that

this RE controls expression of *Wt1* specifically in the (sub)epicardium of the RV. Lastly, transient

overexpression of *WT1* in hiPSC-derived cardiomyocytes resulted in notably reduced sodium

current density. Our study thereby identifies the transcription factor *WT1* as a novel contributor to

the pathophysiology of BrS, at least in part, through *SCN5A*.

Introduction

Brugada Syndrome (BrS) is a heritable cardiac arrhythmia associated with increased risk of sudden cardiac death in young adults[1–3]. The pathophysiological mechanisms underlying BrS remain incompletely understood, hindering development of much needed strategies for risk stratification for life-threatening arrhythmic events and therapy. A long-standing hypothesis revolves around the slowing of conduction of the cardiac electrical impulse, to which the right ventricle (RV) and right ventricular outflow tract (RVOT) are particularly sensitive[4] likely due to their lower conduction reserve[5] and propensity for fibrosis[6]. This premise is supported by the fact that rare pathogenic loss-of-function variants in *SCN5A*, encoding the cardiac sodium channel (Na$_V$1.5), a critical determinant of cardiac conduction, are found in ~20% of BrS patients[7,8] although the vast majority of cases remain genetically elusive[9]. BrS was, until recently, considered a monogenic disease. However, major advances in genetics have uncovered more complex (polygenic) inheritance. Genome-wide association studies (GWAS) have identified 12 genetic loci harboring common variants associated with increased risk for BrS[10,11]. The majority of these disease risk-associated variants are found in noncoding genomic regions, and are presumed to alter the functionality of regulatory elements (REs)[12–14], which in turn lead to changes in gene expression. The disease-associated loci harbor *SCN5A*, genes that converge at Na$_V$1.5 regulation via transcriptional or post transcriptional mechanisms[10,15,16], as well as transcription factors (TFs) that may play a role in patterning of ion channel expression across the ventricular wall[15]. This supports the notion that diminished Na$_V$1.5 function is a central mechanism in BrS pathophysiology. Studies aimed at reliably identifying genes contributing to disease, and exploring how variants affect their expression, are crucial to understand BrS pathophysiology.

GWAS identified a genomic region enriched with noncoding BrS-associated variants in chromosome 11, directly upstream of *WT1* (Wilm's Tumor 1)[10]. This transcription factor is expressed during development and is known to drive epithelial-to-mesenchymal transition (EMT)

to form a population of multipotent epicardial-derived cells[17,18]. Endogenous *Wt1* expression can

be found in cardiomyocytes during development[19–22]; while it diminishes postnatally, it can

become re-expressed upon injury[23–26] or in the setting of disease[27–29]. More recently, two studies

demonstrated that conditional ablation of *Wt1* in cardiomyocytes during development or

postnatally results in morphological abnormalities and *in vivo* electrophysiological changes[22,30].

However, the possible impact of WT1 on ion currents has yet to be explored. Here, we establish

a clear association between this TF and BrS, while demonstrating the direction of change in

expression that may be present in patients.

Results

Wt1[+/-] **RV cardiomyocytes show altered ion channel expression without detectable**

electrophysiological changes

Since GWAS indicated an association with variants directly overlapping *WT1*[10], we studied

the role of this transcription factor using a heterozygous null mouse model (*Wt1*[+/-]). To investigate

how reduced *Wt1* affects the transcriptomes of cardiomyocytes, we performed RNA-sequencing

on the PCM1+ nuclei from RV/RVOT tissue of aged *Wt1*[+/-] (n=4; 15+ months) mice and age-

matched controls (n=4) (Figure 1A). Aged mice were studied in order to resemble the age at which

BrS becomes apparent in patients. Among 10722 transcripts detected (>10 averaged normalized

reads), 466 were upregulated and 405 downregulated in *Wt1*[+/-] cardiomyocyte nuclei (Figure 1B).

Notably, *Wt1* transcripts were almost undetectable in our cardiomyocyte-enriched samples

(Supplemental Table 1). Gene ontology term analysis using PANTHER[31] identified biological

processes including development (*Rbm20*, *Mef2d*, *Tcf7l2* and *Sox6*), cardiac muscle contraction

(*Cacna1c*, *Ryr2*, *Slc9a1* and *Myh7b*), and cell motility (*Actg1*, *Ctnna1*, *Cdh13* and *Sema6d*)

associated with upregulated genes in *Wt1*[+/-]. Cellular respiration (*Mtch2*, *Cox7b*, *Mdh1* and *Dld*),

muscle tissue development (*Tgfb2*, *Myoz2*, *Hand2* and *Gata6*), and intracellular transport (*Dnm1l*,

Mtx2 and *Calm1/2/3*) were associated with the downregulated gene set (Figure 1C). Strikingly,

we found increased *Scn5a* (L2FC 0.2333, p-adj=0.0124) and various deregulated genes involved

in cellular electrophysiology in *Wt1*[+/-] cardiomyocytes (Figure 1B, D).

Given the well-established ventricular transmural ion channel expression gradient for

normal conduction[15,32], we tested the impact of diminished *Wt1* in the (sub)endocardial and

(sub)epicardial sides of the ventricular wall on *Scn5a* by RT-qPCR. As expected, we observed

that in control hearts, *Scn5a* expression was higher in the (sub)endocardial side of controls

(Figure 1E), while *Wt1* showed a clear tendency to higher expression in the (sub)epicardial side

(Figure 1F). We also confirmed that *Wt1* was significantly downregulated in *Wt1$^{+/-}$* ventricles in

both transmural compartments (Figure 1F). Remarkably, we detected increased *Scn5a*

specifically in the (sub)epicardial side of *Wt1$^{+/-}$* ventricles when compared to controls (Figure 1E).

This indicates that the increase in *Scn5a* expression detected in cardiomyocytes of the RV/RVOT

of *Wt1$^{+/-}$* mice is driven by a mechanism primarily present in the (sub)epicardial side of the

ventricle, thus suggesting that epicardial *Wt1* inhibits *Scn5a*.

ECG analyses of adult (3-4 months) mice did not show any differences in PR-, QRS- and

QT duration in *Wt1$^{+/-}$* mice compared to age-matched controls (Supplemental Figure 1A),

indicating no overt differences in cardiac electrophysiological properties. RV and RVOT

(sub)epicardial conduction was measured in more detail in Langendorff-perfused hearts of adult

mice using optical mapping (Supplemental Figure 1B, C). Overall, no differences in conduction

velocity (CV) were observed between *Wt1$^{+/-}$* and control mice in either the RV or the RVOT. To

study the effect of *Wt1* haploinsufficiency on cellular electrical properties we measured action

potentials (AP) and sodium current (I_{Na}) in transmurally unselected cardiomyocytes isolated from

the RV of adult control and *Wt1$^{+/-}$* mice. AP upstroke velocity (V_{max}) was similar in cardiomyocytes

of both groups (Supplemental Figure 2A, B, Supplemental Table 1). In line with this, no differences

were observed in I_{Na} density or voltage dependence of (in)activation (Supplemental Figure 2C-E,

Supplemental Table 2). Resting membrane potential (RMP), AP amplitude (APA) and AP

repolarization parameters APD$_{20}$, APD$_{50}$ and APD$_{90}$ were also not significantly different between

RV cardiomyocytes from adult control and *Wt1$^{+/-}$* mice (Supplemental Figure 2A, B, Supplemental

Table 2). Together, our data indicate that diminished *Wt1* expression results in gene expression

changes which are not sufficient to lead to an electrophysiological phenotype.

Diminished *Wt1* improves conduction in a severely challenged setting

Next, we evaluated the possibility that electrophysiological changes driven by diminished

Wt1 may only become apparent in the context of reduced conduction. To investigate this, we

crossed *Wt1*$^{+/-}$ mice with *Scn5a* haploinsufficient mice (*Scn5a*$^{+/-}$) to generate double

haploinsufficient mice (*Scn5a*$^{+/-}$;*Wt1*$^{+/-}$) and examined whether aging provides a sufficiently

sensitized setting. We therefore aged control, *Scn5a*$^{+/-}$ and *Scn5a*$^{+/-}$;*Wt1*$^{+/-}$ mice to ~15 months,

equivalent to the approximate age at which BrS is diagnosed on average in humans (40-50

years)[2,33]. *In vivo* ECGs of aged mice (Supplemental Figure 3A) revealed that *Scn5a*

haploinsufficiency led to prolonged PR interval (Supplemental Figure 3B), while QRS duration

was unchanged, regardless of *Wt1* loss (Supplemental Figure 3C). Aging, independent of

genotype, did not have an effect on PR duration but did result in QRS interval prolongation in both

Scn5a$^{+/-}$ and *Scn5a*$^{+/-}$;*Wt1*$^{+/-}$, but not in control mice (Supplemental Figure 3). Indeed, an

interaction effect was borderline significant for QRS duration (p=0.0656), suggesting that aging

may unmask a more pronounced ventricular conduction phenotype in mice with only one copy of

Scn5a. We next investigated this in more detail by measuring epicardial CV in the RV and RVOT

in aged mice. We found slower CV in both longitudinal and transverse direction in *Scn5a*$^{+/-}$ and

Scn5a$^{+/-}$;*Wt1*$^{+/-}$ RVs than in controls (Figure 2A). In the RVOT we observed slower conduction in

Scn5a$^{+/-}$ compared to controls, but this difference was not detected between *Scn5a*$^{+/-}$;*Wt1*$^{+/-}$ and

controls (Figure 2B). We then further challenged our experimental model (aged *Scn5a*$^{+/-}$ and

Scn5a$^{+/-}$;*Wt1*$^{+/-}$ animals) by suppressing I_{Na} availability through acute administration of the I_{Na}

blocker ajmaline. Since the relationship of CV and I_{Na} is not linear[34], CV differences are more

easily to detect by the reduced excitability. Optical mapping analysis in the presence of ajmaline

showed a significantly higher CV in the transverse direction of the RV (Figure 2C), and in the

RVOT (Figure 2D) of *Scn5a*$^{+/-}$;*Wt1*$^{+/-}$ compared to *Scn5a*$^{+/-}$ mice. Taken together, our data show

that ageing contributes to ventricular conduction delay, and that decreased *Wt1* expression

improves conduction in *Scn5a* haploinsufficient mice in the setting of severely reduced conduction

reserve.

Inverse correlation between *WT1* and *SCN5A* expression in human transcriptomic datasets

Because diminished *Wt1* leads to increased *Scn5a* in cardiomyocytes, as well as a rescue

effect on conduction slowing in *Scn5a* deficient mice, we explored whether a correlation between

WT1 and *SCN5A* expression is present in human cardiac samples. Using GTEx data, we defined

two groups of samples, classified as "Low" or "High" cardiomyocyte-enriched, based on 10

cardiomyocyte-specific genes[35], which are significantly correlated with each other (Supplemental

Figure 4A-C). In "High" cardiomyocyte-enriched samples, we observed a significant negative

correlation between *SCN5A* and *WT1*, whereas no correlation was present in the "Low", non-

cardiomyocyte-enriched group (Supplemental Figure 4D). This correlation was further supported

by single-cell RNA-seq (snRNAseq) data of human induced pluripotent stem cell-derived

cardiomyocytes (hiPSC-CMs) (Figure 3A, B). We then assessed the relationship between *WT1*

and *SCN5A* expression in all atrial and ventricular cardiomyocytes (Figure 3C), and in atrial and

ventricular cardiomyocytes that express either, or both of these transcripts (Figure 3D). We found

a strong negative correlation, which indicates that cells with higher *WT1* expression have tapered

SCN5A and vice-versa and this may thereby impact on the transmural expression gradient

observed in mice (Figure 1E, F). We confirmed this result in snRNAseq datasets of non-diseased

human hearts[28]. Pseudo-bulk regression analysis across 4 ventricular regions revealed a strong

negative correlation between *WT1* and *SCN5A* (Supplemental Figure 4E, F, Supplemental Table

3).

Several of the conduction-related genes altered in cardiomyocytes of *Wt1*[+/-] mice,

including *Scn5a* (Figure 1D), are similarly correlated with *WT1* expression in human left

ventricular samples (GTEx). *WT1* is negatively correlated with *SCN5A, RYR2, KCNQ1,*

CACNA1C and *KCNIP2,* whereas it is positively correlated with *ATP1B3* (Supplemental Figure

5). This evidence indicates that *Wt1* haploinsufficiency in mice mirrors the transcriptional state

observed in human heart tissue, thereby strengthening the translational relevance of mouse

models.

_WT1_ is increased in pathologic conditions including BrS

In normal hearts, *WT1* expression diminishes after birth, but it can become re-expressed

upon injury[23–26]. Recent snRNAseq studies have shown increased *WT1* in ventricular samples of

patients with congenital cardiac defects and hypertrophic cardiomyopathy (HCM) when compared

to non-diseased hearts[27–29]. Moreover, patients with arrhythmogenic cardiomyopathy (ACM), a

disease exhibiting overlapping pathology with BrS, showed increased *WT1* when compared to

healthy controls[36]. We then considered whether elevated *WT1* expression occurs in the context

of BrS. To test this, we obtained histological ventricular sections from three *SCN5A* mutation-

positive patients. Immunohistological staining for WT1 revealed distinct nuclear signal in 2 out of

3 of these hearts, while non-diseased hearts, neither young (23 years) or old (79 years), showed

barely any detectable staining (Supplemental Figure 6). These data indicate that increased *WT1*,

and not its loss, is consistent with the BrS phenotype.

Next, we considered whether *SCN5A* loss-of-function itself leads to induction of *WT1* at

any point in time. To address this, we performed lineage tracing experiments using

Wt1^{Cre18};R26R-mTmG^{37}, in which the Cre recombinase drives membrane GFP only in the *Wt1*

lineage, and studied this in the context of *Scn5a* haploinsufficiency. We quantified GFP+

cardiomyocytes of the RV, left ventricle (LV), and interventricular septum (IVS) of aged *Scn5a^{+/-}*

and control mice using TnT staining and flow cytometry. We found significantly more GFP+

cardiomyocytes in the IVS and LV of *Scn5a^{+/-}* mice when compared to controls (Figure 4A). We

next analyzed GFP+/cTnT+ cells on histological sections and found increased double positive

cells in mice with reduced *Scn5a* expression (Figure 4B, C). These results indicate that life-long

reduced *Scn5a*, as it occurs in patients with loss-of-function *SCN5A* mutations, leads to induction

of *Wt1* in cardiomyocytes at any point through life, which further solidifies the biological relevance

of the inverse relationship between *Wt1* and *Scn5a*.

The BrS-associated noncoding variant region controls *Wt1* expression in the RV

Given that diminished *Wt1* expression may confer a protective effect, and patients with

SCN5A loss-of-function mutations display elevated WT1, we proceeded to investigate how the

BrS-associated variant region (BAR) in chr11p13 impacts on *WT1.* We first identified candidate

regulatory elements within this region. Human epigenetic datasets, including ventricular-like hiPS

cardiomyocyte ATACseq (a measure of open chromatin regions)[38], CTCF ChIPseq[39], and the

EMERGE enhancer prediction track[40,41] reveal moderate epigenetic signatures overlapping the

BrS-associated variant region (Figure 5A). In the mouse orthologous region, ATACseq of adult

ventricular and neonatal sinus node tissue[42,43] shows open chromatin in the region upstream of

Wt1, with modest H3K27ac signatures[44] (Figure 5B). We therefore selected a 4903 bp region

overlapping these epigenetic signatures for deletion using CRISPR/Cas9 (Figure 5C).

Homozygous deletion mice (*Wt1^BAR/BAR^*) were born at the expected Mendelian ratios and showed

no overt morphological defects. Using RTqPCR, we determined the expression level of *Wt1* in

the (sub)epicardial and (sub)endocardial regions of RVs and LVs of 6-week-old mice. *Wt1^BAR/BAR^*

mice showed a ~45% increase in *Wt1* transcripts in the (sub)epicardial side of the RV, but not in

any other cardiac region assayed (Figure 5D). This observation indicates that the BAR contains

one or more regulatory elements that selectively represses expression of *Wt1* in the

(sub)epicardium of the RV, thus further supporting the notion that increased *WT1* contributes to

BrS pathophysiology.

Overexpression of WT1 in hiPSC-derived cardiomyocytes leads to reduced sodium current

To determine the functional consequences of *WT1* overexpression in single cardiomyocytes, we generated hiPSC-CMs by using an adapted protocol based on small molecule-mediated modulation of the cWnt-pathway[45]. hiPSC-CMs were transfected using Lipofectamine 3000 with plasmids containing either WT1-P2A-GFP or GFP only (which served as control) and measured peak sodium (I_{Na}) in GFP-positive (GFP$^+$) hiPSC-CMs. Peak I_{Na} density was significantly lower in hiPSC-CMs transfected with WT1 (Figure 6A, B). For example, I_{Na} density at –20 mV was -87±10 and -54±7 pA/pF ($p<0.05$; t-test) in GFP$^+$ (n=16) and WT1 (n=12) overexpressing hiPSC-CMs. Neither the voltage dependency of (in)activation nor the time constants of current inactivation, determined at a test potential of -20 mV, were significantly different in hiPSC-cardiomyocytes with WT1 overexpression (Figure 6C, Supplemental Table 4), indicating that WT1 overexpression only affects I_{Na} density and not the gating properties. While a previous study reported that loss of *Wt1* in embryonic cardiomyocytes led to dysregulation of a potassium channel subunit which is an important regulator of the transient outward (I_{to1}) current[22], we did not detect any differences in I_{to1} density (Figure 6D, E) or I_{to1} gating properties (Figure 6F, Supplemental Table 4). AP measurements performed using dynamic clamp[46] demonstrated a significant decrease in V_{max} in hiPSC-CMs overexpressing WT1 compared to controls (Figure 6G, H), in line with the observed reduced I_{Na}. The RMP, APA, and AP repolarization parameters remained unchanged upon WT1 overexpression (Figure 6H, Supplemental Table 4). Taken together, these observations align with the effect of *Wt1* reduction on ventricular conduction demonstrated herein, and suggest a modulatory contribution predominantly on depolarization.

Discussion

Here, we explored the role of *WT1*, a gene at a locus implicated in BrS pathophysiology by GWAS. Our study reveals that diminished *Wt1* expression in mice leads to altered ion channel and cardiac contraction gene expression, including a modest increase in *Scn5a* specifically in subepicardial cardiomyocytes of the right ventricular wall. We further provide evidence supporting an inverse relationship between *WT1* and *SCN5A* expression in humans and identified increased *WT1* in pathological conditions, including in *SCN5A* mutation positive BrS patients. The interaction between these two genes was further explored in a challenged setting provided by *Scn5a* haploinsufficiency. While no differences in electrophysiological properties either *in vivo*, *ex vivo*, or in single cardiomyocytes were detected under baseline conditions, diminished *Wt1* did improve conduction under conditions of suppressed conduction reserve in the setting of advanced age and acute ajmaline exposure. By deleting the BrS-associated noncoding region orthologue, we show that *Wt1* is a target of the REs likely affected by the disease-associated common variants. Lastly, we demonstrate that overexpression of WT1 in hiPSC-derived cardiomyocytes reduces I_{Na} density, thereby establishing *WT1* as a new player in the pathology of BrS.

A recent GWAS proposed *WT1* as the most likely candidate gene implicated in BrS in chromosome 11, as indicated by the association signal in this locus[10]. In our study, we demonstrate that lifelong reduction of *Wt1* leads to an upregulation of *Scn5a* predominantly on the subepicardial side of the cardiac ventricles. This finding points to a potential mechanism by which *WT1* contributes to BrS through regulation of *SCN5A*, while also suggesting that increased-, rather than decreased WT1 expression, is associated with the BrS pathological state. Moreover, this highlights *WT1* as a potential key regulator in maintaining the transmural ion channel gradient, which is essential for proper cardiac function. Correspondingly, we demonstrate that patients with *SCN5A* mutations exhibit elevated WT1 protein levels in cardiac tissue, and that transient overexpression of WT1 in hiPSC-derived cardiomyocytes results in a significant reduction in

sodium current. This is consistent with existing knowledge regarding the loss of $Na_V1.5$ function

in the pathophysiology of BrS.

From an *in vivo* physiological perspective, discerning the impact of reduced *Wt1* is

challenging. In order to observe an electrophysiological phenotype, it was necessary to challenge

WT1 haploinsufficient mice by aging and reducing sodium channel expression and activity

through acute exposure to ajmaline. This could be due to the complexity associated with

accurately measuring increased $Na_V1.5$ function in a laboratory setting. *Scn5a*$^{+/-}$ mice showed

increased QRS duration only with ageing, and not at young adult age as was shown before[47]. A

possible explanation for the mitigated phenotype is differences in used mouse strain, affecting

electrophysiological parameters[48,49]. In this context, a rescue effect provided by *Wt1* reduction

(and thereby increased *Scn5a*) can only be uncovered by aggravating the phenotype even more.

Together, these observations provide strong evidence that *WT1* impacts on SCN5A expression,

and thereby also BrS pathophysiology.

The BrS-associated common variants in chromosome 11 are located in the noncoding

region upstream of *WT1*, and are therefore thought to impact on gene expression via REs[12–14].

RE activity is largely limited to the topologically-associating domain they share with their target

genes[50–52]. We presume that BrS-risk variants disrupt the function of REs that control *WT1*

expression. Consequently, by deleting the mouse orthologue of the candidate RE, we aim to

model the genetic mechanism(s) at play in BrS patients. Indeed, we demonstrated that loss of the

BrS-associated variant-rich RE led to increased *Wt1* specifically in the (sub)epicardial side of the

RV *in vivo*. This finding further supports the notion that common variants associated with

increased BrS risk at this locus likely result in *WT1* upregulation, which results in loss of $Na_V1.5$

function, thereby contributing to increased BrS-risk. However, further research is required to

elucidate the transcriptional regulatory mechanisms driving the function of this and other potential

REs within the locus.

We speculate that decreased or increased *WT1* impacts on *SCN5A* expression in the

opposite direction via a paracrine or cell-autonomous mechanism. Although WT1 is known to

function as either a transcriptional activator or repressor depending on its binding partners

(reviewed in[53]), not all evidence supports a cell-autonomous mechanism. We show that *Wt1*

expression was below detection in the cardiomyocyte-enriched fraction, yet *Scn5a* was

upregulated in *Wt1*[+/-] cardiomyocytes. This result may indicate that deregulation of *WT1* in non-

cardiomyocytes, via a paracrine process, affects gene expression in cardiomyocytes. The specific

cell-type in question has yet to be identified. Alternatively, *WT1* may be expressed at low levels,

in a small proportion, or only transiently in cardiomyocytes, making it difficult to detect any

decrease in expression, and thereby introducing further challenges for the interpretation of data.

Our study, along with others[27–29,36], indicates that elevated *WT1* expression is linked to

pathological states. Therefore, only in the context of increased WT1 we may be able to uncover

a cell-autonomous transcriptional effect on *SCN5A*.

It is also a possibility that WT1 affects SCN5A via a non-transcriptional mechanism.

Recent studies have revealed that WT1 can also control gene expression by modulating the

epigenetic landscape via interacting with TET family proteins[54,55]. WT1 is also known to modulate

topological switch mechanisms involving global changes in chromatin access to provide context

for specific gene activation or repression. For example, it acts as a transcriptional activator of

Wnt4 in the developing kidney mesenchyme undergoing mesenchymal-to-epithelial

transformation, and a repressor of *Wnt4* in epicardial cells poised for EMT[56]. WT1 may also

function post-transcriptionally according to a study which showed that WT1 localizes and interacts

with splicing factors in kidney cells[57]. Furthermore, WT1 has been shown to regulate mitotic

checkpoint via direct interaction with the spindle assembly checkpoint protein MADS2[58]. Thus,

additional research is needed to rule out the possibility of a non-transcriptional contributing

mechanism.

Conversely, decreased *SCN5A/Scn5a* expression also results in induction of *WT1/Wt1*,

as demonstrated by *Wt1*-lineage tracing on *Scn5a$^{+/-}$* mice and histological staining of patient

hearts with an *SCN5A* mutation. This suggests a tightly regulated interplay and the dynamic

balance influences cellular processes including cardiac conduction. Once either of these genes

shifts towards a pathologic state (i.e. increased *WT1* or decreased *SCN5A*), it may trigger a

potential maladaptive cyclic mechanism which progressively exacerbates organ dysfunction.

In conclusion, our data establish a role for the transcription factor *WT1* in the

pathophysiology of BrS, at least in part, through modulating expression and function of the cardiac

sodium channel *SCN5A*. Additionally, we present an explanation for the association of the

noncoding region in chromosome 11 with BrS in patients. Beyond rare coding loss-of-function

variants in *SCN5A*, we expect that the aggregate effect of common variants at one or more

disease-associated loci impact on Na$_V$1.5 function and thereby BrS susceptibility.

Methods

<u>Ethics statement</u>

Housing, husbandry, all animal care and experimental protocols were in accordance with guidelines from the Directive 2010/63/EU of the European Parliament, and Dutch government. Protocols were approved by the Animal Experimental Committee of the Amsterdam University Medical Centers. Animal group sizes were determined based on previous experience.

<u>Mice</u>

$Scn5a^{+/-}$ and $Wt1^{Cre}$ mice were generated as described previously[18,47], respectively. 129S4/SvJae $Wt1^{GFPCre/+}$ and $R26\text{-}mTmG$ mice were obtained from Jackson Laboratories (stock numbers 010911, 007676, respectively). Given that the GFP-Cre insertion in the $Wt1$ allele replaces exon 1 and thereby produces a null allele[59], the mice are herein referred to as $Wt1^{+/-}$. $Wt1^{+/-}$ mice were crossed with $Scn5a^{+/-}$ to generate $Scn5a^{+/-};Wt1^{+/-}$. $Wt1^{Cre}$ mice were crossed first to $R26\text{-}mTmG$ followed by $Scn5a^{+/-}$ for lineage tracing experiments.

$Wt1^{BAR/BAR}$ mice were generated using CRISPR/Cas9. Guide RNA (sgRNA) target sites were designed with PrimerQuest™ program (IDT, Coralville, Iowa, USA. Accessed 12 December, 2018. https://www.idtdna.com/SciTools). The Alt-R® CRISPR-Cas9 crRNA, Alt-R® CRISPR-Cas9 tracrRNA (IDT #1072532) and Alt-R® S.p. Cas9 Nuclease V3 (IDT #1081058) were ordered at IDT. One-cell FVB/NRj zygotes were microinjected via the "Mouse zygote microinjection Alt-R™ CRISPR-Cas9 System ribonucleoprotein delivery protocol" with 20 ng/µl Cas9 Nuclease and 10 ng/µl guide RNA to generate mouse founders. The guide sequences were as follows: 1) GGATTTAGGTCAAGATTCAC, 2) CACCACCATGGGATTCCTAA. Deletions were validated by PCR using the following primers: Forward 1) CTCTGGCTGCTGAAGGGATT, Reverse 1)

CTAGCTCTTACTGGCCACCG, Reverse 2) GAGAACAGAGCCCAGGAAGG and Sanger

sequencing. The deleted region spans chr2:105107107-105112009. To obtain stable lines, the

founders were backcrossed with wild-type FVB/NJ mice and maintained on FVB/NJ background

ordered from the Jackson laboratory (#100800). For tissue harvest, animals were euthanized by

20% CO_2 inhalation followed by cervical dislocation.

<u>Isolation of cardiomyocyte nuclei</u>

Nuclei isolation was performed as previously described[60] with the following changes: snap frozen

adult left and right atria from adult male and female mice were trimmed and homogenized in lysis

buffer containing RNase inhibitor using an Ultra-Turrax homogenizer. Samples were further

homogenized with a loose pestle douncer (10 strokes). After a 10-min incubation in the lysis

buffer, an additional 10 strokes were performed with a tight pestle. The lysis procedure was

monitored by light microscopy to ensure complete tissue and cell lysis and efficient nuclear

extraction. The crude lysate was successively passed through 100 and 30μm mesh filters. The

final lysate was spun at 1000 × g for 5 min and the resulting pellet was resuspended in 500 μl

staining buffer (5% BSA in PBS) supplemented with RNAse inhibitor. Isolated nuclei were

incubated with rabbit polyclonal antibodies specific for pericentriolar material 1 (PCM1) (Sigma-

Aldrich; HPA023370) at a dilution of 1:400 for 1 h rotating at 4 °C. Next, Alexa Fluor 647-

conjugated donkey-anti-rabbit 647 antibodies (ThermoFisher Scientific A-31573; 1:500 dilution),

and DAPI (1:1000 dilution) were added and the incubation was continued for an additional hour.

Samples were spun at 1000 × g for 10 min and washed with 500 μl staining buffer before

resuspension in 500 μl staining buffer supplemented with RNAse inhibitor. Intact cardiomyocyte

nuclei were sorted on a BD Influx FACS on the basis of DAPI and Alexa Fluor 647 positivity into

cold BL+TG buffer from the ReliaPrep RNA Tissue Miniprep System (Promega, Z6112) for RNA

isolation. RNA was isolated following a gDNA depletion step according to the manufacturer's

instructions.

<u>RNAseq</u>

For RNA isolated from cardiac nuclei, 500 pg was used for library generation with the Ovation

RNA-seq v2 kit (NuGEN M01206) kit. Libraries were prepared with the UltraLow V2 (NuGen) kit

and sequenced on the HiSeq4000 system (Illumina) with 50 bp single-end reads. 4 control and 4

Wt1[+/-] samples were sequenced. Reads were mapped to the mm10 build of the mouse

transcriptome using STAR[61]. Differential expression analysis was performed using the DESeq2

package based on a negative binomial distribution model. P-values were corrected for multiple

testing using the false discovery rate (FDR) method of Benjamini-Hochberg. We have used 0.05

as FDR control level. PANTHER32 was used for gene ontology (GO) biological process analysis.

Benjamini–Hochberg correction was performed for multiple testing-controlled P values.

Statistically significant enriched terms were functionally grouped and visualized.

<u>RTqPCR</u>

Total RNA was isolated from (sub) epicardial and (sub) endocardial ventricular sides of 12-16-

(*Wt1+/-*) or 6-week-old (*Wt1BAR/BAR*) male and female mice using ReliaPrep RNA Tissue

Miniprep System (Promega, Z6112) according to the manufacturer's protocol. cDNA was reverse

transcribed with oligo dT primers from 500ng of total RNA, according to the manufacturer's

protocol of the Superscript II Reverse Transcriptase system (Thermo Fisher Scientific, 18064014).

Expression levels of candidate target genes were determined by quantitative real-time PCR using

a LightCycler 480 Instrument II (Roche Life Science, 05015243001). Expression levels were

measured using LightCycler 480 SYBR Green I Master (Roche, 04887352001) and the primers

had a concentration of 10 pmol/L. The amplification protocol consisted of 5 minutes 95 °C followed

by 45 cycles of 10 seconds 95°C, 20 seconds 60°C and 20 seconds 72°C. Relative start

concentration (N0) was calculated using LinRegPCR[62]. Values were normalized to one or two

reference genes per experiment (Hprt and Ppia). The primer sequences are as follows: (all 3' to

5')   Hprt: TGTTGGATATGCCCTTGACT,   GATTCAACTTGCGCTCATCT;   Ppia:

GGGTGGTGACTTTACACGCC,   CTTGCCATCCAGCCATTCAG;   Scn5a:

GGGACTCATTGCCTACATGA,   GCACTGGGAGGTTATCACTG;   Wt1:

TTCAAGGACTGCGAGAGAAGG, TATGAGTCCTGGTGTGGGTCT.

In vivo electrocardiogram

12-16 (adult) or 65-67 week old male mice were anesthetized with 5% Isoflurane (Pharmachemie

B.V. 061756) and placed on thermostated mat (36ºC) with a steady flow of 1.5% isoflurane during

all experiments. Electrodes were inserted subcutaneously in the limbs and connected to an ECG

amplifier (Powerlab 26T, AD Instruments). The electrocardiogram (ECG) was measured for 5

minutes. ECG parameters were determined in Lead I based on 30 seconds of the recording. Only

male mice were used given that BrS is more prevalent in males.

Optical mapping and ajmaline exposure

Following ECG measurement, mice were sacrificed by cervical dislocation. The hearts were

excised, cannulated and mounted on a Langendorff perfusion setup. Hearts were submerged in,

and retrogradely perfused with 37 °C Tyrode's solution containing (in mmol/L): NaCl 128, KCl 4.7,

$CaCl_2$ 1.45, $MgCl_2$ 0.6, $NaHCO_3$ 27, NaH_2PO_4 0.4 and glucose 11 (pH maintained at 7.4 by

equilibration with a mixture of 95% O_2 and 5% CO_2). The excitation-contraction uncoupler

blebbistatin (10 µmol/L, Bio-Techne Ltd) was added in order to prevent movement artifacts.

During the optical mapping experiments, pseudo-electrograms were recorded by placing three

electrodes at ~0.5 cm distance of the heart in the Einthoven configuration. After ~10 minutes

recovery period, hearts received a 0.4 ml bolus of 20 µmol/L Di-4 ANEPPS (Molecular Probes),

the dye used for visualizing optical action potentials. Only hearts showing a regular sinus rhythm

that was stabilized within the first two minutes were included for analyses. Conduction velocities

were measured while pacing at the RV and the RVOT at a cycle length of 120 ms at twice the

stimulation threshold. Hereafter, hearts were treated continuously with ajmaline (1.5 µmol/L,

Carinopharm) and after a 10 minutes incubation period, the protocol was repeated. The average

length of the full protocol was 66.7± 1.1 minutes. Optical signals were analyzed with previously

published software [63], using Matlab2018. In short, optical signals were spatially (7x7 bin) and

temporally (0-150 Hz) filtered and local activation was defined as the maximum positive dV/dt of

the depolarization phase of the action potential. Conduction velocity was analyzed in two

transversal- (RV and RVOT) and two longitudinal (RV) directions and averaged for each direction.

<u>Generation, dissociation and transfection of hiPSC-CMs</u>

The hiPSCs (LUMC0099iCTRL04 line) were maintained on growth factor reduced matrigel

(Corning, USA) -coated plates in mTeSR1 (Stem Cell Technologies, Canada) medium at 37°C

(with 5% CO_2 and 20% O_2) and passaged every 3-4 days.

Cardiomyocytes were derived by using an adaptation of a published method[45]. Briefly, upon

reaching 85-95% confluence differentiation of the hiPSCs started by switching the medium to

RPMI 1640 (ThermoFisher Scientific, USA) supplemented with B27 minus insulin (ThermoFisher

Scientific, USA) and 213ug/ml L-Ascorbic Acid 2-phosphate (hereunder referred to as -I+AA

medium) and containing 6 µM CHIR99021 (Selleckchem, USA) for 72 hours. The medium was

than replaced with -I+AA medium containing 2 µM Wnt-C59 (MedChemExpress, USA) for Wnt

signaling inhibition and kept for other 72 hours on the cells. From day 8 onward, hiPSC-CMs were

cultured in RPMI 1640 supplemented with B27 plus insulin (ThermoFisher Scientific, USA),

without ascorbic acid and refreshed every 3-4 days until day 30.

For transfection and subsequent single-cell electrophysiological analysis, hiPSC-CMs cultures

were dissociated with 1× TrypLE Select Enzyme (ThermoFisher Scientific, USA) for 10 minutes

at 37ºC, collected, centrifuged and reseeded at a density of approximately 4×10^3 cells per

coverslip in presence of RPMI 1640 supplemented with B27 plus insulin and L-Ascorbic Acid 2-

phosphate (+I+AA medium). The glass coverslips were precoated with GFR matrigel/gelatin at

50/50 ratio.

Transfection was performed using Lipofectamine 3000 reagent (ThermoFisher) according to

manufacturer's protocol. Two pcDNA3.1 plasmids with CMV promotor and a P2A-eGFP site,

containing one of two common *WT1* isoforms were used together in 1:1 ratio: isoform D

(NM_024426.4, OHu27351, Genscript) and isoform B (NM_024424.3, OHu27419, Genscript). For

the control, the same plasmid without *WT1* was used ("GFP-only"). Amount of plasmid (in ng) was

chosen based on plasmid size (WT1-GFP 7.7kb, GFP 6.1kb), to ensure equal plasmid copies per

condition. For each well in a 24-well plate, 0.75µl (for WT1) or 0.59µL (for GFP-only) lipofectamine

was dissolved in 25µl RPMI (without glucose) and 500ng WT1 or 396 ng GFP-only plasmid with

1µl (for WT1) or 0.79µl (for GFP) P3000 in 25µl RPMI, after which these two were gently mixed

(i.e. lipofectamine µl to DNA µg ratio 1.5) and incubated 10-15 minutes. Mixes were added to the

cells and removed and replaced by culture medium after 3 days, and patched the next day.

<u>RV cardiomyocyte isolation</u>

Freshly excised hearts of male mice were cannulated and mounted on a Langendorff setup,

retrogradely perfusing oxygenated modified Tyrode's solution at 37 °C containing (in mmol/L):

NaCl 140, KCl 5.4, $CaCl_2$ 1.8, $MgCl_2$ 1.0, glucose 5.5, and HEPES 5.0; pH 7.4 (adjusted using

NaOH). After 10 minutes, hearts were perfused with nominally calcium-free solution containing

(in mmol/L): $CaCl_2$ 0.01, creatine 10.7. After a consecutive 10 minutes, hearts were digested by

addition of Liberase TM (0.032 mg/mL, Roche) and elastase (1.6 U/mL, SERVA Electrophoresis

GmbH), while perfusing for ~15 minutes. Hearts were washed in calcium-free solution containing

1% bovine serum albumin fraction V, fatty acid free (Roche). The RV was removed and cut into

small pieces after further dissociating by pipette trituration to obtain single cardiomyocytes, which

were washed in perfusion solution while up-titrating $CaCl_2$ to 1.8 mmol/L. Isolated RV

cardiomyocytes were stored at room temperature in modified Tyrode's solution and used for AP

and I_{Na} recordings within 2-3 hours.

<u>Cellular electrophysiology</u>

Data acquisition and analysis: APs and I_{to1} were recorded with the amphotericin-B perforated

patch-clamp technique while I_{Na} was recorded with ruptured patch-clamp technique, using an

Axopatch 200B patch-clamp amplifier (Molecular Devices, San Jose, CA, USA). Voltage control,

data acquisition, and analysis were done using custom software[64] and pClamp10.6/Clampfit

(Molecular Devices, San Jose, CA, USA). Borosilicate glass patch pipettes (GC100F-10; Harvard

Apparatus, Waterbeach, UK) with a tip resistance of 2-2.5 MΩ were used. Signals were low-pass-

filtered with a cutoff of 5 kHz and digitized at 20 kHz for I_{Na}, 30 kHz for I_{to1}, and 40 kHz for APs.

Cell membrane capacitance was calculated by dividing the time constant of the decay of the

capacitive transient after a −5 mV voltage step from −40 mV by the series resistance[46]. Series

resistance was compensated for >80%. I_{to1} and APs were corrected for the calculated liquid

junction potential. Following the procedure of Veerman et al.[65], we selected single, spontaneously

beating, GFP[+] or WT1 overexpressing hiPSC-CMS showing regular, synchronous contractions in

modified Tyrode's solution, after which this extracellular solution was switched to a solution

suitable for specific measurements as indicated below. For mouse, recordings were made from

quiescent cardiomyocytes.

Action potential measurement: APs were measured in hiPSC-cardiomyocytes and mouse

cardiomyocytes at 37°C in modified Tyrode's solution, containing (in mmol/L): NaCl 140, KCl 5.4,

$CaCl_2$ 1.8, $MgCl_2$ 1.0, glucose 5.5, HEPES 5.0; pH 7.4 (NaOH). Patch pipettes were filled with a

solution containing (in mmol/L): K-gluconate 125, KCl 20, NaCl 5 (hiPSC-CMs) or 10 (mouse),

amphotericin-B 0.44, HEPES 10; pH of 7.2 (KOH). APs were elicited at 4 Hz in mouse

cardiomyocytes and at 1 Hz in hiPSC-CMs by 3 ms, 1.2-1.4 x threshold current pulses through

the patch pipette. To overcome the lack of the inward rectifying potassium current (I_{K1}) in hiPSC-

cardiomyocytes, which limits the functional availability of I_{Na} and I_{to}[66], we injected an 2 pA/pF in

silico I_{K1} with kinetics of Kir2.1 channels through dynamic clamp, as previously described and

validated[67]. AP parameters were analyzed from 10 consecutive APs and averaged.

Sodium current measurement: I_{Na} was measured at room temperature (~20°C). The bath solution

for I_{Na} recordings in mouse cardiomyocytes contained (in mmol/L): NaCl 7.0, CsCl 133, $CaCl_2$ 1.8,

$MgCl_2$ 1.2, glucose 11.0, nifedipine 0.005, HEPES 5.0; pH 7.4 (adjusted with CsOH), while hiPSC-

CMs were perfused with an external solution containing (in mmol/L): NaCl 20, CsCl 120, $CaCl_2$

1.8, $MgCl_2$ 1.0, glucose 5.5, nifedipine 0.010, HEPES 5.0; pH 7.4 (adjusted with CsOH). Pipette

solution was composed of (mmol/L): NaCl 3.0, CsCl 133, $MgCl_2$ 2.0, Na_2ATP 2.0, TEACl 2.0,

EGTA 10, HEPES 5.0; pH 7.2 (adjusted with CsOH). I_{Na} was recorded with a double-pulse

protocol from a holding potential of −120 mV (see inset of Supplemental Figure 2C and Figure

6A; cycle length of 5 seconds). During the first 500 ms depolarizing pulses, I_{Na} activates and the

currents analyzed here are used to determine I-V relationships and the voltage dependency of

activation. For the latter, the I-V relationships were corrected for driving force and normalized to

maximal current. The second pulse is used to determine the voltage dependency of inactivation

and currents were normalized to the largest I_{Na}. I_{Na} was defined as the difference between peak

and steady-state current and densities were calculated by dividing current amplitude by C_m.

Voltage dependency curves of activation and inactivation were fitted with Boltzmann function:

$I/I_{max}=A/\{1.0+\exp[(V_{1/2}-V)/k]\}$, where $V_{1/2}$ is the half-maximal voltage of (in)activation, and k is the

slope factor (in mV). The time course of current inactivation was fitted by a double-exponential

equation: $I/I_{max}=A_f \times \exp(-t/\tau_f)+A_s \times \exp(-t/\tau_s)$, where A_f and A_s are the fractions of the fast and slow

inactivation components, and τ_f and τ_s are the time constants of the fast and slow inactivating

components, respectively.

Transient outward K⁺ current measurements. I_{to1} was measured under conditions similar as

mentioned for APs, but 0.25 mmol/L $CdCl_2$ was added to the modified Tyrode's to blocks $I_{Ca,L}$[68]

which also strongly inhibits I_{Na}[69]. Suppression of these inward currents allows a more accurate

determination of I_{to1}. I_{to1} was measured using a double pulse protocol from a holding potential of

-80 mV (see inset Figure 6D; cycle length of 5 seconds). The first pulse was used to determine

the I-V relationship; the second pulse was used to analyze the voltage dependency of inactivation.

I_{to1} was defined as the difference between peak and steady state current and densities were

calculated by dividing current amplitude by C_m. Voltage dependence of inactivation curves was

fitted with the above-mentioned Boltzmann function and time course of current inactivation was

fitted by a mono-exponential equation: $I/I_{max}=A_f \times \exp(-t/\tau_f)$.

Analysis of human GTEx and snRNAseq data

The Genotype-Tissue Expression (GTEx) data were obtained from left ventricle *Transcripts Per*

Million (TPM) on 2022-06-06 v10. Samples were enriched in cardiomyocytes using the *Gene Set*

Variation Analysis (GSVA) tool[70] (RRID:SCR_021058). The heatmap was generated from

centered and scaled data, and a Pearson correlation was performed on TPM values, with p-values

adjusted for multiple hypothesis testing (number of tests > 2) using *Benjamini-Hochberg*

correction.

For the correlation between *WT1* and *SCN5A* in single-cell RNA-seq data from hiPSC-CMs, we

used the dataset published by Canac *et al.* (2022)[71]. Cells were differentiated following the

Matrigel sandwich method[72] and sequenced at day 30 using the *Multi Tissue dissociation kit 3*

(Miltenyi Biotec) in duplicate. For these analyses, cells were filtered with a minimum of 250

features and a maximum of 8200, based on the sample distribution, as well as a maximum of 7%

mitochondrial gene. Doublets were removed using *DoubletFinder*[73] (RRID:SCR_018771). Data

were then combined into a single SeuratV5 object[74] (RRID:SCR_016341). Normalization with

regression of cell cycle, number of features, and mitochondrial percentage was performed using

SCTransform[75] (RRID:SCR_022146). Cells were subsequently integrated using the *RPCA*

method and annotated via *Azimuth* from SeuratV5 package with reference of human embryonic

heart cell atlas at 7 PCW published by Asp *et al.* (2019)[76]. Correlation analysis was performed

similarly as for the GTEx data, using the Seurat data matrix.

To assess the relationship between *SCN5A* and *WT1* expression in human heart tissue, we

leveraged the Reichart et al. dataset. Reichart et al. performed snRNAseq on 61 explanted end-

stage cardiomyopathy hearts, as well as 18 hearts from non-failing donors. The authors

performed snRNAseq across various ventricular regions, including the left ventricle, the right

ventricle, the interventricular septum, and the apex. For our analyses, we restricted to the 18 non-

failing hearts and removed nuclei with 'unknown' annotations, leaving a total of 266,219 high-

quality nuclei. For the regression analyses, we performed pseudo-bulking on the count matrix,

where counts for each gene were collapsed per donor/region combination. Lowly-expressed

genes were removed using the function *filterByExpr()* from the *edgeR* package (using design: ~0

+ region + sex + development_stage), after which *SCN5A* and *WT1* remained in the dataset.

Count   data   were   normalized   using   the   *DESeq2*   package,   using   functions

DESeqDataSetFromMatrix(),   *estimateSizeFactors()*,   and   finally   *counts()*   with   flag

'normalized=TRUE'. To regress the normalized expression of *WT1* on the normalized expression

of *SCN5A*, we utilized a mixed negative binomial regression model, implemented in R-package

glmmTMB, with the formula: *WT1 ~ SCN5A* + sex + development_stage + region + (1 | donor).

Histology

Human RV, LV and interventricular septum 7 μm histological sections were obtained from the

department of pathology at Amsterdam UMC. Sections were stained with a WT1-antibody

(Abcam/Epitomics AC9115, clone EP122) diluted 1:50 using HIER (heat-induced epitope

retrieval) (Tris-EDTA, pH 9.0) at 100°C. Sections were examined with a Leica imaged on a

DSM5000 widefield microscope.

Adult mouse hearts were fixed overnight in 4% paraformaldehyde in PBS, dehydrated in a

ascending ethanol series, paraffin embedded and sectioned at 8 µm. Sections were

deparaffinized with xylene, rehydrated in descending ethanol series and treated with HIER

(citrate-based antigen unmasking solution, Vector, H-3300-250) at 100°C. After permeabilization

(Triton-X 0.5%) and blocking (BSA 5%), sections were stained with anti-cTnT (1:400, HyTest Ltd

4T19/2), anti-GFP (1:300, Abcam Cat ab13970), anti-WGA-555 (1:200, ThermoFisher W32464)

and DAPI. Stained sections were imaged and photographed with a Leica DM6 microscope.

Flow cytometry

Cell isolations for flow cytometry experiments were carried out according to a previously published

protocol[77]. In short, after extracting the heart, the ascending aorta was clamped and a collagenase

buffer was injected into the apex of the left ventricle. Then the atria were removed and the right

ventricular free wall, left ventricular free wall and septum were separately dissociated by

trituration. After suspending cells in stop solution (containing FBS), they were passed a 100µm

strainer, centrifuged at low speed and stained for the cardiac marker cTnT. For the latter, cells

were incubated 15 minutes in permeabilizing-blocking buffer (PBS with 0.1% saponin and 5%

FBS), incubated with anti-cTnT (HyTest Ltd Cat 4T19/2, 1:400) for 1 hour and with secondary

antibody (AF647 donkey anti-goat IgG H+L, Life A21447, 1:200) for 15 minutes, all at room

temperature. Cells were confirmed to be all cardiomyocytes using a negative control without

primary antibody. For compensation control, littermates without Cre were used. Cells were

dissolved in FACS buffer (PBS with 0.5% BSA and 2mM EDTA) and measured with a

LSRFortessa Flow Cytometer (BD Biosciences). FlowJo was used for analysis. Forward and side

scatter information was used for gating of non-debris single cells and from these cells, the amount

of endogenous tdT positive and GFP positive cells was quantified. For each experiment $Scn5a^{+/+}$

and $Scn5a^{+/-}$ hearts were used in parallel, with researchers being blinded for the genotype during

experiment and analysis.

Statistics

The experimenters were blind to genotypes during all measurements and outcome assessment.

Datasets were tested for normality using Shapiro-Wilk test unless specified otherwise. Whole

isolated tissue RT-qPCR were analyzed with ANOVA with Welch's correction followed by

unpaired t-tests with Welch's correction (Figure 1), or 2-way ANOVA followed by Šídák's multiple

comparisons test and adjusted p-values are shown (Figure 5). *In vivo* electrophysiology and

optical mapping measurements were analyzed with unpaired t tests for 2 group comparisons, and

2-way ANOVA followed by Tukey's multiple comparison test for aging experiments comparing 3

groups. For single cell electrophysiology, data are expressed as mean±SEM. Statistical analysis

was carried out with SigmaStat 3.5 software (Systat Software, Inc., San Jose, CA, USA).

Normality and equal variance assumptions were tested with the Kolmogorov–Smirnov and the

Levene median test, respectively. Significance between parameters of two groups was tested

using unpaired Student's t-test, or in case of a failed normality and/or equal variance test, by

Mann-Whitney Rank Sum Test. Data of I-V relationships was tested for significance using the

two-way repeated measures ANOVA followed by pairwise comparison using the Student–

Newman–Keuls test. $P < 0.05$ defines statistical significance.

For the flow cytometry data, a general linear mixed model was used after rank transformation of

GFP percentages due to non-normally distributed residuals. Bonferroni post-hoc pairwise

comparisons of regions and Mann-whitney U pairwaise tests of genotypes were used.

Data are presented as individual data points and mean or mean ± standard error of the mean

(SEM) and $p < 0.05$ defines statistical significance. Statistical analysis was performed using

GraphPad Prism 10 and SPSS.

 Figure Legends

 Figure 1. <u>Diminished *Wt1* leads to increased *Scn5a*.</u> A) Schematic depicts approach for isolating

 cardiomyocyte (CM) nuclei of the RV and RVOT. B) Volcano plot shows differentially expressed

 transcripts between control (n=4) and *Wt1*$^{+/-}$ (n=4) cardiomyocytes. Dark red and light red dots

 indicate significantly upregulated genes by raw p-value and p-adjusted for multiple testing,

 respectively. Correspondingly, dark blue and light blue dots indicate downregulated genes. C)

 Gene ontology (GO) analysis of up-and downregulated genes in *Wt1*$^{+/-}$ cardiomyocytes. P-values

 were adjusted for multiple testing using the false discovery rate (FDR) method of Benjamini-

 Hochberg. D) Heatmap depicting deregulated genes and their adjusted p-values (all p<0.05)

 involved in electrical conduction. Scale denotes fold change. E, F) *Hprt*-corrected expression of

 Scn5a (E) and *Wt1* (F) from adult control (n=6) and *Wt1*$^{+/-}$ (n=6) ventricular (sub)endocardium and

 (sub)epicardium determined by RT-qPCR. Significant p-values were determined with Welch's

 ANOVA followed by unpaired *t*-tests with Welch's correction.

 Figure 2. <u>Reduced *Wt1* expression improves conduction in the setting of severely reduced</u>

 <u>conduction in aged *Scn5a*$^{+/-}$ mouse hearts</u>. A) Typical examples of RV longitudinal and

 transversal- and B) RVOT conduction with respective average conduction velocity (CV) of aged

 control (n= 10), *Scn5a*$^{+/-}$ (n= 9) and *Scn5a*$^{+/-}$;*Wt1*$^{+/-}$ (n= 9). RV, right ventricle; RVOT, right

 ventricular outflow tract. C) Typical examples and average CV of RV longitudinal and transversal

 direction, and D) RVOT in aged *Scn5a*$^{+/-}$ and *Scn5a*$^{+/-}$;*Wt1*$^{+/-}$ mice after acute administration of

 ajmaline (n=7). Statistical significance was determined with ANOVA followed by pairwise Tukey's

 multiple comparison tests for A and B, and unpaired *t*-tests for C and D.

Figure 3. <u>A negative correlation between *WT1* and *SCN5A* expression.</u> A) UMAP of single-cell

RNA-seq data from control hiPSC-CMs, annotated using a 7 post-conception week (PCW) human

embryonic heart reference dataset, published by[76]. Cardiomyocytes (CMs) are displayed on the

UMAP as ventricular (vCMs) or atrial (aCMs). B) UMAP of vCMs and aCMs. Cells expressing

SCN5A only, *WT1* only, and both *SCN5A* and *WT1* are displayed in pink, blue, and black,

respectively. C) Contingency table showing the Chi-square test residuals for cells expressing (1)

or not expressing (0) *SCN5A* and *WT1*. Negative and positive residual values indicate that the

corresponding cell proportions are under- or over-represented compared to the expected

distribution, respectively. D) Pearson correlation analysis between *SCN5A* and *WT1* expression

in all *WT1*+ cells (top), all *SCN5A*+ cells (middle), and cells expressing at least one of the two

genes (*bottom*). Cells are ordered by increasing *WT1* expression: blue, indicates absence of

expression, red, indicates highest level of expression and pink, indicates intermediate level of

expression.

Figure 4. <u>Increased *Wt1*-lineage cells in *Scn5a* haploinsufficient hearts</u>. A) Percent GFP+/cTnT+

cells in LV, IVS, and RV in control (n= 5) and *Scn5a*[+/-] (n=6) mice as determined by FACS.

Significance determined by a general linear mixed model and post-hoc tests. B-D) Representative

images of histological sections of control and *Scn5a*[+/-] hearts. LV, left ventricle; RV, right ventricle;

IVS, interventricular septum.

Figure 5. <u>Deletion of BrS-associated region in mouse orthologue leads to increased *Wt1*</u>

<u>selectively in (sub) epicardium of RV.</u> A) UCSC browser view of the human BrS-associated locus

in chr11 including ATACseq from hiPS ventricular cardiomyocytes (black), EMERGE (red), CTCF

ChIPseq (red bars), and conservation track. B) Mouse orthologue of the human region shown in

panel B including ATACseq of ventricular cardiomyocytes and Hcn4+ sinoatrial node (SAN) cells,

H3K27ac ChIPseq from whole adult hearts, and EMERGE. C) Zoom-in of panel B depicting the

deleted region (BAR) in blue. D) Reference gene-corrected expression of *Wt1* from isolated (sub)

epicardial and (sub)endocardial sides of the RV and LV determined by RTqPCR of *Wt1$^{BAR/BAR}$*

(n=9) and control littermates (n=9). Statistical significance was determined with 2-way ANOVA

followed by pairwise comparisons with Tukey's multiple comparison tests. P values are shown.

Figure 6. WT1 overexpression in hiPSC-cardiomyocytes reduces sodium current (I_{Na}) and action

potential (AP) upstroke velocity. A) Voltage clamp protocol and representative traces of I_{Na}

activated during depolarizing steps from -120 to -20 mV. B) Average current-voltage (I-V)

relationships of I_{Na}. C) Average voltage dependence of (in)activation curves. Solid lines

represents the fitted Boltzmann function ($I/I_{max}=A/\{1.0+exp[(V_{1/2} −V)/k]\}$). D) Voltage clamp

protocol and representative traces of the transient outward potassium current (I_{to1}) activated

during depolarizing steps from -80 to 60 mV. E) Average I-V relationships of I_{to1}. F) Average

voltage dependence of (in)activation curves. Solid lines represent the fitted Boltzmann function

for inactivation. G) Typical APs measured at 1 Hz. The inset shows the AP upstroke velocity

(V_{max}). **H.** Average AP parameters. RMP, resting membrane potential; APA, AP amplitude; APD_{20}

APD_{50}, and APD_{90}, AP duration at 20, 50 and 90% of repolarization, respectively; * Indicates

P<0.05 by Two-Way ANOVA or unpaired t-test.

References

1. Marsman, E. M. J., Postema, P. G. & Remme, C. A. Brugada syndrome: update and future

perspectives. *Heart* **108**, 668–675 (2022).

2. Offerhaus, J. A., Bezzina, C. R. & Wilde, A. A. M. Epidemiology of inherited arrhythmias. *Nat. Rev.*

Cardiol. **17**, 205–215 (2020).

3. Priori, S. G. *et al.* 2015 ESC Guidelines for the management of patients with ventricular arrhythmias

and the prevention of sudden cardiac death: The Task Force for the Management of Patients with

Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death of the European Society of

Cardiology (ESC)Endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC).

Eur. Heart J. **36**, 2793–2867 (2015).

4. Miles, C. *et al.* Subepicardial Cardiomyopathy: A Disease Underlying J-Wave Syndromes and

Idiopathic Ventricular Fibrillation. *Circulation* **147**, 1622 (2023).

5. Boukens, B. J. *et al.* Reduced Sodium Channel Function Unmasks Residual Embryonic Slow

Conduction in the Adult Right Ventricular Outflow Tract. *Circ. Res.* **113**, 137–141 (2013).

6. Miles, C. *et al.* Biventricular Myocardial Fibrosis and Sudden Death in Patients With

Brugada Syndrome. *J. Am. Coll. Cardiol.* **78**, 1511–1521 (2021).

7. Chen, Q. *et al.* Genetic basis and molecular mechanism for idiopathic ventricular fibrillation. *Nature*

392, 293–296 (1998).

8. Le Scouarnec, S. *et al.* Testing the burden of rare variation in arrhythmia-susceptibility genes

provides new insights into molecular diagnosis for Brugada syndrome. *Hum. Mol. Genet.* **24**, 2757–

2763 (2015).

9. Hosseini, S. M. *et al.* Reappraisal of Reported Genes for Sudden Arrhythmic Death: Evidence-Based

Evaluation of Gene Validity for Brugada Syndrome. *Circulation* **138**, 1195 (2018).

10. Barc, J. *et al.* Genome-wide association analyses identify new Brugada syndrome risk loci and

highlight a new mechanism of sodium channel regulation in disease susceptibility. *Nat. Genet.* 1–8

(2022) doi:10.1038/s41588-021-01007-6.

11. Bezzina, C. R. *et al.* Common variants at SCN5A-SCN10A and HEY2 are associated with Brugada

syndrome, a rare disease with high risk of sudden cardiac death. *Nat. Genet.* **45**, 1044–1049 (2013).

12. Deplancke, B., Alpern, D. & Gardeux, V. The Genetics of Transcription Factor DNA Binding Variation.

Cell **166**, 538–554 (2016).

13. Maurano, M. T. *et al.* Systematic localization of common disease-associated variation in regulatory

DNA. *Science* **337**, 1190–5 (2012).

14. Schaub, M. A., Boyle, A. P., Kundaje, A., Batzoglou, S. & Snyder, M. Linking disease associations with

regulatory information in the human genome. *Genome Res.* **22**, 1748–59 (2012).

15. Veerman, C. C. *et al.* The Brugada Syndrome Susceptibility Gene HEY2 Modulates Cardiac

Transmural Ion Channel Patterning and Electrical Heterogeneity. *Circ. Res.* **121**, 537–548 (2017).

16. Chiang, D. Y. *et al.* The Role of MAPRE2 and Microtubules in Maintaining Normal Ventricular

Conduction. *Circ. Res.* https://doi.org/10.1161/CIRCRESAHA.123.323231 (2024)

doi:10.1161/CIRCRESAHA.123.323231.

17. Zhou, B. *et al.* Epicardial progenitors contribute to the cardiomyocyte lineage in the developing

heart. *Nature* **454**, 109–113 (2008).

18. Wessels, A. *et al.* Epicardially derived fibroblasts preferentially contribute to the parietal leaflets of

the atrioventricular valves in the murine heart. *Dev. Biol.* **366**, 111–124 (2012).

19. Duim, S. N., Goumans, M.-J. & Kruithof, B. P. T. WT1 in Cardiac Development and Disease. in *Wilms*

Tumor (ed. van den Heuvel-Eibrink, M. M.) (Codon Publications, Brisbane (AU), 2016).

20. Wagner, N. *et al.* Implications of the Wilms' Tumor Suppressor Wt1 in Cardiomyocyte

Differentiation. *Int. J. Mol. Sci.* **22**, 4346 (2021).

21. Tyser, R. C. V. *et al.* Characterization of a common progenitor pool of the epicardium and

myocardium. *Science* **371**, eabb2986 (2021).

22. Díaz del Moral, S. *et al.* Deletion of the Wilms' Tumor Suppressor Gene in the Cardiac Troponin-T

Lineage Reveals Novel Functions of WT1 in Heart Development. *Front. Cell Dev. Biol.* **9**, (2021).

23. van Wijk, B., Gunst, Q. D., Moorman, A. F. M. & van den Hoff, M. J. B. Cardiac Regeneration from

Activated Epicardium. *PLoS ONE* **7**, e44692 (2012).

24. Zhou, B. *et al.* Adult mouse epicardium modulates myocardial injury by secreting paracrine factors.

J. Clin. Invest. **121**, 1894–1904 (2011).

25. Wagner, K.-D. *et al.* The Wilms' tumor suppressor Wt1 is expressed in the coronary vasculature after

myocardial infarction. *FASEB J.* **16**, 1117–1119 (2002).

26. Braitsch, C. M., Kanisicak, O., van Berlo, J. H., Molkentin, J. D. & Yutzey, K. E. Differential expression

of embryonic epicardial progenitor markers and localization of cardiac fibrosis in adult ischemic

injury and hypertensive heart disease. *J. Mol. Cell. Cardiol.* **65**, 10.1016/j.yjmcc.2013.10.005 (2013).

27. Hill, M. C. *et al.* Integrated multiomic characterization of congenital heart disease. *Nature* 1–3

(2022) doi:10.1038/s41586-022-04989-3.

28. Reichart, D. *et al.* Pathogenic variants damage cell composition and single cell transcription in

cardiomyopathies. *Science* **377**, eabo1984 (2022).

29. Chaffin, M. *et al.* Single-nucleus profiling of human dilated and hypertrophic cardiomyopathy.

Nature **608**, 174–180 (2022).

30. Díaz del Moral, S. *et al.* Cardiomyocyte-Specific Wt1 Is Involved in Cardiac Metabolism and Response

to Damage. *J. Cardiovasc. Dev. Dis.* **10**, 211 (2023).

31. Mi, H., Muruganujan, A., Ebert, D., Huang, X. & Thomas, P. D. PANTHER version 14: more genomes,

a new PANTHER GO-slim and improvements in enrichment analysis tools. *Nucleic Acids Res.* **47**,

D419–D426 (2019).

32. Boukens, B. J., Walton, R., Meijborg, V. M. & Coronel, R. Transmural electrophysiological

heterogeneity, the T-wave and ventricular arrhythmias. *Prog. Biophys. Mol. Biol.* **122**, 202–214

(2016).

33. Flurkey, K., M. Currer, J. & Harrison, D. E. Chapter 20 - Mouse Models in Aging Research. in *The*

Mouse in Biomedical Research (Second Edition) (eds Fox, J. G. et al.) 637–672 (Academic Press,

Burlington, 2007). doi:10.1016/B978-012369454-6/50074-1.

34. Shaw, R. M. & Rudy, Y. Ionic Mechanisms of Propagation in Cardiac Tissue. *Circ. Res.* **81**, 727–741

(1997).

35. Poulsen, P. C. *et al.* Quantitative proteomics characterization of acutely isolated primary adult rat

cardiomyocytes and fibroblasts. *J. Mol. Cell. Cardiol.* **143**, 63–70 (2020).

36. Kohela, A. *et al.* Epicardial differentiation drives fibro-fatty remodeling in arrhythmogenic

cardiomyopathy. *Sci. Transl. Med.* **13**, eabf2750 (2021).

37. Muzumdar, M. D., Tasic, B., Miyamichi, K., Li, L. & Luo, L. A global double-fluorescent Cre reporter

mouse. *Genes. N. Y. N 2000* **45**, 593–605 (2007).

38. Protze, S. I. *et al.* Sinoatrial node cardiomyocytes derived from human pluripotent cells function as a

biological pacemaker. *Nat. Biotechnol.* **35**, 56–68 (2017).

39. Davis, C. A. *et al.* The Encyclopedia of DNA elements (ENCODE): data portal update. *Nucleic Acids*

Res. **46**, D794–D801 (2018).

40. Van Duijvenboden, K., De Boer, B. A., Capon, N., Ruijter, J. M. & Christoffels, V. M. EMERGE: A

flexible modelling framework to predict genomic regulatory elements from genomic signatures.

Nucleic Acids Res. **44**, (2015).

41. van Ouwerkerk, A. F. *et al.* Identification of atrial fibrillation associated genes and functional non-

coding variants. *Nat. Commun.* **10**, (2019).

42. Fernandez-Perez, A. *et al.* Hand2 Selectively Reorganizes Chromatin Accessibility to Induce

Pacemaker-like Transcriptional Reprogramming. *Cell Rep.*

https://doi.org/10.1016/j.celrep.2019.04.077 (2019) doi:10.1016/j.celrep.2019.04.077.

43. van Duijvenboden, K. *et al.* Conserved NPPB+ Border Zone Switches From MEF2- to AP-1-Driven

Gene Program. *Circulation* **140**, 864–879 (2019).

44. Gilsbach, R. *et al.* Distinct epigenetic programs regulate cardiac myocyte development and disease

in the human heart in vivo. *Nat. Commun.* **9**, (2018).

45. Feyen, D. A. M. *et al.* Metabolic Maturation Media Improve Physiological Function of Human iPSC-

Derived Cardiomyocytes. *Cell Rep.* **32**, 107925 (2020).

46. Verkerk, A. O. & Wilders, R. Injection of IK1 through dynamic clamp can make all the difference in

patch-clamp studies on hiPSC-derived cardiomyocytes. *Front. Physiol.* **14**, 1326160 (2023).

47. Man, J. C. K. *et al.* An enhancer cluster controls gene activity and topology of the SCN5A-SCN10A

locus in vivo. *Nat. Commun.* **10**, 4943 (2019).

48. Marchal, G. A. *et al.* Genetic background determines the severity of age-dependent cardiac

structural abnormalities and arrhythmia susceptibility in Scn5a-1798insD mice. *Eur. Eur. Pacing*

Arrhythm. Card. Electrophysiol. J. Work. Groups Card. Pacing Arrhythm. Card. Cell. Electrophysiol.

Eur. Soc. Cardiol. **26**, euae153 (2024).

49. Xing, S. *et al.* Genetic influence on electrocardiogram time intervals and heart rate in aging mice.

Am. J. Physiol. - Heart Circ. Physiol. **296**, H1907 (2009).

50. de Wit, E. *et al.* CTCF Binding Polarity Determines Chromatin Looping. *Mol. Cell* **60**, 676–684 (2015).

51. Dixon, J. R. *et al.* Topological domains in mammalian genomes identified by analysis of chromatin

interactions. *Nature* **485**, 376–80 (2012).

52. Nora, E. P. *et al.* Spatial partitioning of the regulatory landscape of the X-inactivation centre. *Nature*

485, 381–385 (2012).

53. Toska, E. & Roberts, S. G. E. Mechanisms of transcriptional regulation by WT1 (Wilms' tumour 1).

Biochem. J. **461**, 15–32 (2014).

54. Rampal, R. *et al.* DNA Hydroxymethylation Profiling Reveals that WT1 Mutations Result in Loss of

TET2 Function in Acute Myeloid Leukemia. *Cell Rep.* **9**, 1841–1855 (2014).

55. Wang, Y. *et al.* WT1 Recruits TET2 to Regulate Its Target Gene Expression and Suppress Leukemia

Cell Proliferation. *Mol. Cell* **57**, 662–673 (2015).

56. Essafi, A. *et al.* A Wt1-Controlled Chromatin Switching Mechanism Underpins Tissue-Specific Wnt4

Activation and Repression. *Dev. Cell* **21**, 559–574 (2011).

57. Larsson, S. H. *et al.* Subnuclear localization of WT1 in splicing or transcription factor domains is

regulated by alternative splicing. *Cell* **81**, 391–401 (1995).

58. Shandilya, J., Toska, E., Richard, D. J., Medler, K. F. & Roberts, S. G. E. WT1 interacts with MAD2 and

regulates mitotic checkpoint function. *Nat. Commun.* **5**, 4903 (2014).

59. von Gise, A. *et al.* WT1 regulates epicardial epithelial to mesenchymal transition through β-catenin

and retinoic acid signaling pathways. *Dev. Biol.* **356**, 421–431 (2011).

60. Bergmann, O. & Jovinge, S. Isolation of Cardiomyocyte Nuclei from Post-mortem Tissue. *J. Vis. Exp.*

JoVE 4–9 (2012) doi:10.3791/4205.

61. Dobin, A. *et al.* STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

62. Ruijter, J. M. *et al.* Amplification efficiency: Linking baseline and bias in the analysis of quantitative

PCR data. *Nucleic Acids Res.* **37**, (2009).

63. Doshi, A. N. *et al.* Feasibility of a semi-automated method for cardiac conduction velocity analysis of

high-resolution activation maps. *Comput. Biol. Med.* **65**, 177–183 (2015).

64. ten Hoope, W. *et al.* Pharmacodynamics and Pharmacokinetics of Lidocaine in a Rodent Model of

Diabetic Neuropathy. *Anesthesiology* **128**, 609 (2018).

65. Veerman, C. C. *et al.* hiPSC-derived cardiomyocytes from Brugada Syndrome patients without

identified mutations do not exhibit clear cellular electrophysiological abnormalities. *Sci. Rep.* **6**,

30967 (2016).

66. Verkerk, A. O. & Wilders, R. Dynamic Clamp in Electrophysiological Studies on Stem Cell–Derived

Cardiomyocytes—Why and How? *J. Cardiovasc. Pharmacol.* **77**, 267 (2021).

67. Meijer van Putten, R. M. E. *et al.* Ion channelopathies in human induced pluripotent stem cell

derived cardiomyocytes: a dynamic clamp study with virtual IK1. *Front. Physiol.* **6**, (2015).

68. Hobai, I. A., Bates, J. A., Howarth, F. C. & Levi, A. J. Inhibition by external Cd2+ of Na/Ca exchange

and L-type Ca channel in rabbit ventricular myocytes. *Am. J. Physiol.-Heart Circ. Physiol.* **272**,

H2164–H2172 (1997).

69. Sheets, M. F. & Hanck, D. A. Mechanisms of extracellular divalent and trivalent cation block of the

sodium current in canine cardiac Purkinje cells. *J. Physiol.* **454**, 299–320 (1992).

70. Hänzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for microarray and RNA-

Seq data. *BMC Bioinformatics* **14**, 7 (2013).

71. Canac, R. *et al.* Deciphering Transcriptional Networks during Human Cardiac Development. *Cells* **11**,

3915 (2022).

72. Es-Salah-Lamoureux, Z. *et al.* HIV-Tat induces a decrease in I Kr and I Ks via reduction in

phosphatidylinositol-(4,5)-bisphosphate availability. *J. Mol. Cell. Cardiol.* [Epub ahead of print]

(2016) doi:10.1016/j.yjmcc.2016.08.022.

73. McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: Doublet Detection in Single-Cell RNA

Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst.* **8**, 329-337.e4 (2019).

74. Hao, Y. *et al.* Dictionary learning for integrative, multimodal, and massively scalable single-cell

analysis. *Nat. Biotechnol.* **42**, 293–304 (2024).

75. Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-seq data

using regularized negative binomial regression. *Genome Biol.* **20**, 296 (2019).

76. Asp, M. *et al.* A Spatiotemporal Organ-Wide Gene Expression and Cell Atlas of the Developing

Human Heart. *Cell* **179**, 1647-1660.e19 (2019).

77. Ackers-Johnson, M. *et al.* A Simplified, Langendorff-Free Method for Concomitant Isolation of Viable

Cardiac Myocytes and Nonmyocytes From the Adult Mouse Heart. *Circ. Res.* **119**, 909 (2016).

Acknowledgements

We acknowledge the resources and support from the Core Facility Genomics of the Amsterdam UMC for the RNA sequencing experiments. We are thankful to Jaël Copier, Caroline Pham, and Aya Aliawi for their technical assistance. We thank Vincent Christoffels for insightful discussions which were helpful in the advancement of this work.

882 **Funding**

883 TS, NG and GL were funded by Agence Nationale pour la Recherche (ANR-22-CE17-

884 0051) and Fédération Française de Cardiologie (FIBRIL R21173NN).

A Adult 15+ months

B *Control* vs *Wt1+/-* PCM1+

Down in *Wt1+/-* 405 by P-adj
Up in *Wt1+/-* 466 by P-adj

C

GO biological process upregulated in *Wt1+/-* CM	Fold Enrichment	Raw P-value	FDR
cellular component organization (GO:0016043)	1.75	1.31E-15	4.14E-12
developmental process (GO:0032502)	1.71	1.89E-15	4.97E-12
regulation of cell communication (GO:0010646)	2.02	7.85E-15	1.03E-11
cell differentiation (GO:0030154)	1.78	1.35E-10	6.43E-08
regulation of gene expression (GO:0010468)	1.62	9.85E-10	3.98E-07
regulation of cardiac muscle contraction (GO:0055117)	7.61	2.09E-06	2.59E-04
regulation of transport (GO:0051049)	1.84	2.55E-06	3.04E-04
cell motility (GO:0048870)	1.90	2.30E-04	1.25E-02

GO biological process downregulated in *Wt1+/-* CM	Fold Enrichment	Raw P-value	FDR
cellular respiration (GO:0045333)	6.67	3.63E-10	1.43E-06
metabolic process (GO:0008152)	1.41	1.66E-08	3.73E-05
cellular localization (GO:0051641)	1.81	2.34E-07	3.35E-04
muscle tissue development (GO:0060537)	3.59	9.88E-07	1.04E-03
intracellular transport (GO:0046907)	1.99	3.86E-05	1.17E-02
electron transport chain (GO:0022900)	5.67	5.42E-05	1.50E-02
apoptotic process (GO:0006915)	2.07	1.47E-04	2.61E-02

D

Kcnip2	3,1E-05
Cacna1s	0,00881
Clcn7	0,04144
Kcnq1	0,00109
Chrna10	0,01176
Cacna1c	1,6E-07
Abcc8	0,00025
Itpr2	1,1E-06
Scn5a	0,01242
Ryr2	3E-05
Atp1b3	0,01465
Slc22a4	0,0014
Kcnh7	0,03188

E W ANOVA p=0.0114

F W ANOVA p=0.0009

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Supplemental Data

Supplemental figure legends

Supplemental Figure 1. <u>Heterozygous loss of *Wt1* (*Wt1*^{+/-}) does not affect cardiac conduction in</u>

<u>adult mice</u>. A) Typical ECGs and average ECG parameters of control (n = 7-12) and *Wt1*^{+/-} (n =

6-8) mice. B) Typical examples and average conduction velocities of longitudinal and transversal

conduction in RV, and C) transversal conduction in RVOT in control (n = 9-10) and *Wt1*^{+/-} (n = 9)

mice. RV, right ventricle; RVOT, right ventricular outflow tract. Statistical significance was

determined with unpaired t-tests.

Supplemental Figure 2. <u>Heterozygous loss of *Wt1* does not affect electrical properties of single</u>

<u>adult RV cardiomyocytes.</u> A, B) Typical action potentials (AP; A) measured at 4 Hz and average

AP characteristics (B) of control (n = 26) and *Wt1*^{+/-} (n = 22) RV cardiomyocytes. C-E) Typical

sodium currents (I_{Na}; C), average I_{Na} current-voltage relationships (D), and voltage dependence

of (in)activation (E) of control (n = 6) and *Wt1*^{+/-} (n = 6) isolated RV cardiomyocytes. Dotted lines

are Boltzmann fits. The inset in panel C shows the used voltage-clamp protocol to measure I_{Na}.

RMP, resting membrane potential; APA, AP amplitude; V_{max}, maximal AP upstroke velocity; APD,

AP duration; RV, right ventricle. Statistical significance was determined with two way RM ANOVA

or unpaired t-tests.

Supplemental Figure 3. <u>Effect of aging and diminished *Wt1* on ECG parameters of *Scn5a*^{+/-}</u>

<u>mice.</u> A) Typical ECG traces and average duration of B) PR- and C) QRS duration of adult and

aged control (n= 12 and 17, respectively) *Scn5a*^{+/-} (n=7 and 11, respectively) and *Scn5a*^{+/-};*Wt1*^{+/-}

(n=7, 9 respectively) mice. Significance was determined using 2-way ANOVA shown in the tables

below each set of graphs, followed by pairwise Tukey's multiple comparisons tests (shown in

graphs).

Supplemental Figure 4. <u>High *WT1* expression correlates with decreased *SCN5A* expression.</u> A) Pearson correlation analysis of left ventricular samples from GTEx dataset using a selection of a CM 10-gene signature, as reported by[35] (***: p-value<0.001). B) Distribution of GTEx sample based on their CM enrichment scores, using the 10-gene signature (Gene Set Variation Analysis method, GSVA R package). Samples were classified into "high" (orange) and "low" (yellow) CM-enrichment groups based on the bimodal distribution of scores. C) Heatmap displaying the expression values of left ventricular samples from GTEx dataset for each of the 10 genes that compose the CM signature. Blue and red indicate low and high expression values, as compared to the mean for each gene, respectively. Dendrogram was generated using Ward's hierarchical clustering method with Euclidean distances. D) Pearson correlation of *SCN5A* and *WT1* expression for the "high" and the "low" CM-enrichment groups. E) Partial regression plot for mixed negative binomial model between *WT1* and *SCN5A* from 18 healthy donors across 4 ventricle regions in pseudo-bulk. Axes represent the covariate-adjusted log-transformed expression of *WT1* and *SCN5A.* Corresponding statistics are added, where r_s represents the Spearman's correlation coefficient, while z and p represent the Z-value and P-value from the mixed negative binomial regression respectively. F) Univariate negative binomial model plot with untransformed data from E.

Supplemental Figure 5. <u>Conduction-related genes altered in cardiomyocytes of *Wt1*[+/-] mice, also correlate with *WT1* expression in human left ventricular samples</u>. Pearson correlation analysis of *WT1* expression and of the conduction-related genes altered in cardiomyocytes of *Wt1*[+/-] mice (Figure 1D), using human left ventricular samples from GTEx database. *: p-value<0.05; ***: p-value<0.001. All p-values were adjusted for multiple hypothesis testing using Benjamini-Hochberg correction.

Supplemental Figure 6. <u>Immunohistochemistry for WT1 in cardiac sections of patients with *SCN5A* mutation and non-diseased hearts.</u> A-K) Representative images of *SCN5A* mutation

carriers (A-G) and non-diseased (H-K) left ventricles (LV (A, D, F, H, J), RVs (B, E, G, I, K), and

interventricular septum (IVS) (C) histological sections. L) Staining control in human kidney.

Arrowheads denote nuclear staining. Asterisk denotes non-specific signal.

Supplemental Table 1. DEseq2 analysis of control vs $Wt1^{+/-}$ cardiomyocytes.

Supplemental Table 2. Average sodium current (I_{Na}) characteristics of cardiomyocytes isolated

from the right ventricle of control and $Wt1^{+/-}$ mice.

	Sodium Current		
Parameter	Control (n=6)	$WT1^{+/-}$(n=6)	P value
Maximal peak I_{Na} (pA/pF)	-52.5 ± 8.6	-46.9± 4.8	0.58
$V_{1/2}$ activation (mV)	-45.55 ± 1.26	-46.25 ± 1.28	0.706
k activation (mV)	4.91± 0.32	5.21± 0.31	0.519
$V_{1/2}$ inactivation (mV)	-83.23 ± 1.98	-84.21 ± 0.89	0.661
k inactivation (mV)	-6.20± 0.32	-6.11± 0.46	0.885

56   **Supplemental Table 3.** Region-specific $WT1$-$SCN5A$ correlation analysis in all cell-types.

Region	N donors	N cells	N donor region combinations	Spearman Rho	Beta	SE	Z	P
All	18	266219	51	-0,4699548	-4,93E-04	1,00E-04	-4,9260962	8,39E-07
IVS	12	51299	12	-0,3286713	-5,22E-04	5,71E-04	-0,9148981	3,60E-01
RV	15	58779	15	-0,5214286	-7,25E-04	3,86E-04	-1,8766413	6,06E-02
LV	18	132742	18	-0,246646	-2,40E-04	7,16E-05	-3,3486163	8,12E-04
Apex	6	23399	6	-0,8857143	-9,32E-04	1,78E-04	-5,246449	1,55E-07

57   Model: expression_gene1~expression_gene2+gene+sex+age

58   **Supplemental Table 4**. I_{Na} and I_{to1} parameters in hiPSC-cardiomyocytes.

		GFP	WT1-P2A-GFP
I_{Na}		16 cells, 4 diff.	12 cells, 5 diff.
	Density at -20 mV (pA/pF)	-86.8 ± 10.3	-54.0 ± 7.2
	$V_{1/2}$ of activation (mV)	-33.6 ± 1.5	-35.2 ± 1.8
	k of activation (mV)	6.7 ± 0.3	6.0 ± 0.5
	$V_{1/2}$ of inactivation (mV)	-82.9 ± 2.4	-82.4 ± 2.3
	k of inactivation (mV)	-5.9 ± 0.3	-6.2 ± 0.3
	τ_{slow} at -20 mV (ms)	7.1 ± 0.5	6.2 ± 0.7
	τ_{fast} at -20 mV (ms)	0.94 ± 0.07	0.92 ± 0.05
	$A_{slow}/(A_{slow}+A_{fast})$ at -20 mV (%)	8.8 ± 0.8	8.6 ± 0.5
I_{to1}		7 cells, 3 diff.	7 cells, 3 diff.
	Density at 60 mV (pA/pF)	15.8 ± 5.8	17.9 ± 2.9
	$V_{1/2}$ of inactivation (mV)	-43.9 ± 6.0	-45.0 ± 7.6
	k of inactivation (mV)	-13.0 ± 4.1	-11.4 ± 3.5
	τ (ms)	25.5 ± 6.1	20.4 ± 5.3

59 Data are average$\pm$SEM. I_{Na} = Na$^+$ current, I_{to1} = transient outward K$^+$ current, diff.= differentiations,

60 $V_{1/2}$ = half-maximal voltage of (in)activation, k = the slope factor of (in)activation, τ = time constant

61 of inactivation, and A = Amplitude. * P<0.05.

Supplemental Figure 1

Supplemental Figure 2

Supplemental Figure 3

Supplemental Figure 4

A

Supplemental Figure 5

Supplemental Figure 6