A new deep-branching environmental lineage of algae

Mahwash Jamy[1], Thomas Huber[2], Thibault Antoine[3], Hans-Joachim Ruscheweyh[4], Lucas Paoli[5], Eric

Pelletier[3], Tom O. Delmont[3]*, Fabien Burki[2]*

[1]Department of Aquatic Sciences and Assessment, Swedish University of Agricultural Sciences,
Uppsala, Sweden
[2]Department of Organismal Biology (Systematic Biology), Uppsala University, Uppsala, Sweden
[3]Génomique Métabolique, Genoscope, Institut François Jacob, CEA, CNRS, Univ Evry, Université
Paris-Saclay, 91057 Evry, France
[4]Department of Biology, Institute of Microbiology and Swiss Institute of Bioinformatics, ETH Zürich,
8093, Zürich, Switzerland
[5]Global Health Institute, School of Life Sciences, École Polytechnique Fédérale de Lausanne (EPFL),
1015 Lausanne, Switzerland
14

*co-corresponding authors

Abstract
Marine algae support the entire ocean ecosystem and beyond. Algae in culture poorly represents their
large environmental diversity, and we still have a limited understanding of their convoluted evolution
by endosymbiosis. Here, we performed a phylogeny-guided plastid genome-resolved metagenomic
survey of *Tara* Oceans expeditions. We present a curated resource of 660 new non-redundant plastid
genomes of environmental pelagic algae. This catalogue vastly expands the plastid genome diversity
within major algal groups, often corresponding to algae without closely related reference genomes.
Notably, we recovered four plastid genomes, including one near complete, forming a deep-branching
plastid lineage of nano-size algae that we informally named leptophytes. This group is globally
distributed and generally rare, although it can reach relatively high abundance at least in the Arctic. A
mitochondrial contig including 62 genes showing strong read coverage correlation with leptophytes
was also recovered from these Arctic samples and assigned to this group. Leptophytes encompass the
enigmatic marine plastid group DPL2, one of the very few known plastid groups not clearly belonging
to any major algal groups and for which only 16S rDNA amplicon data is available. Extensive gene
content comparison and organellar phylogenomics support the view that leptophytes are sister to
haptophytes, and raise the intriguing possibility that cryptophytes acquired their plastids from
haptophytes. Collectively, our study demonstrates that metagenomics can reveal currently hidden
diversity of organellar genomes, and shows the importance of including this diversity to improve
models for plastid evolution.

Key words: Plastid evolution, endosymbiosis, plankton, *Tara* Oceans, genome-resolved
metagenomics, ptMAGs, leptophytes, cryptophytes, haptophytes, red alga
39

Main text

Eukaryotic algae are essential to the success of life on earth, performing nearly 40% of primary production and forming the basis of many food webs[1]. Algae are incredibly diverse in forms and functions, ranging from some of the smallest known eukaryotes in the open ocean phytoplankton to undersea forests of giant coastal seaweed. The phylogenetic position of algae in the eukaryotic tree of life shows that they have evolved in multiple supergroups, owing their origins to a complex and largely unresolved history of endosymbioses. The first eukaryotic algae originated from the endosymbiotic integration of cyanobacteria into predatory host cells. This primary endosymbiosis led to the diversification of eukaryotes with photosynthetic organelles called primary plastids in the Archaeplastida supergroup, containing major algal groups such as red algae and green algae (including land plants)[2,3]. Subsequently, plastids were transferred from red and green algae to unrelated groups of eukaryotes by further endosymbioses, spreading photosynthetic capabilities throughout the tree of eukaryotes[4]. Major questions related to plastid evolution remain open, notably about the number of endosymbioses and partners involved to explain the observed phylogenetic distribution of plastids. The evolutionary history of complex plastids with red algal origin has been among the most difficult to resolve, in spite of the ecological importance of some of these algae in modern oceans, such as diatoms, dinoflagellates, and coccolithophores[5,6]. Multiple hypotheses have been proposed, many involving successive endosymbioses between eukaryotes over vast evolutionary distances, but there is no consensus and very little empirical evidence to support one or another hypothesis[2,7–9].

A more complete catalogue of plastid genomes across the algal tree is needed to better understand the evolution of plastid endosymbiosis. In recent years, comparatively more data has been produced for nuclear genomes of algae—including many transcriptomes—than for the smaller and in principle easier to sequence plastid genomes[10]. There are currently several major groups of algae that have poor representation of plastid data. For example, only 17 plastid genomes are available for haptophyte species, one of the most abundant and bloom-forming marine algal groups containing as many as 1000 described species and a much larger environmental diversity[5]. Beyond known algae, a few so-called deep-branching plastid groups, *i.e.,* plastids that do not belong to any established algal groups, have been reported but currently lack any genomic or morphologic description. The discovery of deep-branching plastid groups is extremely rare, comparable to the discovery of new animal groups[11], owing to a good understanding of what the major algal groups are. The most notable cases of new deep-branching groups in the past 20 years include the rappemonads—but they have since been formally assigned to haptophytes into the new class Rappephyceae[12]—or the enigmatic environmental plastid lineages DPL1 and DPL2 for which only 16S rDNA amplicon sequences are available[13]. While DPL1 was weakly placed within haptophytes but outside of known classes, DPL2 branched

deeper, possibly as sister to haptophytes but without statistical support[13]. Finally, the Picozoa

(previously picobiliophytes), which were originally described as a new algal group with unknown

affinities to eukaryotes, are now considered aplastidic members of Archaeplastida[14].

Here, we used 280 billion metagenomic reads covering a wide range of *Tara* Oceans planktonic size

fractions[15,16] as well as additional resources[17,18] to assemble and manually curate 660 non-redundant

environmental plastid genomes (ptMAGs). We show that genome-resolved metagenomics is a

powerful approach for reconstructing organellar genomes and use this unprecedented plastid genomic

resource to determine the phylogeny and geographical distribution of planktonic algae. Strikingly, we

report a generally rare but widespread and deep-branching clade of plastid genomes encompassing the

enigmatic DPL2 that we informally name leptophytes. From the few Arctic samples where

leptophytes were the most abundant, we also recovered a mitochondrial contig that was assigned to

this group based on its phylogenetic position and highly significant positive correlation in read

coverage. Based on the phylogenetic position of both plastid and mitochondrial genomes, we discuss

the implications of leptophytes for our understanding of plastid endosymbioses.

Results

A large resource of environmental plastid genomes from the sunlit ocean

To characterize the diversity of eukaryotic algae in the global sunlit ocean, we surveyed large *Tara*

Oceans metagenomic co-assemblies (~12 million contigs >2.5 kb) covering plankton size fractions

ranging from 0.8 µm to 2mm (Supplementary Table 1)[15,16]. These co-assemblies, each deriving from

samples from a particular oceanic basin, have been shown to improve the recovery of genomes that

are too scarce to be assembled from individual samples[16,19]. Briefly, we used a phylogeny-guided

metagenomic approach based on the RNApolB gene, and complementary genomic and environmental

information to characterize and manually curate an initial set of 1,448 plastid metagenome-resolved

genomes (ptMAGs) with anvi'o[20,21] (see Methods). Two additional single-contig ptMAGs related to

the focal clade of our study (see subsequent sections) were subsequently added from mOTUs-db[18].

After filtering for redundancy (average nucleotide identity >98%), our final plastid genome database

derived from all oceanic regions consisted of 660 new ptMAGs, as well as 166 reference plastid

genomes (Supplementary Fig. 1, Supplementary Table 1). The occurrence of 44 single copy plastid

core genes indicated that the ptMAGs are 62.9% complete on average, with low redundancy (average:

2.5%) (Supplementary Fig. 2). 126 ptMAGs contained at least 40 of the 44 core plastid genes and

were thus deemed highly complete. However, no ptMAG mapped as a circular sequence due to the

absence of the ribosomal RNA operon which is often missing from MAGs. The ptMAGs had a

median size of 65.7 kbp, with GC content ranging between 24.9% and 46.3%, and encoded up to 168

genes (Supplementary Table 1; Supplementary Figs. 3-5). The ptMAG abundance, as estimated by sequencing depth, varied by more than four orders of magnitude, ranging from a chlorophyte ptMAG with ~21,000× coverage to a diatom ptMAG with only ~2× coverage across all *Tara* Oceans metagenomes (Supplementary Table 1). Collectively, the ptMAGs were far more abundant in the sunlit oceans compared to selected culture representative genomes (82.5% of the total signal), in line with a previous comparison of nuclear genomes among the same metagenomes[16] (Supplementary Fig. 6).

We reconstructed a broad multigene phylogeny of our plastid genome database to assess its taxonomic diversity. Given the large taxon sampling, we inferred a Maximum Likelihood phylogeny using the relatively simple site-homogeneous LG+F+I+G4 model based on 93 plastid-encoded genes. This phylogeny recovered all major groups of algae and is overall in good agreement with previously published plastid trees (Fig. 1, Supplementary Fig. 7)[9,22]. The ptMAGs occurred in most, but not all, major groups of algae. We did not recover ptMAGs for red algae and glaucophytes, which are known to be rare or absent in the open ocean[23], nor for the divergent peridinin plastids of dinoflagellates, which are highly fragmented and lack the *rpoB* gene[24]. Conversely, the ptMAGs greatly expanded the known plastid genome diversity of ochrophytes (n=375, including 304 diatoms displaying a very broad cellular size range), and haptophytes (n=196, including Phaeocystales found in large size fractions likely due to their distinct colony-forming lifestyle[5]). A smaller number of ptMAGs were also recovered for cryptophytes (n=16, restricted to the small size fractions), and green algae (n=45, predominantly found in the smallest size fraction). In multiple cases, the ptMAGs represented deep-branching novel genome diversity within established algal groups where cultured references are lacking. For instance, one ptMAG (REFM_CHLORO_00002) deeply branched as sister to Pavlovales (Fig. 1 and Supplementary Fig. 7). Eight chlorophyte ptMAGs recovered from different oceanic regions formed a monophyletic clade distantly related to a clade of Chloropicophyceae. Within ochrophytes, 15 ptMAGs with a broad geographic distribution were closely related but clearly different from the reference pelagophyte plastid genomes (Supplementary Fig. 8). Finally, among green algae, 13 ptMAGs with a large variation in genome sizes formed a distinct clade related to pedinophyte endosymbionts of dinoflagellates, which include the undescribed MGD and TGD[25] as well as *Lepidodinium chlorophorum*[26] (Supplementary Fig. 9). This group might represent a larger diversity of endosymbionts than previously recognized. Taken together, these examples demonstrate the usefulness of metagenomics to recover a broad diversity of plastid genomes among multiple plastid major groups, in particular filling critical evolutionary gaps with environmental genomic diversity for clades lacking reference genomes.

Leptophytes: a new deep branching plastid group

Aside from the extended genomic diversity within clearly defined plastid groups, we also recovered a

fully supported deep-branching clade formed by four red algal-derived ptMAGs that is related to, but

distinct from all major groups with complex red plastids, *i.e.,* haptophytes, cryptophytes, ochrophytes,

and myzozoans (Fig. 1). To facilitate discussion, we propose the informal name leptophytes for this

new group of plastid genomes. The name refers to the putative small size (*lepto-* indicates something

small) of the corresponding algal cells, which we infer to be below 5µm based on genomic size

fraction distributions across the *Tara* Oceans metagenomes (Supplementary Fig. 10). Leptophytes

were collectively detected in the surface waters of most oceanic basins (Supplementary Fig. 11). At

the level of individual genomes, Lepto-01 and Lepto-02 were only detected in the Arctic Ocean

characterized by cooler, less saline, and more nutrient-rich waters, while Lepto-03 and Lepto-04

showed a broader and non-polar distribution associated with warmer, more saline, and nutrient-poor

waters (Supplementary Figs. 12-13). Although leptophyte genomes were usually found in relatively

low abundance (representing 0.14% of the total plastid signal, and detected in 104 out of 147 *Tara*

stations with a median cumulative coverage of 2.3×), Lepto-01 stood out as one of the most detected

plastid genomes in two Arctic stations (up to 124x mean coverage in the 0.8-2000µm size fraction),

providing evidence that leptophytes can experience localised increases in abundance under certain

conditions (Supplementary Table 2). Interestingly, two 16S rDNA gene fragments were recovered in

Lepto-01 and Lepto-04, respectively, which revealed that leptophytes are most similar to a few

amplicon sequences of the enigmatic DPL2 plastid group[13]. The 16S rDNA of Lepto-04 was 100%

similar to DPL2, while Lepto-01 shared 91% similarity to DPL2. The relatedness of DPL2 to

leptophytes was confirmed by 16S rDNA phylogeny, which also showed a weak affinity of the group

to haptophytes (Supplementary Fig. 14). The reported sub-tropical distribution and pico-size of

DPL2[13] largely overlap with Lepto-03 and Lepto-04 but not with the Arctic preference of Lepto-01

and Lepto-02, thus altogether indicating that leptophytes form a more diverse and widespread group

than the narrower DPL2 lineage.

Lepto-01 is the best representative genome for the group. This assembly had the highest coverage

(detected in 20 stations with an average mean coverage of 6.3×) but it was also near complete and

contiguous with a single contig of 104,203 bp. This ptMAG contained 144 genes with 118 proteins

(no detectable introns) and 24 tRNAs (Supplementary Fig. 15). Interestingly, leptophyte plastid

genomes seem to lack inverted repeats commonly found in plastid genomes[27], although whether they

instead contain direct repeats such as in some haptophytes[28], or lack repeat regions altogether remains

unresolved (Supplementary Fig. 15). The functional repertoire of Lepto-01 shows that it is a

photosynthetic plastid genome, retaining genes for the core components of both photosystem I and II

(*psa-* and *psb-* genes), carbon fixation (*rbcL* and *rbcS*), cytochrome b6/f complex (*pet-* genes), ATP

synthase (*atp-* genes), and chlorophyll biosynthesis (*chlI*) (Fig. 2, Supplementary Fig. 15). While less

complete and more fragmented, the three other leptophyte ptMAGs are largely syntenic with Lepto-01

and support these functional trends (Supplementary Fig. 16). The near complete assembly of Lepto-01

allowed us to compare its gene content with that of red algae and red algal-derived plastids. UPGMA-

clustering of homologous gene occurrence showed that the gene content of leptophytes is most similar

to that of haptophytes (Fig. 2). In particular, leptophytes are lacking 10 genes (including *dnaB*, *ftsH*,

petF, *psbW*, *rpl4*, *rpl18*, *rpl29*, *rpl35*, *sufC*, *ycf33*) that are among the 18 genes almost completely

absent in haptophytes, but mostly present in all other algal groups with plastids of red-algal origin

(Fig. 2, Supplementary Fig. 17).

Leptophyte plastids have phylogenetic affinities with haptophytes and cryptophytes

To gain further insight into the position of leptophytes in the plastid tree, we reconstructed and

evaluated phylogenies using several complex site-heterogeneous mixture models in Bayesian and ML

implementations. These models, which consider site-specific amino-acid frequencies, have been

shown to generally improve tree estimations but they come at heavier computational costs[29]. We thus

produced a reduced dataset based on the same 93 genes as in the full phylogeny but containing 107

plastid genomes to speed up the analyses while maintaining the phylogenetic breadth for red algae and

red algal plastid-containing lineages. Due to overall higher completeness, reference genomes were

preferred over ptMAGs when possible, and the least complete leptophyte ptMAG (Lepto-02) was

excluded. The battery of advanced phylogenetic analyses all unambiguously confirmed that

leptophytes form a novel group of plastid genomes related to haptophytes and cryptophytes (Fig. 3a).

This affinity is also supported by a shared rare bacterial gene replacement by horizontal transfer of the

ribosomal protein gene *rpl36* into the plastid genomes of all three groups (Fig. 3b). We will refer to

this group as the Cryptophytes-Haptophytes-Leptophytes (CHL) group.

Leptophytes were recovered as sister to both haptophytes and cryptophytes (HC-sister topology) with

the Bayesian model CAT+GTR+G4 (PP=0.99), as well as the ML models GTR+CAT-PMSF+G4

(BS=88) and LG+MEOW80+G4 (UFB=61) in which site profiles are estimated directly from the

data[30,31]. The same position for leptophytes was recovered with the empirical mixture model

cpREV+C60+G4 model and in an analysis that removed the 10 fastest-evolving taxa (Supplementary

Fig. 18-19). Because of the relatively modest bootstrap support for the position of leptophytes in ML

inferences, we compared the likelihood of the three possible topologies within the CHL group,

enforced as topological constraints but letting other branches and parameters to be optimised by ML.

Under the GTR+CAT-PMSF model, both alternative topologies were rejected by a likelihood chi-

squared test with a Bonferroni correction for multiple comparisons[32], and the sister position to

cryptophytes was additionally rejected by an AU test (Fig 3c). The likelihood values under the

LG+MEOW80+G4 model had less than four points difference across the three topologies, and none

were statistically rejected (Fig. 3c).

We then evaluated whether the position of leptophytes was influenced by two common types of

systematic errors in phylogenetic reconstruction: heterotachy of evolutionary rates, and compositional

heterogeneity across tree branches. To do this, we calculated the likelihoods of the three alternate

phylogenies as fixed topologies under newly developed complex models as additional parameters to

the LG+MEOW80+G4 model. Heterotachy was modeled with GHOST[33], while compositional

heterogeneity was accounted for with the GFmix model[31,32]. The GHOST model continued to favour

the HC-sister topology, but the best topology under the GFmix model changed to support the

monophyly of leptophytes and haptophytes (H-sister topology) (Fig 3a and c). With both models, the

alternative tested topologies were rejected by the Bonferroni-corrected likelihood chi-squared test (Fig

3c). Given the observation that compositional bias may have impacted our phylogenetic

reconstructions, we attempted to minimize this artefact by recoding the 20 amino acids to a reduced 4-

class alphabet (SR4 recoding)[34], and by removing the most heterogeneous sites (~25% of positions in

the alignment) with a stationary-based trimmer[35]. In agreement with the GFmix model, the recoded

dataset recovered the H-sister topology, albeit with low support (PP=0.75; Supplementary Fig 20).

However, the stationary-trimmed alignment recovered the HC-sister topology, also strong support

under the CAT+GTR+G model (PP=0.99, but low support with the LG+MEOW80+G model

(UFB=72) (Supplementary Fig. 21). Overall, these results indicate that resolving the topology of the

CHL group in general and placement of leptophytes in particular is not trivial, as both HC-sister and

H-sister topologies remain credible (Supplementary Note 1).

A mitochondrial genome places leptophytes as sister to haptophytes

Plastid genomes do not necessarily mirror the long-term evolutionary trajectory of their host cell

because of the complex history of endosymbioses[2,4]. In an effort to identify potential nuclear-encoded

sequences for leptophytes, we searched the Arctic stations data—where the leptophyte abundance was

the highest—for *psbO* and 18S rDNA candidate genes (see Methods). The *psbO* gene is essential for

photosynthesis, it is always host-encoded and never found in non-photosynthetic organisms[36].

Although it is derived from the endosymbiont and thus would not be a good phylogenetic marker for

the host, our rationale was that it could pinpoint to host contigs that may contain additional genes.

These analyses did not provide promising candidates for either gene.

We then turned to mitochondria, as mitochondrial genomes have the double advantage of typically

being multi-copy and in theory mirroring the evolution of the host as this organelle traces back to the

origin of eukaryotes[31,37]. From the six Arctic samples where the Lepto-01 ptMAG was the most

abundant, we performed a targeted metagenomic co-assembly (total of 2.54 billion metagenomic

reads providing a cumulative coverage of 299× for the Lepto-01 ptMAG) that produced 56,103

contigs larger than five kb. Functional annotations based on reference mitochondrial genomes resulted

in the identification of 34 mitochondrial contigs (mtMAGs, Supplementary Table 5).

To find a possible correspondence with the Lepto-01 ptMAG, mtMAGs needed to fulfil two

complementary requirements. Most importantly, their read coverage should show strong correlation

with the Lepto-01 ptMAG across the *Tara* Oceans metagenomes. Secondly, their phylogenetic

position among reference mitochondrial genomes should not be placed within known major algal

groups. Among the 34 mtMAGs, one genome (Lepto-01_mtMAG_004) showed highly significant

read coverage correlation with the Lepto-01 ptMAG (R^2=0.96, p-value < 0.01) (Fig. 4, Supplementary

Fig. 22). Most interestingly, Lepto-01_mtMAG_004 also corresponded to a novel deep-branching

lineage firmly placed as sister to haptophyte mitochondrial genomes (Fig. 4; UFB=100%, PP= 1.0).

This mtMAG is 37,793 bp long and contains 37 protein coding genes as well as 22 tRNA genes

(Supplementary Fig. 23), with the presence of various marker genes indicating that it is nearly

complete. Based on both criteria of abundance and phylogeny, we infer that Lepto-01_mtMAG_004

and the Lepto-01 ptMAG are from the same novel host lineage related to haptophytes.

Discussion

The vast majority of organellar genomic knowledge has been gained from cultured organisms. We

demonstrate that plastid and mitochondrial genomes can readily be recovered from large metagenomic

assemblies using phylogeny-guided genome-resolved metagenomics. This study represents the first

global-scale survey of environmental plastid genomes (ptMAGs), providing a culture-independent

view of the diversity, functioning and evolutionary history of plastid genomes occurring in marine

surface water. ptMAGs better represent the diversity and abundance of planktonic algae than

reference genomes, spanning most major marine phytoplanktonic groups and often filling in

evolutionary gaps in areas of the tree poorly represented by references.

One very interesting new group of plastid genomes are the leptophytes, a globally distributed and

generally rare deep-branching lineage formed by four ptMAGs in our data. This is remarkable

because novel plastid diversity at this taxonomic depth is very rarely reported. Currently, only the

environmental plastid lineages DPL1 and especially DPL2 are still considered possible deep-

branching phytoplankton lineages outside of eukaryotic supergroups[13]. Based on 16S rDNA data

recovered from the ptMAGs and biogeographical comparison, we show not only that one of the

leptophytes (Lepto-04) corresponds to DPL2, thus representing the first genomic data available for

this enigmatic group, but also that leptophytes form a more diverse, widespread, and abundant group

than previously known. Part of our findings have now been independently recapitulated, with the

notable recovery of the Lepto-01 ptMAG from a different metagenomic dataset[38].

The plastids of leptophytes are evolutionarily related to cryptophytes and haptophytes, indicating that

they possess red-algal derived plastids. This is firmly supported by phylogenetic analyses as well as

gene content comparison and the exclusive presence of the c-type paralog of the ribosomal protein

gene *rpl36* in cryptophytes, haptophytes and leptophytes (the CHL group). The exact phylogenetic

position of leptophyte plastids within the CHL group remains ambiguous in spite of the use of a dense

taxon-sampling, 93 genes, and a range of sophisticated evolutionary models and data treatments.

Taking into account this phylogenetic uncertainty, we propose two competing hypotheses for the

evolutionary origin of leptophyte plastids: 1) the HC-sister topology (leptophytes are sister to both

cryptophytes and haptophytes), which was favoured by all site-heterogeneous models, or 2) the H-

sister topology (leptophytes are sister to haptophytes, to the exclusion of cryptophytes), which was

favoured only after specifically taking into consideration amino acid compositional heterogeneity

across taxa, and also by the 16S rDNA phylogeny.

In contrast to the robust support for most bipartitions in the plastid tree (Fig 3a), the ambiguity in the

placement of leptophytes likely resulted from weak stochastic phylogenetic signals due to the ancient

origin of the plastids and closely spaced speciation events early in the CHL group. Although a precise

timeframe for the origin of the CHL plastids is unavailable, it has been suggested that both

cryptophytes and haptophytes acquired their plastids more than 1 billion years ago[39]. Based on the

phylogenetic position of leptophytes, it is reasonable to assume that this timeframe represents a

minimum age estimate for the divergence of this group. This ancient age, combined with the

extremely short internal branch uniting haptophytes and cryptophytes (Fig 3) and the limited

phylogenetic signal in plastid genomes, means that it is probably not possible with our current

phylogenetic models and data to better resolve this part of the plastid tree.

Unlike the unsettled phylogenetic position of the leptophyte plastids, we retrieved a near complete

mitochondrial contig that we have assigned to leptophytes based on highly significant abundance

correlation and strongly supported sister relationship to haptophytes. This is particularly interesting

because unlike plastids, mitochondria have been vertically transmitted throughout the evolution of

eukaryotes and thus show the history of the hosts[31]. This phylogenetic position bears important

implications on our understanding of the origin and spread of complex red plastids across the

eukaryotic tree. To explain the patchy distribution of these plastids in the tree, the existing models of

serial endosymbiosis all identify cryptophytes as the unique recipient of secondary red plastids

(plastids derived directly from red algae), mostly because cryptophytes are the only known algae that

have retained a red algal nucleomorph[40,41]. Most models also specifically propose a plastid transfer

between cryptophytes and haptophytes to take into account the rare *rpl36* gene replacement into their

plastid genomes[42] and affinities of both groups in several plastid phylogenies[22,43]. Our new plastid and

mitochondrial phylogenies are consistent with available data, but adding leptophytes allows us to

make new predictions on the relative timing of events and possible direction of plastid transfer.

Reconciling our two hypotheses for the placement of leptophytes in the plastid phylogeny with the

best current estimate of eukaryotic phylogeny[45], we propose two models for the evolution of complex

red plastids, focusing here on the CHL group for clarity (Fig 5). Both models assume a sister

relationship of leptophytes to haptophytes in the eukaryote tree based on our mitochondrial

phylogeny. Whether this means that haptophytes should be expanded to include leptophytes as a new

deeply diverging class, or that leptophytes is a distinct but related lineage, will require detailed

morphological information (such as the presence of a haptonema) not available at the moment. The

first model presumes that leptophytes are sister to haptophytes (H-sister) in the plastid phylogeny as

well (Fig 5a). Leptophytes would have inherited their plastids, including the *rpl36*-c paralog, in a

common ancestor with haptophytes by tertiary endosymbiosis from cryptophytes and would lack a

nucleomorph. The second model reconciles the HC-sister plastid topology with the position of

leptophytes as sister to haptophytes in the host phylogeny (Fig 5b). In this scenario, the secondary

plastid from red algae was established in the common ancestor of leptophytes and haptophytes, where

the ancestral *rpl36*-p gene was replaced, before passing that plastid further to cryptophytes from a

stem haptophyte. To our knowledge, this model is the first to propose the haptophyte ancestor as the

host of the secondary endosymbiosis with red algae, and is in line with the general older origin of

haptophytes over cryptophytes inferred by molecular clock analysis[39]. The model places the loss of

nucleomorph in haptophytes subsequent to the tertiary transfer to cryptophytes, but leaves open the

possibility that leptophytes have also retained this organelle. To weigh in on both models, we

searched for evidence of red algal nucleomorph in the same metagenomic data used to reconstruct the

ptMAGs. While we found ample evidence of cryptophyte nucleomorphs, we found no evidence of

putative leptophyte nucleomorphs (see Methods). Without a more conclusive plastid phylogeny, or

other supporting evidence, it is currently not possible to decide which model, if any, is correct.

However, we note that the H-sister model is more parsimonious as it requires fewer parallel gene

losses based on our plastid gene content comparison (5 vs 33; Supplementary Fig. 24), and it is also

more consistent with phylogenies of the SELMA proteins involved in protein translocation from the

cytoplasm to the plastid[46]. More generally, our advanced phylogenetic analyses and broad taxon

sampling continue to recover—albeit with only moderate support—the monophyly of all red algal-

derived plastids, unlike in a recent proposal that argues for a separate secondary plastid acquisition in

ochrophytes[47]. The origin of the ochrophyte plastid is a pressing issue to address in the future.

Ultimately, deciding between the origin of the leptophyte plastid and order of transfer to or from
cryptophytes and haptophytes will rest on characterizing the nuclear and putative nucleomorph
genomes of leptophytes, ideally in the context of cultivation, in order to determine their specific
position in the eukaryotic tree and investigate their cell biology. This will be challenging, as our data
suggest that leptophytes are generally rare and it remains unclear if they occur in easily accessible
coastal regions. Despite the fundamental importance of algae with complex red plastid as marine
primary producers, we still understand little of how this great diversity of algae came to be. Our data
show that players still hidden in the rare biosphere hold important clues that could bring new critical
evidence. We demonstrated that organellar genome-resolved metagenomics allows access to this
hidden diversity and should be considered in the future as a valuable approach to survey not only
plastids but also mitochondria across biomes.
382
383   **Material and Methods**
384
385   *Tara* **Oceans metagenomes.** The 937 metagenomes from *Tara* Oceans used in the study are publicly
386   available at the EBI under project PRJEB402 (https://www.ebi.ac.uk/ena/browser/view/PRJEB402).
387
Phylogeny-guided genome-resolved metagenomics for plastids. We used two complementary
datasets (metagenomic assemblies and evolutionary-informative proteins) to characterize ptMAGs.
On the one side, we used 11 large metagenomic co-assemblies from *Tara* Oceans (~12 million contigs
from 798 eukaryote-enriched metagenomes) which were organized into 2,550 metabins using
constrained automatic binning and processed with anvi'o[20,21] v7 for manual binning purposes[16]. On
the other side, we used proteins corresponding to the DNA-dependent RNA polymerase B subunit
(RNApolB) and found in those *Tara* Oceans metabins, in the context of a phylogeny of representative
sequences (amino acid level with sequence similarity <90%)[48]. Until now, the two datasets had only
been used to characterize genomes corresponding to RNApolB clades of giant viruses and
mirusviruses. Here, we focused on a large plastid RNApolB clade most closely related to that of
Cyanobacteria and used this signal as guidance for genome-resolved metagenomics using the anvi'o
interactive interface. Briefly, we mainly used sequence composition and differential coverage across
metagenomes of the corresponding co-assembly to characterize and manually curate 1,448 ptMAGs in
the corresponding metabins. The strong correlation between initial signal (number of plastid
RNApolB genes) and outcome (number of characterized ptMAGs) across metabins (Supplementary
Fig. 1) demonstrated the effectiveness of this method to survey the environmental genomics of
plastids in complex marine metagenomes. Preliminary phylogenies (see sections below) with 32 genes
indicated a deep-branching lineage (the leptophytes) composed of three ptMAGs.
406

Recovery of additional ptMAGs from the mOTU database. To broaden the scope of our survey,
we exploited 85,123 metagenomic assemblies from the mOTU global metagenomic resource covering
a wide range of environmental samples[18]. A total of 1,969,342 RNApolA genes were previously
characterized from this resource[49]. Note that RNApolA displays a very similar evolutionary signal
compared to RNApolB[50]. Here, we identified 6,954 contigs >50 kbp that contained a RNApolA gene
with relatively high sequence similarity to that of our characterized ptMAGs (DIAMOND blast, with
percent identify >70% at the amino acid level). Based on preliminary phylogenies, we found one
contig characterized from a *Tara* Oceans metagenome and corresponding to a leptophyte (contig ID
Lepto-01_REFM_CHLORO_00001 with a length of 104,203 nt). In addition, we also found one
contig (also characterized from a *Tara* Oceans metagenome) and corresponding to a deep-branching
clade of haptophytes (contig ID REFM_CHLORO_00002 with a length of 84,869 nt). We integrated
these two contigs into our database of ptMAGs.

Creation of a non-redundant plastid genomic database. We collected all the manually curated
Tara Oceans ptMAGs as well as the two ptMAGs from mOTUS-db and included 166 manually
selected reference plastid genomes from cultured organisms. We determined the average nucleotide
identity (ANI) of each pair of genomes using skani[51] v0.2.1. Genomes were considered redundant
when their ANI was >98% (minimum alignment of >25% of the smaller genome in each comparison).
For genomes found to be redundant, reference chloroplast genomes were selected over ptMAGs, and
in most cases where only MAGs were redundant, the longest one was selected. We estimated the
completeness and redundancy of each genome based on the occurrence of 44 core-plastid genes[52]
(Supplementary Table 6), and removed ptMAGs with a redundancy level higher than 15%. This
analysis provided a non-redundant database containing the 166 reference plastid genomes and 660
ptMAGs.

To assess potential mitochondrial contamination, all ptMAGs were screened for the presence of 25
canonical mitochondrial genes (Supplementary Table 7), as listed in [53]. Five non-redundant ptMAGs
were found to contain genes annotated as typical mitochondrial genes, which were confirmed by
BLAST searches. Further manual inspection indicated that these instances of contamination were not
due to chimeric misassemblies. Instead, they resulted from inadvertent co-binning of mitochondrial
and plastid contigs. The contaminated contigs were removed from their respective ptMAGs.

Biogeography of the GOEV database. We performed a mapping of 937 metagenomes from *Tara*
Oceans to calculate the mean coverage (vertical coverage) and detection (horizontal coverage) of each
genome in the non-redundant plastid genomic database. Briefly, we used BWA v0.7.15 (minimum
identity of 95%) and a FASTA file containing all contigs from the database to recruit short reads from
each metagenome[54]. We considered a genome to be detected in a metagenome when >25% of its

length was covered by reads. The number of recruited reads below this cut-off was set to 0 before

determining the mean coverage of genomes, as an effort to minimize non-specific read recruitments.

Genome annotation of ptMAGs. Plastid annotation was done with MFannot[55] v1.3.6 using the

genetic code 11. The resulting ASN format files were converted to GenBank format using the *asn2gb*

utility (https://ftp.ncbi.nlm.nih.gov/asn1-converters/by_program/asn2gb/) from the NCBI toolkit. We

then extracted all annotated proteins from the GenBank files using the script *gb_to_prot.py* (available

at https://github.com/burki-lab/beta-Cyclocitral/blob/master/src/gb_to_prot.py). For the leptophyte

ptMAGs, intron prediction was carried out using RNAweasel

(https://megasun.bch.umontreal.ca/apps/rnaweasel/), the 5S rRNA gene was detected with Infernal

Cmscan (https://www.ebi.ac.uk/jdispatcher/rna/infernal_cmscan), and maps were generated using

OGDraw version 1.3.1 (https://chlorobox.mpimp-golm.mpg.de/OGDraw.html)[56]. Downstream

analyses with OrthoFinder (see **Comparative analysis of plastid gene content in leptophytes and**

related lineages) revealed that MFannot did not detect the *rpl22* gene, labelling it as a "hypothetical

protein" instead. The annotations of the leptophyte ptMAGs were manually updated to include this

gene.

Taxonomic annotation of ptMAGs. We manually determined the taxonomy of ptMAGs based on

the global phylogenetic analysis (see section **Phylogenomic analyses of plastid dataset**), using

guidance from the reference plastid genomes.

Phylogenomic plastid dataset assembly

We began by using the two plastid gene sets originally defined by Janouškovec et al 2010[57]: a larger

set of 68 plastid genes, and a more conserved subset of 34 genes. These served as the foundation for

constructing preliminary phylogenies. During dataset assembly (see below), two genes were excluded:

acsF due to low taxon occupancy, and *psbH* due to anomalously long branches in its single-gene tree.

This reduced the gene sets to 32 and 66 genes, respectively. The resulting preliminary phylogenies

identified leptophytes as a novel clade, but could not confidently resolve their position using

maximum-likelihood analyses (Supplementary Figs. 25-26). Therefore, we attempted to increase the

phylogenetic signal, by including additional genes used in Ponce-Toledo et al 2017[58] and Pietluch et

al 2024[47], provided they were also present in leptophytes, resulting in a final set of 93 genes

(Supplementary Table 8).

We first built a dataset of 93 protein-coding genes from the 179 reference taxa (166 plastid genomes

and 13 cyanobacterial genomes). This was done as follows: (1) Homologous sequences from all

reference taxa were retrieved with BLASTP[59] v2.15.0 searches using sequences from *Fucus*

vesiculosus (which has a well-annotated plastid genome; NC_016735) as queries (e-value: 1e-02), (2)

For each gene, sequences were aligned with MAFFT[60] v.7.407 using the -auto option, and then

trimmed with trimAL[61] v1.4.1 using a gap threshold of 0.8, (3) Single-gene trees were inferred with

raxml-ng[62] v.1.2.0 using the LG4X model and 100 rapid bootstrap searches, (4) Gene trees were

manually parsed to identify duplicates and extremely long branches corresponding to sequences from

another gene, which were then removed from the dataset.

We then proceeded to construct a dataset initially including the 1448 ptMAGs identified (including

redundant ptMAGs). Using reference sequences from the previous step as queries, we retrieved

homologous sequences from the ptMAGs using BLASTP (e-value: 1e-02) and the gene annotation

from MFannot. The 93 single-gene dataset was aligned individually using MAFFT-G-INS-I (using the

--unalign 0.6 option to avoid over-alignment), and gently trimmed with trimAL (gap threshold: 0.9).

Single-gene trees were inferred with IQ-TREE[63] v2.2.2.6 using the best fitting model as determined

by ModelFinder Plus[64]. We manually parsed the single gene trees to check for putative chimeras and

generated our final dataset comprising 93 genes and 839 taxa (660 ptMAGs and 179 references) with

> 4% data.

The concatenated alignment was generated as follows: for each curated gene, we first used Prequal[65]

v1.02 to filter sequence stretches with no clear homology using a posterior probability threshold of

0.95, then aligned with MAFFT-G-INS-I (--unalign 0.6), and subsequently trimmed with BMGE

v1.12[35] (BLOSUM35 matrix, gap threshold: 0.8). After concatenating, we obtained a supermatrix

(839-taxon dataset) with 18,759 aligned amino acid sites, which was used for broad-scale

phylogenetic analyses. Finally, we also generated a smaller, 107-taxon dataset to enable analyses

under more complex and computationally demanding models. For this dataset, raw sequences for the

93 genes were again filtered, aligned, trimmed and concatenated as previously described, yielding a

supermatrix with 20,292 amino acid sites.

Phylogenomic analyses of plastid datasets

Site-homogeneous models

To assess the phylogenetic diversity of the ptMAGs, the full 839-taxon supermatrix was used to

compute a maximum-likelihood tree using the best-fitting site-homogeneous model, LG+F+I+G4 in

IQ-TREE v.2.2.2.6. Branch support was assessed with 1000 ultrafast bootstrap replicates.

Site-heterogeneous models

We used the smaller, 107-taxon dataset to perform analyses with the more complex, site-

heterogeneous models in order to determine the position of leptophytes in the plastid tree. An initial

tree was inferred using the best-fitting cpREV+C60+G4 model in IQ-TREE with 1000 ultrafast

bootstrap replicates. This phylogeny was used as a guide tree to compute custom site-profiles directly

from the input alignment using the programme MEOW[31] (-ri -p R -C 5 -f H -l; custom version of

MEOW used is provided at https://github.com/burki-lab/beta-

Cyclocitral/blob/master/src/meow_custom.R), an approach which often outperforms generic mixture

models (such as the C-series)[66]. MEOW (MAMMaL Extension On Whole alignments) extends the

MAMMaL approach by using all variable sites in the alignment to estimate custom site-profile

classes. Following Williamson et al[31], we estimated three sets of profile classes using varying

proportions of high- and low rate classes respectively: (1) MEOW(80,0) indicates 80 classes from

high rate sites and none from low rate sites, (2) MEOW(60,20) represents 60 classes from high rate

sites and 20 classes from low rate sites, and (3) MEOW(40,40) corresponds to 40 classes each from

the two partitions. We inferred a maximum likelihood phylogeny under the best fitting model based

on the BIC criterion, LG+MEOW(60,20)+G4, with 1000 UFBOOT replicates, and for brevity, we

refer to this model as LG+MEOW80+G throughout the manuscript.

Another Maximum-likelihood phylogeny was inferred with the 107-taxon dataset with the CAT-

PMSF approach[30]. Briefly, a phylogeny inferred under the site-homogeneous LG+G model in IQ-

TREE was used as a fixed tree for analysis with the CAT+GTR+G model in PhyloBayes-MPI[67] v1.8.

Two Markov Chain Monte Carlo (MCMC) chains were run for more than 5,000 cycles until the

effective sample sizes of nearly all parameters (seven out of eight) were above 100 (Supplementary

Table 9). The posterior mean exchangeabilities and the custom site-frequency profiles were extracted

for analysis in IQ-TREE and support was assessed with 100 non-parametric bootstrap replicates.

Finally, we also analysed the 107-taxon dataset in PhyloBayes under the CAT+GTR+G model via

three independent MCMC chains. The chains were run for almost 9,000 cycles each. After discarding

the first 2,000 cycles as burn-in, convergence was assessed (with maxdiff reaching 0.047) and

consensus trees were generated using the bpcomp command.

Derivatives of 107-taxon dataset

First, we tested the impact of fast-evolving taxa in our dataset by removing 10 taxa (all ochrophytes)

with the largest root to tip distances, generating a 97-taxon dataset. Second, we dealt with possible

compositional heterogeneity across taxa by using BMGE's stationary-based trimmer to remove sites

based on Stuart's test of marginal homogeneity (BMGE -s FAST -h 0:1 -g 1; 4752 out of 20292 sites

= 23.4% sites removed). Finally, we dealt with possible compositional heterogeneity and sequence

saturation by recoding the dataset into SR4 categories[34]. Both the non-recoded datasets were analysed

in IQ-TREE with the LG+MEOW80+G model with 1,000 ultrafast bootstraps. Additionally, the

stationary-trimmed alignment was analysed in PhyloBayes using the CAT+GTR+G model, with three

MCMC chains runs for 7,000 cycles after a burn-in of 5,000 cycles, achieving a maxdiff of 0.104. The

SR4 recoded dataset was also analysed under the CAT+GTR+G model in PhyloBayes, using three

MCMC chains run for 22,000 cycles, with a burn-in of 4,000, reaching a maxdiff of 0.101.

Assessing alternative topologies

Site-heterogeneous models

To assess alternate positions of leptophytes, we performed constrained tree searches in IQ-TREE. We

considered three possible positions in the algal tree: (1) sister to haptophytes and cryptophytes (HC-

sister), (2) sister to haptophytes (H-sister), and (3) sister to cryptophytes (C-sister). For each of these

three positions, we considered two topologies; one in which the complex plastids are monophyletic,

and the other where the complex plastids are non-monophyletic (as recently recovered by [47]), bringing

the total number of tested topologies to six. Constrained topologies were inferred with IQ-TREE

under the GTR+CAT-PMSF+G and LG+MEOW80+G models for the 107-taxon dataset, and the

xmC60SR4 model for the SR4 recoded dataset, with all parameters optimized freely. In the case of

the LG+MEOW80+G model, ten independent runs in IQ-TREE were conducted for each constraint to

avoid likelihood scores from searches stuck in local optima[68]. In all cases, the topologies were

evaluated with the Approximately Unbiased (AU)[69] test implemented in IQ-TREE. Additionally, the

best scoring topology under each model was compared with alternative topologies using the

Bonferroni-corrected chi-squared test[31,32]. Briefly, the likelihood ratio test statistic (LRS; two times

the difference between the maximised log-likelihood of the two topologies under consideration) is

first calculated. The p-value for the chi-squared test ($\square$) is calculated as the probability that a chi-

squared random variable would be larger than the LRS with the degrees of freedom defined as the

number of branches collapsed in the strict consensus tree of the two trees. The Bonferroni-corrected p-

value is calculated as $(1 - \{1 - p\}^A)$ where A is the set of trees compatible with the strict consensus

tree.

GHOST and GF-mix models

We additionally tested the six alternative topologies under GHOST[33] and GFmix[31,32] as additional

parameters to the LG+MEOW80 base model. For GHOST, the log-likelihood scores of the six

topologies (inferred under the LG+MEOW80+G model) were calculated under the

LG+MEOW80+H8 model in IQ-TREE, using the topologies as fixed trees. The LG+MEOW80+H8

indicates a model with eight linked heterotachy classes and was the best fit model using the BIC

criterion when compared to models with four or six heterotachy classes. Topologies were tested using

the AU test and the Bonferroni-corrected chi-squared test as described previously.

The GFmix model models compositional heterogeneity by modifying the vector of amino acid

frequencies in a branch-specific manner to account for shifts in the relative frequencies of amino acids

in different branches of the tree. While prior studies using this model have divided their alignment

into multiple partitions to account for varying compositional heterogeneity in mitochondrial and

nuclear encoded genes (e.g., [31,32]), we opted to use a single partition. This is because we do not expect

substantial discrepancy in compositional heterogeneity between the only plastid-encoded genes

considered in this study. With no *a priori* knowledge of how taxa might be compositionally biased,

we first estimated the groups of amino acids that are depleted and enriched in different lineages from

our data as done in [31]. Briefly, we used a custom R script (C. McCarthy, Dalhousie University,

Canada, available at https://github.com/burki-lab/beta-Cyclocitral/blob/master/src/quickBinomial.R)

to carry out the following steps: (1) tabulate the counts of all 20 amino acids for each taxon, (2)

perform a chi-squared test on these counts and use the Pearson's residuals to construct a UPGMA tree

of taxa, (3) separate taxa into two groups based on the UPGMA tree, (4) conduct a binomial test for

each amino acid and calculate a score reflecting the differences in counts between the two groups, and

(5) classify amino acids into "enriched," "depleted," or "other" categories based on their scores

(Supplementary Note 1). The enriched and depleted classes of amino acids were used by GFmix v1.2

to compute the log-likelihoods of the six alternative topologies, which were inferred using the

LG+MEOW80+G model and provided as fixed input trees.

16S rDNA phylogeny

Barrnap v0.9 (https://github.com/tseemann/barrnap) was used to detect 859 bp and 808 bp fragments

of the 16S rDNA gene in the Lepto-01 and Lepto-04 ptMAGs respectively. A BLAST search of the

Lepto-01 16S rDNA sequence revealed that it was most similar to the DPL2[13] group with ~91%

sequence similarity (accession numbers EF574856 and KX935025). However, the Lepto-04 16S

rDNA sequence best BLAST hits included DPL2 as well as cyanobacteria. Manually BLASTing

fragments of the Lepto-04 16S rDNA sequence against nt indicated that it was likely chimeric, with

the first ~440 bp corresponding to DPL2 (100% similarity), and the latter half originating from

cyanobacteria. We therefore trimmed the Lepto-04 16S rDNA sequence to only the first 440 bp. To

confirm the association of leptophytes with DPL2, we constructed a 16S rDNA dataset including

Lepto-01, Lepto-04, DPL2, and representatives of each eukaryotic algal clade. The 65-taxon dataset

was aligned with MAFFT-L-INS-I, and trimmed with trimAL (-gt 0.1) to get an alignment with 1496

sites. A Maximum-Likelihood phylogeny was inferred with raxml-ng with 20 ML searches and non-

parametric bootstrapping until convergence using the GTR+G model.

Rpl36 alignment and phylogeny

To investigate whether leptophytes possessed the *rpl36*-p type or *rpl36*-c type gene [42], we extracted

the L36 amino acid sequences from the 107 taxa in the subset phylogeny when available. The

sequences were aligned with MAFFT-L-INS-I, trailing ends trimmed manually, and the alignment

was visualised alongside the 93-gene PhyloBayes phylogeny using the R package ggtree[70] v3.6.2.

Synteny analysis of leptophyte ptMAGs

Synteny analysis of the leptophyte ptMAGs was performed using the GenBank files generated during genome annotation using PyGenomeViz v1.0.0 via the PyGenomeViz Streamlit Web Application (https://pygenomeviz.streamlit.app/). Sequence similarity of protein coding genes was estimated using MMseqs2[71] v15-6f452 for reciprocal best-hit CDS search.

Comparative analysis of plastid gene content in leptophytes and related lineages

To gain insights into the evolutionary dynamics and functional capabilities of the leptophyte plastid, we compared its gene content with that of red algae and other red algal-derived plastids. As ptMAGs can have missing genes due to incompleteness, we opted to use only reference plastid genomes where possible. For this analysis, we selected 15 rhodophyte, seven haptophyte, eight cryptophyte and 21 ochrophyte reference plastomes from our plastid genomic database. Similarly, we used the Lepto-01 ptMAG, the only near-complete leptophyte ptMAG, as the sole representative of leptophytes. We extracted amino-acid sequences from all selected plastid genomes and used OrthoFinder[72] v2.5.5 with default settings to detect homologous proteins. This analysis yielded 281 phylogenetic hierarchical orthogroups (HOGs) containing 7,648 protein-coding genes (98.5% of the total). However, as no orthology-predictor is perfect, we manually refined the HOGs by inspecting gene annotations, single-gene trees, and performing BLAST and InterProScan searches (Supplementary Table 10). This step yielded 237 HOGs containing sequences from at least three taxa. Heatmaps depicting gene presence/absence were plotted using the R package pheatmap v1.0.12 (https://github.com/raivokolde/pheatmap), and species and orthologs clustered using UPGMA clustering.

Leptophyte associations with environmental parameters

We assessed correlations between leptophyte abundance and eight physiochemical parameters: sea surface temperature, salinity, dissolved silica, nitrate, phosphate, iron, and seasonality indices of nitrate and sea surface temperature. These parameters were obtained from Delmont et al 2022 [16], representing data pulled from climatology and biogeochemical modelling data (World Ocean Atlas 2013 and PISCES v2)[73,74] to account for missing physio-chemical samples in the *Tara* Oceans *in-situ* dataset. Seasonality indices were defined as the range of the nitrate and temperature in one grid cell divided by the total range of that variable across all *Tara* sampling stations.

We calculated Pearson's correlation coefficients between the abundance of each leptophyte ptMAG and the environmental parameters using the R package, corrgram v1.14.

Survey for putative nucleomorphs of leptophytes

664   We searched for genomic signal corresponding to red-algal nucleomorphs associated with leptophytes
665   across the 11 Tara Oceans metagenomic co-assemblies. Specifically, we surveyed the assembled
666   RNApolB genes, following the protocol used to characterize ptMAGs but this time using publicly
667   available cryptophyte nucleomorph RNApolB genes as a reference. With this approach, we did not
668   identify any nucleomorph-affiliated RNApolB genes co-occurring with the ptMAGs of leptophytes.
669

Survey for host nuclear sequences of leptophytes
Survey for leptophyte psbO sequences
We considered the *psbO* gene as a good candidate to attempt finding contigs corresponding to the
nuclear genome of leptophytes, as it is encoded by all photosynthetic eukaryotes, usually, in a single
copy, in their nuclear genome, and is absent from non-photosynthetic eukaryotes[36]. We identified
psbO sequences in the Arctic metagenomic co-assembly (available at:
https://www.genoscope.cns.fr/tara/; where Lepto-01 is abundant) through an HMM search based on
the PFAM accession PF01716, carried out using HMMER v3.4 as implemented in anvi'o v8. A total
of 51 hits were detected which were added to 346 reference *psbO* sequences from different algal
groups. Sequences were aligned (MAFFT-L-INS-I), trimmed (trimAL, -gt 0.1), and then used to infer
a tree with raxml-ng (20 searches, non-parametric bootstrapping until convergence) to determine the
phylogenetic affiliation of the 51 *psbO* sequences. However, all *psbO* candidates fell within
established algal groups, and no putative *psbO* sequence corresponding to leptophytes could be
identified. We hypothesized that this result could be due to low coverage of the leptophyte nuclear
genome. We investigated this by focusing on the two filters (both belonging to station 194) where
Lepto-01 was among the most abundant eukaryotic plastids (124× and 70× coverage; Supplementary
Table 1). We extracted the top 50 most abundant *psbO* sequences in these two filters from the recently
assembled *psbO* database[36]. We generated an alignment and inferred a phylogeny as before, but again,
did not identify any strong candidates for leptophytes. Most *psbO* sequences had only a small number
of metagenomic reads mapping to them per sample (median number of mapped reads = 3), except for
those from highly abundant chlorophytes and ochrophytes. This suggests that nuclear genomes
generally have substantially lower coverage than the plastid genomes, making it challenging to
retrieve nuclear genomic sequences from eukaryotes other than the most abundant ones.
693

Survey for leptophyte 18S rDNA sequences
We reasoned that a putative leptophyte 18S rDNA sequence would branch outside known algal
lineages, and would correlate in abundance with the ptMAG abundance. We started with a targeted
approach by generating a co-assembly of the six samples where Lepto-01 was most abundant using
Megahit[75] v1.2.9. We then extracted 18S sequences from the co-assembly using Barrnap (length
cutoff: 0.2) with the aim of getting a small, manageable pool of candidate sequences for further
investigation. These 18S sequences were BLASTed (e-value: 1e-10) against PR2[76] v5.0.0 and the nt

database to identify them, however, all sequences were linked to well established clades and no

candidate leptophyte 18S sequence could be identified.

We then turned to metabarcoding data from the top 15 filters where Lepto-01 was most abundant [77].

We attempted to match the 18S-V9 ASVs (n=5,336) to Lepto-01 based on correlation of abundances

following the workflow established by Zavadska et al 2024 [78]. Briefly, we estimated a correlation

coefficient (Spearman's correlation) between ASV relative abundance and the relative, and absolute

abundance of the Lepto-01 ptMAG. We identified 13 and 7 candidate ASVs, respectively, that

showed a Spearman's correlation coefficient greater than 0.7. These ASVs were further analyzed by

performing BLAST searches against the PR2 database and nt database. However, none of the

candidates matched leptophytes and the top hits were to cercozoan sequences in each case. This may

be due to variation in plastid genome copy number under different conditions, leading to an

inconsistent ratio between plastid and nuclear genome abundance.

Survey for leptophyte mitochondrial sequences

Extracting mitochondrial contigs from targeted samples

We first identified core mitochondrial marker genes by functionally annotating a set of seven publicly

available haptophyte and cryptist mitochondrial genomes in anvi'o (*anvi-run-ncbi-cogs*). The following

mitogenomes were used: NC_005332.1 (*Gephyrocapsa huxleyi*), OL703630.1 (*Gephyrocapsa*

oceanica), OL703631.1 (*Gephyrocapsa muellerae*), AB930144.1 (*Chrysochromulina* sp. NIES-1333),

AF288090.1 (*Rhodomonas salina*), NC_010637.1 (*Hemiselmis andersenii*), and NC_031832.1

(*Palpitomonas bilix*). This step revealed 13 core genes present in these mitochondrial genomes that

were functionally annotated as: Cytochrome *c* oxidase subunit III, Cytochrome *c* and quinol oxidase

polypeptide I, Cytochrome *c* oxidase subunit II (periplasmic domain), Cytochrome b/b_6, Proton-

conducting membrane transporter, NADH-ubiquinone/plastoquinone oxidoreductase chain 4L,

NADH-ubiquinone/plastoquinone oxidoreductase chain 6, NADH-ubiquinone/plastoquinone

oxidoreductase chain 3, ATP synthase subunit □, ATP synthase subunit c, Ribosomal protein

L16p/L10e, Ribosomal protein S12/S23, ATP synthase F_0 subunit.

We then turned to the co-assembly of the six filters where Lepto-01 was most abundant and extracted

all mitochondrial contigs that were at least 5 kbp long. To do so, we first searched for contigs

containing the *cox3* gene which encodes the cytochrome c oxidase subunit III enzyme, using an HMM

search (based on the PFAM accession PF00510) in anvi'o v8, and functionally annotated the contigs

as before (*anvi-run-ncbi-cogs*). We then selected contigs containing at least six of the 13 core genes

present in the reference genomes, which yielded a total of 34 contigs representing partial

mitochondrial genomes or mtMAGs (Supplementary Table 5).

Abundance correlation analyses

We expected the putative mtMAG of Lepto-01 to show a correlated abundance pattern with the

Lepto-01 ptMAG. To test this, we mapped all 34 mtMAGs against 937 metagenomes from *Tara*

Oceans, using the same approach applied to the plastid genomes (see **Biogeography of the GOEV**

database. Abundance between each mtMAG and the Lepto-01 ptMAG was compared and plotted in

R using the ggplot2[79] v3.5.1 and ggpmisc[80] 0.6.0 packages. Abundance correlations were tested using

two ways (1) applying a detection threshold where at least 25% of a genome's length had to be

covered by reads for it to be considered present in a metagenome, and (2) without applying this

threshold. Both approaches produced highly consistent results, thus, only results from the analyses

without the coverage threshold are presented here.

Phylogenetic inference

We reasoned that a putative leptophyte mitochondrial genome would be expected to form a distinct

branch from established clades in a mitochondrial phylogeny. To assemble a mitochondrial

phylogenetic dataset, we used the publicly available dataset of Williamson et al 2025 [31] with 93

protein-coding genes and 100 taxa as a starting point. We subset this dataset to retain 50 eukaryotic

taxa and the genes present in their corresponding mitochondrial genomes (40 protein-coding genes).

To this dataset, we added our 34 mtMAGs and 18 additional reference genomes from various sources

to increase the taxon sampling of cryptophytes and haptophytes as well as under-represented lineages

such as Picozoa (Supplementary Table 11). This was done as follows: (1) the tool Codetta[81] v2.0 was

used to determine the genetic code of all mtMAGs and additional reference genomes, (2) mtMAGs

and additional reference genomes were annotated MFannot using the standard or mold mitochondrial

genetic code as appropriate, (3) The resulting ASN files were converted to GenBank format using the

asn2gb tool, and all amino acid sequences were extracted using the custom script *gb_to_prot.py*

(available at https://github.com/burki-lab/beta-Cyclocitral/blob/master/src/gb_to_prot.py), (4)

Homologous sequences from mtMAGs and additional references were retrieved by BLASTP searches

using the 40 genes as queries, (5) Sequences for each gene were aligned using MAFFT-L-INS-I, and

trimmed with trimAL using a gap threshold of 0.8, (6) Single-gene-trees were inferred with IQ-TREE

using the best-fitting model, (7) Gene trees manually parsed to check for problematic sequences. 12

genes were excluded from the dataset at this point due to low taxon occupancy, and the final dataset

comprised 28 protein-coding genes (Supplementary Table 12).

To generate the concatenated alignment, we aligned each gene with MAFFT-G-INS-I (--unalign 0.6),

and gently trimmed alignments with BMGE (BLOSUM30 matrix, gap threshold: 0.8). We

concatenated the genes to obtain an alignment with 102 taxa, 28 genes, and 6,302 sites. The dataset

was used for ML analysis in IQ-TREE using the site-heterogeneous LG+C60+G model with 1,000

ultrafast bootstraps. Additionally, we analysed the dataset in a Bayesian framework with PhyloBayes

under the CAT+GTR+G model. Three independent MCMC chains were run for 15,000 cycles each,

with a burn-in of 4,500 cycles, and achieving a maxdiff value of 0.23.

Data availability. The 937 metagenomes from *Tara* Oceans used in the study are publicly available

at the EBI under project PRJEB402 (https://www.ebi.ac.uk/ena/browser/view/PRJEB402). Data our

study generated has been deposited in an online repository:

https://doi.org/10.17044/scilifelab.28212173. This link provides access to the individual FASTA files

from each plastid and mitochondrial genome used in our study (including the 660 non-redundant

ptMAGs and 34 mtMAGs), the co-assembly of the top six samples where Lepto-01 was most

abundant, individual gene alignments, concatenated and trimmed alignments, and maximum-

likelihood and Bayesian tree files for the phylogenomic dataset.

Code availability. All scripts used for genome annotation and phylogenetic analyses are available on

GitHub: https://github.com/burki-lab/beta-Cyclocitral.

Acknowledgements

We thank Shinichi Sunagawa for having facilitated the recovery of relevant data from the mOTU

metagenomic database maintained by his research group at the Department of Biology at ETH Zürich.

We thank A. Roger, H. Baños, and C. McCarthey for discussions, and for kindly providing custom

scripts for running the phylogenetic models MEOW and GF-MIX. We thank J.E. Dharamshi for

discussions about phylogenetic analyses. Our survey was made possible by the sampling and

sequencing efforts of the *Tara* Oceans Project. *Tara* Oceans (which includes the *Tara* Oceans and

Tara Oceans Polar Circle expeditions) would not exist without the leadership of the *Tara* Oceans

Foundation and the continuous support of 23 institutes (https://oceans.taraexpeditions.org/).

Phylogenetic analyses were enabled by resources provided by the National Academic Infrastructure

for Supercomputing in Sweden (NAISS 2024/5-197), partially funded by the Swedish Research

Council through grant agreement no. 2022-06725. M.J. was supported by the Swedish Research

Council (International Postdoc grant 2022-00351). FB's research is supported by grants from the

European Research Council (ERC consolidator grant 101044505), the Swedish Research Council VR

(2021-04055), and Science for Life Laboratory. TD's research is supported by a grant from the

l'Agence Nationale de la Recherche (ANR-23-CE02-0022).

We also thank the commitment of the CNRS and Genoscope/CEA. Some of the computations were

performed using the platine, titane and curie HPC machine provided through GENCI grants

(t2011076389, t2012076389, t2013036389, t2014036389, t2015036389 and t2016036389).

Author contributions

F.B. and T.O.D. conceived the project. T.O.D. characterized the ptMAGs. M.J., F.B. and T.H.,

performed phylogenetic analyses. T.A., E. P. and T.O.D. created the plastid genomic database and

performed surveys for nucleomorphs. H.J.R. and L.P. retrieved relevant data from mOTUs. M.J. and

T.H. annotated the ptMAGs, and E.P. performed mapping analyses. F.B., M.J. and T.O.D. wrote the

manuscript with input from all the authors.

1.   Andersen, R. A. Diversity of eukaryotic algae. *Biodivers. Conserv.* **1**, 267–292 (1992).

2.   Archibald, J. M. The puzzle of plastid evolution. *Curr. Biol.* **19**, pR81-R88 (2009).

3.   Irisarri, I., Strassert, J. F. H. & Burki, F. Phylogenomic Insights into the Origin of Primary

Plastids. *Syst. Biol.* **71**, 105–120 (2022).

4.   Keeling, P. J. The endosymbiotic origin, diversification and fate of plastids. *Philos. Trans. R. Soc.*

B Biol. Sci. **365**, 729–748 (2010).

5.   Penot, M., Dacks, J. B., Read, B. & Dorrell, R. G. Genomic and meta-genomic insights into the

functions, diversity and global distribution of haptophyte algae. *Appl. Phycol.* **3**, 340–359 (2022).

6.   Malviya, S. *et al.* Insights into global diatom distribution and diversity in the world's ocean. *Proc.*

Natl. Acad. Sci. **113**, E1516–E1525 (2016).

7.   Dorrell, R. G. *et al.* Chimeric origins of ochrophytes and haptophytes revealed through an ancient

plastid proteome. *eLife* **6**, e23717 (2017).

8.   Stiller, J. W. *et al.* The evolution of photosynthesis in chromist algae through serial

endosymbioses. *Nat. Commun.* **5**, 5764 (2014).

9.   Ševčíková, T. *et al.* Updating algal evolutionary relationships through plastid genome

sequencing: did alveolate plastids emerge through endosymbiosis of an ochrophyte? *Sci. Rep.* **5**,

10134 (2015).

10. Keeling, P. J. *et al.* The Marine Microbial Eukaryote Transcriptome Sequencing Project

(MMETSP): Illuminating the Functional Diversity of Eukaryotic Life in the Oceans through

Transcriptome Sequencing. *PLOS Biol.* **12**, e1001889 (2014).

11. Moreira, D. & López-García, P. The rise and fall of Picobiliphytes: How assumed autotrophs

turned out to be heterotrophs. *BioEssays* **36**, 468–474 (2014).

12. Kawachi, M. *et al.* Rappemonads are haptophyte phytoplankton. *Curr. Biol.* **31**, 2395–2403

(2021).

13. Choi, C. J. *et al.* Newly discovered deep-branching marine plastid lineages are numerically rare

but globally distributed. *Curr. Biol.* **27**, R15–R16 (2017).

14. Schön, M. E. *et al.* Single cell genomics reveals plastid-lacking Picozoa are close relatives of red

algae. *Nat. Commun.* **12**, 6651 (2021).

15. Sunagawa, S. *et al.* Tara Oceans: towards global ocean ecosystems biology. *Nat. Rev. Microbiol.*

18, 428–445 (2020).

16. Delmont, T. O. *et al.* Functional repertoire convergence of distantly related eukaryotic plankton

lineages abundant in the sunlit ocean. *Cell Genomics* **2**, 100123 (2022).

17. Ruscheweyh, H.-J. *et al.* Cultivation-independent genomes greatly expand taxonomic-profiling

capabilities of mOTUs across various environments. *Microbiome* **10**, 212 (2022).

18. Dmitrijeva, M. *et al.* The mOTUs online database provides web-accessible genomic context to

taxonomic profiling of microbial communities. *Nucleic Acids Res.* **53**, D797–D805 (2025).

19. Delmont, T. O. *et al.* Nitrogen-fixing populations of Planctomycetes and Proteobacteria are

abundant in surface ocean metagenomes. *Nat. Microbiol.* **3**, 804–813 (2018).

20. Eren, A. M. *et al.* Anvi'o: an advanced analysis and visualization platform for 'omics data. *PeerJ*

3, e1319 (2015).

21. Eren, A. M. *et al.* Community-led, integrated, reproducible multi-omics with anvi'o. *Nat.*

Microbiol. **6**, 3–6 (2021).

22. Kim, J. I. *et al.* Evolutionary Dynamics of Cryptophyte Plastid Genomes. *Genome Biol. Evol.* **9**,

1859–1872 (2017).

23. de Vargas, C. *et al.* Eukaryotic plankton diversity in the sunlit ocean. *Science* **348**, 1261605

(2015).

24. Muñoz-Gómez, S. A. & Slamovits, C. H. Plastid Genomes in the Myzozoa. in *Advances in*

Botanical Research (eds Chaw, S.-M. & Jansen, R. K.) vol. 85 55–94 (Academic Press, 2018).

25. Matsuo, E. *et al.* Comparative Plastid Genomics of Green-Colored Dinoflagellates Unveils

Parallel Genome Compaction and RNA Editing. *Front. Plant Sci.* **13**, 918543 (2022).

26. Kamikawa, R. *et al.* Plastid Genome-Based Phylogeny Pinpointed the Origin of the Green-
Colored Plastid in the Dinoflagellate *Lepidodinium chlorophorum*. *Genome Biol. Evol.* **7**, 1133–
1140 (2015).

27. de Vries, J. & Archibald, J. M. Plastid genomes. *Curr. Biol.* **28**, R336–R337 (2018).

28. Ha, J.-S. *et al.* Plastid Genome Evolution of Two Colony-Forming Benthic *Ochrosphaera*
neapolitana Strains (Coccolithales, Haptophyta). *Int. J. Mol. Sci.* **24**, 10485 (2023).

29. Baños, H., Susko, E. & Roger, A. J. Is Over-parameterization a Problem for Profile Mixture
Models? *Syst. Biol.* **73**, 53–75 (2024).

30. Szánthó, L. L., Lartillot, N., Szöllősi, G. J. & Schrempf, D. Compositionally Constrained Sites
Drive Long-Branch Attraction. *Syst. Biol.* **72**, 767–780 (2023).

31. Williamson, K. *et al.* A robustly rooted tree of eukaryotes reveals their excavate ancestry. *Nature*
640, 974–981 (2025).

32. Muñoz-Gómez, S. A. *et al.* Site-and-branch-heterogeneous analyses of an expanded dataset
favour mitochondria as sister to known Alphaproteobacteria. *Nat. Ecol. Evol.* **6**, 253–262 (2022).

33. Crotty, S. M. *et al.* GHOST: Recovering Historical Signal from Heterotachously Evolved
Sequence Alignments. *Syst. Biol.* **69**, 249–264 (2020).

34. Susko, E. & Roger, A. J. On Reduced Amino Acid Alphabets for Phylogenetic Inference. *Mol.*
Biol. Evol. **24**, 2139–2150 (2007).

35. Criscuolo, A. & Gribaldo, S. BMGE (Block Mapping and Gathering with Entropy): a new
software for selection of phylogenetic informative regions from multiple sequence alignments.
BMC Evol. Biol. **10**, 210 (2010).

36. Pierella Karlusich, J. J. *et al.* A robust approach to estimate relative phytoplankton cell
abundances from metagenomes. *Mol. Ecol. Resour.* **23**, 16–40 (2023).

37. Hirakawa, Y. & Ishida, K.-I. Polyploidy of Endosymbiotically Derived Genomes in Complex
Algae. *Genome Biol. Evol.* **6**, 974–980 (2014).

38. Shrestha, B. *et al.* Global metagenomics reveals plastid diversity and unexplored algal lineages.
bioRxiv 2025.03.28.644651 (2025) doi:10.1101/2025.03.28.644651.

39. Strassert, J. F. H., Irisarri, I., Williams, T. A. & Burki, F. A molecular timescale for eukaryote

evolution with implications for the origin of red algal-derived plastids. *Nat. Commun.* **12**, 1879

(2021).

40. McFadden, G. I. The cryptomonad nucleomorph. *Protoplasma* **254**, 1903–1907 (2017).

41. Tanifuji, G. *et al.* Complete Nucleomorph Genome Sequence of the Nonphotosynthetic Alga

Cryptomonas paramecium Reveals a Core Nucleomorph Gene Set. *Genome Biol. Evol.* **3**, 44–54

(2011).

42. Rice, D. W. & Palmer, J. D. An exceptional horizontal gene transfer in plastids: gene replacement

by a distant bacterial paralog and evidence that haptophyte and cryptophyte plastids are sisters.

BMC Biol. **4**, 31 (2006).

43. Khan, H. *et al.* Plastid Genome Sequence of the Cryptophyte Alga *Rhodomonas salina*

CCMP1319: Lateral Transfer of Putative DNA Replication Machinery and a Test of Chromist

Plastid Phylogeny. *Mol. Biol. Evol.* **24**, 1832–1842 (2007).

44. Keeling, P. J. & Eglit, Y. Openly available illustrations as tools to describe eukaryotic microbial

diversity. *PLOS Biol.* **21**, e3002395 (2023).

45. Burki, F., Roger, A. J., Brown, M. W. & Simpson, A. G. B. The New Tree of Eukaryotes. *Trends*

Ecol. Evol. **35**, 43–55 (2020).

46. Ponce-Toledo, R. I., Moreira, D., López-García, P. & Deschamps, P. Molecular phylogeny of the

SELMA translocation machinery recounts the evolution of complex photosynthetic eukaryotes.

Mol. Biol. Evol. msaf167 (2025) doi:10.1093/molbev/msaf167.

47. Pietluch, F., Mackiewicz, P., Ludwig, K. & Gagat, P. A New Model and Dating for the Evolution

of Complex Plastids of Red Alga Origin. *Genome Biol. Evol.* **16**, evae192 (2024).

48. Gaïa, M. *et al.* Mirusviruses link herpesviruses to giant viruses. *Nature* **616**, 783–789 (2023).

49. Gaïa, M. *et al.* Closest relatives of poxviruses are spread in the gut of humans and animals

worldwide: the egoviruses. 2024.03.23.586382 Preprint at

https://doi.org/10.1101/2024.03.23.586382 (2025).

50. Martinez-Gutierrez, C. A. & Aylward, F. O. Phylogenetic Signal, Congruence, and Uncertainty

across Bacteria and Archaea. *Mol. Biol. Evol.* **38**, 5514–5527 (2021).

51. Shaw, J. & Yu, Y. W. Fast and robust metagenomic sequence comparison through sparse

chaining with skani. *Nat. Methods* **20**, 1661–1665 (2023).

52. Puerta, M. V. S., Bachvaroff, T. R. & Delwiche, C. F. The Complete Plastid Genome Sequence of

the Haptophyte *Emiliania huxleyi*☐: a Comparison to Other Plastid Genomes. *DNA Res.* **12**, 151–

156 (2005).

53. Butenko, A., Lukeš, J., Speijer, D. & Wideman, J. G. Mitochondrial genomes revisited: why do

different lineages retain different genes? *BMC Biol.* **22**, 15 (2024).

54. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform.

Bioinformatics **25**, 1754–1760 (2009).

55. Lang, B. F. *et al.* Mitochondrial genome annotation with MFannot: a critical analysis of gene

identification and gene model prediction. *Front. Plant Sci.* **14**, 1222186 (2023).

56. Greiner, S., Lehwark, P. & Bock, R. OrganellarGenomeDRAW (OGDRAW) version 1.3.1:

expanded toolkit for the graphical visualization of organellar genomes. *Nucleic Acids Res.* **47**,

W59–W64 (2019).

57. Janouškovec, J., Horák, A., Oborník, M., Lukeš, J. & Keeling, P. J. A common red algal origin of

the apicomplexan, dinoflagellate, and heterokont plastids. *Proc. Natl. Acad. Sci.* **107**, 10949–

10954 (2010).

58. Ponce-Toledo, R. I. *et al.* An Early-Branching Freshwater Cyanobacterium at the Origin of

Plastids. *Curr. Biol.* **27**, 386–391 (2017).

59. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search

tool. *J. Mol. Biol.* **215**, 403–410 (1990).

60. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

61. Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973 (2009).

62. Kozlov, A., Darriba, D., Flouri, T., Morel, B. & Stamatakis, A. RAxML-NG: a fast, scalable and

user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* **35**, 4453–4455

(2019).

63. Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in

the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

64. Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.

ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods* **14**, 587–589

(2017).

65. Whelan, S., Irisarri, I. & Burki, F. PREQUAL: detecting non-homologous characters in sets of

unaligned homologous sequences. *Bioinformatics* **34**, 3929–3930 (2018).

66. Susko, E., Lincker, L. & Roger, A. J. Accelerated Estimation of Frequency Classes in Site-

Heterogeneous Profile Mixture Models. *Mol. Biol. Evol.* **35**, 1266–1283 (2018).

67. Lartillot, N., Rodrigue, N., Stubbs, D. & Richer, J. PhyloBayes MPI: phylogenetic reconstruction

with infinite mixtures of profiles in a parallel environment. *Syst. Biol.* **62**, 611–615 (2013).

68. Liu, C. *et al.* The Influence of the Number of Tree Searches on Maximum Likelihood Inference in

Phylogenomics. *Syst. Biol.* **73**, 807–822 (2024).

69. Shimodaira, H. An Approximately Unbiased Test of Phylogenetic Tree Selection. *Syst. Biol.* **51**,

492–508 (2002).

70. Yu, G. Using ggtree to Visualize Data on Tree-Like Structures. *Curr Protoc Bioinform* **69**, e96

(2020).

71. Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence searching for the

analysis of massive data sets. *Nat. Biotechnol.* **35**, 1026–1028 (2017).

72. Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative

genomics. *Genome Biol.* **20**, 238 (2019).

73. Boyer, T. P. *et al.* World ocean database 2013. National Oceanographic Data Center (U.S.),

Ocean Climate Laboratory https://doi.org/10.7289/V5NZ85MT (2013).

74. Aumont, O., Ethé, C., Tagliabue, A., Bopp, L. & Gehlen, M. PISCES-v2: an ocean

biogeochemical model for carbon and ecosystem studies. *Geosci. Model Dev.* **8**, 2465–2513

(2015).

75. Li, D., Liu, C.-M., Luo, R., Sadakane, K. & Lam, T.-W. MEGAHIT: an ultra-fast single-node

solution for large and complex metagenomics assembly via succinct de Bruijn graph.

Bioinformatics **31**, 1674–1676 (2015).

76. Guillou, L. *et al.* The Protist Ribosomal Reference database (PR2): a catalog of unicellular

eukaryote Small Sub-Unit rRNA sequences with curated taxonomy. *Nucleic Acids Res.* **41**,

D597–D604 (2013).

77. Mahé, F., Henry, N., de Vargas, C., Tara Oceans Consortium, C. & Tara Oceans Expedition, P.

rDNA 18S V9 metabarcoding tables (Swarm) for Tara Oceans Expedition (2009-2013), including

Tara Polar Circle Expedition (2013). Zenodo (2022).

78. Zavadska, D., Henry, N., Auladell, A., Berney, C. & Richter, D. J. Diverse patterns of

correspondence between protist metabarcodes and protist metagenome-assembled genomes.

PLOS ONE **19**, e0303697 (2024).

79. Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag, New York).

80. Aphalo, P. ggpmisc: Miscellaneous Extensions to 'ggplot2'. (2025).

81. Shulgina, Y. & Eddy, S. R. Codetta: predicting the genetic code from nucleotide sequence.

Bioinformatics **39**, btac802 (2022).

994
995
Fig. 1. Global phylogenetic analysis of plastid genomes from the sunlit ocean.
Maximum-Likelihood phylogeny inferred using 93 plastid-encoded genes and the LG+F+I+G4 model in IQ-
TREE. The phylogeny contains 179 manually selected reference genomes (166 plastid and 13 cyanobacterial
genomes) and 660 plastid MAGs (ptMAGs) from various algal groups; cyanobacteria (including the *Paulinella*
plastid) were set as outgroup. Ring 1 around the phylogeny indicates algal groups; Ring 2 depicts whether the
taxon is a reference sequence or a MAG; Ring 3 depicts the estimated size of each marine taxon as determined
by metagenomic mapping of 937 *Tara* Ocean metagenomes to each genome (see Methods). Colpodellid
branches have been shortened to 40% of their original length to improve visual clarity. The full phylogeny with
taxon labels is shown in Supplementary Fig. 7.

1005
Fig. 2. Distribution patterns of plastid-encoded genes across algae with red plastids
A binary heatmap and dendrograms of genes and taxa generated using the UPGMA algorithm, based on the
presence or absence of 237 plastid protein-coding genes. Blue and grey boxes represent gene presence and

absence, respectively. For leptophytes, present genes are highlighted in red for improved clarity. A larger

version of this figure, including gene names, is presented in Supplementary Fig. 17.

Fig. 3. Phylogenomic analyses based on a reduced taxon-sampling.

(a) Bayesian inference based on 20,292 amino acid sites across 107 taxa under the CAT+GTR+G4 model.

Branch support values are given in the following order: Bayesian posterior probabilities (PPs), non-parametric

bootstrap support (BS; 100 replicates) under the GTR+CAT-PMSF model, and ultrafast bootstrap support

(UFB; 1,000 replicates) under the LG+MEOW80+G4 model. The inset shows the alternative topology of the

CHL clade. **(b)** Alignment of the *rpl36* gene across corresponding taxa in the tree. **(c)** Maximum likelihood

estimates under different models for alternate positions of leptophytes within the CHL clade. Likelihoods in

bold text represent the highest scoring topology. Difference in log-likelihood scores between each topology and

the best-scoring topology is shown by data bars in blue. A single asterisk indicates topologies rejected by the

Bonferroni-corrected chi-squared test, while two asterisks indicate topologies also rejected by an Approximately

Unbiased (AU) test. See also Supplementary Tables 3 and 4.

1022

Fig. 4. Phylogeny and abundance patterns of a putative leptophyte mitochondrial genome

Maximum-Likelihood phylogeny inferred using 28 mitochondrial genes in IQ-TREE. Support values on

branches indicate Bayesian posterior probabilities (PPs) under the CAT+GTR+G model and ultrafast bootstrap

support (UFB; 1,000 replicates) under the LG+C60+G4 model. The phylogeny contains 34 mitochondrial

MAGs (mtMAGs, highlighted in bold) recovered from a coassembled metagenomic dataset of the six *Tara*

Oceans samples with the highest Lepto-01 abundance, and 68 manually selected reference genomes. The

phylogeny was arbitrarily rooted with jakobids (discobans). The inset shows the abundance correlation between

the putative Lepto-01 mtMAG and the Lepto-01 ptMAG across *Tara* Oceans metagenomes from the 0.22-3µm

size fraction.

1032

1033 **Fig. 5. Main proposed models for evolution of complex red plastids**

1034 The trees depict hypothetical host relationships with red algae, cryptophytes, haptophytes, and leptophytes

1035 highlighted. Numbers on the tree represent the level of endosymbiosis events. Inset phylogenies depict the

1036 corresponding plastid phylogenies. Plastid transfers to ochrophytes and myzozoans are not depicted for clarity.

1037 Illustrations of algae are from [44].

Pelagophytes + Dictyochophytes

Diatoms + Bolidophytes

Other Ochrophytes

Colpodellids

Haptophytes

Cyanobacteria + *Paulinella*
Glaucophytes
Streptophytes
Prasinodermophytes
Chlorophytes

Euglenophytes

Chlorarachniophytes

Dinophytes

Rhodophytes

Cryptophytes

Leptophytes

1   2   3

0.4

1   Main Group

Ochrophytes

Cryptophytes

Haptophytes

Leptophytes

Red algae

Viridiplantae and secondary green lineages

Glaucophytes

Cyanobacteria

2   Type

MAG   Reference

3   Size fraction

pico-nano: 0.2-5 µm   micro: 20-180 µm

nano: 3-20 µm   meso: 180-2,000 µm

Galdieria sulphuraria
Cyanidium caldarium
Cyanidiococcus yangmingshanensis
Cyanidioschyzon merolae
Bulboplastis apyrenoidosa
Gracilaria changii
Chondrus crispus
Thaumatella adunca
Balbiania investiens
Gelidium coulteri
Calliarthron tuberculosum
Porphyra purpurea
Compsopogon caeruleus
Bangiopsis subsimplex
Porphyridium sordidum

Rhodophytes

Cryptomonas paramecium
Cryptomonas curvata
Guillardia theta
Baffinella frigidus
Storeatula sp. CCMP1868
Chroomonas mesostigmatica
Rhodomonas salina
Teleaulax amphioxeia

Cryptophytes

Lepto-01   **Leptophytes**

Diacronema lutheri
Pavlomulina ranunculiformis
Phaeocystis antarctica
Phaeocystis globosa
Chrysochromulina parva
Gephyrocapsa huxleyi
Isochrysis galbana

Haptophytes

Proboscia sp.
Rhizosolenia fallax
Pseudo−nitzschia cuspidata
Thalassionema frauenfeldii
Fistulifera solaris
Actinocyclus subtilis
Cylindrotheca closterium
Asterionella formosa
Chaetoceros laevisporus
Nitzschia supralitorea
Eunotogramma sp.
Thalassiosira nordenskioeldii
Rhizochromulina marina
Octactis speculum
Pseudopedinella elastica
Vaucheria litorea
Dictyopteris divaricata
Fucus vesiculosus
Characiopsis acuta
Ochromonas sp. CCMP1393
Synura uvella

Ochrophytes

Missing in
haptophytes

Core genes

Rhodophytes and
cryptophytes

a

b

c

	ML analyses using fixed topology trees				
Dataset	Untreated				Stationary trimmed
Model	GTR-CAT-PMSF	LG-MEOW80	LG-MEOW80 + GHOST (8 cat)	LG-MEOW80 + GF-MIX †	LG-MEOW80
Topology					
Leptophytes sister to haptophytes and cryptophytes	**-931909.236**	**-1056271.087**	**-1045458.696**	-1055620.569 *	**-592787.090**
Leptophytes sister to haptophytes	-931920.074 *	-1056274.766	-1045464.078 *	**-1055586.193**	-592793.164 *
Leptophytes sister to cryptophytes	-931924.826 **	-1056273.024	-1045463.152 *	-1055602.069 *	-592795.016 *

* Rejected by Bonferroni corrected χ2 test

** Rejected by Bonferroni corrected χ2 test and AU test

† AU test not possible under GF-MIX model

Abundance correlation across *Tara* Oceans metagenomes

$R^2 = 0.96$, $P < 0.001$

a. Plastid transfer from cryptophytes to ancestor of haptophytes and leptophytes

H-sister

Haptophytes

Leptophytes

Cryptophytes

Ochrophytes

non-photosynthetic Stramenopiles

Myzozoans

non-photosynthetic Alveolates

Rhizarians

Telonemids

Haptophytes

gene loss

Leptophytes

Centrohelids

nuc. loss

Red algae (Cyanodiophytina)

③

Red algae (Rhodophytina)

Green plants

②

Glaucophytes

①

rpl36

Cryptophytes

non-photosynthetic cryptistans

b. Plastid transfer from haptophytes to cryptophytes

HC-sister

Haptophytes

Cryptophytes

Leptophytes

Ochrophytes

non-photosynthetic Stramenopiles

Myzozoans

non-photosynthetic Alveolates

Rhizarians

Telonemids

nuc. loss gene loss

Haptophytes

rpl36

gene loss

Leptophytes

②

Centrohelids

Red algae (Cyanodiophyta)

Red algae (Rhodophytina)

Green plants

③

Glaucophytes

①

Cryptophytes

non-photosynthetic cryptistans

 rpl36 replacement

(nuc. loss) nucleomorph loss

 gene loss from plastid genome