# The inflammasome adaptor protein ASC triggers immunometabolic dysregulation induced by dysbiosis during obesity

Martinez-Garcia, JJ[1,2*]; Rodríguez, S[1]; Guijarro, A[1]; Quevedo-Romero, I[1], Mateo, SV[1]; Molina-López, C[1]; Hurtado-Navarro, L[1]; Baroja-Mazo, A[1]; Pelegrín, P[1,2*].

[1]Molecular Inflammation Group, Biomedical Research Institute of Murcia (IMIB), Murcia, Spain

[2]Department of Biochemistry and Molecular Biology B and Immunology, Faculty of Medicine, University of Murcia, Murcia, Spain

**Correspondence to**: jjmartinezgarciaio@gmail.com and pablopel@um.es

**Running title**: ASC controls immunometabolism in obesity

**Abstract**

The innate immune system oversees the response to danger signals. One of the most complexes organs that develops this response is the gut, which is extremely sensitive to the microbial changes in response to high saturated fat diets (HFD), leaving the space to be colonised by opportunistic bacteria, and triggering the innate immune response. Different evidence suggests the potential of the inflammasome in diet-induced inflammation, but little is known about its impact on the gut immunometabolic dysregulation. In this work, we report that the inflammasome adaptor apoptosis speck-like protein with a caspase activation domain (ASC) is essential for the development of obesity and insulin resistance during a high-fat diet (HFD). In ASC-deficient mice (*Pycard*$^{-/-}$), lower weight gain and improved glucose tolerance were observed in comparison to wild type mice. Therefore, the absence of ASC reduces intestinal permeability and improves the benefitial effects of the gut microbiota, decreasing dysbiosis, making the intestinal environment healthier. Additionally, *Pycard*$^{-/-}$ mice presented an enhanced glycolysis and aerobic metabolism, contributing to reduced inflammation, fat accumulation in the liver and improved hepatic metabolic function. Developing therapies targeting ASC could be beneficial to treat obesity, type II diabetes, and metabolism-associated liver diseases.

**Keywords:** ASC, inflammasome, microbiota, metabolism, glucose tolerance, macrophages, intestinal permeability, high-fat-diet

**Abbreviations list:** ASC: Apoptosis-associated Speck-like protein containing a CARD; cDNA: Complementary Deoxyribonucleic Acid; DEG: Differential Expressed Gene; DNA: Deoxyribonucleic Acid; FDR: False Discovery Rate; GO: Gene Ontology; GSDMD: Gasdermin D; GTT: Glucose Tolerance Test; HFD: High-Fat-Diet; IL: Interleukin; LPS: Lipopolysaccharide; mRNA: Messenger Ribonucleic Acid; NCD: Normal-Chow-Diet; NIN1: Ninjurin-1; NLR: Nucleotide-binding domain Leucine-rich Repeat receptor; NLRP: Nucleotide-binding domain Leucine-rich Repeat Pyrin domain-containing receptor; NOD: Nucleotide-binding and Oligomerization Domain; PCA: Principal Components Analysis; PCR: Polymerase Chain Reaction; PBS: Phosphate Saline Buffer; RNA: Ribonucleic Acid; TCA: Tricarboxylic Acids cycle; TLR: Toll-Like Receptor.

## Introduction

Nucleotide-binding domain leucine-rich repeat receptors (NLRs) constitute an important part of the inflammatory response, responding to a wide variety of danger and pathogens associated molecular patterns, especially those ones presented by bacteria[1,2]. NLRs, including the most studied NLRP3, are highly expressed in myeloid cells from the immune system as well as in epithelial cells[3]. NLRP3 and other NLRs can oligomerize forming inflammasomes that activate caspase-1 to mature the proinflammatory cytokines interleukin (IL)-1β and IL-18[4,5]. Caspase-1 also processes the protein gasdermin D (GSDMD), which N-terminal fragment form pores in the plasma membrane initiating the release of IL-1β and IL-18 and an immunogenic programmed cell death termed as pyroptosis[6]. During pyroptotic cell death the plasma membrane protein ninjurin-1 (NINJ1) induces membrane rupture[7]. Some inflammasomes require interaction and oligomerization of the apoptosis-associated speck-like protein containing a caspase-1 activating domain (ASC) to activate caspase-1. Large inflammasome oligomers of ASC are externalized to the extracellular space during pyroptosis to amplify the inflammatory response[8–13].

The intestinal microbiota plays a crucial role contributing to the host's health and survival[14]. This makes the gut extremely sensitive to microbiota changes, which controls colonisation by opportunistic bacteria and triggering innate immune responses, including inflammation[15,16] Subsequently, the effect of the gut microbiome controlling the activity of the inflammasome has been reported with both antibiotics and germ-free (GF) conditions[17,18]. The participation of the different inflammasomes, including the NLRP3, has been tightly associated with diet-related metabolic diseases, such as obesity and type II diabetes [19]. In this direction, the consume of diets with high saturated fats (HFD) induces an inflammatory reaction and potentiates IL-1β production [20]. Furthermore, the NLRP6 inflammasome, which is present in the intestinal epithelial cells, is capable of maintaining the gut microbiota homeostasis through the production of IL-18 [21]. Other studies demonstrate the participation of pyroptosis induced by GSDMD as a protective agent that controls the intestinal microbiota and repairs the intestinal mucosa [22]. However, there is controversy with the role of pyroptosis in obesity, as it is not only associated with an increase in obesity but is also capable of inducing metabolic dysregulations, such as blockage of cholesterol transport for subsequent catabolic processes, the accumulation of atheroma plaques, and the development of inflammatory diseases such as steatohepatitis [23–25]. This disparity highlights that the molecular

mechanism by which inflammation is induced by the inflammasome and pyroptosis can affect

glucose homeostasis and the development of metabolic pathologies. However, is still not well

defined the role of the common inflammasome adaptor protein ASC in these diseases, as

oligomers of ASC could be contributing to metabolic dysregulation, being a potential target

of study on the field of the immunometabolic cues regulating gut physiopathology.

In this study we identify that the deletion of the common inflammasome assembly adaptor

protein ASC protects mice in response to a HFD. This effect increases tolerance to glucose

and exacerbated glycolytic and aerobic metabolism in HFD conditions, with a decrease of the

inflammatory response. It also involves a change of the gut epithelial integrity and

permeability, and regulates gut microbiome. This work suggests ASC as a potential therapy

to treat obesity, metabolic dysfunction-associated steatotic liver disease, and type II diabetes,

which could positively impact on the lifestyle of those patients.

**Results**

*ASC deficiency prevents gain of weight and glucose intolerance under high-fat-diet conditions*

To monitor potential weight changes during the time dependent on the inflammasome, weight from C57/BL6 background wild type (*Pycard*[+/+]) mice togetherand *Pycard*[-/-] mice from same age was monitored during 5 months in normal chow diet. We observed that the *Pycard*[-/-] tended to gain less weight than the *Pycard*[+/+], but it was not significant **(Figure 1A)**. This body weight gain control was found in both males and females, revealing that *Pycard*[-/-] mice can prevent body weight gain independently on the sex differences **(Suppl. Figure 1A)**. Therefore, we next asked whether ASC deficiency can modulate weight gain in obesity after feeding with a HFD. Weigh of *Pycard*[-/-] tended to decrease when compared to wild type from the third month of HFD, but it was not significant **(Figure 1B)**. HFD-dependent weight gain was confirmed comparing HFD to normal chow diet (NCD) fed wild type mice **(Suppl. Figure 1B).** To explore if ASC oligomers were functioning extracellularly during HFD feeding, we rescue the ASC-dependent phenotype in *Pycard*[-/-] with intraperitoneal administration of recombinant ASC oligomers, but this treatment did not increase weight gain to the levels of the wild type **(Figure 1B)**. We next asked whether ASC is involved in the pathologic phenotype observed in metabolic syndrome. We then measured glucose levels in mice feed with HFD and observed an increase of glucose tolerance in the ASC-deficient mice respect to the wild type either at 3 and 5 months after HFD **(Figure 1C,D)**. Glucose tolerance of *Pycard*[-/-] mice under HFD was independent on the administration of recombinant ASC oligomers **(Figure 1C,D)**. However, ASC oligomers induced a tendency to increase the maximum peak of glycemia at 20 minutes after glucose injection **(Figure 1C,D)**.

*ASC controls intestinal permeability and peritoneal cell infiltration in high-fat-diet*

As expected, HFD induced intestinal permeability respect NCD fed mice **(Figure 2A)**. However, *Pycard*[-/-]mice fed with HFD presented a significant decrease of intestinal permeability, which was rescued by administration of recombinant ASC oligomers **(Figure 2B)**. In parallel to intestinal permeability increase, we also observed an increase of the number of infiltrated cells in the peritoneum upon HFD respect NCD fed wild type mice **(Figure 2C)**. Surprisingly the *Pycard*[-/-] mice presented a higher increase of peritoneal cells after 5 months of HFD when compared to wild type, which tended to decrease with the addition of the intraperitoneal recombinant ASC oligomers, but it was not significant **(Figure 2D)**. There was a positive correlation between

intestinal permeability and the number of peritoneal cells in *Pycard*[+/+] mice **(Figure 2E)**, but

intestinal permeability did not correlated to the increase of peritoneal cells in *Pycard*[-/-] mice

**(Figure 2F)**, suggesting independent processes.

To investigate inflammasome function in infiltrated peritoneal cells during HFD conditions, we

cultured adherent peritoneal cells from mice and measure IL-1β in cell supernatants. Peritoneal

cells from wild type mice fed with HFD presented IL-1β release by LPS treatment and after

classical NLRP3 inflammasome activation **(Figure 2G)**. As expected, IL-1β release induced by

nigericin in both NCD and HFD conditions presented a dependency on the NLRP3 inhibitor

MCC950 **(Figure 2G)**. However, LPS induced IL-1β release in HFD was also dependent on

NLRP3 activity **(Figure 2G)**, suggesting increase sensitivity to activate NLRP3 in peritoneal

infiltrated cells after HFD. As expected, IL-1β release in peritoneal cells from *Pycard*[-/-] mice fed

with HFD was decreased when treated with LPS and nigericin, independently of the intraperitoneal

injection of recombinant ASC oligomers **(Figure 2H, Suppl. Figure 2A)**. This effect could be due

to the metabolization of ASC oligomers in vivo or little capture of these oligomers by peritoneal

cells during the 5 months of HFD. As control, IL-6 release from peritoneal cells was independent

on the genotype **(Suppl. Figure 2B,C)**. These data suggest that ASC deficiency could protect

intestinal permeability during obesity.

***ASC deficiency increases gut microbiota biodiversity during high-fat-diet***

We next wondered if ASC could control gut dysbiosis induced by 5 months of HFD. We

determined 16S RNA in faeces and found that the microbiota composition changed in *Pycard*[-/-]

mice fed with HFD in comparison with the *Pycard*[+/+] mice **(Figure 3A)**. Intraperitoneal

administration of recombinant ASC oligomers reverted gut microbiota composition of *Pycard*[-/-] to

the dysbiosis microbial composition induced by the HFD **(Figure 3A)**. This suggests that during

HFD, inflammasome oligomers might be eliminated by the intestine where it could impact in gut

microbiota composition. Within the most abundant bacteria that was influenced by the deficiency

of ASC, we encounter mainly *Akkermansia muciniphila, Blautia coccoides, Romboutsia ilealis*

and *Bacteroides acidifaciens*, which constitute approximately 90% of the microbiota biodiversity

**(Figure 3A).** In particular, ASC deficiency increase the proportion of *B. coccoides* and decrease

*A. muciniphila* when compared to wild type mice, while administration of recombinant ASC

oligomers restored this ratio, and also increased *R. ilealis* and *B. acidifaciens* **(Figure 3A)**. We

also found that the abundance of Verrucomicrobiota is about the 25% of the microbiota
composition in wild type mice under HFD **(Figure 3B)**. However, the ratio of Verrucomicrobiota
respect to the other main phylum, Firmicutes and Bacteroidota, decreased in *Pycard*[-/-] with HFD
**(Figure 3C)**. Intraperitoneal administration of recombinant ASC oligomers also rescued phylum
ratio, increasing Verrucomicrobiota phyla **(Figure 3D)**. As a quality control, we observed in a
different batch of mice that the NCD ASC[+/+] group presented a very few proportions of
Verrucomicrobiota **(Suppl. Figure 3A)**, which strongly increased in HFD conditions **(Suppl.**
**Figure 3B)**. Regarding the importance of the low abundance of Verrucomicrobiota to control
dysbiosis, and maintain the microbiota biodiversity, *A. muciniphila*, the main bacteria from the
Verrucomicrobiota phyla, tended to decrease in *Pycard*[-/-] mice fed with HFD **(Figure 3E)**.
However, other top 5 abundant bacteria, *Bacteroides* did not change the relative abundance
**(Figure 3F)**. The decrease of these bacteria left the niche to the colonization of new bacteria that
incremented the microbiota abundance, such as *Enteractinococcus* and *Apostotipes* **(Figure 3G**
**and 3H)**. For these bacteria, the intraperitoneal administration of recombinant ASC oligomers in
*Pycard*[-/-] mice under HFD rescued dysbiosis, and in the case of *Bacteroides*, it strongly increased
the abundance contributing to decrease the microbiota biodiversity **(Figure 3E-H)**. As a control
quality, we verified that the abundance of *A. muciniphila* in wild type HFD conditions tended to
increase respect the NCD fed controls **(Suppl. Figure 3C)**. These data corroborated that *Pycard*[-/-]
mice prevent the dysbiosis, creating a benefitial environment in the gut by the lack of extracellular
ASC oligomers.

### *ASC induces bacteria attachment in intestinal villi during high-fat-diet*

Since the absence of ASC reduces intestinal permeability and prevents the dysbiosis, we next
studied if the intestinal mucosa was involved in the regulation of this phenomenon under HFD.
We observed that wild type mice fed with HFD presented a severe bacterial colonization in the
small intestinal *villi* **(Figure 4A)**. Bacteria attached to the *villi* decreased in ASC-deficient mice,
and mucus was present in the border of each intestinal *villi* **(Figure 4B)**. When recombinant ASC
oligomers were intraperitoneally injected on *Pycard*[-/-] mice with HFD, the number of bacteria
within the intestinal *villi* was not rescued **(Figure 4C)**, but mucus was absent in comparison with
the *Pycard*[-/-] mice without recombinant ASC oligomers administration **(Figure 4C)**. This suggests
that extracellular ASC oligomers partially rescue the protective effect induced by the absence of

endogenous ASC. As a control, when the intestinal *villi* of wild type mice fed with HFD were
compared with the NCD fed mice, we observed a normal intestinal histology with low bacteria in
*villi* and abundant mucus under NCD **(Suppl. Figure 4A,B)**.

Bulk RNA sequencing in the small intestine of mice fed with HFD demonstrated that the *Pycard*
*-/-* mice have increased expression of metabolic, insulin degrading enzyme and protein synthesis
transcript genes (*Gdpd3* and *Ide*) **(Figure 4D)**. ASC deficiency also resulted in a decreased
expression of genes related with intestinal inflammation and cancer (e.g. *Ern2)*, epithelial integrity
(e.g. *Slc26a6 or Nts)*, hypoxia (e.g. *Ddit4)*, and chemokines produced in response against bacteria
(e.g. *Ccl7)* **(Figure 4D)**. Principal component analysis (PCA) was conducted to visualize sample-
level clustering based on the expression of the top 20 genes **(Suppl. Figure 4E)**. Administration
of recombinant ASC specks in *Pycard-/-* mice during HFD partially rescued gene expression profile
in the gut towards wild type genotype **(Figure 4D)**. Gene Ontology (GO) analysis from the
differential expressed genes in the small intestine revealed that meanwhile the most enriched
pathways in the HFD fed *Pycard-/-* mice were the regulation of the T cell response and the
regulation of the MHC class I and II activity, on *Pycard+/+* mice, the AIM2 and NLRP3
inflammasome, hormonal activity and positive regulation of the humoral immune response were
between the top 10 highly expressed pathways **(Figure 4E)**. Supporting this, the concentration of
IL-1β in small intestine decreased in *Pycard-/-* feed with a HFD compared with wild type **(Figure**
**4F)**. Intraperitoneal administration of recombinant ASC oligomers reverted IL-1β production in
the intestine **(Figure 4F)**. However, IL-6 production did not experimented significant changes in
*Pycard-/-* mice **(Suppl. Figure 4C)**. We confirmed that there is not ASC gene expression during
the administration of recombinant ASC oligomers **(Suppl. Figure 4D)**. All together summarizes
that small intestine mucosa in absence of extracellular ASC oligomers initiates a protective
mechanism under HFD reducing innate immunity, increasing epithelial integrity and the aerobic
environment, together with the decrease of bacteria and the reduced intestinal permeability.

***ASC deficiency increases the glycolytic and tricarboxylic acids pathway in the small intestine***
***under high-fat-diet***

As the RNA sequencing revealed a tendency to increase some metabolic genes under HFD
treatment, which could also explain the elevation of IL-1β expression, we then analyzed in the
small intestine the expression of key metabolic genes related with the glycolysis and the

tricarboxylic acids (TCA) cycle. Interestingly, we found that the glycolytic *Pkm2* gene coding for

the pyruvate kinase enzyme was highly expressed in the *Pycard*[-/-] HFD fed mice **(Figure 5A)**.

Contrarily, the anaerobic respiration-related gene *Ldha*, corresponding to lactate dehydrogenase,

was decreased in the *Pycard*[-/-] mice **(Figure 5B).** Consequently, we observed an increase

expression of *Pdha* gene, which encodes the pyruvate dehydrogenase enzyme **(Figure 5C)**. This

observation could indicate that the high glycolytic state on the *Pycard*[-/-] mice small intestine under

HFD drives to the aerobic metabolism, and not to the anaerobic pathway. To verify this switch,

we also analyzed the intestinal gene expression of the enzyme that belonged to the TCA cycle. We

reported that *Glud1* and *Sdha* genes expression generally was increased in the *Pycard*[-/-] HFD mice

**(Figure 5D,E)**. We also demonstrated that the β-oxidation-related gene *Cpt1a* expression

increased in *Pycard*[-/-] HFD mice respect the wild type **(Figure 5F)**. These results aligns with the

fact that the *Pycard*[-/-] have decreased weight and increased glucose tolerance independently on the

HFD due to a high catabolism and energy uptake. This effect was dependent on the extracellular

function of oligomers of ASC, as the intraperitoneal administration of recombinant ASC oligomers

tended to revert the elevated catabolic environment towards the levels of the wild type mice

**(Figure 5A-F)**.

### *ASC deficiency enhances hepatic metabolism and reduces inflammation, bile acids and fat accumulation under high-fat-diet*

We finally investigated if *Pycard*[-/-] mice have an altered liver physiopathology and metabolism

during HFD. Initially we observed that wild type mice fed with HFD presented an exacerbated fat

accumulation in the liver **(Figure 6A)**. Liver fat accumulation was absent in ASC-deficient mice

under HFD **(Figure 6B)**. Differentially to the effects observed in the small intestine, intraperitoneal

administration of recombinant ASC oligomers did not rescue wild type phenotype **(Figure 6C)**,

suggesting that extracellular ASC oligomers might impact small intestine, but not hepatic

physiopathologic consecuences, that might be controlled by intracellular ASC. The accumulation

of hepatic fat under HFD in wild type was corroborated when compared with NCD fed mice

**(Suppl. Figure 5A,B)**. Moreover, we observed increased expression of metabolic genes in the

liver of *Pycard*[-/-]HFD fed mice, in particular genes related with the PPARα signaling pathway

**(Figure 6D)**, the electron transport chain from the mitochondrial respiration **(Figure 6E)**, and the

synthesis of aminoacids derived from the TCA **(Figure 6F)**. Interestingly, we observed a decrease

expression of the classical bile acid synthesis enzyme *Cyp7a1* gene **(Figure 6G)**, which was confirmed when we measured the opposite effect on the *Shp* gene expression **(Figure 6H)**. Similarly to the small intestinal mucosa, in the liver there was also a tendency to decrease IL-1β production in the *Pycard*[-/-] mice fed with HFD **(Figure 6I)**. Similar to hepatic fat accumulation, the administration of recombinant ASC oligomers were not able to rescue gene expression in the liver of *Pycard*[-/-] mice **(Figure 6D-I)**, supporting the idea that extracellular ASC oligomers do not impact liver during HFD.

Overall, ASC is important to control metabolic syndrome induced by HFD, with a dual role, being intracellular ASC intestine permeability, obesity, glucose tolerance and hepatic inflammation, metabolism and steatosis; while extracellular ASC oligomers modulate dysbiosis and metabolism and inflammation in the small intestine.

**Discussion**

In this study we found that ASC deficiency causes a protective effect during HFD on associated obesity and metabolic syndrome effect. This protection was observed at different levels, including the decrease of intestinal permeability, dysbiosis, changes in the integrity, metagenomic, metabolism and physiology of the small intestine epithelia, and a robust decrease of the IL-1β production. This protective effect was also found in the hepatic metabolism, inflammation and steatosis. Dysbiosis, intestinal metabolism and IL-1β production were reverted in the *Pycard*[-/-] mice when recombinant ASC oligomers were intraperitoneally administrated, suggesting a differential role of extracellular ASC oligomers for microbiota, intestine and hepatic functions. Recent developments of nanobodies-targeting extracellular ASC oligomers (11) might be beneficial to treat dysbiosis associated to HFD.

We initially observed that the *Pycard*[-/-] mice fed with a HFD reduced obesity and increased glucose tolerance. But even the weight gain did not increase with the intraperitoneal addition of recombinant ASC oligomers, glucose tolerance tended to increase during the maximum glycemia peak, at 20 minutes after the glucose injection. This effect let us conclude that the protective effect of ASC depletion could be mediated by a mechanism of regulation which partially depends on the inflammasome induced ASC oligomerization and release. Extracellular ASC oligomers have been detected extracellularly in different pathologies, including autoinflammatory syndromes, amyloidosis, Alzheimer disease, and HIV infection (8,9,11,12,26). However, pathological consequences of the metabolic syndrome appearing at long term, including obesity and the steatosis (27,28), are not affected by the presence of extracellular ASC oligomers, being probably dependent on the intracellular functions of ASC.

Previous studies have reported a relation between intestinal permeability and immune function of peritoneal cells (29), which can be supported by the fact that during HFD intestinal permeability positively correlated with the number of infiltrated peritoneal cells found in our study. Conversely, we found an increase of peritoneal cells in the absence of ASC, probably as consequence of the reduced cell death of peritoneal cells upon inflammasome activation (30). The decrease in intestinal permeability might be related to the gut microbiota (31). Indeed, our results revealed that a HFD impact intestinal microbiota biodiversity, increasing the abundance of species of the Verrucomicrobiota Phylum, including *A. muciniphila*, a mucin-degrading bacteria which normally is benefitial, but in excess, it can dysregulate gut homeostasis (32–34). It is important to consider

that the pathological consequences of HFD could not be only attributed to *A. muciniphila*, but
maybe to a role of a homeostatic regulator of *A. muciniphila,* when either the excess or the low
abundance of this bacteria can be prejudicial. Moreover, the decrease of this bacteria in the gut
free a niche that can be colonised by less abundant bacteria, reducing dysbiosis and increasing the
microbial biodiversity as we observe for the *Pycard*[-/-] HFD fed mice group. As an example, the
decreasing mucus can reduce the attachment of segmented filamentous bacteria, which induces an
immune response to maintain homeostasis of the intestinal mucosa (35). The protection of
dysbiosis by the deficiency of ASC was also corroborated at histological level, where ASC
deficiency prevented the inclusion of bacteria in the intestinal mucosa, with an increase of mucus
between the intestinal *villi*. Additionally, administration of recombinant ASC oligomers rescue the
dysbiosis found in the wild type mice induced by HFD and also depleted the mucus formation,
which is mediated by the goblet cells from the intestinal mucosa (36). Concomitantly with this
effect, ASC controlled intestinal epithelia gene expression related with the innate immune response
against pathogens, inflammation, and the epithelial integrity, some of them are considered as
markers for obesity and cancer (37–39). Moreover, the increasing expression of the insulin
degrading enzyme (*Ide*) gene that we observed in *Pycard*[-/-] mice under HFD could explain the
increasing tolerance to glucose, as this enzyme protects from insulin resistance, and obesity in
response to high glycemia levels (40,41). Additionally, the intestinal production of IL-1β was
reduced in ASC deficient mice could be attributed to the NLRP3 inflammasome activation, as
previous studies have found (17,42). We do not discard that other inflammasomes, which are
highly expressed in intestinal epithelia can contribute either directly or indirectly to IL-1β
production (43–45).

Our study also found that small intestinal cells presented an exacerbated catabolic pathway in
*Pycard*[-/-] mice. This might be due because the increase glucose tolerance in an environment with
high energy can be a consequence of the augmented glycolytic pathway and TCA through the
enzyme's pyruvate kinase and pyruvate dehydrogenase, and decreasing anaerobic lactate
dehydrogenase activity. It is important to highlight that even if there is a controversy with the
PKM2 isoform, as it is described as an inflammatory potentiator and as inducer of lactate
dehydrogenase activity (characteristic of the Warburg effect), this is the PK isoform which is
predominant in the intestinal epithelia, and its presence could bypass between the high glycolysis
state and the elevated production of IL-1β that we observed in the *Pycard*[+/+] HFD fed mice, or the

increasing TCA, depending on the dimeric or tetrameric forms of PK respectively (46,47). This

regulation of glycolysis could reduce fat accumulation in the liver, and might increase hepatic

metabolism (48). In the liver of *Pycard*[-/-] HFD fed mice we also observed a reduction of IL-1β

production. The increased metabolism found in the liver could be the consequence to reduce fat

accumulation, in a very similar manner as occurs for the small intestine.

In summary, we report a mechanism of protection to the hyperglycaemic, westernized and high-

saturated fat diets which is dependent on the deficiency of the inflammasome adaptor protein ASC.

This mechanism is characterized by an exacerbated glucose metabolism together an increasing

inflammatory phenotype and a decreasing intestinal permeability by the prevention of dysbiosis

and mucus degradation. This mechanism has positive physiological consequences, reducing

obesity and increasing glucose tolerance and reducing type II diabetes, and reducing metabolic

dysfunction-associated steatosis liver disease, . The presence of extracellular ASC oligomers

partially restored the beneficial effects of ASC deficiency on the metabolic syndrome, particularly

at intestinal level and dysbiosis, but not in the liver. Further studies are warranted to evaluate

potential therapeutic approximations based in the inhibition of ASC oligomerization.

**Materials and Methods**

*Animal Models.* Animal studies were approved by the regional statement, the Animal Health Service of the General Directorate of Farming, Fishing and Aquiculture of the Council of Murcia (CARM, ID code Nº A13211201) once were positively valuated by the Animal Experimentation Ethics Committee (CEEA) of the University of Murcia (#738/2021) according to the 53/2013 Royal Decree and the 32/2007 National Law. Mice were bred in specific pathogen-free conditions with a 12:12 hr light-dark cycle in the animal facility from the University of Murcia, according to governmental guidelines No. 86/609/CEE. Age and gender-matched ASC-deficient (PYCARD-/-) and wild type C57BL6/J (*Pycard*$^{+/+}$) backgrounds mice were provided to a free access to a standard laboratory chow diet. For the obesity model, mice were fed with a 60% Fat HFD (D12492, Research Diets). Mice were autopsied after euthanasia with Isoflurane and cervical dislocation according to the ethical protocols of the national and international animal experimentation guidelines. After sacrifice, feces, peritoneal exudates, the right lobule of the liver, 1 cm of the proximal small intestine and blood samples were recovered. All samples were recovered strictly in the morning between Zeitgeber Time (ZT)0 and ZT4, to avoid the interference of the circadian changes. Histological samples were collected on Paraformaldehyde 4% diluted from the 32% stock (Sigma-Aldrich, Burlington, MA) at room temperature. For RTqPCR samples, tissues were collected on RNA later (Qiagen, Germany) and frozen on liquid Nitrogen. Feces were collected in a dry tube and frozen on liquid Nitrogen. Peritoneal cells were collected on Phosphate Saline Buffer (PBS) and kept at 4ºC. Plasma was collected in Heparin tubes (Microvette® CB 300 LH, Sarstedt, Germany). For longer tissue storage, samples were stored at -80ºC.

*In vivo treatments and tests*. Counted oligomers were diluted into 1x PBS at a concentration of $4 \times 10^6$ oligomers per 100µl, ready to use it in vivo. ASC Oligomers were weekly injected intraperitoneally at a concentration of $4 \times 10^6$ oligomers per mouse. Glucose Tolerance Test (GTT) was done at 3 and 5 months after the administration of the HFD upon gender and sex matched animals. Prior the GTT assay, animals were fasted the day before during 14 hours to avoid the food interference on the glucose levels. Basal fasted glycemia was measured before the Glucose treatment. Following, 1g/Kg of D-Glucose (Sigma-Aldrich) was intraperitoneally injected in each mouse, and glycemia was measured every 20 minutes, minimum during one hour. The last point was measured at 90 minutes after the glucose injection. At the end of the experiment, mice were

fed back again. For the intestinal permeability assay, mice were deprived of drinking water during one hour at ZT0. Immediately, 4KDa-FITC were administered orally by gavage at a concentration of 600mg/Kg of mouse. The following hour, blood was collected in the moment of the sacrifice, and plasma was obtained by centrifugation at 2000rcf during 8 minutes.

***ASC oligomers purification***. ASC Oligomers were produced and purified as we previously described in Alarcón-Vila et al. 2024 (doi:10.1038/s44321-024-00176-1) (10).

***Cells and Treatments.*** $5x10^5$ peritoneal myeloid cells were seeded overnight into each well from a 24 wells plate at 1 million/mL concentration (lymphocytes also were added but were not included into the count). The following day, supernatant was removed, leaving the attached cells for the treatment. *Escherichia coli* LPS 055:B5 (Sigma-Aldrich) was added at 1µg/mL during 4 hours. After priming, supernatant was removed and Nigericin (Sigma-Aldrich) was added at 10µM in E-total buffer (147 mM NaCl, 10 mM HEPES, 13 mM glucose, 2mM CaCl2, 1mM MgCl2, and 2mM KCl, pH 7.4) during 30 minutes. MCC950 (Sigma-Aldrich) was added simultaneously to the treatments at 10µM.

***Tissue lysates preparation.*** Liver and Small intestines previously stored at -80ºC were mechanically lysed and kept on ice. After being centrifuged at 13200rpm during 10 minutes. Supernatant was recovered. Protein concentration was measured using the Bradford reagent (Sigma Aldrich).

***ELISA determinations.*** After In vitro stimulations, supernatants from the peritoneal cells were collected and centrifuged at 13200rpm for 30 seconds. IL-1β and IL-6 were measured in the supernatants using the mouse Quantikine™ ELISA kits, catalog number MLB00C-1 and M6000B respectively (R and D Systems, Minneapolis, MN) following the manufacturer indications, and results were analysed in a Sinergy MX plate reader (BioTek, Winooski, VT).

***Histological samples preparation and analysis.*** Liver and small intestine were fixed in 4% p-formaldehyde for 24 hr, processed, paraffin-embedded, and sections stained with haematoxylin

and eosin using standard methods to evaluate damage. Slides were examined using a Zeiss Axio

Scope AX10 microscope with an AxioCamICC3 (Carl Zeiss, Germany).

***RNA extraction and RTqPCR determination.*** RNA was isolated using the RNA mini-extraction

kit according the manufacturer guidelines (Qiagen). Cell lysis was performed with the addition of

RLT buffer supplemented with β-mercaptoethanol. RNA integrity and concentration were

assessed using the NanoDrop spectrophotometer (Thermo Fisher Scientific, Waltham, MA)

Isolated RNA was reverse-transcribed with the iScript™ cDNA Synthesis Kit (BioRad, Hercules,

CA) according to the manufacturer's instructions. The resulting complementary DNA (equivalent

to 500 ng of total RNA) was amplified using the mix SYBR Green Premix ExTaq (Takara) and

detected at real time on the iCyclerMyiQ thermocycler (BioRad). Quantitative PCR was performed

using forward and reverse primers (sequences available in the web of Sigma-Aldrich). On

completion of the PCR amplification, a DNA melting curve analysis was carried out to confirm

the presence of a single amplicon. *Actb* gene was used as a reference gene to normalize the

transcript levels. Relative mRNA levels (2- DDCt) were determined by comparing the DCt values

for treated and control groups (DDCt). Primers for each gene were purchased from Sigma-Aldrich.

***RNA Library preparation and sequencing.*** RNA libraries were prepared using the QuantSeq 3′

mRNA-Seq Library Prep Kit (Tecan Genomics, Redwood City, CA), with Unique Dual Indices

(12nt), CA, following the manufacturer's protocol. Briefly, 500 ng of total RNA per sample was

used as input for poly(A) mRNA capture using oligo-dT beads. Captured mRNA was fragmented

and reverse-transcribed into cDNA. Second-strand synthesis and adapter ligation were performed

to generate double-stranded cDNA libraries. Libraries were indexed with 12-nucleotide unique

dual indices for multiplexing and amplified using polymerase chain reaction (PCR). The quality

and quantity of the final libraries were evaluated using the Agilent 2100 Bioanalyzer and the Qubit

4 Fluorometer (Thermo Fisher Scientific). Libraries were sequenced on a NextSeq 2000 system

(Illumina, San Diego, CA) in paired-end mode with a read length of 2 × 100 bp to a target depth

of 30 million reads per sample.

***RNAseq analysis.*** For RNAseq-Data processing and alignment, Raw sequencing data were processed using the bcl2fastq software (Illumina) to demultiplex and convert base call files into FASTQ format. Quality control of the FASTQ files was performed using FastQC (v0.11.9) to assess read quality, adapter content, and GC distribution. Low-quality reads and adapter sequences were trimmed using Trimmomatic (v0.39). Cleaned reads were aligned to the mouse reference genome (GRCm39) using STAR (v2.7.10a) with default parameters, generating BAM files for each sample. For the differential gene expression, Gene expression quantification was performed using featureCounts (v2.0.3) to obtain raw read counts per gene, based on the GENCODE M28 annotation. Count data were provided in a tab-delimited file containing unique read counts for each gene across samples. The dataset was preprocessed using R (R Core Team, 2023). Gene count matrices were normalized using the DESeq2 package (49), which applies a variance-stabilizing transformation to adjust for sequencing depth and RNA composition biases. Differential gene expression analyses were performed using DESeq2's generalized linear model framework, considering the experimental condition as the primary variable. To identify the top differentially expressed genes, the normalized counts were z-score transformed. Heatmaps of the top 20 differentially expressed genes were generated using the pheatmap package (50), with sample group annotations included for clarity. Gene identifiers were converted to gene symbols using the org.Mm.eg.db annotation database (51). The PCA was performed using the prcomp function in R, and the resulting components were plotted using ggplot2 (52). Pathway enrichment analysis was performed on differentially expressed genes (DEGs) using the Cluster Profiler package (v4.8.1) in R. Gene Ontology (GO) biological processes were analyzed, with pathways considered significant at a false discovery rate (FDR) < 0.05. The selected resulting pathways were visualized using dot plots generated with ggplot2 (v3.4.0). Pathway enrichment analysis was performed on DEGs using the Cluster Profiler package (v4.8.1) in R. Gene GO biological processes were analyzed. The resulting pathways were visualized using dot plots generated with ggplot2 (v3.4.0).

***Bacterial DNA extraction.*** Fecal samples were collected directly from the bedding-free cage floor to avoid contamination. Immediately after collection, fecal samples were frozen at −80 °C until further processing. Bacterial DNA from feces was extracted using the Purelink™ Microbiome DNA Purification Kit (Thermo Fisher Scientific), according to the manufacturer instructions. The

concentration and purity of extracted DNA were assessed with a NanoDrop spectrophotometer
(Thermo Fisher Scientific).

***16S ribosome RNA gene amplification, sequencing, and metagenomic analysis.*** For the
amplification of the 16S rRNA gene, primers targeting the V3-V4 hypervariable regions were
designed to cover the bacterial domain (84.2%), archaea (75.8%), based on the SILVA database
(March 2020). PCR reactions were performed using 10 ng of DNA as input, combining targeted
amplification with barcode addition for sample multiplexing. The resulting amplicons were
approximately 606 base pairs in length. The amplification process utilized the Quick-16S Plus
NGS Library Prep Kit (Zymo Research) according to the manufacturer's instructions. Following
amplification, libraries were pooled in equimolar volumes and purified using magnetic beads to
remove residual primers and small fragments, as specified by the kit's protocol. Sequencing of the
libraries was performed on the NextSeq 2000 platform (Illumina, San Diego, CA) using a paired-
end (2x300 bp) approach. This setup ensured high-resolution reads suitable for downstream
taxonomic and functional analyses. A minimum of 50,000 reads per sample was targeted to
achieve sufficient depth for robust microbial community profiling. For the bioinformatic analysis,
the raw data was first quality controlled by trimming low-quality bases and removing adapters
using Trimmomatic. Next, the sequences were denoised and chimeras were removed using the
DADA2 algorithm implemented in QIIME 2. Following denoising, the sequences were assigned
taxonomy at the genus level using the SILVA v138 database and a Naïve Bayes classifier, which
classifies each ASV based on its sequence similarity to known reference sequences. Phyloseq
package was used to create a phyloseq object from the ASV table, taxonomic classification, and
metadata. The data was then aggregated at the genus level using the tax_glom() function, and
relative abundances were calculated using the transform_sample_counts() function in the phyloseq
package. The entire workflow was performed using the phyloseq, ggplot2, dplyr, and writexl R
packages (52,53).

***Statistical analysis.*** For two variables analysis, normality tests Shapiro-Wilk and Kolmogorov-
Smirnov were used to elucidate parametric and non-parametric groups. Where at least, one of the
independent groups was non-parametric, non-parametric analysis was used. For testing differences
between two independent groups, Mann-Whitney test was used for non-parametric data, and

Student T test was used for parametric data. For testing differences between 3 or more independent groups, Kruskal-Wallis's test was used for non-parametric data, and One-Way-ANOVA test was used for parametric data. For three or more variables analysis, Two-Way-ANOVA test was used. Between each ANOVA test, Multivariant test analysis was used to compare each bidirectional comparison between single groups (Dunn´s test for the non-parametric Kruskal-Wallis multivariant analysis, Holm-Sidak´s test for the parametric One-Way ANOVA multivariant analysis, Bonferroni´s test for Two-Way ANOVA multivariant analysis). For correlations, non-parametric Spearman test was used. Outliers were discarded when were significant for the Grubb's test.

**Disclosure Statement and conflict of interest**

PP and JJMG declare that they are inventors in a patent filed on March 2020 by the Fundación para la Formación e Investigación Sanitaria de la Región de Murcia (PCT/EP2020/056729) for a method to identify NLRP3-immunocompromised septic patients. PP, JJMG and ABM are co-founders of Viva in vitro diagnostics SL, but declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest. The remaining authors declare no competing interests.

**Funding**

This work was supported by grants from MCIN/AEI/10.13039/501100011033 and European Union «Next Generation EU/PRTR» (grants PID2020-116709RB-I00, CNS2022-135105, PID2023-147531OB-I00 and RED2022-134511-T to PP), the Instituto Salud Carlos III (grant AC22/00009 to PP) and the Fundación Séneca (grant grant 21897/PI/22 to PP and 21967/JLI/22 to JJMG). JJGM was supported by a Maria Zambrano fellowship at University of Murcia. AG was supported by an FPI doctoral fellowship from *Ministerio de Ciencia, Innovación y Universidades* (PRE2021-100356). SVM was supported by a fellowship from Instituto Salud Carlos III (FI21/00073). CM-L was funded by the fellowship PRE2018-086824 (Ministerio economía y competitividad). LH-N was supported by the fellowship 21214/FPI/19 (Fundación Séneca, Región de Murcia, Spain).

**Acknowledgements**

517  We thank A Gomez for cell culture and lab assistance, and the members of Dr. Pelegrin's
518  laboratory for all comments and suggestions. We also acknowledge to the animal facility from the
519  University of Murcia, as well as the SPF-animal house, Pathology, Genomics and Bioinformatics
520  platforms from the Pascual Parrilla-Biomedical Research Institute of Murcia (IMIB-PP).
521

522  **Authors Contribution**

523  JJMG visualized and administered the research. JJMG and PP conceptualized, provided funding,
524  and wrote the manuscript. JJMG and SR did the formal analysis, wrote, and edited the manuscript.
525  ABM provided funding. JJMG, AG, IQR, SVM. CML and LHN performed the experimental
526  research.
527

REFERENCES

1. Medzhitov R. Origin and physiological roles of inflammation. *Nature* **(2008)** 454:428–435.
doi: 10.1038/nature07201

2. Sharma D, Kanneganti T-D. The cell biology of inflammasomes: Mechanisms of
inflammasome activation and regulation. *Journal of Cell Biology* **(2016)** 213:617–629. doi:
10.1083/jcb.201602089

3. Tartey S, Kanneganti T. Differential role of the NLRP3 inflammasome in infection and
tumorigenesis. *Immunology* **(2019)** 156:329–338. doi: 10.1111/imm.13046

4. Akira S, Uematsu S, Takeuchi O. Pathogen Recognition and Innate Immunity. *Cell* **(2006)**
124:783–801. doi: 10.1016/j.cell.2006.02.015

5. Martinon F, Burns K, Tschopp J. The Inflammasome: a molecular platform triggering
activation of inflammatory caspases and processing of proIL-beta. *Mol Cell* **(2002)** 10:417–
426. doi: 10.1016/S1097-2765(02)00599-3

6. Shi J, Zhao Y, Wang K, Shi X, Wang Y, Huang H, Zhuang Y, Cai T, Wang F, Shao F.
Cleavage of GSDMD by inflammatory caspases determines pyroptotic cell death. *Nature*
**(2015)** 526:660–665. doi: 10.1038/nature15514

7. Kayagaki N, Kornfeld OS, Lee BL, Stowe IB, O'Rourke K, Li Q, Sandoval W, Yan D,
Kang J, Xu M, et al. NINJ1 mediates plasma membrane rupture during lytic cell death.
*Nature* **(2021)** 591:131–136. doi: 10.1038/s41586-021-03218-7

8. Losa M, Emmenegger M, De Rossi P, Schürch PM, Serdiuk T, Pengo N, Capron D, Bieli
D, Bargenda N, Rupp NJ, et al. The ASC inflammasome adapter governs SAA-derived
protein aggregation in inflammatory amyloidosis. *EMBO Mol Med* **(2024)** 16:2024–2042.
doi: 10.1038/s44321-024-00107-0

9. Baroja-Mazo A, Martín-Sánchez F, Gomez AI, Martínez CM, Amores-Iniesta J, Compan
V, Barberà-Cremades M, Yagüe J, Ruiz-Ortiz E, Antón J, et al. The NLRP3 inflammasome
is released as a particulate danger signal that amplifies the inflammatory response. *Nat*
*Immunol* **(2014)** 15:738–748. doi: 10.1038/ni.2919

10. Alarcón-Vila C, Hurtado-Navarro L, Mateo S V, Peñín-Franch A, Martínez CM, Molina-
López C, Baños MC, Gómez AI, Gómez-Román J, Baroja-Mazo A, et al. The
inflammasome adaptor protein ASC promotes amyloid deposition in cryopyrin-associated
periodic syndromes. *EMBO Mol Med* **(2024)** 17:41–53. doi: 10.1038/s44321-024-00176-1

11.     Bertheloot D, Wanderley CW, Schneider AH, Schiffelers LD, Wuerth JD, Tödtmann JM,

560           Maasewerd S, Hawwari I, Duthie F, Rohland C, et al. Nanobodies dismantle post-pyroptotic

561           ASC specks and counteract inflammation *in vivo*. *EMBO Mol Med* **(2022)** 14: doi:

562           10.15252/emmm.202115415

563     12.     Venegas C, Kumar S, Franklin BS, Dierkes T, Brinkschulte R, Tejera D, Vieira-Saecker A,

564           Schwartz S, Santarelli F, Kummer MP, et al. Microglia-derived ASC specks cross-seed

565           amyloid-β in Alzheimer's disease. *Nature* **(2017)** 552:355–361. doi: 10.1038/nature25158

566     13.     Franklin BS, Bossaller L, De Nardo D, Ratter JM, Stutz A, Engels G, Brenker C, Nordhoff

567           M, Mirandola SR, Al-Amoudi A, et al. The adaptor ASC has extracellular and "prionoid"

568           activities that propagate inflammation. *Nat Immunol* **(2014)** 15:727–737. doi:

569           10.1038/ni.2913

570     14.     Henry LP, Bruijning M, Forsberg SKG, Ayroles JF. The microbiome extends host

571           evolutionary potential. *Nat Commun* **(2021)** 12:5141. doi: 10.1038/s41467-021-25315-x

572     15.     Kudelka MR, Stowell SR, Cummings RD, Neish AS. Intestinal epithelial glycosylation in

573           homeostasis and gut microbiota interactions in IBD. *Nat Rev Gastroenterol Hepatol* **(2020)**

574           17:597–617. doi: 10.1038/s41575-020-0331-7

575     16.     Muller PA, Matheis F, Schneeberger M, Kerner Z, Jové V, Mucida D. Microbiota-

576           modulated CART+ enteric neurons autonomously regulate blood glucose. *Science* **(2020)**

577           370:314–321. doi: 10.1126/science.abd6176

578     17.     Seo S-U, Kamada N, Muñoz-Planillo R, Kim Y-G, Kim D, Koizumi Y, Hasegawa M,

579           Himpsl SD, Browne HP, Lawley TD, et al. Distinct Commensals Induce Interleukin-1β via

580           NLRP3 Inflammasome in Inflammatory Monocytes to Promote Intestinal Inflammation in

581           Response to Injury. *Immunity* **(2015)** 42:744–755. doi: 10.1016/j.immuni.2015.03.004

582     18.     Feng Y, Huang Y, Wang Y, Wang P, Song H, Wang F. Antibiotics induced intestinal tight

583           junction barrier dysfunction is associated with microbiota dysbiosis, activated NLRP3

584           inflammasome and autophagy. *PLoS One* **(2019)** 14:e0218384. doi:

585           10.1371/journal.pone.0218384

586     19.     Sokolova M, Yang K, Hansen SH, Louwe MC, Kummen M, Hov JER, Sjaastad I, Berge

587           RK, Halvorsen B, Aukrust P, et al. NLRP3 inflammasome deficiency attenuates metabolic

588           disturbances involving alterations in the gut microbial profile in mice exposed to high fat

589           diet. *Sci Rep* **(2020)** 10:21006. doi: 10.1038/s41598-020-76497-1

590     20.     Finucane OM, Lyons CL, Murphy AM, Reynolds CM, Klinger R, Healy NP, Cooke AA,
591             Coll RC, McAllan L, Nilaweera KN, et al. Monounsaturated Fatty Acid–Enriched High-Fat
592             Diets Impede Adipose NLRP3 Inflammasome–Mediated IL-1β Secretion and Insulin
593             Resistance Despite Obesity. *Diabetes* **(2015)** 64:2116–2128. doi: 10.2337/db14-1098

594     21.     Levy M, Thaiss CA, Zeevi D, Dohnalová L, Zilberman-Schapira G, Mahdi JA, David E,
595             Savidor A, Korem T, Herzig Y, et al. Microbiota-Modulated Metabolites Shape the
596             Intestinal Microenvironment by Regulating NLRP6 Inflammasome Signaling. *Cell* **(2015)**
597             163:1428–1443. doi: 10.1016/j.cell.2015.10.048

598     22.     Zhang J, Yu Q, Jiang D, Yu K, Yu W, Chi Z, Chen S, Li M, Yang D, Wang Z, et al.
599             Epithelial Gasdermin D shapes the host-microbial interface by driving mucus layer
600             formation. *Sci Immunol* **(2022)** 7: doi: 10.1126/sciimmunol.abk2092

601     23.     Miao P, Ruiqing T, Yanrong L, Zhuwen S, Huan Y, Qiong W, Yongnian L, Chao S.
602             Pyroptosis: A possible link between obesity-related inflammation and inflammatory
603             diseases. *J Cell Physiol* **(2022)** 237:1245–1265. doi: 10.1002/jcp.30627

604     24.     Opoku E, Traughber CA, Zhang D, Iacano AJ, Khan M, Han J, Smith JD, Gulshan K.
605             Gasdermin D Mediates Inflammation-Induced Defects in Reverse Cholesterol Transport
606             and Promotes Atherosclerosis. *Front Cell Dev Biol* **(2021)** 9: doi:
607             10.3389/fcell.2021.715211

608     25.     Xu B, Jiang M, Chu Y, Wang W, Chen D, Li X, Zhang Z, Zhang D, Fan D, Nie Y, et al.
609             Gasdermin D plays a key role as a pyroptosis executor of non-alcoholic steatohepatitis in
610             humans and mice. *J Hepatol* **(2018)** 68:773–782. doi: 10.1016/j.jhep.2017.11.040

611     26.     Ahmad F, Mishra N, Ahrenstorf G, Franklin BS, Latz E, Schmidt RE, Bossaller L. Evidence
612             of inflammasome activation and formation of monocyte-derived ASC specks in HIV-1
613             positive patients. *AIDS* **(2018)** 32:299–307. doi: 10.1097/QAD.0000000000001693

614     27.     Eckel RH, Grundy SM, Zimmet PZ. The metabolic syndrome. *The Lancet* **(2005)**
615             365:1415–1428. doi: 10.1016/S0140-6736(05)66378-7

616     28.     Hamblin PS, Russell AW, Talic S, Zoungas S. The growing range of complications of
617             diabetes mellitus. *Trends in Endocrinology & Metabolism* **(2025)** doi:
618             10.1016/j.tem.2024.12.006

619     29.     Oami T, Abtahi S, Shimazui T, Chen C-W, Sweat YY, Liang Z, Burd EM, Farris AB,
620             Roland JT, Tsukita S, et al. Claudin-2 upregulation enhances intestinal permeability,

immune activation, dysbiosis, and mortality in sepsis. *Proceedings of the National Academy of Sciences* **(2024)** 121: doi: 10.1073/pnas.2217877121

30. Honda M, Kadohisa M, Yoshii D, Komohara Y, Hibi T. Directly recruited GATA6 + peritoneal cavity macrophages contribute to the repair of intestinal serosal injury. *Nat Commun* **(2021)** 12:7294. doi: 10.1038/s41467-021-27614-9

31. Markovich Z, Abreu A, Sheng Y, Han SM, Xiao R. Deciphering internal and external factors influencing intestinal junctional complexes. *Gut Microbes* **(2024)** 16: doi: 10.1080/19490976.2024.2389320

32. Greer RL, Dong X, Moraes ACF, Zielke RA, Fernandes GR, Peremyslova E, Vasquez-Perez S, Schoenborn AA, Gomes EP, Pereira AC, et al. Akkermansia muciniphila mediates negative effects of IFNγ on glucose metabolism. *Nat Commun* **(2016)** 7:13329. doi: 10.1038/ncomms13329

33. Geach T. Mucin-munching bacteria modulate glucose metabolism. *Nat Rev Endocrinol* **(2017)** 13:66–66. doi: 10.1038/nrendo.2016.201

34. Matsumoto H, Shiotani A, Katsumata R, Fukushima S, Handa Y, Osawa M, Murao T, Handa O, Umegaki E, Inoue R, et al. Mucosa-Associated Microbiota in Patients with Irritable Bowel Syndrome: A Comparison of Subtypes. *Digestion* **(2021)** 102:49–56. doi: 10.1159/000512167

35. Brooks JF, Behrendt CL, Ruhn KA, Lee S, Raj P, Takahashi JS, Hooper L V. The microbiota coordinates diurnal rhythms in innate immunity with the circadian clock. *Cell* **(2021)** 184:4154-4167.e12. doi: 10.1016/j.cell.2021.07.001

36. Matsuzawa M, Ando T, Fukase S, Kimura M, Kume Y, Ide T, Izawa K, Kaitani A, Hara M, Nakamura E, et al. The protective role of conjunctival goblet cell mucin sialylation. *Nat Commun* **(2023)** 14:1417. doi: 10.1038/s41467-023-37101-y

37. Yu K, Li Q, Sun X, Peng X, Tang Q, Chu H, Zhou L, Wang B, Zhou Z, Deng X, et al. Bacterial indole-3-lactic acid affects epithelium–macrophage crosstalk to regulate intestinal homeostasis. *Proceedings of the National Academy of Sciences* **(2023)** 120: doi: 10.1073/pnas.2309032120

38. Tirado-Hurtado I, Fajardo W, Pinto JA. DNA Damage Inducible Transcript 4 Gene: The Switch of the Metabolism as Potential Target in Cancer. *Front Oncol* **(2018)** 8: doi: 10.3389/fonc.2018.00106

652    39.    Luo Z-H, Niu M-W, Gong S-H, Wu G-Y, Wang T, Wang F-Z, Wei G-Q, He Z-K, Jiang Y,
653           Chen P. Intestinal epithelial chemokine (C-C motif) ligand 7 overexpression protects
654           against high fat diet-induced obesity and hepatic steatosis in mice. *Chin Med J (Engl)* **(2020)**
655           133:1805–1814. doi: 10.1097/CM9.0000000000000915

656    40.    Merino B, Fernández-Díaz CM, Parrado-Fernández C, González-Casimiro CM, Postigo-
657           Casado T, Lobatón CD, Leissring MA, Cózar-Castellano I, Perdomo G. Hepatic insulin-
658           degrading enzyme regulates glucose and insulin homeostasis in diet-induced obese mice.
659           *Metabolism* **(2020)** 113:154352. doi: 10.1016/j.metabol.2020.154352

660    41.    Su Q, Huang J, Chen X, Wang Y, Shao M, Yan H, Chen C, Ren H, Zhang F, Ni Y, et al.
661           Long-Term High-Fat Diet Decreases Renal Insulin-Degrading Enzyme Expression and
662           Function by Inhibiting the PPARγ Pathway. *Mol Nutr Food Res* **(2023)** 67: doi:
663           10.1002/mnfr.202200589

664    42.    Bauer C, Duewell P, Mayer C, Lehr HA, Fitzgerald KA, Dauer M, Tschopp J, Endres S,
665           Latz E, Schnurr M. Colitis induced in mice with dextran sulfate sodium (DSS) is mediated
666           by the NLRP3 inflammasome. *Gut* **(2010)** 59:1192–1199. doi: 10.1136/gut.2009.197822

667    43.    Frühbeck G, Gómez-Ambrosi J, Ramírez B, Becerril S, Rodríguez A, Mentxaka A, Valentí
668           V, Moncada R, Reina G, Baixauli J, et al. Decreased expression of the NLRP6
669           inflammasome is associated with increased intestinal permeability and inflammation in
670           obesity with type 2 diabetes. *Cellular and Molecular Life Sciences* **(2024)** 81:77. doi:
671           10.1007/s00018-024-05124-3

672    44.    Tye H, Yu C-H, Simms LA, de Zoete MR, Kim ML, Zakrzewski M, Penington JS, Harapas
673           CR, Souza-Fonseca-Guimaraes F, Wockner LF, et al. NLRP1 restricts butyrate producing
674           commensals to exacerbate inflammatory bowel disease. *Nat Commun* **(2018)** 9:3728. doi:
675           10.1038/s41467-018-06125-0

676    45.    Mullins B, Chen J. NLRP9 in innate immunity and inflammation. *Immunology* **(2021)**
677           162:262–267. doi: 10.1111/imm.13290

678    46.    Patel S, Das A, Meshram P, Sharma A, Chowdhury A, Jariyal H, Datta A, Sarmah D, Nalla
679           LV, Sahu B, et al. Pyruvate kinase M2 in chronic inflammations: a potpourri of crucial
680           protein–protein interactions. *Cell Biol Toxicol* **(2021)** 37:653–678. doi: 10.1007/s10565-
681           021-09605-0

47.   Palsson-McDermott EM, Curtis AM, Goel G, Lauterbach MAR, Sheedy FJ, Gleeson LE,

van den Bosch MWM, Quinn SR, Domingo-Fernandez R, Johnston DGW, et al. Pyruvate

Kinase M2 Regulates Hif-1α Activity and IL-1β Induction and Is a Critical Determinant of

the Warburg Effect in LPS-Activated Macrophages. *Cell Metab* **(2015)** 21:347. doi:

10.1016/j.cmet.2015.01.017

48.   Sunny NE, Bril F, Cusi K. Mitochondrial Adaptation in Nonalcoholic Fatty Liver Disease:

Novel Mechanisms and Treatment Strategies. *Trends in Endocrinology & Metabolism*

**(2017)** 28:250–260. doi: 10.1016/j.tem.2016.11.006

49.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol* **(2014)** 15:550. doi: 10.1186/s13059-014-0550-

8

50.   Kolde R. pheatmap: Pretty Heatmaps. *CRAN: Contributed Packages* **(2019)** doi:

10.32614/CRAN.package.pheatmap

51.   Carlson M. Genome wide annotation for Mouse, primarily based on mapping using Entrez

Gene identifiers. *Bioconductor Package Maintainer* **(2023)**

52.   Wickham H. *ggplot2: Elegant Graphics for Data Analysis.* 2nd edition. Springer. **(2016).**

53.   McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis

and Graphics of Microbiome Census Data. *PLoS One* **(2013)** 8:e61217. doi:

10.1371/journal.pone.0061217

**Figure legends**

**Figure 1. ASC deficiency prevents gain of weight and glucose intolerance during high fat diet.**

**A.** Body weight from *Pycard*$^{+/+}$ (blue, n=4), *Pycard*$^{-/-}$ (green, n=2) mice fed with a normal chow diet (NCD).

**B.** Body weight from *Pycard*$^{+/+}$ (red, n=5), *Pycard*$^{-/-}$ (green, n=4), and *Pycard*$^{-/-}$ with intraperitoneal injection of recombinant ASC oligomers (+ipASC) (purple, n=5) mice fed with a high fat diet (HFD).

**C,D.** Glucose tolerance on mice at 3 months **(C)** and 5 months **(D)** of NCD or HFD.

Each dot represents a distinct animal, bars represent the mean, error bars represent SEM. Two-way ANOVA together multivariant Bonferroni's test was used for **A-D** (*$p \leq 0.05$; **$p \leq 0.01$; *** $p \leq 0.001$).

**Figure 2. ASC-deficiency reduces intestinal permeability and increases peritoneal cell infiltration in high-fat-diet.**

**A,B.** FITC-dextran concentration in plasma of *Pycard*$^{+/+}$ or *Pycard*$^{-/-}$ mice fed with a normal chow diet (NCD) or high fat diet (HFD) for 5 months; *Pycar*$^{-/-}$ mice with intraperitoneal injection of recombinant ASC oligomers (+ipASC).

**C,D**. Number of peritoneal cells of mice treated as in A,B.

**E,F.** Correlation between FITC-dextran plasma concentration and peritoneal cells of mice treated as in A,B.

**G,H.** IL-1β release from peritoneal cells collected from mice treated as in as in A,B and then *in vitro* stimulated as described in the legend.

*Pycard*$^{+/+}$ mice fed with NCD (n=4) in comparison with HFD (n= 5) **(A, C, E, G)**, and on HFD fed ASC$^{+/+}$ (n=4-8) in comparison with *Pycard*$^{-/-}$ mice with (n=5) or without ipASC (n=4) **(B, D, F, H)**.

Each dot represents a distinct animal, bars represent the mean, error bars represent SEM. *t*-student test was used for **A**. One-way ANOVA and multivariant Holm-Sidak´s test was done for **B**. Mann-Whitney's test was done for **C**. Kruskal Wallis and multivariant Dunn's test was used for **D**. Spearman's correlation test was used for **E, F**. Two-way ANOVA and multivariant Bonferroni test was used for **G, H** (ns, p>0,05; *$p \leq 0.05$; **$p \leq 0.01$; *** $p \leq 0.001$).

**Figure 3. Extracellular ASC oligomers modulates microbiota biodiversity during high-fat-diet.**

**A-D.** Relative microbiota abundance species **(A)** or class levels **(B-D)** on high fat diet (HFD) fed for 5 months *Pycard*$^{+/+}$ (n=5) **(A, B)**, *Pycard*$^{-/-}$ mice (n=5) **(A, C),** or *Pycard*$^{-/-}$ mice with intraperitoneal injection of recombinant ASC oligomers (ipASC) (n=4) **(A, D)**.

**E-H**. Relative abundance of *Akkermansia* **(E)**, *Bacteroides* **(F)**, *Enteractinococcus* **(G)**, and *Atopostipes* **(H)** in the same conditions as in A-D.

Each dot represents a distinct animal, bars represent the mean, error bars represent SEM. One-way ANOVA together multivariant Holm-Sidak's test was used for **E, F**. Kruskal-Wallis together multivariant Dunn's test was done for **G, H** (**p≤0.01).

**Figure 4. ASC induces bacteria attachment in small intestinal *villi* during high-fat-diet and alters epithelial integrity-related genes and increase innate immune response.**

**A,B,C.** Hematoxilin and eosin histology of the small intestine of mice fed with a high fat diet (HFD) for 5 months, *Pycard*$^{+/+}$ (**A**), *Pycard*$^{-/-}$ mice (**B**), or *Pycard*$^{-/-}$ mice with intraperitoneal injection of recombinant ASC oligomers (ipASC) **(C)**. Scale bar = 200 μm.

**D**, Differential expressed genes in the small intestine of mice treated as in A-C.

**E**, Gene Ontology differential expressed pathways in the small intestine of mice treated as in A,B.

**F,** Interleukin (IL)-1β detection in the small intestine of mice treated as in A-C

*Pycard*$^{+/+}$ (n=3-5), *Pycard*$^{-/-}$ mice (n=3), *Pycard*$^{-/-}$ mice with intraperitoneal injection of recombinant ASC oligomers (ipASC) (n=5).

Black arrows indicate presence of bacteria in **(A)**, blue arrows indicate presence of mucus in **(B)**, red arrows indicate absence of mucus in **(C)**.

Each dot represents a distinct animal, bars represent the mean, error bars represent SEM. One Way ANOVA together multivariant Holm-Sidak's test was used for **F.** (*p≤0.05).

**Figure 5. ASC deficiency increase glycolytic and tricarboxylic acids related genes in high-fat-diet.**

**A-F.** Relative mRNA expression of *Pkm2* **(A)**, *Ldha* **(B)**, *Pdha* **(C)**, *Glud1* **(D)**, *Sdha* **(E)**, and *Cpt1a* **(F)** on the small intestine of mice fed with a HFD.

*Pycard*^+/+^ (n=5), *Pycard*^-/-^ mice (n=4) or *Pycard*^-/-^ mice with intraperitoneal injection of

recombinant ASC oligomers (ipASC) (n=5).

Bars represent the mean, error bars represent SEM. Kruskal Wallis together multivariant Dunn's

test was used for all panels (*$p \leq 0.05$; **$p \leq 0.01$).

**Figure 6. ASC deficiency enhance hepatic metabolism under high-fat-diet and reduce**

**steatosis.**

**A-C.** Haematoxylin and eosin histology of the liver of mice fed with a high fat diet (HFD) for 5

months, *Pycard*^+/+^ (**A**), *Pycard*^-/-^ mice (**B**), or *Pycard*^-/-^ mice with intraperitoneal injection of

recombinant ASC oligomers (ipASC) (**C**). Scale bar = 200 μm.

**D-H.** Relative mRNA expression of *Pgc1a* (**D**)*, Nnt* (**E**)*, Glud1* (**F**)*, Cyp7a1* (**G**)*,* and *Shp* (**H**) in

the liver of mice treated as in C-D.

**I.** Interleukin (IL)-1β production in the liver of mice treated as in C-D.

HFD fed *Pycard*^+/+^ (n=5), *Pycard*^-/-^ mice (n=4), *Pycard*^-/-^ mice with intraperitoneal injection of

recombinant ASC oligomers (ipASC) (n=5).

Each dot represents a distinct animal, bars represent the mean, error bars represent SEM. One way

ANOVA together multivariant Holm-Sidak's test was used for (**D-I**) (*$p \leq 0.05$; **$p \leq 0.01$; ***

$p \leq 0.001$).

**Supplemental information**

**Supplemental Figure 1. Body weight of different mice genotypes.**

**A.** Body weight on normal chow diet (NCD) fed *Pycard*$^{+/+}$ (blue, n=3), and *Pycard*$^{-/-}$ (green, n=2), male mice, and on NCD fed wild type (blue, n=4), and *Pycard*$^{-/-}$ (green, n=2), female mice.

**B-C**. Body weight on *Pycard*$^{+/+}$ mice fed with high fat diet (HFD) (n=3) in comparison with normal chow diet fed (NCD) mice (n=4) **(B)**, and on HFD fed wild type in comparison with *Gsdmd*$^{-/-}$ **(C)**. Values represent the mean of each condition and error bars represent SEM. Two-way ANOVA together multivariant Bonferroni's test was used for all panels (*p≤0.05; **p≤0.01; *** p≤0.001).

**Supplemental Figure 2. Cytokines produced from peritoneal cells and intestinal permeability in *Pycard*$^{-/-}$ mice in comparison with wild type mice.**

**A.** IL-1β release from peritoneal cells treated *in vitro* with the treatments indicated in the legend. Peritoneal cells were isolated from normal chow diet (NCD) fed *Pycard*$^{-/-}$ (n=4) and *Pycard*$^{+/+}$ (n=4) mice.

**B,C.** IL-6 production on mice fed with high fat diet (HFD) for 5 months, *Pycard*$^{+/+}$ (n=5), *Pycard*$^{-/-}$ mice (n=4), *Pycard*$^{-/-}$ mice with intraperitoneal injection of recombinant ASC oligomers (ipASC) (n=5) **(B)** and on wild type mice fed with NCD or HFD **(C)**..

Each dot represents a distinct animal, bars represent the mean, error bars represent SEM. Kruskal-Wallis together multivariant Dunn's test was used for **A-C**. (*p≤0.05; **p≤0.01; *** p≤0.001).

**Supplemental Figure 3. Analysis of the intestinal microbiota of mice fed with normal chow diet or high fat diet.**

**A-C.** Relative abundance of the total faecal microbiota at class level **(A, B)**, and of *Akkermansia* **(C)** on *Pycard*$^{+/+}$ mice fed with a normal chow diet (NCD) (n=4) in comparison with high fat diet (HFD) (n= 5).

Each dot represents a distinct animal, bars represent the mean, error bars represent SEM. *t*-student test was used for **C**.

**Supplemental Figure 4. Histological analysis of the small intestine from mice fed with normal**

**chow diet or with high fat diet.**

**A-B.** Haematoxylin and eosin histology of the small intestine of mice fed with a normal chow diet

(NCD) representative of n=4 **(A)**, or a high fat diet (HFD) representative of n= 5 **(B)** for 5 months.

Scale bar = 200 μm.

**C-D.** Interleukin (IL)-6 production **(C),** mRNA relative expression of *Pycard* **(D),** and principal

components analysis **(E)** in the small intestine of mice fed with a HFD, *Pycard*$^{+/+}$ (n=4), *Pycard*$^{-/-}$

mice (n=4), *Pycard*$^{-/-}$ mice with intraperitoneal injection of recombinant ASC oligomers (ipASC)

(n=5).

Each dot represents a distinct animal, bars represent the mean, error bars represent SEM. One way

ANOVA together multivariant Holm-Sidak's test was used for **C**. Kruskal-Wallis together

multivariant Dunn's test was used for **D** (*** p≤0.001).

**Supplemental Figure 5. Histological analysis of the liver from mice fed with normal chow**

**diet compared with high fat diet.**

**A, B.** Haematoxylin and eosin histology of the liver of *Pycard*$^{+/+}$mice fed with a normal chow diet

(NCD) representative of n=4 **(A)**, or a high fat diet (HFD) representative of n= 5 for 5 months **(B)**.

Scale bar = 200 μm.

**Figure 1**

# Figure 2

# Figure 3

# Figure 4

**A** *Pycard*+/+HFD

**B** *Pycard*-/-HFD

**C** *Pycard*-/-HFD+ipASC

**D**

*Pycard*-/- HFD    *Pycard*-/-HFD+ipASC    *Pycard*+/+HFD    condition

Gdpd3
Ide
Spink1
Nts
Prr13
Chordc1
Mertk
Smpdl3b
Slc35b1
Ksr1
Pmaip1
Erbb3
Slc26a6
Sertad1
Npas2
Vps4b
Atp6v0e
Tm4sf20
Ccl7
Ern2
Slx1b
Ddit4
Fkbp5

**E**

*Pycard*-/- HFD

Inhibitory MHC Class I receptor activity

Regulation of T cell costimulation

Regulation of antigen processing and presentation of peptide or polysaccharide antigen via MHC II

MHC Class I activity

*Pycard*+/+HFD

AIM2 Inflammasome complex

Positive regulation of humoral immune response

Hormone activity

NLRP3 inflammasome complex

p.adjust
0.002
0.004
0.006
0.008

Count
○ 10
○ 20

GeneRatio

**F**

IL1β (pg/mg total protein)

* *

*Pycard*+/+HFD    *Pycard*-/-HFD    *Pycard*-/-HFD+ipASC

# Figure 5

# Figure 6

# Supplementary Figure 1

**A**

**B**

**A**

**B**

**C**

# Supplementary Figure 3

**A**

*Pycard*<sup>+/+</sup>NCD

**B**

*Pycard*<sup>+/+</sup>HFD

**C** **Akkermansia**

# Supplementary Figure 4

# Supplementary Figure 5

**A** *Pycard*+/+ NCD

**B** *Pycard*+/+ HFD