The evolution of mutation rates in the light of development and cell-lineage selection

Paco Majic[1,*], Malvika Srivastava[2] & Justin Crocker[1]

[1]European Molecular Biology Laboratory, 69117 Heidelberg, Germany

[2]Department of Ecology and Evolution, University of Lausanne, 1015 Lausanne, Switzerland

*Paco Majic

Email: paco.majic@embl.de

Abstract

Mutation rates drive the pace and potential of evolutionary change. However, to better understand the evolutionary implications of mutation rates, there is a need to uncover the causes of their diversification. In multicellular organisms, all mutations first arise in a single cell in a developmental context. Whether a mutation enters a population's gene pool can therefore depend on developmental events that affect the likelihood of mutant cell lineages of producing gametes. For this reason, the evolution of mutation rates in populations is governed not only by changes in the rates at which mutations occur at the molecular level, but also by changes in developmental features of organisms and how mutations impact cellular fitness during development. We present a theoretical framework that, supported by empirical data from mammals and experiments on the fruit fly, demonstrate how generational mutation rates can be shaped by changes in developmental parameters and intraorganismal selective processes even when molecular mutation rates are presumed constant. Our model highlights how the diversity of mutation rates observed across animals may be the byproduct of organismal development rather than the result of direct selection against mutator alleles. As such, development not only introduces phenotypic biases, it also shapes the rates and trajectories of genetic diversity and is thus at the core of evolutionary theory.

Main Text

"Seen from this point of view, life appears as a current which goes from

germ to germ through the intermediation of a developed organism"

- Henri Bergson, *Creative Evolution*

Introduction

How genetic variation arises within a population shapes evolutionary trajectories (1-8). The rates of mutations in a population may affect how promptly adaptive genotypes may be encountered, while also defining how deleterious mutations may burden populations. Thus, mutation rates are closely tied to the pace and potential of evolutionary change. Although classical models presumed substitution and mutation rates to be relatively constant across the tree of life (9, 10), advances in molecular biology have progressively challenged this view (11-14). The depth and granularity at which we can now explore genetic diversity suggest that mutation rates are an evolvable trait, varying not only between clades (15-17), but also within species (18) and populations (19), underscoring their dynamic role in shaping genetic diversity.

The diversification of mutation rates can be shaped by the evolution of mutator alleles, defined as "any genomically encoded modifier of the mutation rate" (20). These alleles are typically considered to operate at the molecular scale, altering the balance between DNA damage and repair through variations in genes involved in these processes (14, 17, 20-23). Mutator alleles that promote hypermutation may be favored under strong adaptive pressures (19, 24), while those causing excessive deleterious mutations would be selected against (25, 26). Because most mutations tend to be deleterious, the mutation rates of species are thought to be dominated by the efficacy of selection in purging alleles that increase the rates of mutations (26). This is why the drift-barrier hypothesis posits that the efficiency of selection against mutator alleles depends on population size. Larger populations can better counteract genetic drift, resulting in lower mutation rates, whereas smaller populations experience weaker selection against mutators and, thus, higher mutation rates (20, 26, 27). This view suggests that mutation rates evolve primarily through changes in molecular mechanisms constrained by selection. However, mutation rates

may also evolve through non-molecular causes. A relevant factor in this regard is generation time

and age at reproduction, which influence the number of mutations germlines accumulate before

reproduction (28-31).

Another non-molecular factor that may influence mutation rate evolution is that, in multicellular

organisms, new mutations necessarily arise in a single cell in the organismal context of each

parent. For a mutation to enter the population's gene pool, it must propagate within the organism

and be part of the germline, meaning developmental traits have the potential to shape the

likelihood of heritable mutations. For instance, the timing of germline determination (32), and

differences in gamete production between sexes or species can influence mutation rates (16, 30,

31, 33). Moreover, mutations may result in differences in cellular fitness, which can greatly

influence the genetic diversity existing within organisms. In the germline context, mutations that

affect the fitness of cells relative to their sisters may bias their own heritability if they occur on

the lineage of cells that go from zygote to gamete (34-41). Such bias can have important

evolutionary repercussions (40): it may affect the probability of fixation of variants that alter cell

fitness, it allows for a "cheap" purging of deleterious variants, and it can also promote evolvability

when the impact of mutations on cellular fitness and on individual fitness align. Nevertheless, cell-

lineage selection tends to be overlooked in discussions of mutation rate evolution (17, 22, 30),

and it is generally unaccounted for in mutation rate estimations, since whenever selection is

considered there is a tendency to focus exclusively at the level of individuals - a perspective which

biases causal explanations of mutation rate variation (36).

In this work, we emphasize the view that differences in population-level mutation rates across

species may result from causes other than molecular changes mediated by mutator alleles. Using

analytical models, data analysis and experiments in *Drosophila,* our work consolidates a view of

mutation rate evolution, which demonstrates how the interplay of developmental and life-history

parameters with intraorganismal selection on mutant cells can majorly affect the rate of new

mutations in a population. This perspective has important conceptual implications for

understanding the evolution of mutation rates. Unlike explanations like the drift-barrier

hypothesis, our model highlights how developmental processes may influence generational

mutation rates, positioning them as a byproduct of organismal development rather than direct

selection at the level of individuals in a population. Furthermore, this model broadens the role of

organismal development in population genetics: development not only introduces phenotypic

biases, but it also shapes the rates and trajectories of genetic diversity, which places development

at the core of evolutionary theory.

Results

<u>A theoretical framework to assess the influence of developmental features and cell-lineage</u>

<u>selection on mutation rate diversification</u>

To distinguish between the molecular mutation rate, μ_{mol}, and the mutation rate per generation

at the population level, μ_{pop}, we revisited the theoretical framework of Otto & Orive (35), who

explored how the mutation load of a population is affected by intraorganismal selection within

individuals (see Methods). Conceptually, development is modeled as a branching tree of cell

lineages, with μ_{pop} partly determined by the number of cells carrying a mutation at locus i at the

end of development, M_i (Figure 1A). M_i is a function of:

i)   a molecular variable μ_{mol}, which is the mutation probability per cell per locus per unit of

developmental tempo, which can be either per cell division when mutations are of

replicative nature (35), or per time unit when they result from other sources of DNA

lesions $(29, 42)$,

ii)   developmental parameters such as the duration of development (τ), and the rate of cell

proliferation (C), and

iii)   the effect of mutations at locus i on cell fitness (β_i), which alters the rate of cell

proliferation of mutant cells relative to nonmutant cells in the developmental context. β_i

< 1 indicates deleterious effects that decrease cell proliferation, β_i = 1 neutrality, and β_i

> 1 beneficial effects that increase proliferation.

With an estimation of M_i and assuming that germline determination is independent of mutational

events, it is then possible to calculate μ_{pop} at locus i as the probability of sampling a mutant cell

among the total pool of cells, which includes all M_i mutated cells as well as all cells that completed

development without mutations, N_i. Such that:

111
$$\mu_{pop_i} = \frac{M_i}{M_i + N_i}$$

By defining μ_{pop} as the probability of randomly sampling mutant cells at the end of development,

we are implying that this probability also encompasses the probability that sampled cells are of

the germline lineage that goes from the zygote to differentiated gametes. This captures aspects

of development such as how body size, which scales with the total number of cells, generally

reflects germline size (43). However, our model does not consider many nuances of actual

development. Development is highly structured with different cell lineages proliferating and

differentiating at different tempos. Furthermore, overproliferation of cells by means of increases

in β_i could result in tumorigenesis, which would affect μ_{pop} through deleterious effects on

individual fitness. However, more often than not, mutant alleles spread selectively within healthy

tissues without causing tumors or deleterious effects on organismal fitness (44-50). This is not

only true for somatic tissues, but also for the germline (41). Our formalization is therefore

sufficient to assess how fundamental developmental features in combination with cell-lineage

selection can influence the rates at which mutations enter a population's gene pool, regardless of

their effect on organismal fitness and even if the molecular mutation rate μ_{mol} is constant.

<u>Developmental parameters can influence the diversification of mutation rates</u>

As per our model, under neutral conditions (β_i = 1), evolutionary lineages that increase τ, also

increase the expected μ_{pop} for locus i (Fig. 1B). This is also the case for shifts in C, but only if μ_{mol}

is considered to track cell divisions (Fig. 1B). When considering μ_{mol} to track mutations per unit

time, μ_{pop} also increases as a function of τ, but it is only affected by changes in C only if changes

in the rate of cell proliferation also result in changes in τ (Fig. S1A). To evaluate whether these

developmental parameters may explain interspecific differences in μ_{pop}, we analyzed estimates of

μ_{pop} (16) for a number of amniote species. Focusing on the mammalian clade, which has the most

comprehensive phylogenetic sampling (Fig. 1C, Table S1) and developmental information (51), we

focused on gestation time as a proxy for τ, as well as on the body mass as a proxy for total cell

count, proportional to $2^{C\tau}$ (52). As predicted by our model, the μ_{pop} of mammals positively

correlated with the proxy τ (Fig. 1D, Pearson's $\rho = 0.49$, $p = 0.0025$), as well as $2^{C\tau}$ (Fig. S1B,

Pearson's $\rho = 0.48$, $p = 0.0029$).

To more directly address the question of whether developmental parameters could account for

the interspecific diversification of μ_{pop}, we compared these variables between sister clades in our

phylogeny by calculating the ratio of μ_{pop} and body mass (Methods). We found that the majority

of the comparisons between relatively recently diverged species (Fig 1C, black bars) yielded a

correlation such that the larger of both species tended to have higher μ_{pop} (Fig. 1E, black dots). In

79% of recently diverged species pairs the larger of both species also had a higher μ_{pop}, which was

significantly different from random pairs (Fig. S1C, p = 0.01, one-sample permutation test). For

example, the estimated μ_{pop} of domestic cats (*Felis catus*) is ~2 times lower than that of tigers

(*Panthera tigris*) and ~2.5 times lower than that of leopards (*Panthera pardus*), while the μ_{pop} of

walruses (*Odobenus rosmarus*) is ~1.95 times that of the much smaller Antarctic fur seals

(*Arctocephalus gazella*)*. Interestingly, this held not only for mammals but also for birds, even if

the phylogenetic granularity of estimated μ_{pop} is lower (Fig. S2, in 74% of comparisons of related

bird species pairs the larger of both species also had a higher μ_{pop}, p = 0.02, one-sample

permutation test). Therefore, as lineages diverge, we expect that if either lineage tends to

increase in size, its mutation rate will also tend to increase. We do note that body size is also

correlated with generation time, and that the correlation between generation time and μ_{pop} (16),

as well as the influence of parental age at reproduction (53), may bias our interpretation. But, if

developmental parameters affecting body mass or other correlated life history traits are indeed

affecting μ_{pop}, then the diversification of developmental parameters may offer a non-molecular

explanation for observed diversification patterns of mutation rates.

As mentioned above, according to the drift-barrier hypothesis, the negative correlation existing

between the generational mutation rates of species and their effective population sizes could be

due to the efficacy of selection against alleles affecting μ_{mol} (20, 26). However, that relationship

could also be the neutral byproduct of changes in developmental parameters affecting body size,

rather than the selective purging of mutator alleles. This is because mammals follow a global

scaling relationship between body size and population density described by Damuth's rule (54),

as well as with the effective population size (17). Considering that body mass is proportional to

the total number of cells in a body, then τ and C not only affect μ_{pop}, but also the demographic

properties of a population. Therefore, simply considering the predicted effective population size

for different values of $C\tau$ based on this proportionality produces an inverse relationship between

the mutation rate of a species and its effective population size (Fig. 1F), as would also be expected

from the drift-barrier hypothesis. Thus, the differences in mutation rates observed in populations

of different sizes may be the result of changes in developmental parameters that, because of

ecological and energetic reasons, may also relate to changes in population sizes (55).

<u>The embryonic selective environment and its effect on mutation rates</u>

So far, we have been considering the case in which mutations are selectively neutral at the cellular

level. In other words, we have been considering the case where β_i = 1. However, if we take into

account how mutations may impact their own heritability by affecting cellular fitness, a selective

regime in the embryo or body of an organism can then also affect μ_{pop} without changes in μ_{mol}

(Fig. 2A). Cellular fitness could be affected by various processes, including the migration of cells,

proliferation, and, in the specific case of germline cells, the determination and maintenance of

the germline cell fate. Mutations on genes involved in these processes may end up under- or over-

represented in the gamete pool produced by an organism (41, 56).

To explore how mutations affecting β_i as in our model may impact their own heritability, we

considered the *Drosophila melanogaster* gene *trithorax-related (trr)*, which influences cell

proliferation (57). We generated mutant clones by inducing mitotic recombination through the

FLP/FRT technique (58) on flies that carried both a sub-functional allele of *trr* (TrrCA,(59)) on an

unmarked chromosome and a wild-type version of the gene (TrrWT) on a chromosome marked

with a *red fluorescent protein, mRFP* (see Methods). As a result, within a single embryo, some cells

are homozygotic for TrrCA, homozygotic for TrrWT, or heterozygotic with a copy of each allele

(Fig. 2B). Because an impaired *trr* activity increases the rates of cell proliferation (57), we then

expected that $\beta_{trrCA/trrCA} > \beta_{trrWT/trrCA} > \beta_{trrWT/trrWT}$ and, therefore, that the TrrCA allele would be

overrepresented in the germline and thus have a higher heritability than TrrWT. Consistent with

this hypothesis, when mosaicism was induced early in development, a median of ~55% of

offspring carried the mutant allele, a fraction that was significantly higher than the cases when

mosaicism was not induced on the TrrCA/TrrWT heterozygotes or when it was induced in the lines

with two copies of TrrWT (Fig. 2C). This experiment illustrates how a mutation inducing a greater

proliferation of cells that carry it can increase its own heritability, representing a case of what

Otto and Hastings referred to as "mitotic drive" (40).

In our model, varying the proliferative effect β_i of a mutation on a cell can drastically change the

effective μ_{pop} even when molecular mutation rates are constant, such that higher values of β_i

imply higher μ_{pop} (Fig. 2D). A relevant insight from this model is that even mild consequences for

cellular proliferation may have major consequences for the rate at which new mutations enter

the gene pool of populations. For example, in the case of $C\tau = 20$ (representing 2^{20} cells, roughly

the number of ovarian follicles an infant is born with (60) or half of the number of germ cells in

the ovary of cows by the third trimester of pregnancy (61)), a mere 1% increase ($\beta_I = 1.01$) or

decrease ($\beta_i = 0.99$) in cell fitness may result in a ~7% change in μ_{pop} (Fig. 2D).

The effect of β_i on μ_{pop} also depends on the developmental parameters C and τ (Fig. 2D). Higher

values of $C\tau$ tend to produce larger differences in μ_{pop} for large β_i . On the other hand, when β_i is

small, $C\tau$ does not have a strong effect on the mutation rate. We note that when $\beta_i \neq 1$, changes

in C and τ, also affect μ_{pop} when μ_{mol} is considered as mutations per unit of time, rather than per

cell division (Fig. S3A). This contrasts with the aforementioned neutral case, where changes in C

did not change μ_{pop} for constant τ (Fig. S1A).

Our model, therefore, suggests that: i) the selective consequences of mutations at the cellular

level may influence generational mutation rates, and ii) the magnitude of this influence depends

on developmental parameters. Specifically, our model predicts that organisms of larger size and

slower development should be more efficient in purging mutations that are deleterious for cells

and more prone to facilitate the fixation of mutations that are beneficial to cells (Fig. 2D). This

shows how differences in mutation rates between species may be enhanced by developmental

differences, and it highlights how the relevance of cell-lineage selection as a "cheap" way to filter

deleterious mutations (40) may be evolve by shifting developmental parameters.

<u>The interplay between cellular distribution of fitness effects and developmental parameters</u>

The effect of mutations on cellular fitness is expected to vary across genes, depending on their

involvement on cellular processes, and even across loci within genes, depending on their

relevance for gene function. For example, in our *Drosophila* case study, despite carrying a

mutation in the gene *white* (*w*) (Methods), the TrrWT allele did not lead to differences in

heritability in our FLP/FRT experiments (Fig. 2C). This suggest that even though mutations on *trr*

can influence β_i, mutations on *w* do not, which implies that *trr* and *w* likely contribute differentially

to μ_{pop}.

To explore how differences in β_i at different loci affects genome-wide mutation rate estimations,

we adjusted our developmental model such that the overall genomic mutation rate would be

proportional to the fraction of sites that have either deleterious ($F_{\beta < 1}$), neutral ($F_{\beta = 1}$) or

advantageous ($F_{\beta > 1}$) fitness effects on cells (Methods). Moving from a case in which mutations

on all sites are neutral for cellular fitness ($F_{\beta = 1} = 1,$), to a case in which all mutations are strongly

deleterious ($F_{\beta = 0} = 1$), there is a steady drop in the expected μ_{pop}, tending towards μ_{mol} as most

mutations become deleterious (Fig. 2E). Considering cases in which most mutations are either

neutral or deleterious for cellular fitness would be in accordance with what is expected (25). In

contrast, scenarios in which many mutations tend to increase cellular fitness are presumably rare.

However, our model does suggest that even if infrequent, mutations with strong beneficial effects

on the fitness of cells can drastically influence the genome-wide estimations of μ_{pop}. Depending

on the developmental parameters $C\tau$, this is true if mutations that increase cellular fitness are 1

in 1000 ($F_{\beta > 1} = 0.001$), or even 1 in a million ($F_{\beta > 1} = 1x10^{-6}$) (Fig.2F). Notably, variations in

the ratios of $F_{\beta = 1}$, $F_{\beta > 1}$ and $F_{\beta < 1}$ also vary the relationship between μ_{pop} and developmental

parameters (Fig. 1E,F).

Furthermore, because β_i affects mutated cells by altering their proliferative capacity, the number

of genome-wide mutations capable of increasing or decreasing cellular fitness may depend on the

rate of cell proliferation of wild-type cells. In other words, $F_{\beta = 1}$, $F_{\beta > 1}$ and $F_{\beta < 1}$ may depend on

baseline *C,* such that for greater *C* there should be a smaller number of mutations for which $\beta_i >$

1. Such would represent a case of diminishing returns epistasis, according to which the fitness

effect of a mutation is milder when it occurs on a fit genotype than when it occurs on an unfit

genotype (62-64). Epistatic effects of this kind could explain why there is a negative correlation

between our estimates of *C* for a subset of mammalian species (Methods) and μ_{pop} (Fig. S3B,

Pearson's $\rho = -0.38, p = 0.044$), since greater $F_{\beta > 1}$ imply greater μ_{pop} (Fig. 2F). This would mean

that whenever species evolve a greater size through increases in *C*, the cellular distribution of

fitness effect could change as well, thus affecting μ_{pop}.

Therefore, the genome-wide distribution of cellular fitness effects of mutations interacts with and

may be defined by the developmental background of an organism to jointly influence μ_{pop}. A

factor that needs consideration in any interpretation of differences in mutation rates between

species using genomic data.

<u>Selection in the mature germline stem cell niche</u>

In addition to embryonic development, the lifetime of organisms could also shape μ_{pop}. This could

take place, as mentioned earlier, because longer generations imply a greater accumulation of new

mutations before reproduction (30, 65, 66). Here, we argue that this could also be because of the

dynamic competition between germline stem cells of different genotypes. During an individual's

lifetime, germline stem cell niches can undergo a cellular turnover. In mice, for example,

spermatogenic stem cells sometimes cease to be stem cells and are replaced by neighboring stem

cells, keeping the number of stem cells relatively constant (67). Although the repopulation of

these cleared niches by neighboring cells could be a stochastic process (68), mutations that affect

the rate of stemness loss (either through death or differentiation) might eventually create a

mosaic of spermatic stem cells with an overall higher fitness (41, 56). Thus, a selective cell

turnover might impact the repertoire of mutations that a single individual can introduce into a

population's gene pool.

To incorporate the effect of selection during the lifespan of an organism, we built on our

developmental model described above and included a maturation and reproductive phase (Fig.

3A, Methods). In this adapted version, at the end of development, there is a final and constant

pool of stem cells that can produce gametes, $M_{t=0}$ of which are mutated. These cells can undergo

a turnover, either because of a death-birth dynamic in the stem cell niche or because of

differentiation. This turnover of cells in the stem cell niche occurs at a rate d over an arbitrary unit

of time t subdividing the total generational time l, and a new selective variable β_m which describes

the fitness effect of mutation on cells related to stem cell turnover during the lifetime of an

organism. In this model, in the case when there is no turnover of cells during the lifetime of an

individual (d = 0) or when mutations are neutral ($\beta_m = 1$), μ_{pop} only changes due to the

mutational accumulation of new mutations. Whenever there is a turnover of cells and mutations

do affect the proliferative capacities of cells, a positive effect of mutations on the fitness of cells

increases the overall μ_{pop} while deleterious mutations decrease it closer to μ_{mol} (Fig. 3B). This

effect is more pronounced for individuals that take longer to reproduce (Fig. 3C).

The intraorganismal selective process over the lifespan of a mature organism may interplay with

the intraorganismal selective process taking place earlier during development to shape an

effective μ_{pop} (Fig. 3D). For example, mutations on genes involved in basic cellular functions

would likely have fitness consequences for cells at any stage of an individual's life, thus affecting

cell proliferation during development, but also the rates of germline stem cell turnover ($\beta > 1$, β_m

> 1). On the other hand, there may be genes that are irrelevant for cellular fitness in the germline

stem cell niche that may be important during development ($\beta \neq 1$, $\beta_m = 1$) and vice versa ($\beta = 1$,

$\beta_m \neq 1$). The latter would be the case of mutations on the *chimno* mutants in *Drosophila*, which

have been reported to evict wild-type cells from the stem cell niche of the testis (38). It is also

possible that the selective regime during development and maturity point in opposite directions,

which is what we observed in the case of the TrrCA mutation presented above: even though

mosaic females introduced in Fig. 2C produced a progeny that was biased in favor of the *trr*

mutant allele right after pupation, as those same females aged, the biased shifted in favor of

TrrWT (Fig. 3E), suggesting that for this allele $\beta > 1$ and $\beta_m < 1$. On the other hand, as expected,

TrrWT retained a $\beta = \beta_m = 1$ through the development and lifespan of each female fly (Fig. 3F).

These results reinforce the notion that the estimation of global genome-wide mutation rates may

lead to biased interpretations of the molecular rates because of the fitness consequences of

mutations on different loci. More generally, our results also support that the overall expected

μ_{pop} is likely the composite outcome of the selective effect of mutations at the cellular level

during development and maturity, as well as the life history traits of the organism.

Discussion

Mutations arise in organismal contexts that vary across individuals and species. As our results

demonstrate, these contexts significantly influence how genetic variation is produced and

incorporated into gene pools. Therefore, differences in population-level mutation rates may not

solely result from changes in molecular mutation rates or changes in generation time, but could

arise from evolutionary shifts in developmental parameters. These include the rate of cell

proliferation, developmental duration, germline differentiation modes, germline dynamics during

maturity, and—crucially—selective dynamics at the cellular level.

This perspective implies that it is not essential to invoke direct selection on mutation rates to

account for the observed diversity of mutation rates across multicellular lineages. This is not to

say that selection at the level of individuals plays no role in modulating mutation rates. Molecular

rates might indeed evolve, for example, to mitigate the burden of deleterious mutations,

particularly somatic mutations that can cause cancer in the case of larger and longer-lived species

(65, 69-71). However, because developmental parameters alone can produce substantial shifts in

mutation rates, as species evolve different morphologies and life history strategies, their rates of

genetic diversification can also diverge as a secondary outcome that may or may not be neutral.

Our results illustrate how the diversification of mutation rates can arise as a consequence of

changes in developmental parameters associated with body size evolution. This is supported both

by our theoretical model and by a general pattern in which the larger of two closely related

species tends to exhibit a higher generational mutation rate. However, this trend is not universal

and can be influenced- or even reversed- by other factors. Although body size typically correlates

with germline size (43), deviations from this relationship, such as disproportionate germline cell

numbers or developmental heterochronies in the gonads, can alter expected generational

mutation rates, as suggested to explain higher rates in chimpanzees relative to humans (72).

Additionally, developmental changes tied to body size may interact in complex ways with

intraorganismal selection. For instance, in the leopard-tiger and goat-dwarf musk deer species

pairs (Fig. 1C), the larger species shows a lower estimated generational mutation rate (red

arrowheads in Fig. 1E, Table S1). We speculate that this misalignment reflects a smaller fraction

of mutations capable of further increasing cellular proliferation ($F_{\beta > 1}$) in species with an

inherently higher proliferative capacity of wild-type cells. We infer tigers and goats to have a

higher rate of cell proliferation per unit time (greater C) than leopards and dwarf musk deer,

respectively. This is evidenced by the fact that tigers, despite reaching larger birth sizes than

leopards, have similar developmental times, while dwarf musk deer, though significantly smaller

than goats, possess markedly longer developmental times (Table S1).

The biological causes of mutation rate diversification

The diversification of developmental parameters not only provides a basis for hypotheses about

species-specific shifts in shifts in mutation rates, but also offers potential explanations for well-

established patterns of mutation rate diversification. Our results suggest that the prevalent

negative correlation between generational mutation rates and population size (16, 17, 20, 26), is

more parsimoniously explained by developmental variations accompanying lineage divergence

than by recurrent selective adjustments of molecular mutation rates in response to demographic

changes, as proposed by the drift barrier hypothesis (26). Given that ecological drivers may

influence body size evolution (55) and that developmental parameters that increase body size

also increase generational mutation rates, the co-adjustment of population size and

developmental properties (or correlated life history traits) could underlie the diversification of

generational mutation rates. Although the drift barrier hypothesis is supported by various studies

that show a correlation between mutation rates and effective population size across clades (17),

we are here suggesting that, at least for multicellular organisms, there are other biological

properties that may explain the same trend, or at least reinforce it. This is consistent with our

findings and supported by evidence that diversification in mutation spectra, often attributed to

molecular changes, primarily occurs between distantly related lineages (73) and may also be

influenced by environmental factors (74) or cell cycle dynamics (23).

The developmental influence on generational mutation rates also offers alternative explanations

to other patterns of mutation rate variability that have been attributed to changes in molecular

mutation rates. An example is the differential accumulation of mutations during different stages

of mouse development, which has been interpreted to be the result of a stage-specificity of DNA

repair systems (75), but that could be explained by a stage-specificity of any of the developmental

parameters we explored. Another example is the negative correlation between the generation

times of species and their estimated yearly mutation rates (76). A proposed explanation aligning

with the drift barrier hypothesis is that gamete mutation rate modifiers are constrained by the

product of population size and generation time by selection at the individual level (65). Our intra-

organismal selection model predicts similar trends, albeit through a different mechanism: longer-

lived species allow more efficient elimination of variants that are deleterious the cellular level and

a greater retention of beneficial ones due to extended selective dynamics. Thus, if most mutations

tend to negatively affect cellular fitness, which, as discussed above, may be the case for larger

organisms that also have longer generation times, intraorganismal selection would produce lower

estimations of yearly mutation rates for species with longer generation times.

Developmental parameters and intraorganismal selection shape not only interspecific differences

in mutation rates, but may also help explain variation within species. A well-established pattern

is the higher mutation rates in males compared to females in the germlines of mammals and birds,

a bias largely absent in reptiles and fish, where reproductive biology is more similar between sexes

(16). The male-bias in birds and mammals has been attributed to a greater cell division number

during the lifetime of males (30), but also to imbalances in the DNA repair mechanisms between

the sexes (77). But sex-specific biases could also be caused by intraorganismal selection (40, 41,

56). The selective expansions of mutant lineages in the male germline may exacerbate the

differences in generational mutation rates, including the paternal age effect (78), as has been

reported from clinical cases (41, 79, 80). In terms of our model, male-specific expansions are

facilitated by a greater stem cell turnover over the lifetime of organisms, with females having $d \sim$

0 and males $d > 0$ (Fig. 3B). Interestingly, if intraorganismal selection is at least partially driving

the differences between male and female mutation rates, we expect not only a quantitative bias,

but also a qualitative bias: whereas mutations inherited from females would be enriched in

mutations that are deleterious to cells and would likely get purged if that mutation arose in the

male germline, mutations inherited from fathers are likely enriched in effects increasing cell

fitness.

Although our argument broadly aligns with the idea that non-molecular causes may underlie the

diversification of generational mutation rates (28, 29, 33), we do not rule out the evolutionary

relevance of molecular mechanisms. Alterations to baseline molecular mutation rates are well-

documented (81) and likely coexist with intra-organismal selection and developmental influences

(36, 82). Disentangling whether a particular molecular, developmental or life history parameter

has driven the diversification of generational mutation rates between species may be challenging,

as many relevant factors tend to be correlated. But ultimately, the causes of mutation rate

diversification are likely multifaceted. In this context, we propose an expansion of the definition

of "mutator alleles" (20) beyond molecular mechanisms that affect DNA repair and damage.

Instead, mutator alleles may also expand to genes that influence developmental processes, life

history traits, or organismal physiology.

<u>Considerations about the place of development in evolution</u>

The role of development in evolution is increasingly recognized, but much of the discussion has

focused on how developmental systems constrain or drive phenotypic evolution (83-86). Our

findings emphasize an additional role: by influencing mutation rates, development can shape the

evolutionary pace and direction at both phenotypic and genotypic scales.

In the conceptual framework of genotype space (87), populations could, in principle, explore the

space equally in all directions (Fig. 4A). However, mutational biases—such as the well-

documented transition versus transversion bias—skew this exploration (Fig. 4B), affecting the

spectrum of adaptive substitutions available to populations (4). Beyond molecular biases,

developmental processes can further constrain or promote specific directions of exploration (Fig.

4C) and influence the breadth of genotype space sampled within a generation (Fig. 4D). This has

quantitative and qualitative implications for the genetic diversity of populations. Quantitively,

populations of larger organisms like rheas, for example, can explore a broader portion of genotype

space compared to those of smaller organisms like gulls, which have a considerably smaller

generational mutation rate (16). While qualitatively, this broader exploration may occasionally

introduce deleterious variants, many of these—particularly those disrupting basic cellular

functions or germline potential—are more efficiently pruned in larger species due to their

developmental and intraorganismal selective dynamics. Thus, organisms are not merely vehicles

for genes, as major evolutionary paradigms suggest (88). Instead, development actively

participates in shaping genotypic changes arising in the gene pools of populations and, by

extension, the evolutionary trajectories of species.

Overall, by highlighting the interplay between development and selection at the cellular level, we

provide a broader framework for understanding the diversification of mutation rates. Taking this

framework into account, we suggest that future studies exploring the diversification of mutation

rates should account for the developmental contexts in which mutations arise in each species and

the ecological interactions between cells of different genotypes.

Materials and Methods

<u>Developmental model for estimation of μ_{pop}</u>

We define the generational mutation rate μ_{pop} as the probability of sampling a mutant cell among

all cells in an organism that are capable of producing gametes at the time of reproduction of an

individual. To calculate this probability, it is necessary to first calculate M_i, which represents the

total number of cells that are expected to carry a mutation at a given locus i. We adapted a model

from Otto & Orive (35), wherein M_i is a function of the molecular mutation rate μ_{mol}, the total

developmental time τ, the rate of cell divisions during development C, and a factor β_i, which

represents the effect of a mutation on the fitness of a cell carrying a mutation on locus i. At

developmental time x, one would expect $2^x(1 - \mu_{mol})^{x-1}$ cells to not carry any mutations. Each

of those cells can mutate with a probability given by μ_{mol} and the mutated lineage will then

undergo $\beta_i(C\tau - x)$ cell divisions more until the conclusion of development. Thus, the number of

mutant cells resulting from mutations at time x is:

439
$$2^x(1 - \mu_{mol})^{x-1} * \mu_{mol} * 2^{\beta_i(C\tau-x)}$$

440   Summing over all x gives us M_i at the conclusion of development:

441
$$M_i = \sum_{x=1}^{C\tau} 2^x(1 - \mu_{mol})^{x-1} * \mu_{mol} * 2^{\beta_i(C\tau-x)}$$

442   For estimating μ_{pop} it is also necessary to calculate the expected number of cells that do not

443   undergo mutations at locus i through the course of development i.e., N_i. This can be calculated

444   as:

445
$$N_i = 2^{C\tau}(1 - \mu_{mol})^{C\tau}$$

446   The total number of cells at the end of development is $(M_i + N_i)$ and thus μ_{pop_i} approximates the

447   probability of sampling a cell mutated at locus i from all cells in the pool of gametes, such that

448
$$\mu_{pop_i} = \frac{M_i}{M_i + N_i}$$

449　The calculations shown above correspond to those considering μ_{mol} to track cell divisions. In the

450　cases in which mutation rates are taken to track time rather than cell divisions, the calculations

451　for M_i and N_i were modified such that,

452
$$M_i = \sum_{x=1}^{\tau} 2^{Cx}(1 - \mu_{mol})^{x-1} * \mu_{mol} * 2^{\beta_i C(\tau - x)}$$

453　And,

454
$$N_i = 2^{C\tau}(1 - \mu_{mol})^{\tau}$$

To estimate μ_{pop} across the entire length of a genome, as a first order approximation, we assumed

that no cell obtains mutations at more than one locus during the entire course of development.

In other words, we assume $\mu_{mol}^2 \approx 0$. This assumption would hold for very small μ_{mol}. Using this

simplification, we can estimate μ_{pop} across the entire length of a genome by averaging μ_{pop} across

all sites with different values of β_i. In our model, we considered $F_{\beta i = f}$ to be the fraction of the

total sites in a genome that have a $\beta_i = f$, giving:

462
$$\mu_{pop} = \sum_{f} \mu_{pop_i}(\beta_i = f) \cdot F_{\beta i = f}$$

<u>Expansion of the model to account for maturation and time to reproduction</u>

To consider the role of lifetime of an organism in the calculation of μ_{pop}, we considered the pool

of cells resulting from the developmental process to be constant in number. Immediately at the

end of development, there are $M_{t=0}$ cells in the germline stem cell niche that are mutated and $N_{t=0}$

that are not. During an organism lifetime l, germline stem cells can undergo a turnover (due to

differentiation or death) at a rate d over an arbitrary unit of time t. The rate at which mutant cells

replace lost cells during this turnover is given by a selective variable β_m. As in the case of the

developmental selective parameter β, when $\beta_m < 1$ mutations are deleterious, when $\beta_m = 1$

mutations are neutral and when $\beta_m > 1$ mutations are advantageous to cells. In our model,

during the lifetime of an organism before reproduction, it is possible for the total pool of mutated

cells to decrease either due to reverse mutations (which would occur at a low rate, $M_{t-1}\mu_{mol}$) or

loss due to death or differentiation (dM_{t-1}). Furthermore, mutant cells can increase in proportion

whenever non-mutant cells in the stem cell niche mutate ($N_{t-1}\mu_{mol}$) or whenever mutant cells

proliferate to replenish the stem cell niche after the loss of sibling cells. The number of mutated

cells at time *t* during the lifetime of an organism can thus be calculated as,

478
$$M_t = M_{t-1} + N_{t-1}\mu_{mol} - dM_{t-1} - M_{t-1}\mu_{mol} + \frac{\beta_m d(M_{t-1} + N_{t-1})M_{t-1}}{\beta_m M_{t-1} + N_{t-1}}$$

479   And the number of non-mutant cells would be given by

480
$$N_t = N_{t-1} + M_{t-1}\mu_{mol} - dN_{t-1} - N_{t-1}\mu_{mol} + \frac{d(M_{t-1} + N_{t-1})N_{t-1}}{\beta_m M_{t-1} + N_{t-1}}$$

<u>Comparison of μ_{pop} for different mammalian species</u>

To corroborate the predictions of our model we considered estimations of the μ_{pop} of each of 36

species of mammals. We considered the μ_{pop} of each species to be the mean of the number of de

novo mutations per base pair across different numbers of parent-offspring trios (16). We collected

the other biological parameters used, namely the body mass of each species and the gestation

time, from the Encyclopedia of Life (51). The phylogenetic relationships we considered were

based on recent phylogenetic assessments of mammals (89), while the estimated divergence

times were collected from various sources listed in Table S1. To estimate *C*, we considered

gestation times as a proxy for τ and the mass of neonates as a proxy for $2^{C\tau}$ for those species for

which this information was available in the Encyclopedia of Life.

To consider how the mutation rates of recently diverged species have evolved in relation to

developmental parameters, we compared the ratios of the body mass for pairs of species with

the ratios of mutations rates measured by Bergeron et al (16). The pairs we used for our

comparisons consisted of pairs of the two most closely related species (Fig. 1C). In cases like those

of *Sus scrofa* that have more than one species that are equally closely related, we considered all

possible species pairs (Fig. 1C, black bars). For these comparisons, we considered the ratio of the

mutation rate or body size of the smaller of both species against that of the larger one. Thus, the

ratio of body sizes should be greater than 1 for all comparisons. If the ratio of mutation rates is

also greater than one, body size and mutation rates are aligned, while if it is smaller than one, the

smaller species have a greater mutation rate. To test the significance of our result, we compared

the alignment of the μ_{pop} and body size ratios (the percentage of comparisons for which μ_{pop}

ratios were greater than 1) with the expectation from 1000 random permutations of μ_{pop}. Pairs

of species were considered to be recently diverged if their common ancestor dates back to less

than 65 mya (black bars in Fig. 1C, older pairs are not shown in the phylogeny but we used them

for our comparisons, Fig. 1E, gray point). The same analysis was carried out for the birds species

for which mutation rates were available (16), but no phylogenetic aging was assigned since all

extant birds diverged after the Cretaceous-Tertiary boundary of 65 my.

<u>Estimation of effective population size (N_e) and the relationship between μ_{pop} and N_e</u>

Organisms comprising of $2^{C\tau}$ cells, have a mass $m = m_{cell} \, 2^{C\tau}$, where $m_{cell} = 10^{-15} g$ is the

mass of a single cell (90). We used this mass estimate to compute the effective population size

N_e. This was done by using the scaling of effective population size and organism size (adult dry

weight) that was reported by Lynch et al. (17) for organisms across the tree of life. The fitted log–

log regression for the data resulted in the line: $y = 6.717 - 0.199x$. The above mass estimate

was also used to calculate the mutation rate μ_{pop} using our equation. Finally, we also compared

our predicted curve with mammalian weight and mutation rate data (see section above). Setting

μ_{mol} to $10^{-9.74}$ gave the best agreement with the empirical data.

<u>Flippase assay to assess cell competition in fly germlines</u>

We developed heat shock-induced mitotic clones in fruit flies following a standard protocol (58).

Briefly, a fly line (# 31418 from the Bloomington Drosophila Stock Center) containing a red

fluorescent protein (mRFP), a flippase-encoding gene induced by a heat-shock promoter (hsFLP)

and a FRT site for the binding of flippases was crossed with another line carrying a single-amino

acid substitution in the *trithorax-related* gene (*trr*) and the w^{1118} allele for the gene white. We thus

created a *trr*CA,w^{1118},FRT,hsFLP and a w^{1118},FRT,hsFLP lines. For our experiments we crossed these

lines with the parental mRFP,FRT,hsFLP lines and we exposed the heterozygotic embryos

collected during a window of four hours to a heat shock of 37°C for one hour.

We collected mosaic virgin females that developed from these heat-shocked embryos as well as

virgin females from embryos that were not heat-shocked (control). Each of these virgins were

individually paired to two w^{1118} males and we then counted the fraction of offspring that were

positive for red fluorescence, which are the ones that inherited the mRFP+ allele and not the

mRFP- (trr^{CA},w^{1118}) allele. For the same set of pairs, we evaluated the fraction of mRFP+ pupae

from offspring produced during the first week immediately after mothers were collected, as well

as during the second and third weeks.

Acknowledgments
534
535   We are grateful to Lautaro Gandara, Anna-Lena Vigil, David Porubsky and Cecilia Padilla Iglesias for their
536   useful feedback on our manuscript.
537

References

1.   G. G. Simpson, *The major features of evolution*, Columbia biological series,
(Columbia University Press, New York,, 1953), pp. xx, 434 p.
2.   L. Y. Yampolsky, A. Stoltzfus, Bias in the introduction of variation as an orienting
factor in evolution. *Evolution and Development* **3**, 73-83 (2001).
3.   A. Stoltzfus, L. Y. Yampolsky, Climbing Mount Probable: Mutation as a Cause of
Nonrandomness in Evolution. *J Hered* **100**, 637-647 (2009).
4.   A. V. Cano, H. Rozhoňová, A. Stoltzfus, D. M. McCandlish, J. L. Payne, Mutation
bias shapes the spectrum of adaptive substitutions. *Proc Natl Acad Sci U S A* **119**
(2022).
5.   A. V. Cano, J. L. Payne, Mutation bias interacts with composition bias to influence
adaptive evolution. *Plos Comput Biol* **16** (2020).
6.   M. Nei, *Mutation-driven evolution* (Oxford University Press, Oxford, ed. 1st,
2013), pp. xi, 244 p.
7.   N. S. Martin, S. Schaper, C. Q. Camargo, A. A. Louis, Y. Kim, Non-Poissonian
Bursts in the Arrival of Phenotypic Variation Can Strongly Affect the Dynamics of
Adaptation. *Mol Biol Evol* **41** (2024).
8.   D. Agashe, M. Sane, S. Singhal, Revisiting the Role of Genetic Variation in
Adaptation. *The American Naturalist* **202**, 486-502 (2023).
9.   M. Kimura, The Rate of Molecular Evolution Considered from the Standpoint of
Population Genetics. *Proc Natl Acad Sci U S A* **63**, 1181-1188 (1969).
10.   E. Zuckerkand, L. Pauling, Molecules as documents of evolutionary history. *J
Theor Biol* **8**, 357-366 (1965).

11. M. Kimura, Molecular evolutionary clock and the neutral theory. *J Mol Evol* **26**,
24-33 (1987).
12. T. H. Jukes, R. Holmquist, Evolutionary Clock: Nonconstancy of Rate in Different
Species. *Science* **177**, 530-532 (1972).
13. J. H. Gillespie, *The causes of molecular evolution*, Oxford series in ecology and
evolution (Oxford University Press, New York, 1991), pp. xiv, 336 p.
14. R. J. Britten, Rates of DNA Sequence Evolution Differ Between Taxonomic
Groups. *Science* **231**, 1393-1398 (1986).
15. Y. Wang, D. J. Obbard, Experimental estimates of germline mutation rate in
eukaryotes: a phylogenetic meta-analysis. *Evolution Letters* **7**, 216-226 (2023).
16. L. A. Bergeron *et al.*, Evolution of the germline mutation rate across vertebrates.
Nature **615**, 285-291 (2023).
17. M. Lynch *et al.*, The divergence of mutation rates and spectra across the Tree of
Life. *Embo Rep* **24** (2023).
18. K. Harris, Evidence for recent, population-specific evolution of the human
mutation rate. *Proc Natl Acad Sci U S A* **112**, 3439-3444 (2015).
19. P. D. Sniegowski, P. J. Gerrish, R. E. Lenski, Evolution of high mutation rates in
experimental populations of E. coli. *Nature* **387**, 703-705 (1997).
20. M. Lynch *et al.*, Genetic drift, selection and the evolution of the mutation rate.
Nat Rev Genet **17**, 704-714 (2016).
21. P. D. Sniegowski, P. J. Gerrish, T. Johnson, A. Shaver, The evolution of mutation
rates: separating causes from consequences. *Bioessays* **22**, 1057-1066 (2000).
22. A. C. Beichman, L. Zhu, K. Harris, The Evolutionary Interplay of Somatic and
Germline Mutation Rates. *Annual Review of Biomedical Data Science* **7**, 83-105
(2024).
23. N. Spisak, M. de Manuel, W. Milligan, G. Sella, M. Przeworski, The clock-like
accumulation of germline and somatic mutations can arise from the interplay of
DNA damage and repair. *PLOS Biology* **22** (2024).
24. F. Taddei *et al.*, Role of mutator alleles in adaptive evolution. *Nature* **387**, 700-
702 (1997).
25. M. Kimura, On the evolutionary adjustment of spontaneous mutation rates.
Genetical Research **9**, 23-34 (1967).
26. M. Lynch, Evolution of the mutation rate. *Trends in Genetics* **26**, 345-352 (2010).
27. W. Sung, M. S. Ackerman, S. F. Miller, T. G. Doak, M. Lynch, Drift-barrier
hypothesis and mutation-rate evolution. *Proc Natl Acad Sci U S A* **109**, 18488-
18492 (2012).
28. G. W. C. Thomas *et al.*, Reproductive Longevity Predicts Mutation Rates in
Primates. *Current Biology* **28**, 3193-3197.e3195 (2018).
29. Z. Gao, M. J. Wyman, G. Sella, M. Przeworski, Interpreting the Dependence of
Mutation Rates on Age and Time. *PLOS Biology* **14** (2016).
30. C. F. Baer, M. M. Miyamoto, D. R. Denver, Mutation rate variation in multicellular
eukaryotes: causes and consequences. *Nat Rev Genet* **8**, 619-631 (2007).

31.   G. Amster, G. Sella, Life history effects on the molecular clock of autosomes and
604   sex chromosomes. *Proc Natl Acad Sci U S A* **113**, 1588-1593 (2016).
605   32.   L. W. Buss, Evolution, development, and the units of selection. *Proc Natl Acad Sci
606   U S A* **80**, 1387-1391 (1983).
607   33.   P. Moorjani, Z. Gao, M. Przeworski, Human Germline Mutation and the Erratic
608   Evolutionary Clock. *PLOS Biology* **14** (2016).
609   34.   I. M. Hastings, Potential Germline Competition in Animals and Its Evolutionary
610   Implications. *Genetics* **123**, 191-197 (1989).
611   35.   S. P. Otto, M. E. Orive, Evolutionary Consequences of Mutation and Selection
612   within an Individual. *Genetics* **141**, 1173-1187 (1995).
613   36.   P. Majic, J. L. Payne, Developmental selection and the perception of mutation
614   bias. *Mol Biol Evol* 10.1093/molbev/msad179 (2023).
615   37.   C. Extavour, A. García-Bellido, Germ cell selection in genetic mosaics in
616   Drosophila melanogaster. *P Natl Acad Sci USA* **98**, 11341-11346 (2001).
617   38.   C.-Y. Tseng *et al.*, chinmo-mutant spermatogonial stem cells cause mitotic drive
618   by evicting non-mutant neighbors from the niche. *Dev Cell* **57**, 80-94.e87 (2022).
619   39.   D. Haig, Mutational selection: fragile sites, replicative stress, and genome
620   evolution. *Qeios* 10.32388/9eryw5 (2023).
621   40.   S. P. Otto, I. M. Hastings, Mutation and selection within the individual. *Genetica*
622   **102-3**, 507-524 (1998).
623   41.   N. Arnheim, P. Calabrese, Germline Stem Cell Competition, Mutation Hot Spots,
624   Genetic Disorders, and Older Fathers. *Annual Review of Genomics and Human
625   Genetics* **17**, 219-243 (2016).
626   42.   F. L. Wu *et al.*, A comparison of humans and baboons suggests germline
627   mutation rates do not track cell divisions. *PLOS Biology* **18** (2020).
628   43.   G. J. Kenagy, S. C. Trombulak, Size and Function of Mammalian Testes in Relation
629   to Body Size. *Journal of Mammalogy* **67**, 1-22 (1986).
630   44.   B. Colom *et al.*, Spatial competition shapes the dynamic mutational landscape of
631   normal esophageal epithelium. *Nature Genetics* **52**, 604-614 (2020).
632   45.   A. R. J. Lawson *et al.*, Extensive heterogeneity in somatic mutation and selection
633   in the human bladder. *Science* **370**, 75-+ (2020).
634   46.   I. Martincorena, P. J. Campbell, Somatic mutation in cancer and normal cells.
635   *Science* **349**, 1483-1489 (2015).
636   47.   I. Martincorena *et al.*, Somatic mutant clones colonize the human esophagus
637   with age. *Science* **362**, 911-+ (2018).
638   48.   I. Martincorena, P. H. Jones, P. J. Campbell, Constrained positive selection on
639   cancer mutations in normal skin. *P Natl Acad Sci USA* **113**, E1128-E1129 (2016).
640   49.   I. Martincorena *et al.*, Universal Patterns of Selection in Cancer and Somatic
641   Tissues. *Cell* **171**, 1029-1041.e1021 (2017).
642   50.   I. Martincorena *et al.*, High burden and pervasive positive selection of somatic
643   mutations in normal human skin. *Science* **348**, 880-886 (2015).

644   51.   C. S. Parr *et al.*, The Encyclopedia of Life v2: Providing Global Access to
645   Knowledge About Life on Earth. *Biodiversity Data Journal* **2** (2014).

646   52.   P. Alberch, S. J. Gould, G. F. Oster, D. B. Wake, Size and shape in ontogeny and
647   phylogeny. *Paleobiology* **5**, 296-317 (1979).

648   53.   D. Porubsky *et al.*, Human de novo mutation rates from a four-generation
649   pedigree reference. *Nature* 10.1038/s41586-025-08922-2 (2025).

650   54.   J. Damuth, Population density and body size in mammals. *Nature* **290**, 699-700
651   (1981).

652   55.   J. D. Yeakel, C. P. Kempes, S. Redner, Dynamics of starvation and recovery
653   predict extinction risk and both Damuth's law and Cope's rule. *Nat Commun* **9**
654   (2018).

655   56.   R. A. Hodge, E. A. Bach, Mechanisms of Germline Stem Cell Competition across
656   Species. *Life* **14** (2024).

657   57.   H. Kanda, A. Nguyen, L. Chen, H. Okano, I. K. Hariharan, The Drosophila Ortholog
658   of MLL3 and MLL4, trithorax related, Functions as a Negative Regulator of Tissue
659   Growth. *Mol Cell Biol* **33**, 1702-1710 (2023).

660   58.   K. G. Golic, S. Lindquist, The FLP recombinase of yeast catalyzes site-specific
661   recombination in the drosophila genome. *Cell* **59**, 499-509 (1989).

662   59.   L. Gandara *et al.*, Developmental phenomics suggests that H3K4
663   monomethylation confers multi-level phenotypic robustness. *Cell Reports* **41**
664   (2022).

665   60.   E. R. te Velde, P. L. Pearson, The variability of female reproductive ageing.
666   *Human Reproduction Update* **8**, 141-154 (2002).

667   61.   K. C. Silva-Santos *et al.*, Estimate of the population of preantral follicles in the
668   ovaries of Bos taurus indicus and Bos taurus taurus cattle. *Theriogenology* **76**,
669   1051-1057 (2011).

670   62.   B. Griffing, Analysis of Quantitative Gene Action by Constant Parent Regression
671   and Related Techniques. *Genetics* **35**, 303-321 (1950).

672   63.   D. Aggeli, Y. Li, G. Sherlock, Changes in the distribution of fitness effects and
673   adaptive mutational spectra following a single first step towards adaptation. *Nat*
674   *Commun* **12** (2021).

675   64.   E. R. Jerison, M. M. Desai, Genomic investigations of evolutionary dynamics and
676   epistasis in microbial evolution experiments. *Curr Opin Genet Dev* **35**, 33-39
677   (2015).

678   65.   L. Zhu, A. Beichman, K. Harris, 10.1101/2023.12.06.570457 (2024).

679   66.   E. Witt, C. B. Langer, N. Svetec, L. Zhao, Transcriptional and mutational
680   signatures of the Drosophila ageing germline. *Nat Ecol Evol* **7**, 440-449 (2023).

681   67.   T. Nakagawa, Y.-i. Nabeshima, S. Yoshida, Functional Identification of the Actual
682   and Potential Stem Cell Compartments in Mouse Spermatogenesis. *Dev Cell* **12**,
683   195-206 (2007).

68. A. M. Klein, T. Nakagawa, R. Ichikawa, S. Yoshida, B. D. Simons, Mouse Germ Line
Stem Cells Undergo Rapid and Stochastic Turnover. *Cell Stem Cell* **7**, 214-224
(2010).
69. M. Lynch, The Cellular, Developmental and Population-Genetic Determinants of
Mutation-Rate Evolution. *Genetics* **180**, 933-943 (2008).
70. A. Cagan *et al.*, Somatic mutation rates scale with lifespan across mammals.
Nature **604**, 517-524 (2022).
71. G. W. C. Thomas, M. W. Hahn, The Human Mutation Rate Is Increasing, Even As It
Slows. *Mol Biol Evol* **31**, 253-257 (2014).
72. A. Wong, Covariance between Testes Size and Substitution Rates in Primates.
Mol Biol Evol **31**, 1432-1436 (2014).
73. A. C. Beichman *et al.*, Evolution of the Mutation Spectrum Across a Mammalian
Phylogeny. *Mol Biol Evol* **40** (2023).
74. L. B. Alexandrov *et al.*, Mutational signatures associated with tobacco smoking in
human cancer. *Science* **354**, 618-622 (2016).
75. S. J. Lindsay, R. Rahbari, J. Kaplanis, T. Keane, M. E. Hurles, Similarities and
differences in patterns of germline mutation between mice and humans. *Nat
Commun* **10** (2019).
76. L. Lewin, A. Eyre-Walker, K. Harris, Estimates of the Mutation Rate per Year Can
Explain Why the Molecular Clock Depends on Generation Time. *Mol Biol Evol* **42**
(2025).
77. F. L. Wu, M. de Manuel, M. Przeworski, A paternal bias in germline mutation is
widespread in amniotes and can arise independently of cell division numbers.
eLife **11** (2022).
78. J. M. Goldmann, J. A. Veltman, C. Gilissen, De Novo Mutations Reflect
Development and Aging of the Human Germline. *Trends in Genetics* **35**, 828-839
(2019).
79. S.-R. Yoon *et al.*, Age-Dependent Germline Mosaicism of the Most Common
Noonan Syndrome Mutation Shows the Signature of Germline Selection. *The
American Journal of Human Genetics* **92**, 917-926 (2013).
80. S.-K. Choi, S.-R. Yoon, P. Calabrese, N. Arnheim, A germ-line-selective advantage
rather than an increased mutation rate can explain some unexpectedly common
human disease mutations. *Proc Natl Acad Sci U S A* **105**, 10143-10148 (2008).
81. T. A. Sasani *et al.*, A natural mutator allele shapes mutation spectrum variation in
mice. *Nature* **605**, 497-502 (2022).
82. B. Xia *et al.*, Widespread Transcriptional Scanning in the Testis Modulates Gene
Evolution Rates. *Cell* **180**, 248-262.e221 (2020).
83. J. Maynard Smith *et al.*, Developmental Constraints and Evolution. *Q Rev Biol* **60**,
265-287 (1985).
84. W. Arthur, Developmental drive: an important determinant of the direction of
phenotypic evolution. *Evolution and Development* **3**, 271-278 (2001).

85.　P. Majic, E. Y. Erten, J. L. Payne, The adaptive potential of nonheritable somatic
mutations. *The American Naturalist* 10.1086/721766 (2022).
86.　R. Galupa *et al.*, Enhancer architecture and chromatin accessibility constrain
phenotypic space during Drosophila development. *Dev Cell* **58**, 51-62.e54 (2023).
87.　S. Wright, The roles of mutation, inbreeding, crossbreeding, and selection in
evolution. 355-366 (1932).
88.　R. Dawkins, *The selfish gene* (Oxford University Press, Oxford, 1976), pp. xii, 224
p.
89.　N. M. Foley *et al.*, A genomic timescale for placental mammal evolution. *Science*
380 (2023).
90.　R. Sender, S. Fuchs, R. Milo, Revised Estimates for the Number of Human and
Bacteria Cells in the Body. *PLOS Biology* **14** (2016).
91.　B. Milholland *et al.*, Differences between germline and somatic mutation rates in
humans and mice. *Nat Commun* **8** (2017).
92.　X. Wang, "Diversity: Who Is Who in the Dog Family" in Dogs: Their Fossil
Relatives and Evolutionary History. (Columbia University Press, 2008),
10.7312/wang13528-004, pp. 23-68.
93.　S. J. Rossiter *et al.*, Phylogenomic Resolution of the Cetacean Tree of Life Using
Target Sequence Capture. *Systematic Biology* **69**, 479-501 (2020).
94.　A. Hassanin, E. J. P. Douzery, J. Thorne, Molecular and Morphological
Phylogenies of Ruminantia and the Alternative Position of the Moschidae.
Systematic Biology **52**, 206-228 (2003).
95.　W. E. Johnson *et al.*, The Late Miocene Radiation of Modern Felidae: A Genetic
Assessment. *Science* **311**, 73-77 (2006).
96.　C. J. Law, G. J. Slater, R. S. Mehta, Lineage Diversity and Size Disparity in
Musteloidea: Testing Patterns of Adaptive Radiation Using Molecular and Fossil-
Based Methods. *Systematic Biology* **67**, 127-144 (2018).
97.　C. Gilbert, A. Ropiquet, A. Hassanin, Mitochondrial and nuclear phylogenies of
Cervidae (Mammalia, Ruminantia): Systematics, morphology, and biogeography.
Molecular Phylogenetics and Evolution **40**, 101-117 (2006).

     Figures and Tables

757
Figure 1. Changes in developmental parameters can causally explain diversification of mutation
rates. (A) Illustrated summary of the core model consisting of an individual developmental phase and
a reproductive phase. In the developmental phase a lineage starting from a single cell goes through
successive cell divisions at a rate C over a developmental time τ. During development, mutations can

occur at a rate μ_{mol}. Cell proliferation could be affected by the fitness effect of mutations on cells β.
For β = 1 mutations are neutral. At the end of development, during the reproductive phase, the
fraction of germline-capable cells that carry mutations define the rate at which new alleles are added
to the gene pool μ_{pop}. (B) Population-level mutation rate μ_{pop} at a single locus i as a function of τ. Each
curve represents a different value of C. μ_{mol} was held constant at a value of 1×10^{-10} mutations per cell
division, approximating empirical estimates for human and mouse (91). (C) Phylogeny of mammals for
which estimations for μ_{pop} are available (16). Black bars indicate recently diverged species pairs for
which body size and μ_{pop} were considered. (D) μ_{pop} for each mammalian species in (C) as a function of
their gestation time, a proxy for τ. (E) Ratio of the μ_{pop} of species pairs as a function of the ratio of the
mass of species pairs. Black dots indicate branchings younger than 65 my, while gray dots represent
older branchings. Notice how, for the vast majority of comparisons between younger branches μ_{pop}
ratios are above 1 (red line), which means that larger clades tend to have higher mutation rates, as
predicted by our model. Red arrowheads indicate the species pairs corresponding to the tiger
(*Panthera tigris*) and leopard (*Panthera pardus*) and goat (*Capra hircus*) and dwarf musk deer (*Moschus*
berezovskii) (F) Datapoints represent the expected relationship between the predicted μ_{pop} of each of
the studied mammalian species and their effective population size N_e based on their body mass
(Methods). The green line represents the prediction of this relationship based on our developmental
model for μ_{pop}.

Figure 2. The impact of mutations on cell fitness during development can repercuss on mutation rates. (A) In the model introduced in Fig. 1A, when $\beta \neq 1$, mutations can affect the total number of mutated cells at the end of development M. (B) Induced mitotic recombination can generate mosaic organisms with cells with different genotypes. The shown specimen corresponds to a female *Drosophila melanogaster* on which mosaicism can be easily observed in the eyes. Red, white, and orange patches represent each of the possible genotypes thanks to the w^{1118} allele. (C) Fraction of offspring carrying the mRFP(-) allele from crossing a female of interest with a w^{1118} male. TrrCA refers to females of the trr^{CA}, w^{1118} genotype, while TrrWT refers to the w^{1118} genotype that is wild-type for *trr*. "HS" refers to females that developed from embryos that were heat-shocked, while "Control" females developed from embryos that were not heat-shocked. Each point represents the fraction of offspring from a different mother. * indicates $p < 0.05$, ** indicates $p < 0.005$ using a Wilcoxon test. (D) Population-level mutation rate μ_{pop} as a function of the effect of mutations on the fitness of cells β. Each curve represents a different value of developmental time τ with a constant value for $C = 1$. (E) Population-level mutation rate μ_{pop} as a function of developmental parameters $C\tau$. Each curve represents a different value proportion of neutral or deleterious mutations. (F) Population-level mutation rate μ_{pop} as a function of developmental parameters $C\tau$. Each curve represents a different proportion of mutations that are advantageous for the cell ($\beta = 1.5$) in the context of an otherwise fully neutral sequence.

801
Figure 3. Selection within the germline stem cell niche can also impact mutation rates over an
organism's lifetime. (A) Illustrated summary of the model integrating a maturation phase before
reproduction. Adding this phase incorporates parameters such as the time to reproduction of an
organism l, a turnover rate of germline stem cells d, and a cellular fitness variable for the stem cell
niche β_l. (B) Population-level mutation rate μ_{pop} as a function of the rate of stem cell turnover d for a
constant generation time of $l = 50$. Each curve represents a different proportion of mutations that are
advantageous for the cell ($\beta_l = 1.5$) in the context of an otherwise fully neutral sequence (gray to purple
lines), a fully neutral case (black) or a fully deleterious case (green). (C) Population-level mutation rate
μ_{pop} as a function of generation time l for a constant rate of stem cell turnover $d = 0.1$. The different
lines follow the same color code as in (B). (D) μ_{pop} as a function of cellular fitness variable for the stem
cell niche β_l for different values of l and the fitness effect of mutation on cells during development β.
(E and F) Fraction of offspring carrying the mRFP(-) allele from the same set of mothers
reproducing at the beginning of weeks one, two or three post-hatching. TrrCA refers to females of
the trr^{CA}, w^{1118} genotype, while TrrWT refers to the w^{1118} genotype that is wild-type for trr. "HS" refers
to females that developed from embryos that were heat-shocked. * indicates $p < 0.05$.

Figure 4. How development can influence the trajectories and pace of genotypic evolution. In a
theoretical model of evolution, it could be considered that given a starting genotype, GTC, all
mutational steps are equally likely (A). However, if molecular realism is considered there are certain
mutations that are more likely than others, as represented by arrows of different thickness going to
CTC and GAC. (B). Furthermore, as we argue in this work, development can act as a selective filter for
mutations which increases or decreases the likelihood of mutational steps depending on how a
mutation affects the cell lineage it belongs to (C). Developmental and life-history differences can also
impact how far mutational steps can take an organism. As per our model, larger organisms should be
capable of more broadly exploring genotypic spaces within a generation (D).

Supplementary materials

Table S1. Empirical parameters for 36 mammalian species.

Bergeron et al (2023) estimated the number of de novo mutations in parent-offspring trios of each of the studied species. With those estimations we calculated μ_{pop} as the average of all replicates reported. The mass and τ (gestation time) values were collected from the Encyclopedia of Life (Parr et al 2014). Branch ages representing the estimated divergence time from the closest relative in the phylogeny (Fig. 1C) were gathered from several sources indicated by the superindeces in the Branch Age column: [1](89), [2](92) [3](93) [4](94) [5](95) [6](96) [7](97).

Species	μ_{pop}	Mass (grams)	τ (months)	Branch age (my)
Vicugna pacos	9.44 x10^{-9}	60000	10.79	66[1]
Cavia aperea	8.67 x10^{-9}	442.8	2.07	56[1]
Rousettus aegyptiacus	8.11 x10^{-9}	135.5	4.1	80[1]
Saimiri boliviensis boliviensis	6.90 x10^{-9}	615	5.23	37.3[1]
Felis catus	3.49 x10^{-9}	2884	2.09967	33.9
Pan troglodytes	1.25 x10^{-8}	49896	7.36	7.246[1]
Sarcophilus harrisii	5.95 x10^{-9}	8109.3	0.7	160[1]
Canis lupus familiaris	8.68 x10^{-9}	32183.3	2.16	33[2]
Tursiops truncatus	8.83 x10^{-9}	209791	12.12	19.5[3]
Mandrillus leucophaeus	6.33 x10^{-9}	18250	6.19	37.71[1]
Vulpes vulpes	4.46 x10^{-9}	5603.75	1.73	33[2]
Hylobates lar	7.05 x10^{-9}	5930	7.08	27[1]
Giraffa camelopardalis	1.01 x10^{-8}	932324.5	14.89	45[4]
Capra hircus	5.26 x10^{-9}	61000	5.22	27.8[1]
Hippopotamus amphibius	1.19 x10^{-8}	1417490	7.75	61[1]
Homo sapiens	1.17 x10^{-8}	58540	9	7.246[1]
Procavia capensis	9.20 x10^{-9}	3029.3	7.44	100[1]
Panthera pardus	8.84 x10^{-9}	52038.3	3.22	10[5]
Neovison vison	4.64 x10^{-9}	834.67	1.19	33[6]
Fukomys damarensis	7.41 x10^{-9}	88.5	3	56[1]
Mus musculus	5.35 x10^{-9}	10.1	0.69	66[1]
Moschus berezovskii	1.00 x10^{-8}	12477	6.1413	27.8[1]
Monodelphis domestica	4.60 x10^{-9}	66.5	0.49	160[1]
Orcinus orca	1.41 x10^{-8}	4300000	14.89	19.5[3]
Sus scrofa	4.32 x10^{-9}	86250	3.95	60[1]
Ailurus fulgens	1.31 x10^{-8}	4900	4.3	33[6]
Rangifer tarandus	5.90 x10^{-9}	94744.67	7.24	20.2[7]
Cervus elaphus yarkandensis	6.71 x10^{-9}	165016	8.03	6.8[7]
Ceratotherium simum simum	9.56 x10^{-9}	2591659.5	15.87	61[1]
Pithecia pithecia	9.94 x10^{-9}	1900	5.32	37.3[1]
Arctocephalus gazella	6.78 x10^{-9}	143500	9	33.9[1]
Tupaia chinensis belangeri	7.50 x10^{-9}	200	1.5	77.69[1]
Cervus nippon	4.71 x10^{-9}	74750	7.59	6.8[7]
Tapirus indicus	4.80 x10^{-9}	296250	12.76	61[1]
Panthera tigris	6.99 x10^{-9}	141132	3.51	10[5]
Odobenus rosmarus	1.32 x10^{-8}	825000	11.5	33.9[1]

837
838

839
Figure S1. (A) Population-level mutation rate μ_{pop} at a single locus *i* as a function of τ. μ_{mol} was held constant
at a value of 1×10^{-10} mutations per developmental time unit. Each curve represents a different value of *C*,
but they are all collapsed on the same line. (B) μ_{pop} for each mammalian species in Fig. 1C as a function of
$C\tau$, which we estimated as the log2 of the number of cells of an organism, measured as the ratio between
body mass and the mass of a single cell (Methods). (C) Barplot showing the fraction of species pairs for
which the larger of both species also a higher μ_{pop}. The green bar shows the mean fraction of 1000 random
permutations of the values with error bars indicating the standard deviation, the gray bar indicates the
fraction for species pairs that branched more than 65 mya, while the black bar shows the fraction for
recently diverged species pairs.

Figure S2. (A) Phylogeny of birds for which estimations for μ_{pop} are available (16). (B) Ratio of the μ_{pop}
of species pairs as a function of the ratio of the mass of species pairs. Notice how, for the vast majority
of comparisons between younger branches μ_{pop} ratios are above 1 (red line), which means that larger
clades tend to have higher mutation rates, as predicted by our model. (C) Barplot showing the fraction
of species pairs for which the larger of both species also a higher μ_{pop}. The green bar shows the mean
fraction of 1000 random permutations of the values with error bars indicating the standard deviation, while
the black bar shows the fraction for all considered empirical species pairs.

Figure S3. (A) Population-level mutation rate μ_{pop} at a single locus i as a function of β_i for $\tau = 20$. μ_{mol} was held constant at a value of 1×10^{-10} mutations per developmental time unit. Each curve represents a different value of C. (B) μ_{pop} for each mammalian species in Table S1 for which neonate mass was available as a function of an estimated C (Methods).