ADAR1 Regulates Lipid Remodeling through MDM2 to Dictate Ferroptosis Sensitivity

Che-Pei Kung[1], Nick D. Terzich[2], Ma Xenia G. Ilagen[3,4], Michael J. Prinsen[3], Madhurima

Kaushal[5], Raleigh D. Kladney[1], Jade H. Weber[1], Alex R. Mabry[1,6], Luisangely Soto Torres[1,6],

Emily R. Bramel[1], Eric C. Freeman[1], Thwisha Sabloak[1], Kyle A. Cottrell[1], Sua Ryu[1], William

M. Weber[1], Leonard Maggi Jr.[1], Leah P. Shriver[2,7], Gary J. Patti[2,8,9] and Jason D. Weber[1,4,9,10]

[1]Department of Medicine, Division of Molecular Oncology, Washington University School of

Medicine, Saint Louis, MO, USA

[2]Department of Chemistry, Washington University, St. Louis, MO, USA

[3]High Throughput Screening Center, Washington University School of Medicine, Saint Louis,

MO, USA

[4]Department of Biochemistry and Molecular Biophysics, Washington University School of

Medicine, Saint Louis, MO, USA

[5]Institute for Informatics, Data Science and Biostatistics, Washington University School of

Medicine, Saint Louis, MO, USA

[6]Cancer Biology Graduate Program, Washington University School of Medicine, Saint Louis,

Missouri, USA

[7]Department of Medicine, Division of Nutritional Science and Obesity Medicine, Washington

University School of Medicine, Saint Louis, MO, USA

[8]Department of Genetics, Washington University, St. Louis, MO, USA

[9]Siteman Cancer Center, Washington University, St. Louis, MO, USA

[10]Department of Cell Biology and Physiology, Washington University School of Medicine, Saint

Louis, MO, USA

Running title: ADAR1-mediated ferroptosis resistance in triple-negative breast cancer

26 Co-correspondence:
27 Che-Pei Kung, Ph.D.
28 Department of Medicine, Division of Molecular Oncology
29
Washington University School of Medicine
660 South Euclid Avenue
Campus Box 8069
St. Louis, MO 63110 USA
Email: patkung@wustl.edu
Telephone: 314-747-3898
Fax: 314-362-0152
37
38 and
39
40 Jason D. Weber, Ph.D.
41 Department of Medicine, Division of Molecular Oncology
42 Department of Cell Biology and Physiology
43
Washington University School of Medicine
660 South Euclid Avenue
Campus Box 8069
St. Louis, MO 63110 USA
Email: jweber@wustl.edu
Telephone: 314-747-3896
Fax: 314-362-0152
51

Abstract

Triple-negative breast cancer (TNBC), lacking expression of estrogen, progesterone, and HER2 receptors, is aggressive and lacks targeted treatment options. An RNA editing enzyme, adenosine deaminase acting on RNA 1 (ADAR1), has been shown to play important roles in TNBC tumorigenesis. We posit that ADAR1 functions as a homeostatic factor protecting TNBC from internal and external pressure, including metabolic stress. We tested the hypothesis that the iron-dependent cell death pathway, ferroptosis, is a ADAR1-protected metabolic vulnerability in TNBC by showing that ADAR1 knockdown sensitizes TNBC cells to GPX4 inhibitors. By performing single-reaction monitoring-based liquid chromatography coupled to mass spectrometry (LC-MS) to measure intracellular lipid contents, we showed that ADAR1 loss increased the abundance of polyunsaturated fatty acid phospholipids (PUFA-PL), of which peroxidation is the primary driver of ferroptosis. Transcriptomic analyses led to the discovery of the proto-oncogene MDM2 contributing to the lipid remodeling in TNBC upon ADAR1 loss. A phenotypic drug screen using a ferroptosis-focused library was performed to identify FDA-approved cobimetinib as a drug-repurposing candidate to synergize with ADAR1 loss to suppress TNBC tumorigenesis. By demonstrating that ADAR1 regulates the metabolic fitness of TNBC through desensitizing ferroptosis, we aim to leverage this metabolic vulnerability to inform basic, pre-clinical, and clinical studies to develop novel therapeutic strategies for TNBC.

Introduction

Triple-negative breast cancer (TNBC), defined by the lack of estrogen receptor (ER),

progesterone receptor (PR) and HER2 expression, accounts for about 15 percent of all breast

cancer diagnoses in the United States each year[1]. Despite recent development of PARP

inhibitors, immunotherapy and antibody-drug conjugates, there remains a significant gap for

targeted therapies against TNBC[2]. It leaves surgery, radiation and chemotherapy as the main

treatment options, leading to generally poor prognosis. A multi-prong approach to investigate the

immunogenic, epigenetic and metabolic landscapes of TNBC as well as its tumor

microenvironment (TME) is underway to identify points of vulnerability to develop novel and

effective therapies[3-5].

One such vulnerability is iron-mediated cell death, or ferroptosis. Ferroptosis is a distinctive

form of programmed cell death depending on iron-mediated peroxidation of long

polyunsaturated fatty acids (PUFA) to incorporate into and break down cell membranes[6]. TNBC

cells have been found to be sensitive to ferroptosis through multiple mechanisms. TNBC cell

lines have been found to be glutamine auxotrophs, resulting in increased levels of environmental

cystine acquisition and heightened sensitivity to inhibition of the glutamate/cystine antiporter

xCT[7]. The xCT-cystine axis operates upstream of glutathione, which counters oxidative stress

through its antioxidant activity. It has been shown that cellular levels of glutathione are lower in

TNBC cell lines, rendering them more vulnerable to ferroptosis inducers that function through

inhibiting GPX4, the catalyzer of glutathione's antioxidant function[8,9]. Dysregulation of iron

metabolism has also been noted in TNBC, expressing lower levels of iron exporter ferroportin

leading to increased cellular labile iron pools[9,10]. Moreover, a subset of TNBC cell lines

expressed higher level of ACSL4, which activates long-chain PUFA for phospholipid

biosynthesis to enrich cellular membranes[11]. It has been suggested that TNBC's levels of

sensitivity to ferroptosis are correlated with abundance of specific PUFA species, partially

dictated by expression levels of enzymes contributing to PUFA biosynthesis, such as Acyl-CoA

synthetase long-chain family member 1 (ACSL1) and fatty acid desaturases 1 and 2

(FADS1/2)[12,13].

Adenosine Deaminase Acting on RNA (ADAR1, encoded by *ADAR*) can convert adenosine to

inosine within cellular dsRNA, in a process known as A-to-I editing. Elevated expression of

ADAR1 is prevalent in breast cancer and ADAR1-mediated A-to-I editing has been shown to

contribute to tumor progression and therapy resistance in breast cancer, including TNBC[14-18]. We

have previous reported a TNBC-centric ADAR1 dependency phenomenon, in which ADAR1

protects TNBC cells from translational stress and type I interferon (IFN) activation predisposed

by TNBC cells' elevated level of interferon stimulated gene (ISG) signatures[19]. ADAR1-

mediated regulation of innate immune response has been noted in several cancer types, and

ADAR1 loss has been shown to sensitize cancer cells to immunotherapy[20-23]. Through its ability

to prevent cytotoxic inflammatory response, ADAR1 also protects cancer cells from other anti-

tumorigenic stresses, such as necroptosis and epigenetic modulation[24,25]. Few studies, however,

have investigated ADAR1's ability to regulate cancer cells' metabolic fitness and their

sensitivity to metabolic stresses. Considering ADAR1's known role in metabolic regulations and

the importance of cancer metabolism, it is important to fill such knowledge gaps[26,27].

In the current study, we demonstrate that ADAR1 protects TNBC cells from ferroptotic cell

death through regulating MDM2 expression and lipid profile reprogramming. Our observations

indicate broader applications of ADAR1-targeting therapies and open opportunities to repurpose

ferroptosis-inducing drugs for synergistic effects against TNBC.

Results

ADAR1 protects TNBC cells from ferroptosis induction

To determine if ADAR1 inhibits ferroptosis directly, we knocked down ADAR1 expression with

a short-hairpin RNA (shRNA) in TNBC cell line MDA-MB-231 (**Figure S1A**). The level of

malondialdehyde (MDA), a byproduct of lipid peroxidation, was measured to show that cells

with ADAR1 knockdown (ADAR1-Deficient cells) did not display increased ferroptosis activity

compared to cells transduced with non-targeting shRNA (shNT), or ADAR1-Intact cells (**Figure**

S1B). Similarly, by staining both the reduced and oxidized forms of the lipid peroxidation probe

BODIPY^{TM} 581/591 C11 to compare their intensity, it was revealed that reduced expression of

ADAR1 did not enhance the cells' ability to oxidize polyunsaturated fatty acids (**Figure S1C-D**).

To corroborate these observations, cell viability analysis was conducted to show that reduced

proliferation and viability of MDA-MB-231 cells resulting from ADAR1 knockdown could be

partially rescued by either an apoptosis inhibitor (Z-VAD-FMK) or an eIF2α inhibitor (ISRIB),

but not by ferroptosis inhibitors Ferrostatin-1 and Liproxstatin-1 (**Figure S1E**). These results

suggest that altering ADAR1 expression does not overtly impact ferroptosis activity in cells.

ADAR1-Intact and ADAR1-Deficient MDA-MB-231 cells then were tested to determine if

ADAR1 expression influences cells' sensitivity to ferroptosis induction. Ferroptosis in MDA-

MB-231 cells can be induced by GPX4 inhibitor RSL3, whose effect can be rescued by

ferroptosis inhibitor Ferrostatin-1 or UAMC, but not Z-VAD-FMK or necroptosis inhibitor

Necrostatin-1 (**Figure S1F**). It was first observed that ADAR1-Deficient MDA-MB-231 cells

became much more sensitive, compared to ADAR1-Intact cells, to RSL3 at concentrations as

low as 50nM (**Figure 1A**). Cell viability assay was performed to determine IC$_{50}$ (half-maximal

inhibitory concentration) of RSL3, and ADAR1-Defective cells (IC$_{50}$ = ~0.11μM) were shown to

be more than 25-fold more sensitive compared to ADAR1-Intact cells (IC_{50} = ~2.9µM) (**Figure**

1B). Similar level of sensitization was observed when using another GPX4 inhibitor, ML162, to

treat ADAR1-Deficient (IC_{50} = ~0.08µM) and ADAR1-Intact cells (IC_{50} = ~2.27µM) (**Figure**

1C).

To evaluate if this sensitization effect occurs in TNBC cells besides MDA-MB-231, three other

TNBC cells lines (MDA-MB-468, MDA-MB-436, HCC1806) were transduced with shNT and

shADAR1 before being subjected to RSL3 treatment. In all three TNBC cell lines tested,

ADAR1 knockdown resulted in increased sensitivity to RSL3 ranging from 3.5-fold (HCC1806)

to over 25-fold (MDA-MB-468) (**Figure 1D-F**). Four non-TNBC cell lines (MCF7, T47D,

ZR75.1, SKBR3) were also examined for their sensitivity to RSL3 with or without ADAR1

knockdown. Interestingly, none of the non-TNBC cell lines tested acquired higher sensitivity to

RSL3 upon ADAR1 knockdown (**Figure S1G-J**). It indicates that ADAR1 regulates TNBC cell

lines' metabolic fitness by protecting them from ferroptosis induction.

An orthotopic implantation mouse model was established to test if these *in vitro* results can be

recapitulated *in vivo*. MDA-MB-231 cells stably transduced with Tet-inducible shADAR1 were

implanted into the mammary fat pad of female immunocompromised NSG mice (**Figure 1G and**

1H). Upon formation of palpable tumors 10 days post implantation, half of the mice were

switched to doxycycline-containing diet to induce expression of shADAR1 to reduce ADAR1

levels (**Figure S1K**). Both control- and doxycycline-diet mice were subjected to i.p. injection of

RSL3 (100mg/kg weight; twice a week) for 5 weeks before tumors were resected. Weekly health

and weight monitoring suggested no additional toxicity associated with RSL3 treatment (**Figure**

S1L). Doxycycline diet or RSL3 treatment alone reduced tumor burdens, and mice that received

the combination treatment (doxycycline diet + RSL3) experienced significantly slower tumor

progression (**Figure 1I-K, Figure S1M-O**). Immunohistochemical staining of 4-

Hydroxynonenal (4HNE), a product of lipid peroxidation, showed that the combination treatment

enhances ferroptosis phenotype in MDA-MB-231-formed mammary tumors (**Figure 1L**). These

results suggest that ADAR1 reduction and ferroptosis induction form synergy to suppress

tumorigenesis.

Knockdown of ADAR1 leads to lipid remodeling

Multiple pathways contribute to ferroptosis sensitivity, including ferrous iron availability,

glutathione activity, and relative abundance of polyunsaturated fatty acid (PUFA) over

monounsaturated fatty acid (MUFA)[6] (**Figure 2A**). Focusing on a cell line that can be highly

sensitized to ferroptosis induction by ADAR1 knockdown, MDA-MB-231, these elements were

evaluated to test the hypothesis that ADAR1-Deficient cells are primed to the cytotoxic effects of

ferroptosis induction.

In ADAR1-Defective cells, both the level of total glutathione and ratio between reduced and

total glutathione remain the same (**Figure S2A-B**). FerroOrange staining was used to detect

ferrous iron (Fe^{2+}) to show that ADAR1-Deficient MDA-MB-231 cells also do not display

significantly higher level of ferrous iron (**Figure S2C-D**). It suggests that the elevated sensitivity

to ferroptosis cannot be attributed to either alteration of glutathione antioxidant activity or

abundance of ferrous iron upon ADAR1 knockdown.

A non-targeted lipidomic analysis was performed to identify potential changes in lipid profiles

between ADAR1-Intact and ADAR1-Deficient cells. Interestingly, a clear pattern switch was

detected with the species of phosphatidylcholine (PC), the most abundant phospholipid in

mammalian cells and the most impactful phospholipid in controlling ferroptosis sensitivity

(**Figure 2B**). The observed lipid reprogramming disproportionally enriched PC species of longer

carbon chains with higher degrees of unsaturation indicated by the number of double bonds

within the carbon chain (**Figure 2C-F**). Some of the longest and most highly unsaturated PC

species, such as PC38:4, are among the most significantly enriched PCs in ADAR1-Defective

cells (**Figure 2G, Figure S2E**).

To investigate if similar lipid reprogramming occurs in TNBC cells sensitized by ADAR1

knockdown to a lesser extent, lipidomic analysis was performed to compare between ADAR1-

Intact and ADAR1-Deficient MDA-MB-436 cells (**Figure S2F**). Several highly unsaturated PCs,

such as PC 34:3, 35:3, and 44:6, were found to be enriched in ADAR1-Deficient MDA-MB-436

cells (**Figure 2H-I**). The enrichment upon ADAR1 knockdown, however, was not limited to

PUFA, as some MUFA PC species (i.e. PC36:1, 37:1, 38:1) were also upregulated in ADAR1-

Deficient cells. Other PUFA PC species, meanwhile, appeared to be reduced in ADAR1-

Deficient cells (**Figure S2G**).

Interestingly, PUFAs of other phospholipid families, such as phosphatidylserine (PS) and

phosphatidylethanolamine (PE), were also found to be disproportionally enriched in ADAR1-

Deficient MDA-MB-436 cells (**Figure 2J-L**, **Figure S2H-I**). These results showed that ADAR1

reduction leads to lipid reprogramming, specifically enrichment of PUFA over MUFA and

saturated fatty acid (SFA). It suggests that in TNBC cells, ADAR1 helps maintaining FA

homeostasis favoring MUFA and SFA. Interestingly, expression levels of ADAR1 strongly

correlate with abundance of SFA (PC32:0) and MUFA (PC32:1 and PC34:1) species in TNBC,

but not non-TNBC, cells according to the publicly available metabolomics dataset (**Figure S2J-**

L).

ADAR1 regulates MDM2 expression to dictate lipid profiles and ferroptosis sensitivity

To identify potential mechanisms that contribute to ADAR1-regulated lipid reprogramming in TNBC cells, four TNBC cell lines (MDA-MB-231, MDA-MB-468, HCC1806, BT549) were subjected to RNA-seq analyses to compare transcriptomic profiles between ADAR1-Intact and ADAR1-Deficient cells (**Figure S3A**).

Pathway analyses were performed to provide a glimpse of cellular functions impacted upon ADAR1 knockdown. Genes that were significantly altered in at least three cell lines tested can be categorized into a variety of pathways, including metabolic regulations in cancer cells as well as lipid and fatty acid metabolism (**Figure 3A-D, Figure S3B-C, Table S1-4**). Gene set enrichment analysis (GSEA) with RNA-seq data of MDA-MB-231 cells also suggests that ADAR1 reduction resulted in significant alterations in transcriptomes involved in fatty acid metabolism, but not in glutathione activity, iron metabolism, or direct ferroptosis regulation (**Figure 3E-F, Figure S3D-E**). GSEA results from MDA-MB-231 cells also suggest a lack of clear effect on interferon signaling gene sets, while cell cycle related transcriptome is significantly affected (**Figure S3F-G**). Interestingly, when other metabolic pathways were evaluated, ADAR1 knockdown was found to have no effect on the genet set associated with glycolysis, while significantly alter the transcriptome involved in amino acid metabolism (**Figure S3H-I**). It suggests that ADAR1's ability to regulate metabolic fitness in TNBC might extend beyond lipid programming.

To narrow down genetic targets for further investigation, more stringent cut-offs were applied to identify individual genes that are altered the most between ADAR1-Intact and ADAR1-Deficient cells. Fourteen genes (including *ADAR*, which encodes ADAR1) were downregulated in all four

TNBC cell lines tested, while 4 genes were upregulated (**Figure 3G-H**). Among these genes,

MDM2 has been shown to be subjected to ADAR1-mediated transcriptional regulation[28].

Induction of MDM2 protein level in ADAR1-Deficient cells was confirmed using western blot

analysis, and reduced RNA editing at the reported 3'UTR double-stranded editing site was

observed using both Sanger sequencing and RNA editing site-specific quantitative PCR (RESS-

qPCR) (**Figure 3I-K, Figure S3J-K**).

It was recently shown that MDM2 promotes ferroptosis sensitivity through mediating lipid

remodeling, including induction of PUFA levels[29]. To determine if MDM2 is also capable of

promoting ferroptosis sensitivity in TNBC cells, MDA-MB-231 cells were stably transfected

with a MDM2-overexpressing construct before being subjected to RSL3 treatment (**Figure 3L**).

Overexpression of MDM2 is sufficient to sensitize MDA-MB-231 cells to RSL3 treatment,

albeit to a lesser degree compared to the effect mediated by ADAR1 knockdown (**Figure 3M**).

To determine if MDM2 promotes enrichment of PUFAs in breast cancer cells, MDA-MB-231

cells overexpressing MDM2 were subjected to untargeted lipidomic analysis. A Tet-inducible

MDM2 construct was used in this experiment to identify profiles of PC PUFA at both 48 and 96

hours after doxycycline treatment (**Figure S3L**). At 48 hours post treatment, a few PC PUFAs

(i.e. PC34:3) were seen enriched (**Figure S3M**). After 96 hours of MDM2 overexpression,

however, a broader enrichment profile of PC PUFAs became prominent, including significant

inductions of several highly unsaturated PC species, such as PC42:7 and PC42:8, among others

(**Figure 3N-O, Figure S3N**).

To determine if the elevated ferroptosis sensitivity upon ADAR1 reduction can be indeed

attributed to MDM2 induction, a rescue experiment was conducted with RSL3-treated ADAR1-

Intact and ADAR1-Deficient MDA-MB-231 cells. The inflection-point concentration of RSL3

(100nM) distinguishing between ADAR1-Intact and ADAR1-Deficient MDA-MB-231 cells was

used in combination with MDM2 inhibitors. Cell viability analysis was performed to detect

reduced viability in RSL3-treated ADAR1-Deficient cells and effect of MDM2 inhibition. All

three MDM2 inhibitors used (MX69, SP-141, MEL23) were shown to partially rescue RSL3-

treated ADAR1-Deficient cells (**Figure 3P**). An agonist of peroxisome proliferator-activated

receptor alpha (PPAR-α), pirinixic acid, was also shown to confer mild yet significant rescue

effect, consistent with the hypothesis that MDM2 promotes ferroptosis sensitivity through

suppressing PPAR-α activity[29]. These results collectively indicate that ADAR1 regulates lipid

remodeling through MDM2 to protect TNBC cells from ferroptosis induction.

ADAR1 loss synergizes with cobimetinib to suppress tumorigenesis

The most commonly used ferroptosis inducers in preclinical studies, such as RSL3, remain

distant from clinical applications due to poor pharmacological profiles and excessive toxicity[30].

Meanwhile, many drugs currently being used in clinics or tested in clinical trials have been

shown to possess ferroptosis-inducing activities[31]. The opportunity exists, therefore, to harness

the lipid-remodeling effect of ADAR1 inhibition to enhance efficacy of treatments readily

available to patients.

To identify drug candidates synergizing with ADAR1 inhibition, a high-throughput screening

was conducted. Cell viability analysis was used to assess the effect of ADAR1 status on MDA-

MB-231 cells' sensitivity to drugs included in a ferroptosis-focused library (**Figure 4A**). Drugs

that synergize with ADAR1 inhibition (ADAR1-sensitized) were expected to suppress ADAR1-

Deficient cells more effectively than ADAR1-Intact cells similarly to RSL3, which was used as a

positive control in the screening (**Figure 4B-C**).

By comparing the relative viability between ADAR1-Deficient and -Intact cells, several drugs

were identified to react at least twice as strong with ADAR1-Deficient MDA-MB-231 cells

(**Figure 4D-E**). Siramesine is a lysosome disrupting agent that has been shown to induce

ferroptosis by promoting ROS (reactive oxygen species) – mediated oxidative stress and

reducing expression of ferroportin-1, an iron transport protein[32,33]. SB590885 is a potent inhibitor

of B-Raf, whose inhibition has been shown to induce oxidative stress and accumulation of

ROS[34,35]. UC2288 is a derivative of Sorafenib, a known ferroptosis inducer, and has been shown

to promote ferroptosis through regulating p21 activity[36,37]. Alpha-Mangostin is a dietary

xanthone that has also been shown to induce ROS activity[38]. Considering the effect of ADAR1

reduction on lipid remodeling, however, the synergy might work through compensating α-

Mangostin-mediated suppression of lipogenesis[39]. Similar mechanisms might be attributed to the

sensitization to Nordihydroguaiaretic acid (NDGA), a natural product inhibitor of 5-

lipoxygenase (5-LOX) that promotes the peroxidation of PUFA[40,41]. GW284543 and cobimetinib

are both MEK inhibitors, and MEK inhibition has been recently shown to induce ferroptosis in

lung and ovarian cancers[42,43].

Among the hits identified, cobimetinib is the only drug clinically available to treat FDA-

approved indications, including metastatic melanoma and histiocytic neoplasms[44,45]. Cobimetinib

is being evaluated in clinical trials for its efficacy in other cancers and has shown potential to

prolong survival of TNBC patients when combined with chemotherapy and immunotherapy[46].

Interestingly, publicly available drug response data show that cobimetinib displays dramatically

stronger activity towards TNBC compared to non-TNBC cell lines, similarly to carboplatin that

is known to improve TNBC treatment outcome[47] (**Figure S4A-B**).

The orthotopic implantation mouse model described earlier was performed to test the potential

synergistic effect between ADAR1 loss and cobimetinib *in vivo* (**Figure S4C**). MDA-MB-231

cells expressing Tet-inducible shADAR1 were implanted into the mammary fat pad of female

NSG mice. Upon formation of palpable tumors, mice were fed control or doxycycline-containing

diet before being subjected to cobimetinib treatment (oral administration at 20mg/kg weight;

twice a week) for 5 weeks before tumors were resected. Weekly health and weight monitoring

suggested no additional toxicity associated with cobimetinib treatment (**Figure S4D**). The

combination treatment of doxycycline diet and cobimetinib significantly reduce the tumor

burden compared to doxycycline diet or cobimetinib alone (**Figure 4F-H, Figure S4E-G**). It

indicates a strong synergistic effect between ADAR1 reduction and cobimetinib to suppress

tumorigenesis.

Discussion

Justified by data shown in this study, we expanded our knowledge of ADAR1-associated

vulnerabilities in TNBC by demonstrating their increased sensitivity to ferroptosis upon ADAR1

loss. Interestingly, two recent studies showed that ADAR1 loss can lead to overt ferroptosis

phenotypes, including in breast cancer cells[48,49]. It is unclear what caused the disparate

observations. Different elements of experimental methodology, such as cell culture condition,

can affect detection of ferroptosis phenotypes[50,51]. Notably, CRISPR (clustered regularly

interspaced short palindromic repeats) – mediated gene editing used to reduce ADAR1

expression in these studies has been shown to alter signaling pathways of tumor suppressor p53,

which can promote ferroptotic cell death[52,53]. Regardless of the reasons for these inconsistencies,

our results agree with the premise that ADAR1 plays a protective role against ferroptosis and

represents a novel target to sensitize cancer cells to ferroptosis-promoting agents.

ADAR1-mediated desensitization to ferroptosis was attributed to reprogramming of lipid

profiles, as ADAR1 reduction led to increased PUFA abundance and PUFA/MUFA ratio.

Transcriptomic analysis revealed protooncogene MDM2 as one of the commonly regulated

genes in ADAR1-Deficient TNBC cell lines that are sensitized to ferroptosis. Consistent with its

recently identified function in promoting ferroptosis through lipid reprogramming, alterations of

expression and activity of MDM2 were shown to connect to ADAR1-regulated ferroptosis

sensitivity[29]. Results presented in this manuscript depict a novel mechanism by which ADAR1

protects TNBC cells from stresses that could lead to ferroptosis (**Figure 4I**).

Sensitivity of TNBC cells to all three ferroptosis inducers used in this study (RSL3, ML162,

cobimetinib), interestingly, are inversely correlated with their dependency on ADAR1 (**Figure**

S4H-J). It suggests the possibility that TNBC cells' dependency on ADAR1 could be attributed

to its ability to protect them from ferroptosis induction, akin to the phenomenon of oncogene

addiction[54]. Although recent studies questioned ADAR1's status as a *bona fide* oncogene, it

undoubtably deserves to be considered a critical non-oncogene target for its ability to suppress

multiple stresses directly and indirectly[55,56]. We posit that the induced ferroptosis sensitivity in

ADAR1 knockdown TNBC cells is attributed to their intrinsic cellular environment taking

advantage of MDM2-facilitated remodeling of PUFA/MUFA ratio upon GPX4 inhibition. The

lack of ferroptosis induction with ADAR1 knockdown alone suggests that TNBC cells' lipid

profile is not the limiting factor, and the observed lipid remodeling serves as a priming event.

Further supporting the potential of this approach, recent studies revealed that TNBC are

protected from ferroptosis by MUFAs generated both intracellularly and by their originating

environment[57-59].

Preliminary results showed that non-TNBC cell lines are not sensitized to GPX4 inhibitor RSL3

upon ADAR1 knockdown. It will be interesting to investigate if the ADAR1-MDM2-PUFA axis

is also altered in these cells without the sensitization effect. It has been shown previously that

multiple mechanisms contributing to resistance of ferroptosis are elevated in non-TNBC breast

cancer cells, suggesting higher thresholds to sensitize them to ferroptosis inducers[60-64]. It is also

worth noting that TNBC disproportionally possess deactivating p53 mutations that abrogate

MDM2 interactions[65]. The ability of MDM2 to promote lipid remodeling and ferroptosis

sensitivity, therefore, could be dictated by its dynamic relationship with p53. The recent

discovery of mutant p53's role in protecting TNBC from ferroptosis indicates that MDM2 might

play an antagonistic role in promoting ferroptosis sensitivity in TNBC[66]. Moreover, Adams *et al.*

recently suggested that MDM2 is required for p53-inactivated TNBC cell survival[67]. While it is

not surprising with MDM2's many p53-independent pro-tumorigenic functions, it signals the

need to understand MDM2's complex role in ADAR1-targeting strategy against TNBC[68].

Exciting opportunities are highlighted by the synergistic effect between ADAR1 loss and

ferroptosis to identify novel combinatory cancer treatments. Besides the lack of clinically proven

ADAR1 and ferroptosis modulators, both ADAR1 inactivation and ferroptosis have been

associated with pathological consequences, handicapping the drug development progress[30,69]. In

addition to development of effective ADAR1 inhibitors, functional drug repurposing offers a

derisking strategy to bring biological insights closer to clinical applications[70,71]. A ferroptosis-

focused high-throughput drug screen identified cobimetinib, an FDA-approved MEK inhibitor

with ferroptosis-inducing activity, to be sensitized with ADAR1 loss. Cobimetinib and other

MEK inhibitors have been shown in both preclinical and clinical settings to inhibit TNBC

tumorigenesis[46,72]. In addition to its ability to promote ferroptosis, cobimetinib has been shown

to induce a type I interferon response[73]. It demonstrated the potential of repurposing drugs

coordinating with multiple signaling pathways downstream of ADAR1 inhibition to result in

superior efficacy against TNBC. It will be interesting to investigate if other existing TNBC

treatments, such as platinum-based agents, also synergize with ADAR1 inhibition with their

ferroptosis-promoting activity[74].

Raised by the new finding are many questions about the implication of ADAR1's ability to

regulate lipid metabolism and ferroptosis sensitivity. Does ADAR1 expression or its editing

profile have prognostic value, like recently shown androgen receptor, to predict TNBC patients'

response to ferroptosis inducing agents[75]? As both ferroptosis and ADAR1 inhibition can

sensitize "immune cold" tumors to immunotherapy, does a sequential or combined approach

targeting ADAR1, ferroptosis, and immune check points have the potential to minimize

treatment resistance[76,77]? Considering ADAR1's known role in regulating systemic metabolic

functions, does it protect TNBC from other metabolic stresses, such as amino acid

dysregulation[27]? And how do ADAR1-regulated metabolic adaptations fit into the development

of dietary interventions to improve the treatment of cancers, including TNBC[78-80]? Possessing

the capability to alter a quarter of the entire RNA species, ADAR1 holds immerse potential for

exploitation to develop innovative therapies against tough-to-treat cancers like TNBC.

Materials and Methods
384
Cell lines, reagents, and plasmids

Breast cancer cell lines (MDA-MB-231, MDA-MB-436, MDA-MB-468, HCC1806, BT549,

BT20, MCF7, T47D, ZR751, SKBR3) and human embryonic kidney 293T (HEK293T) cells

were obtained from American Tissue Cells Consortium (ATCC, Manassas, VA USA). All cancer

cell lines were maintained in Dulbecco's Modification of Eagle's Medium (DMEM, GE Life

Sciences, Pittsburgh, PA USA) supplemented with 10% fetal bovine serum (FBS, Gibco, 10091-

148), Sodium Pyruvate (Cellgro, Waltham, MA USA, 30-002-CI), Non-Essential Amino Acids

(NEA, Cellgro, 25-030-CI), and L-glutamine (Cellgro, 25-005-CI). Sensitivity of cells to

ferroptosis inducing agents were tested in defined medium with low serum using DMEM

supplemented with 2% FBS and L-glutamine. LipoFexin (Lamda Biotech, St. Louis, MO USA,

TS310) was used to generate lentivirus. Fugene 6 transfection reagent (Promega, E2692) was

used for other transfection experiments.

Tetracycline-inducible shADAR1 was generated by cloning shADAR1 oligonucleotides into

Tet-pLKO.1-puro construct with AgeI and EcoRI restriction enzyme sites. Tet-inducible MDM2-

overexpressing construct was generated by subcloning MDM2 open-reading frame (ORF) from

pCMV-myc3-MDM2 (Addgene 20935) into pSBtet-GP sleeping beauty vector (Addgene 60495)

using the Sfil restriction enzyme site. pSBtet-GP-MDM2 and pCMV(CAT)T7-SB100

transposase construct (Addgene 34879) were co-transfected into MDA-MB-231 cells to generate

a puromycin-resistant stable cell line. Commercially available chemicals and plasmids, as well as

primers and oligonucleotides used are listed in the Resources Table.

Lentiviral production and transduction

To generate lentivirus, HEK293T cells were transfected using LipoFexin with pCMV-VSV-G,

pCMV-ΔR8.2, and expression constructs (pLKO.1-puromycin backbone for short-hairpin

RNAs). Growth medium was replaced with fresh medium 24hr after transfection, and

supernatants containing lentivirus were harvested 24hr later. For transduction, one million cells

were infected with lentivirus for 24hr in the presence of 10 µg/ml protamine sulfate to facilitate

viral entry. Transduced cells were selected with puromycin (2 µg/ml) for at least 72hr. Sequences

of shRNAs are listed in the Resource table.

Cell viability assay

For cell viability assays, cells were plated at a density of 1,000 to 3,000 cells per well on a 96-

well plate and left to grow overnight in complete growth medium at 37°C. Cells were then

treated with the indicated drugs in equal volume at 2× desired concentration and left to incubate

for 72 h. Cells were then incubated with CellTiter-Glo® 2.0 Reagent (Promega, Madison, WI,

USA, G924C) for 1h at room temperature, and viability was read out according to the

manufacturer's protocol on a GloMax Navigator Microplate Luminometer (Promega, GM2000).

The half-maximal inhibitory concentration (IC_{50}) was determined using a nonlinear regression

model using Prism 10 software (GraphPad).

Lipid peroxidation assay; Glutathione activity assay; Ferrous iron assay

Biochemical assays were performed to evaluate ferroptosis phenotypes or molecular mechanisms

contributing to cells' sensitivity to ferroptosis induction. Level of lipid peroxidation was

determined by 1) measuring its natural byproduct malondialdehyde (MDA) using MDA

Colorimetric/Fluorometric Assay (Abcam, Cambridge, UK, ab118970), or by 2) staining and

comparing levels of reduced and oxidized forms of lipid peroxidation probe BODIPY^{TM} 581/591

C11 (Image-iT^{TM} Lipid Peroxidation Kit, Thermo Fisher Scientific, Waltham, MA USA),

according to manufacturer's instruction. Intracellular oxidative stress level was determined by

measuring the ratio between reduced (GSH) and oxidized (GSSG) forms of Glutathione using

GSH/GSSG Ratio Detection Assay (Abcam, ab205811) according to manufacturer's instruction.

Ferrous iron, which drives ferroptosis, was measured by FerroOrange staining (Dojindo, Japan,

F374) according to the manufacturer's instructions. Quantification of signal intensity was

performed using ImageJ software (NIH, Bethesda, MD USA).

Untargeted lipidomic analysis with reversed-phase liquid chromatography (RPLC)

Cells were washed with cold PBS 3 times, cold molecular biology grade water (Corning,

Glendale, AZ, 46-000-CM) 3 times, metabolically quenched with cold methanol for 5 min,

scraped harvested, and centrifuged at $10,000 \times g$ for 5 min to remove supernatant. Pellets were

dried using CentriVap Concentrator (Labconco, Kansas City, MO, 79700-00) and stored in -

80°C before extraction of metabolites. Cold extraction buffer containing

Methanol:Acetonitrile:Water at 2:2:1 ratio was added to the pellets. Samples were then subjected

to 3 times of 30-sec vortex → 1-min liquid nitrogen bath → 10-sec thaw → 10-min bath

sonication at 25°C (Fischer Scientific, CPXH Series Model 3800 Heated Ultrasonic Cleaner),

before being stored at -20°C for 2 hours to overnight. After centrifuging at 14000 RPM at 4°C

for 10 min, sample supernatants were transferred to new Eppendorf tubes and speed dried for 2

hours. BCA Protein Assay was performed on cell pellets in advance to determine protein

concentration. Isopropanol:Methanol at 1:1 ratio was added at 1µL per 2.5µg of proteins to the

residue and subjected to 2 times of 5-min bath sonication at 25°C → 1-min vortex, before being

stored at 4°C for 1 hour to overnight. After centrifuging at 14000 RPM at 4°C for 10 min,

supernatants were transferred to LC vials for LC/MS analysis.

No plastics were used to minimize palmitate contamination. An aliquot (4µL) of the extracted

sample was injected onto a Waters HSS T3 1.8µM, 2.1x150mm column with a 1.8µM,

2.1x50mm guard column connected to 1290 Infinity II LC. The columns were used with the

following buffers and gradients: Mobile Phase A = 40% water, 60% Acetonitrile, 0.1% Formic

Acid, 10mM Ammonium Formate, 2.5μM Medronic Acid; Mobile Phase B = 90% Isopropanol,

10% Acetonitrile, 0.1% Formic Acid, 10mM Ammonium Formate; Needle Wash = 70% Mobile

Phase A solution + 30% Mobile Phase B solution; Seal Wash = 9:1 Water:Isopropanol. All

solutions were mixed and degassed for 10 min before use. The binary pump was operated at 70%

A + 30% B from 0 - 17 min, 25% A + 75% B 0% B from 17 - 20 min, 15% A + 85% B from 20 -

23 min, 100% B from 23 - 27 min, and 70% A + 30% B thereafter at constant 0.25 mL/min flow

rate. MS detection was carried out using Agilent 6545 Q-TOF with 150-1,200 Da mass range.

An internal standard sample (Avanti Research, SPLASH® LIPIDOMIX® Mass Spec Standard,

330707) was used to normalize for injection variability. Lipid Annotator (Agilent) was used to

generate a target list for all lipids detected, and DecolD was used to identify polar lipids[81].

Bulk RNA sequencing and analysis

Total RNA was isolated from cells using miRNeasy Mini Kit (Qiagen, Hilden, Germany,

217004) including on-column DNase digestion following the manufacturer's protocol. Samples

were prepared according to RiboErase library kit (Roche) manufacturer's protocol, indexed,

pooled, and sequenced on an Illumina NovaSeq S4. Base calls and demultiplexing were

performed with Illumina's bcl2fastq software and a custom python demultiplexing program with

a maximum of one mismatch in the indexing read. RNA-seq reads were then aligned to the

Ensembl release 76 primary assembly with STAR version 2.5.1a1. Gene counts were derived

from the number of uniquely aligned unambiguous reads by Subread:featureCount version 1.4.6-

p52. Sequencing performance was assessed for the total number of aligned reads, total number of

uniquely aligned reads, and features detected. All gene counts were then imported into the

R/Bioconductor package EdgeR and TMM normalization size factors were calculated to adjust

for samples for differences in library size. Ribosomal genes and genes not expressed in the

smallest group size minus one sample greater than one count-per-million were excluded from

further analysis. The TMM size factors and the matrix of counts were then imported into the

R/Bioconductor package Limma. Weighted likelihoods based on the observed mean-variance

relationship of every gene and sample were then calculated for all samples with the

voomWithQualityWeights. The performance of all genes was assessed with plots of the residual

standard deviation of every gene to their average log-count with a robustly fitted trend line of the

residuals. Differential expression analysis was then performed to analyze for differences between

conditions and raw p-values and Benjamini-Hochberg false-discovery rate adjusted p-values

were determined to filter results based on cut-off. Biological and functional pathways analyses,

as well as gene set enrichment analyses were performed with the Enrichr platform, the

GeneOntology enrichment tools, and the Gene Set Enrichment Analysis (GSEA) software[82-87].

Immunoblotting analysis

Cell lysates were extracted from cells at ~90% confluence. Cells were washed with phosphate-

buffered saline (PBS, GE Life Sciences), scrape harvested, centrifuged at $1000 \times g$ for 5 min,

and lysed with RIPA buffer (20mM Tris-HCl pH7.5, 150mM NaCl, 1mM EDTA, 1% NP-40,

0.1% SDS, 0.1% Deoxycholate) supplemented with 10mM PMSF and HALT protease inhibitor

cocktail (Thermo Fisher Scientific, Waltham, MA USA). Lysates were clarified by

centrifugation and the protein concentration was determined using DC protein assay system

(Bio-Rad, Hercules, CA USA). Equal amount of protein was resolved by sodium dodecyl

sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) using Criterion TGX Stain-Free Precast

Gels (Bio-Rad) and transferred onto Immobilon-P membranes (MilliporeSigma, St. Louis, MO

USA). Primary antibodies used are listed in the Resource table. Secondary antibodies conjugated

to Horseradish peroxidase were used at a dilution of 1:5-10,000 (Jackson Immunochemicals,

West Grove, PA USA). Clarity Western ECL Substrate (Bio-Rad) was then applied to blots and

protein levels were detected using autoradiography with ChemiDoc XRS+ Imager (Bio-Rad).

RNA editing site-specific quantitative PCR (RESS-qPCR)

RNA editing fingerprint assay using RESS-qPCR was performed as previously described[88]. Total

RNA was isolated from lentivirus-transduced cells using RNeasy Plus Mini Kit including on-

column DNase digestion following the manufacturer's protocol. To eliminate any residual

genomic DNA contamination, TURBO DNA-free^{TM} Kit (ThermoFisher Scientific) was used to

further clean up isolated RNAs. RNAs free of genomic DNA were reverse transcribed to cDNAs

using the High-Capacity cDNA Reverse Transcription Kit (Life technologies, CA, USA).

Primers used for RESS-qPCR were previously published and are listed in the Resource table[19].

Quantitative RT-PCR was performed using iTaq Universal SYBR Green Supermix (Bio-Rad) on

the C1000 Thermal Cycler (Bio-Rad). The relative RNA editing level (Edit/Wild-Type RNA

ratio) was calculated as $2^{-(Ct\ Edit\ -\ Ct\ WT)}$.

Sanger sequencing to detect RNA editing

Isolated RNAs free of genomic DNA were used for Sanger sequencing analysis to verify RNA

editing events. PCR was performed using the high-fidelity Phusion PCR kit according to the

manufacturer's instructions (New England Biolabs, Ipswich, MA, E0553). "Outer" primers listed

in the Resource table were used to produce amplicons flanking the editing site about 75-125 bp

on either side to ensure successful sequencing analysis. Successful PCRs were verified by DNA

gel electrophoresis and amplicons were extracted from the gel using Monarch gel extraction kit

(BioRad, T1020S) according to manufacturer's instructions and submitted for sequencing

analysis (IDT Technologies). Sanger sequencing was performed using the outer primers (A/G

were read for WT/Edited if forward primers were used; T/C were read if reverse primers were

used) used for PCR, and sequence chromatograms were analyzed using Chromas software.

Drug screening with ferroptosis-focused library

MDA-MB-231 cells treated with shNT (ADAR1-Intact) or shADAR1 (ADAR1-Deficient) were

screened against the ferroptosis-focused drug library (MedChemExpress, Monmouth Junction,

NJ, HY-L051) at the High Throughput Screening Center (HTSC) at Washington University in

St. Louis. CellTiter-Glo® Luminescent Cell Viability Assay (Promega) was utilized to determine

cell viability upon drug treatment as previously described. Cells were prepared in defined

medium with low serum using DMEM supplemented with 2% FBS and L-glutamine.

All drugs in the library were pre-dispensed into 96-well sterile polypropylene plates (Corning,

Axyen) using a Hummingbird and subsequently diluted with media to produce intermediate

plates at 20µM and 2µM. Diluted compounds were then transferred using a BiomekFX

(Beckman Coulter) to 96-well white cell culture plates (Greiner Bio-One, Monroe, NC, 655083)

(50µl/well, 18 plates total with 9 plates for each experimental group. Select wells containing

DMSO alone served as controls. MDA-MB-231 cells treated with shNT (ADAR1-Intact) or

shADAR1 (ADAR1-Deficient) were seeded (15,000 cells in 50µl cell culture media per well)

using the Multidrop 384 dispenser (Thermo Fisher Scientific, 24291). Plates were kept at room

temperature for 30 min before being moved to the incubator (37°C, 5% CO_2) for 24h. On day 2,

CellTiter-Glo® Viability Assays were performed using the integrated and automated platform

(Beckman Coulter), with library plate-matched shNT and shADAR1 samples alternating

between screening steps: Cell plates were equilibrated at room temperature for 30 min in a

Cytomat MPH incubator. CellTiter-Glo® reagent (100µl/well) was added with a Multidrop 384

dispenser. Plates were shaken at 250 rpm for 2 min with a Teleshake followed by room

temperature incubation for 1h in a Cytomat MPH incubator. Luminescence was measured (1

sec/well after a 1 min delay) with a Perkin Elmer Envision plate reader. To monitor assay

performance, Coefficient of variations (CVs) and Z'-factors for DMSO-treated and no-cell

control samples were determined. All screening plates were confirmed to display high quality

assay metrics (CVs < 10%; Z'-factors > 0.8). All test wells were normalized to no-treatment

samples on the same plate. Relative cell viability was compared between shADAR1 and shNT

samples, and the cut-off was set at 51% to discriminate hits from no-hits.

Mammary fat pad orthotopic implantation

Immuno-deficient NOD scid gamma (NSG) female mice at 6-8 weeks old were purchased from

Jackson Laboratory (Bar Harbor, ME, USA, 005557) and placed on Teklad Global 18% protein

rodent diet (Envigo, Indianapolis, USA, T.2918.15) 3 days before implantation. 1.5×10^5 MDA-

MB-231 cells were harvested and resuspended in PBS, mixed with Matrigel® Matrix Basement

Membrane (Corning, Waltham, USA, 354230) at 1:1 volume ratio, and kept at 4°C until

implantation. In total 100µl of cells-Matrigel solutions were injected into the right inguinal

mammary glands of NSG mice. Ten days after the implantation, palpable tumors were detected

and half of the mice were switched to doxycycline-containing diet (Envigo, 625mg/kg,

TD.01306). Tumors were measured using a caliper once per week, and tumor volumes were

calculated using the following formula: $L \times W^2 \times 0.52$ (L and W represents the large and small

diameter of the tumor, respectively). Mice were euthanized 6 weeks post implantation or before

tumors reached 2 cubic cm in size. Tumors were dissected from the mice for weight

measurement and subsequently fixed in 10% formalin, before transferred to 70% ethanol for

processing to prepare formalin-fixed paraffin-embedded (FFPE) tissue blocks. All animal-related

experimental procedures were performed in compliance with the guidelines given by the

American Association for Accreditation for Laboratory Animal Care and the U.S. Public Health

Service Policy on Human Care and Use of Laboratory Animals. All animal studies were

approved by the Washington University Institutional Animal Care and Use Committee (IACUC)

in accordance with the Animal Welfare Act and NIH guidelines (Protocol 20220144).

Immunohistochemistry

Tumor FFPE tissue blocks were sectioned at 5μm thickness. Hematoxylin and eosin (H&E)

staining was performed for histological evaluations. For immunohistochemistry, sections were

stained using a Bond RXm autostainer (Leica, Buffalo Grove, IL USA). Primary antibodies for

ADAR1-p150 (Abcam, ab126745) and 4-Hydroxynonenal (4HNE, ThermoFisher Scientific,

MA5-27570) were diluted 1:100 in Antibody Diluent (Leica). Briefly, slides were baked at 65°C

for 15min and automated software performed dewaxing, rehydration, antigen retrieval, blocking,

primary antibody incubation, post primary antibody incubation, detection (DAB), and

counterstaining using Bond reagents (Leica). Samples were then removed from the machine,

dehydrated through ethanols and xylenes, mounted and cover slipped before imaging.

Bioinformatic analysis and data availability

Bioinformatic analyses to assess relationships among gene expression, drug response, gene

dependency and metabolomics in breast cancer cell lines were performed using the DepMap

public data (depmap.org) with the TNBC (HCC1187, HS578T, HCC1599, DU4475, BT549,

HCC1143, BT20, MDA-MB-436, HCC1806, HCC70, HCC2157, HCC1395, MDA-MB-231,

MDA-MB-468, MDA-MB-453) and non-TNBC (SKBR3, MCF7, KPL1, MDA-MB-134VI,

ZR751, EFM192A, T47D, AU565, HCC1419, EFM19, HCC1500, HCC1428, UACC893,

UACC812, HCC202, HCC2218, MDA-MB-175VII, BT483, ZR7530, MDA-MB-415, BT474,

HCC1569, MDA-MB-361) groups[89]. The results published here are in whole or part based upon

data generated by Cancer Target Discovery and Development (CTD²) Network

(https://ccg.cancer.gov/ccg/research/functional-genomics/ctd2/data-portal) established by the

National Cancer Institute's Center for Cancer Genomics. RNA-seq data from this study are

available at the Gene Expression Omnibus under accession number GSE286185.

Statistical analysis

All *in vitro* and *in vivo* data are reported as the mean ± SD unless stated otherwise, and statistical

analyses were performed using GraphPad Prism 10. The two-tailed unpaired student t test was

performed for experiments comparing between two experimental groups. One-way ANOVA

with Tukey's multiple comparisons was used for statistical analysis of all other experiments

unless stated otherwise. Tumor volume analyses from *in vivo* studies were performed by 1)

comparing AUC (area under curve) between experimental groups using one-way ANOVA, and

2) comparing weekly measured tumor volumes between groups using two-way ANOVA with

Tukey's multiple comparison. Pearson correlation coefficient and/or Spearman's rank correlation

coefficient were calculated to assess correlations between two data sets. Statistical significance

was evaluated using p-value, which is shown in the numerical format. Lower p-values reject the

null hypothesis more strongly, and p-value <0.05 is generally considered statistically significant.

Acknowledgements

We thank the Division of Comparative Medicine at Washington University in St. Louis for

housing the mouse colony. We thank the Denardo lab for processing the fixed tumor tissues. We

thank the Tissue Histology Core, Department of Obstetrics and Gynecology, for sectioning the

tumor samples and performing H&E staining. We thank the support by the Genome Technology

Access Center (GTAC) at McDonnell Genome Institute (MGI) Symposium Pilot Project

Funding to perform RNA-seq analyses (Catrina Fronick and Michael Heinz for logistics; Toni

Sinnwell, Elliott Klotz, and Eric Tycksen for quality control, sequencing and analyses). We

acknowledge the Just-In-Time (JIT) Core Usage Funding Program to support the collaboration

with HTSC for drug screening. Both GTAC@MGI Symposium Pilot Project and JIT programs

are funded through the Institute of Clinical and Translational Sciences (ICTS), Washington

University in St. Louis. ICTS was supported by the National Center for Advancing Translational

Sciences (NCATS) grant UL1TR002345. This work was supported by R01CA262804 (JDW)

from the National Cancer Institute, and W81XWH-21-1-0476 and W81XWH-21-1-0466 from

the Congressionally Directed Medical Research Programs (CDMRP), Department of Defense

(JDW). L.S.T. is supported by R25HG006687 and T32GM148405 from National Human

Genome Research Institute and National Institute of General Medical Sciences, respectively.

Author Contributions

C-P. Kung: Conceptualization, investigation, methodology, formal analysis, project

administration, writing–original draft, writing–review and editing. **N.D. Terzich:** investigation,

methodology, formal analysis. **MXG. Ilagen:** investigation, methodology, formal analysis,

writing–review and editing. **M.J. Prinsen:** Investigation, methodology, formal analysis. **M.**

Kaushal: Investigation, methodology, formal analysis. **R.D. Kladney:** Investigation,

methodology. **J.H. Weber:** Investigation. **A.R. Mabry:** Investigation. **L.S. Torres:**

Investigation. **E.R. Bramel:** Investigation. **E.C. Freeman:** Investigation. **T. Sabloak:**

Investigation. **K.A. Cottrell:** Investigation. **S. Ryu:** Investigation. **W.M. Weber:** Investigation.

L. Maggi Jr.: investigation, writing–review and editing. **L.P. Shriver:** investigation,

methodology, writing–review and editing. **G.J. Patti:** funding acquisition, investigation,

writing–review and editing. **J.D. Weber:** Conceptualization, resources, supervision, funding

acquisition, writing–original draft, writing–review and editing.

Declaration of Interests

G.J. Patti is the Chief Scientific Officer of Panome Bio, a scientific advisory board member for

Cambridge Isotope Laboratories, and has a collaborative research agreement with Agilent

Technologies.

References

1. Ademuyiwa, F.O., Tao, Y., Luo, J., Weilbaecher, K., and Ma, C.X. (2017). Differences in the mutational landscape of triple-negative breast cancer in African Americans and Caucasians. Breast Cancer Res Treat *161*, 491-499. 10.1007/s10549-016-4062-y.

2. Obidiro, O., Battogtokh, G., and Akala, E.O. (2023). Triple Negative Breast Cancer Treatment Options and Limitations: Future Outlook. Pharmaceutics *15*. 10.3390/pharmaceutics15071796.

3. Loizides, S., and Constantinidou, A. (2022). Triple negative breast cancer: Immunogenicity, tumor microenvironment, and immunotherapy. Front Genet *13*, 1095839. 10.3389/fgene.2022.1095839.

4. Zhou, L., and Yu, C.W. (2024). Epigenetic modulations in triple-negative breast cancer: Therapeutic implications for tumor microenvironment. Pharmacol Res *204*, 107205. 10.1016/j.phrs.2024.107205.

5. Munkacsy, G., Santarpia, L., and Gyorffy, B. (2023). Therapeutic Potential of Tumor Metabolic Reprogramming in Triple-Negative Breast Cancer. Int J Mol Sci *24*. 10.3390/ijms24086945.

6. Stockwell, B.R. (2022). Ferroptosis turns 10: Emerging mechanisms, physiological functions, and therapeutic applications. Cell *185*, 2401-2421. 10.1016/j.cell.2022.06.003.

7. Timmerman, L.A., Holton, T., Yuneva, M., Louie, R.J., Padro, M., Daemen, A., Hu, M., Chan, D.A., Ethier, S.P., van 't Veer, L.J., et al. (2013). Glutamine sensitivity analysis identifies the xCT antiporter as a common triple-negative breast tumor therapeutic target. Cancer Cell *24*, 450-465. 10.1016/j.ccr.2013.08.020.

8. Beatty, A., Fink, L.S., Singh, T., Strigun, A., Peter, E., Ferrer, C.M., Nicolas, E., Cai, K.Q., Moran, T.P., Reginato, M.J., et al. (2018). Metabolite Profiling Reveals the Glutathione Biosynthetic Pathway as a Therapeutic Target in Triple-Negative Breast Cancer. Mol Cancer Ther *17*, 264-275. 10.1158/1535-7163.MCT-17-0407.

9. Verma, N., Vinik, Y., Saroha, A., Nair, N.U., Ruppin, E., Mills, G., Karn, T., Dubey, V., Khera, L., Raj, H., et al. (2020). Synthetic lethal combination targeting BET uncovered intrinsic susceptibility of TNBC to ferroptosis. Sci Adv *6*. 10.1126/sciadv.aba8968.

10. Pinnix, Z.K., Miller, L.D., Wang, W., D'Agostino, R., Jr., Kute, T., Willingham, M.C., Hatcher, H., Tesfay, L., Sui, G., Di, X., et al. (2010). Ferroportin and iron regulation in breast cancer progression and prognosis. Sci Transl Med *2*, 43ra56. 10.1126/scitranslmed.3001127.

11. Doll, S., Proneth, B., Tyurina, Y.Y., Panzilius, E., Kobayashi, S., Ingold, I., Irmler, M., Beckers, J., Aichler, M., Walch, A., et al. (2017). ACSL4 dictates ferroptosis sensitivity by shaping cellular lipid composition. Nat Chem Biol *13*, 91-98. 10.1038/nchembio.2239.

12. Beatty, A., Singh, T., Tyurina, Y.Y., Tyurin, V.A., Samovich, S., Nicolas, E., Maslar, K., Zhou, Y., Cai, K.Q., Tan, Y., et al. (2021). Ferroptotic cell death triggered by conjugated linolenic acids is mediated by ACSL1. Nat Commun *12*, 2244. 10.1038/s41467-021-22471-y.

13. Lorito, N., Subbiani, A., Smiriglia, A., Bacci, M., Bonechi, F., Tronci, L., Romano, E., Corrado, A., Longo, D.L., Iozzo, M., et al. (2024). FADS1/2 control lipid metabolism and ferroptosis susceptibility in triple-negative breast cancer. EMBO Mol Med *16*, 1533-1559. 10.1038/s44321-024-00090-6.

14. Fumagalli, D., Gacquer, D., Rothe, F., Lefort, A., Libert, F., Brown, D., Kheddoumi, N., Shlien, A., Konopka, T., Salgado, R., et al. (2015). Principles Governing A-to-I RNA

699   Editing in the Breast Cancer Transcriptome. Cell Rep *13*, 277-289.
700   10.1016/j.celrep.2015.09.032.

701 15. Peng, X., Xu, X., Wang, Y., Hawke, D.H., Yu, S., Han, L., Zhou, Z., Mojumdar, K., Jeong,
702   K.J., Labrie, M., et al. (2018). A-to-I RNA Editing Contributes to Proteomic Diversity in
703   Cancer. Cancer Cell *33*, 817-828 e817. 10.1016/j.ccell.2018.03.026.

704 16. Han, L., Diao, L., Yu, S., Xu, X., Li, J., Zhang, R., Yang, Y., Werner, H.M.J., Eterovic,
705   A.K., Yuan, Y., et al. (2015). The Genomic Landscape and Clinical Relevance of A-to-I
706   RNA Editing in Human Cancers. Cancer Cell *28*, 515-528. 10.1016/j.ccell.2015.08.013.

707 17. Paz-Yaacov, N., Bazak, L., Buchumenski, I., Porath, H.T., Danan-Gotthold, M.,
708   Knisbacher, B.A., Eisenberg, E., and Levanon, E.Y. (2015). Elevated RNA Editing
709   Activity Is a Major Contributor to Transcriptomic Diversity in Tumors. Cell Rep *13*, 267-
710   276. 10.1016/j.celrep.2015.08.080.

711 18. Baker, A.R., Miliotis, C., Ramirez-Moya, J., Marc, T., Vlachos, I.S., Santisteban, P., and
712   Slack, F.J. (2022). Transcriptome Profiling of ADAR1 Targets in Triple-Negative Breast
713   Cancer Cells Reveals Mechanisms for Regulating Growth and Invasion. Mol Cancer Res
714   *20*, 960-971. 10.1158/1541-7786.MCR-21-0604.

715 19. Kung, C.P., Cottrell, K.A., Ryu, S., Bramel, E.R., Kladney, R.D., Bao, E.A., Freeman,
716   E.C., Sabloak, T., Maggi, L., Jr., and Weber, J.D. (2021). Evaluating the therapeutic
717   potential of ADAR1 inhibition for triple-negative breast cancer. Oncogene *40*, 189-202.
718   10.1038/s41388-020-01515-5.

719 20. Ishizuka, J.J., Manguso, R.T., Cheruiyot, C.K., Bi, K., Panda, A., Iracheta-Vellve, A.,
720   Miller, B.C., Du, P.P., Yates, K.B., Dubrot, J., et al. (2019). Loss of ADAR1 in tumours
721   overcomes resistance to immune checkpoint blockade. Nature *565*, 43-48.
722   10.1038/s41586-018-0768-9.

723 21. Gannon, H.S., Zou, T., Kiessling, M.K., Gao, G.F., Cai, D., Choi, P.S., Ivan, A.P.,
724   Buchumenski, I., Berger, A.C., Goldstein, J.T., et al. (2018). Identification of ADAR1
725   adenosine deaminase dependency in a subset of cancer cells. Nat Commun *9*, 5450.
726   10.1038/s41467-018-07824-4.

727 22. Liu, H., Golji, J., Brodeur, L.K., Chung, F.S., Chen, J.T., deBeaumont, R.S., Bullock, C.P.,
728   Jones, M.D., Kerr, G., Li, L., et al. (2019). Tumor-derived IFN triggers chronic pathway
729   agonism and sensitivity to ADAR loss. Nat Med *25*, 95-102. 10.1038/s41591-018-0302-5.

730 23. Karki, R., Sundaram, B., Sharma, B.R., Lee, S., Malireddi, R.K.S., Nguyen, L.N.,
731   Christgen, S., Zheng, M., Wang, Y., Samir, P., et al. (2021). ADAR1 restricts ZBP1-
732   mediated immune response and PANoptosis to promote tumorigenesis. Cell Rep *37*,
733   109858. 10.1016/j.celrep.2021.109858.

734 24. Zhang, T., Yin, C., Fedorov, A., Qiao, L., Bao, H., Beknazarov, N., Wang, S., Gautam, A.,
735   Williams, R.M., Crawford, J.C., et al. (2022). ADAR1 masks the cancer
736   immunotherapeutic promise of ZBP1-driven necroptosis. Nature *606*, 594-602.
737   10.1038/s41586-022-04753-7.

738 25. Mehdipour, P., Marhon, S.A., Ettayebi, I., Chakravarthy, A., Hosseini, A., Wang, Y., de
739   Castro, F.A., Loo Yau, H., Ishak, C., Abelson, S., et al. (2020). Epigenetic therapy induces
740   transcription of inverted SINEs and ADAR1 dependency. Nature *588*, 169-173.
741   10.1038/s41586-020-2844-1.

742 26. McGuirk, S., Audet-Delage, Y., and St-Pierre, J. (2020). Metabolic Fitness and Plasticity
743   in Cancer Progression. Trends Cancer *6*, 49-61. 10.1016/j.trecan.2019.11.009.

27. Kung, C.P., Maggi, L.B., Jr., and Weber, J.D. (2018). The Role of RNA Editing in Cancer
Development and Metabolic Disorders. Front Endocrinol (Lausanne) *9*, 762.
10.3389/fendo.2018.00762.
28. Yang, C.C., Chen, Y.T., Chang, Y.F., Liu, H., Kuo, Y.P., Shih, C.T., Liao, W.C., Chen,
H.W., Tsai, W.S., and Tan, B.C. (2017). ADAR1-mediated 3' UTR editing and expression
control of antiapoptosis genes fine-tunes cellular apoptosis response. Cell Death Dis *8*,
e2833. 10.1038/cddis.2017.12.
29. Venkatesh, D., O'Brien, N.A., Zandkarimi, F., Tong, D.R., Stokes, M.E., Dunn, D.E.,
Kengmana, E.S., Aron, A.T., Klein, A.M., Csuka, J.M., et al. (2020). MDM2 and MDMX
promote ferroptosis by PPARalpha-mediated lipid remodeling. Genes Dev *34*, 526-543.
10.1101/gad.334219.119.
30. Liu, J., Kang, R., and Tang, D. (2024). Adverse effects of ferroptotic therapy: mechanisms
and management. Trends Cancer *10*, 417-429. 10.1016/j.trecan.2024.01.002.
31. Zhao, L., Zhou, X., Xie, F., Zhang, L., Yan, H., Huang, J., Zhang, C., Zhou, F., Chen, J.,
and Zhang, L. (2022). Ferroptosis in cancer and cancer immunotherapy. Cancer Commun
(Lond) *42*, 88-116. 10.1002/cac2.12250.
32. Ma, S., Henson, E.S., Chen, Y., and Gibson, S.B. (2016). Ferroptosis is induced following
siramesine and lapatinib treatment of breast cancer cells. Cell Death Dis *7*, e2307.
10.1038/cddis.2016.208.
33. Ostenfeld, M.S., Fehrenbacher, N., Hoyer-Hansen, M., Thomsen, C., Farkas, T., and
Jaattela, M. (2005). Effective tumor cell death by sigma-2 receptor ligand siramesine
involves lysosomal leakage and oxidative stress. Cancer Res *65*, 8975-8983.
10.1158/0008-5472.CAN-05-0269.
34. Gagliardi, M., Cotella, D., Santoro, C., Cora, D., Barlev, N.A., Piacentini, M., and
Corazzari, M. (2019). Aldo-keto reductases protect metastatic melanoma from ER stress-
independent ferroptosis. Cell Death Dis *10*, 902. 10.1038/s41419-019-2143-7.
35. Haq, R., Shoag, J., Andreu-Perez, P., Yokoyama, S., Edelman, H., Rowe, G.C., Frederick,
D.T., Hurley, A.D., Nellore, A., Kung, A.L., et al. (2013). Oncogenic BRAF regulates
oxidative metabolism via PGC1alpha and MITF. Cancer Cell *23*, 302-315.
10.1016/j.ccr.2013.02.003.
36. Gao, F., Zhang, B., Sun, Z., Gao, Y., Liu, C., Dou, X., Tong, H., and Wang, R. (2023).
Regulation of endothelial ferroptosis by SESN1 in atherosclerosis and its related
mechanism. Aging (Albany NY) *15*, 5052-5065. 10.18632/aging.204777.
37. Dixon, S.J., Patel, D.N., Welsch, M., Skouta, R., Lee, E.D., Hayano, M., Thomas, A.G.,
Gleason, C.E., Tatonetti, N.P., Slusher, B.S., and Stockwell, B.R. (2014). Pharmacological
inhibition of cystine-glutamate exchange induces endoplasmic reticulum stress and
ferroptosis. Elife *3*, e02523. 10.7554/eLife.02523.
38. Lee, C.H., Ying, T.H., Chiou, H.L., Hsieh, S.C., Wen, S.H., Chou, R.H., and Hsieh, Y.H.
(2017). Alpha-mangostin induces apoptosis through activation of reactive oxygen species
and ASK1/p38 signaling pathway in cervical cancer cells. Oncotarget *8*, 47425-47439.
10.18632/oncotarget.17659.
39. Shi, Y., Fan, Y., Hu, Y., Jing, J., Wang, C., Wu, Y., Geng, Q., Dong, X., Li, E., and Dong,
D. (2020). alpha-Mangostin suppresses the de novo lipogenesis and enhances the
chemotherapeutic response to gemcitabine in gallbladder carcinoma cells via targeting the
AMPK/SREBP1 cascades. J Cell Mol Med *24*, 760-771. 10.1111/jcmm.14785.

40. Gilbert, N.C., Gerstmeier, J., Schexnaydre, E.E., Borner, F., Garscha, U., Neau, D.B.,
790   Werz, O., and Newcomer, M.E. (2020). Structural and mechanistic insights into 5-
791   lipoxygenase inhibition by natural products. Nat Chem Biol *16*, 783-790. 10.1038/s41589-
792   020-0544-7.

793   41. Sun, Q.Y., Zhou, H.H., and Mao, X.Y. (2019). Emerging Roles of 5-Lipoxygenase
794   Phosphorylation in Inflammation and Cell Death. Oxid Med Cell Longev *2019*, 2749173.
795   10.1155/2019/2749173.

796   42. Bhatt, V., Lan, T., Wang, W., Kong, J., Lopes, E.C., Wang, J., Khayati, K., Raju, A.,
797   Rangel, M., Lopez, E., et al. (2023). Inhibition of autophagy and MEK promotes ferroptosis
798   in Lkb1-deficient Kras-driven lung tumors. Cell Death Dis *14*, 61. 10.1038/s41419-023-
799   05592-8.

800   43. Yin, J., Chen, J., Hong, J.H., Huang, Y., Xiao, R., Liu, S., Deng, P., Sun, Y., Chai, K.X.Y.,
801   Zeng, X., et al. (2024). 4EBP1-mediated SLC7A11 protein synthesis restrains ferroptosis
802   triggered by MEK inhibitors in advanced ovarian cancer. JCI Insight *9*.
803   10.1172/jci.insight.177857.

804   44. Larkin, J., Ascierto, P.A., Dreno, B., Atkinson, V., Liszkay, G., Maio, M., Mandala, M.,
805   Demidov, L., Stroyakovskiy, D., Thomas, L., et al. (2014). Combined vemurafenib and
806   cobimetinib in BRAF-mutated melanoma. N Engl J Med *371*, 1867-1876.
807   10.1056/NEJMoa1408868.

808   45. Diamond, E.L., Durham, B.H., Ulaner, G.A., Drill, E., Buthorn, J., Ki, M., Bitner, L., Cho,
809   H., Young, R.J., Francis, J.H., et al. (2019). Efficacy of MEK inhibition in patients with
810   histiocytic neoplasms. Nature *567*, 521-524. 10.1038/s41586-019-1012-y.

811   46. Brufsky, A., Kim, S.B., Zvirbule, Z., Eniu, A., Mebis, J., Sohn, J.H., Wongchenko, M.,
812   Chohan, S., Amin, R., Yan, Y., et al. (2021). A phase II randomized trial of cobimetinib
813   plus chemotherapy, with or without atezolizumab, as first-line treatment for patients with
814   locally advanced or metastatic triple-negative breast cancer (COLET): primary analysis.
815   Ann Oncol *32*, 652-660. 10.1016/j.annonc.2021.01.065.

816   47. Han, H.S., Vikas, P., Costa, R.L.B., Jahan, N., Taye, A., and Stringer-Reasor, E.M. (2023).
817   Early-Stage Triple-Negative Breast Cancer Journey: Beginning, End, and Everything in
818   Between. Am Soc Clin Oncol Educ Book *43*, e390464. 10.1200/EDBK_390464.

819   48. Yin, C., Zhang, M.M., Wang, G.L., Deng, X.Y., Tu, Z., Jiang, S.S., Gao, Z.D., Hao, M.,
820   Chen, Y., Li, Y., and Yang, S.Y. (2024). Loss of ADAR1 induces ferroptosis of breast
821   cancer cells. Cell Signal *121*, 111258. 10.1016/j.cellsig.2024.111258.

822   49. He, D., Niu, C., Bai, R., Chen, N., and Cui, J. (2024). ADAR1 Promotes Invasion and
823   Migration and Inhibits Ferroptosis via the FAK/AKT Pathway in Colorectal Cancer. Mol
824   Carcinog. 10.1002/mc.23818.

825   50. Murray, M.B., Leak, L.B., Lee, W.C., and Dixon, S.J. (2023). Protocol for detection of
826   ferroptosis in cultured cells. STAR Protoc *4*, 102457. 10.1016/j.xpro.2023.102457.

827   51. Leu, J.I., Murphy, M.E., and George, D.L. (2019). Mechanistic basis for impaired
828   ferroptosis in cells expressing the African-centric S47 variant of p53. Proc Natl Acad Sci
829   U S A *116*, 8390-8396. 10.1073/pnas.1821277116.

830   52. Enache, O.M., Rendo, V., Abdusamad, M., Lam, D., Davison, D., Pal, S., Currimjee, N.,
831   Hess, J., Pantel, S., Nag, A., et al. (2020). Cas9 activates the p53 pathway and selects for
832   p53-inactivating mutations. Nat Genet *52*, 662-668. 10.1038/s41588-020-0623-4.

53. Gnanapradeepan, K., Basu, S., Barnoud, T., Budina-Kolomets, A., Kung, C.P., and
Murphy, M.E. (2018). The p53 Tumor Suppressor in the Control of Metabolism and
Ferroptosis. Front Endocrinol (Lausanne) *9*, 124. 10.3389/fendo.2018.00124.

54. Pagliarini, R., Shao, W., and Sellers, W.R. (2015). Oncogene addiction: pathways of
therapeutic response, resistance, and road maps toward a cure. EMBO Rep *16*, 280-296.
10.15252/embr.201439949.

55. Nagel, R., Semenova, E.A., and Berns, A. (2016). Drugging the addict: non-oncogene
addiction as a target for cancer therapy. EMBO Rep *17*, 1516-1531.
10.15252/embr.201643030.

56. Mendez Ruiz, S., Chalk, A.M., Goradia, A., Heraud-Farlow, J., and Walkley, C.R. (2023).
Over-expression of ADAR1 in mice does not initiate or accelerate cancer formation in
vivo. NAR Cancer *5*, zcad023. 10.1093/narcan/zcad023.

57. Xie, Y., Wang, B., Zhao, Y., Tao, Z., Wang, Y., Chen, G., and Hu, X. (2022). Mammary
adipocytes protect triple-negative breast cancer cells from ferroptosis. J Hematol Oncol *15*,
72. 10.1186/s13045-022-01297-1.

58. Ackermann, T., Shokry, E., Deshmukh, R., Anand, J., Galbraith, L.C.A., Mitchell, L.,
Rodriguez-Blanco, G., Villar, V.H., Sterken, B.A., Nixon, C., et al. (2024). Breast cancer
secretes anti-ferroptotic MUFAs and depends on selenoprotein synthesis for metastasis.
EMBO Mol Med *16*, 2749-2774. 10.1038/s44321-024-00142-x.

59. Puente-Cobacho, B., Esteo, C., Altea-Manzano, P., Garcia-Perez, J.L., Quiles, J.L.,
Sanchez-Rovira, P., Martín-Salvago, M.D., Molina-Jiménez, L., Luque, R.J., Fendt, S.-M.,
and Vera-Ramirez, L. (2024). De novo lipogenesis protects dormant breast cancer cells
from ferroptosis and promotes metastasis. Redox Biology, 103480.
https://doi.org/10.1016/j.redox.2024.103480.

60. Park, S.Y., Jeong, K.J., Poire, A., Zhang, D., Tsang, Y.H., Blucher, A.S., and Mills, G.B.
(2023). Irreversible HER2 inhibitors overcome resistance to the RSL3 ferroptosis inducer
in non-HER2 amplified luminal breast cancer. Cell Death Dis *14*, 532. 10.1038/s41419-
023-06042-1.

61. Bajbouj, K., Shafarin, J., and Hamad, M. (2019). Estrogen-dependent disruption of
intracellular iron metabolism augments the cytotoxic effects of doxorubicin in select breast
and ovarian cancer cells. Cancer Manag Res *11*, 4655-4668. 10.2147/CMAR.S204852.

62. Li, X., Rhee, D.K., Malhotra, R., Mayeur, C., Hurst, L.A., Ager, E., Shelton, G., Kramer,
Y., McCulloh, D., Keefe, D., et al. (2016). Progesterone receptor membrane component-1
regulates hepcidin biosynthesis. J Clin Invest *126*, 389-401. 10.1172/JCI83831.

63. Liu, L., Zhang, C., Qu, S., Liu, R., Chen, H., Liang, Z., Tian, Z., Li, L., Ma, S., and Liu,
X. (2022). ESR1 inhibits ionizing radiation-induced ferroptosis in breast cancer cells via
the NEDD4L/CD71 pathway. Arch Biochem Biophys *725*, 109299.
10.1016/j.abb.2022.109299.

64. Wang, H., Hou, M.J., Liao, L., Li, P., Chen, T., Wang, P., and Zhu, B.T. (2024). Strong
Protection by 4-Hydroxyestrone against Erastin-Induced Ferroptotic Cell Death in
Estrogen Receptor-Negative Human Breast Cancer Cells: Evidence for Protein Disulfide
Isomerase as a Mechanistic Target for Protection. Biochemistry *63*, 984-999.
10.1021/acs.biochem.3c00261.

65. Cancer Genome Atlas, N. (2012). Comprehensive molecular portraits of human breast
tumours. Nature *490*, 61-70. 10.1038/nature11412.

66. Dibra, D., Xiong, S., Moyer, S.M., El-Naggar, A.K., Qi, Y., Su, X., Kong, E.K., Korkut,
A., and Lozano, G. (2024). Mutant p53 protects triple-negative breast adenocarcinomas
from ferroptosis in vivo. Sci Adv *10*, eadk1835. 10.1126/sciadv.adk1835.
67. Adams, C.M., Mitra, R., Xiao, Y., Michener, P., Palazzo, J., Chao, A., Gour, J., Cassel, J.,
Salvino, J.M., and Eischen, C.M. (2023). Targeted MDM2 Degradation Reveals a New
Vulnerability for p53-Inactivated Triple-Negative Breast Cancer. Cancer Discov *13*, 1210-
1229. 10.1158/2159-8290.CD-22-1131.
68. Kung, C.P., and Weber, J.D. (2022). It's Getting Complicated-A Fresh Look at p53-
MDM2-ARF Triangle in Tumorigenesis and Cancer Therapy. Front Cell Dev Biol *10*,
818744. 10.3389/fcell.2022.818744.
69. Cottrell, K.A., Torres, L.S., Dizon, M.G., and Weber, J.D. (2021). 8-azaadenosine and 8-
chloroadenosine are not selective inhibitors of ADAR. Cancer Res Commun *1*, 56-64.
10.1158/2767-9764.crc-21-0027.
70. Weth, F.R., Hoggarth, G.B., Weth, A.F., Paterson, E., White, M.P.J., Tan, S.T., Peng, L.,
and Gray, C. (2024). Unlocking hidden potential: advancements, approaches, and obstacles
in repurposing drugs for cancer therapy. Br J Cancer *130*, 703-715. 10.1038/s41416-023-
02502-9.
71. Rehwinkel, J., and Mehdipour, P. (2024). ADAR1: from basic mechanisms to inhibitors.
Trends Cell Biol. 10.1016/j.tcb.2024.06.006.
72. Lee, J., Lim, B., Pearson, T., Choi, K., Fuson, J.A., Bartholomeusz, C., Paradiso, L.J.,
Myers, T., Tripathy, D., and Ueno, N.T. (2019). Anti-tumor and anti-metastasis efficacy of
E6201, a MEK1 inhibitor, in preclinical models of triple-negative breast cancer. Breast
Cancer Res Treat *175*, 339-351. 10.1007/s10549-019-05166-3.
73. Lulli, D., Carbone, M.L., and Pastore, S. (2017). The MEK Inhibitors Trametinib and
Cobimetinib Induce a Type I Interferon Response in Human Keratinocytes. Int J Mol Sci
18. 10.3390/ijms18102227.
74. Jennis, M., Kung, C.P., Basu, S., Budina-Kolomets, A., Leu, J.I., Khaku, S., Scott, J.P.,
Cai, K.Q., Campbell, M.R., Porter, D.K., et al. (2016). An African-specific polymorphism
in the TP53 gene impairs p53 tumor suppressor function in a mouse model. Genes Dev *30*,
918-930. 10.1101/gad.275891.115.
75. Yang, F., Xiao, Y., Ding, J.H., Jin, X., Ma, D., Li, D.Q., Shi, J.X., Huang, W., Wang, Y.P.,
Jiang, Y.Z., and Shao, Z.M. (2023). Ferroptosis heterogeneity in triple-negative breast
cancer reveals an innovative immunotherapy combination strategy. Cell Metab *35*, 84-100
e108. 10.1016/j.cmet.2022.09.021.
76. Bhate, A., Sun, T., and Li, J.B. (2019). ADAR1: A New Target for Immuno-oncology
Therapy. Mol Cell *73*, 866-868. 10.1016/j.molcel.2019.02.021.
77. Deng, J., Zhou, M., Liao, T., Kuang, W., Xia, H., Yin, Z., Tan, Q., Li, Y., Song, S., Zhou,
E., and Jin, Y. (2022). Targeting Cancer Cell Ferroptosis to Reverse Immune Checkpoint
Inhibitor Therapy Resistance. Front Cell Dev Biol *10*, 818453. 10.3389/fcell.2022.818453.
78. Ferrer, M., Mourikis, N., Davidson, E.E., Kleeman, S.O., Zaccaria, M., Habel, J., Rubino,
R., Gao, Q., Flint, T.R., Young, L., et al. (2023). Ketogenic diet promotes tumor ferroptosis
but induces relative corticosterone deficiency that accelerates cachexia. Cell Metab *35*,
1147-1162 e1147. 10.1016/j.cmet.2023.05.008.
79. Salvadori, G., Zanardi, F., Iannelli, F., Lobefaro, R., Vernieri, C., and Longo, V.D. (2021).
Fasting-mimicking diet blocks triple-negative breast cancer and cancer stem cell escape.
Cell Metab *33*, 2247-2259 e2246. 10.1016/j.cmet.2021.10.008.

80. Martinez-Garay, C., and Djouder, N. (2023). Dietary interventions and precision nutrition in cancer therapy. Trends Mol Med *29*, 489-511. 10.1016/j.molmed.2023.04.004.

81. Koelmel, J.P., Li, X., Stow, S.M., Sartain, M.J., Murali, A., Kemperman, R., Tsugawa, H., Takahashi, M., Vasiliou, V., Bowden, J.A., et al. (2020). Lipid Annotator: Towards Accurate Annotation in Non-Targeted Liquid Chromatography High-Resolution Tandem Mass Spectrometry (LC-HRMS/MS) Lipidomics Using A Rapid and User-Friendly Software. Metabolites *10*. 10.3390/metabo10030101.

82. Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., and Mesirov, J.P. (2005). Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A *102*, 15545-15550. 10.1073/pnas.0506580102.

83. Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S., Eppig, J.T., et al. (2000). Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. Nat Genet *25*, 25-29. 10.1038/75556.

84. Gene Ontology, C., Aleksander, S.A., Balhoff, J., Carbon, S., Cherry, J.M., Drabkin, H.J., Ebert, D., Feuermann, M., Gaudet, P., Harris, N.L., et al. (2023). The Gene Ontology knowledgebase in 2023. Genetics *224*. 10.1093/genetics/iyad031.

85. Xie, Z., Bailey, A., Kuleshov, M.V., Clarke, D.J.B., Evangelista, J.E., Jenkins, S.L., Lachmann, A., Wojciechowicz, M.L., Kropiwnicki, E., Jagodnik, K.M., et al. (2021). Gene Set Knowledge Discovery with Enrichr. Curr Protoc *1*, e90. 10.1002/cpz1.90.

86. Chen, E.Y., Tan, C.M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G.V., Clark, N.R., and Ma'ayan, A. (2013). Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. BMC Bioinformatics *14*, 128. 10.1186/1471-2105-14-128.

87. Mootha, V.K., Lindgren, C.M., Eriksson, K.F., Subramanian, A., Sihag, S., Lehar, J., Puigserver, P., Carlsson, E., Ridderstrale, M., Laurila, E., et al. (2003). PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. Nat Genet *34*, 267-273. 10.1038/ng1180.

88. Crews, L.A., Jiang, Q., Zipeto, M.A., Lazzari, E., Court, A.C., Ali, S., Barrett, C.L., Frazer, K.A., and Jamieson, C.H. (2015). An RNA editing fingerprint of cancer stem cell reprogramming. J Transl Med *13*, 52. 10.1186/s12967-014-0370-3.

89. Tsherniak, A., Vazquez, F., Montgomery, P.G., Weir, B.A., Kryukov, G., Cowley, G.S., Gill, S., Harrington, W.F., Pantel, S., Krill-Burger, J.M., et al. (2017). Defining a Cancer Dependency Map. Cell *170*, 564-576 e516. 10.1016/j.cell.2017.06.010.

Figure 1. ADAR1 protects TNBC cells from ferroptosis induction

A) Light micrographs of ADAR1-Intact (shNT) and ADAR1-Deficient (shADAR1) MDA-MB-231 TNBC cells treated with different concentrations of RSL3 for 24h. Cell viability analyses were performed to determine IC_{50} values of ferroptosis inducers **B)** RSL3 and **C)** ML162 for ADAR1-Intact and ADAR1-Deficient MDA-MB-231 cells. IC_{50} values of RSL3 were then determined for ADAR1-Intact and ADAR1-Deficient **D)** MDA-MB-436, **E)** HCC1806, and **F)** MDA-MB-468 TNBC cells. Cell viability graphs are representatives of three independent experiments. FC, fold-change. **G)** Immunoblot analysis showed similar levels of ADAR1 knockdown in doxycycline-treated MDA-MB-231 cells possessing Tet-inducible shADAR1 compared to MDA-MB-231 cells treated with shADAR1-containing lentivirus. Images are representative of three replicates. GAPDH, loading control. **H)** MDA-MB-231 cells (1.5 X 10^5 per mouse) possessing Tet-inducible shADAR1 were injected into mammary fat pads of female NSG mice through orthotopic implantations. Ten days post implantation, half of the mice were switched to doxycycline-containing (625mg/kg) diet. Two days after the diet switch, both control- and doxycycline-diet mice were subjected to RSL3 treatments through i.p. injection (100mg/kg weight; twice a week) for 5 weeks before tumors were resected. **I)** Progression of overall tumor volumes, n=5-9 each group. Error bars mark standard deviation. **J)** Quantification of areas under curve (AUC) for overall tumor volumes from I). Error bars mark standard deviation. **K)** Two-way ANOVA analysis to compare weekly measured overall tumor volumes. Statistically significant p-values (<0.05) were highlighted in red. **L)** Representative images of IHC staining of H&E, ADAR1 (p150 isoform), and 4HNE in tumor samples resected from each treatment group. Scale bar, 200μm. See also **Figure S1**.

Figure 2: ADAR1 reduction leads to lipid remodeling

A) A simplified illustration highlighting key regulators of ferroptosis pathway. GSH, Glutathione. PL, phospholipid. MUFA, monounsaturated fatty acid. PUFA, polyunsaturated fatty acid. PM, plasma membrane. **B)** Heatmap of lipidomic analysis showed that ADAR1-knockdown results in lipid remodeling in MDA-MB-231 cells. PC, phosphatidylcholines. **C)** Relative abundance of PC species shown in B), listed in the order of carbon chain length. Values were log10 transformed, mean centered, and shown as mean over standard deviation ratio for each PC species. n=5. Red dots denote ADAR1-Intact (shNT), and blue dots denote ADAR1-Deficient (shADAR1) MDA-MB-231 samples. **D)** Relative abundance of PC species listed in the order of lipid unsaturation. n=5. **E)** Enrichment of PC species with different carbon chain lengths (28-30, 31-33, and >33 carbons) in ADAR1-Deficient over ADAR1-Intact MDA-MB-231 cells. **F)** Enrichment of PC species with different unsaturation levels (SFA, 0 double-bond; MUFA, 1 double-bond; PUFA, 2-3 double-bond; PUFA2, at least 4 double-bond) in ADAR1-Deficient over ADAR1-Intact MDA-MB-231 cells. **G)** Relative abundance of PC38:4 between ADAR1-Intact and -Deficient MDA-MB-231 cells. Left, detected peak intensity normalized to internal standard (ITSD). Right, log10 transformed and mean normalized peak intensity. n=5 **H)** Relative abundance of selected PC species in ADAR1-Intact and -Deficient MDA-MB-436 cells. Values were log10 transformed, mean centered, and shown as mean over standard deviation ratio. n=3. **I)** Relative abundance of PC44:6 between ADAR1-Intact and -Deficient MDA-MB-436 cells.

Detected peak intensity was log10 transformed and mean normalized. n=3. **J)** Relative
abundance of PE and PS species in ADAR1-Intact and -Deficient MDA-MB-436 cells, listed in
the order of lipid unsaturation. Values were log10 transformed, mean centered, and shown as
mean over standard deviation ratio. n=3. **K)** Enrichment of PE and PS species with different
unsaturation levels (MUFA, 1 double-bond; PUFA, 2-3 double-bond; PUFA2, at least 4 double-
bond) in ADAR1-Deficient over ADAR1-Intact MDA-MB-436 cells. **L)** Relative abundance of
PE38:5 between ADAR1-Intact and -Deficient MDA-MB-436 cells. Detected peak intensity was
log10 transformed and mean normalized. n=3. See also **Figure S2**.

Figure 3: ADAR1 regulates MDM2 expression to dictate ferroptosis sensitivity

A) Venn diagram of significantly downregulated genes in ADAR1-Deficient over -Intact TNBC
cell lines (MDA-MB-231, MDA-MB-468, HCC1806, BT549). **B)** Venn diagram of significantly
upregulated genes in ADAR1-Deficient over -Intact TNBC cell lines (MDA-MB-231, MDA-
MB-468, HCC1806, BT549). **C)** The ten most significantly altered signaling pathways
(BioPlanet_2019) in ADAR1-Deficient over -Intact TNBC cell lines. Genes significantly altered
(both up- and down-regulated) in at least 3 out of 4 cell lines tested are included in the analysis.
D) The ten most significantly altered disease-related biological functions (PheWeb_2019) in
ADAR1-Deficient over -Intact TNBC cell lines. Genes significantly altered in at least 3 out of 4
cell lines tested are included in the analysis. **E)** GSEA enrichment plot of ferroptosis related
genes in ADAR1-Deficient MDA-MB-231 cells. **F)** GSEA enrichment plot of fatty acid
metabolism genes in ADAR1-Deficient MDA-MB-231 cells. **G)** Venn diagram of the most
significantly downregulated genes (log2<-0.4) in ADAR1-Deficient over -Intact TNBC cell
lines. Genes altered in all 4 cell lines are listed on the right. **H)** Venn diagram of the most
significantly upregulated genes (log2>0.4) in ADAR1-Deficient over -Intact TNBC cell lines.
Genes altered in all 4 cell lines are listed on the right. **I)** Immunoblot analysis showed significant
induction of MDM2 protein levels in ADAR1-Deficient (shADAR1) MDA-MB-231 cells.
Images are representative of three replicates. GAPDH, loading control. **J)** Reported ADAR1
editing site (bolded and underlined) in MDM2 3'UTR. Numbers mark the location within
MDM2 open reading frame. **K)** Reduced editing of MDM2 in ADAR1-Deficient TNBC cells.
T/C were read with reverse primers in Sanger sequencing to indicate WT (T) and edited (C)
RNA. Ratio of C/T peak heights indicates level of editing. Sequence chromatograms showed are
representative of 3 independent experiments. **L)** Immunoblot analysis showed overexpression of
MDM2 in MDA-MB-231 cells. Anti-MDM2 antibody detected both endogenous and
overexpressed MDM2, and anti-Myc-tag antibody detected overexpressed MDM2 containing
Myc-tag. GAPDH, loading control. ADAR1 level was not affected by MDM2 overexpression.
M) Cell viability analysis was performed to determine IC_{50} values of ferroptosis inducers RSL3
for MDA-MB-231 cells with (MDM2) or without (Ctrl) MDM2 overexpression. Cell viability
graph is representative of three independent experiments. FC, fold-change. **N)** Relative
abundance of PUFA PC species, listed in the order of lipid unsaturation, in MDA-MB-231 cells
with (red dots) or without (blue dots) MDM2 overexpression for 96 h. Values were log10
transformed, mean centered, and shown as mean over standard deviation ratio for each PC
species. n=5. **O)** Relative abundance of PC42:7 between control (Ctrl) and MDM2-
overexpressing MDA-MB-231 cells. Detected peak intensity was log10 transformed and mean

normalized. n=5. **P)** Relative cell viability within each of ADAR1-Intact (shNT) and -Deficient
(shADAR1) MDA-MB-231 cells treated with RSL3 (100nM), as well as ADAR1-Deficient
MDA-MB-231 cells co-treated with RSL3 and MDM2 inhibitors MX69 (10µM), MEL23
(10µM), and SP-141 (10µM), or PPAR-α agonist pirinixic acid (PA; 10µM) for 72h. n=3. Error
bars mark standard error. See also **Figure S3 and Table S1-4**.

Figure 4: ADAR1 loss synergizes with cobimetinib to suppress tumorigenesis

A) Drug screen designed to identify ferroptosis modulators synergizing with ADAR1 loss. **B)**
Anticipated results in identifying ADAR1-sensitized ferroptosis modulators. Small molecules
(Exampled by blue dots) that display ADAR1-sensitized activity against cancer cells will reduce
viability of ADAR1-Deficient (shADAR1) cells more strongly compared to ADAR1-Intact
(shNT) cells. **C)** Screening result with a ferroptosis-focused library (654 small molecules) in
MDA-MB-231 cells. X and Y axis represent cell viability for ADAR1-Intact (shNT) and -
Deficient (shADAR1) cells after treatment, respectively. Data were normalized to cells treated
with DMSO. RSL3 treatment was included as positive control for ADAR1-sensitized hits (red
dots). Dashed line denotes expected locations for molecules without discriminating activity
based on ADAR1 expression. **D)** Cell viability ratio of 0.51 (horizontal red line) between
shADAR1 and shNT cells was used as the cutoff to identify ADAR1-sensitized hits (red dots).
RSL3 treatment was included as positive control. **E)** List of hits identified from C) and D),
ranked by cell viability ratio between treated shADAR1 and shNT MDA-MB-231 cells. **F)**
Progression of overall tumor volumes throughout the treatment regimen, n=5-10 each group.
Error bars mark standard deviation. **G)** Quantification of areas under curve (AUC) for overall
tumor volumes from F). Error bars mark standard deviation. **H)** Two-way ANOVA analysis to
compare weekly measured overall tumor volumes. Statistically significant p-values (<0.05) were
highlighted in red. **I)** ADAR1 protects TNBC cells from ferroptosis through regulating MDM2
and lipid compositions between MUFA and PUFA. MUFA, monounsaturated fatty acid. PUFA,
polyunsaturated fatty acid. FIN, ferroptosis inducer. See also **Figure S4**.

Figure S1. ADAR1 protects TNBC cells from ferroptosis induction

A) Immunoblot analysis showed significant reduction of ADAR1 protein levels in MDA-MB-
231 cells treated with shADAR1-containing lentivirus. Images are representative of three
replicates. GAPDH, loading control. **B)** Light and fluorescent images showed the morphology
(Bright Field) and reduced and oxidized forms of the lipid peroxidation probe BODIPY^{TM}
581/591 C11 in ADAR1-Intact (shNT) and ADAR1-Deficient (shADAR1) MDA-MB-231 cells.
C) Quantification of ratios of reduced over oxidized forms of BODIPY^{TM} 581/591 C11 in shNT
and shADAR1 MDA-MB-231 cells. n=3. Error bars mark standard error. **D)** Concentration of
the lipid peroxidation marker MDA in ADAR1-Intact and -Deficient MDA-MB-231 cells. n=3.
Error bars mark standard deviation. **E)** Relative cell viability of ADAR1-Intact and -Deficient
MDA-MB-231 cells, as well as ADAR1-Deficient MDA-MB-231 cells treated with apoptosis
inhibitor Z-VAD-FMK (50µM), eIF2α inhibitor ISRIB (10µM), and ferroptosis inhibitors Fer-1
(Ferrostatin-1, 1µM) and Lip-1 (Liproxstatin-1, 1µM) for 72h. n=3 independent experiments.
Error bars mark standard error. **F)** Light micrographs of MDA-MB-231 cells treated with RSL3
(10µM), in combination with DMSO, apoptosis inhibitor Z-VAD-FMK (5µM), necroptosis

inhibitor Nec-1 (Necrostatin-1, 5µM), and ferroptosis inhibitors Fer-1 (Ferrostatin-1, 5µM) and
UAMC (5µM) for 24h. IC_{50} values of RSL3 were then determined for ADAR1-Intact and
ADAR1-Deficient **G)** MCF7, **H)** ZR75.1, **I)** T47D and **J)** SKBR3 non-TNBC cells. Cell
viability graphs are representatives of two independent experiments. **K)** Immunoblot analysis
showed reduction of ADAR1 protein levels in lysates extracted from newly formed tumors from
MDA-MB-231 cells possessing Tet-inducible shADAR1, 7 days after implanted mice (M1 and
M2) were switched to doxycycline diet. GAPDH, loading control. **L)** Progression of average
mouse weight between treatment groups, n=5-9 each group. Error bars mark standard deviation.
M) Progression of relative tumor volumes, normalized to tumor volumes measured at week 0,
throughout the treatment regimen, n=5-9 each group. Error bars mark standard deviation. **N)**
Two-way ANOVA analysis compares weekly measured relative tumor volumes. Statistically
significant p-values (<0.05) were highlighted in red. **O)** Representative images of tumors
resected after the treatment regimens concluded. Scale bar, 5mm.

Figure S2: ADAR1 reduction leads to lipid remodeling

A) Concentration of Glutathione (GSH) in ADAR1-Intact (shNT) and -Deficient (shADAR1)
MDA-MB-231 cells. n=3. Error bars mark standard deviation. **B)** Ratio of reduced/total levels of
GSH to indicate GSH activity in ADAR1-Intact and -Deficient MDA-MB-231 cells. **C)** Light
and fluorescent images showed the morphology (Bright Field) and ferrous ion detection
(FerroOrange) in ADAR1-Intact and -Deficient MDA-MB-231 cells, respectively. **D)**
Quantification of ferrous ion detection shown in C), normalized to cell numbers. n=3. Error bars
mark standard error **E)** Significantly altered PC species (red dots) in ADAR1-Deficient
compared to ADAR1-Intact MDA-MB-231 cells. PUFA PC species are marked. **F)** Immunoblot
analysis showed significant reduction of ADAR1 protein levels in MDA-MB-436 cells treated
with shADAR1-containing lentivirus. Images are representative of three replicates. GAPDH,
loading control. **G)** Relative abundance of all detected PC species in ADAR1-Intact (red dots)
and -Deficient (blue dots) MDA-MB-436 cells. Values were log10 transformed, mean centered,
and shown as mean over standard deviation ratio. n=3. **H)** Relative abundance of PE and PS
species in ADAR1-Intact and -Deficient MDA-MB-436 cells, listed in the order of carbon chain
length. Values were log10 transformed, mean centered, and shown as mean over standard
deviation ratio. n=3**. I)** Relative abundance of all detected PC, PE, and PS species in ADAR1-
Intact and -Deficient MDA-MB-436 cells, listed in the order of lipid unsaturation. Values were
log10 transformed, mean centered, and shown as mean over standard deviation ratio. n=3.
Abundance of **J)** PC32:0, **K)** PC32:1, and **L)** PC34:1 positively correlate with *ADAR* (encoding
ADAR1) expression level in TNBC (n=13), but not non-TNBC (n=23), cell lines. Data are
extracted from publicly available metabolomics data set from DepMap (depmap.com).

Figure S3: ADAR1 regulates MDM2 expression to dictate ferroptosis sensitivity

A) RNA sequencing to identify significantly altered genes in ADAR1-Deficient TNBC cells. **B)**
GeneOntology enrichment analysis for genes significantly altered in ADAR1-Deficient MDA-
MB-231 cells, using PANTHER Overrepresentation Fisher's Exact Test. **C)** The ten most
significantly altered biological pathways (Elsevier Collection Pathway) in ADAR1-Deficient
over -Intact TNBC cell lines. Genes significantly altered (both up- and down-regulated) in at

least 3 out of 4 cell lines tested are included in the analysis. GSEA enrichment analyses were performed to assess regulated pathways such as **D)** Glutathione metabolism; **E)** Iron uptake and transport; **F)** Cell cycle; **G)** Interferon signaling; **H)** Glycolysis; **I)** Amino acid metabolism in ADAR1-Deficient MDA-MB-231 cells. **J)** Predicted RNA secondary structure of the entire MDM2 3'UTR and the reported ADAR1 editing site (circled in red in the zoomed-in square). **K)** RESS-qPCR showed that editing of MDM2 3'UTR site was reduced in ADAR1-Deficient MDA-MB-231 and MDA-MB-468 cells. n=3. Error bars mark standard error. **L)** Immunoblot analysis showed doxycycline-induced overexpression of MDM2 in MDA-MB-231 cells after 48 and 96 h. GAPDH, loading control. **M)** Relative abundance of PUFA PC species, listed in the order of lipid unsaturation, in MDA-MB-231 cells with (red dots) or without (blue dots) MDM2 overexpression for 48 h. Values were log10 transformed, mean centered, and shown as mean over standard deviation ratio for each PC species. n=5. **N)** Significantly enriched PUFA PC species (red dots) in MDM2-overexpressing (96h) compared to control MDA-MB-231 cells.

Figure S4: ADAR1 loss synergizes with cobimetinib to suppress tumorigenesis

A) Cobimetinib shows superior efficacy against TNBC (n=7) over non-TNBC (n=10) cell lines. Efficacy numbers (PRISM Repurposing Public 24Q2) <0 indicate reduced viability upon treatment. Data are extracted from publicly available data set from DepMap (depmap.com). **B)** Cobimetinib (TNBC: -0.72; non-TNBC: 0.71) shows superior preference, compared to carboplatin (TNBC: -0.75; non-TNBC: -0.18), against TNBC (n=7) over non-TNBC (n=10) cell lines. **C)** MDA-MB-231 cells (1.5×10^5 per mouse) possessing Tet-inducible shADAR1 were injected into mammary fat pads of female NSG mice through orthotopic implantations. Ten days post implantation, half of the mice were switched to doxycycline-containing (625mg/kg) diet. Two days after the diet switch, both control- and doxycycline-diet mice were subjected to cobimetinib treatments through oral administration (20mg/kg weight; twice a week) for 5 weeks before tumors were resected. **D)** Progression of average mouse weight between treatment groups, n=5-10 each group. Error bars mark standard deviation. **E)** Progression of relative tumor volumes, normalized to tumor volumes measured at week 0, throughout the treatment regimen, n=5-10 each group. Error bars mark standard deviation. **F)** Two-way ANOVA analysis to compare weekly measured relative tumor volumes. Statistically significant p-values (<0.05) were highlighted in red. **G)** Representative images of tumors resected after the treatment regimens concluded. Scale bar, 5mm. Drug sensitivity of TNBC, but not non-TNBC, cell lines to **H)** RSL3 (TNBC, n=9; non-TNBC, n=10), **I)** ML162 (TNBC, n=9; non-TNBC, n=11), and **J)** cobimetinib (TNBC, n=7; non-TNBC, n=7) negatively correlate with *ADAR* (encoding ADAR1) dependency (Chronos CRISPR Gene Effect). Lower AUC (RSL3, ML162) and efficacy (cobimetinib) numbers indicate higher drug sensitivity. Lower CRISPR Gene Effect numbers indicate higher *ADAR* dependency. Data are extracted from publicly available drug sensitivity (RSL3, ML162: Drug sensitivity AUC, CTD^2; cobimetinib: PRISM Repurposing Public 24Q2) and gene dependency (Public 24Q2+Score, Chronos) data sets from DepMap (depmap.com).

Figure 1

Figure 3

Figure 4