Rhizosphere-colonizing bacteria persist in the protist microbiome

Stephen. J Taerum[a], Ravikumar R. Patel[a], Justin E. Alamo[a], Daniel Gage[b], Blaire Steven[c], and Lindsay R.

Triplett[a]#

a. Department of Plant Pathology and Ecology, the Connecticut Agricultural Experiment Station, New

Haven, CT 06511, U.S.A.

b. Department of Molecular and Cell Biology, the University of Connecticut, Storrs, CT 06269, U.S.A.

c. Department of Environmental Science and Forestry, the Connecticut Agricultural Experiment Station,

New Haven, CT 06511, U.S.A.

Running title: Rhizosphere colonization by protist-associated bacteria

#Address correspondence to Lindsay R. Triplett, Lindsay.Triplett@ct.gov.

Abstract

Soils contain diverse predatory protists that affect the abundance and behavior of rhizosphere

bacteria, including bacteria that may benefit plant health. Protists also harbor their own bacterial

microbiomes, including transient and extracellular associates, but it is not known whether the protist

microbiome affects the plant rhizosphere. To address this question, we profiled the bacterial

microbiomes of eight evolutionarily diverse rhizosphere protist isolates after two years of continuous

laboratory culture. We then compared the protist culture microbiomes to maize rhizosphere

communities six weeks after protist inoculation. Introduction of protists enriched 13 protist-associated

bacterial amplicon sequence variants (ASVs) in the rhizosphere, which comprised ~10% of the

rhizosphere bacterial community. An additional 17 bacterial ASVs ranked highly in abundance in both

rhizosphere and protist microbiomes; together, a median 47% of the protist culture bacterial community

was enriched or abundant in the rhizosphere. Inoculation with some of the protist cultures positively

affected root biomass traits, but a protist mixture had no effect, indicating that the impact of protist

holobionts on plant growth is context-dependent. Isolates of protist-associated bacteria had both

positive and negative effects on protist growth in culture, suggesting that the bacteria use multiple

strategies to survive in proximity to predators. This study demonstrates that even after prolonged

laboratory culture, evolutionarily diverse rhizosphere protists host bacterial microbiomes dominated by

plant-colonizing bacteria that impact the rhizosphere microbiome after inoculation. The findings suggest

that protists may contribute to the rhizosphere as part of the soil microbial seedbank, and identify

bacterial groups that may be important to the plant-protist interaction.

Importance

Understanding the impact of predatory protists on the plant microbiome will be essential to

deploy protists in sustainable agriculture. This study shows that eight rhizosphere protist isolates hosted

diverse and distinct bacterial communities, and that a large proportion of these bacteria could be found

colonizing the maize root environment long after protists were introduced. This study demonstrates that

maize rhizosphere bacteria can persist for years in the protist microbiome, indicating that extremely

diverse eukaryotes could help select and maintain rhizosphere bacteria in the absence of the plant.

Introduction

Microbial eukaryotes including fungi, nematodes, and protists are integral to the plant microbiome.

Single-celled protists- eukaryotes that cannot be classified as plant, animal, or fungus- are ubiquitous in

soil, and are often predators of bacteria and fungi (1). Protists may stimulate plant growth, and this was

historically often attributed to the stimulation of nutrient cycling (2). Consensus has grown that

predators also benefit plant growth and disease resilience through selective predation. Culling prey

bacteria can increase the competitive fitness of predation-resistant bacteria, including plant growth

promoting strains of *Bacillus*, *Sphingomonas, Pseudomonas*, and *Azospirillum* (3). Although few species

of rhizosphere bacteria are known to benefit from protists, network association studies have identified

hundreds of positive correlations among protist and bacterial groups in the rhizosphere (4–6). This

suggests that a much wider diversity of rhizosphere bacteria may have positive ecological associations

with predators than is currently established. Given that correlations could arise from niche constraints,

environmental factors, or random chance (7), systematic studies are needed to understand which

rhizosphere-colonizing bacteria are positively selected by protists.

Rhizosphere eukaryotes interact with bacteria, but they also host their own bacterial microbiomes

that shape their interaction with plants. For example, nematode endosymbionts detoxify plant

secondary metabolites, endofungal *Rhizobium* and *Burkholderia* spp. promote plant beneficial and

pathogenic fungi, and *Fusarium*-colonizing *Pantoea* inhibit virulence by producing antifungal compounds

(8–11). Highly cosmopolitan multi-kingdom colonists, including some strains of *Bacillus* and

Pseudomonas bacteria, may even be transferred from fungi or nematodes to plants (12, 13). Protists

host intracellular and extracellular bacterial communities through a wide range of mutualistic and

antagonistic mechanisms (14). The model amoebae *Dictyostelium discoideum* hosts both prey and

predation-resistant bacteria, which may be obligate or transient residents (15). Green algae

microbiomes parallel those of plant roots, and include many bacteria that can colonize and benefit both

organisms (16). No studies to our knowledge have characterized bacteria associated with other groups

of rhizosphere protists, but an understanding of such communities could point to synergisms that

benefit the plant. For example, nitrogen-fixing *Sinorhizobium meliloti* is a transient resident of ciliate

cells, and association with protists increases the bacteria's distribution and nodulation capacity (17, 18).

We previously observed that inoculating plants with a defined protist consortium resulted in a

rhizosphere bacterial community composed of bacteria similar to those in the protist inoculum (19).

Protist inoculation also alleviated a negative growth effect of field soil bacteria, and this impact was

dependent on bacteria in the protist culture. These findings led us to hypothesize that the protist

microbiome could help establish a beneficial rhizosphere microbiome. In this study, we identify bacterial

taxa found in long-term culture with rhizosphere protist isolates. (Although the nature of their

interaction with protists is not known, we term these "protist-associated bacteria" for brevity). Second,

we tracked abundance of those bacterial taxa after protist inoculation onto plants. We asked, 1) Are

protist-associated bacteria enriched in the rhizosphere after protist inoculation? 2) Do the microbiome

effects of protist inoculation vary among different protists? 3) Is bacterial establishment in the

rhizosphere affected by protist inoculum diversity? Finally, 4) how do protist-associated bacteria affect

protist growth? We find that bacterial taxa associated with evolutionarily diverse protists can colonize

the rhizosphere, suggesting that protists could affect rhizosphere health through shared microbiota.

Results

Characterization of bacteria associated with rhizosphere protist isolates

Protists were previously isolated from independent maize root samples at two sites (20), and

maintained in laboratory culture for over two years without addition of live bacteria. Eight protist

isolates were selected for microbiome characterization, of which five are known to colonize the maize

rhizosphere after inoculation to roots (*Colpoda*, *Cercomonas, Ochromonas*, *Thaumatomonas*, and

Chrysophyceae sp., Table S1) (19). Three isolates were selected to represent the groups Amoebozoa and

Rhizaria (*Flamella* and *Nuclearia* sp.), and *Allapsa*, a genus with significance to plant health (5). Sequence

profiling of 16S rRNA genes in twice-washed protist cultures identified 30 to 75 bacterial genera in each

protist microbiome (Table S1). Protist culture microbiomes had lower Shannon diversity indices than

bacteria cultured from bulk and rhizosphere soils collected at the original protist isolation sites, which

we sequenced concurrently and used as bacterial inoculum for plants (Table S1).

We identified the 30 most abundant bacterial genera, belonging to 15 orders, across inoculum

samples (henceforth referred to as "dominant genera", Fig. 1). While each microbiome was distinct,

there were five core dominant genera detected in all protist microbiomes (*Pseudomonas*, *Variovorax*,

Sphingomonas, *Reyranella*, and *Bradyrhizobium*), and eight additional genera represented in six of eight

communities. However, there were no core ASVs of *Sphingomonas* or *Reyranella*, and two core ASVs of

Variovorax (Table S2), showing that the core genera include bacterial strain diversity. Members of

Burkholderiales and Hyphomicrobiales were ubiquitously abundant in the protist microbiomes; nine of

the dominant bacterial genera represented these two orders. The results indicate that while each protist

culture enriched a distinct subset of bacteria after isolation from rhizosphere soil, certain bacterial taxa

broadly persisted in and dominated microbiomes of diverse rhizosphere protists.

Establishment of protists and protist-associated bacteria in the rhizosphere

To study the fate of protist-associated bacteria in the maize rhizosphere, we inoculated maize

with the eight protist cultures singly and in mixture, and co-inoculated these with a bulk soil bacterial

community cultured from our Lockwood Farm field site (BSC1, Fig. 2). To determine the effects of the

bacterial community, additional plants were inoculated with bulk soil communities from Griswold (BSC2)

or a rhizosphere soil community from the Lockwood site (RSC, Fig. 2), and co-inoculated with the eight-

protist mixture. Rhizosphere microbial communities were profiled by amplicon sequencing of the 16S

and 18S rRNA genes after six weeks.

<u>*Colpoda* sp., but not other inoculated protists, were consistently abundant after six weeks.</u>

ASVs corresponding to four of the eight inoculated protists were detected in the rhizosphere

samples after six weeks (Table S3); all were found to establish in the maize rhizosphere three weeks after

inoculation in our previous growth chamber study: *Cercomonas* sp. (UC19), *Ochromonas* sp. (UC48),

Thaumatomonas sp. (UC70), and *Colpoda* sp. (UC22) (19). In this study, the former three protists were

detected inconsistently at six weeks, and at low read frequency (<0.01% of eukaryotic reads). In contrast,

the *Colpoda* isolate ASVs consistently dominated protist sequence in inoculated pots, comprising an

average of 20% of the eukaryotic reads (Table S3). Maize protist communities are highly temporally

dynamic (21), this result suggests that *Colpoda sp*. is uniquely persistent as a plant inoculant.

Bacterial communities were highly variable within each bacterial treatment, and bacterial

inoculation did not significantly affect rhizosphere diversity or plant biomass (Fig. S1). The lack of effect

of bacterial inoculation suggested that bacteria from non-inoculum sources, such as the open

greenhouse environment or maize seed, primarily determined the rhizosphere bacterial community

composition.

<u>The maize rhizosphere consistently recruited specific protist-associated bacteria.</u>

Thirteen bacterial ASVs were significantly enriched in the rhizosphere after inoculation with

individual protist cultures, even when the corresponding protists were not detected (Fig. 3A, Fig. S2).

Bacterial taxa enriched in the rhizosphere typically comprised greater than 0.5% of the bacterial

community in the corresponding protist cultures (Fig. 3A, Table S4), and nine of the enriched ASVs

represented dominant culture genera identified in Fig. 1. A median 8.4% of the reads in each protist

microbiome corresponded to ASVs enriched in the rhizosphere (0.8%-44%); over 17% of the microbiome

associated with *Thaumatomonas*, *Flamella,* and *Allapsa* were rhizosphere-enriched. Some ASVs that

were relatively rare in the protist microbiomes were also enriched, including *Brevundimonas* and an

unclassified Burkholderiaceae. Many highly abundant protist-associated microbiome members, such as

Variovorax (Fig. 1), were not enriched in the rhizosphere, indicating that high abundance in association

with the protist was not predictive of rhizosphere enrichment.

Eight (61%) of the enriched bacterial ASVs were enriched by only one protist treatment. In

contrast, a *Sphingomonas* ASV (*Sphingomonas*-α) was broadly enriched by four of the eight protist

cultures and comprised up to 14.3% of the rhizosphere microbiome after inoculation. *Sphingomonas*-α

was detected in all other protist microbiomes except that of *Nuclearia*. Collectively, these results

demonstrate that a substantial proportion of the rhizosphere protist microbiome can colonize the

rhizosphere, even after prolonged laboratory cultivation of protist isolates.

<u>Inoculating with multiple protists increased the abundance of protist-associated bacteria in the</u>

<u>rhizosphere.</u>

Nine of the thirteen protist-enriched bacterial ASVs were also enriched in the rhizosphere after

inoculation with an eight-protist mixture (Fig. 3A), indicating that inoculating diverse protist cultures has

an additive effect in enriching protist-associated bacteria. The abundance of eight protist-associated

bacterial taxa did not significantly differ whether single or mixed protists were inoculated (Fig. S2), even

though the mixture contained a one-eighth dilution of each culture. However, the more diverse

inoculum had a negative effect on the relative abundances of *Sphingomonas*-α, *Mucilaginibacter*,

unclassified Burkholderiaceae sp., and two uncultured Planctomycetales sp., which were significantly

lower in the mixed than single treatments (Fig. S2). Mixed protist inoculation resulted in an increased

overall abundance of protist-associated bacteria in the rhizosphere (Fig. 3B). After inoculation with

mixed protists, the thirteen protist-enriched bacterial ASVs comprised a median 9.9% of the bacterial

reads in the rhizosphere, compared with 3.5% of reads after inoculation with single protists.

<u>Treatment with some protist cultures benefited root growth when inoculated individually, but</u>

<u>not in mixture.</u>

We previously observed that plants inoculated with an antibiotic-treated protist culture had

reduced root biomass compared with plants treated with a microbiome-replete protist (19), indicating

that protist-associated bacteria were required to increase root growth. Therefore, we hypothesized that

inoculation with mixed protist cultures would have a greater biomass effect than individual cultures due

to the greater number and diversity of protist-associated bacteria enriched on the root. Among all data

combined, plants inoculated with single protists had greater root:shoot biomass ratio than uninoculated

plants, and showed a nonsignificant trend toward greater root biomass, but plants inoculated with the

protist mixture did not (Fig. 4A-B). Among single protist treatments, *Thaumatomonas* culture-treated

plants had increased root:shoot biomass ratios compared to the negative control, and *Flamella* and

Allapsa treatments had increased root biomass (Fig. 4C-D). *Flamella* and *Allapsa* had the greatest

bacterial richness among the protist cultures cultures (Table S1) and with *Thaumatomonas* had the

greatest abundance of rhizosphere-enriched bacteria (Table S4), suggesting that bacterial richness or

dominance of rhizosphere bacteria in the protist microbiome could be associated with greater biomass

impact. However, no significant biomass phenotype was observed in plants treated with the protist

mixture (Fig. 4C-D). Hence, the increased microbiome effect of the protist mixture did not translate to an

increased plant biomass effect, and instead may have precluded the growth effects associated with

Thaumatomonas, *Flamella*, and *Allapsa* treatments. Given the complex interactions among protists and

bacteria, the taxonomic composition of enriched bacteria, rather than the overall richness or proportion

of rhizosphere colonizers, may be more important for plant growth.

Protist inoculation was less impactful in the presence of a rhizosphere bacterial community

To determine whether the microbiome effect of protists was influenced by the surrounding

bacterial community, we co-inoculated protist mixtures with bacterial communities cultured from three

soil sources. Bacterial enrichment patterns following protist inoculation were similar among plants

treated with bulk soil communities or no bacterial treatments; seven ASVs were enriched in multiple

treatments (Fig. 5A). However, when protists were co-inoculated with rhizosphere bacterial community

obtained from mature maize plants (RSC), only one ASV was enriched. We hypothesized that the

rhizosphere soil community was a source of some protist-associated bacteria, and therefore they were

not enriched by the protist inoculations. By comparing RSC-inoculated plants with controls, we

confirmed that inoculation with RSC was sufficient to increase the rhizosphere abundance of four protist-

associated ASVs in the rhizosphere (Fig. S3A). RSC-treated plants also had a higher combined abundance

of protist-enriched taxa than bulk soil communities, and protist inoculation of RSC-treated plants did not

increase the total abundance of protist-associated bacteria their abundance as observed in other

treatments (Fig. S3B). This supports our previous finding that protist inoculation may substitute for

rhizosphere soil in enriching certain bacterial taxa in the environment (19).

We asked whether the rhizosphere environment was required for enrichment of protist-

associated bacteria, or if protists alone were sufficient to enrich their abundance in soil. In unplanted

pots, only *Brevundimonas* and *Pedobacter* ASVs showed increased abundance after protist inoculation,

while others were only enriched in the presence of the plant (Fig. 5B). This result indicates that the

protist-associated bacteria are specifically enriched by the rhizosphere. Unclassified Burkholderiaceae α

was not consistently detected in unplanted pots (Fig. 5B, Fig. S3C), indicating that this taxon may

obligately require either plant or protist hosts for environmental persistence.

Seventeen additional bacterial ASVs were abundant in both rhizosphere soil and protist cultures

We detected diverse rhizosphere bacterial communities on both inoculated and untreated plants

(Fig S1, Table S5), which could mask potential enrichment of some bacteria by protist inoculation. To

determine whether other bacterial taxa colonized both rhizosphere and protist microbiomes, we

identified ASVs in high rank abundance in both protist culture (top 20) and rhizosphere samples of the

plants inoculated with the corresponding single protist (top 100). Thirty ASVs met these criteria, of which

17 had not been enriched by protist inoculation (Table 1, Table S6). Shared abundant ASVs included

representatives of dominant genera that were abundant in protist cultures, including *Azospirillum*,

Pseudomonas, and *Bradyrhizobium*. Notably, a *Variovorax* sp. ASV that was highly abundant in most

protist microbiomes was also usually among the top 20 most abundant taxa on maize (Table S6).

Together, ASVs that were enriched or abundant on plants comprised a median 46.8% of the reads in the

protist culture microbiomes (Table S4, range 6.6%-67.8%). This shows that several rhizosphere protist

cultures were dominated by candidate rhizosphere-colonizing bacteria, including a wider range of taxa

than was enriched by protist inoculation.

Rhizosphere-enriched bacterial genera may promote or suppress protist growth

We lastly wondered why many rhizosphere colonizing bacteria are able to persist for long

periods in culture with diverse types of protists. We hypothesized that the bacteria are not a food

source for the protists, or alternatively, that they could provide a benefit to the protists. To test this, we

isolated four bacteria from the protist cultures having 16S rRNA gene similarity to rhizosphere-enriched

ASVs *Sphingomonas*-α (100% similarity), *Pedobacter*-β (98.4%), *Caulobacter* (98.0%), and

Mucilaginibacter (96%). These genera were all enriched in the rhizosphere after inoculation with *Allapsa*

(Fig. 3), which had a positive effect on root biomass. We measured the effects of these bacterial isolates

on antibiotic-treated cultures of *Allapsa* as well as two other isolates from phylum Cercozoa,

Cercomonas and *Thaumatomonas*, in which the four bacterial genera were detected (Fig. 1) but variably

enriched (Fig. 3). All protists grew when cultures were supplemented with *Caulobacter* or

Sphingomonas, but not when *Pedobacter, Mucilaginibacter,* or no bacteria were provided (Fig. S4); this

indicates that *Pedobacter* and *Mucilaginibacter* isolates were not a source of prey (15). To determine

whether the bacterial isolates suppress growth, we measured the effects of the bacteria on protists

supplemented with an excess of inert prey (heat-killed *E. coli*). At 3 days, *Sphingomonas* increased

growth of all three protists, increasing density of the cultures by 2.4 to 3.3-fold compared to *E. coli*

alone. *Caulobacter sp.* had a strong positive effect on growth of *Allapsa* (2.4x, p<0.001), but a smaller

effect on *Cercomonas* (1.5x, p=0.066), and no effect on *Thaumatomonas* (Fig. 6)*.* Conversely, *Pedobacter*

and *Mucilaginibacter* were strongly antagonistic to *Allapsa* growth*,* had differing positive and negative

effects on *Cercomonas*, and had no significant effect on *Thaumatomonas*. Taken together, the results

show that protist-associated rhizosphere bacteria use differing strategies to persist under predation

pressure, supporting or inhibiting protist growth in broad or isolate-specific ways.

Discussion

Green algae in soil recruit bacterial communities that have strong structural and functional

parallels to plant root communities (16), and the algal microbiome is a proposed source of plant

beneficial bacteria (22). By surveying the microbiomes of diverse protist cultures and tracking their fates

in the rhizosphere, this study demonstrates that a wide diversity of rhizosphere microeukaryotes outside

the Chloroplastida have microbiomes largely comprised of plant-colonizing bacterial taxa. Diverse

rhizosphere bacteria persisted for years in active cultures of protists from SAR, Excavata, and Amoebozoa

supergroups, and maintained the capacity to colonize roots in a complex microbial environment. The

findings support the hypothesis that, in addition to their roles imparting selective and behavioral

advantages to rhizosphere bacteria (23, 24), predatory soil protists could recruit or promote a

microbiome that contributes to rhizosphere establishment. Understanding how these bacteria interact

with protists could help identify defense and signaling mechanisms important to bacterial survival, and

may ultimately help enable the design of improved microbial consortia to benefit agriculture.

Profiling the protist microbiomes revealed broad taxonomic overlap in among protists

independently isolated from different plants and field sites, indicating the broad prevalence of

Burkholderiales and Hyphomicrobiales. They also included groups previously documented to survive

protist internalization: members of *Sphingomonas*, *Burkholderia*, *Pseudomonas*, *Pseudoxanthomonas*,

Pedobacter, *Mesorhizobium, and Acidovorax* have been identified in protist cells (Karaś et al., 2015; Lin

et al., 2023; Morón et al., 2024; Zhang et al., 2023). *Variovorax* spp. are common epibionts and

endosymbionts of protists (28, 29), and *Bosea* and *Reyranella* sp. are free-living protist mutualists that

can be isolated using a strategy of protist co-culture (30, 31). While these studies support protist

selection of these taxa, it's also important to note that protist microbiome composition is likely shaped in

part by the local soil microbiome, stochastic events during isolation, culture conditions, and bacteria-

bacteria interactions. The microbiome communities identified here likely only represent a portion of true

protist interactors in soil, and culture conditions could enhance associations that are rare or weak in

nature. More work is needed to confirm which features of protist microbiomes represent replicable

selection patterns and true symbioses. Enriched bacteria included two ASVs associated with uncultured

Planctomyces species, indicating that protist inoculation could be a means for manipulation and

rhizosphere introduction of some unculturable bacteria on the plant. However, we did not observe

rhizosphere enrichment of the foodborne human pathogens hypothesized to be vectored by amoebae

on produce (32, 33).

Protist-associated bacteria were enriched in the maize rhizosphere after six weeks, persisting

even after the corresponding protists could no longer be detected. Some enriched genera identified here

were previously correlated with protist abundance in plot studies, indicating that protist-bacterial

network connections might reflect microbiome associations that are robust in the field. For example,

Luteimonas, *Brevundimonas*, *Pseudomonas* and *Pseudoxanthomonas* were positively correlated with a

Colpoda sp. in tomato rhizosphere (34). In this study the ASV *Sphingomonas-α* was enriched to high

levels by the widest spectrum of protists; we previously identified the same ASV as the largest protist

network hub on several field grown solanaceous crops (35). *Sphingomonas* abundance was previously

linked to increased plant growth following *Cercomonas* inoculation on cucumber seedlings, and

Cercomonas triggered biofilm formation of the bacteria *in vitro* (36). Our own observations that

Sphingomonas-α is correlated with protists in the field, found in microbiomes of diverse protist cultures,

enriched in the rhizosphere after protist inoculation, and broadly promotes protist growth *in vitro*,

suggest that *Sphingomonas* could be a partner in widespread synergistic associations with protists and

plants.

The effects of protist diversity on microbial communities is variable, and can shift dynamically

under different environmental and biotic stress conditions (37, 38). We observed that inoculation with a

mixture of protist cultures enriched a greater number and abundance of rhizosphere bacteria compared

with single protist inoculation, but only a few single protists affected biomass traits. The increase in

inoculum diversity may have affected the activity and distribution of bacteria through intermicrobial

interactions, resulting in decreased abundance of potentially beneficial bacteria such as *Sphingomonas*

and *Mucilaginibacter* (Fig. S1). Even when reducing inoculum diversity, changing community composition

greatly shapes protist effects on the microbiome. We previously found that an 18-protist mixture had a

biocontrol effect and greatly enriched the dominance of the genus *Duganella* in the rhizosphere after 3

weeks (19). The eight-protist mixture tested here did not enrich *Duganella*, which we later determined is

found specifically in the microbiome of an *Ochrophyta* sp. isolate that was excluded from this study

(UC29, Patel *et al*, in preparation). As we observed in this study, inclusion of a protist isolate hosting even

a few rhizosphere-colonizing bacteria can have a significant impact on the microbiome. The observation

of parallels among plant and protist microbiomes raises many questions: What is the mechanism and

nature of protist interactions with its microbiome? Do protists provide the bacteria a fitness or other

plant colonization advantage in the rhizosphere, or induce the bacteria to benefit the plant? The

interactions identified here provide a starting point for understanding the ecology and agricultural

significance of protist holobionts on plants.

Methods

Culture preparation

We selected protist cultures from a collection isolated from maize roots in 2020 (39). Protist

cultures were maintained in PAGE's amoebal saline (i.e., PAGE's solution; 0.12 g of NaCl, 0.004 g of

$MgSO_4 \cdot H_2O$, 0.004 g of $CaCl_2$, 0.142 g of Na_2HPO_4, and 0.136 g of KH_2PO_4 in 1 liter of dH_2O) amended

with heat-killed *E. coli* as previously described (19). Protists had been maintained with monthly

passaging for two years at the time of DNA extraction and inoculation onto plants. To prepare cultures

for plant inoculation, protists were transferred to sterile PAGE's solution and fed with heat-killed *E. coli*

weekly for one month, grown just until encystment, centrifuged for 30 minutes at 2000 x g, resuspended

in 4 mL PAGE's solution, and pelleted again for 15 minutes at 2,000 x g before resuspension in 500 uL

PAGE's solution. Protist concentration was determined by counting under a microscope, then protists

were diluted in PAGE's solution to a concentration of 1000 cells/mL. Cultures were mixed in equal

proportion to create the eight-protist mixture.

To generate protist-free soil bacteria, we collected 10 cm deep soil cores from five random

locations in maize fields from Lockwood Farm (Hamden, CT) and Griswold, CT. We suspended 10 g of soil

in 100 mL of sterile phosphate-buffered saline (PBS; pH 7.4; 8 g of NaCl, 1.44 g of Na_2HPO_4, 0.2 g of KCl,

and 0.24 g of KH_2PO_4 in 1 liter of dH_2O). Rhizosphere soil was collected from 5 plants of maize (*Zea mays*

L.) inbred line B73 (stage V5) collected from the Lockwood field on the same day as the bulk soil. Roots

were shaken to remove excess soil, and 5 roots per plant were agitated in sterile PBS to collect

rhizosphere soil. Soil samples were plated in serial dilutions on both tryptic soya agar and Reasoner's 2

agar plates (0.25 g of casein digest peptone, 0.25 g of peptic digest of animal tissue, 0.5 g of $MgSO_4 \cdot 7H_2O$,

0.3 g of $C_3H_3NaO_3$, 0.5 g of casein acid hydrolysate, 0.3 g of K_2HPO_4, 0.5 g yeast extract, 0.5 g soluble

starch, 0.5 g glucose, 15 g agar in 1 L of dH_2O), which were incubated at 28°C for 72 hours. After

incubation, the bacterial colonies from each plate were scraped and collected in PBS containing 20%

glycerol and mixed by vortexing. Suspensions were stored at -80°C until use. One aliquot was used for

colony enumeration. Bacterial inoculum for plant experiments was then prepared by thawing the tubes

on ice and resuspending to 10^6 cells/mL in sterile PBS.

Plant inoculation and growth

We prepared a soil mixture containing 53% sand, 37% vermiculite, and 10% field soil from

Lockwood farm, Hamden CT, USA, which is the same field from which all but two the protists were

originally isolated. The mixture was wetted and autoclaved twice in 12 cm pots. Seeds of B73 maize were

surface sterilized in 3% NaOCL for 1 min, then washed 10 times in sterilized dH_2O. The seeds were

germinated on sterile filter paper in the dark at 28°C for 2 days. In all but 20 pots, germinated seeds

were placed into the middle of the pots at a depth of 4 cm and inoculated with 1 mL of protist inoculum

(at a concentration of 10^3 cells/mL) or sterile PAGE's solution, then with 1 mL of bacteria (10^6 cells/mL) or

sterile phosphate buffered saline (pH 7.4), depending on the experimental treatment. Germinated seeds

were inoculated by slowly pipetting the solution directly onto the emerging radicle before covering with

the soil mixture. No germinated seeds were placed in the remaining 20 pots; instead, soil was removed

to 4 cm in the center of the pot, and the soil in this location was inoculated with 1 mL of bulk soil

bacteria and 1 mL of mixed protist inoculum (10 pots), or 1 mL of bulk soil bacteria alone (10 pots),

before being covered by soil. Pots were arranged in a randomized block design and maintained in the

greenhouse for six weeks during June and July, 2021 (16h days, temperatures 25-30°C). Plants were

watered with autoclaved distilled water when pot weights decreased. At collection, roots were shaken

vigorously to remove excess soil, and then placed in plastic bags containing 35 mL PBS. Roots were

agitated for 30 seconds to remove attached soil, after which the rhizosphere suspensions were

transferred to 50 mL tubes and stored at -80°C until DNA extraction. Root and shoot biomass were

recorded after drying for one week at 50°C. For the plant-free replicates, soil was removed to a depth of

4 cm, and ~1 g of soil was removed and transferred to a 50 mL tube containing 35 mL PBS.

16S and 18S rRNA gene library preparation

DNA extractions, PCR amplifications and sequencing was conducted following previously

published protocols (19). The Illumina sequences can be found in NCBI GenBank under BioSample

PRJNA950058.

Read assembly and classification

We assembled the reads using mothur v. 1.44.0 (40). 16S contigs > 300 bp and 18S contigs < 90

bp or > 250 bp were removed, as were contigs containing ambiguous bases, or homopolymers greater

than eight bases. Chimeras were removed in mothur using VSEARCH (41). 16S amplicon sequence

variants (ASVs) were classified using the SILVA v. 132 database, while 18S ASVs were classified using the

Protist Ribosomal Reference (PR^2) v. 4.12 database (42). Both classifications used the naïve Bayesian

classifier implemented in mothur. Libraries with fewer than 2000 reads were excluded from further

analyses.

The sterilized PAGE's solution sample was analyzed as a negative control. For the 16S analyses

the control run totaled 7.9% reads of the next smallest library, while for the 18S analyses the control run

totaled 7.7% reads of the next smallest library.

Diversity and enrichment analyses

We calculated diversity metrics of inoculum samples and rhizosphere communities after

subsampling to the smallest read number among inoculum samples or rhizosphere sample datasets

using the phyloseq v. 1.38.0 package (43) implemented in R v. 4.1.2. The 16S communities were

subsampled to 2195 reads for the protist inocula, 21547 for the bacterial inocula, and 2218 for the

rhizosphere samples, while the 18S communities were subsampled to 14666 reads for the rhizosphere

samples. Heatmap generation was performed using the MicEco v. 0.9.14 package (44), after removing

ASVs with fewer than 5 reads in the dataset. Bacterial ASV enrichment between rhizosphere samples was

performed using the DESeq2 v. 1.34.0 package (45) implemented in R. Adjusted P values for DESeq2

were calculated, and false positives were estimated using the Benjamini-Hochberg method.

Bacterial isolation and identification

Protist culture solutions were diluted onto plates of TSA agar and incubated at 28° C for 3 days.

96 colonies were selected and cultured at random. Bacterial genomic DNA was extracted and amplified

using the primers 27F (5' AGAGTTTGATCCTGGCTCAG 3') and 1492R (5' GGTTACCTTGTTACGACTT 3') (46).

Partial 16S sequences were sequenced using Sanger sequencing at the Keck DNA Sequencing Core at

Yale University, New Haven, USA, and identified using Blast search in NCBI database. Strains identified as

Mucilaginibacter, *Caulobacter*, *Sphingomonas*, and *Pedobacter* were respectively isolated from cultures

of UC70, UC70, UC122, and UC225 and chosen for study. Sequence was deposited as Genbank

accessions PQ741793-PQ741796.

Protist growth assay

Cultures of *Allapsa* isolate UC225, *Cercomonas* isolate UC19, and *Thaumatomonas* isolate UC70

were treated with antibiotics to reduce endogenous bacteria as previously described (19). Protist

cultures were grown in PAGE's solution amended with heat-killed *E. coli* (O.D. 0.1) until the cells were

freshly encysted. Cysts were centrifuged twice at 4000 g and resuspended in fresh PAGE's solution to

remove excess remaining killed *E. coli* cells, then enumerated and resuspended in PAGE's solution to a

concentration of 1000 cysts/mL. Bacterial cultures were grown for 72 h in Reasoner's 2 broth media at

28°C.

Assays were conducted in 24-well cell culture plates. Protist cells were diluted to 100 cells in 1

mL of sterile PAGE's solution, while bacteria were diluted to an O.D. of 0.07. In addition, wells were

amended with heat-killed *E. coli* to an O.D. of 0.07. Three replicates each were set up of all protist-

bacteria pairwise combinations, while three replicates each were set up of protists grown without added

bacteria as controls. After the plates were set up, cells were visualized at 200 x magnification using a

Zeiss ID02 Invertoscope inverted Microscope and photographed using an Axiocam 305 color camera.

Three photos were taken of each well in a triangular pattern. Cells were photographed every 24 hours

for the following 6 days. Cells were enumerated using ImageJ, after which the number of cells in 1 mL

were calculated for each well and timepoint.

Statistical Analysis

Data were graphed using ggplot2 v. 3.4.0 (47), implemented in R version 4.3.0. Statistical

differences between treatments and controls were determined using ANOVA followed by Dunnett's

multiple comparison test using the DescTools package in R. Cell count data were analyzed in R using a

Dunnett Test to compare with the controls.

Acknowledgements

This research was funded by AFRI Foundational Program grants from the United States Department of

Agriculture National Institute of Food and Agriculture (2022-67013-37144 to L. R. Triplett, B. Steven, and

S. J. Taerum). We thank Regan Huntley and Joseph Liquori for technical assistance.

References

1. Oliverio AM, Geisen S, Delgado-Baquerizo M, Maestre FT, Turner BL, Fierer N. 2020. The global-

scale distributions of soil protists and their contributions to belowground systems. Sci Adv

6:eaax8787.

2. Gao Z, Karlsson I, Geisen S, Kowalchuk G, Jousset A. 2019. Protists: puppet masters of the

rhizosphere microbiome. Trends Plant Sci 24:165–176.

3. Martins SJ, Taerum SJ, Triplett L, Emerson JB, Zasada I, de Toledo BF, Kovac J, Martin K, Bull CT.

2022. Predators of soil bacteria in plant and human health. Phytobiomes J 6:184–200.

4. Nguyen TBA, Chen Q-L, Yan Z-Z, Li C, He J-Z, Hu H-W. 2023. Trophic interrelationships of bacteria

are important for shaping soil protist communities. Environ Microbiol Rep 15:298–307.

5. Xiong W, Song Y, Yang K, Gu Y, Wei Z, Kowalchuk GA, Xu Y, Jousset A, Shen Q, Geisen S. 2020.

Rhizosphere protists are key determinants of plant health. Microbiome 8:27.

6. Yue Y, Liu C, Xu B, Wang Y, Lv Q, Zhou Z, Li R, Kowalchuk GA, Jousset A, Shen Q, Xiong W. 2023.

Rhizosphere shapes the associations between protistan predators and bacteria within microbiomes

through the deterministic selection on bacterial communities. Environ Microbiol 25:3623–3629.

7. Röttjers L, Faust K. 2018. From hairballs to hypotheses–biological insights from microbial networks.

FEMS Microbiol Rev 42:761–780.

8. Cheng X-Y, Tian X-L, Wang Y-S, Lin R-M, Mao Z-C, Chen N, Xie B-Y. 2013. Metagenomic analysis of

the pinewood nematode microbiome reveals a symbiotic relationship critical for xenobiotics

degradation. Sci Rep 3:1869.

9. Guo H, Glaeser SP, Alabid I, Imani J, Haghighi H, Kämpfer P, Kogel K-H. 2017. The abundance of

endofungal bacterium *Rhizobium radiobacter* (syn. *Agrobacterium tumefaciens*) increases in its

fungal host *Piriformospora indica* during the tripartite sebacinalean symbiosis with higher plants.

Front Microbiol 8.

10. Partida-Martinez LP, Hertweck C. 2005. Pathogenic fungus harbours endosymbiotic bacteria for

toxin production. Nature 437:884–888.

11. Xu S, Liu Y-X, Cernava T, Wang H, Zhou Y, Xia T, Cao S, Berg G, Shen X-X, Wen Z, Li C, Qu B, Ruan H,

Chai Y, Zhou X, Ma Z, Shi Y, Yu Y, Bai Y, Chen Y. 2022. *Fusarium* fruiting body microbiome member

Pantoea agglomerans inhibits fungal pathogenesis by targeting lipid rafts. Nat Microbiol 7:831–

843.

12. Kjeldgaard B, Listian SA, Ramaswamhi V, Richter A, Kiesewalter HT, Kovács ÁT. 2019. Fungal hyphae

colonization by *Bacillus subtilis* relies on biofilm matrix components. Biofilm 1:100007.

13. Yergaliyev T, Alexander-Shani R, Dimerets, H. 2020. Bacterial community structure dynamics in

Meloidogyne incognita-infected roots and its role in worm-microbiome interactions.

https://journals.asm.org/doi/epdf/10.1128/msphere.00306-20. Retrieved 4 September 2024.

14. Amin SA, Hmelo LR, Van Tol HM, Durham BP, Carlson LT, Heal KR, Morales RL, Berthiaume CT,

Parker MS, Djunaedi B, Ingalls AE, Parsek MR, Moran MA, Armbrust EV. 2015. Interaction and

signalling between a cosmopolitan phytoplankton and associated bacteria. Nature 522:98–101.

15.   Steele MI, Peiser JM, Shreenidhi PM, Strassmann JE, Queller DC. 2023. Predation-resistant

Pseudomonas bacteria engage in symbiont-like behavior with the social amoeba *Dictyostelium*

discoideum. ISME J 17:2352–2361.

16.   Durán P, Flores-Uribe J, Wippel K, Zhang P, Guan R, Melkonian B, Melkonian M, Garrido-Oter R.

2022. Shared features and reciprocal complementation of the *Chlamydomonas* and *Arabidopsis*

microbiota. Nat Commun 13:406.

17.   Hawxhurst CJ, Micciulla JL, Bridges CM, Shor M, Gage DJ, Shor LM. 2022. Soil protists can actively

redistribute beneficial bacteria along *Medicago truncatula* roots. bioRxiv

https://doi.org/10.1101/2021.06.16.448774.

18.   Micciulla JL, Shor LM, Gage DJ. 2024. Enhanced transport of bacteria along root systems by protists

can impact plant health. Appl Environ Microbiol 90:e02011-23.

19.   Taerum SJ, Patel RR, Micciulla J, Steven B, Gage D, Triplett LR. 2024. Establishment and effect of a

protist consortium on the maize rhizosphere. Phytobiomes J PBIOMES-04-24-0035-R.

20.   Taerum SJ, Micciulla J, Corso G, Steven B, Gage DJ, Triplett LR. 2022. 18S rRNA gene amplicon

sequencing combined with culture-based surveys of maize rhizosphere protists reveal dominant,

plant-enriched and culturable community members. Environ Microbiol Rep 14:110–118.

21.   Ceja-Navarro JA, Wang Y, Ning D, Arellano A, Ramanculova L, Yuan MM, Byer A, Craven KD, Saha

MC, Brodie EL, Pett-Ridge J, Firestone MK. 2021. Protist diversity and community complexity in the

rhizosphere of switchgrass are dynamic as plants develop. Microbiome 9:96.

22. Zheng Q, Hu Y, Kosina SM, Van Goethem MW, Tringe SG, Bowen BP, Northen TR. 2023.

Conservation of beneficial microbes between the rhizosphere and the cyanosphere. New Phytol

240:1246–1258.

23. Dumack K, Feng K, Flues S, Sapp M, Schreiter S, Grosch R, Rose LE, Deng Y, Smalla K, Bonkowski M.

2022. What drives the assembly of plant-associated protist microbiomes? Investigating the effects

of crop species, soil type and bacterial microbiomes. Protist 173:125913.

24. Yue Y, Liu C, Xu B, Wang Y, Lv Q, Zhou Z, Li R, Kowalchuk GA, Jousset A, Shen Q, Xiong W. 2023.

Rhizosphere shapes the associations between protistan predators and bacteria within microbiomes

through the deterministic selection on bacterial communities. Environ Microbiol 25:3623–3629.

25. Karaś MA, Turska-Szewczuk A, Trapska D, Urbanik-Sypniewska T. 2015. Growth and survival of

Mesorhizobium loti Inside *Acanthamoeba* enhanced its ability to develop more nodules on *Lotus*

corniculatus. Microb Ecol 70:566–575.

26. Lin C, Li L-J, Ren K, Zhou S-Y-D, Isabwe A, Yang L-Y, Neilson R, Yang X-R, Cytryn E, Zhu Y-G. 2023.

Phagotrophic protists preserve antibiotic-resistant opportunistic human pathogens in the vegetable

phyllosphere. ISME Commun 3:94.

27. Morón Á, Tarhouchi AE, Belinchón I, Valenzuela JM, de Francisco P, Martín-González A, Amaro F.

2024. Protozoan predation enhances stress resistance and antibiotic tolerance in *Burkholderia*

cenocepacia by triggering the SOS response. ISME J wrae014.

28. Zhang X, Bi L, Gentekaki E, Zhao J, Shen P, Zhang Q. 2023. Culture-independent single-cell PacBio

sequencing reveals epibiotic *Variovorax* and nucleus associated *Mycoplasma* in the microbiome of

the marine benthic protist *Geleia* sp. YT (Ciliophora, Karyorelictea). Microorganisms 11.

29.   Corsaro D, Michel R, Walochnik J, Müller K-D, Greub G. 2010. *Saccamoeba lacustris*, sp. nov.

509   (Amoebozoa: Lobosea: Hartmannellidae), a new lobose amoeba, parasitized by the novel

510   chlamydia '*Candidatus Metachlamydia lacustris*' (Chlamydiae: *Parachlamydiaceae*). Eur J Protistol

511   46:86–95.

512   30.   Pagnier I, Croce O, Robert C, Raoult D, La Scola B. 2012. Genome sequence of *Reyranella*

513   *massiliensis*, a bacterium associated with amoebae. J Bacteriol 194:5698–5698.

514   31.   Thomas V, Casson N, Greub G. 2007. New *Afipia* and *Bosea* strains isolated from various water

515   sources by amoebal co-culture. Syst Appl Microbiol 30:572–579.

516   32.   Vaerewijck M, Houf K. 2015. 4 - The role of free-living protozoa in protecting foodborne pathogens,

517   p. 81–101. *In* Sofos, J (ed.), Advances in Microbial Food Safety. Woodhead Publishing, Oxford.

518   33.   Moreno-Mesonero L, Hortelano I, Moreno Y, Ferrús MA. 2020. Evidence of viable *Helicobacter*

519   *pylori* and other bacteria of public health interest associated with free-living amoebae in lettuce

520   samples by next generation sequencing and other molecular techniques. Int J Food Microbiol

521   318:108477.

522   34.   Guo S, Jiao Z, Yan Z, Yan X, Deng X, Xiong W, Tao C, Liu H, Li R, Shen Q, Kowalchuk GA, Geisen S.

523   2024. Predatory protists reduce bacteria wilt disease incidence in tomato plants. Nat Commun

524   15:829.

525   35.   Taerum SJ, Steven B, Gage D, Triplett LR. 2022. Dominance of Ciliophora and Chlorophyta among

526   ohyllosphere protists of solanaceous plants. Phytobiomes J PBIOMES-04-22-0021-FI.

36.   Guo S, Geisen S, Mo Y, Yan X, Huang R, Liu H, Gao Z, Tao C, Deng X, Xiong W, Shen Q, Kowalchuk

GA, Li R. 2024. Predatory protists impact plant performance by promoting plant growth-promoting

rhizobacterial consortia. ISME J 18:wrae180.

37.   Berlinches De Gea A, Li G, Chen JO, Wu W, Kohra A, Aslan SK, Geisen S. 2023. Increasing soil protist

diversity alters tomato plant biomass in a stress-dependent manner. Soil Biol Biochem 186:109179.

38.   Xiong W, Li R, Guo S, Karlsson I, Jiao Z, Xun W, Kowalchuk GA, Shen Q, Geisen S. 2019. Microbial

amendments alter protist communities within the soil microbiome. Soil Biol Biochem 135:379–382.

39.   Taerum SJ, Micciulla J, Corso G, Steven B, Gage DJ, Triplett LR. 2022. 18S RRNA gene amplicon

sequencing combined with culture-based surveys of maize rhizosphere protists reveal dominant,

plant-enriched and culturable community members. Environ Microbiol Rep 14:110–118.

40.   Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB,

Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ, Weber CF. 2009. Introducing

mothur: open-source, platform-independent, community-supported software for describing and

comparing microbial communities. Appl Environ Microbiol 75:7537–7541.

41.   Rognes T, Flouri T, Nichols B, Quince C, Mahé F. 2016. VSEARCH: a versatile open source tool for

metagenomics. PeerJ 4:e2584.

42.   Guillou L, Bachar D, Audic S, Bass D, Berney C, Bittner L, Boutte C, Burgaud G, de Vargas C, Decelle J,

del Campo J, Dolan JR, Dunthorn M, Edvardsen B, Holzmann M, Kooistra WHCF, Lara E, Le Bescot N,

Logares R, Mahé F, Massana R, Montresor M, Morard R, Not F, Pawlowski J, Probert I, Sauvadet A-L,

Siano R, Stoeck T, Vaulot D, Zimmermann P, Christen R. 2013. The Protist Ribosomal Reference

database (PR2): a catalog of unicellular eukaryote Small Sub-Unit rRNA sequences with curated

taxonomy. Nucleic Acids Res 41:D597–D604.

43.   McMurdie PJ, Holmes S. 2013. phyloseq: An R package for reproducible interactive analysis and

graphics of microbiome census data. PLoS ONE 8:e61217.

44.   Russel J. 2021. Russel88/MicEco: v0.9.15. Zenodo.

45.   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-

seq data with DESeq2. Genome Biol 15:550.

46.   Lane DJ. 1991. 16S/23S sequencing, p. 115–175. *In* Nucleic Acid Technologies in Bacterial

Systematics. Wiley, New York, NY, U.S.A.

47.   Wilkinson L. 2011. ggplot2: elegant graphics for data analysis by WICKHAM, H. Biometrics 67:678–

679.

Tables

Table 1. Protist culture microbiome bacterial ASVs that were enriched or abundant in the rhizosphere

after protist inoculation.

ASV[a]	Genus	Protist culture ID	Enriched in rhizosphere[b]	Abundant in rhizosphere[c]
1	*Azospirillum* sp.	UC19, UC48, UC74, UC122		X
2	*Bosea* sp.	UC74		X
3	*Bosea* sp.	UC19, UC22, UC74	X	X
4	*Bradyrhizobium* sp.	UC19, UC22, UC48, UC70, UC74, UC122		X
5	*Brevundimonas* sp.	none	X	
6	*Brevundimonas sp.*	UC242		X
7	Burholderiaceae uncl.	UC22		X
8	Burkholderia uncl.	UC74		X
9	Burkholderiaceae uncl.	UC122	X	X
10	*Caulobacter* sp.	UC225, UC242	X	X
11	*Caulobacter* sp.	UC19, UC225		X
12	*Cupriavidus* sp.	UC225		X
13	Enterobacteriaceae uncl.	UC48		X
14	*Luteimonas* sp.	UC70, UC122	X	X
15	*Mucilaginibacter* sp.	UC19, UC22		X
16	*Mucilaginibacter* sp.	none	X	
17	*Paenarthrobacter sp.*	UC242		X
18	*Pedobacter* sp.-α	UC22, UC70	X	X
19	*Pedobacter* sp.-β	UC225	X	X
20	*Pseudomonas* sp.	UC19, UC48, UC74		X
21	*Pseudoxanthomonas* sp.	UC122, UC19		X
22	*Reyranella* sp.	UC225, UC242	X	X
23	*Reyranella* sp.	none		X

24	*Rhodococcus* sp.	UC19, UC22, UC48, UC74, UC122, UC242		X
25	*Sphingomonas* sp.-α	UC22, UC48, UC70, UC122	X	X
26	*Sphingomonas* sp.-β	none	X	
27	uncl. Planctomycetales	UC122	X	X
28	uncl. Xanthobacteraceae	UC48, UC70, UC122, UC225		X
29	uncultured gen. SH-PL14	none	X	
30	*Variovorax* sp.	UC19, UC22, UC48, UC70, UC74, UC122, UC242		X

a.   ASV identifiers are found in Table S5.

b.   Significantly enriched in maize rhizosphere 42 days after single protist inoculation in greenhouse, relative

to uninoculated plants

c.   Among the 100 most abundant bacterial ASVs in the rhizosphere 42 days after protist inoculation,

regardless of enrichment.

d.   X indicates that a matching ASV was enriched in maize rhizosphere samples 21 days after inoculation with

18-member protist consortium in the growth chamber. (g) indicates that other ASVs from the same genus

were enriched in the growth chamber.

Figure captions:

Figure 1: Bacterial microbiome composition of the protist-free bacterial inocula and the eight protist

isolates from the maize rhizosphere. Genera having the highest mean abundance across samples are

shown. Bacterial order is indicated after the colon. Shade corresponds to proportional abundance in the

culture sample.

Figure 2: Experimental design showing eight combinations of protist and bacterial treatments. Numbers

of planted replicates are listed under each circled number. Six plants failed to grow and eleven 16S

rhizosphere libraries had fewer than 2000 reads, so these samples were removed from the study.

Figure 3: Enrichment of bacterial ASVs in the maize rhizosphere six weeks after protist inoculation onto

germinated seeds. **A.** Enrichment by single protists. Diameter of diamonds and dots indicates the mean

relative abundance of ASVs in the protist culture microbiome (n = 1) and the rhizosphere samples (n = 9

or 10), respectively. Blue color indicates statistical enrichment in the protist treatment relative to protist-

uninoculated controls, as determined by DeSeq2. Red stars indicate that the bacteria were enriched

when the eight protists were inoculated as a consortium. **B.** The proportion of rhizosphere reads

representing protist-enriched ASVs after plants were inoculated with single protist cultures or mixed

protists, or uninoculated. Asterisks indicate significant differences from the mixed protist treatment

using Dunnett's Multiple Comparison test (*** = p<0.0005).

Figure 4: Biomass traits of maize plants inoculated with single protists and an eight-member consortium

in a greenhouse study. **A-B.** Root biomass (**A**) and root-shoot mass ratio (**B**) six weeks after inoculation

with zero (n = 29), one (n = 78), or a mixture of eight protists (n = 29) onto germinated maize seeds.

Filled asterisks indicate significant differences from the single protist treatment at p < 0.05. **C-D.** Root

biomass (**C**) and root-shoot mass ratio (**D**) after individual protist inoculation in the presence of protist-

free soil bacteria (n = 10 for all treatments except UC22 and UC74, which had one plant fail to grow).

Asterisks indicate significant differences from the no-protist treatment at p < 0.05.

Figure 5: Effect of bacterial source (**A**) and rhizosphere environment (**B**) on protist enrichment of soil

bacteria. **A.** The eight-protist consortium was co-inoculated with bulk soil bacteria from Lockwood Farm

(BSC1), Griswold Farm (BSC2), maize rhizosphere soil bacteria from Lockwood Farm (RSC), or no bacteria.

B. Enrichment after the eight-protist consortium was inoculated into the center of unplanted pots. Dot

diameter indicates the mean relative abundance of the ASV (n = 9 or 10). Blue color indicates statistical

enrichment in the protist mixture treatment relative to protist-uninoculated controls within each

column.

Figure 6: Cell counts of three protist taxa (*Allapsa*, *Cercomonas*, and *Thaumatomonas*) after being fed

with heat-killed *E. coli* alone, or with heat-killed *E. coli* in addition to one of four bacterial cultures

isolated from protist cultures (n = 3 for each treatment). Cells were counted at 0, 3 and 6 days after the

cells were initially fed. Black asterisks indicate significantly more protists than those fed with heat-killed

E. coli alone, while red asterisks indicate significantly fewer (p < 0.05).

Supplementary figure captions:

Figure S1: Effect of bacterial treatment on (**A**) the β-diversity of rhizosphere bacterial communities and

(**B**) the biomass of maize plants. **A.** NMDS plot showing the clustering of 16S communities of maize

rhizospheres after inoculation with bulk soil bacteria from Lockwood Farm (BSC1), Griswold Farm (BSC2),

maize rhizosphere soil bacteria from Lockwood Farm (RSC), or no bacteria, and either no protists or the

eight-protist consortium. Dashed lines represent 95% confidence intervals. **B.** Root biomass of maize

plants six weeks after inoculation with the different bacterial treatments, with or without the eight-

protist mixture.

Figure S2: Relative abundances of the 13 enriched bacteria in maize rhizospheres six weeks after

inoculation with zero, one, or a mixture of eight protists onto germinated maize seeds. Asterisks indicate

that relative abundances were significantly greater (p < 0.05) than in the mixture-inoculated plants

based on a Dunnett's Multiple Comparison Test.

Figure S3: Effect of rhizosphere on bacterial enrichment and abundance. **A.** Enrichment of five bacterial

ASVs in maize rhizospheres inoculated with bulk soil bacteria from Lockwood Farm (BSC1) alone or maize

rhizosphere soil bacteria from Lockwood Farm (RSC) alone compared with uninoculated controls. Circle

diameters indicate mean relative abundances of bacteria. **B.** Proportion of reads of the 13 enriched

bacterial ASVs in the plants inoculated with no protists, but different bacterial inocula. Asterisks indicate

that the read proportions are significantly greater when plants were inoculated with protists compared

with the plants that were not inoculated with protists at p < 0.005 (**) and p < 0.001 (***). **C.** Relative

abundance of the bacterial ASV Burkholderiaceae unclassified β in soil inoculated with the protist

mixture or no protist, with or without the plant.

Figure S4: Cell counts of three protist taxa (*Allapsa*, *Cercomonas*, and *Thaumatomonas*) after being fed

with no bacteria, or one of four bacterial isolates from protist cultures (n = 3 for each treatment). Cells

were counted at 0, 3 and 6 days after the cells were initially fed. Black asterisks indicate significantly

more protists than those fed with no bacteria (p < 0.05) based on a Dunnett's Multiple Comparison Test.