Isometric handgrip contraction increases tibialis anterior intrinsic motoneuron excitability in a dose-dependent manner

Lucas Ugliara[1], Lucas B. R. Orssatto[2], Amilton Vieria[1] and Gabriel S. Trajano[3].

[1]Faculty of Physical Education, University of Brasilia (UnB), Brasilia, Distrito Federal, Brazil.

[2]Centre for Sensorimotor Performance, School of Human Movement and Nutrition Sciences, The University of Queensland, Brisbane, Queensland, Australia.

[3]Faculty of Health, School of Exercise and Nutrition Sciences, Queensland University of Technology (QUT), Brisbane, Queensland, Australia.

Abstract

Persistent inward currents (PICs) contribution to motoneuron firing in the lower limb typically increase after a remote handgrip contraction, believed to result from diffuse increases of serotonergic input on the spinal cord. We investigated whether handgrip contraction intensity, duration, and/or impulse would affect tibialis anterior estimates of PICs. Multi-channel electromyograms were recorded from the tibialis anterior of 21 participants (18-40 years), during dorsiflexions at 20% of individual's maximal torque, before and after four handgrip conditions: i) *80%15s*, 80% of their maximal handgrip strength sustained for 15s; ii) *40%15s*, 40% sustained for 15s; iii) *40%30s*, 40% sustained for 30s; and iv) *Control* (no handgrip). PICs contribution to self-sustained motoneuron firing was estimated with the delta frequency (ΔF) using the paired motor unit analysis. The 'brace height', normalised as a percentage of a right triangle (%rTri), was used to quantify the effects of PICs on the non-linearity of firing patterns, representing the neuromodulatory drive (metabotropic regulation of motoneuron excitability) onto the motoneurons. ΔF increased by 0.33 pulses per second (pps; 95%CI 0.16–0.49, *d*=0.47) after *40%30s* and by 0.24 pps (0.09–0.38, *d*=0.34) after *80%15s* but remained unchanged after *40%15s* and *Control*. Similarly, brace height increased by 2.24 %rTri (0.18–4.30, *d*=0.20) after *40%30s* and by 2.45 %rTri (0.64–4.25, *d*=0.22) after *80%15s;* remaining unchanged after *40%15s* and

31 *Control*. The increase in PICs contribution to motoneuron firing induced by a remote

32 handgrip contraction is impulse-dependent rather than intensity or duration. The

33 parallel increases in ΔF and brace height suggest augmented neuromodulatory input

34 onto the spinal cord.

Introduction

Motoneurons rely on a complex control system for excitability, with

neuromodulation as a key component. Unlike ionotropic systems, which directly

convert synaptic inputs into action potentials via ion channel opening,

neuromodulation operates through neurotransmitters that bind to specific receptors,

initiating intracellular signalling pathways that regulate motoneuron responsiveness

(Heckman et al., 2009). This neuromodulatory influence facilitates the generation of

strong persistent inward currents (PICs) in spinal motoneurons, which increase cell

excitability, accelerating, amplifying, and prolonging their discharge output activity for

a given excitatory input (Heckman, Johnson, et al., 2008). This input–output gain

mechanism can be adjusted based on the level of serotonin (5HT) and noradrenaline

(NE) released from the raphe nuclei and locus coeruleus, respectively, which travel

through the monoaminergic projections in the spinal cord (Heckman, Hyngstrom, et

al., 2008), and activate their respective receptors located largely on the

motoneurons' dendrites. This activation promote an intracellular response via second

messengers, leading to the opening of L-type Ca^{2+} and Na^+ channels and facilitating

PICs in a dose-dependent manner (Johnson & Heckman, 2014). It has been

suggested that the level of neuromodulatory input onto the motoneurons could be

adjusted according to the physical tasks demand (Heckman et al., 2009). Lower

levels of monoaminergic drive would reduce PICs magnitude, allowing the motor

system to perform delicate muscular actions with accuracy (e.g., threading a needle).

Alternatively, moderate-to-maximal motor tasks (e.g., lifting heavy weights) would

demand higher monoaminergic drive and an increased state of PIC facilitation

(Johnson & Heckman, 2014). This theory is supported by the observation that the

firing rate of neurons in the caudal raphe nuclei is influenced by the intensity of motor

outflow, such that 5HT release is proportional to locomotion demands

(Fenstermacher et al., 2024; Veasey et al., 1995). Importantly, much of current

knowledge about the effects of neuromodulation on PICs and the role of 5HT in
motoneuron gain control comes from invasive animal experiments and computational
simulations (Heckman et al., 2009; Hounsgaard et al., 1988; Lee & Heckman, 2000).
However, translating these findings to humans can be challenging, requiring non-
invasive techniques.

In humans, researchers have used different strategies to investigate the
contribution of 5HT and NE on PICs neuromodulation assessed with indirect
measures. Pharmacological trials involving drugs that can change the concentration
of 5HT and NE at the synaptic cleft have demonstrated their influence on estimates
of PICs and different variables linked to motoneuron excitability (e.g., discharge rate,
recruitment threshold) (D'Amico et al., 2013; Goodlich et al., 2023, 2024; Udina et
al., 2010). Additionally, some studies have used muscle contraction of different
intensities to theoretically increase monoaminergic input onto the spinal cord to
understand its effects on PICs neuromodulation. These studies show that PICs
contribution to motoneuron firing increase with contraction intensity levels of the
same muscle group (Mackay et al., 2023; Orssatto et al., 2021). Other studies used
remote handgrip contractions to, theoretically, increase 5HT availability via diffused
serotonergic projections on the spinal cord and; thus, increase estimates of PICs on
lower limb muscles (Mackay Phillips et al., 2023; Orssatto et al., 2022). This
paradigm was supported by Wei et al. (2014), who showed that remote leg
contractions increased force variance during a precision task with the palm or index
finger. This effect was interpreted as 5HT-mediated, as force variation decreased in
the presence of selective 5HT receptor blockers and reuptake inhibitors. These
findings aligns with evidence that the firing rate of neurons in the caudal raphe nuclei
and locus coeruleus is influenced, respectively, by the intensity of motor outflow and
arousal state, such that 5HT and NE release is proportional to motor task or
locomotion demands (Fenstermacher et al., 2024; Heckman, Johnson, et al., 2008;
Veasey et al., 1995). Collectively, these findings align with evidence from animal
experiments and support the assumption that it is feasible to investigate indirectly the
contribution of serotonin, using contraction of remote muscle groups, to promote
changes in motoneuron firing pattern in humans.

After establishing the efficacy of using remote handgrip tasks to increase 5HT
input onto the spinal cord and facilitate tibialis anterior and soleus PICs (Mackay

Phillips et al., 2023; Orssatto et al., 2022), the logical next step is to understand whether the PICs response could be differently influenced by mechanical aspects of the remote contraction (i.e., force intensity, duration, and impulse). This rationale is supported by evidence that higher contraction intensities increase estimates of PIC contribution to motoneuron self-sustained firing (e.g., ΔF) in muscles like the soleus, gastrocnemius (Orssatto et al., 2021) and tibialis anterior (Mackay et al., 2023), suggesting that intensity modulates PIC activation. Additionally, serotonergic neuron activity in the raphe nuclei of cats rises and maintains during prolonged locomotion until fatigue (*i.e,* the inability of the cat to maintain pace) (Jacobs et al., 2002) implying that sustained motor activity may also enhance PIC levels. To investigate this phenomenon, the present study used multi-channel electromyography to assess MU discharge rates, and the paired MU technique to estimate PICs contribution to motoneuron firing (Mesquita et al., 2024) following handgrip contractions at different intensity levels and contraction durations. In addition, we estimated the non-linearity caused by the effect of monoaminergic drive onto motoneuron discharge patterns using a quasi-geometric approach to calculate the 'brace height' of the ascending phase of the MU firing rates (Beauchamp et al., 2023). Given the evidence of the importance of the intensity of both muscle contraction for the estimates of PICs of motoneurons (Mackay et al., 2023; Orssatto et al., 2021) and greater levels of motor output inducing greater concentration of 5HT in the spinal cord (Jacobs et al., 2002), potentially leading to increasing motoneuron excitability, we hypothesize that the combination of both greater intensity and duration (i.e., impulse) of remote contraction will induce higher estimates of PIC activity.

Methods

Participants and ethical procedures

Twenty-three young adults participated in this study (Table 1). Inclusion criteria required volunteers to be within the age range of 18–40 years, have no history of musculoskeletal injuries in the tested limbs, and not be using medications that could affect the monoaminergic system (e.g., beta-blockers and selective serotonin reuptake inhibitors) (Thorstensen et al., 2024). Furthermore, participants were instructed to abstain from strenuous physical activities and alcohol

128 consumption for 48 hours before the testing session. The Queensland University of
129 Technology Human Research Ethics Committee approved the study (Reference
130 number: 6770), and all participants signed a written informed consent prior to their
131 participation.

Study design and testing procedures

The dorsiflexion and handgrip contractions were performed using the
"preferred leg for kicking a ball" and the corresponding ipsilateral hand, respectively.
For the dorsiflexion tasks, participants were positioned upright in an isokinetic
dynamometer (Biodex System 4, Biodex Medical system, Shirley, NY) with the knee
fully extended (0˚), seat reclined at 70˚ of hip flexion and ankle in anatomical
position. For the handgrip tasks, a handgrip dynamometer (model MLT004/ST,
ADinstruments, Australia) was held with shoulder in the anatomic position at 0°
flexion/extension, 0° abduction/adduction, and 10-15° external rotation, with the
elbow flexed at 90˚, while seated on the isokinetic dynamometer. A warm-up was
performed alternating between handgrip and ankle dorsiflexion tasks, and consisted
of progressive contraction levels (20%, 40%, 60%, 80% of perceived maximal effort)
for ~5 s each. Two minutes after, they performed two 4-s handgrip maximal voluntary
contractions, with 60 s of interval between attempts. They were given two minutes of
resting and then also performed two 4-s dorsiflexor maximal voluntary contractions,
with 60 s of interval between attempts. Peak handgrip force and dorsiflexion torque
were considered as the maximum value achieved during the maximal voluntary
contractions. Thereafter, participants were familiarized with triangular-shaped
ramped contractions to 20% of their dorsiflexion peak torque at 2%/s rate of torque
increase and decrease (10-s up and 10-s down). Visual feedback was provided
through a 23" computer monitor, guiding the participants to closely follow the real-
time torque trajectory. This relative torque level was chosen for based on previous
studies investigating the effects of a remote handgrip contraction on lower limbs ΔF
(Mackay Phillips et al., 2023; Orssatto et al., 2022).

Five minutes after familiarization, four clusters of two sets of triangular-shaped
contractions, interspersed with either a resting control condition or three distinct
handgrip contractions were performed in a randomized order. During the control
condition, the participants were asked to remain quiet and relaxed, avoiding to

move/contract any muscle for 60 s in between the two triangular-shaped

contractions. For the three conditions involving a remote handgrip contraction,

participants were requested to quickly reach a relative grip force level and sustain it

for a given time; the three conditions were i) handgrip contraction at 40% of their

peak force sustained for 15 s (i.e., 40%15s); ii) handgrip contraction at 40% of their

peak force sustained for 30 s (i.e., 40%30s), and iii) handgrip contraction at 80% of

their peak force sustained for 15 s (i.e., 80%15s). Conditions 80%15s and 40%30s

were impulse matched, 80%15s and 40%15s were time-matched, and 40%30s and

40%15s were intensity matched. This approach was adopted to allow us to

investigate whether the increases in ΔF are affected by remote contraction time

(40% sustained for 15 s vs 30 s), intensity (40% vs 80% sustained for 15 s), or

impulse (40% sustained for 30 s and 80% sustained for 15 s vs 40% sustained for 15

s). The time between the two triangular-shaped contractions, performed before and

after the control or handgrip conditions, was standardized at 60 s. Thus, a 30-s

waiting period preceded the handgrip contraction lasting 30 s, while a 45-s waiting

period preceded the handgrip contraction lasting 15 s. The second triangular-shaped

contractions were performed immediately after the handgrip conditions. A 5-min rest

interval was adopted between conditions. Figure 1 illustrates the study protocol

design. In case abrupt/steep increases or decreases (>5% of peak torque) of torque

were observed during the ascending or descending phase of the triangular-shaped

contractions, the whole trial for the specific condition was excluded and repeated

after 5 minutes. The participants were given only one extra attempt for each

condition, when necessary. No further attempts were given to avoid the presence of

fatigue, which can affect the outcomes obtained from the triangular contractions

(Mackay et al., 2023). If the participant was not able to maintain consistency in the

before and after triangular ramp-shaped contractions (i.e., avoid abrupt/steep

increases or decreases in torque) for both attempts in a given condition, they were

excluded from the analysis for that condition. Finally, the impulse (area under the

force curve) performed at the handgrip conditions was calculated using the

trapezoidal rule in LabChart software (version 7.3, ADinstruments, Australia).

Figure 1. Study testing protocol.

Blue triangles represent the triangular ramp-shaped dorsiflexion contraction with the time between these two tasks set at 60 s for all conditions; the dashed orange rectangle represents half handgrip impulse as the other two solid-lined orange rectangles; conditions were performed randomly with 5 min rest interval.

Multi-channel Electromyography Recordings and Analyses

The tibialis anterior skin area was prepared by shaving and cleansing with 70% isopropyl alcohol. A semi-disposable 64-channel electrode grid with an interelectrode distance of 8 mm (GR08MM1305, OT Bioelettronica, Turin, Italy), attached to a bi-adhesive foam layer and coated with a conductive paste (Ten20, Weaver and Company, Aurora, CO, USA) was positioned over the most prominent part of the tibialis anterior. The grounding electrode comprised dampened strap (WS2, OTBioelettronica, Torino, Italy) placed around the ankle joint.

Multi-channel electromyograms were recorded during the triangular ramp-shaped

contractions using monopolar mode, amplified (256×), subjected to band-pass

filtering (10–500 Hz), and digitally sampled at 2048 Hz, with a 16-bit wireless

amplifier (Sessantaquattro, OT Bioelettronica, Turin, Italy), interfacing with OTBioLab

+ software (version 1.3.0., OT Bioelettronica, Turin, Italy) and were stored for

subsequent offline analysis. The recorded data was processed offline using the

DEMUSE software (Holobar & Zazula, 2007). The signals were band-pass filtered

(20–500 Hz) through a second-order, zero-lag Butterworth filter. Subsequently, MU

decomposition was performed using the convolutive kernel compensation (CKC)

method (Holobar et al., 2014; Holobar & Zazula, 2007). Following decomposition, the

same MUs were tracked across the before and after handgrip or control conditions,

but not across conditions. The files for each contraction were concatenated and

separation vectors (i.e., MU filters) were used to identify and generate the MU spike

trains across the two contractions (Del Vecchio et al., 2019; Holobar et al., 2014;

Holobar & Zazula, 2007). Then, a trained investigator (LU) examined the MU spike

trains, and, when necessary, edited the discharge patterns of the MUs (Del Vecchio

et al., 2020). Only MUs with a pulse-to-noise ratio ≥ 30 dB were retained for

presenting higher reliability (sensitivity > 90% and false alarm rates < 2%) (Holobar

et al., 2014). The edited MUs were quality-checked by a researcher, blinded for each

condition, (LBRO) with extensive experience in decomposing, tracking and editing of

MUs. When no MU was identified in both contractions for a given condition, the

participant was not included in the respective condition analysis.

Estimation of PIC contribution to motoneuron self-sustained firing (ΔF) and peak

discharge rate

The MU discharge events were converted into instantaneous discharge rates

and smoothed with support vector regression machine learning fit (Beauchamp et al.,

2022). PIC contribution to motoneuron firing was estimated using the paired MU

analysis (Afsharipour et al., 2020). MUs with lower recruitment thresholds (control

units) were paired with units of higher recruitment threshold (test units). ΔF was

calculated as the change in discharge rates of the control MU from the onset of

recruitment to the point of de-recruitment of the test unit (i.e., discharging hysteresis)

(Afsharipour et al., 2020). MUs were paired when the following criteria was obtained

between control and test units i) a rate-to-rate correlation threshold of r ≥ 0.7, ii) test

– control unit recruitment time >1 s, iii) the control unit's discharge rate at test unit

recruitment minus the control unit's peak discharge rate was > 0.5 pps, iv) control

unit derecruitment time > test unit derecruitment time (Afsharipour et al., 2020;

Hassan et al., 2020; Vandenberk & Kalmar, 2014). ΔFs calculated for each individual

test unit were averaged across control units to yield a singular ΔF value for each

corresponding test MU. In the case that no MU pair was identified in both

contractions for a given condition, the participant was not included in the analysis for

that respective condition(s). The highest value derived from the support vector

regression fit curve was determined as peak discharge rate. The relative torque (%)

produced during the recruitment time of each MU was identified as MU recruitment

threshold.

Quantification of neuromodulatory and inhibitory drive in MUs firing changes

We used the quasi-geometric approach to quantify the neuromodulatory and

inhibitory drive onto the MUs, as proposed by Beauchamp et al. (2023). Firstly, we

generated a straight line (hypothenuse) from the discharge rate at MU recruitment to

the peak discharge rate of the support vector regression (SVR) smoothed curve.

Then, we calculated the 'brace height', which represents the maximum deviation of

the smoothed discharge rates trace from linearity (hypothenuse). The maximal

orthogonal distance between the hypothenuse and the smoothed MU discharge

trace was considered the brace height. Thereafter, brace height was normalized as a

percentage of the maximal orthogonal distance between the straight line and a right

triangle in which sides originate from the hypotenuse. We also calculated the

'attenuation slope', which is associated with the inhibitory input effect on MU

discharge (Beauchamp et al., 2023). Attenuation slope was calculated from the

brace height insertion on the smoothed discharge rates to the peak discharge rates

of the ascending phase.

Figure 2. A, delta frequency (ΔF) calculation from a single participant for triangular-shaped contraction to 20% of maximal their maximal peak torque.

Red was used to represent firing of the test unit (top), paired with other three control units (green, blue and purple). Solid black lines, SVR smoothed curve of the MUs; dashed vertical black lines, moment of recruitment and decruitment of the test unit; dashed horizontal grey lines, firing of control units at the moment of recruitment and derecruitment of the test unit, used to calculate ΔF.

 B, brace height and attenuation slope calculation for the same MUs in A.

Solid coloured lines, SVR smoothed curve of the MUs; dashed coloured lines, brace height; solid black line, orthogonal distance originating from the hypotenuse (blue dot) to the right triangle (black dot) formed by the moment of recruitment to derecruitment of each MU (dashed black lines); dashed grey lines, attenuation slope.

Data and statistical analyses

Linear-mixed effects models were investigated utilizing the *robustlmm* package (Kuznetsova et al., 2017). We estimated marginal mean differences between handgrip conditions in ΔF, peak discharge rate, brace height and attenuation slope along with 90% and 95% confidence intervals (CI), using the emmeans package (Lenth et al., 2023). Each MU was treated as repeated measure and nested for each participant, including a random intercept for each participant to consider for the correlation between repeated observations on each individual (i.e., 1| participant/MU ID). For ΔF, four distinct models were fitted: 1) time and condition were included as fixed effects and a random intercept and slope (ΔF) for each participant; 2) similar to model 1, but included peak discharge rate as covariate; 3) similar to model 1, but included recruitment threshold as covariate; 4) similar to model 1 but included both peak discharge and recruitment threshold as covariates. All the four models resulted in similar outcomes; thus we presented the findings derived from Model 4, which exhibited superior fit (as indicated by the lowest Akaike's information criteria and Bayesian information criteria). We performed an additional analysis by adding sex as fixed effect to ΔF to investigate a possible effect of sex on results, but no time by condition by sex interaction effect [β = 0.32 (-0.29, 0.92) pps, SE = 0.32; t = 0.31] and main effect of sex [β = -0.80 (-1.76, 0.16) pps, SE = 0.49; t = -1.64] were observed, so sex was removed from the model. For all other variables except by recruitment thresholds (i.e., peak discharge rate, brace height and attenuation slope), a model with time and condition as fixed effects and recruitment thresholds as covariate was employed. For recruitment thresholds, the model included time and condition as fixed effects. In addition, the mean impulse for each condition were compared using the *lmerTest* package for linear mixed effects models analysis. Pairwise *post hoc* analysis was used for significant effects observed. Cohen's d effect sizes were computed for the differences between after and before handgrip conditions, leveraging the population standard deviation (σ) estimated from corresponding robust linear mixed-effects models as the denominator (i.e, d = mean difference / σ). For interpretation, we considered Cohen's d <0.20 trivial; 0.20–0.49 small; 0.50–0.79 moderate; ≥0.80 large. A significance threshold of 5% (α level) was adhered to for all tests. All the analyses were conducted in a free

software environment (RStudio, version 2024.12.1). The complete dataset and corresponding R script are accessible at https://github.com/lugliara/PICs.

Results

Participant characteristics and MU identification

One female participant was excluded from all the analyses due to inconsistencies in performing the ramp-shaped triangle contractions in all the conditions and another female participant was excluded because no MUs were identified at any condition. The final analysis included data from 21 participants. Among these 21 participants, all of them were included in the control and 80% of peak force for 15 s conditions (80%15s). For the 40% of peak force for 15 s (40%15s), data from five participants were excluded from the analyses because they were unable to consistently follow the force trace path during the triangular-shaped contractions without steep increases and decreases of force production (N = 3), or MUs did not meet essential prerequisites to form pairs (N = 2). For the 40% of peak force for 30 s (40%30s), data from three participants were excluded from the analyses because they were unable to follow the feedback path (i.e., detected abrupt/steep increases or decreases >5% of peak torque) during the triangular-shaped contractions (N = 2) or no MUs were detected (N = 1).

341, 249, 245 and 316 MUs which were matched before and after the control (mean (95% CI) = 16.2 (13.9, 18.6), N = 21), 40%15s (15.6 (13.3, 17.8), N = 16), 40%30s (13.6 (10.8, 16.4), N = 18) and 80%15s (15.0 (12.2, 17.9), N = 21) conditions, respectively. This resulted in 199 (9.5 (7.8, 11.2), N = 21), 146 (9.1 (7.2, 11.1), N = 16), 137 (7.6 (5.7, 9.6), N = 18) and 191 (9.1 (6.8, 11.4), N = 21) test units, respectively.

Handgrip contraction impulse

The impulse generated during the handgrip contractions during 40%30s (estimated marginal mean = 5639 (95%CI: 5054, 6224) N·s) and 80%15s (5667 (95%CI: 5082, 6252) N·s) were similar (p = 0.868), but superior than 40%15s (2933 (95%CI: 2348, 3519) N·s, p < 0.001).

Table 1. Participant characteristics

Participant characteristics	
Age (years)	31 (29, 33)
Sex (n)	
Male	15
Female	6
Body mass (kg)	79 (73, 86)
Height (cm)	177(173, 181)
BMI (kg/m^2)	25 (24, 27)
Handgrip peak force (N)	437 (386, 488)
Dorsiflexion peak torque (N·m)	54 (49, 60)

Participant characteristics data are presented as means with 95% confidence interval.

MU outcomes

ΔF

A time by condition interaction effect was observed β = -0.40 (-0.62, -0.18)
pps, SE = 0.11; t = -3.60]. ΔF increased from before to after the intervention on
40%30s [0.33 (0.16, 0.49) pps, *d* = 0.47 (0.23, 0.72)] and 80%15s [0.24 (0.09, 0.38)
pps, *d* = 0.34 (0.14, 0.55)], but remained unchanged on 40%15s [0.10 (−0.06, 0.26)
pps, *d* = 0.15 (-0.09, 0.38)] and control [-0.07 (-0.21, 0.06) pps, *d* = -0.11 (-0.31,
0.09)] conditions (Figure 3).

Brace height

A time by condition interaction effect was observed β = -2.56 (-5.06, -0.06) %
rTri, SE = 1.27; t = -2.01]. Brace height increased from before to after the
intervention on 40%30s [2.24 (0.18, 4.30) % rTri, *d* = 0.20 (0.02, 0.39)] and 80%15s
[2.45 (0.64, 4.25) % rTri, *d* = 0.22 (0.06, 0.39)], but remained unchanged on 40%15s
[1.86 (-0.15, 3.87) % rTri, *d* = 0.17 (-0.01, 0.35)] and control [-0.11 (-1.84, 1.61) %
rTri, *d* = -0.01 (-0.17, 0.15)] conditions.

Attenuation slope

A time by condition interaction effect was observed β = 0.06 (0.02, 0.10)

pps/%MVT, SE = 0.02; t = 2.73]. Attenuation slope decreased from before to after the

intervention on 40%30s [-0.05 (-0.08, -0.01) pps/%MVT, *d* = -0.27 (-0.46, -0.08)]and

80%15s [-0.03 (-0.06, -0.00) pps/%MVT, *d* = -0.17 (-0.34, -0.01)], but remained

unchanged on 40%15s [-0.01 (-0.04, 0.02) pps/%MVT, *d* = -0.07 (-0.26, 0.11)] and

control [0.01 (-0.01, 0.04) pps/%MVT, *d* = 0.07 (-0.08, 0.23)].

Peak discharge rates

A time by condition interaction was observed β = -0.50 (-0.62, -0.37) pps, SE

= 0.06; t = -7.64]. Peak discharge rates decreased from before to after the

intervention only on control [-0.29 (−0.38, -0.21) pps, *d* = -0.52 (-0.67, -0.36)] and

increased on 80%15s [0.20 (0.11, 0.29) pps, *d* = 0.35 (0.19, 0.51)] condition, but

remained unchanged on 40%15s [0.06 (-0.04, 0.16) pps, *d* = 0.10 (-0.08, 0.28)] and

40%30s [-0.09 (-0.19, 0.02) pps, *d* = -0.16 (-0.34, 0.03)].

Recruitment thresholds

A time by condition interaction effect was observed β = -0.89 (-1.17, -0.60) %

of peak torque, SE = 0.15; t = -6.01]. Recruitment thresholds increased from before

to after the intervention only on 40%15s [0.51 (0.29, 0.73) % of peak torque, *d* = 0.42

(0.24, 0.60)] and 40%30s [0.31 (0.09, 0.54) % of peak torque, *d* = 0.26 (0.07, 0.44)]

conditions, and decreased on control [-0.38 (-0.57, -0.19) % of peak torque, *d* = -0.31

(-0.47, -0.16)] condition, but remained unchanged on 80%15s [-0.06 (-0.26, 0.14) %

of peak torque, *d* = -0.05 (-0.21, 0.11)] condition.

Table 2 describes the before and after estimated marginal means and mean

differences for each MU outcome and tested conditions.

Table 2. Estimated marginal means and mean differences (95% confidence interval lower and upper limits) for ΔF, peak discharge rates, recruitment thresholds, brace height, and attenuation slope for handgrip and control conditions.

	ΔF (pps)	Brace height (% rTri)	Attenuation slope (pps/%MVT)	Peak discharge rate (pps)	Recruitment thresholds (% of peak torque)
Control					
Before	5.29 (4.90, 5.69)	38.2 (36.1, 40.3)	0.40 (0.36, 0.44)	17.2 (16.1, 18.2)	8.54 (7.16, 9.92)
After	5.22 (4.82, 5.62)	38.1 (36.0, 40.2)	0.41 (0.38, 0.45)	16.9 (15.8, 17.9)	8.16 (6.79, 9.54)
After-before	-0.07 (−0.21, 0.06)	-0.11 (−1.84, 1.61)	0.01 (−0.01, 0.04)	**-0.29 (−0.38, −0.21)**	**-0.38 (−0.57, −0.19)**
40%15s					
Before	5.64 (5.22, 6.05)	39.3 (37.0, 41.6)	0.40 (0.36, 0.44)	17.1 (16.0, 18.1)	8.08 (6.66, 9.49)
After	5.73 (5.32, 6.15)	41.2 (38.9, 43.4)	0.39 (0.35, 0.42)	17.1 (16.1, 18.2)	8.58 (7.17, 10.00)
After-before	0.10 (−0.06, 0.26)	1.86 (−0.15, 3.87)	−0.01 (−0.04, 0.02)	0.06 (−0.04, 0.16)	**0.51 (0.29, 0.73)**
40%30s					
Before	5.49 (5.07, 5.91)	35.4 (33.1, 37.7)	0.46 (0.42, 0.49)	17.4 (16.3, 18.4)	8.32 (6.91, 9.74)
After	5.81 (5.39, 6.24)	37.7 (35.4, 39.9)	0.41 (0.37, 0.45)	17.3 (16.2, 18.3)	8.63 (7.22, 10.05)
After-before	**0.33 (0.16, 0.49)**	**2.24 (0.18, 4.30)**	**−0.05 (−0.08, −0.01)**	-0.09 (−0.19, 0.02)	**0.31 (0.09, 0.54)**
80%15s					
Before	5.46 (5.05, 5.86)	37.3 (35.2, 39.5)	0.412 (0.38, 0.45)	17.4 (16.4, 18.5)	9.00 (7.61, 10.40)
After	5.69 (5.29, 6.10)	39.8 (37.6, 42.0)	0.39 (0.35, 0.42)	17.6 (16.6, 18.7)	8.94 (7.55, 10.33)
After-before	**0.24 (0.09, 0.38)**	**2.45 (0.64, 4.25)**	**−0.03 (−0.06, −0.00)**	**0.20 (0.11, 0.29)**	-0.06 (−0.26, 0.14)

389 ΔF, Δ frequency; % rTri, percentage of the right triangle; MVT, maximum voluntary torque. Bold values indicate significant changes
390 (95%CI not including zero).

Figure 3. A, ΔF before and after the Control, handgrip to 40% of maximal isometric force for 15 s (40%15s) or 30 s (40%30s), and to 80% of maximal isometric force for 15 s (80%15s) conditions.

The estimated marginal means (black circles) and respective 95% confidence intervals are offset to the middle. Individual data points (averaged ΔF per participant) are coloured by participants and individual test units value are plotted in light grey. pps, pulses per second. ΔF remained unchanged before and after Control and 40%15s conditions but increased at 40%30s and 80%15s.

B, Estimated marginal mean differences (After – Before each condition) for ΔF

in the control, handgrip at 40% of maximal isometric force for 15 s or 30 s, and

at 80% of maximal isometric force for 15 s conditions.

Significant increases were observed on the handgrip at 40%30s and 80%15s

conditions (not crossing the "zero" red line. The thick inner line and the thin outer line

represent the 90% and 95% confidence intervals, respectively.

Figure 4. Top, brace height; bottom, attenuation slope; before and after the Control, handgrip to 40% of maximal isometric force for 15 s (40%15s) or 30 s (40%30s), and to 80% of maximal isometric force for 15 s (80%15s) conditions.

The estimated marginal means (black circles) and respective 95% confidence intervals are offset to the middle. Individual data points (averaged brace height and attenuation slope per participant) are coloured by participants and individual motor units value are

plotted in light grey. pps, pulses per second; MVT, maximum voluntary torque; % rTri,

percentage of the right triangle. Both brace height and attenuation slope remained

unchanged before and after Control and 40%15s conditions but, respectively,

increased and decreased at 40%30s and 80%15s. Note: the y-axis for attenuation

slope has been limited to 2 pps/%MVT to enhance the visualization of data points and

their 95% confidence intervals. A full-range version of the figure is available at

https://github.com/lugliara/PICs.

Discussion

We investigated whether some characteristics of the remote handgrip

contraction, including duration (40%15s vs 40%30s), the intensity of effort (40%15s

vs 80%15s), and the impulse (40%30s and 80%15s vs 40%15s) affect tibialis

anterior PICs. Additionally, quasi-geometric analyses were used to estimate the

contribution of neuromodulatory (i.e., brace height) and inhibitory (i.e., attenuation

slope) inputs to PIC responses. The main findings of this study were: 1) ΔF

increased after the matched higher handgrip impulse conditions; 2) brace height also

increased under the same conditions; 3) attenuation slope decreased consistently

after the higher handgrip impulse conditions. These findings suggest that the

mechanical impulse of a remote contraction was the primary driver of increased

tibialis anterior ΔF, likely due to enhanced neuromodulation and a shift in the

inhibitory pattern.

The changes we observed in ΔF, brace height, and attenuation slope between

the two highest handgrip impulse conditions (80%15s and 40%30s), compared to

40%15s and Control, support our hypothesis, indicating that a given impulse

threshold might be necessary to influence increases in motoneuron recruitment-

derecruitment hysteresis of tibialis anterior MUs. The small increase in ΔF after

handgrip contractions at the 40%30s ($d = 0.47$) and 80%15s ($d = 0.34$) conditions

were similar to the changes observed by Mackay et al., (2023) in soleus

motoneurons, $d = 0.30$ and Orssatto et al., (2022) in tibialis anterior motoneurons, d

$= 0.55$, both using a similar protocol to our 40%30s condition. Such changes on ΔF

can be influenced by either neuromodulatory and/or inhibitory input onto the

motoneurons (Beauchamp et al., 2023). The neuromodulatory input caused by the

presence of monoamines is the main mechanism controlling PICs activity. In

animals, voltage-clamp experiments provided strong evidence that PICs are led by

the activation of voltage-gated L-type Ca^{2+} (Eckert & Lux, 1976; Svirskis &

Hounsgaard, 1997) and Na^+ (Harvey et al., 2006; Schwindt & Crill, 1977) channels.

Subsequent studies showed the input of 5HT and NE activating specific G-protein

coupled receptors at motoneurons' dendrites and soma were responsible for

triggering intracellular signalling cascade that modulate the properties of these ion

voltage-gated channels (Harvey et al., 2006; Heckman et al., 2009; Perrier &

Hounsgaard, 2003). In humans, oral administration of amphetamine, which is

assumed to enhance the presynaptic release of NE, also increased estimates of ΔF

(Udina et al., 2010), suggesting the neuromodulation mechanism observed in

animals could also be present in humans. Moreover, Wei et al. (2014) demonstrated

that 5HT modulates motoneurons input-output gain by using selective 5HT reuptake

inhibitors and receptor blockers. More recent studies used multi-channel

electromyography (EMG) during voluntary isometric ramp-shaped contractions and

showed evidence of decreased MU excitability in presence of 5HT blockers by

measuring ΔF and/or others estimates of PIC activity (Goodlich et al., 2023, 2024).

Furthermore, studies found higher ΔF during ramped-shaped contractions at greater

intensity (Mackay et al., 2023; Orssatto et al., 2021), suggesting higher voluntary

muscle activity in humans is associated with higher PIC activity and, likely, higher

5HT and NE concentration into the spinal cord. Therefore, it is reasonable to assume

that the increase in ΔF observed in this study after handgrip contractions with higher

impulse could have been caused by a transient increase in monoamines

concentration within the spinal cord; however, this mechanism was not directly tested

in this study and future studies with drugs affecting monoamines concentration might

be able to address this mechanism.

We used brace height to estimate neuromodulation contribution to

motoneurons firing changes, since it was proposed to be influenced by

neuromodulation, likely resulting from increased serotonergic and/or noradrenergic

input (Beauchamp et al., 2023). Thus, the increases we observed in brace height at

the 40%30s (d = 0.20) and 80%15s (d = 0.22) conditions theoretically suggest

remote handgrip contraction induce greater neuromodulation. In addition, PICs are

highly sensitive to inhibition (Hultborn et al., 2003; Kuo et al., 2003), an important

feature to control unwanted movement that could occur as a consequence of

neuromodulation by the diffuse descending projections of the monoaminergic system

(Heckman et al., 2009). Previous studies observed inhibition responses at spinal

level through different stimuli, such as small changes in antagonist muscles length

(Hultborn et al., 2003; Hyngstrom et al., 2007), tendon vibration (Matthews, 1966;

Orssatto et al., 2022; Pearcey et al., 2022), passive muscle stretching (Trajano et al.,

2014), and voluntary co-contraction (Gomes et al., 2024), likely induced by activation

of a disynaptic inhibitory circuit activated by muscle spindle primary Ia afferents

(Crone et al., 1987; Kuffler et al., 1951). However, it is unlikely that changes in Ia

inhibition would elicit inhibitory inputs in the different conditions used in the present

study due to the consistency of the isometric ramp-shaped contraction we used

before and after the remote contraction. Inhibition input might also originates at

supraspinal or segmental neural mechanisms, like the observed by Renshaw cell

activity (Hultborn & Pierrot-Deseilligny, 1979) and those generated by nociceptive

inputs (Heckman et al., 2009). We used attenuation slope to estimate the influence

of inhibitory input since the inhibition pattern was identified as a significant predictor

of this measure (Beauchamp et al., 2023). The small and trivial decrease in

attenuation slope at 40%30s (d = -0.27) and 80%15s (d = -0.17) might indicate a

transition from a tonic inhibitory pattern to inhibitory commands that are more

reciprocal to excitation (i.e., push-pull excitation-inhibition synaptic control) (Škarabot

et al., 2023). However, caution is warranted before attributing this decrease solely to

reduced inhibitory input, as ΔF sensitivity to inhibition patterns depends on

neuromodulation levels (Beauchamp et al., 2023). Additionally, MUs exhibiting larger

brace height would theoretically be associated with lower attenuation slope

(Mesquita et al., 2024). Thus, our findings suggest that, once a certain level of

impulse is reached, remote contractions performed at either lower or higher

intensity/duration result in similar increase in the estimated contribution of PICs to

motoneuron self-sustained firing (ΔF). This effect was likely driven by increased

neuromodulation onto the spinal cord, as indicated by the small increase observed in

brace height, while attenuation slope might reflect a shift in the pattern of inhibitory

input.

Complementarily, we investigated peak discharge rates and recruitment

threshold since both of which are associated with motoneuron excitability and might

also be influenced by PIC activity (Mesquita et al., 2024; Orssatto et al., 2022).

Moderate and small decreases in peak discharge rates (d = -0.52) and recruitment

threshold (d = -0.38) were observed under the Control condition, contrasting with

previous studies using a similar protocol, where no changes in Control were reported

for tibialis anterior (Orssatto et al., 2022) or soleus motoneurons (Mackay Phillips et

al., 2023). Unlike these studies, participants in our study performed three distinct

handgrip tasks in a randomized order with 5-minute intervals, which may have

mitigated task carryover effects on ΔF but not on peak discharge rates or recruitment

threshold. Furthermore, we found a small increase (d = 0.35) in peak discharge rates

only on 80%15s condition, whereas recruitment threshold increased in both the

40%15s (d = 0.42) and 40%30s (d = 0.26) conditions. Previous studies reported

small increases in peak discharge rates under the 40%30s condition (Orssatto et al.,

(2022): d = 0.36; Mackay Phillips et al., (2023): d = 0.37), but while Orssatto et al.

(2022) observed no changes in recruitment threshold, Mackay et al. (2023) did not

report recruitment threshold results. These findings suggest that PICs may

differentially modulate MU ΔF, peak discharge rates, and recruitment threshold, with

these variables providing distinct insights into motoneuron excitability.

Final considerations and future directions

This study combined robust non-invasive methods to estimate the influence of

motor task on PIC activity in humans, likely via increased 5HT input to the spinal

cord. Using an innovative EMG protocol allowed us to non-invasively assess key

metrics of motoneuron excitability (e.g., ΔF, brace height and attenuation slope)

without pharmacological intervention (Beauchamp et al., 2023). We also minimized

the potential for type I error through rigorous statistical analysis (Yu et al., 2022). On

the other hand, we assessed tibialis anterior MUs recruited during a ramp-shaped

contraction to 20% of maximal force. Thus, caution should be taken when

extrapolating our findings beyond lower-threshold units, at lower intensity

contractions, and from the tibialis anterior muscle. Additionally, direct measurements

of 5HT concentration in the spinal cord in vivo remain technically impossible, and we

must still rely on indirect methods to infer its effects. Lastly, while we focused on 5HT

and NE contributions to PICs, the potential influence of other neuromodulators

should be considered in future studies. For example, evidence suggests that spinal

interneurons may modulate motoneuron excitability via muscarinic m2 receptor

activation, affecting resting K^+ conductance (Miles et al., 2007).

Future studies could incorporate additional methods alongside EMG to clarify

findings further. For instance: (1) given that both 40%30s and 80%15s conditions

successfully increased ΔF in young adults, similar protocols could assess

neuromodulatory capacity in populations with potentially reduced 5HT input or

impaired PIC activation, such older adults, individuals with neurodegenerative

diseases, or partial spinal cord injuries; (2) further investigations are desirable

whether impulse-specific effects on PIC modulation differ across muscle groups, as

variations in motoneuron distribution may affect responsiveness to remote

contraction-induced neuromodulation;

Conclusion

This study provides novel insights into the mechanisms through which remote

voluntary muscle contractions influence motoneuron excitability, estimated by ΔF in

tibialis anterior motor units. By manipulating both the intensity and duration of a

handgrip task, we identified contraction impulse as a critical factor driving changes in

motoneuron recruitment-derecruitment hysteresis, which likely reflects in increased

PICs activity. Specifically, higher ΔF values in remote task conditions with matched

contraction impulse suggest that both higher and lower intensity/duration conditions

can lead to increased PIC modulation, provided that a sufficient impulse threshold is

met. The increased brace height observed in the impulse-matched conditions

support our hypothesis that increased tibialis anterior ΔF following handgrip remote

contraction may result from increased neuromodulation, possibly by augmented

serotonergic input onto the spinal cord.

References

Afsharipour, B., Manzur, N., Duchcherer, J., Fenrich, K. F., Thompson, C. K., Negro,

F., Quinlan, K. A., Bennett, D. J., & Gorassini, M. A. (2020). Estimation of self-

sustained activity produced by persistent inward currents using firing rate

profiles of multiple motor units in humans. *Journal of Neurophysiology*, *124*(1),

63–85. https://doi.org/10.1152/jn.00194.2020

Beauchamp, J. A., Khurram, O. U., Dewald, J. P. A., Heckman, C., & Pearcey, G. E. P. (2022). A computational approach for generating continuous estimates of motor unit discharge rates and visualizing population discharge characteristics. *Journal of Neural Engineering*, *19*(1), 016007. https://doi.org/10.1088/1741-2552/ac4594

Beauchamp, J. A., Pearcey, G. E. P., Khurram, O. U., Chardon, M., Wang, Y. C., Powers, R. K., Dewald, J. P. A., & Heckman, C. (2023). A geometric approach to quantifying the neuromodulatory effects of persistent inward currents on individual motor unit discharge patterns. *Journal of Neural Engineering*, *20*(1), 016034. https://doi.org/10.1088/1741-2552/acb1d7

Crone, C., Hultborn, H., Jespersen, B., & Nielsen, J. (1987). Reciprocal Ia inhibition between ankle flexors and extensors in man. *The Journal of Physiology*, *389*(1), 163–185. https://doi.org/10.1113/jphysiol.1987.sp016652

D'Amico, J. M., Murray, K. C., Li, Y., Chan, K. M., Finlay, M. G., Bennett, D. J., & Gorassini, M. A. (2013). Constitutively active 5-HT$_2$ /α_1 receptors facilitate muscle spasms after human spinal cord injury. *Journal of Neurophysiology*, *109*(6), 1473–1484. https://doi.org/10.1152/jn.00821.2012

Del Vecchio, A., Holobar, A., Falla, D., Felici, F., Enoka, R. M., & Farina, D. (2020). Tutorial: Analysis of motor unit discharge characteristics from high-density surface EMG signals. *Journal of Electromyography and Kinesiology*, *53*. https://doi.org/10.1016/j.jelekin.2020.102426

Del Vecchio, A., Negro, F., Holobar, A., Casolo, A., Folland, J. P., Felici, F., & Farina, D. (2019). You are as fast as your motor neurons: Speed of recruitment and maximal discharge of motor neurons determine the maximal rate of force development in humans. *The Journal of Physiology*, *597*(9), 2445–2456. https://doi.org/10.1113/JP277396

Eckert, R., & Lux, H. D. (1976). A voltage-sensitive persistent calcium conductance in neuronal somata of Helix. *The Journal of Physiology*, *254*(1), 129–151. https://doi.org/10.1113/jphysiol.1976.sp011225

Fenstermacher, S. J., Vonasek, A., Gattuso, H., Chaimowitz, C., Dymecki, S. M., Jessell, T. M., & Dasen, J. S. (2024). *Potentiation of active locomotor state by spinal-projecting serotonergic neurons*. https://doi.org/10.1101/2024.09.26.615260

Gomes, M. M., Jenz, S. T., Beauchamp, J. A., Negro, F., Heckman, C. J., & Pearcey, G. E. P. (2024). *Voluntary co-contraction of ankle muscles alters motor unit discharge characteristics and reduces estimates of persistent inward currents*. https://doi.org/10.1101/2024.02.28.582534

Goodlich, B. I., Del Vecchio, A., Horan, S. A., & Kavanagh, J. J. (2023). Blockade of 5-HT$_2$ receptors suppresses motor unit firing and estimates of persistent inward currents during voluntary muscle contraction in humans. *The Journal of Physiology*, *601*(6), 1121–1138. https://doi.org/10.1113/JP284164

Goodlich, B. I., Pearcey, G. E. P., Del Vecchio, A., Horan, S. A., & Kavanagh, J. J. (2024). Antagonism of 5-HT$_2$ receptors attenuates self-sustained firing of human motor units. *The Journal of Physiology*, *602*(8), 1759–1774. https://doi.org/10.1113/JP285867

Harvey, P. J., Li, X., Li, Y., & Bennett, D. J. (2006). Endogenous Monoamine Receptor Activation Is Essential for Enabling Persistent Sodium Currents and Repetitive Firing in Rat Spinal Motoneurons. *Journal of Neurophysiology*, *96*(3), 1171–1186. https://doi.org/10.1152/jn.00341.2006

Hassan, A., Thompson, C. K., Negro, F., Cummings, M., Powers, R. K., Heckman, C. J., Dewald, J. P. A., & McPherson, L. M. (2020). Impact of parameter selection on estimates of motoneuron excitability using paired motor unit analysis. *Journal of Neural Engineering*, *17*(1). https://doi.org/10.1088/1741-2552/ab5eda

Heckman, C. J., Hyngstrom, A. S., & Johnson, M. D. (2008). Active properties of motoneurone dendrites: Diffuse descending neuromodulation, focused local inhibition. *Journal of Physiology*, *586*(5), 1225–1231. https://doi.org/10.1113/jphysiol.2007.145078

Heckman, C. J., Johnson, M., Mottram, C., & Schuster, J. (2008). Persistent Inward Currents in Spinal Motoneurons and Their Influence on Human Motoneuron Firing Patterns. *The Neuroscientist*, *14*(3), 264–275. https://doi.org/10.1177/1073858408314986

Heckman, C. J., Mottram, C., Quinlan, K., Theiss, R., & Schuster, J. (2009). Motoneuron excitability: The importance of neuromodulatory inputs. *Clinical Neurophysiology*, *120*(12), 2040–2054. https://doi.org/10.1016/j.clinph.2009.08.009

Holobar, A., Minetto, M. A., & Farina, D. (2014). Accurate identification of motor unit discharge patterns from high-density surface EMG and validation with a novel signal-based performance metric. *Journal of Neural Engineering*, *11*(1). https://doi.org/10.1088/1741-2560/11/1/016008

Holobar, A., & Zazula, D. (2007). Multichannel blind source separation using convolution Kernel compensation. *IEEE Transactions on Signal Processing*, *55*(9), 4487–4496. https://doi.org/10.1109/TSP.2007.896108

Hounsgaard, J., Hultborn, H., Jespersen, B., & Kiehn, O. (1988). Bistability of alpha-motoneurones in the decerebrate cat and in the acute spinal cat after intravenous 5-hydroxytryptophan. *The Journal of Physiology*, *405*(1), 345–367. https://doi.org/10.1113/jphysiol.1988.sp017336

Hultborn, H., Denton, M. E., Wienecke, J., & Nielsen, J. B. (2003). Variable amplification of synaptic input to cat spinal motoneurones by dendritic persistent inward current. *The Journal of Physiology*, *552*(3), 945–952. https://doi.org/10.1113/jphysiol.2003.050971

Hultborn, H., & Pierrot-Deseilligny, E. (1979). Changes in recurrent inhibition during voluntary soleus contractions in man studied by an H-reflex technique. *The Journal of Physiology*, *297*(1), 229–251. https://doi.org/10.1113/jphysiol.1979.sp013037

Hyngstrom, A. S., Johnson, M. D., Miller, J. F., & Heckman, C. J. (2007). Intrinsic electrical properties of spinal motoneurons vary with joint angle. *Nature Neuroscience*, *10*(3), 363–369. https://doi.org/10.1038/nn1852

Jacobs, B. L., Martín-Cora, F. J., & Fornal, C. A. (2002). Activity of medullary serotonergic neurons in freely moving animals. *Brain Research Reviews*, *40*(1–3), 45–52. https://doi.org/10.1016/S0165-0173(02)00187-X

Johnson, M. D., & Heckman, C. J. (2014). Gain control mechanisms in spinal motoneurons. *Frontiers in Neural Circuits*, *8*(JULY). https://doi.org/10.3389/fncir.2014.00081

Kuffler, S. W., Hunt, C. C., & Quilliam, J. P. (1951). Function of medullated small-nerve fibers in mammalian ventral roots: Efferent muscle spindle innervation. *Journal of Neurophysiology*, *14*(1), 29–54. https://doi.org/10.1152/jn.1951.14.1.29

Kuo, J. J., Lee, R. H., Johnson, M. D., Heckman, H. M., & Heckman, C. J. (2003). Active Dendritic Integration of Inhibitory Synaptic Inputs In Vivo. *Journal of Neurophysiology*, *90*(6), 3617–3624. https://doi.org/10.1152/jn.00521.2003

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in Linear Mixed Effects Models. *Journal of Statistical Software*, *82*(13), 1–26. https://doi.org/10.18637/JSS.V082.I13

Lee, R. H., & Heckman, C. J. (2000). Adjustable Amplification of Synaptic Input in the Dendrites of Spinal Motoneurons *In Vivo*. *The Journal of Neuroscience*, *20*(17), 6734–6740. https://doi.org/10.1523/JNEUROSCI.20-17-06734.2000

Lenth, R. V., Bolker, B., Buerkner, P., Giné-Vázquez, I., Herve, M., Jung, M., Love, J., Miguez, F., Riebl, H., & Singmann, H. (2023). *Emmeans: Estimated marginal means, aka Least-Squares Means—R package*. https://github.com/rvlenth/emmeans

Mackay, K., Orssatto, L. B. R., Polman, R., Van Der Pols, J. C., & Trajano, G. S. (2023). Caffeine does not influence persistent inward current contribution to motoneuron firing. *Journal of Neurophysiology*, *130*(6), 1529–1540. https://doi.org/10.1152/jn.00350.2023

Mackay Phillips, K., Orssatto, L. B. R., Polman, R., Van Der Pols, J. C., & Trajano, G. S. (2023). The effects of α-lactalbumin supplementation and handgrip contraction on soleus motoneuron excitability. *European Journal of Applied Physiology*, *123*(2), 395–404. https://doi.org/10.1007/s00421-022-05101-3

Matthews, P. B. C. (1966). The reflex excitation of the soleus muscle of the decerebrate cat caused by vibration applied to its tendon. *The Journal of Physiology*, *184*(2), 450–472. https://doi.org/10.1113/jphysiol.1966.sp007926

Mesquita, R. N. O., Taylor, J. L., Heckman, C. J., Trajano, G. S., & Blazevich, A. J. (2024). Persistent inward currents in human motoneurons: Emerging evidence and future directions. *Journal of Neurophysiology*, *132*(4), 1278–1301. https://doi.org/10.1152/jn.00204.2024

Miles, G. B., Hartley, R., Todd, A. J., & Brownstone, R. M. (2007). Spinal cholinergic interneurons regulate the excitability of motoneurons during locomotion. *Proceedings of the National Academy of Sciences*, *104*(7), 2448–2453. https://doi.org/10.1073/pnas.0611134104

Orssatto, L. B. R., Fernandes, G. L., Blazevich, A. J., & Trajano, G. S. (2022). Facilitation–inhibition control of motor neuronal persistent inward currents in

young and older adults. *Journal of Physiology*.

https://doi.org/10.1113/JP283708

Orssatto, L. B. R., Mackay, K., Shield, A. J., Sakugawa, R. L., Blazevich, A. J., &

Trajano, G. S. (2021). Estimates of persistent inward currents increase with

the level of voluntary drive in low-threshold motor units of plantar flexor

muscles. *Journal of Neurophysiology*, *125*(5), 1746–1754.

https://doi.org/10.1152/jn.00697.2020

Pearcey, G. E. P., Khurram, O. U., Beauchamp, J. A., Negro, F., & Heckman, C. J.

(2022). *Antagonist tendon vibration dampens estimates of persistent inward*

currents in motor units of the human lower limb.

https://doi.org/10.1101/2022.08.02.502526

Perrier, J.-F., & Hounsgaard, J. (2003). 5-HT$_2$ Receptors Promote Plateau Potentials

in Turtle Spinal Motoneurons by Facilitating an L-Type Calcium Current.

Journal of Neurophysiology, *89*(2), 954–959.

https://doi.org/10.1152/jn.00753.2002

Schwindt, P., & Crill, W. E. (1977). A persistent negative resistance in cat lumbar

motoneurons. *Brain Research*, *120*(1), 173–178. https://doi.org/10.1016/0006-

8993(77)90510-8

Škarabot, J., Beauchamp, J. A., & Pearcey, G. E. (2023). *Human motor unit*

discharge patterns reveal differences in neuromodulatory and inhibitory drive

to motoneurons across contraction levels [Preprint]. Neuroscience.

https://doi.org/10.1101/2023.10.16.562612

Svirskis, G., & Hounsgaard, J. (1997). Depolarization-Induced Facilitation of a

Plateau-Generating Current in Ventral Horn Neurons in the Turtle Spinal Cord.

Journal of Neurophysiology, *78*(3), 1740–1742.

https://doi.org/10.1152/jn.1997.78.3.1740

Thorstensen, J. R., Henderson, T. T., & Kavanagh, J. J. (2024). Serotonergic and

noradrenergic contributions to motor cortical and spinal motoneuronal

excitability in humans. *Neuropharmacology*, *242*, 109761.

https://doi.org/10.1016/j.neuropharm.2023.109761

Trajano, G. S., Seitz, L. B., Nosaka, K., & Blazevich, A. J. (2014). Can passive

stretch inhibit motoneuron facilitation in the human plantar flexors? *Journal of*

Applied Physiology, *117*(12), 1486–1492.

https://doi.org/10.1152/japplphysiol.00809.2014

Udina, E., D'Amico, J., Bergquist, A. J., & Gorassini, M. A. (2010). Amphetamine
increases persistent inward currents in human motoneurons estimated from
paired motor-unit activity. *Journal of Neurophysiology*, *103*(3), 1295–1303.
https://doi.org/10.1152/jn.00734.2009

Vandenberk, M. S., & Kalmar, J. M. (2014). An evaluation of paired motor unit
estimates of persistent inward current in human motoneurons. *J Neurophysiol*,
111, 1877–1884. https://doi.org/10.1152/jn.00469.2013.-Persistent

Veasey, S., Fornal, C., Metzler, C., & Jacobs, B. (1995). Response of serotonergic
caudal raphe neurons in relation to specific motor activities in freely moving
cats. *The Journal of Neuroscience*, *15*(7), 5346–5359.
https://doi.org/10.1523/JNEUROSCI.15-07-05346.1995

Wei, K., Glaser, J. I., Deng, L., Thompson, C. K., Stevenson, I. H., Wang, Q., Hornby,
T. G., Heckman, C. J., & Kording, K. P. (2014). Serotonin affects movement
gain control in the spinal cord. *Journal of Neuroscience*, *34*(38), 12690–
12700. https://doi.org/10.1523/JNEUROSCI.1855-14.2014

Yu, Z., Guindani, M., Grieco, S. F., Chen, L., Holmes, T. C., & Xu, X. (2022). Beyond
t test and ANOVA: Applications of mixed-effects models for more rigorous
statistical analysis in neuroscience research. *Neuron*, *110*(1), 21–35.
https://doi.org/10.1016/j.neuron.2021.10.030

760