Histone deacetylase-1 is required for epigenome stability in *Neurospora crassa*

Felicia Ebot-Ojong[1], Aileen R. Ferraro[1], Farh Kaddar[2], Clayton Hull-Crew[2], Ashley W. Scadden[2],

Andrew D. Klocko[2], and Zachary A. Lewis[1‡]

[1]Department of Microbiology, University of Georgia, Athens, GA, 30602 USA

[2]Department of Chemistry & Biochemistry, University of Colorado Colorado Springs, Colorado

Springs, CO, 80918, USA

[‡]Corresponding Author (zlewis@uga.edu)

Keywords: constitutive heterochromatin, facultative heterochromatin, Polycomb Repressive Complex

2, H3K27me3, histone deacetylase, histone post-translational modifications

Abstract

Polycomb group (PcG) proteins form chromatin modifying complexes that stably repress

lineage- or context-specific genes in animals, plants, and some fungi. Polycomb Repressive Complex 2

(PRC2) catalyzes trimethylation of lysine 27 on histone H3 (H3K27me3) to assemble repressive

chromatin. In the model fungus *Neurospora crassa,* H3K27me3 deposition is controlled by the H3K36

methyltransferase ASH1 and components of constitutive heterochromatin including the H3K9me3-

binding protein HETEROCHROMATIN PROTEIN 1 (HP1). Hypoacetylated histones are a defining

feature of both constitutive heterochromatin and PcG-repressed chromatin, but how histone

deacetylases (HDACs) contribute to normal H3K27me3 and transcriptional repression within PcG-

repressed chromatin is poorly understood. We performed a genetic screen to identify HDACs required

for repression of PRC2-methylated genes. In the absence of HISTONE DEACETYLASE-1 (HDA-1),

PRC2-methylated genes were activated and H3K27me3 was depleted from typical PRC2-targeted

regions. At constitutive heterochromatin, HDA-1 deficient cells displayed reduced H3K9me3,

hyperacetylation, and aberrant enrichment of H3K27me3 and H3K36me3. CHROMODOMAIN

PROTEIN-2 (CDP-2) is required to target HDA-1 to constitutive heterochromatin and was also

required for normal H3K27me3 patterns. Patterns of aberrant H3K27me3 were distinct in isogenic

Δhda-1 strains, suggesting that loss of HDA-1 causes stochastic or progressive epigenome dysfunction.

To test this, we constructed a new *Δhda-1* strain and performed a laboratory evolution experiment.

Deletion of *hda-1* led to progressive epigenome decay over hundreds of nuclear divisions. Together,

our data indicate that HDA-1 is a critical regulator of epigenome stability in *N. crassa.*

Introduction

Formation of specialized heterochromatin is directed by histone post-translational modifications (PTMs) (1). In some cases, repressive chromatin states can be epigenetically inherited over the lifetime of an organism. Different types of repressive chromatin domains are enriched for characteristic histone modifications and have distinct functions. Constitutive heterochromatin (cHet) is found at repeat-rich regions of the genome and is typically enriched for di- or tri-methylated lysine-9 on histone H3 (H3K9me2/3) (1). A second type of repressed chromatin is assembled and maintained by Polycomb Group (PcG) proteins, which interact to form Polycomb Repressive Complex (PRC) 1 and PRC2 (2–4). PRC2 methylates histone H3 lysine 27 in PcG-repressed chromatin, which is often referred to as facultative heterochromatin (fHet) because PcG-repressed genes are expressed in context- or cell-type-specific manner (5–7). Notably, hypoacetylated histones are a characteristic feature of both cHet and fHet domains (1, 8).

Defective assembly or maintenance of heterochromatin is associated with human diseases including cancer, highlighting the critical need to define genes and mechanisms that structure the epigenome (9, 10). The PcG proteins have been intensely studied in diverse experimental systems, revealing that numerous interconnected regulatory mechanisms establish and maintain H3K27me3-enriched domains. In plants and mammals, accessory proteins can recruit PRC1 or PRC2 to target sites (2, 3, 11–15). In addition, the activity of PRC2 is modulated by chromatin features, including activating chromatin modifications (16–18), inhibitory modifications (19–21), RNA (22–25), and densely packed chromatin (26).

The fungus *Neurospora crassa* encodes conserved components of both cHet and fHet pathways and is a well-established model system to investigate assembly and maintenance of heterochromatin (5, 27). Prior studies with *N. crassa* showed that normal H3K27 methylation depends on core PRC2 components (28) and a variety of additional factors including H2A.Z (29), accessory proteins (CAC-

3/NPF, EPR-1, and PAS) (28, 30, 31), the chromatin remodeler IMITATION SWITCH (ISWI) (32,

33), and other histone modifications such as H3K36me3 (34), and H3K9me3 (35, 36). In contrast to

mammalian systems, H3K36 methylation frequently co-occurs with and can promote H3K27me3 in

fungi (37, 34, 38, 39). Thus, PRC2 localization and activity is influenced by diverse histone

modifications in *N. crassa*, like in higher eukaryotes, but much remains unknown about how various

modifications cooperate to structure the epigenome.

In plants, animals, and fungi, molecular features of cHet antagonize H3K27me3 (40). In wild

type *N. crassa,* cHet domains are defined as repeat-rich, gene-poor DNA sequences enriched for

H3K9me3, HP1, DNA cytosine methylation (5mC), and hypoacetylated histones (41, 27, 42). cHet

domains depend on multiple HP1-containing complexes that bind to H3K9me3 and modulate cHet

structure by directing histone deacetylation, 5mC deposition, or by recruiting the anti-silencing factor

DNA Methylation Modulator 1 (DMM-1) (43–47, 42, 48). In *N. crassa* mutants lacking H3K9me3 or

HP1, H3K27me2/3 is lost from native PRC2 target domains and aberrantly gained at cHet regions (35,

36). Similar redistribution of H3K27me3 is observed in mammalian cells depleted for H3K9me3 or

5mC (49–52). In mutant mouse embryonic stem cells (mESCs) lacking 5mC, a non-canonical PRC1

complex (ncPRC1.1) binds to hypomethylated CpG-rich DNA in pericentromeric cHet regions and

deposits ubiquitylated H2A (H2Aub), leading to PRC2 recruitment and H3K27 methylation of

H3K9me3-depleted histones (53, 54). Thus, features of mammalian cHet domains modulate

H3K27me3 localization in at least two ways; 5mC prevents aberrant recruitment of PRC1 and PRC2 to

mammalian cHet domains, whereas H3K9me3 inhibits PRC2 activity at these regions. In contrast to

mammals, loss of 5mC does not impact H3K27me2/3 patterns in *N. crassa* (35, 36), and how

H3K9me3 and HP1 exclude H3K27me3 from cHet regions is not understood. In addition, it is unclear

if other components of cHet domains, such as histone deacetylases, impact H3K27me3 localization in

any system.

Here, we show that HDA-1, a cHet-specific histone deacetylase, is required for repression of

PRC2-methylated genes in *N. crassa.* HDA-1 is a component of the <u>H</u>P1/<u>C</u>DP-2/<u>H</u>DA-1/<u>C</u>HAP

(HCHC) complex, which limits DNA accessibility at cHet domains and is critical for normal three-

dimensional (3D) chromosome conformation (42, 48, 55, 56). In cells lacking HCHC components, we

found that H3K27me3 is lost from typical PRC2-methylated genes (fHet) and gained at cHet regions.

We observed distinct patterns of H3K27me3 in strains lacking individual HCHC components and in

individual Δ*hda-1* strain isolates, raising the possibility that loss of HDA-1 targeting or activity leads

to stochastic or progressive changes to the epigenome. To test this hypothesis, we constructed a new

Δ*hda-1* strain and performed a lab evolution experiment. In this new Δ*hda-1* strain, changes in the

pattern of H3K27me3 were progressive, taking place over hundreds of nuclear divisions, and were

accompanied by altered lysine acetylation, H3K36me3, and H3K9me3. Together, our results show that

HCHC is a critical regulator of epigenome stability in *N. crassa*.

Materials and Methods

Strains, strain construction, and growth conditions

Strains and primers are listed in Tables S1 and S2, respectively. Strains were grown in Vogel's

minimal medium (VMM) with 1.5% sucrose or with 0.05% fructose, 0.05% glucose, and 2% sorbose

as described (57). Media was supplemented with 25 mg/mL histidine or 200µg/mL hygromycin B

where appropriate. Gene deletion strains were obtained from the *Neurospora* knockout collection (58),

unless otherwise indicated. The NKL9 and NKL10 Δ*hda-1* strains were generated in the Klocko lab by

backcrossing an Δ*hda-1* strain (FGSC 12003) to wild type and passaging the resulting progeny. A new

hda-1 knockout mutant, strain S863, was constructed for this study using a split marker approach (59).

Partial deletion fragments were generated by polymerase chain reaction (PCR) using genomic DNA

from an existing Δ*hda-1* strain (S297) as a template using primer pairs *hda*-1 5' flank FP + *hph* SM-1

and *hda-1* 3' flank RP + *hph* SM-2. Fragments were transformed into the wild type strain S2 by

electroporation as described (60). To perform complementation experiments, we first generated

plasmid pBM61_HDA-1-3xFLAG by cloning synthetic gene fragments (Twist Biosciences)

corresponding to the wild type *hda-1-3xflag* sequence under control of the native *hda-1* promoter

(546bp upstream of the annotated transcription start site) into the plasmid backbone of pBM61-

3xFLAG amplified by PCR using primers pBM61_Inverse P3 + pBM61_Inverse P2. A plasmid

containing a presumptive catalytic dead *hda-1^{h225A}-3xflag* allele was constructed in the same way.

Plasmid sequences were verified by sequencing (Plasmidsaurus). Both constructs were introduced into

the *his-3* locus of strain S188 (*his-3; Δhda-1*) as described (60).

***Chromatin Immunoprecipitation (ChIP), CUT&RUN,* RNA-seq, and Hi-C**

All antibodies used in this study are reported in Table S3. ChIP-seq was performed as described (56,

61, 62). For RNA-seq experiments, RNA isolation and RNA-seq library preparation was performed as

described (Kamei et al., 2021; Zhou et al., 2018). We adapted CUT&RUN for *N. crassa* by modifying

a previously published protocol (64). For Neurospora CUT&RUN, cells were grown in 5mL VMM +

1.5% sucrose for 16-18 hours. Mycelial mats were harvested using a Buchner funnel and washed with

10mL of PBS. Washed mycelial mats were transferred to a clean petri dish, flooded with 1mL of

nuclear isolation buffer (20 mM HEPES pH 7.5, 0.5 mM Spermine tetrahydrochloride, 80 mM KCl, 20

mM NaCl, 0.1 mM CaCl$_2$, 15 mM 2-Mercaptoethanol, 0.3% Triton X-100), and chopped with a razor

blade for 2 minutes. The suspension of mycelial fragments was filtered through Miracloth (Millipore-

Sigma, Catalog # 475855) into a 1.5 mL microfuge tube and centrifuged for 5 minutes at 1000 x G to

remove cell debris. The nuclei-containing supernatant was subject to CUT&RUN using an established

procedure (65).

In situ Hi-C was performed using crosslinked nuclei. Briefly, 25mL cultures of 1x Vogel's and

1.5% sucrose were started from ~1.0 x 10^8 conidia and grown overnight at 32ºC with shaking at 200

rpm. Cultures were crosslinked with 1% [final] formaldehyde, quenched with 125mM Tris-HCl, pH

8.0, and the mycelial mat was isolated by vacuum filtration and placed in a petri dish on ice. Nuclear

isolation buffer (5mL; 20mM HEPES pH 7.5, 80mM KCl, 20mM NaCl, 0.1mM $CaCl_2$, 15mM β-

mercaptoethanol, 0.3% Triton, and 0.5mM Spermine tetrahydrochloride [freshly added]) was added

and the mycelial mat was rapidly cut with a new razor blade. The lysate was filtered through

autoclaved Miracloth (Millipore Sigma, cat# 475855-1R) lining a funnel into a 15mL conical tube, and

nuclei were collected by centrifugation at 4,500 rpm for 5 minutes. The pellet was washed with an

additional 1mL nuclear isolation buffer, and nuclei pellets were stored at -80ºC. Nuclei comprising at

least 3.5 ug of DNA were then used for the remainder of the *in situ* Hi-C protocol using the restriction

enzyme *Mse*I to more efficiently capture AT-rich genomic loci, starting with 6.25% SDS treatment to

make the nuclei porous, as previously described (56). All DNA sequencing data was generated as

paired-end reads on a NextSeq 2000, NovaSeq 6000, or a NovoSeq X instrument.

Analysis of High-throughput sequencing data

For ChIP-seq, fastq files were trimmed to remove adaptor sequences using TrimGalore!

V0.6.(66), and then mapped to *N.crassa* assembly 12 (GCA_000182925.2) or assembly 14 for

comparison to Hi-C data (67) with BWA v0.7.17 mem (option -M) (68) or bowtie2 (69), respectively.

Reads were sorted and indexed using SAMtools v1.16.1 (70). Bigwig files for visualization were made

using deepTools v3.5.2 'bamCoverage' module (71), normalized by either BPM (bins per million

mapped reads) or RPKM (reads per kilobase per million mapped reads), while heatmaps of ChIP-seq

data were created using deepTools 'computeMatrix' and 'plotHeatmap' modules. Resulting bigwig

files were visualized with the Integrative Genome Viewer (72), and saved images were used for figure

creation. The 'csaw' and 'Complex Heatmap' packages in R were used to compute and plot

H3K27me3, H3K9me3, H3K36me3, or lysine acetylation (Kac) ChIP-seq enrichment values in wild

type and passaged Δ*hda-1* strains (73). Previously published ChIP-seq datasets used for this

manuscript include wild type H3K9me3 (NCBI GEO submission # GSE232933) and wild type

H3K27me2/3 (merged files from GSE68897 and GSE100770) (36, 56, 74).

For RNA-seq, reads were trimmed with TrimGalore v0.6.5 (66) and mapped using STAR

v2.7.10 (75). Output files in BAM format were indexed using SAMtools v1.9, and alignment counts

were determined using subRead v2.0.1 'featureCounts' (76). Differential expression was calculated

using DESeq2 (77). PRC2-targeted genes defined previously and are listed in (**Dataset S1**) (32).

For Hi-C data processing, output fastq datasets were mapped to the *N. crassa* reference

genome, version 14 (nc14) (67) with Bowtie2 (69), and contact matrices were built at a high (1 kb)

resolution using the "hicBuildMatrix" command of the hiCExplorer software package; lower

resolution (10 kb or 20 kb bin) matrices were generated with the hicMergeMatrixBins command. (78).

To compare to the previously published wild type *Mse*I Hi-C dataset, a number of total reads were

extracted from the NKL9 *Mse*I dataset that would give approximately equal numbers of valid reads for

comparison to a wild type N150 *Mse*I Hi-C dataset (or reads were extracted from the wild type *Mse*I

dataset when comparing to NKL10) using the sed command (67). After mapping reads and building

the contact matrix with the selected fastq files, the resulting contact matrices were compared using the

hicCompareMatrices command of hiCExplorer. Images were generated with the hicPlotMatrix

command from hiCExplorer and used for figure construction.

Results

HDA-1 is required for repression of PRC2-targeted genes

To determine if any histone deacetylases are required for transcriptional repression of genes in

facultative heterochromatin (fHet; i.e. H3K27me3-enriched domains), we tested available HDAC-

deficient mutants for defective gene silencing. We performed RNA-seq to quantify mRNA levels in

wild type and ten gene deletion strains lacking predicted histone deacetylase enzymes (79). The *N.*

crassa hda-3 gene is essential and was not included in this study (80). For all HDAC-deficient strains

tested, reproducible results were obtained with replicate RNA-seq samples (**Figure S1**). For each gene

deletion strain, we first examined average expression levels of previously defined PRC2-methylated

genes (32), which are tightly repressed in vegetative mycelia (Deaven et al. in preparation;

Supplemental Dataset 1). The Δ*hda-1* strain showed striking upregulation of PRC2-targeted genes

relative to wild type or other HDAC-deficient mutants (**Figure 1A**). The extent of upregulation in

Δ*hda-1* was similar to that observed for the Δ*set-7* control strain, which lacks all H3K27me3. A

heatmap plotting relative expression of each PRC2-targeted gene confirmed broad upregulation of this

gene set in Δ*hda-1* but not in other HDAC-deficient mutants (**Figure 1B**). In total, 2263 genes had

significant expression changes in the Δ*hda-1* mutant compared to wild type (**Figure 1C, Dataset S2;**

+/- 2-fold [Δ*hda-1/*WT], adjusted p-value of < 0.05). Of these, 1493 genes displayed increased

expression whereas 770 genes showed reduced expression levels compared to wild type. More than

30% of PRC2 target genes were significantly upregulated in the Δ*hda-1* by more than 2-fold (**Dataset**

S2). Importantly, genes encoding PRC2 complex members or other known components of the fHet

pathway did not exhibit altered expression in the Δ*hda-1* strain (**Dataset S2**). We conclude that HDA-1

is required for normal repression of ~15% of *N. crassa* genes, including a significant fraction of silent

PRC2-methylated genes located in fHet regions.

Histone modification patterns are altered in **the Δ*hda-1* strain**

To determine if HDA-1 is required for normal chromatin structure at fHet domains, we performed CUT&RUN to examine the distribution of H3K27me3 across the *N. crassa* genome. Because HDA-1 is reportedly targeted to cHet regions by HP1 (42), we also examined the distribution of H3K9me3. **Figure 2A** shows the distributions of H3K27me3 and H3K9me3 across Linkage Group III for wild type and Δ*hda-1*. We observed substantial reduction of H3K27me3 in fHet regions (i.e. domains enriched for H3K27me3 in a wild type strain). Loss of H3K27me3 at fHet domains was accompanied by gains of H3K27me3 in cHet regions (i.e. domains enriched for H3K9me3 in a wild type strain). For cHet domains that gained H3K27me3, we observed lower levels of H3K9me3 enrichment compared to cHet regions that did not gain H3K27me3 (**Figure 2A-B**). Replicate experiments produced similar results (**Figure S2A**). To examine the global changes in the repressive chromatin landscape, we generated heatmaps to visualize H3K27me3 and H3K9me3 levels across all wild type fHet and cHet domains (**Figure 2C-D**). This confirmed widespread loss of H3K27me3 from fHet domains, while ~ 20% of cHet regions gained H3K27me3 and lost H3K9me3.

To confirm that changes in histone PTMs are caused by loss of HDA-1, we introduced an *hda-1-3xflag* construct into an Δ*hda-1* deletion strain and performed ChIP-seq. Wild-type patterns of H3K27me3 and H3K9me3 were restored in this complemented strain (**Figure 2E-F**). Importantly, ChIP-seq data validated results obtained using CUT&RUN (**Figure 2A-D)**, which had not previously been adapted for *N. crassa.* We next asked if HDA-1 catalytic activity is required for normal patterns of H3K27me3 and H3K9me3. We generated a mutant *hda-1^{H225A}-3xflag* construct in which a histidine required for catalytic activity is mutated to alanine (81) and introduced this allele into Δ*hda-1*. This mutant *hda-1* allele failed to rescue H3K9me3 or H3K27me3 patterning defects (**Figure 2E,G**). Both epitope-tagged proteins were expressed at similar levels (**Figure S2B**), and replicate ChIP-seq experiments produced similar results (**Figure S2C**). We conclude that *hda*-1 is required for normal

patterns of both H3K27me3 and H3K9me3 in *N. crassa.*

HDA-1-deficient cells exhibit hyperacetylation at constitutive heterochromatin domains (42,

48, 56). To explore the relationship between lysine acetylation and altered fHet/cHet histone PTMs in

the Δ*hda-1* strain, we performed ChIP-seq using antibodies to pan acetyl-lysine (Kac), H3K27me3,

and H3K9me3. We used a pan-acetyl lysine antibody because previous work revealed a striking

increase in H2B acetylation in Δ*hda-1* strain (82), but no H2Bac antibodies are available for *N. crassa.*

In WT, low acetylation levels were observed in cHet domains marked by H3K9me3 and to a lesser

extent, in fHet domains marked by H3K27me3 (**Figures 3A; S3A**). These results are similar to those

obtained in prior studies using antibodies to acetylated H3 residues (42, 56, 80). In the Δ*hda-1* strain,

we observed hyperacetylation in cHet regions, such as the centromere, consistent with previous reports

of the HCHC targeting these loci (**Figure 3A, S3A**) (42).

In *N. crassa*, H3K27me3 co-occurs with ASH1-dependent H3K36 methylation. In fHet

regions, ASH-1 deposits H3K36me2, which is efficiently converted to H3K36me3 by SET-2 (34, 83).

SET-2 also methylates H3K36me1/2/3 in the coding sequences of actively transcribed genes within

euchromatin. We performed ChIP-seq to determine if HDA-1 is required for normal H3K36me3

localization (**Figure 3B**). Consistent with previous results, wild type H3K36me3 patterns were distinct

in euchromatin, fHet, and cHet regions. In euchromatin, H3K36me3 was restricted to gene bodies. In

fHet domains, H3K36me3 overlapped with H3K27me3 covering both promoters and gene bodies

across large, multi-gene regions. In cHet regions, no H3K36me3 enrichment was observed in wild

type. In the Δ*hda-1* strain, H3K36me3 was enriched in a subset of cHet domains, similar to the

aberrant pattern of H3K27me3 in this mutant. We examined the interrelationships of Kac, H3K27me3,

H3K36me3, and H3K9me3 by plotting enrichment of these PTMs across all wild type cHet domains

(**Figure 3C**). As expected, wild-type cHet regions were heavily enriched for H3K9me3 and devoid of

Kac, H3K36me3, and H3K27me3, except for small domains adjacent to each telomere that exhibit co-

enrichment of fHet- and cHet-specific PTMs (84). In Δ*hda-1*, individual cHet domains gained either

Kac or both H3K27me3 and H3K36me3. The subset of cHet domains that gained H3K27me3 and

H3K36me3 exhibited concomitant loss of H3K9me3 (~20% of cHet regions). In contrast, most cHet

domains (~80%) displayed typical enrichment of H3K9me3 accompanied by aberrant

hyperacetylation. The patterns of Kac and H3K36me3 in euchromatin were unaffected in the Δ*hda-1*

mutant, consistent with the known role of HDA-1 at cHet (**Figure S3C-D**). Replicate ChIP-seq

samples produced similar results (**Figure S3E**). Thus, loss of HDA-1 leads to aberrant deposition of

two modifications typically found at fHet regions, H3K27me3 and H3K36me3, in a subset of cHet

loci. In addition, reduced acetylation at these cHet loci suggests these fHet modifications can direct

recruit a compensatory deacetylase to cHet loci of Δ*hda-1* strains.

HCHC-deficient mutants exhibit distinct patterns of aberrant H3K27me3

HDA-1 is a component of the HP1-containing HCHC complex. HP1 was previously shown to exclude

H3K27me3 from cHet regions (35, 36). We next asked if other components of HCHC are required for

normal H3K27me3 localization. We performed CUT&RUN to map enrichment of H3K27me2/3 in

strains lacking the individual HCHC components HP1, CDP2, or CHAP (**Figure 4A**). HP1-deficient

(Δ*hpo)* and CDP-2-deficient (Δ*cdp-2)* strains exhibited widespread re-localization of H3K27me2/3

from typical PRC2-target domains to cHet domains, while the CHAP-deficient (Δ*chap*) strain had a

wild type H3K27me2/3 pattern. Interestingly, the global relocalization pattern of H3K27me2/3 was

distinct in the Δ*hpo,* Δ*hda-1,* and Δ*cdp-2* strains. This was evident when H3K27me2/3 enrichment was

plotted across all seven *N. crassa* centromeres (**Figure 4A-B**). We also plotted H3K27me2/3

enrichment across all wild type cHet domains in each of the HCHC-deficient mutants, which

confirmed that the pattern of aberrant H3K27me2/3 enrichment was distinct in each of these strains

(**Figure 4C**). Replicate experiments produced similar results (**Figure S4**).

Δhda-1 strains have increased intra- and inter-heterochromatin interactions in the nucleus

To assess if altered enrichment of histone PTMs in Δ*hda-1* influences the 3D organization of

the *Neurospora* genome, we performed cHet-specific chromosome conformation capture coupled with

high-throughput sequencing (*in situ* Hi-C); here, the use of the restriction enzyme *Mse*I (recognition

sequence: T^TAA) effectively captures contacts of the AT-rich cHet regions (67, 85). The *Neurospora*

genome forms a Rabl chromosome conformation characterized by a centromere cluster and multiple

telomere bundles (85–87). The other ~200 cHet regions strongly associate over Megabases (Mb) of

genomic distance. Previously, it was shown that intra- and inter-heterochromatic region interactions

are strongly increased in a Δ*cdp-2* strain due to histone hyperacetylation (56). To determine if similar

changes are observed in another HCHC-deficient mutant, we performed *in situ* Hi-C using *Mse*I in two

independent strain isolates of Δ*hda-1* (NKL9, NKL10). We also performed ChIP-seq for H3K9me3

and H3K27me2/3 in both strains.

As previously shown, *Mse*I Hi-C of a wild type strain produced 20 kilobase (kb) resolution

contact probability heatmaps in which the interspersed cHet regions show strong intra-region (on-

diagonal) interactions (**Figure 5A**) (67). Moreover, long-range interactions between cHet regions are

readily observed, as shown for two individual chromosomes (**Figure 5A,E**; Linkage Groups [LGs] II

and V). Similar contact probability heatmaps are observed for the NKL9 Δ*hda-1* strain, except that

strong inter-region interactions between cHet domains are more apparent (**Figure 5B,F**), especially for

the centromeres of each LG, which are more apt to interact with distal cHet regions. Comparison of the

wild type contact probability heatmap to that of the Δ*hda-1* strains showed increased interactions at the

center of each cHet region, as well as between cHet regions, with the contacts between centromeres

and interspersed heterochromatic regions having the most prominent gains (**Figure 5C,G**). Similar

results were previously reported for a Δ*cdp-2* strain (56). These increased cHet domain contacts of the

LG II and LGV centromeres in the Δ*hda-1* strain NKL9, relative to a wild type strain, are readily

apparent even at a higher resolution (10 kb bins; **Figure 5D,H**). A replicate Hi-C experiment with a

second Δ*hda-1* strain isolate, NKL10, showed similar cHet contact probability gains (**Figure S5**).

These data suggest that loss of HCHC leads to promiscuous interactions between distant cHet regions,

presumably due to hyperacetylation at these regions.

Epigenomic instability in Δhda-1 leads to progressive decay of histone PTM patterns

The two Δ*hda-1* strain isolates examined by Hi-C produced distinct patterns of aberrant

H3K27me3. In the Δ*hda-1* NKL9 isolate, we did not observe aberrant H3K27me2/3 enrichment at the

centromere of LGV (**Figure 5H, S6**). In contrast, the Δ*hda-1* NKL10 isolate displayed striking

enrichment of H3K27me2/3 at this centromere (**Figure S5H, S6**). These data suggest that

redistribution of H3K27me3 may occur in a stochastic or progressive manner, giving rise to unique

patterns of aberrant H3K27me3 enrichment in isogenic Δ*hda-1* strains. This could also explain the

distinct patterns of H3K27me3 enrichment observed in different HCHC-deficient mutants (**Figure 4**).

If epigenomic defects observed in Δ*hda-1* occurred slowly over many rounds of mitosis, we

reasoned that a newly constructed Δ*hda-1* strain might have a moderated phenotype. To test this, we

generated a new Δ*hda-1* strain by split-marker transformation of a wild-type recipient strain. Primary

transformants were selected, cultured, and screened by PCR to identify a homokaryotic Δ*hda-1*

knockout. This homokaryotic transformant (Δ*hda-1* T0) was then passaged by streaking cells on a

plate, isolating a single colony, and then expanding that colony on a slant for a total of three passages

(**Figure 6A**). We saved each slant and refer to the sequentially passaged strains as Δ*hda-1* P1, P2, and

P3. Cells from each slant were then grown for an additional ~18 hours in liquid medium, and ChIP-seq

was performed using antibodies to H3K27me3, H3K36me3, H3K9me3, and Kac. We estimate that

each passage corresponds to ~25 - 40 rounds of mitosis, based on the number of conidiospores in a

typical slant (~10^8/slant).

H3K27me3 enrichment patterns for the primary transformant and the passaged strains are

shown in **Figure 6**. The phenotype of the Δ*hda-1* T0 strain closely resembled the wild-type strain with

respect to H3K27me3 enrichment patterns (**Figure 6A**). With each passage, we observed progressive

loss of H3K27me3 from native fHet domains accompanied by gains of H3K27me3 within typical cHet

domains (**Figure 6B-C**). However, even after three passages, some cHet domains that were enriched

for H3K27me3 in our original Δ*hda-1* lab strain did not gain H3K27me3 in the new Δ*hda-1* P1 – P3

isolates (**Figure 6B,C;** see CENVII)

We next compared H3K27me3, H3K36me3, H3K9me3, and Kac enrichment patterns in our

passaged Δ*hda-1* isolates. Enrichment levels of each modification were quantified in 300 basepair

windows covering typical fHet and cHet loci. Enrichment values were then plotted in a heatmap for

wild type, the Δ*hda-1* P1, P2, P3 isolates, and our original Δ*hda-1* lab strain (S297; obtained from the

Fungal Genetics Stock Center) (**Figure 6D**). After the first passage (P1), both H3K27me3 and

H3K36me3 were retained at native fHet domains, but modest gains of H3K27me3 and H3K36me3

were detected in cHet regions. In addition, the Δ*hda-1* P1 isolate displayed increased Kac and

relatively normal H3K9me3 enrichment in cHet domains. With each sequential passage, the number of

fHet windows with significant enrichment of H3K27me3 and H3K36me3 decreased, accompanied by

an increase in the number of cHet windows enriched for both modifications. Progressive gains of

H3K27me3/H3K36me3 were correlated with loss of H3K9me3, suggesting that H3K27me3 ultimately

leads to displacement of H3K9me3 or that progressive loss of H3K9me3 allows H3K27me3

deposition. As previously observed (**Figure 3**), cHet regions enriched for H3K27me3 in Δ*hda-1* strains

showed reduced levels of Kac compared to cHet regions that did not gain H3K27me3, implying that

H3K27me2/3 or H3K36me3 directs deacetylation at these regions. We conclude that loss of HDA-1

leads to progressive epigenomic dysfunction.

Discussion

In this study, we found that HDA-1, a cHet-specific histone deacetylase, is required for repression of PRC2-methylated genes and for maintaining global epigenome stability in *N. crassa*. In HDA-1-deficient cells, cHet domains displayed hyperacetylation, reduced H3K9me3 and abnormal enrichment of two modifications typically enriched in fHet regions, H3K27me2/3 and H3K36me3. Aberrant gains of H3K27me2/3 and H3K36me3 were accompanied by loss of H3K27me2/3 from typical fHet regions and activation of genes normally repressed by PRC2 (**Figure 6E**). These findings extend previous reports of altered H3K27me2/3 in mutants lacking H3K9me3 or HP1, two components required for targeting HDA-1 to cHet domains (35, 36). Notably, our findings differ from those of Jamieson and colleagues, who reported that HCHC components HDA-1 and CDP-2 were not required for normal H3K27me3 patterns (36). This discrepancy likely reflects a difference in the "ages" of Δ*hda-1* and Δ*cdp-2* strains investigated in each study. Here, we showed that redistribution of H3K27me2/3 in a newly constructed Δ*hda-1* mutant occurred gradually, taking place over hundreds of nuclear divisions. It is likely that Jamieson and colleagues performed experiments with Δ*hda-1* and Δ*cdp-2* isolates that were recently backcrossed. If so, relatively normal H3K27me2/3 patterns are expected due the delayed onset of epigenomic defects in the absence of HDA-1 activity. We found that CHAP, an additional HCHC component, was not required for proper H3K27me3. We do not know how many times the Δ*chap* mutant strain obtained from the Neurospora knockout project has been passaged. It is therefore possible that continued propagation of this strain will reveal gradual epigenome instability as observed in Δ*hda-1* and Δ*cdp-2* strains. However, this is unlikely given that previous studies of Δ*chap* strains reported only modest effects on Kac and 3D genome structure compared to other HCHC components, consistent with a non-essential role for CHAP in targeting HDA-1 (42, 56). More generally, our data highlight the importance of controlling for strain age, especially for the study of self-reinforcing repressive chromatin states.

Our results suggest that the aberrant H3K36me3 or H3K27me3 observed at cHet regions of

HDA-1-deficient strains directs a compensatory deacetylase activity. In Δ*hda-1* strains, acetylation

levels were high in cHet regions that retained normal H3K9me3 enrichment and low in cHet regions

enriched for H3K36me3/H3K27me3. This compensatory HDAC activity is likely provided by HDA-3.

The *hda-3* gene is essential and therefore was not tested in our screen; however, the HDA-3-containing

RPD3L complex was recently identified as a key repressor of PRC2 target genes (80). Thus, we

propose that aberrant H3K27me2/3 and/or H3K36me3 targets RPD3L deacetylase activity to a subset

of cHet regions in HDA-1-deficient cells. Perhaps depletion of RPD3L from normally repressed fHet

is responsible for increased transcription of fHet genes in the Δ*hda-1* mutant. Future experiments are

needed to establish how these fHet modifications target the RPDL3 complex.

Our data indicate that HDA-1 normally suppresses off-target activity of PRC2 and ASH-1 at

cHet loci. It is possible that HDA-1 functions to restrict DNA accessibility in cHet domains and that

loss of HDA-1 leads to transient exposure of cryptic PRC2 or ASH1 recruitment sites. Prior work in *N.*

crassa showed that HDA-1 limits DNA methyltransferase accessibility in cHet regions (42). In *S.*

pombe, histone deacetylation by the HDA-1 homolog Clr3 is critical for inhibiting chromatin

remodeling activity and suppressing nucleosome turnover (88, 89). How PCR2 is recruited to target

loci is not well understood in *N. crassa,* but H3K27me2/3 can be directed by telomere repeat

sequences in the fungus (90). Future work is needed to determine if the fHet machinery is recruited to

cHet regions by DNA sequences or structures that are transiently unmasked in the absence of HDA-1.

Alternatively, HDA-1-deficient cells may express cHet-dervied RNAs that recruit PRC2 or ASH1. The

ability of PRC2 to bind RNA is conserved from fungi to humans (91), but RNA-dependent recruitment

of PRC2 remains controversial (92, 93). A third possibility is that HDA-1 loss may produce changes in

the localization or biophysical properties of cHet-associated molecular condensates, leading to off-

target activity of PRC2 or ASH1. Phase-separated condensates modulate the activity of PRC2 in

Cryptococcus neoformans. In *Ccc1Δ* mutants, H3K27me3 accumulates at centromeric cHet domains,

where HP1 is proposed to form condensates that concentrate PRC2 to drive off-target methylation in

this mutant background (94, 95). In contrast to the situation in *C. neoformans, N. crassa* HP1 acts to

prevent aberrant H3K27me3 deposition (35, 36). In *Δhda-1,* cHet regions that gain H3K27me3 are

depleted for H3K9me3 and presumably for HP1. Thus, changes to, or loss of, HP1 condensates may be

a precursor to aberrant H3K27me3/H3K36me3 deposition at cHet domains. Additional work is needed

to determine how H3K27me3/H3K36me3 are excluded from cHet in wild-type cells and how these

PTMs are aberrantly targeted to cHet regions in *Δhda-1* strains.

Our findings add to the growing body of evidence that HDACs are critical for maintaining

epigenome stability. HDACs are required for epigenetic inheritance of H3K9me3 in *S. pombe* (96). It

is possible that cHet-specific HDACs maintain proper H3K27me3 patterns in higher eukaryotes, as

observed here for *N. crassa*. In mammals, patterns of H3K27me3 are influenced by key features of

cHet in the early embryo (97) and by 5mC in a variety of developmental contexts (98). In mESCs,

5mC depletion drives aberrant binding of ncPRC1.1 and deposition of H3K27me3 at pericentromeric

heterochromatin and other CpG-rich regions (53, 54). Although DNA methylation is not required for

normal H3K27me3 patterns in *N. crassa* (35, 36), antagonism between cHet and H3K27me3 is a

shared feature of fungal and mammalian cells. In mESCs, acute 5mC inhibition leads to gradual gains

in H3K27me3 at transposable elements and pericentromeric heterochromatin over a two-week period

(51). The progressive epigenomic decay observed in 5mC-depleted mESC cells is similar to that

observed for HDA-1-deficient cells of *N. crassa*. We speculate that hypoacetylated histones are

important for maintaining proper H3K27me3 patterns in both systems. In mammals, it is known that

methyl binding domain proteins (MBDs) direct HDAC activity to methylated CpGs (99–101). As such,

5mC depletion in mESCs may lead to hyperacetylation at CpG-dense cHet regions, which could

contribute to H3K27me3 redistribution. More generally, data from diverse systems point to HDACs as

critical regulators of epigenome stability. Elucidating HDAC-dependent mechanisms that maintain the

epigenome may ultimately provide new insights into the origins of epigenetic disorders, including

cancer, and could lead to improvements in the detection and treatment of diseases associated with PcG

dysfunction.

Data Availability

All sequencing data are available through the NCBI Gene Expression at the National Center for

Biotechnology Information (NCBI) under accession numbers GSE286914 and GSE286535 (ChIP-seq),

GSE286916 (CUT&RUN), GSE286538 (Hi-C). Accession numbers for RNA-seq data are included in

Supplementary Dataset S2.

Funding

This work was supported by grants from National Institutes of Health to ZAL

(1R01GM132644, 1R35GM152134) or ADK (1R15GM140396-01). The work (proposal:

10.46936/10.25585/60001136) conducted by the U.S. Department of Energy Joint Genome Institute

(https://ror.org/04xm1d337), a DOE Office of Science User Facility, is supported by the Office of

Science of the U.S. Department of Energy operated under Contract No. DE-AC02-05CH11231.

Acknowledgements

The authors would like to Dr. Mary Goll (University of Georgia) and members of the Lewis

and Klocko labs for constructive comments on the manuscript. We also thank colleagues at the

University of Georgia and the UCCS Department of Chemistry & Biochemistry for helpful

discussions.

References

1.   S. I. S. Grewal, The molecular basis of heterochromatin assembly and epigenetic inheritance.
Mol. Cell **83**, 1767–1785 (2023).

2.   B. Schuettengruber, H.-M. Bourbon, L. Di Croce, G. Cavalli, Genome Regulation by Polycomb
and Trithorax: 70 Years and Counting. *Cell* **171**, 34–57 (2017).

3.   N. P. Blackledge, R. J. Klose, The molecular principles of gene regulation by Polycomb
repressive complexes. *Nat. Rev. Mol. Cell Biol.* **22**, 815–833 (2021).

4.   S. Tamburri, S. Rustichelli, S. Amato, D. Pasini, Navigating the complexity of Polycomb
repression: Enzymatic cores and regulatory modules. *Mol. Cell* **84**, 3381–3405 (2024).

5.   A. A. Erlendson, S. Friedman, M. Freitag, A matter of scale and dimensions: Chromatin of
chromosome landmarks in the fungi. *Microbiol. Spectr.* **5**, 10.1128/microbiolspec.FUNK-0054–
2017 (2017).

6.   R. Margueron, D. Reinberg, The Polycomb complex PRC2 and its mark in life. *Nature* **469**,
343–349 (2011).

7.   P. Trojer, D. Reinberg, Facultative Heterochromatin: Is There a Distinctive Molecular
Signature? *Mol. Cell* **28**, 1–13 (2007).

8.   M. Matsuwaka, M. Kumon, A. Inoue, H3K27 dimethylation dynamics reveal stepwise
establishment of facultative heterochromatin in early mouse embryos. *Nat. Cell Biol.* 1–11
(2024). https://doi.org/10.1038/s41556-024-01553-1.

9.   A. Janssen, S. U. Colmenares, G. H. Karpen, Heterochromatin: Guardian of the Genome. *Annu.*
Rev. Cell Dev. Biol. **34**, 265–288 (2018).

10.   A. Piunti, A. Shilatifard, The roles of Polycomb repressive complexes in mammalian
development and cancer. *Nat. Rev. Mol. Cell Biol.* **22**, 326–345 (2021).

11.   J. W. Højfeldt, *et al.*, Non-core Subunits of the PRC2 Complex Are Collectively Required for Its
Target-Site Specificity. *Mol. Cell* **76**, 423-436.e3 (2019).

12.   J. I. Barrasa, T. G. Kahn, M. J. Lundkvist, Y. B. Schwartz, DNA elements tether canonical
Polycomb Repressive Complex 1 to human genes. *Nucleic Acids Res.* **51**, 11613–11633 (2023).

13.   E. Glancy, *et al.*, PRC2.1- and PRC2.2-specific accessory proteins drive recruitment of different
forms of canonical PRC1. *Mol. Cell* **83**, 1393-1411.e7 (2023).

14.   H. Bharti, S. Han, H.-W. Chang, D. Reinberg, Polycomb repressive complex 2 accessory
factors: rheostats for cell fate decision? *Curr. Opin. Genet. Dev.* **84**, 102137 (2024).

15.   J. Xiao, *et al.*, Cis and trans determinants of epigenetic silencing by Polycomb repressive
complex 2 in Arabidopsis. *Nat. Genet.* **49**, 1546–1552 (2017).

16.   R. Kalb, *et al.*, Histone H2A monoubiquitination promotes histone H3 methylation in Polycomb
472   repression. *Nat. Struct. Mol. Biol.* **21**, 569–571 (2014).

473   17.   H. Barbour, S. Daou, M. Hendzel, E. B. Affar, Polycomb group-mediated histone H2A
474   monoubiquitination in epigenome regulation and nuclear processes. *Nat. Commun.* **11**, 5947
475   (2020).

476   18.   H. Ohtomo, *et al.*, H2A Ubiquitination Alters H3-tail Dynamics on Linker-DNA to Enhance
477   H3K27 Methylation. *J. Mol. Biol.* **435**, 167936 (2023).

478   19.   F. W. Schmitges, *et al.*, Histone Methylation by PRC2 Is Inhibited by Active Chromatin Marks.
479   *Mol. Cell* **42**, 330–341 (2011).

480   20.   W. Yuan, *et al.*, H3K36 Methylation Antagonizes PRC2-mediated H3K27 Methylation*. *J. Biol.
481   Chem.* **286**, 7983–7989 (2011).

482   21.   T. Cookis, A. Lydecker, P. Sauer, V. Kasinath, E. Nogales, Structural basis for the inhibition of
483   PRC2 by active transcription histone posttranslational modifications. *bioRxiv*
484   2024.02.09.579730 (2024). https://doi.org/10.1101/2024.02.09.579730.

485   22.   M. M. Balas, *et al.*, Establishing RNA-RNA interactions remodels lncRNA structure and
486   promotes PRC2 activity. *Sci. Adv.* **7**, eabc9191 (2021).

487   23.   M. Uckelmann, C. Davidovich, Not just a writer: PRC2 as a chromatin reader. *Biochem. Soc.
488   Trans.* **49**, 1159–1170 (2021).

489   24.   A. K. Braceros, *et al.*, Proximity-dependent recruitment of Polycomb repressive complexes by
490   the lncRNA *Airn*. *Cell Rep.* **42**, 112803 (2023).

491   25.   J. Song, *et al.*, Structural basis for inactivation of PRC2 by G-quadruplex RNA. *Science* **381**,
492   1331–1337 (2023).

493   26.   W. Yuan, *et al.*, Dense chromatin activates Polycomb repressive complex 2 to regulate H3
494   lysine 27 methylation. *Science* **337**, 971–975 (2012).

495   27.   R. Aramayo, E. U. Selker, Neurospora crassa, a model system for epigenetics research. *Cold
496   Spring Harb. Perspect. Biol.* **5**, a017921 (2013).

497   28.   K. Jamieson, M. R. Rountree, Z. A. Lewis, J. E. Stajich, E. U. Selker, Regional control of
498   histone H3 lysine 27 methylation in Neurospora. *Proc. Natl. Acad. Sci.* **110**, 6027–6032 (2013).

499   29.   A. J. Courtney, *et al.*, Normal Patterns of Histone H3K27 Methylation Require the Histone
500   Variant H2A.Z in Neurospora crassa. *Genetics* **216**, 51–66 (2020).

501   30.   K. J. McNaught, E. T. Wiles, E. U. Selker, Identification of a PRC2 Accessory Subunit
502   Required for Subtelomeric H3K27 Methylation in Neurospora crassa. *Mol. Cell. Biol.* **40**,
503   e00003-20 (2020).

31. E. T. Wiles, *et al.*, Evolutionarily ancient BAH–PHD protein mediates Polycomb silencing.
Proc. Natl. Acad. Sci. **117**, 11614–11623 (2020).

32. M. Kamei, *et al.*, IMITATION SWITCH is required for normal chromatin structure and gene
repression in PRC2 target domains. *Proc. Natl. Acad. Sci.* **118**, e2010003118 (2021).

33. E. T. Wiles, C. C. Mumford, K. J. McNaught, H. Tanizawa, E. U. Selker, The ACF chromatin-
remodeling complex is essential for Polycomb repression. *eLife* **11**, e77595 (2022).

34. V. T. Bicocca, T. Ormsby, K. K. Adhvaryu, S. Honda, E. U. Selker, ASH1-catalyzed H3K36
methylation drives gene repression and marks H3K27me2/3-competent chromatin. *eLife* **7**,
e41497 (2018).

35. E. Y. Basenko, *et al.*, Genome-wide redistribution of H3K27me3 is linked to genotoxic stress
and defective growth. *Proc. Natl. Acad. Sci.* **112**, E6339–E6348 (2015).

36. K. Jamieson, *et al.*, Loss of HP1 causes depletion of H3K27me3 from facultative
heterochromatin and gain of H3K27me2 at constitutive heterochromatin. *Genome Res.* **26**, 97–
107 (2016).

37. L. R. Connolly, K. M. Smith, M. Freitag, The Fusarium graminearum Histone H3 K27
Methyltransferase KMT6 Regulates Development and Expression of Secondary Metabolite
Gene Clusters. *PLOS Genet.* **9**, e1003916 (2013).

38. M. Möller, J. B. Ridenour, D. F. Wright, F. A. Martin, M. Freitag, H4K20me3 is important for
Ash1-mediated H3K36me3 and transcriptional silencing in facultative heterochromatin in a
fungal pathogen. *PLOS Genet.* **19**, e1010945 (2023).

39. M. Xu, *et al.*, A repressive H3K36me2 reader mediates Polycomb silencing. *Nat. Commun.* **15**,
7287 (2024).

40. E. T. Wiles, E. U. Selker, H3K27 methylation: a promiscuous repressive chromatin mark. *Curr.*
Opin. Genet. Dev. **43**, 31–37 (2017).

41. Z. A. Lewis, *et al.*, Relics of repeat-induced point mutation direct heterochromatin formation in
Neurospora crassa. *Genome Res.* **19**, 427–437 (2009).

42. S. Honda, *et al.*, Heterochromatin protein 1 forms distinct complexes to direct histone
deacetylation and DNA methylation. *Nat. Struct. Mol. Biol.* **19**, 471–477, S1 (2012).

43. H. Tamaru, *et al.*, Trimethylated lysine 9 of histone H3 is a mark for DNA methylation in
Neurospora crassa. *Nat. Genet.* **34**, 75–79 (2003).

44. M. Freitag, P. C. Hickey, T. K. Khlafallah, N. D. Read, E. U. Selker, HP1 Is Essential for DNA
Methylation in *Neurospora*. *Mol. Cell* **13**, 427–434 (2004).

45. S. Honda, E. U. Selker, Direct Interaction between DNA Methyltransferase DIM-2 and HP1 Is
Required for DNA Methylation in Neurospora crassa. *Mol. Cell. Biol.* **28**, 6044–6055 (2008).

46.   Z. A. Lewis, *et al.*, DNA Methylation and Normal Chromosome Behavior in Neurospora
Depend on Five Components of a Histone Methyltransferase Complex, DCDC. *PLOS Genet.* **6**,
e1001196 (2010).

47.   S. Honda, *et al.*, The DMM complex prevents spreading of DNA methylation from transposons
to nearby genes in Neurospora crassa. *Genes Dev.* **24**, 443–454 (2010).

48.   S. Honda, *et al.*, Dual chromatin recognition by the histone deacetylase complex HCHC is
required for proper DNA methylation in Neurospora crassa. *Proc. Natl. Acad. Sci. U. S. A.* **113**,
E6135–E6144 (2016).

49.   J. P. Reddington, *et al.*, Redistribution of H3K27me3 upon DNA hypomethylation results in de-
repression of Polycomb target genes. *Genome Biol.* **14**, R25 (2013).

50.   N. Saksouk, *et al.*, Redundant Mechanisms to Form Silent Chromatin at Pericentromeric
Regions Rely on BEND3 and DNA Methylation. *Mol. Cell* **56**, 580–594 (2014).

51.   M. Walter, A. Teissandier, R. Pérez-Palacios, D. Bourc'his, An epigenetic switch ensures
transposon repression upon dynamic loss of DNA methylation in embryonic stem cells. *eLife* **5**,
e11418 (2016).

52.   H. Wu, *et al.*, Dnmt3a-Dependent Nonpromoter DNA Methylation Facilitates Transcription of
Neurogenic Genes. *Science* **329**, 444–448 (2010).

53.   N. P. Blackledge, *et al.*, Variant PRC1 Complex-Dependent H2A Ubiquitylation Drives PRC2
Recruitment and Polycomb Domain Formation. *Cell* **157**, 1445–1459 (2014).

54.   S. Cooper, *et al.*, Targeting polycomb to pericentric heterochromatin in embryonic stem cells
reveals a role for H2AK119u1 in PRC2 recruitment. *Cell Rep.* **7**, 1456–1470 (2014).

55.   J. D. Gessaman, E. U. Selker, Induction of H3K9me3 and DNA methylation by tethered
heterochromatin factors in Neurospora crassa. *Proc. Natl. Acad. Sci.* **114**, E9598–E9607 (2017).

56.   A. W. Scadden, *et al.*, Histone deacetylation and cytosine methylation compartmentalize
heterochromatic regions in the genome organization of Neurospora crassa. *Proc. Natl. Acad. Sci.*
120, e2311249120 (2023).

57.   R. H. Davis, F. J. de Serres, "[4] Genetic and microbiological research techniques for
Neurospora crassa" in *Methods in Enzymology*, Metabolism of Amino Acids and Amines Part
A., (Academic Press, 1970), pp. 79–143.

58.   H. V. Colot, *et al.*, A high-throughput gene knockout procedure for Neurospora reveals
functions for multiple transcription factors. *Proc. Natl. Acad. Sci.* **103**, 10352–10357 (2006).

59.   R. S. Goswami, "Targeted Gene Replacement in Fungi Using a Split-Marker Approach" in
Plant Fungal Pathogens: Methods and Protocols, M. D. Bolton, B. P. H. J. Thomma, Eds.
(Humana Press, 2012), pp. 255–269.

60. B. Margolin, M. Freitag, E. Selker, Improved plasmids for gene targeting at the his-3 locus of
Neurospora crassa by electroporation. *Fungal Genet. Rep.* **44**, 34–36 (1997).

61. M. Seymour, *et al.*, Histone H1 Limits DNA Methylation in Neurospora crassa. *G3
GenesGenomesGenetics* **6**, 1879–1889 (2016).

62. A. R. Ferraro, Z. A. Lewis, "ChIP-Seq Analysis in Neurospora crassa" in *Fungal Genomics:
Methods and Protocols*, Methods in Molecular Biology., R. P. de Vries, A. Tsang, I. V.
Grigoriev, Eds. (Springer, 2018), pp. 241–250.

63. Z. Zhou, Y. Dang, M. Zhou, H. Yuan, Y. Liu, Codon usage biases co-evolve with transcription
termination machinery to suppress premature cleavage and polyadenylation. *eLife* **7**, e33569
(2018).

64. P. J. Skene, S. Henikoff, An efficient targeted nuclease strategy for high-resolution mapping of
DNA binding sites. *eLife* **6**, e21856 (2017).

65. M. P. Meers, T. D. Bryson, J. G. Henikoff, S. Henikoff, Improved CUT&RUN chromatin
profiling tools. *eLife* **8**, e46314 (2019).

66. F. Krueger, TrimGalore: A wrapper around Cutadapt and FastQC to consistently apply adapter
and quality trimming to FastQ files, with extra functionality for RRBS data. (2019). Available
at: https://github.com/FelixKrueger/TrimGalore [Accessed 11 November 2024].

67. S. Rodriguez, *et al.*, The genome organization of Neurospora crassa at high resolution uncovers
principles of fungal chromosome topology. *G3 GenesGenomesGenetics* **12**, jkac053 (2022).

68. H. Li, R. Durbin, Fast and accurate short read alignment with Burrows–Wheeler transform.
Bioinformatics **25**, 1754–1760 (2009).

69. B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**, 357–
359 (2012).

70. P. Danecek, *et al.*, Twelve years of SAMtools and BCFtools. *GigaScience* **10**, giab008 (2021).

71. F. Ramírez, *et al.*, deepTools2: a next generation web server for deep-sequencing data analysis.
Nucleic Acids Res. **44**, W160–W165 (2016).

72. J. T. Robinson, *et al.*, Integrative genomics viewer. *Nat. Biotechnol.* **29**, 24–26 (2011).

73. A. T. L. Lun, G. K. Smyth, csaw: a Bioconductor package for differential binding analysis of
ChIP-seq data using sliding windows. *Nucleic Acids Res.* **44**, e45–e45 (2016).

74. A. D. Klocko, *et al.*, Selection and Characterization of Mutants Defective in DNA Methylation
in Neurospora crassa. *Genetics* **216**, 671–688 (2020).

75. A. Dobin, *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

76. Y. Liao, G. K. Smyth, W. Shi, featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

77. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

78. F. Ramírez, *et al.*, High-resolution TADs reveal DNA sequences underlying genome
organization in flies. *Nat. Commun.* **9**, 189 (2018).

79. K. A. Borkovich, *et al.*, Lessons from the Genome Sequence of Neurospora crassa: Tracing the
Path from Genomic Blueprint to Multicellular Organism. *Microbiol. Mol. Biol. Rev.* **68**, 1–108
(2004).

80. C. C. Mumford, *et al.*, The RPD3L deacetylation complex is required for facultative
heterochromatin repression in Neurospora crassa. *Proc. Natl. Acad. Sci.* **121**, e2404770121
(2024).

81. I. Bauer, *et al.*, A Class 1 Histone Deacetylase with Potential as an Antifungal Target. *mBio* **7**,
10.1128/mbio.00831-16 (2016).

82. K. M. Smith, *et al.*, H2B- and H3-Specific Histone Deacetylases Are Required for DNA
Methylation in Neurospora crassa. *Genetics* **186**, 1207–1216 (2010).

83. K. K. Adhvaryu, S. A. Morris, B. D. Strahl, E. U. Selker, Methylation of Histone H3 Lysine 36
Is Required for Normal Development in Neurospora crassa. *Eukaryot. Cell* **4**, 1455–1464
(2005).

84. A. D. Klocko, *et al.*, Normal chromosome conformation depends on subtelomeric facultative
heterochromatin in Neurospora crassa. *Proc. Natl. Acad. Sci.* **113**, 15048–15053 (2016).

85. S. S. P. Rao, *et al.*, A 3D map of the human genome at kilobase resolution reveals principles of
chromatin looping. *Cell* **159**, 1665–1680 (2014).

86. E. Lieberman-Aiden, *et al.*, Comprehensive mapping of long range interactions reveals folding
principles of the human genome. *Science* **326**, 289–293 (2009).

87. D. E. Torres, A. T. Reckard, A. D. Klocko, M. F. Seidl, Nuclear genome organization in fungi:
from gene folding to Rabl chromosomes. *FEMS Microbiol. Rev.* **47**, fuad021 (2023).

88. J. F. Garcia, P. A. Dumesic, P. D. Hartley, H. El-Samad, H. D. Madhani, Combinatorial, site-
specific requirement for heterochromatic silencing factors in the elimination of nucleosome-free
regions. *Genes Dev.* **24**, 1758–1771 (2010).

89. R. K. Sahu, *et al.*, Nucleosome remodeler exclusion by histone deacetylation enforces
heterochromatic silencing and epigenetic inheritance. *Mol. Cell* **84**, 3175-3191.e8 (2024).

90. K. Jamieson, *et al.*, Telomere repeats induce domains of H3K27 methylation in Neurospora.
eLife **7**, e31216 (2018).

91. Y. Long, *et al.*, Conserved RNA-binding specificity of polycomb repressive complex 2 is
achieved by dispersed amino acid patches in EZH2. *eLife* **6**, e31558 (2017).

92. J. K. Guo, *et al.*, Denaturing purifications demonstrate that PRC2 and other widely reported
chromatin proteins do not appear to bind directly to RNA in vivo. *Mol. Cell* **84**, 1271-1289.e12
(2024).

93. Y. Lee, R. Blum, M. Rosenberg, J. T. Lee, Re-analysis of CLAP data affirms PRC2 as an RNA
binding protein. [Preprint] (2024). Available at:
https://www.biorxiv.org/content/10.1101/2024.09.19.613009v1 [Accessed 24 December 2024].

94. P. A. Dumesic, *et al.*, Product binding enforces the genomic specificity of a yeast polycomb
repressive complex. *Cell* **160**, 204–218 (2015).

95. S. Lee, *et al.*, Intrinsic mesoscale properties of a Polycomb protein underpin heterochromatin
fidelity. *Nat. Struct. Mol. Biol.* **30**, 891–901 (2023).

96. M. Zofall, R. Sandhu, S. Holla, D. Wheeler, S. I. S. Grewal, Histone deacetylation primes self-
propagation of heterochromatin domains to promote epigenetic inheritance. *Nat. Struct. Mol.*
Biol. **29**, 898–909 (2022).

97. M. Tardat, *et al.*, Cbx2 Targets PRC1 to Constitutive Heterochromatin in Mouse Zygotes in a
Parent-of-Origin-Dependent Manner. *Mol. Cell* **58**, 157–171 (2015).

98. J. J. Kim, R. E. Kingston, Context-specific Polycomb mechanisms in development. *Nat. Rev.*
Genet. **23**, 680–695 (2022).

99. Q. Du, P.-L. Luu, C. Stirzaker, S. J. Clark, Methyl-CpG-Binding Domain Proteins: Readers of
the Epigenome. *Epigenomics* **7**, 1051–1073 (2015).

100. X. Nan, *et al.*, Transcriptional repression by the methyl-CpG-binding protein MeCP2 involves a
histone deacetylase complex. *Nature* **393**, 386–389 (1998).

101. H.-H. Ng, *et al.*, MBD2 is a transcriptional repressor belonging to the MeCP1 histone
deacetylase complex. *Nat. Genet.* **23**, 58–61 (1999).

   Figure Legends

Figure 1: HDA-1 is required for the repression of PRC2-methylated genes. (**A**) The boxplots show

the average level of expression for all PRC2-target genes (n= 679) for the indicated strains, with the x-

axis values corresponding to log2-transformed transcripts per million (TPM) read counts. Lines

represents the median expression value, and the notches indicate the 95% confidence interval. (**B**) The

heatmap shows relative expression levels of each PRC2-target gene for the indicated strains (n= 537

genes expressed in at least one strain). (**C**) The volcano plot shows differentially expressed genes for

wild type and Δ*hda-1*. Dark green spots correspond to PRC2-methylated genes that meet fold-change

and statistical thresholds (log2[fold change] +/- 1.5 and an adjusted p-value < 0.05). Black spots

correspond to non-PRC2 methylated genes that meet the same fold-change and statistical threshold.

Genes that do not meet statistical or fold-change thresholds are colored in pale green for PRC2-target

genes or grey for non-PRC2 targets.

Figure 2: H3K27me3 aberrantly localizes to heterochromatin in the Δ*hda-1* mutant. (**A-B**) The

genome browser image shows the distribution of H3K9me3 (blue) and H3K27me3 (green), as assessed

by CUT&RUN, across (**A**) Linkage Group III or (**B**) three representative regions shown at higher

magnification. The scale bars indicate genomic distance in kilobases (kb). (**C-D**) Heatmaps depicting

H3K27me3 (green) or H3K9me3 (blue) enrichment across (**C**) all wild-type fHet domains or (**D**) all

wild-type cHet domains for the indicated strains. The heatmaps are sorted such that the first 14 rows in

each heatmap correspond to the 14 subtelomeric regions located at the chromosome ends. The

remaining rows are sorted by the level H3K27me3 enrichment in Δ*hda-1* in descending order from

highest to lowest. Each peak size is normalized to 100% with 1,000 bp shown up/downstream. (**E**) The

genome browser tracks show the distribution of H3K9me3 (blue) and H3K27me3 (green), as in A, for

the indicated strains. (**F-G**) The heatmaps show enrichment of H3K27me3 (green) or H3K9me3 (blue)

across (**F**) all wild-type fHet domains or (**G**) all wild type cHet domains for the indicated strains,

ordered as in panels C-D.

Figure 3: **Acetylation and H3K36me3 patterns are altered in Δ*hda-1*.** (**A-B**) The genome browser

tracks show ChIP-seq enrichment of (**A**) H3K9me3 (blue), H3K27me3 (green), and pan-acetyl lysine

(Kac; purple) or (**B**) H3K36me3 across Linkage Group VII, as in Figure 2A. (**C**) The heatmaps show

ChIP-seq enrichment of H3K9me3 (blue), H3K27me3 (green), H3K36me3 (orange), or pan-acetyl

lysine (H3Kac; purple) for wild type (left) or Δ*hda-1* (right) across all cHet domains. The heatmaps are

scaled and sorted as in Figure 2C-D, with subtelomeres at the top followed by the descending

H3K27me3 signal in fHet regions in a Δ*hda-1* strain.

Figure 4: HP1 and CDP-2, but not CHAP, are required for normal H3K27me3 distribution. (A-

B) The genome browser tracks show CUT&RUN enrichment of H3K9me3 (blue) and H3K27me2/3

(green) across (**A**) Linkage Group IV, or (**B**) the centromeres of the remaining chromosomes, for the

indicated strains, as in Figure 2A-B. (**C**) The heatmaps show H3K27me2/3 (green) or H3K9me3 (blue)

CUT&RUN enrichment for the indicated strains across all wild type cHet regions, except for 14 sub-

telomeric domains adjacent to the chromosome ends. Each cHet region is normalized to 100% size,

and no flanking sequences are plotted.

Figure 5: Inter-heterochromatic region contact gains and variable H3K27me2/3 enrichment

occur in a Δ*hda-1* strain. Heterochromatin-specific (*Mse*I) *in situ* Hi-C and H3K27me2/3 or

H3K9me3 ChIP-seq of wild type or Δ*hda-1* (NKL9) strains are displayed across two Linkage Groups

(LG), LG II (**A-D**) or LG V (**E-H**). Contact probability heatmaps of raw *in situ* Hi-C reads at 20 kb

resolution of a wild-type strain (**A, E**) or the NKL9 Δ*hda-1* strain (**B, F**) are shown; the heatmap

scalebar is below, while the genomic distance is indicated on the x-axis and the number of bins

indicated on the right. **(C, G**) Heatmaps showing the change in contact probability ($\log_2$) between wild

type and $\Delta hda-1$ strains, with the heatmap scalebar on the right. The purple line shows the centromeric

region highlighted in panels D and H. **(D, H**) Heatmaps of the contact probability of wild type (top) or

$\Delta hda-1$ (NKL9) strains (middle), or the change in contact probability between wild type and $\Delta hda-1$

strains (bottom), between the centromeres and right chromosomal arm, displayed at 10 kb resolution.

IGV images of H3K27me2/3 (green) or H3K9me3 (blue) ChIP-seq tracks of wild type and $\Delta hda-1$

(NKL9) strains are shown below. Genomic distances of LG II (panel **D**) and LG V (panel **H**) are

shown above.

Figure 6. H3K27me3 relocalization is progressive in a laboratory evolution experiment. (A) The

schematic diagram illustrates the procedure for generating and passaging a new $\Delta hda-1$ strain (left).

The genome browser images (right) show ChIP-seq enrichment of H3K9me3 (blue) or H3K27me3

(green) ChIP-seq enrichment across Linkage Group III for the indicated strains, as in Figure 2A-B. **(B-**

C) The genome browser images show H3K27me3 (green) ChIP-seq enrichment across (**B**) Linkage

Groups III, VI, and VII for the indicated strains, or (**C**) enhanced images of a cHet region that exhibits

progressive gains of H3K27me3 with each passage (left), an fHet region that exhibits progressive loss

of H3K27me3 (middle), and the centromere of LGVII (right), which is enriched for H3K27me3 in our

original $\Delta hda-1$ strain but not in the newly constructed and passaged $\Delta hda-1$ strain. (**D**) The heatmap

shows relative enrichment (log2[IP/input]) of H3K27me3 (green), H3K9me3 (blue), or H3K36me3

(orange) in 300bp windows spanning all wildtype fHet and cHet domains for the indicated strains.

Relative enrichment of Kac (purple; log2[$\Delta hda-1$ IP / WT IP]) is shown for the same regions.

Enrichment values that were negative or did not meet a statistical threshold (FDR < 0.1) are shaded in

white. (**E**) Model of the relationship between fHet (left) and cHet (right) repression in wild type (top)

or Δ*hda-1* (bottom) strains.

A LGVII H3K9me3 H3K27me2/3 H3K36me3 Kac

500 kb

WT

Δhda-1

genes

B WT

Δhda-1

genes

C WT Δhda-1

H3K9me3 Kac H3K27me3 H3K36me3 H3K9me3 Kac H3K27me3 H3K36me3

cHet domains

Figure 4

Figure 5

Figure 6