The Genome of *Chenopodium ficifolium*: Developing Genetic Resources and a Diploid Model System for Allotetraploid Quinoa

Clayton D. Ludwig[1], Peter J. Maughan[2], Eric N. Jellen[2], Thomas M. Davis[1]

Affiliations:

[1]Department of Agriculture, Nutrition, and Food Systems, University of New Hampshire, Durham, NH 03824, USA

[2]Brigham Young University, Department of Plant and Wildlife Sciences, College of Life Sciences, Provo, UT 84602, USA.

Abbreviations:

AED, annotation edit distance; EST, expressed sequence tag; GO, gene ontology; GWAS, genome wide association study; HCGS, UNH Hubbard Center for Genome Studies; HMW, high molecular weight; MSA, multiple sequence alignment; ONT, Oxford Nanopore Technology; PC, pseudochromosome

Abstract

High-quality nuclear, chloroplast, and preliminary mitochondrial genomes have been assembled and annotated for the B-genome diploid (BB: $2n = 2x = 18$) figleaf goosefoot (*Chenopodium ficifolium*). The primary objective was to advance a simplified model system for genetic characterization and improvement of allotetraploid (AABB: $2n = 4x = 36$) quinoa (*Chenopodium*

quinoa), a nutritionally valuable, halophytic orphan crop. In addition to its diploidy and

favorably small genome size, the *C. ficifolium* model provides a shorter generational period and

smaller overall plant size as compared to *C. quinoa*, while displaying relevant agronomic trait

variations amenable to gene-trait association studies. The *C. ficifolium* 'Portsmouth' nuclear

genome was sequenced using PacBio HiFi Long Read technology and assembled using Hifiasm.

After manual adjustments, the final ChenoFicP_1.0 assembly consisted of nine

pseudochromosomes spanning 711.5 Mbp, while 22,617 genes were identified and annotated.

BUSCO analyses indicated a nuclear genome completeness of 97.5%, and a proteome and

transcriptome completeness of 98.4 percent. The chloroplast genome assembly detected two

equally represented structural haplotypes differing in the orientation of the Short Single Copy

region relative to the Long Single Copy region. Phylogenetic and parentage analyses pointed to

an unspecified AA diploid species and away from *C. ficifolium* as the likely maternal chloroplast

and mitochondrial genome donor(s) during the initial tetraploidization event in the *C. quinoa*

lineage. Using the new ChenoFicP_1.0 reference genome, a GWAS was performed on a

previously studied *C. ficifolium* F2 population to further define region(s) implicated in the

control of three key agronomic traits: days to flowering, plant height, and branch number. This

analysis localized control of all three traits to a 7 Mb interval on pseudochromosome Cf4. This

region contains approximately 770 genes, including the *FTL1* locus, thus confirming and

extending our prior, single-marker analysis showing association of these three traits with an

FTL1 amplicon length polymorphism. The use of these data to further develop *C. ficifolium* as a

model species for genetics and breeding of quinoa serves to expand knowledge and germplasm

resources for quinoa improvement.

Introduction

Chenopodium ficifolium (figleaf goosefoot) is a weedy, Old-World B-genome diploid (BB: $2n = 2x = 18$) relative of the cultivated allotetraploid (AABB: $2n = 4x = 36$) pseudocereal crop quinoa (*Chenopodium quinoa* Willd.) (Walsh *et al.*, 2015; Kolano *et al.*, 2016) and its wild sister AABB tetraploid *Chenopodium berlandieri* (Walsh *et al.*, 2015). The genus *Chenopodium* now resides in the family Amaranthaceae, its former Chenopodiaceae family home having recently been subsumed into the Amaranthaceae (Chase *et al.*, 2016), which also hosts the relevant comparator *Beta vulgaris* (sugar beet). Putatively originating from Eurasia (Walsh *et al.*, 2015), figleaf goosefoot is now widespread in North America, both as a weed of human disturbance and as a naturalized part of disturbed ecosystems. It is one of two recognized potential contributors, or a close relative thereof, of the B subgenome to the *C. quinoa* lineage during its initial tetraploidization event, the other B subgenome candidate being diploid *C. suecicum* (Jellen *et al.*, 2011; Mandák *et al.*, 2012; Walsh *et al.*, 2015; Jarvis *et al.*, 2017). Because figleaf goosefoot potentially contributed the B subgenome to quinoa, or is an extant sister species of the contributor, it is a highly relevant candidate to explore as a diploid model species in relation to quinoa's genetic characterization and breeding, and its study may shed light on specific genes governing domestication-related traits in quinoa (Jellen *et al.*, 2013; Walsh *et al.*, 2015; Kolano *et al.*, 2016; Subedi, Neff and Davis, 2021).

Absent a history of intensive breeding, *C. quinoa* has been categorized as an 'orphan crop' (Lemmon *et al.*, 2018; Kumar and Bhalothia, 2020). However, quinoa is receiving increasing interest from consumers due to its high nutritional value and from agronomists and breeders due to its drought and salt tolerance (Gorinstein *et al.*, 2002; Rojas *et al.*, 2009; Vega-Gálvez *et al.*, 2010; Jacobsen, 2011). These and other attributes make quinoa an attractive candidate for

production in regions that cannot sustain the major grain crops. Furthermore, quinoa has the

potential to serve as a valuable secondary crop, thereby contributing to food security. Thus, it is

an attractive candidate for increased production, justifying investment in development of novel

genetics and methods to streamline the breeding process (Jacobsen, 2003; Bazile, Jacobsen and

Verniau, 2016).

Using *C. ficifolium* as a diploid model species has the potential to accelerate trait dissection and

gene identification due to its comparative genetic simplicity; shorter generation time of ~40 days

compared to quinoa's 90-120 days (Jacobsen, 1997); profuse flower, pollen, and seed production;

and smaller plant size facilitating growth chamber cultivation (Subedi et al., 2021). Building

upon prior, foundational studies (Štorchová *et al.*, 2015, 2019), the targeted development of *C.*

ficifolium as a diploid model species was initiated (Subedi, 2020; Subedi, Neff and Davis, 2021)

with a focus on: i) producing intraspecific hybrids and segregating progenies; ii) investigating

segregation in domestication-related traits of relevance to *C. quinoa* breeding programs; and iii)

detecting gene-trait associations for key genes and traits.

In an F_2 segregating population from an initial cross involving *C. ficifolium* accessions from

Portsmouth ('P') NH USA, and Quebec City ('QC') Quebec Canada, allelic variation in the

Flowering Locus T-Like 1 (*FTL1*) marker locus was found to be associated with variation in days

to flower, plant height, and branch number (Subedi, 2020; Subedi, Neff and Davis, 2021). Using

previously designed primers (Cháb *et al.*, 2008; Štorchová *et al.*, 2015) to target and

conveniently genotype an indel polymorphism within the *FTL1* gene, it was found that F2 plants

homozygous for the 'P' allele flowered on average 12 days earlier than plants homozygous for the

'QC' allele, and two days earlier than heterozygotes (Subedi, 2020; Subedi, Neff and Davis,

2021). As a likely consequence of their decreased maturation time, 'P' allele homozygotes were

also significantly shorter and had fewer branches than plants of the alternate genotypes. These

results strongly implicated the *FTL1* chromosomal region in the control of three agronomically

important traits, but did not rule out the possibility of influence from other genomic regions.

While genotype-phenotype relationships involving *FTL1* must be further studied at both the

diploid and the tetraploid level, this research demonstrates the potential value of *C. ficifolium* as

a model system in which to accelerate discovery and characterization of gene-trait relationships.

The goal of the work reported here is to expand knowledge and establish foundational genomic

resources for the *C. ficifolium* model system.

Materials and Methods

Germplasm

All plant cultivation and phenotyping activities were conducted at the New Hampshire

Agricultural Experiment Station (NHAES) at the University of New Hampshire (UNH). Two

figleaf goosefoot accessions were employed in this study, both of which have been deposited as

seed samples with the USDA North Central Plant Introduction Station (NCRPIS) at Ames Iowa.

The 'Portsmouth' ('P') accession (PI 698433) was collected by Erin Neff and Thomas Davis in

Portsmouth, New Hampshire, USA (Neff, 2017), while the 'Quebec City' ('QC') accession (PI

698434) was collected by Thomas Davis from Quebec City, Quebec, Canada. Crosses between

the 'P' and 'QC' accessions to generate F_1 and F_2 populations, and confirmation of parentage

using informative *FTL1* amplicons, were as previously described (Subedi, Neff and Davis,

2021). For purposes of comparison, we also employed in-house-numbered accessions 302-A of

AA diploid *C. foggii*, and RB6 of AABB allotetraploid *C. berlandieri* var. *macrocalycium*, which

had been collected by Erin Neff and Thomas Davis in southern Maine (Neff, 2017; Neff,

Sullivan and Davis, 2018) and in Rye Beach New Hampshire, respectively; and *C. quinoa*

accession QQ065 (PI 614880), obtained from USDA NPGS.

DNA Isolation and Genome Sequencing

The figleaf goosefoot 'Portsmouth' ('P') accession was used to construct the primary *de novo*

reference genome assembly. Using leaf tissue provided by the UNH investigators, DNA from the

'P' accession was isolated at Brigham Young University (BYU) using methods designed to yield

high molecular weight (HMW) DNA from plant tissues (Vaillancourt and Buell, 2019). The 'P'

accession HMW DNA was sequenced at the BYU DNA Sequencing Center (DNASC) using the

PacBio HiFi platform (Eid *et al.*, 2009). In addition, this DNA was used at UNH to generate

Nanopore sequence on an Oxford Nanopore Technologies (ONT) R9.4.1 flow cell using a library

prepared with a rapid sequencing kit (SQK-RAD004) following methods described by Nanopore

entitled "Rapid sequencing gDNA – whole genome amplification".

Illumina sequence was generated at the UNH Hubbard Center for Genome Studies (HCGS) for

the following: for the 'P' and 'QC' accessions of *C. ficifolium*; for 35 F2 generation individuals

descended from a 'P' x 'QC' cross that was previously studied and described (Subedi et al., 2021);

the RB6 accession of *C. berlandieri* var. *macrocalycium*; and for the 302-A accession of *C.*

foggii. DNA for Illumina sequencing was isolated at UNH (Subedi et al., 2021) using a CTAB

method described by Torres *et al.* (1993).

Nuclear Genome Assembly

PacBio reads were assembled at BYU using Hifiasm Version v0.18.5-r499 (Cheng *et al.*, 2021,

2022) with default settings. Subsequent correction, annotation, and analyses were conducted at

UNH. Inspector (Chen *et al.*, 2021) was used to detect and correct errors in the Hifiasm assembly

followed by manual examination and adjustments, yielding the *C. ficifolium* Version 1.0 nuclear

reference genome designated as ChenoFicP_1.0. Finally, BUSCO (Manni *et al.*, 2021) and

dependencies (Stanke *et al.*, 2008; Camacho *et al.*, 2009; Eddy, 2011; Mirarab, Nguyen and

Warnow, 2012; Levy Karin, Mirdita and Söding, 2020; Li, 2023) using default settings and the

embryophyta_odb10 database was used to calculate nuclear genome completeness. Telomeres

were identified within pseudochromosomes (PCs) with the quarTeT toolkit program

TeloExplorer (Lin *et al.*, 2023) and within unplaced contigs using tidk (Brown, la Rosa and

Mark, 2023). Centromeric regions were identified using the tool CentroMiner which is supplied

with the quarTeT toolkit (Lin *et al.*, 2023). Nuclear genome assembly data was visualized using

Circos (Krzywinski *et al.*, 2009) in combination with the deepStats toolkit (Richard, 2019) and

BEDOPS (Neph *et al.*, 2012) for track generation.

Nuclear Genome Annotation

Once ChenoFicP_1.0 was finalized, repetitive elements were identified using RepeatModeler

(Flynn *et al.*, 2020). Those repetitive elements were then masked using RepeatMasker (Tarailo-

graovac and Chen, 2009; Flynn *et al.*, 2020; Storer *et al.*, 2021) as the first step in an annotation

pipeline patterned after Card *et al.* (2019). The RepeatMasker scripts

calcDivergenceFromAlign.pl and createRepeatLandscape.pl were used to analyze output files,

determine repetitive element content, and produce data visualizations. A BRAKER IsoSeq

compatible singularity container (Bruuna, Gabriel and Hoff, 2024) was passed the fasta file

containing soft-masked repetitive elements and Pacific Biosciences IsoSeq data aligned to the

genome using minimap2 (Li, 2018). BRAKER flags were set to use the Viridiplantae OrthoDB

database (Kuznetsov *et al.*, 2023) for protein sequence, and the Embryophyta odb10 database

(Manni *et al.*, 2021) as the Busco lineage.

The resulting GFF3 annotation file was passed to MAKER (Holt and Yandell, 2011) as a

prediction GFF file, as were the cds-transcripts and proteins files which were used for EST

evidence and protein homology evidence, respectively. MAKER was used to update annotations

and to provide Annotation Edit Distance (AED) scores. AED values for the final annotation file

were produced using AED_cdf_generator.pl (https://github.com/mscampbell). Once complete,

the output files for each pseudochromosome were merged to create one contiguous file. This

output file contained gene predictions, however without functional annotations. To integrate

functional details and GO terms, BLASTP (Johnson *et al.*, 2008) using the UniProtKB/Swiss-

Prot database (Bateman *et al.*, 2015), as well as InterProScan (Jones *et al.*, 2014) were passed the

protein file produced by MAKER, and the resulting information was integrated into a single file

using the maker_functional_gff and ipr_update_gff commands, and sorted with AGAT (Dainat,

Hereñú and Pucholt, 2020), which yielded the final GFF3 annotation file.

A genome-wide synteny analysis was performed comparing the ChenoFicP_1.0 assembly to the

B subgenome of the *C. quinoa* V2 reference genome (Rey *et al.*, 2023) and to the *Beta vulgaris*

EL10.1 reference genome (McGrath *et al.*, 2023). As a co-member of the Amaranthaceae family,

diploid (2n = 2x = 18) *B. vulgaris* (sugar beet) has been used in previous genomic comparisons

with quinoa (Jarvis *et al.*, 2017; Rey *et al.*, 2023). To produce the synteny analysis, *C. quinoa*

annotations and protein fasta files were downloaded from CoGe (id60716), and *B. vulgaris* files

from NCBI (GCF_002917755.1). Annotation files were converted to BED format via

agat_convert_sp_gff2bed.pl (Dainat, Hereñú and Pucholt, 2020), before being manually

rearranged to comply with requirements for MCScanX_h (Wang *et al.*, 2012). BlastP was used to

compare the proteome of *C. ficifolium* with those of *B. vulgaris* and *C. quinoa* (Camacho *et al.*,

2009), yielding two blast output files which were merged and passed to MCScanX_h, along with

BED files containing positional information for the genes in the blast files. For this analysis,

MCScanX used default filtering parameters. SynVisio (Bandi and Gutwin, 2020) was used to

visualize the resulting files, illuminating rearrangements and structural variations.

Chloroplast Genome Assembly and Annotation

The 'P' chloroplast genome was initially assembled from extracted PacBio reads using

GetOrganelle v1.7.6.1 (Jin *et al.*, 2020) with dependencies SPAdes (Bankevich *et al.*, 2012),

Bowtie2 (Langmead and Salzberg, 2012), BLAST+ (Camacho *et al.*, 2009), and Bandage (Wick

et al., 2015), and annotated with the online tool GeSeq (Tillich *et al.*, 2017). GetOrganelle was

called using get_organelle_from_reads.py and was supplied with raw PacBio reads using the

unpaired '-u' flag. Embplant_pt was selected as the genome type, and the target genome size was

set to 150 kb which is the approximate size of the *C. quinoa* chloroplast genome (Hong *et al.*,

2017). For the purpose of determining the relative abundance of each detected haplotype of the

'P' chloroplast genome based on the numbers of supporting Nanopore reads, Cp-hap (Wang,

Lanfear and Gaut, 2019) was passed reads produced via an ONT R9.4.1 flow cell as previously

noted. The final output file produced from GetOrganelle was passed to GeSeq

(https://chlorobox.mpimp-golm.mpg.de/geseq.html) (Tillich *et al.*, 2017) for visualization and

annotation.

Mitochondrial Genome Assembly and Annotation

A contig from the Hifiasm assembly was identified as mitochondrial by aligning all unplaced

contigs to the quinoa mitochondrial reference sequence NC_041093.1 (Maughan *et al.*, 2019)

with BWA MEM (Li and Durbin, 2009), and manually selecting a long, well-aligned contig. This

contig was used as a seed sequence and passed to a *de novo* organelle assembly pipeline

described by Jarvis et al., (2022) to assemble the mitochondrial genome. The pipeline was used

in two iterations, initially using the HiFi dataset which produced a contig that was passed to the

mitochondrial assembly pipeline in conjunction with the long read Nanopore dataset. The

Nanopore dataset was then aligned to this contig using minimap2 (Li, 2018), reads matching GC

content and length thresholds were stripped from the alignment file via SeqKit (Shen *et al.*,

2016), and these reads were passed to Canu (Koren *et al.*, 2017) for *de novo* assembly. Nanopore

reads were later used to resolve assembly structure via comparing alignments to the draft

mitogenome assembly. The output contig from Canu was manually circularized by locating

regions at the start and end of the file that shared 100 percent sequence identity. The circularized

version of the assembly was polished using NextPolish (Hu *et al.*, 2020) and the PacBio read set

to correct single nucleotide errors. Annotation was performed by GeSeq. Importantly, GeSeq was

set to pass third party mitochondrial NCBI RefSeqs from both *C. quinoa* (NC_041093.1) and

Beta vulgaris subsp. *vulgaris* (NC_002511.2) to BLAT (Kent, 2002) which used annotations

from those genomes to locate and annotate homologs within the *C. ficifolium* mitogenome.

Chloroplast and Mitochondrial Inheritance Patterns in *C. ficifolium*

Chloroplast and mitochondrial reference-guided assemblies for 'QC', one F1 individual, and four F2 individuals, were constructed and aligned to the respective 'P' organelle genomes using BWA mem (Li and Durbin, 2009) to identify heritable organelle polymorphisms differentiating the two parents. The resulting SAM files were then converted to BAM, sorted, indexed, and a consensus was called using the respective samtools modules (Danecek *et al.*, 2021). Once BAM files for each individual were converted to fasta, freebayes-parallel (Garrison and Marth, 2012; Tange, 2018) was used to call variants. Freebayes-parallel was invoked with options to produce a gVCF file, include genotype qualities, use best n alleles set to 4, and cap read depth to 5000. VCF files were filtered to remove low quality calls using vcffilter from vcflib (Garrison *et al.*, 2022) and bcftools (Danecek *et al.*, 2021). Sites or calls were pruned from the chloroplast multisample vcf file if: a) the average depth across all samples was below 100; b) or the site quality was below 998; c) or the genotype quality phred score was below 30; d) or depth for a given sample was below 50; e) or more than half of individuals were not represented; f) or the allele balance was below 25 percent or greater than 75 percent; g) or there was a discrepancy between the number of paired and unpaired reads aligned. The vcf file generated for the mitochondrial genome using the same samples was filtered using the same criteria with the addition of a cap of 3000 on the average depth per sample. The filtering criteria and script were based on those provided by http://www.ddocent.com/filtering/. The resulting filtered vcf files were visualized with IGV (Thorvaldsdóttir, Robinson and Mesirov, 2013).

Chloroplast and Mitochondrial Genome Ancestries

To illuminate the possible role of *C. ficifolium* in the cpDNA ancestries of allotetraploids *C. quinoa* and *C. berlandieri*, a minimal phylogenetic tree was generated using six chloroplast genome (cpDNA) assemblies representing four *Chenopodium* species and including beet (*Beta vulgaris*) as an outgroup. The *C. quinoa* and *B. vulgaris* chloroplast genomes used were NCBI accessions MK159176.1 and OU343016.1, respectively. All other accessions included in this analysis were assembled by us *de novo* from raw reads using GetOrganelle (Jin *et al.*, 2020). We assembled the *C. berlandieri* chloroplast genome from Illumina 2x150 PE reads previously uploaded to NCBI by Mark Samuels from the University of Montreal (Accession: PRJNA895488) (Samuels *et al.*, 2023). The *C. ficifolium* 'P' chloroplast genome was *de novo* assembled using sequence data extracted from the PacBio dataset as described above, while the *C. ficifolium* 'QC' and the *C. foggii* 302-A chloroplast genomes were assembled from Illumina data produced at the UNH-HCGS. These assemblies were aligned using Mafft (Katoh and Standley, 2013) with default (auto) settings, and a tree was produced using RAxML V. 8.2.12 with the GTRCAT model (Stamatakis, 2014). Trees were visualized with TreeViewer (Bianchini and Sánchez-Baracaldo, 2023).

For a parallel purpose, a minimal mitochondrial phylogeny was produced by aligning raw reads, either generated by the HCGS for *C. foggii* 302-A*,* and 'P' and 'QC' *C. ficifolium* accessions or downloaded from NCBI, to the quinoa mitochondrial reference genome NC_041093.1 (Maughan *et al.*, 2019). *B. vulgaris* (SRR6305245) (McGrath *et al.*, 2023) and *C. berlandieri* (PRJNA895488) (Samuels *et al.*, 2023) read sets were downloaded from NCBI. All read data used in this analysis were trimmed and cleaned with Trimmomatic (Bolger, Lohse and Usadel, 2014) using the primer sequence file "TruSeq3-PE-2.fa" invoked with the following settings:

ILLUMINACLIP:TruSeq3-PE-2.fa:2:30:10 LEADING:3 TRAILING:3

SLIDINGWINDOW:4:15 MINLEN:36. FastQC

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) was used to visualize and validate

the Trimmomatic output. These reads were then aligned to the quinoa mitochondrial reference

genome (Maughan *et al.*, 2019) using BWA mem (Li and Durbin, 2009). Using Samtools view,

sort, index, and consensus (Danecek *et al.*, 2021), the BWA produced SAM file was converted to

a fasta file containing the consensus sequence. These six fasta files, those produced from reads

plus the quinoa mitogenome file used as a reference in the alignment process, were merged into a

multi-fasta and Mafft (Katoh and Standley, 2013) was used with default (auto) settings to

produce an MSA. A tree was produced using RAxML V. 8.2.12 (Stamatakis, 2014) with the

GTRCAT model and *B. vulgaris* specified as the outgroup.

Genome Wide Association Studies

A subset of 21 'P' x 'QC' F2 individuals chosen from the 35 sequenced at the HCGS on the basis

of high read depth were used in a GWAS analysis to locate regions throughout the genome

implicated in the control of branch angle, internode length, flowering time, plant height, and

number of lateral branches. BWA MEM was used to align reads from F2 individuals to the *C.*

ficifolium 'P' V1.0 reference genome. Resulting sam files were converted to bam, sorted, and

indexed via samtools (Danecek *et al.*, 2021), and read depths visualized with

WGSCoveragePlotter (Lindenbaum, 2015). GATK AddOrReplaceReadGroups (Van der Auwera

and O'Connor, 2020) was called to change read group IDs to reflect the sample name to make

variant calling possible. Freebayes-parallel (Garrison and Marth, 2012; Tange, 2018) was used to

generate a vcf file containing both InDel and SNP variants for all 35 initial individuals. The

resulting vcf file was filtered to remove low quality calls following guidelines from

http://www.ddocent.com/filtering/. The filtered vcf data in conjunction with phenotypic data

collected by Subedi (2020) (Supplemental Table 1) , and name and positional data extracted from

the GFF3 annotation file were supplied to the package vcf2gwas (Vogt, Shirsekar and Weigel,

2022) which produced Manhattan plots and positional data for putatively causative SNPs.

A region of interest on Cf4 identified via vcf2gwas was then closely examined with respect to

gene content and local synteny and collinearity with the *C. quinoa* V2 assembly (Rey *et al.*,

2023). For this purpose, this approximately 7 Mb region of interest was manually excised from

the full annotation file associated with our ChenoFicP_1.0 assembly, Revigo (Supek *et al.*, 2011)

was used to analyze pathways and relationships of genes with attached GO terms, and gene IDs

were used to subset the proteome fasta file. BlastP and MCScanX were used to compare this

protein subset to the complete proteome of *C. quinoa* and create a homology file. This file, plus

the required concatenated BED files were passed to MCScanX_h, which used default filtering

criteria with the exception of MATCH_SIZE, which was set to 9 to only include matches with 10

or more collinear genes. The MCScanX pipeline developed by BDX consulting (Wang *et al.*,

2024) was used to convert data to a format compatible with Circos (Krzywinski *et al.*, 2009).

Results

Nuclear Genome Assembly

The PacBio HiFi read set from *C. ficifolium* 'P' comprised a total of 26.5 Gb of sequence with an

N50 of ~14 Kb. These data were supplied to Hifiasm Version 18, yielding 781 contigs of

cumulative length 759,656,873 bp (~760 Mb) with an average read depth of 34.8x (Table 1).

Table 1. Hifiasm assembly statistics.

Statistics of contigs:		Read to Contig alignment:	
Number of contigs	780	Mapping rate /%	100.00
Number of contigs > 10000 bp	780	Split-read rate /%	0.47
Number of contigs >1000000 bp	9	Depth	34.87
Total length	759656810	Mapping rate in large contigs /%	92.05
Total length of contigs >1000000bp	729997343	Split-read rate in large contigs /%	0.46
Longest contig	92986964	Depth in large contigs	33.45
Second longest contig length	85613771	Error and Quality Metrics	
N50	79863811	Small-scale assembly error /per Mbp	0.30
L50	5	QV	62.60

Of these contigs, the largest nine ranged in size from a maximum of 92,986,964 bp down to

70,697,261 bp, while the tenth largest (contig ptg000009l) was 8,313,269 bp (Table 2). The

remaining 771 contigs were all much smaller: less than 1 Mbp. On the basis of these size

discontinuities, and the prior expectation based on the *C. ficifolium* chromosome number (n = x =

9) (Walsh *et al.*, 2015), the nine largest Hifiasm contigs were defined as comprising the initial

pseudochromosome (PC) assembly. These nine PCs were then numbered and oriented to

Table 2. Original contig names, pseudochromosome designations, and contig lengths of the
primary assembly plus the two largest unplaced contigs after ptg000009l were appended to Cf7.

Name	Renamed	Initial Length	Final Length
ptg000002l	Cf6	92986964	92986964
ptg000006l	Cf5	85613789	85613789
ptg000005l	Cf3	82834534	82834534
ptg000008l	Cf7	74073315	82386584
ptg000003l	Cf2	79863818	79863818
ptg000001l	Cf8	79853931	79853931
ptg000011l	Cf1	77970451	77970451
ptg000004l	Cf4	77790002	77790002
ptg000007l	Cf9	70697261	70697261
ptg000009l	(Merged with Cf7)	8313269	------------
ptg000020l		871519	871519
ptg000025l		721674	721674

maximize correspondence with the *Beta vulgaris* EL10.1 reference genome (McGrath *et al.*,

2023), as was done previously in the assembly of the quinoa Version 1 assembly (Jarvis *et al.*,

2017). Inspector subsequently identified and corrected three structural errors, and 228 small-

scale assembly errors, reporting a QV score of 63.39 after correction for this assembly (Figure

1). The G/C content of this genome is 36.9 percent, and varies between approximately 20 and 80

percent per 100 Kb window (Figure 1 Ring D). G/C rich regions are often centrally located

within pseudochromosomes; however, centromeric regions on the whole have an average G/C

content, with pseudochromosome Cf1 having the highest content with 45.47 percent, and the

average across all centromeres being 37.75 percent. When comparing the *C. ficifolium* 'QC'

accession to the *C. ficifolium* 'P' assembly (Figure 1 Ring C), the number of SNPs differentiating

the two per 100 Kb window ranges from as low as 12 to a maximum of 2,321, or approximately

one SNP per 43 bases, as found on PC Cf1 from base 11,800,000 to base 11,900,000.

The initial Hifiasm assembly was telomere-to-telomere for PCs Cf3, Cf4, Cf5, Cf6, Cf8,

and Cf9, with Cf1 and Cf2 having only one telomere each, and Cf7 having a partially assembled

telomere which was oriented in a backwards direction. A homology search of contig ptg0000009l,

the only unplaced contig larger than 1 Mbp, relative to the quinoa B subgenome (Rey *et al.*,

2023) revealed that this contig was syntenic to the distal end of quinoa chromosome Cq7B,

including a telomeric sequence, and thus directed its placement onto Cf7. Following this

adjustment, the final "ChenoFicP_1.0" assembly of nine pseudochromosomes had a total length

of 729,997,451 bp (~730 Mb). The CentroMiner package identified one region per chromosome

as being centromeric (Supplemental Table 2) and produced positional data. Centromeric and

telomeric positional data are provided in Supplemental Table 2. BUSCO completeness of the

344   reference genome without unplaced contigs is 96.7 percent, and with unplaced contigs is 97.2

345   percent (Figure 2B).

*Figure 1. This assembly of the C. ficifolium 'UNH_ChenoFicP_1.0' nuclear genome comprises nine pseudochromosomes and spans 729.9 Mb. **A.** Gene density ranging from 20 to 3000 genes represented per 100 kb window. **B.** Repetitive element density ranging from 20 to 300 elements per 100 kb window **C.** SNPs present compared to C. ficifolium accession 'QC' per 100 kb window. **D.** G/C content. Graph represents 20 - 55 percent G/C.*

The remaining 771 unplaced contigs comprised a total of about 29.6 Mb, of which 594 contigs

(19.6 Mb) and 6 contigs (626 Kb) were subsequently assigned on the basis of sequence

homology to the chloroplast and mitochondrial genomes, respectively, leaving a final total of 171

contigs (9.4 Mb) unplaced. Further investigation into unplaced contigs using tidk indicated that

contigs ptg000020 (871,519 bp) and ptg000025 (721,674 bp) both contain telomeric sequence.

These contigs were found by BWA to likely belong on the upstream ends of Cf2 and Cf7

respectively; however, they have not been included in the ChenoFicP_1.0 assembly pending

further confirmation based upon anticipated long read data.

Repetitive Element Identification and Genome Annotation

Data produced by RepeatModeler indicates that 69.9 percent of the ChenoFicP_1.0 genome is

comprised of repetitive elements, with the largest contributors being LTR/Gypsy (33.8 percent)

and LTR/Copia (11.2 percent) type repetitive sequences, while unknown repeats account for 17.5

percent of the total genome fraction. These repetitive elements tend to cluster towards telomeres

in (Figure 1 Ring B). Non-repetitive sequence accounts for 30.1 percent of the genome (Figure

2A).

*Figure 2 **A**. Repetitive element analysis and summary graph produced by RepeatModeler. Green: LTR elements, Grey: Unknown repetitive elements, Black: Non-repetitive sequence. **B**. BUSCO analyses of completeness for the masked genome used in the annotation process (top), the proteome (middle), and transcriptome (bottom). **C**. AED (annotation edit distance) scores of gene reannotations produced by Maker. Lower AED values correspond with higher levels of agreement between the annotation and evidence, with 0 representing perfect agreement between the two.*

The final annotation file was used to calculate relative gene density (Figure 1 Ring A). The

BUSCO analysis yielded a transcriptome and proteome completeness of 98.4 percent (Figure

2B). Resultant AED values for the final annotation are graphed in Figure 2C. In total, 22,617

genes were identified and annotated with the average gene length (including introns and UTRs,

as defined by Maker utilizing unprocessed mRNA transcripts) being 3,958 bp.

Synteny with Quinoa B-subgenome Chromosomes

The MCScanX synteny analysis of the ChenoFicP_1.0 assembly relative to *B. vulgaris* EL10.1

(McGrath *et al.*, 2023) and the quinoa V2 assembly B-subgenome (Rey *et al.*, 2023) revealed

numerous structural variations. In both comparisons, translocational differences tended to

involve small distal regions near telomeres. Large inversional differences were revealed in

ChenoFicP_1.0 chromosomes Cf1, Cf3, and Cf4 relative to the quinoa B-subgenome

homologues, the largest of which was on Cf3 and spanned approximately 63.2 Mb (Figure 3).

These inversions were positioned centrally, and in every case for chromosomes Cf1, Cf3, and

Cf4 were pericentric.

*Figure 3. Synteny conservation diagram comparing the assembled pseudochromosomes of C.
ficifolium (center) to the B subgenome of quinoa (top), and B. vulgaris EL10.1 (bottom).
Chromosomes 1, 4, 5, 7, and 8 of the quinoa B subgenome were manually reoriented (flipped –
indicated by arrows) to match those of C. ficifolium for illustrative purposes.*

Chloroplast Genome Assembly

GetOrganelle produced two versions of the *C. ficifolium* chloroplast genome assembly, termed "Path 1" and "Path 2", both 151,826 bp in length, but differing in their orientations of the Short Single Copy (SSC) region relative to the Long Single Copy (LSC) region (Figure 4A). When supplied with approximately 263 Mb of Nanopore reads, Cp-hap (Wang, Lanfear and Gaut, 2019) returned approximately equal long read support for both Path 1 (248 reads) and Path 2 (242 reads) orientations, where the Path 2 orientation (Figure 4A, lower) coincided with that of the quinoa chloroplast assembly (MK159176.1) reported by Maughan et al. (2019).

*Figure 4 **A**. Both assembly versions of the C. ficifolium chloroplast genome represented by 'Path 1' and 'Path 2'. These assemblies differ only in the direction of the Short Single Copy region. 'Path 2' (lower) was concordant with the C. quinoa chloroplast genome (MK159176). Genes inside the circle are transcribed in the clockwise direction, genes outside are transcribed counterclockwise. **B**. Gene map of the C. ficifolium mitogenome. The mitochondrial genome contains 118 protein coding genes, 23 tRNA genes, and is ~304 Kb in length.*

In both Path 1 and Path 2 orientations, the large single copy region is 83,900 bp, the short single

copy is 18,168 bp, and each of the inverted repeat regions are 15,109 bases in length. The

chloroplast genome regional boundaries are specified in Table 3. The assembly, including both

inverted repeat regions, has a GC content of 37.3 percent and comprises 131 genes, of which

there are 86 protein-encoding, 37 tRNA, and 8 rRNA genes. For comparison, the previously

published (Kim et al., 2019) *C. ficifolium* chloroplast genome assembly (MK182725) had 129

genes, including 84 protein-encoding, 37 tRNA, and 8 rRNA.

Table 3. Start and end positions of the four regions of the chloroplast assembly. Note that these
positions apply to both Path 1 and Path 2 haplotypes; however, the SSC region is inverted in
Path 1 relative to Path 2.

Cp Region	Start	End	No. Genes	Protein Encoding
LSC	1	83670	85	63
IR	83671	108779	16	5
SSC	108780	126717	14	13
IR	126718	151826	16	5
		Total	131	86

Mitochondrial Genome Assembly

A 275 kb contig containing a partially assembled mitochondrial genome from the 'P' accession

was initially identified from a preliminary Hifiasm assembly. In an iterative sequence seeding

process, Nanopore reads were aligned to this contig, extracted from the resulting alignment,

assembled *de novo*, and reads were once again aligned. After two rounds of this process, a contig

was produced that was manually circularized, and polished with NextPolish using HiFi reads.

The resulting 'P' mitochondrial genome assembly has a GC content of 43.8 percent and

comprises 90 genes, of which 33 are protein-coding (Figure 4B). The *C. ficifolium* mitochondrial

genome assembly is similar in length to that of *C. quinoa* (MK182703), at 303,742 bp versus

319,505 bp respectively, and in protein-coding genes, with 33 and 30 respectively (Maughan *et*

al., 2019), although these two assemblies display several structural differences.

Organelle Genomes' Modes of Transmission

When plastid genome mode of transmission was examined using multisample vcf files

containing both parents, 'P' and 'QC', an F1, and four F2 individuals, two distinct sequence

haplotypes were revealed. Of the respective polymorphisms, two were single nucleotide indels,

while the remaining 17 were SNPs (Supplemental Table 3). The F1 and all F2 individuals share

the same sequence haplotype as the maternal parent ('P') as distinct from the 'QC' paternal

haplotype (Figure 5A), thereby demonstrating maternal transmission of the chloroplast genome

in the 'P' x 'QC' cross used to generate the respective F1 and F2 progenies.

To determine the transmission pattern of the mitochondrial genome in the 'P' x 'QC' cross, 'QC'

short read alignment to the 'P' mitochondrial reference genome detected nine polymorphisms, of

which eight were informative, with the uninformative polymorphism being a SNP for which all

individuals had a non-reference nucleotide. Seven of those informative polymorphisms were

SNPs, while the eighth was a 22 bp indel (Figure 5B, Supplemental Table 3). These data

demonstrate that transmission of the mitochondrial genome in the 'P' x 'QC' cross was maternal.

*Figure 5 **A**. Chloroplastic 'Path 2' polymorphisms found among three F2 individuals, one F1 individual, and the maternal ('QC') and paternal ('P') parents in the cross that yielded the F1 individual. **B**. Mitochondrial polymorphisms found among three F2 individuals, one F1 individual, and the male and female within the cross that yielded the F1 individual, 'QC' and 'P' respectively. Interestingly one polymorphism shared across all individuals is a heterozygous A/C SNP.*

Chloroplast and Mitochondrial Genome Ancestries

A phylogeny based on *de novo* assembled chloroplast genomes from *B. vulgaris, C. quinoa, C. berlandieri* var. *macrocalycium, C. foggii,* and both the 'P' and 'QC' *C. ficifolium* accessions (Figure 6A) revealed a close affinity of the two *Chenopodium* allotetraploids with the A-genome diploid *C. foggii*, and a more distant relationship with B-genome diploid *C. ficifolium*. Similarly, mitochondrial reference-based assemblies utilizing the same set of germplasm (Figure 6

B) in a narrowly focused phylogeny detected the same pattern of affinities. These outcomes

support the prior hypothesis (Maughan *et al.*, 2019) that the organelle genomes of the AABB

allotetraploids were both contributed by the AA ancestral diploid.

*Figure 6 **A**. Phylogenetic tree illustrating chloroplast genome ancestral relationships among*
Chenopodium AA and BB diploids and AABB tetraploids in our study. Phylogenetic tree
illustrating chloroplast genome ancestral relationships among Chenopodium AA and BB diploids
and AABB tetraploids in our study. This phylogenetic tree is based on chloroplast assemblies
from C. foggii, both C. ficifolium accessions, C. quinoa, C. berlandieri, and Beta vulgaris which
*is set as the outgroup. **B**. Phylogenetic tree illustrating mitochondrial genome ancestral*
relationships among Chenopodium AA and BB diploids and AABB tetraploids. This phylogenetic
tree was constructed using reference guided assemblies with the C. quinoa mitogenome
(NC_041093.1) selected as the reference genome. Beta vulgaris is set as the outgroup.

Genome Wide Association Study

Of the five trait-specific GWAS analyses performed, two (branch angle and internode length)

failed to detect genomic associations. The remaining three traits – days to flowering, plant

height, and branch number – were found by GWAS to correlate with a single common region of

chromosome Cf4 (Figure 7 A-C). Of these three traits, the correlated SNPs for branch number

covered the largest region and fully contained those implicated in the plant height and days to

flower analyses. The implicated region spans 7,020,561 bp between positions 70,765,247 and

77,785,808 on Cf4. Features within this region were pulled from the annotation file to create a

list of gene names and GO terms associated with those genes. There are 770 annotated genes

within the defined region. Of those, 132 were of unknown function. GO terms were assigned to

403 genes, including *FTL1*, which at position 74,807,401 – 74,811,571 is number 425 on the list

of 770 genes, and is annotated as *Vernalization 3*. *FTL1* had one product classification identified

by Interpro (Paysan-Lafosse *et al.*, 2023), being a phosphatidylethanolamine-binding protein

(IPR008914).

*Figure 7 **A-C**. GWAS Manhattan plots indicating regions of the genome implicated in the control*
*of plant height, number of branches, and flowering time. **D**. Synteny analysis of the ~7 Mb region*
found in C. ficifolium Cf4 (Left) implicated by the genome wide association study as compared to
C. quinoa chromosomes Cq4A and Cq4B as oriented in Rey et al., (2023). Both quinoa
chromosomes Cq4A and Cq4B are in reverse orientation compared to C. ficifolium chromosome
Cf4.

A microsynteny analysis focusing on the ~7 Mb window implicated in this GWAS was

performed to identify homeologs within both *C. quinoa* subgenomes and their locations (Figure 7

D). MCScanX identified nine collinear blocks of 10 or more genes from this region in *C.*

ficifolium that also occur, but in reverse orientation, on quinoa chromosomes Cq4A and Cq4B of

the Rey et al., (2023) assembly, as will be discussed later. The analysis of these data indicates

that for the 770 genes found in *C. ficifolium* annotation file at this ~7 Mb region, there were 581

homoeologous genes found within the *C. quinoa* A subgenome and 610 found within the B

subgenome. In total, 107 of the *C. ficifolium* genes within this region are of unknown function

and of those, only six have associated GO terms.

Discussion

Nuclear Genome

As described here the ChenoFicP_1.0 assembly provides a valuable component to the *C.*

ficifolium diploid (BB) model system, as will be expanded upon in subsequent reports on

ongoing investigations at the diploid and allotetraploid (AABB) levels in *Chenopodium*.

Pseudochromosomes within this *C. ficifolium* assembly have been numbered to correspond with,

and oriented to be syntenic and collinear with, the previously published *Beta vulgaris* (2n = 2x =

18) EL10.1 genome (McGrath *et al.*, 2023), thereby following the example of Jarvis et al.,

(2017) in which the quinoa V1 assembly was similarly oriented. The fact that the quinoa V2

assembly (Rey *et al.*, 2023) did not follow this orientation pattern opens the possibility for some

confusion, especially with respect to chromosomes Cq1B, Cq4B, Cq5B, Cq7B and Cq8B, which

are presented and numbered in reverse orientation in quinoa V2 as compared with quinoa V1 and

beet EL10.1. We have followed the initial orientation approach in conjunction with the recently

published *Chenopodium* pangenome analysis (Jaggi *et al.*, 2024), which employs the original

orientation pattern.

Beet was initially selected as the genome comparator for *Chenopodium* because it was the first

species to have a published reference genome within the subfamily Chenopodioideae. While the

diploid *Spinacia oleracea* (2n = 2x = 12) now has a published genome (Cai *et al.*, 2021), its

nuclear genome has only six chromosomes, making comparisons with it suboptimal for

identifying syntenic relationships and orienting chromosomes of newly sequenced species.

Genome assemblies have been reported for three other *Chenopodium* diploids: the AA diploids

C. pallidicaule (Jarvis *et al.*, 2017), *C. watsonii* (Young *et al.*, 2023), and the BB diploid *C.*

suecicum versions 1 (Jarvis et al., 2017) and 2 (Rey et al., 2023). However, to our knowledge, no

gene-trait association studies have yet been conducted in these species.

The ChenoFicP_1.0 nine-pseudochromosome assembly size of ~730 Mbp incorporates 96% of

the 760 Mbp total assembly for the 781 contigs generated by Hifiasm. Cytometrically determined

C-values of *C. ficifolium* 'Portsmouth' predict a genome size in the range of 821 to 831 Mbp

(Neff, 2017). For comparison, the nine B-subgenome pseudochromosomes of the quinoa

assembly sum to 670 Mbp in length (Rey *et al.*, 2023), or approximately 8% smaller than the

ChenoFicP_1.0 assembly. Similarly, the nine pseudochromosomes of the quinoa A subgenome

sum to 531 Mbp (Rey *et al.*, 2023), which is 3% smaller than the diploid *C. watsonii* 548 Mbp A-

genome assembly (Young *et al.*, 2023). An earlier, broad survey of *Chenopodium* germplasm

reported C values of 893 Mbp for a Eurasian accession of *C. ficifolium*, 645 Mbp for *C. watsonii*,

and 1.545 Gbp for *C. quinoa* (Mandák *et al.*, 2016) Thus, all three of the aforementioned

Chenopodium genome assemblies are somewhat smaller than the size predicted by cytometric

assay. It has been speculated that this pattern of discrepancy may be due to incomplete assembly

coverage of some repetitive regions (Young et al., 2023), or it could be due to a consistent

technical artifact of flow cytometric analyses.

The ChenoFicP_1.0 genome contains 22,617 genes (total genes), of which 5,060 encode proteins

that have yet to be characterized within the UniProtKB/Swiss-Prot database. Gene density is

greatest in the distal regions of chromosomes (Figure 1 Ring A). For comparison, the total gene

counts of the *C. quinoa* A and B subgenomes are 25,913 and 26,942 respectively (Rey *et al.*,

2023), the AA diploid *C. watsonii* count was 30,325 (Young *et al.*, 2023), and the BB diploid *C.*

suecicum count was 29,702 (Rey *et al.*, 2023). Compared with the ChenoFicP_1.0 assembly, the

total number of annotated genes within the *C. quinoa* B subgenome is approximately 19.1

percent higher. Given that different gene count methodologies were used in these various studies,

the observed differences in gene counts among subgenomes and species should be interpreted

with caution.

RepeatModeler indicated that repetitive elements account for approximately 70 percent (510

Mbp) of the *C. ficifolium* genome. For comparison, the quinoa repetitive element content was

approximately 65% for the total genome (Zou *et al.*, 2017; Rey *et al.*, 2023), and approximately

63% (421 Mbp) for the B subgenome, with the largest contributors being long terminal repeat

retrotransposons representing 46 percent of the total genome size. Over half of the *C. ficifolium*

repetitive elements are retrotransposons that cluster towards the center of chromosomes (Figure 1

Ring B). These data indicate that there has been some change in the repetitive element landscape

in *C. ficifolium* and/or the *C. quinoa* B subgenome subsequent to the quinoa allotetraploid

speciation event.

Organelle Genome Assemblies

The generation of two different chloroplast genome assembly versions, designated Paths 1 and 2,

was an unexpected outcome. Prior chloroplast genome assemblies in *Chenopodium*, including *C.*

quinoa (Maughan *et al.*, 2019; Gao *et al.*, 2021) and *C. ficifolium* (Kim, Chung and Park, 2019),

and in the broader Amaranthaceae family (Chaney *et al.*, 2016; Dong *et al.*, 2016; Kim, Chung

and Park, 2019; Ding *et al.*, 2021) including sugar beet (McGrath *et al.*, 2023), have reported

only a single chloroplast genome configuration, corresponding to our Path 2 assembly, although

Kim et al. (2019b) mention the need to reposition the SSC and LSC regions of the quinoa

accession 'Real Blanca' for their alignment procedure, indicating a disagreement in SSC and/or

LSC orientation when compared to other assemblies. However, recent literature (Wang, Lanfear

and Gaut, 2019) (and citations therein) has established that chloroplast genomes do indeed exist

in two equally represented alternate configurations in many if not most plant species, as has now

become evident in *Chenopodium*, and that the detection of these alternate forms is enhanced by

employment of long read sequencing technologies such as PacBio HiFi and Nanopore. The very

long and accurate reads now available from Nanopore technology will be of particular value in

relation to mitochondrial genomes, the potential alternate forms of which are often challenging to

assemble (Kozik *et al.*, 2019). Given the known complexities of mitochondrial genome

organization, the assembly configuration that we present here should be viewed as preliminary

and subject to revision.

Organelle Transmission Patterns

In the absence of evidence to the contrary, organelle genomes are generally assumed to be

transmitted maternally in Angiosperms: however, exceptions have been reported (Davis *et al.*,

2010), including paternal chloroplast transmission in some *Actinidia* hybrids (Testolin and

Cipriani, 1997). Until very recently (Maughan *et al.*, 2024), empirical documentation of modes

of organelle inheritance in *Chenopodium* appears to have been limited to a single study involving

atrazine resistance in *Chenopodium album* (Warwick and Black, 1980; Corriveau and Coleman,

1988; Harris and Ingram, 1991; Kolano *et al.*, 2016). Our analysis agrees with this prior study in

its finding of maternal transmission of both chloroplast and mitochondrial markers in our diploid

level *C. ficifolium* cross. Similarly, Maughan et al. (2024) have documented the maternal

transmission of the organelle genomes in several tetraploid level *Chenopodium* crosses.

Organelle Genome Ancestries

Although *C. ficifolium* is a candidate B nuclear genome donor in the initial hybridization that

formed quinoa 3.3 to 6.3 million years ago (Štorchová *et al.*, 2015; Kolano *et al.*, 2016; Jarvis *et*

al., 2017; Maughan *et al.*, 2019), prior analyses indicate that a BB diploid was not the ancestral

donor of the allotetraploids' organelle genomes (Maughan *et al.*, 2019). Instead, that role has

been assigned to the A nuclear genome donor (Maughan *et al.*, 2019). That assignment is

supported by our chloroplast and mitochondrial minimal phylogenies, which both place *C.*

ficifolium as sister to a clade consisting of AABB tetraploids and an AA diploid.

It is considered likely that, in the initial AABB hybridization event, the ancestral AA

diploid served as female and maternal organelle genome donor (Maughan *et al.*, 2019).

Documentation of such in our *C. ficifolium* cross and in tetraploid level crosses (Maughan *et al.*,

2024) strengthens the thesis that the original hybridization event was of the form AA x BB, i.e.,

with the AA diploid ancestor serving as female. Final validation of this model will await the

analysis of organelle genome transmission in reciprocal crosses between AA and BB diploids.

Genome Wide Association Studies

Genome Wide Association Studies

The data produced from these genome-wide analyses indicate that, in the studied segregating

population, variation in the agronomically important traits of days to flower, plant height, and

branch number were predominantly, if not entirely, under the control of one or more genes within

a single, 7 Mb region located towards the bottom of pseudochromosome Cf4 and containing the

FTL1 locus previously employed as a marker by Subedi et al. (2021). Our genome-wide analysis

extends the findings of Subedi *et al*. (2021) by focusing attention on this Cf4 "hot spot" and the

genes residing therein. Of note, the association of branch angle with a single SNP on Cf9 was not

further investigated due to its presumed spurious nature. Manhattan plots will typically associate

a number of SNPs within and surrounding a causative region, and this questionable SNP did not

have any significantly associated markers adjacent to it, nor was it seen in the other two analyses

investigating plant height or days to flower.

Gene ontology data submitted to Revigo indicated the presence of a myriad of genes responsible

for controlling functions that are not intrinsically associated with these traits of interest; however,

this region contains *Flowering Locus T-Like 1* (*FTL1*), which was automatically annotated as

Vernalization 3 (*VRN3*) within the GFF file. This gene is between positions 74,807,401 and

74,811,571, placing it centrally within the denoted significance region on Cf4. *FTL1* had one

product classification identified by Interpro (Paysan-Lafosse *et al.*, 2023), being a

phosphatidylethanolamine-binding protein (IPR008914).

To further understand relationships between *C. ficifolium* and *C. quinoa,* and functionally utilize

C. ficifolium as a model species, BlastP and MCScanX microsynteny analyses were performed to

investigate relative gene locations across species and subgenomes (Figure 7D). When comparing

the *C. ficifolium* region of interest to both of the quinoa subgenomes, all homologous genes in

syntenic blocks of ten or more genes were found to reside in corresponding distal regions of

quinoa chromosomes Cq4A and Cq4B, reflecting substantial gene order and location

conservation within the respective genomic window. This analysis identified quinoa genes

CQ035263 (A subgenome) and CQ019992 (B subgenome) within the *C. quinoa* V2 annotation

(GFF3) file as being homologous to *C. ficifolium VRN3* (*FTL1*). The product of CQ035263 was

queried with BlastP and has an 83 percent sequence identity match to XP_021772286.1, *C.*

quinoa FLOWERING LOCUS T-like. Similarly, the product of gene CQ019992 is identical to

XP_021734330.1. *C. quinoa FLOWERING LOCUS T-like* has a 97% protein identity with the

annotated *C. ficifolium* gene Cfic_20093. The data produced using *C. ficifolium* as a model

organism for *C. quinoa* via this pipeline has been demonstrated to be an efficacious way of

establishing gene-trait relationships via GWAS and locating syntenic regions across species that

may be worth investigating to understand agronomically relevant traits. Further investigation of

genes within this region is required, perhaps in conjunction with more high-depth sequence data

and additional plants to facilitate a fine-mapping study; however, these data are strongly

indicative of one or more genes within this region as being implicated in the control of these

traits. Further research, likely employing gene editing or transformation methods, is required to

test this hypothesis and firmly establish this relationship.

The results of this study, in particular the synteny and collinearity analysis comparing *C.*

ficifolium and quinoa 'QQ74' chromosomes, presents intriguing hints regarding the potential

usefulness of figleaf goosefoot as a breeding resource for quinoa improvement. The primary

structural variants differentiating the B genome chromosomes in these two species are three large

pericentric inversions. However, the largest of these, on chromosome Cq3B, was a

rearrangement present only in a subset of quinoa lines derived from lowland Chilean germplasm,

with at least 80% of quinoa lines expected to have collinear Cq3B in comparison with figleaf

goosefoot (Rey *et al.* 2023). The remaining two inversions, on Cq1B and Cq4B, appear to have

breakpoints extending less than halfway out into their respective chromosome arms, such that

they do not encompass recombinogenic regions (Maughan *et al.* 2024). Consequently, while

triploid (AB_qB_f) figleaf goosefoot x quinoa hybrids are expected to be highly sterile, backcross

hybrids might theoretically be recoverable and, if so, we might expect a high degree of pairing

and recombination between their B genomes. This approach – crossing of the polyploid crop to

its wild, diploid progenitor - has been successfully employed to introduce numerous resistance

genes from the wild progenitor goatgrass (*Aegilops tauschii*, DD) into bread wheat (*Triticum*

aestivum, AABBDD) while restricting recombination, and consequently undesirable linkage

drag, to a single subgenome (Cox, Raupp and Gill, 1994).

Conclusion

The development of high-quality annotated reference nuclear and chloroplast genomes

and a preliminary mitochondrial genome for *C. ficifolium* serves as a foundational advance in the

development of this species as a BB diploid model system for related allotetraploids of interest,

with a primary focus on *C. quinoa* and *C. berlandieri.* Looking ahead, next steps in the

development and implementation of this model system include: broadening the germplasm base

via plant collection in diverse environments; establishing new hybrids and populations

segregating for additional traits of interest including male sterility (Ward, 1998), and disease

resistance, specifically downy mildew (Nolen *et al.*, 2022); and the establishment of effective

protocols for gene editing and *in vitro* culture and regeneration for both quinoa and its diploid

model(s), with immediate focus on manipulation of *FTL1*. The fact that the organelle genomes

present in allotetraploid quinoa were likely derived via maternal transmission from its AA

diploid ancestor, not from its BB ancestor, points to the need for parallel development of an AA

diploid model system in *Chenopodium*. Foundational steps in this direction have already been

taken via the generation of genome assemblies for AA diploids *C. pallidicaule* (Mangelson *et al.*,

2019) and *C. watsonii* (Young *et al.*, 2023), and useful germplasm collection in *C. foggii* (Neff,

Sullivan and Davis, 2018). As yet, to our knowledge, no genetic crosses or segregating progeny

populations have yet been generated in an AA diploid, so this should be a top priority going

forward. Finally, opportunity exists for breeding of quinoa with figleaf goosefoot as an avenue

for gene introgression from the diploid to the tetraploid level.

Acknowledgements

The research reported in this publication was partially funded by the New Hampshire

Agricultural Experiment Station. This is Scientific Contribution Number 3032. This work is/was

supported by awards to Thomas Davis, UNH, from the USDA National Institute of Food and

Agriculture (Hatch) Project NH00678 (accession number 1019990), and by USDA NIFA project

110390. Ed Wilcox and the BYU DNASC (RRID: SCR_017781) provided PacBio sequencing

services, as compensated by USDA NIFA project 110390.

Conflicts of Interest

These authors have not declared any conflicts of interest.

ORCID iDs

Clayton D. Ludwig https://orcid.org/0009-0004-8921-2479

Peter J. Maughan https://orcid.org/0000-0003-3714-3411

Eric N. Jellen https://orcid.org/0000-0002-7906-4845

Thomas M. Davis https://orcid.org/0000-0001-5455-0524

Data Availability

The reference genome and sequence datasets used to create it are available from NCBI using IDs
SUB12919328 and SRR28959327, respectively. The reference genome can be found using the
search term "UNH_ChenoFicP_1.0". The reference assembly, annotation data in GFF3 format,
and supplemental annotation files can be accessed via figshare.com:
https://doi.org/10.6084/m9.figshare.c.7590620. Sequence data for F2 individuals used in GWAS
analyses are hosted on NCBI: https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1262536. Code
used can be found on GitHub: https://github.com/UNH-DavisLab/Ficifolium-Genome-V1/

References

Van der Auwera, G.A. and O'Connor, B.D. (2020) *Genomics in the cloud: using Docker, GATK,*
and WDL in Terra. O'Reilly Media.

Bandi, V. and Gutwin, C. (2020) 'Interactive exploration of genomic conservation', *Proceedings*
- Graphics Interface, 2020-May.

Bankevich, A. *et al.* (2012) 'SPAdes: A new genome assembly algorithm and its applications to
single-cell sequencing', *Journal of Computational Biology*, 19(5), pp. 455–477. Available at:
https://doi.org/10.1089/cmb.2012.0021.

Bateman, A. *et al.* (2015) 'UniProt: A hub for protein information', *Nucleic Acids Research*,
43(D1), pp. D204–D212. Available at: https://doi.org/10.1093/nar/gku989.

Bazile, D., Jacobsen, S.-E. and Verniau, A. (2016) 'The Global Expansion of Quinoa: Trends
and Limits', *Frontiers in Plant Science*, 7(May), pp. 1–6. Available at:
https://doi.org/10.3389/fpls.2016.00622.

Bianchini, G. and Sánchez-Baracaldo, P. (2023) 'TreeViewer Version 2.1.0'. Zenodo. Available
at: https://doi.org/10.5281/zenodo.7768344.

Bolger, A.M., Lohse, M. and Usadel, B. (2014) 'Trimmomatic: A flexible trimmer for Illumina
sequence data', *Bioinformatics*, 30(15), pp. 2114–2120. Available at:
https://doi.org/10.1093/bioinformatics/btu170.

Brown, M., la Rosa, P.M. and Mark, B. (2023) 'A Telomere Identification Toolkit'. Zenodo.
Available at: https://doi.org/10.5281/zenodo.10091385.

Bruuna, T., Gabriel, L. and Hoff, K.J. (2024) 'Navigating Eukaryotic Genome Annotation
Pipelines: A Route Map to BRAKER, Galba, and TSEBRA', *arXiv preprint arXiv:2403.19416*
[Preprint].

Cai, X. *et al.* (2021) 'Genomic analyses provide insights into spinach domestication and the
genetic basis of agronomic traits', *Nature Communications*, 12(1), p. 7246. Available at:
https://doi.org/10.1038/s41467-021-27432-z.

Camacho, C. *et al.* (2009) 'BLAST+: Architecture and applications', *BMC Bioinformatics*, 10,
pp. 1–9. Available at: https://doi.org/10.1186/1471-2105-10-421.

Card, D.C. *et al.* (2019) 'GBE Genomic Basis of Convergent Island Phenotypes in Boa
Constrictors', 11(11), pp. 3123–3143. Available at: https://doi.org/10.1093/gbe/evz226.

Cháb, D. *et al.* (2008) 'Two FLOWERING LOCUS T (FT) homologs in Chenopodium rubrum
differ in expression patterns', *Planta*, 228(6), pp. 929–940. Available at:
https://doi.org/10.1007/s00425-008-0792-3.

Chaney, L. *et al.* (2016) ' The complete chloroplast genome sequences for four Amaranthus
species (Amaranthaceae) ', *Applications in Plant Sciences*, 4(9), pp. 4–9. Available at:
https://doi.org/10.3732/apps.1600063.

Chase, M.W. *et al.* (2016) 'An update of the Angiosperm Phylogeny Group classification for the

orders and families of flowering plants: APG IV', *Botanical Journal of the Linnean Society*, 181(1), pp. 1–20. Available at: https://doi.org/10.1111/boj.12385.

Chen, Y. *et al.* (2021) 'Accurate long-read de novo assembly evaluation with Inspector', *Genome Biology*, 22(1), pp. 1–21. Available at: https://doi.org/10.1186/s13059-021-02527-4.

Cheng, H. *et al.* (2021) 'Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm', *Nature Methods*, 18(2), pp. 170–175. Available at: https://doi.org/10.1038/s41592-020-01056-5.

Cheng, H. *et al.* (2022) 'Haplotype-resolved assembly of diploid genomes without parental data', *Nature Biotechnology*, 40(9), pp. 1332–1335. Available at: https://doi.org/10.1038/s41587-022-01261-x.

Corriveau, J.L. and Coleman, A.W. (1988) 'Rapid screening method to detect potential biparental inheritance of plastid DNA and results for over 200 angiosperm species', *American Journal of Botany*, 75(10), pp. 1443–1458. Available at: https://doi.org/https://doi.org/10.1002/j.1537-2197.1988.tb11219.x.

Cox, T.S., Raupp, W.J. and Gill, B.S. (1994) 'Leaf rust-resistance genes Lr41, Lr42, and Lr43 transferred from Triticum tauschii to common wheat', *Crop science*, 34(2), pp. 339–343.

Dainat, J., Hereñú, D. and Pucholt, P. (2020) 'AGAT: Another Gff Analysis Toolkit to handle annotations in any GTF', *GFF format. Zenodo*, 431.

Danecek, P. *et al.* (2021) 'Twelve years of SAMtools and BCFtools', *GigaScience*, 10(2), pp. 1–4. Available at: https://doi.org/10.1093/gigascience/giab008.

Davis, T.M. *et al.* (2010) 'Chloroplast DNA inheritance, ancestry, and sequencing in Fragaria', *Acta Hort*, 859, pp. 221–228. Available at: http://www.actahort.org/books/859/859_25.htm.

Ding, D. Bin *et al.* (2021) 'Characterization and phylogenetic analysis of the complete chloroplast genome of Amaranthus viridis (Amaranthaceae)', *Mitochondrial DNA Part B: Resources*, 6(9), pp. 2610–2612. Available at: https://doi.org/10.1080/23802359.2021.1961631.

Dong, W. *et al.* (2016) 'Comparative analysis of the complete chloroplast genome sequences in psammophytic Haloxylon species (Amaranthaceae)', *PeerJ*, 2016(11), pp. 1–21. Available at: https://doi.org/10.7717/peerj.2699.

Eddy, S.R. (2011) 'Accelerated profile HMM searches', *PLoS Computational Biology*, 7(10). Available at: https://doi.org/10.1371/journal.pcbi.1002195.

Eid, J. *et al.* (2009) 'Real-time DNA sequencing from single polymerase molecules', *Science*, 323(5910), pp. 133–138. Available at: https://doi.org/10.1126/science.1162986.

Flynn, J.M. *et al.* (2020) 'RepeatModeler2 for automated genomic discovery of transposable element families', *Proceedings of the National Academy of Sciences of the United States of America*, 117(17), pp. 9451–9457. Available at: https://doi.org/10.1073/pnas.1921046117.

Gao, M.Z. *et al.* (2021) 'Complete chloroplast genome of the grain Chenopodium quinoa Willd., an important economical and dietary plant', *Mitochondrial DNA Part B: Resources*, 6(1), pp. 40–42. Available at: https://doi.org/10.1080/23802359.2020.1845107.

Garrison, E. *et al.* (2022) 'A spectrum of free software tools for processing the VCF variant call
format: vcflib, bio-vcf, cyvcf2, hts-nim and slivar', *PLoS Computational Biology*, 18(5), pp. 1–
14. Available at: https://doi.org/10.1371/journal.pcbi.1009123.

Garrison, E. and Marth, G. (2012) 'Haplotype-based variant detection from short-read
sequencing', pp. 1–9. Available at: http://arxiv.org/abs/1207.3907.

Gorinstein, S. *et al.* (2002) 'Characterisation of pseudocereal and cereal proteins by protein and
amino acid analyses', *Journal of the Science of Food and Agriculture*, 82(8), pp. 886–891.
Available at: https://doi.org/10.1002/jsfa.1120.

Harris, S.A. and Ingram, R. (1991) 'Chloroplast DNA and biosystematics: The effects of
intraspecific diversity and plastid transmission', *TAXON*, 40(3), pp. 393–412. Available at:
https://doi.org/https://doi.org/10.2307/1223218.

Holt, C. and Yandell, M. (2011) 'MAKER2: An annotation pipeline and genome-database
management tool for second-generation genome projects', *BMC Bioinformatics*, 12(1), p. 491.
Available at: https://doi.org/10.1186/1471-2105-12-491.

Hong, S.Y. *et al.* (2017) 'Complete chloroplast genome sequences and comparative analysis of
Chenopodium quinoa and C. Album', *Frontiers in Plant Science*, 8(October), pp. 1–12.
Available at: https://doi.org/10.3389/fpls.2017.01696.

Hu, J. *et al.* (2020) 'NextPolish: A fast and efficient genome polishing tool for long-read
assembly', *Bioinformatics*, 36(7), pp. 2253–2255. Available at:
https://doi.org/10.1093/bioinformatics/btz891.

Jacobsen, S.E. (1997) 'Adaptation of quinoa (Chenopodium quinoa) to Northern European
agriculture: Studies on developmental pattern', *Euphytica*, 96(1), pp. 41–48. Available at:
https://doi.org/10.1023/A:1002992718009.

Jacobsen, S.E. (2003) 'The worldwide potential for quinoa (Chenopodium quinoa Willd.)', *Food
Reviews International*, 19(1–2), pp. 167–177. Available at: https://doi.org/10.1081/FRI-
120018883.

Jacobsen, S.E. (2011) 'The Situation for Quinoa and Its Production in Southern Bolivia: From
Economic Success to Environmental Disaster', *Journal of Agronomy and Crop Science*, 197(5),
pp. 390–399. Available at: https://doi.org/10.1111/j.1439-037X.2011.00475.x.

Jaggi, K.E. *et al.* (2024) 'A Pangenome Reveals LTR Repeat Dynamics as a Major Driver of
Genome Evolution in Chenopodium', *The Plant Genome* [Preprint].

Jarvis, D.E. *et al.* (2017) 'The genome of Chenopodium quinoa', *Nature*, 542(7641), pp. 307–
312. Available at: https://doi.org/10.1038/nature21370.

Jarvis, D.E. *et al.* (2022) 'Chromosome-Scale Genome Assembly of the Hexaploid Taiwanese
Goosefoot "Djulis" (Chenopodium formosanum)', *Genome Biology and Evolution*, 14(8), pp. 1–
7. Available at: https://doi.org/10.1093/gbe/evac120.

Jellen, E.N. *et al.* (2011) 'Chenopodium', in *Wild Crop Relatives: Genomic and Breeding
Resources*. Berlin, Heidelberg, Heidelberg: Springer Berlin Heidelberg, pp. 35–61. Available at:
https://doi.org/10.1007/978-3-642-14387-8_3.

Jellen, E.N. *et al.* (2013) 'Botany , Phylogeny and Evolution', in *State of the Art Report on*
Quinoa around the World. Rome: FAO & CIRAD, pp. 12–23.

Jin, J.J. *et al.* (2020) 'GetOrganelle: A fast and versatile toolkit for accurate de novo assembly of
organelle genomes', *Genome Biology*, 21(1), pp. 1–31. Available at:
https://doi.org/10.1186/s13059-020-02154-5.

Johnson, M. *et al.* (2008) 'NCBI BLAST: a better web interface.', *Nucleic acids research*,
36(Web Server issue), pp. 5–9. Available at: https://doi.org/10.1093/nar/gkn201.

Jones, P. *et al.* (2014) 'InterProScan 5: Genome-scale protein function classification',
Bioinformatics, 30(9), pp. 1236–1240. Available at:
https://doi.org/10.1093/bioinformatics/btu031.

Katoh, K. and Standley, D.M. (2013) 'MAFFT multiple sequence alignment software version 7:
Improvements in performance and usability', *Molecular Biology and Evolution*, 30(4), pp. 772–
780. Available at: https://doi.org/10.1093/molbev/mst010.

Kent, W.J. (2002) ' BLAT —The BLAST -Like Alignment Tool ', *Genome Research*, 12(4), pp.
656–664. Available at: https://doi.org/10.1101/gr.229202.

Kim, Y., Chung, Y. and Park, J. (2019) 'The complete chloroplast genome of Chenopodium
ficifolium Sm. (Amaranthaceae)', *Mitochondrial DNA Part B: Resources*, 4(1), pp. 872–873.
Available at: https://doi.org/10.1080/23802359.2019.1573122.

Kim, Y., Park, J. and Chung, Y. (2019) 'The complete chloroplast genome of Suaeda japonica
Makino (Amaranthaceae)', *Mitochondrial DNA Part B: Resources*, 4(1), pp. 1505–1507.
Available at: https://doi.org/10.1080/23802359.2019.1601039.

Kolano, B. *et al.* (2016) 'Molecular and cytogenetic evidence for an allotetraploid origin of
Chenopodium quinoa and C. berlandieri (Amaranthaceae)', *Molecular Phylogenetics and*
Evolution, 100, pp. 109–123. Available at: https://doi.org/10.1016/j.ympev.2016.04.009.

Koren, S. *et al.* (2017) 'Canu: Scalable and accurate long-read assembly via adaptive κ-mer
weighting and repeat separation', *Genome Research*, 27(5), pp. 722–736. Available at:
https://doi.org/10.1101/gr.215087.116.

Kozik, A. *et al.* (2019) 'The alternative reality of plant mitochondrial DNA: One ring does not
rule them all', *PLoS Genetics*, 15(8), pp. 1–30. Available at:
https://doi.org/10.1371/journal.pgen.1008373.

Krzywinski, M.I. *et al.* (2009) 'Circos: An information aesthetic for comparative genomics',
Genome Research [Preprint]. Available at: https://doi.org/10.1101/gr.092759.109.

Kumar, B. and Bhalothia, P. (2020) 'Orphan crops for future food security', *Journal of*
Biosciences, 45(1). Available at: https://doi.org/10.1007/s12038-020-00107-5.

Kuznetsov, D. *et al.* (2023) 'OrthoDB v11: annotation of orthologs in the widest sampling of
organismal diversity', *Nucleic Acids Research*, 51(1 D), pp. D445–D451. Available at:
https://doi.org/10.1093/nar/gkac998.

Langmead, B. and Salzberg, S.L. (2012) 'Fast gapped-read alignment with Bowtie 2', *Nature*

Methods, 9(4), pp. 357–359. Available at: https://doi.org/10.1038/nmeth.1923.

Lemmon, Z.H. *et al.* (2018) 'Rapid improvement of domestication traits in an orphan crop by
genome editing', *Nature Plants*, 4(10), pp. 766–770. Available at:
https://doi.org/10.1038/s41477-018-0259-x.

Levy Karin, E., Mirdita, M. and Söding, J. (2020) 'MetaEuk-sensitive, high-throughput gene
discovery, and annotation for large-scale eukaryotic metagenomics', *Microbiome*, 8(1), pp. 1–15.
Available at: https://doi.org/10.1186/s40168-020-00808-x.

Li, H. (2018) 'Minimap2: Pairwise alignment for nucleotide sequences', *Bioinformatics*, 34(18),
pp. 3094–3100. Available at: https://doi.org/10.1093/bioinformatics/bty191.

Li, H. (2023) 'Genome analysis Protein-to-genome alignment with miniprot', 39(January), pp.
1–6.

Li, H. and Durbin, R. (2009) 'Fast and accurate short read alignment with Burrows-Wheeler
transform', *Bioinformatics*, 25(14), pp. 1754–1760. Available at:
https://doi.org/10.1093/bioinformatics/btp324.

Lin, Y. *et al.* (2023) 'QuarTeT: A telomere-To-Telomere toolkit for gap-free genome assembly
and centromeric repeat identification', *Horticulture Research*, 10(8), pp. 1–7. Available at:
https://doi.org/10.1093/hr/uhad127.

Lindenbaum, P. (2015) 'JVarkit: java-based utilities for Bioinformatics', *Figshare*, pp. 2–5.
Available at: https://dx.doi.org/10.6084/m9.figshare.1425030.v1.

Mandák, B. *et al.* (2012) 'Is hybridization involved in the evolution of the Chenopodium album
aggregate? An analysis based on chromosome counts and genome size estimation', *Flora:*
Morphology, Distribution, Functional Ecology of Plants, 207(7), pp. 530–540. Available at:
https://doi.org/10.1016/j.flora.2012.03.010.

Mandák, B. *et al.* (2016) 'How genome size variation is linked with evolution within
Chenopodium sensu lato', *Perspectives in Plant Ecology, Evolution and Systematics*, 23, pp. 18–
32. Available at: https://doi.org/10.1016/j.ppees.2016.09.004.

Mangelson, H. *et al.* (2019) 'The genome of Chenopodium pallidicaule: An emerging Andean
super grain', *Applications in Plant Sciences*, 7(11), pp. 1–12. Available at:
https://doi.org/10.1002/aps3.11300.

Manni, M. *et al.* (2021) 'BUSCO Update: Novel and Streamlined Workflows along with Broader
and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes',
Molecular Biology and Evolution, 38(10), pp. 4647–4654. Available at:
https://doi.org/10.1093/molbev/msab199.

Maughan, P.J. *et al.* (2019) 'Mitochondrial and chloroplast genomes provide insights into the
evolutionary origins of quinoa (Chenopodium quinoa Willd.)', *Scientific Reports*, 9(1), pp. 1–11.
Available at: https://doi.org/10.1038/s41598-018-36693-6.

Maughan, P.J. *et al.* (2024) 'North American pitseed goosefoot (Chenopodium berlandieri) is a
genetic resource to improve Andean quinoa (C. quinoa)', *Scientific Reports*, 14(1), pp. 1–13.
Available at: https://doi.org/10.1038/s41598-024-63106-8.

McGrath, J.M. *et al.* (2023) 'A contiguous de novo genome assembly of sugar beet EL10 (Beta
vulgaris L.)', *DNA research : an international journal for rapid publication of reports on genes*
and genomes, 30(1), pp. 1–14. Available at: https://doi.org/10.1093/dnares/dsac033.

Mirarab, S., Nguyen, N. and Warnow, T. (2012) 'SEPP: SATé-enabled phylogenetic placement',
in *Biocomputing 2012*. World Scientific, pp. 247–258.

Neff, E. (2017) *Developing a Molecular Pipeline To Identify Chenopodium Species in New*
England. Master's thesis, University of New Hampshire. University of New Hampshire.

Neff, E., Sullivan, J.R. and Davis, T.M. (2018) 'Enhanced Documentation of Chenopodium
foggii (Amaranthaceae) in Northern New England', *Rhodora*, 120(983), pp. 257–259. Available
at: https://doi.org/10.3119/18-02.

Neph, S. *et al.* (2012) 'BEDOPS: High-performance genomic feature operations',
Bioinformatics, 28(14), pp. 1919–1920. Available at:
https://doi.org/10.1093/bioinformatics/bts277.

Nolen, H. *et al.* (2022) 'Evaluation of Disease Severity and Molecular Relationships Between
Peronospora variabilis Isolates on Chenopodium Species in New Hampshire', *Plant Disease*,
106(2), pp. 564–571. Available at: https://doi.org/10.1094/PDIS-06-21-1150-RE.

Paysan-Lafosse, T. *et al.* (2023) 'InterPro in 2022', *Nucleic Acids Research*, 51(D1), pp. D418–
D427. Available at: https://doi.org/10.1093/nar/gkac993.

Rey, E. *et al.* (2023) 'A chromosome-scale assembly of the quinoa genome provides insights into
the structure and dynamics of its subgenomes', *Communications Biology*, 6(1), pp. 1–14.
Available at: https://doi.org/10.1038/s42003-023-05613-4.

Richard, G. (2019) 'gtrichard/deepStats: deepStats 0.3. 1 (Version 0.3. 1)', *Zenodo. doi*, 10.

Rojas, W. *et al.* (2009) 'From neglect to limelight: issues, methods and approaches in enhancing
sustainable conservation and use of Andean grains in Peru and Bolivia', *Journal of Agriculture*
and Rural Development in the Tropics and Subtropics, 92, p. 87.

Samuels, M.E. *et al.* (2023) 'Genomic Sequence of Canadian Chenopodium berlandieri: A North
American Wild Relative of Quinoa', *Plants*, 12(3), pp. 1–23. Available at:
https://doi.org/10.3390/plants12030467.

Shen, W. *et al.* (2016) 'SeqKit: A cross-platform and ultrafast toolkit for FASTA/Q file
manipulation', *PLoS ONE*, 11(10), pp. 1–10. Available at:
https://doi.org/10.1371/journal.pone.0163962.

Stamatakis, A. (2014) 'RAxML version 8: A tool for phylogenetic analysis and post-analysis of
large phylogenies', *Bioinformatics*, 30(9), pp. 1312–1313. Available at:
https://doi.org/10.1093/bioinformatics/btu033.

Stanke, M. *et al.* (2008) 'Using native and syntenically mapped cDNA alignments to improve de
novo gene finding', *Bioinformatics*, 24(5), pp. 637–644. Available at:
https://doi.org/10.1093/bioinformatics/btn013.

Štorchová, H. *et al.* (2015) 'The introns in FLOWERING LOCUS T-LIKE (FTL) genes are

useful markers for tracking paternity in tetraploid Chenopodium quinoa Willd', *Genetic*
Resources and Crop Evolution, 62(6), pp. 913–925. Available at:
https://doi.org/10.1007/s10722-014-0200-8.

Štorchová, H. *et al.* (2019) 'Chenopodium ficifolium flowers under long days without
upregulation of FLOWERING LOCUS T (FT) homologs', *Planta* [Preprint]. Available at:
https://doi.org/10.1007/s00425-019-03285-1.

Storer, J. *et al.* (2021) 'The Dfam community resource of transposable element families,
sequence models, and genome annotations', *Mobile DNA*, 12(1), p. 2. Available at:
https://doi.org/10.1186/s13100-020-00230-y.

Subedi, M. (2020) *Developing Chenopodium ficifolium as a diploid model system relevant to*
genetic characterization and improvement of allotetraploid C. quinoa Master's Theses and
Capstones. 1407. Available at: https://scholars.unh.edu/thesis/1407.

Subedi, M., Neff, E. and Davis, T.M. (2021) 'Developing Chenopodium ficifolium as a potential
B genome diploid model system for genetic characterization and improvement of allotetraploid
quinoa (Chenopodium quinoa)', *BMC Plant Biology*, 21(1), pp. 1–18. Available at:
https://doi.org/10.1186/s12870-021-03270-5.

Supek, F. *et al.* (2011) 'Revigo summarizes and visualizes long lists of gene ontology terms',
PLoS ONE, 6(7). Available at: https://doi.org/10.1371/journal.pone.0021800.

Tange, O. (2018) *GNU Parallel 2018*, *Isbn 9781387509881*. Available at: http://ole.tange.dk.

Tarailo-graovac, M. and Chen, N. (2009) 'Using RepeatMasker to Identify Repetitive Elements
in Genomic Sequences', (March), pp. 1–14. Available at:
https://doi.org/10.1002/0471250953.bi0410s25.

Testolin, R. and Cipriani, G. (1997) 'Paternal inheritance of chloroplast DNA and maternal
inheritance of mitochondrial DNA in the genus Actinidia', *Theoretical and Applied Genetics*,
94(6–7), pp. 897–903. Available at: https://doi.org/10.1007/s001220050493.

Thorvaldsdóttir, H., Robinson, J.T. and Mesirov, J.P. (2013) 'Integrative Genomics Viewer
(IGV): High-performance genomics data visualization and exploration', *Briefings in*
Bioinformatics, 14(2), pp. 178–192. Available at: https://doi.org/10.1093/bib/bbs017.

Tillich, M. *et al.* (2017) 'GeSeq - Versatile and accurate annotation of organelle genomes',
Nucleic Acids Research, 45(W1), pp. W6–W11. Available at:
https://doi.org/10.1093/nar/gkx391.

Torres, A.M., Weeden, N.F. and Martín, A. (1993) 'Linkage among isozyme, RFLP and RAPD
markers in Vicia faba', *Theoretical and Applied Genetics*, 85(8), pp. 937–945. Available at:
https://doi.org/10.1007/BF00215032.

Vaillancourt, B. and Buell, C.R. (2019) 'High molecular weight DNA isolation method from
diverse plant species for use with Oxford Nanopore sequencing', *bioRxiv*, p. 783159. Available
at: https://doi.org/10.1101/783159.

Vega-Gálvez, A. *et al.* (2010) 'Nutrition facts and functional potential of quinoa (Chenopodium
quinoa willd.), an ancient Andean grain: A review', *Journal of the Science of Food and*

Agriculture, 90(15), pp. 2541–2547. Available at: https://doi.org/10.1002/jsfa.4158.

Vogt, F., Shirsekar, G. and Weigel, D. (2022) 'vcf2gwas: Python API for comprehensive GWAS
analysis using GEMMA', *Bioinformatics*, 38(3), pp. 839–840. Available at:
https://doi.org/10.1093/bioinformatics/btab710.

Walsh, B.M. *et al.* (2015) ' Chenopodium polyploidy inferences from Salt Overly Sensitive 1 (
SOS1) data ', *American Journal of Botany*, 102(4), pp. 533–543. Available at:
https://doi.org/10.3732/ajb.1400344.

Wang, W., Lanfear, R. and Gaut, B. (2019) 'Long-Reads Reveal That the Chloroplast Genome
Exists in Two Distinct Versions in Most Plants', *Genome Biology and Evolution*, 11(12), pp.
3372–3381. Available at: https://doi.org/10.1093/gbe/evz256.

Wang, Y. *et al.* (2012) 'MCScanX: A toolkit for detection and evolutionary analysis of gene
synteny and collinearity', *Nucleic Acids Research*, 40(7), pp. 1–14. Available at:
https://doi.org/10.1093/nar/gkr1293.

Wang, Y. *et al.* (2024) 'Detection of colinear blocks and synteny and evolutionary analyses
based on utilization of MCScanX', *Nature Protocols*, 19(7), pp. 2206–2229. Available at:
https://doi.org/10.1038/s41596-024-00968-2.

Ward, S.M. (1998) 'A new source of restorable cytoplasmic male sterility in quinoa', *Euphytica*,
101(2), pp. 157–163. Available at: https://doi.org/10.1023/A:1018396808387.

Warwick, S.I. and Black, L. (1980) 'Uniparental Inheritance of Atrazine Resistance in
Chenopodium album', *Canadian Journal of Plant Science*, 60(2), pp. 751–753. Available at:
https://doi.org/10.4141/cjps80-108.

Wick, R.R. *et al.* (2015) 'Bandage: Interactive visualization of de novo genome assemblies',
Bioinformatics, 31(20), pp. 3350–3352. Available at:
https://doi.org/10.1093/bioinformatics/btv383.

Young, L.A. *et al.* (2023) 'A chromosome-scale reference of Chenopodium watsonii helps
elucidate relationships within the North American A-genome Chenopodium species and with
quinoa', *Plant Genome*, 16(3): e20(April), pp. 1–17. Available at:
https://doi.org/10.1002/tpg2.20349.

Zou, C. *et al.* (2017) 'A high-quality genome assembly of quinoa provides insights into the
molecular basis of salt bladder-based salinity tolerance and the exceptional nutritional value',
Cell Research, 27(11), pp. 1327–1340. Available at: https://doi.org/10.1038/cr.2017.124.

1026

1027

1028

1029

1030

1031

1032

Supplemental Tables

1033

1034 Supplemental Table 1. Individuals included in GWAS and phenotypic data used in analyses.

1035

Individual	Flowering Time	Height	No. Branches	Branch Angle	Internode Length
4	23	46.5	13	71.9	3.6
5	25	38.1	17	68.2	2.2
6	32	64.8	27	61.6	2.4
7	25	43.4	14	69.4	3.1
8	21	36.6	15	58.9	2.4
9	23	46.7	13	63.3	3.6
10	27	52.1	14	68.8	3.7
11	35	72.4	26	78.2	2.8
12	23	55.6	19	75.4	2.9
13	21	45.7	15	79.5	3
14	39	74.9	27	75.5	2.8
15	23	35.1	12	63.3	2.9
16	21	36.3	14	72.8	2.6
17	23	50	16	66.6	3.1
18	21	30.5	15	64.4	2
9r2	18	54.7	19	71.5	2.9
10r2	20	50.7	17	71.9	3
11r2	22	60.6	23	73.5	2.6
12r2	22	40.5	18	76	2.3
13r2	22	59	19	72	3.1
14r2	29	72.8	29	63	2.5

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046 Supplemental Table 2. Centromere positions and lengths as identified by quarTeT CentroMiner
1047 and telomere lengths and directions as identified by quarTeT TeloExplorer.

1048

ID	Centromere			Telomere				
	Start	End	Length	Present	Upstream Length	Dir.	Downstream Length	Dir.
Cf1	58004969	58247898	242930	left	3668	+	0	
Cf2	79538569	79657117	118549	right	0		3941	-
Cf3	49844598	51718346	1873749	both	9779	+	3549	-
Cf4	39049495	40750452	1700958	both	20412	+	5019	-
Cf5	41797451	42801417	1003967	both	17843	+	3731	-
Cf6	6996291	7223985	227695	both	5397	+	4382	-
Cf7	38832653	41378639	2545987	both	882	-	5054	-
Cf8	34002546	34566367	563822	both	4480	+	5243	-
Cf9	46653552	46799075	145524	both	3710	+	4753	-

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068 Supplemental Table 3. Positions and alleles of SNPs found in mitochondrial and chloroplast
1069 genomes used to establish maternal cytoplasmic inheritance.

1070

Genome	Position	Reference	Alternate
Mitochondria	45844	AAAAGAATTCATTAAGAATTCACA	AA
	73300	A	C
	107870	A	C
	112616	A	C
	112987	A	C
	149592	C	T
	184591	T	A
	192086	G	T
	229731	C	A
Chloroplast	2238	T	G
Path 2	3171	C	A
	12101	C	A
	28316	CTTTTTTTTCTTTGA	CTTTTTTTCTTTGA
	30112	G	C
	30993	C	G
	33109	T	G
	42745	G	T
	49677	T	G
	50161	C	A
	63570	CAAAAAAGAAA	CAAAAAAAAAA
	64005	C	T
	67316	T	C
	81756	ATTTTTTTTTTGTT	ATTTTTTTTTGTT
	108762	G	T
	115629	A	C
	116544	A	T
	118756	C	A
	126735	C	A

1071