Measuring the Strength of Spatial Sorting

Nikunj Goel*[§][1,2], Mattheau S. Comerford*[3], Anastasia Bernat[4], Scott P. Egan[5], Thomas E. Juenger[1], and Timothy H. Keitt[1]

[1]Department of Integrative Biology, The University of Texas at Austin, Austin, Texas, 78712, USA

[2]Department of Statistics and Data Sciences, The University of Texas at Austin, Austin, Texas, 78712, USA

[3]Boston Department of Biology, University of Massachusetts, Boston, Massachusetts, 02125, USA

[4] Pacific Northwest National Laboratory, Richland, WA 99352, USA

[5]Department of BioScience, Rice University, Houston, Texas, 77005, USA

* These authors contributed equally to the work

[§]Corresponding author; email: nikunj410@gmail.com

Open research statement: The data and code has been deposited at Zenodo:

https://zenodo.org/records/15725572.

Abstract

Empirical work suggests that spatial sorting—a mechanism of evolutionary change fueled

by phenotype-dependent dispersal—may lead to phenotypic shifts on ecological timescales.

However, we currently lack theoretical tools to measure the strength of spatial sorting, as we do to

measure the strength of natural selection. To address this gap, we present a quantitative genetics

model and identify an evolutionary parameter in the model to measure the strength of spatial

sorting. This parameter, the standardized sorting gradient, is structurally akin to the standardized

selection gradient, commonly used to measure the strength of natural selection. To show the utility

of our approach, we analyzed wing-morphology data of soapberry bugs (*Jadera haematoloma*)

recolonizing flooded habitats extirpated by a hurricane. We found that the estimated strength of

phenotypic shift ranked in the top ten percentile of documented phenotypic shifts in the scientific

literature. Our results underscore that, like natural selection, spatial sorting, too, can yield rapid

evolution after extreme events.

Introduction

Extreme climatic events are becoming common due to rising carbon emissions and concomitant planetary warming (Solow 2015, Stott 2016). These climatic perturbations may impose strong selection on populations, fueling rapid evolution (Grant et al. 2017). Indeed, many studies have documented strong selection in response to extreme events (Bumpus 1899, Boag and Grant 1981, Grant and Grant 1993, Brown and Brown 1998, Donihue et al. 2018). These studies underscore a common theme—an extreme event yields non-random mortality wherein some individuals live past the event because of a genetically determined trait that confers a survival advantage. When these survivors reproduce, the mean phenotype of the population changes, resulting in evolution by natural selection (Darwin and Wallace 1858).

Recent empirical work suggests that extreme events can also produce drastic phenotypic changes by a relatively lesser known evolutionary mechanism that relies on non-random dispersal (Comerford et al. 2023). Extreme events can extirpate some local populations in a patchy landscape, creating empty patches (van Bergen et al. 2020). These local extinctions create an opportunity for non-random immigration from nearby extant populations wherein a subset of individuals colonize the empty patches because of a genetically determined trait that confers a dispersal advantage. When these immigrants with traits that confer dispersal advantage mate with each other, the mean phenotype in hitherto empty patches changes relative to the nearby extant populations. This mechanism of evolution driven by phenotype-dependent dispersal is referred to as spatial sorting (Shine et al. 2011).

Although a growing number of studies have documented evolution by spatial sorting in both natural (MacKay and Lamb 1979, Peroni 1994, Hill et al. 1999, Hanski et al. 2004, Phillips et al. 2006, Duckworth 2008, Forsman et al. 2011, Lombaert et al. 2014) and experimental settings

(Williams et al. 2016, Ochocki and Miller 2017, Szűcs et al. 2017, Weiss-Lehman et al. 2017, Usui

and Angert 2024), we lack theoretical tools to measure the strength of spatial sorting from

phenotypic data, especially after an extreme event. This gap in the literature stems from two

reasons. First, extreme events are hard to predict, making it challenging to plan longitudinal studies

to quantify their evolutionary impacts (Grant et al. 2017). Second, the theory of spatial sorting is

still in its infancy (Bolnick and Otto 2013, Novak and Kollár 2017, Phillips and Perkins 2019,

Peischl and Gilbert 2020). Unlike the Breeder's equation (Lush 1937) and its extensions (Falconer

1960, Lande and Arnold 1983, Walsh and Lynch 2018) that provide quantitative tools to measure

the strength of selection using standardized selection gradients, we do not have similar quantitative

genetics theory that may allow biologists to study phenotypic evolution by spatial sorting.

Addressing this knowledge gap is crucial for providing quantitative evidence that spatial sorting

can be strong enough to influence ecology, producing novel spatial eco-evolutionary dynamics

(Phillips et al. 2006, Ochocki and Miller 2017, Weiss-Lehman et al. 2017, Miller et al. 2020).

This paper has two major goals. First, we derive a quantitative genetics model of spatial

sorting to quantify evolutionary change during colonization of empty habitats. This model is

analogous to the Breeder's equation and serves several functions, such as facilitating a conceptual

understanding of how spatial sorting drives the evolution of quantitative traits and providing a

standardized measure to quantify the strength of spatial sorting, which we refer to as standardized

sorting gradient. Based on the knowledge gained from the model, biologists can plan what data to

collect, how to analyze it, and how to interpret the results in the broader context of evolutionary

theory. Second, to demonstrate the utility of the quantitative genetics model, we analyze wing

morphology data of red-shoulder soapberry bugs (*Jadera haematoloma*; hereafter soapberry bug)

from a multi-year monitoring experiment (2016-2020) in the Greater Houston area that overlapped

with Hurricane Harvey in 2017 (Comerford et al. 2023). Specifically, using the survey data, we

estimate the standardized sorting gradient to quantify the strength of spatial sorting due to the

phenotype-dependent recolonization of habitats extirpated by the hurricane. We also perform

experimental crosses to evaluate if the wing morphology has a genetic basis and to test if the non-

random recolonization could have resulted in an evolutionary response.

Quantitative genetics model of spatial sorting

Consider a patchy landscape consisting of randomly distributed vacant and occupied

patches. The occupied patches are collectively inhabited by a large population of N parents. We

denote the trait value of the ith parent as ϕ_i, which determines individuals' dispersal capacity and

lifetime reproductive success. We denote the mean trait value of the parents by $\bar{\phi}$. The bar is a

shorthand for the mean. Next, some parents disperse from occupied patches to vacant patches. To

track the dispersing parents, we define an indicator variable V_i, which is one if the ith parent

disperses to a vacant patch and zero otherwise. Following dispersal, the ith parent produces W_i

offspring, such that the mean trait value of offspring, ϕ_i^o, is δ_i more than the trait value of the

parent.

Based on these notations, the mean trait value of offspring in the vacant patches is

$$\bar{\phi}' = \frac{\sum_{i=1}^N V_i W_i (\phi_i + \delta_i)}{\sum_{i=1}^N V_i W_i}. \tag{1}$$

98   This equation is obtained by adding the trait value of offspring in the vacant patches and then

99   dividing this sum by the number of offspring in the vacant patches. We can re-express the above

100 equation compactly as

$$\bar{\phi}' = \frac{\overline{VW\phi} + \overline{VW\delta}}{\overline{VW}}$$

$$= \frac{\mathbb{C}\text{ov}(VW, \phi) + \overline{VW\delta}}{\overline{VW}} + \bar{\phi}. \tag{2}$$

101   By re-arranging the terms in equation (2), we find that the evolutionary change due to spatial

102   sorting ($\Delta\bar{\phi} = \bar{\phi}' - \bar{\phi}$; here defined as, the mean trait value of offspring in the vacant patches

103   minus the mean trait value of parents in the occupied patches before dispersal) is given by

$$\Delta\bar{\phi} = \frac{1}{\overline{VW}}\left[\mathbb{C}\text{ov}(VW, \phi) + \overline{VW\delta}\right]$$

$$= \frac{1}{\overline{VW}}\,\mathbb{C}\text{ov}(VW, \phi^o) + \bar{\delta} \tag{3}$$

104   Assuming that the joint distribution of parent and offspring trait values is a multivariate

105   normal distribution, the mean trait value of offspring of the ith parent can be expressed as

$$\phi_i^o = \bar{\phi} + \beta_{\phi^o,\phi}(\phi_i - \bar{\phi}), \tag{4}$$

106   where $\beta_{\phi^o,\phi}$ is the slope of the regression between mean offspring (ϕ_i^o) and parents'(ϕ_i) trait

107   values (Rice 2004). This regression coefficient, $\beta_{\phi^o,\phi}$, is commonly referred to as heritability (h^2)

108   of the trait and is equal to the ratio of the additive genetic variance (σ_A^2) and phenotypic variance

109   (σ_ϕ^2) in the parent population (Falconer 1960, Walsh and Lynch 2018). Substituting equation (4)

110   in equation (3) yields an evolutionary change of

$$\Delta\bar{\phi} = \frac{1}{\overline{VW}}\,\beta_{\phi^o,\phi}\,\mathbb{C}\text{ov}(VW, \phi)$$

$$= h^2 s_{vw} \tag{5}$$

111   where $h^2 = \beta_{\phi^o,\phi} = \sigma_A^2/\sigma_\phi^2$, $s_{vw} = \overline{VW}^{-1}\,\mathbb{C}\text{ov}(VW, \phi)$, and $\bar{\delta} = 0$ (See Appendix for details).

112   Alternatively, using the identity for covariance, equation (5) can be re-expressed as

$$\Delta\bar{\phi} = \beta_{\phi^o,\phi}\,\beta_{vw,\phi}\,\sigma_\phi^2$$

$$= \sigma_A^2 \beta_{vw,\phi}, \tag{6}$$

where $\beta_{vw,\phi}\left(= \sigma_{\phi}^{-2}s_{vw}\right)$ is the slope of the regression line between $V_iW_i/\overline{VW}$ and ϕ_i. Intuitively,

the equation (6) suggests that the magnitude of evolutionary change due to spatial sorting is

governed by the heritable variation in the trait and the strength of the statistical association between

the number of offspring a parent leaves in the vacant patches and its trait value. This association

arises because parents with traits that confer them a dispersal or reproductive advantage are more

likely to leave offspring in the vacant patches.

Note that the equation of evolution by spatial sorting (Eq. 5) is structurally similar to the

Breeder's equation (Phillips and Perkins 2019). Consequently, we adopt interpretation and the

naming convention of variables in equations (5) and (6) that are analogs to their counterpart

variables in the Breeder's equation. We refer to s_{vw} as the *sorting differential*—phenotypic shifts

in parents due to spatial sorting, $\beta_{vw,\phi}$ as the *sorting gradient*, V_iW_i—the number of offspring the

ith parent leaves in the vacant patches—as the *sorting fitness* of i, $\overline{VW}$ as the *mean population*

sorting fitness, and $V_iW_i/\overline{VW}$ as the *relative sorting fitness* of i.

More importantly, the similarities between the two models offer insights into what measure

to use to quantify the strength of spatial sorting. We propose that the strength of spatial sorting

should be measured using the sorting gradient of the standardized phenotypic trait, which we refer

to as the *standardized sorting gradient*—$\beta'_{vw,\phi} = \sigma_{\phi}\beta_{vw,\phi}$. In the case of a single phenotypic trait,

$\beta'_{vw,\phi} = \sigma_{\phi}\beta_{vw,\phi} = \sigma_{\phi}^{-1}s_{vw} = s'_{vw}$, where s'_{vw} is the *standardized sorting differential*. Using the

standardized sorting gradient to quantify the strength of spatial sorting offers several advantages.

First, the sorting gradient is closely connected to the equation of evolutionary change by spatial

sorting (Eq. 5) and, therefore, has a predictive value in the context of quantifying evolutionary

changes in phenotypic traits. Second, standardizing phenotypic traits provides a measure of the

strength of spatial sorting, which can be compared across biological systems by putting traits on a

common scale (Matsumura et al. 2012). Third, due to the structural congruence between the

equations of evolution by natural selection and spatial sorting, standardized sorting gradients can

be compared with standardized selection gradients to evaluate the relative magnitude of the two

evolutionary forces in generating phenotypic shifts.

Data Analysis

To show the practical utility of our model in quantifying the strength of spatial sorting and

corresponding evolutionary change, we analyze an exemplar of spatial sorting—survey data on

soapberry bugs recolonizing habitats extirpated by a hurricane (Comerford et al. 2023).

Soapberry Bugs

The soapberry bugs' ancestral range in the USA falls along the southern U.S. border with

Mexico and the states abutting the Gulf Coast (Tsai et al. 2013). In humid tropical regions,

including the Greater Houston area, soapberry bugs breed throughout the year. Upon mating, the

females lay eggs in 1 cm deep holes beneath their host plants of the soapberry family (Carroll

1988). It takes about 10 days before the larval nymphs emerge from the eggs. Subsequently, the

nymphs undergo five molting stages (also referred to as instars) over the next 29 ± 4 days before

reaching sexual maturity. These newly molted adult females avoid mating for 6 ± 1 days, after

which they lay eggs throughout most of their 60-day lifespan (Carroll 1991) (see Fig. S1).

Female soapberry bugs generally tend to begin histolysis of flight muscles during

oogenesis (Carroll et al. 2003), and males show prolonged mate guarding behavior (Carroll 1988).

As a result, the potential movement of females between habitats after mating is greatly reduced

(Comerford et al. 2023). These insects have binary wing morphologies—long-winged

'macropterous' and short-winged 'brachypterous' forms (Fig. 1A) (Dingle and Winchell 1997).

These binary forms are strongly associated with dispersal abilities of both male and female

insects—macropterous insects have fully developed wings that allow them to fly long distances,

while flightless brachypterous insects lack flight muscles. Previous studies suggest that this

discrete wing polyphenism is determined by juvenile hormones, which, in turn, are regulated by

many genes with small effects, in addition to resource availability in the environment (Dingle and

Winchell 1997, Dingle 2001).

Hurricane Harvey

Warm sea surface temperatures are perfect incubators for tropical storms in the Gulf of

Mexico (Chen et al. 2008, Risser and Wehner 2017). On 25[th] August 2017, Hurricane Harvey

struck southeastern Texas, releasing in excess of 1500 mm of precipitation in a span of seven days

(Trenberth et al. 2018). This one in ten-thousand-year event led to intense flooding across the

Greater Houston area (Emanuel 2017, Van Oldenborgh et al. 2017). However, not all regions were

affected equally. Variation in ground porosity and subsidence led to prolonged flooding in some

areas while others remained unaffected (Miller and Shirzaei 2019). Fortuitously, the hurricane

occurred during a multi-year population survey by Comerford et al. (2023), which was initiated a

year before the hurricane and continued for an additional three years.

Monitoring Scheme

Comerford et al. (2023) noted the insect abundances and the morphological form of

soapberry bugs every 2 weeks at fifteen sites across the region (Fig. 1A). The survey data revealed

two important observations. First, four of the fifteen sites that did not experience prolonged

flooding (hereafter called unflooded sites) maintained a stable population throughout the survey

(blue points in Fig. 1A). The remaining eleven flooded sites (pink points in Fig. 1A) experienced

a local extinction of insects (Fig. 1 B and C). Extinction was confirmed by a gap of 4.5 to 27

months gap in insect detection at the flooded sites, exceeding the ~100 days life expectancy of

soapberry bugs (from the egg stage to the death of an insect). In addition, inundation experiments

by Comerford et al. (2023) showed that after a week of submergence, egg survival drops

precipitously and is zero after ten days. This provided further corroborative evidence for local

extinction because females lay eggs in the soil beneath the host plants (Carroll 1988), and the

topsoil at the flooded sites was inundated for 7 to 90+ days after the hurricane (Fig. 1C). Second,

before the hurricane, approximately two thirds of sampled male and female individuals in the

unflooded sites were macropterous insects (Fig 2A). In contrast, during the six weeks after the

hurricane, starting from the first co-detection of male and female insects, the survey showed that

the flooded sites were re-established only by macropterous male and female insects (a total of 88

individuals were sampled). We chose this specific sampling time period because it takes about six

weeks for an egg to develop into a penultimate nymph (the last stage before the insect reaches

reproductive maturity). During this development period, the instars can be visually distinguished

from adults, which allows us to avoid individuals who were born in the flooded sites after

recolonization (see Fig. S1). These survey observations are consistent with the theory of spatial

sorting—since macropterous insects are better dispersers than flightless brachypterous insects,

insects recolonizing the flooded sites have a higher proportion of macropterous forms relative to

the insect population in the unflooded sites before recolonization.

203   *Strength of Spatial Sorting*

To calculate the strength of spatial sorting, we assume the visible morphological form of

an insect has a threshold response to an underlying unobserved continuous trait, commonly

referred to as liability (Falconer 1960), such that when the liability of an insect is above (below)

zero, the individual takes the macropterous (brachypterous) form (Fig. 2B; see Roff (1996) for

discussion on thereshold models to describe evolution of disrcrete traits that are regulated by an

underlying continious character). Based on this assumption, the proportion of macropterous insects

in a population is simply the area under the liability distribution from zero to infinity. The variation

in the liability is assumed to be determined by environmental and polygenic factors and can be

thought of as the concentration of some substance, such as the juvenile hormones (Dingle and

Winchell 1997, Dingle 2001).

Based on our derivation, the standardized sorting gradient can be expressed as

$$\beta'_{vw,\phi} = \sigma_\phi^{-1} s_{vw}$$

$$= \sigma_\phi^{-1} \overline{VW}^{-1} \, \mathbb{cov}(VW, \phi) \tag{7}$$

$$= \sigma_\phi^{-1} \frac{\overline{VW\phi}}{\overline{VW}} - \sigma_\phi^{-1} \bar{\phi}$$

215   Because the survey data revealed that only macropterous insects participate in the recolonization

216   of flooded sites, we set the sorting fitness of brachypterous insects to zero and assume that the

217   sorting fitness of macropterous insects is a non-negative fixed value. This allowed us to simplify

218   equation (7) as

$$\beta'_{vw,\phi} = \sigma_\phi^{-1} (\bar{\phi}_m - \bar{\phi}), \tag{8}$$

219   where $\bar{\phi}$ and $\bar{\phi}_m$ are the mean liabilities of all insects and macropterous insects in the unflooded

220   sites before recolonization, respectively (see Appendix for details).

Next, assuming the liability of insects has a normal distribution (Fig. 2B), the mean liability of insects in the unflooded sites (in units of σ_ϕ) is

$$\bar{\phi}\sigma_\phi^{-1} = \sqrt{2}\,\mathrm{Erf}^{-1}(2p-1) \tag{9}$$

and the mean liability of macropterous insects in the unflooded sites (in units of σ_ϕ) is

$$\bar{\phi}_m\sigma_\phi^{-1} = \bar{\phi}\sigma_\phi^{-1} + \sqrt{2\pi p^2}^{-1}\, e^{-\frac{1}{2}\left(\frac{\bar{\phi}}{\sigma_\phi}\right)^2}, \tag{10}$$

where Erf^{-1} is the inverse error function and p is the proportion of macropterous insects in the unflooded sites before recolonization. Intuitively, $\bar{\phi}_m\sigma_\phi^{-1}$ corresponds to the mean of the liability distribution truncated from below at zero (Fig. 2B). By rearranging the terms in equations (8-10), we can show that the standardized sorting gradient is given by

$$\beta'_{vw,\phi} = \sqrt{2\pi p^2}^{-1}\, e^{-\left[\mathrm{Erf}^{-1}(2p-1)\right]^2}. \tag{11}$$

To estimate $\beta'_{vw,\phi}$, we took six weeks of pre-hurricane data from unflooded sites (Comerford et al. 2023). The survey revealed that 180 of the 280 sampled individuals at the unflooded sites had macropterous forms. Since the insects were sampled randomly, we statistically modeled the proportion of macropterous insects at the unflooded sites (p) as

$$180 \sim \mathrm{Binomial}(280, p). \tag{12}$$

Using the posterior samples of p as input in equation (11), we obtained the posterior distribution of $\beta'_{vw,\phi}$ (Carpenter et al. 2017). We found $p = 0.64 \pm 0.03$ and $\beta'_{vw,\phi} = s'_{vw} = 0.58 \pm 0.05$. Here, the first and second terms in the estimate of p and $\beta'_{vw,\phi}$ correspond to the mean and standard deviation of the posterior distribution, respectively (see supplementary code for statistical analysis). When we compared the magnitude of the standardized sorting gradient to the magnitude of standardized selection gradients documented in the scientific literature, it ranked in the top ten percentile among ~1400 previous studies (Kingsolver et al. 2001).

Heritability and Evolutionary Response to Spatial Sorting

To test if spatial sorting could have resulted in an evolutionary change, we performed

multigenerational crosses to estimate the heritability (see Supplementary for details). This

multigenerational breeding experiment was conducted under standardized laboratory conditions,

which allowed us to control for plasticity and maternal effects.

We collected 40 juvenile insects from three geographical locations (Austin, Stock Island,

and Gulfport) separated by more than 350 km. The insects were reared under standardized

laboratory conditions and randomly mated to generate 28 males and 28 females of the F1

generation of insects. Upon maturation, the morphological form of F1 insects was noted: we used

the numerical value one (zero) to represent macropterous (brachypterous) insects. Subsequently,

the insects were randomly paired to generate an F2 generation of insects. Upon eclosion from eggs,

14 F2 nymphs were selected haphazardly from each maternal line. The insects were reared under

the same conditions as F1 insects, and their morphological form was noted upon maturation.

To estimate heritability, we fit a Bayesian animal model (De Villemereuil et al. 2016) with

the morphology of F1 and F2 insects as input for parents and offspring forms, respectively. We

found a statistically significant estimate of heritability of liability ($h^2 = 0.65 \pm 0.13$) (Hadfield

2010), suggesting that wing morphology is highly heritable.

Finally, we used the posterior samples of h^2 and s'_{vw} to calculate the posterior distribution

of standardized evolutionary change ($\Delta\bar{\phi}' = \Delta\bar{\phi}\sigma_\phi^{-1} = h^2 s'_{vw}$). We find that non-random

recolonization of the flooded sites would have resulted in a standardized evolutionary change

($\Delta\bar{\phi}'$) of 0.37 ± 0.08 in insect liability. The large magnitude of standardized evolutionary change

suggests that evolution by spatial sorting may exert a strong influence on the spatial population

dynamics of a species.

264   **Discussion**

We present a quantitative genetics model of spatial sorting to describe evolution in patchy

landscape (Eq. 5). The model suggests that the evolutionary change by spatial sorting can be

partitioned into two components, sorting gradient and additive genetic variance, which can be

estimated independently using dispersal traits of organisms measured during a recolonization event

and an animal model with pedigree data as input, respectively. We propose that, analogous to the

Breeder's equation, the sorting gradient of standardized traits (i.e., $\beta'_{vw,\phi}$) can be used to quantify

the strength of spatial sorting. This choice offers two advantages. First, putting traits on a common

scale (*i.e.*, in units of standardized deviation) allows evolutionary biologists to compare the

strength of spatial sorting across biological systems (Matsumura et al. 2012). Second, biologists

can juxtapose standardized sorting and selection gradients to assess the relative magnitude of

phenotypic shifts arising from spatial sorting and natural selection.

We applied our quantitative genetics model of spatial sorting to study the evolution of

soapberry bugs' liability because of morphology-dependent recolonization of habitats extirpated

by Hurricane Harvey in Greater Houston (Fig. 1; Comerford et al. 2023). We found a statistically

significant standardized sorting gradient ($\beta'_{vw,\phi}$) of 0.58 ± 0.05 (Fig. 2), which ranked in the top

ten percentile of standardized selection gradients documented in the scientific literature

(Kingsolver et al. 2001). Genetic crosses revealed that insect liability is a heritable trait ($h^2 = 0.65$

$\pm\ 0.13$), yielding a standardized evolutionary change in liability ($\Delta\bar{\phi}'$) of 0.37 ± 0.08 due to spatial

sorting.

The estimate of the standardized sorting gradient suggests that non-random colonization of

habitats can produce phenotypic shifts comparable to the most extreme observed phenotypic shifts

in natural populations (Kingsolver et al. 2001). And, since many natural populations have abundant

heritable variation in dispersal-associated traits (Saastamoinen et al. 2018), spatial sorting may

yield evolutionary changes on ecological timescales, which can, in turn, produce novel eco-

evolutionary feedback (Miller et al. 2020).

Because of the structural similarities between the Breeder's equation and the quantitative

genetics model presented in the paper, our model can be extended to incorporate additional

features, such as genetic correlations between multiple traits (Lande and Arnold 1983) arising from

pleiotropy and linkage. These correlations can dramatically change inferences about the direction

and the magnitude of spatial sorting obtained from univariate analysis. For example, conducting

separate univariate analyses on two strongly correlated traits with the same signed phenotypic

shifts will yield the same signed sorting gradients. However, a joint multivariate analysis may

reveal that the traits may have opposite-signed sorting gradients, even though the observed

phenotypic shifts after spatial sorting have the same sign. This situation can arise when the indirect

response to spatial sorting due to genetic correlations on one of the traits is much larger in

magnitude than the direct response.

Genetic correlations can also be important when there is a sex-specific response to spatial

sorting (Lande 1980, McGlothlin et al. 2019). For example, when mating precedes dispersal, the

probability of leaving offspring in the vacant patches depends mainly on the phenotype of females.

Consequently, the evolutionary response in males will be dominated by an indirect response to

spatial sorting due to the shared genome between the two sexes.

Our theoretical framework also has several practical limitations. The model describes

evolution over a single generation in a patchy landscape with local extinction and colonization

dynamics. Although spatial sorting has been documented in many patchily distributed populations

(Hill et al. 1999, Hanski et al. 2004, Forsman et al. 2011), the model may not be appropriate to

describe evolution in spatially structured populations, such as invasion fronts (Phillips et al. 2006)

and range expansions (Duckworth and Badyaev 2007), where phenotypic changes due to spatial

sorting can compound over multiple generations. This is because our model assumes that, during

spatial sorting, individuals disperse to empty habitats, which makes the model formulation

analytically intuitive and simplifies the estimation of sorting gradients using phenotypic data from

patchy populations. However, this assumption is unlikely to hold at invasion fronts, complicating

theoretical modeling of evolutionary dynamics and quantification of sorting gradient in these

natural settings.

Nevertheless, in some special cases, the phenotypic analysis presented in the paper can also

be applied to structured populations. For example, in early stages of population expansion or when

a new satellite colony is formed far from the invasion front, a biogeographer can divide the

landscape into a well-mixed core habitat and a periphery region. The core (periphery) region can

be treated as occupied (unoccupied) patches in our model. Now, by tracking movement and

phenotypes of individuals from core (thorough mark and recapture or animal trackers) to

periphery, one can estimate the sorting fitness, which can, in turn, be used to estimate the sorting

gradient.

To summarize, our quantitative genetics model and empirical analysis provide quantitative

evidence that, like natural selection, spatial sorting can produce phenotypic shifts on ecological

timescales, As a result, spatial sorting is likely to play an important role in the Anthropocene in

determining the distribution of organisms and spatial genetic variation. And, this general approach

will serve as the foundation for additional work quantifying the magnitude of spatial sorting under

different scenarios in an era of global change, where movement is of increasing importance.

333   **Acknowledgment**

NG was supported by the Continuing and Stengl-Wyer fellowships by the University of

Texas at Austin, Timothy Keitt was supported by the Planet Texas 2050 grant from the University

of Texas at Austin. The monitoring survey of soapberry bugs was funded by the National Science

Foundation to SPE (Award Number: 1802715). We thank Robert Heckman for his help with

setting up the animal model. We also thank Dan Bolnick, Mevin Hooten, and Xinyi Yan for

providing valuable feedback, and Dr. Scott Carroll for introducing MSC and SPE to the soapberry

bug system.

342   **Author Contributions**

343 NG conceived and designed the study based on feedback from TEJ, MSC, and THK. MSC led the

344 survey and performed the crosses with feedback from SPE. NG built the model and analyzed the

345 data with feedback from TEJ. NG wrote the paper with feedback from all authors. Authors declare

346 no conflict of interest.

References

Boag, P. T., and P. R. Grant. 1981. Intense natural selection in a population of Darwin's Finches

(Geospizinae) in the Galapagos. Science **214**:82-85.

Bolnick, D. I., and S. P. Otto. 2013. The magnitude of local adaptation under genotype-dependent

dispersal. Ecology and Evolution **3**:4722-4735.

Brown, C. R., and M. B. Brown. 1998. Intense natural selection on body size and wing and tail

asymmetry in cliff swallows during severe weather. Evolution **52**:1461-1475.

Bumpus, H. C. 1899. The elimination of the unfit as illustrated by the introduced sparrow, Passer

domesticus. Pages 209–226 Biological Lectures, Marine Biological Laboratory, Woods

Hall.

Carpenter, B., A. Gelman, M. D. Hoffman, D. Lee, B. Goodrich, M. Betancourt, M. Brubaker, J.

Guo, P. Li, and A. Riddell. 2017. Stan: A probabilistic programming language. Journal of

Statistical Software **76**:1–32.

Carroll, S. P. 1988. Contrasts in reproductive ecology between temperate and tropical populations

of *Jadera haematoloma*, a mate-guarding hemipteran (Rhopalidae). Annals of the

Entomological Society of America **81**:54-63.

Carroll, S. P. 1991. The adaptive significance of mate guarding in the soapberry bug, *Jadera*

haematoloma (Hemiptera: Rhopalidae). Journal of Insect Behavior **4**:509-530.

Carroll, S. P., M. Marler, R. Winchell, and H. Dingle. 2003. Evolution of cryptic flight morph and

life history differences during host race radiation in the soapberry bug, *Jadera*

haematoloma Herrich-Schaeffer (Hemiptera: Rhopalidae). Annals of the Entomological

Society of America **96**:135-143.

Chen, Q., L. Wang, and R. Tawes. 2008. Hydrodynamic response of northeastern Gulf of Mexico

to hurricanes. Estuaries and Coasts **31**:1098-1116.

Comerford, M. S., T. M. La, S. Carroll, and S. P. Egan. 2023. Spatial sorting promotes

instantaneous (mal)adaptation after hurricane driven-local extinctions. Nature Ecology &

Evolution **7**:1856-1868.

Darwin, C., and A. Wallace. 1858. On the tendency of species to form varieties, and on the

perpetuation of varieties and species by natural means of selection. Journal of the Linnean

Society of London **3**:45 – 62.

De Villemereuil, P., H. Schielzeth, S. Nakagawa, and M. Morrissey. 2016. General methods for

evolutionary quantitative genetic inference from generalized mixed models. Genetics

204:1281-1294.

Dingle, H. 2001. The evolution of migratory syndromes in insects. Pages 159-181 *in* I. P. Woiwod,

D. R. Reynolds, and C. D. Thomas, editors. Insect movement: Mechanisms and

Consequences. . CABI Publishers, Wallingford, UK.

Dingle, H., and R. Winchell. 1997. Juvenile hormone as a mediator of plasticity in insect life

histories. Archives of Insect Biochemistry and Physiology **35**:359-373.

Donihue, C. M., A. Herrel, A. C. Fabre, A. Kamath, A. J. Geneva, T. W. Schoener, J. J. Kolbe,

and J. B. Losos. 2018. Hurricane-induced selection on the morphology of an island lizard.

Nature **560**:88-91.

Duckworth, R. A. 2008. Adaptive dispersal strategies and the dynamics of a range expansion. The

American Naturalist **172**:S4-S17.

Duckworth, R. A., and A. V. Badyaev. 2007. Coupling of dispersal and aggression facilitates the

rapid range expansion of a passerine bird. Proceedings of the National Academy of

Sciences **104**:15017-15022.

Emanuel, K. 2017. Assessing the present and future probability of Hurricane Harvey's rainfall.

Proceedings of the National Academy of Sciences **114**:12681-12684.

Falconer, D. S. 1960. Introduction to Quantitative Genetics. Ronald Press, New York, USA.

Forsman, A., J. Merilä, and T. Ebenhard. 2011. Phenotypic evolution of dispersal-enhancing traits

in insular voles. Proceedings of the Royal Society B: Biological Sciences **278**:225-232.

Grant, B. R., and P. R. Grant. 1993. Evolution of Darwin's finches caused by a rare climatic event.

Proceedings of the Royal Society of London. Series B: Biological Sciences **251**:111-117.

Grant, P. R., B. R. Grant, R. B. Huey, M. T. Johnson, A. H. Knoll, and J. Schmitt. 2017. Evolution

caused by extreme events. Philosophical Transactions of the Royal Society B: Biological

Sciences **372**:20160146.

Hadfield, J. D. 2010. MCMC methods for multi-response generalized linear mixed models: the

MCMCglmm R package. Journal of Statistical Software **33**:1-22.

Hanski, I., C. Erälahti, M. Kankare, O. Ovaskainen, and H. Sirén. 2004. Variation in migration

propensity among individuals maintained by landscape structure. Ecology letters **7**:958-

966.

Hill, J., C. Thomas, and O. Lewis. 1999. Flight morphology in fragmented populations of a rare

British butterfly, Hesperia comma. Biological conservation **87**:277-283.

Kingsolver, J. G., H. E. Hoekstra, J. M. Hoekstra, D. Berrigan, S. N. Vignieri, C. Hill, A. Hoang,

P. Gibert, and P. Beerli. 2001. The strength of phenotypic selection in natural populations.

The American Naturalist **157**:245-261.

Lande, R. 1980. Sexual dimorphism, sexual selection, and adaptation in polygenic characters.

Evolution **34**:292-305.

Lande, R., and S. J. Arnold. 1983. The Measurement of Selection on Correlated Characters.

Evolution **37**:1210-1226.

Lombaert, E., A. Estoup, B. Facon, B. Joubard, J. C. Grégoire, A. Jannin, A. Blin, and T.

Guillemaud. 2014. Rapid increase in dispersal during range expansion in the invasive

ladybird Harmonia axyridis. Journal of evolutionary biology **27**:508-517.

Lush, J. L. 1937. Animal Breeding Plans. Iowa State Press, Ames, Iowa.

MacKay, P. A., and R. J. Lamb. 1979. Migratory tendency in aging populations of the pea aphid,

Acyrthosiphon pisum. Oecologia **39**:301-308.

Matsumura, S., R. Arlinghaus, and U. Dieckmann. 2012. Standardizing selection strengths to study

selection in the wild: a critical comparison and suggestions for the future. Bioscience

62:1039-1054.

McGlothlin, J. W., R. M. Cox, and E. D. Brodie III. 2019. Sex-specific selection and the evolution

of between-sex genetic covariance. Journal of Heredity **110**:422-432.

Miller, M. M., and M. Shirzaei. 2019. Land subsidence in Houston correlated with flooding from

Hurricane Harvey. Remote Sensing of Environment **225**:368-378.

Miller, T. E., A. L. Angert, C. D. Brown, J. A. Lee-Yaw, M. Lewis, F. Lutscher, N. G. Marculis,

B. A. Melbourne, A. K. Shaw, and M. Szűcs. 2020. Eco-evolutionary dynamics of range

expansion. Ecology **101**:e03139.

Novak, S., and R. Kollár. 2017. Spatial gene frequency waves under genotype-dependent dispersal.

Genetics **205**:367-374.

Ochocki, B. M., and T. E. Miller. 2017. Rapid evolution of dispersal ability makes biological

invasions faster and more variable. Nature communications **8**:14315.

Peischl, S., and K. J. Gilbert. 2020. Evolution of dispersal can rescue populations from expansion

load. The American Naturalist **195**:349-360.

Peroni, P. A. 1994. Seed size and dispersal potential of Acer rubrum (Aceraceae) samaras produced

by populations in early and late successional environments. American Journal of Botany

81:1428-1434.

Phillips, B. L., G. P. Brown, J. K. Webb, and R. Shine. 2006. Invasion and the evolution of speed

in toads. Nature **439**:803.

Phillips, B. L., and T. A. Perkins. 2019. Spatial sorting as the spatial analogue of natural selection.

Theoretical Ecology **12**:155-163.

Rice, S. H. 2004. Evolutionary Theory: Mathematical and Conceptual Foundations. Sinauer

Associates, MA.

Risser, M. D., and M. F. Wehner. 2017. Attributable human-induced changes in the likelihood and

magnitude of the observed extreme precipitation during Hurricane Harvey. Geophysical

Research Letters **44**:12,457-412,464.

Roff, D. A. 1996. The evolution of threshold traits in animals. The Quarterly Review of Biology

71:3-35.

Saastamoinen, M., G. Bocedi, J. Cote, D. Legrand, F. Guillaume, C. W. Wheat, E. A. Fronhofer,

C. Garcia, R. Henry, and A. Husby. 2018. Genetics of dispersal. Biological Reviews

93:574-599.

Shine, R., G. P. Brown, and B. L. Phillips. 2011. An evolutionary process that assembles

phenotypes through space rather than through time. Proceedings of the National Academy

of Sciences **108**:5708-5711.

Solow, A. R. 2015. Extreme weather, made by us? Science **349**:1444-1445.

Stott, P. 2016. How climate change affects extreme weather events. Science **352**:1517-1518.

Szűcs, M., M. Vahsen, B. Melbourne, C. Hoover, C. Weiss-Lehman, and R. Hufbauer. 2017. Rapid

adaptive evolution in novel environments acts as an architect of population range

expansion. Proceedings of the National Academy of Sciences **114**:13501-13506.

Trenberth, K. E., L. Cheng, P. Jacobs, Y. Zhang, and J. Fasullo. 2018. Hurricane Harvey links to

ocean heat content and climate change adaptation. Earth's Future **6**:730-744.

Tsai, J.-F., Y.-X. Hsieh, and D. Rédei. 2013. The soapberry bug, *Jadera haematoloma* (Insecta,

Hemiptera, Rhopalidae): First Asian record, with a review of bionomics. ZooKeys **297**:1-

47.

Usui, T., and A. L. Angert. 2024. Range expansion is both slower and more variable with rapid

evolution across a spatial gradient in temperature. Ecology letters **27**:e14406.

van Bergen, E., T. Dallas, M. F. DiLeo, A. Kahilainen, A. L. Mattila, M. Luoto, and M.

Saastamoinen. 2020. The effect of summer drought on the predictability of local

extinctions in a butterfly metapopulation. Conservation biology **34**:1503-1511.

Van Oldenborgh, G. J., K. Van Der Wiel, A. Sebastian, R. Singh, J. Arrighi, F. Otto, K. Haustein,

S. Li, G. Vecchi, and H. Cullen. 2017. Attribution of extreme rainfall from Hurricane

Harvey, August 2017. Environmental Research Letters **12**:124009.

Walsh, B., and M. Lynch. 2018. Evolution and selection of quantitative traits. Oxford University

Press, Oxford, UK:.

Weiss-Lehman, C., R. A. Hufbauer, and B. A. Melbourne. 2017. Rapid trait evolution drives

increased speed and variance in experimental range expansions. Nature communications

8:14303.

Williams, J. L., B. E. Kendall, and J. M. Levine. 2016. Rapid evolution accelerates plant population

spread in fragmented experimental landscapes. Science **353**:482-485.

Figure 1: Map of survey sites across the Greater Houston area in Texas where Comerford et al. (2023) sampled soapberry bugs (**A**). Out of the fifteen sampling sites, eleven (pink) experienced local extinction due to prolonged flooding, while four sites (blue) maintained a stable population throughout the survey period. The photographs on the right show a flooded site (marked with an asterisk in **A**) before (**B**) and after (**C**) Hurricane Harvey from the same vantage point. The image of the macropterous and brachypterous forms of soapberry bug in **A** was taken from Tsai et al. (2013).

496

Figure 2: (A) The survey data of proportion (± 1 sd.) of macropterous insects suggests that post-hurricane, immigrants in the flooded sites had a higher proportion of macropterous insects than in the unflooded sites before the hurricane. **(B)** We find that the mean liability of insects in the flooded sites after the hurricane (right triangle) is 0.58 ± 0.05 ($\beta'_{vw,\phi}$) standard deviation higher than that of insects in unflooded sites before the hurricane (left triangle). The estimated standardized sorting gradient ($\beta'_{vw,\phi}$) is in the top 10 percentile of standardized selection coefficients observed in natural populations (Kingsolver et al. 2001). Genetic crosses show that the liability is a heritable trait ($h^2 = 0.65 \pm 0.13$). Consequently, spatial sorting resulted in a standardized evolutionary change of 0.37 ± 0.08. The variation in the parameter estimates corresponds to standard deviation of the posterior distribution. For details on the statistical analysis, refer to the supplementary R codes.