Transcriptional Landscape of THAP9 across Oligodendrocyte Developmental Stages: Molecular Mechanisms and Network Analysis

Tanuja Bhardwaj, Dhrumi Patel, Sharmistha Majumdar*

Department of Biological Sciences and Engineering, Indian Institute of Technology Gandhinagar

Author1 email: bhardwajtanuja@iitgn.ac.in,

Author 2 email: dhrumi.patel@iitgn.ac.in,

*Corresponding Author email: sharmistham@iitgn.ac.in

Abstract:

Oligodendrocyte maturation and myelination are critical processes in human neurodevelopment, and their dysregulation is linked to numerous neurological disorders. While model organisms have provided insight into these processes, human-specific regulatory mechanisms remain poorly understood. This study investigated human THAP9, a protein homologous to the *Drosophila* P-element transposase, whose function in oligodendrocytes remains unknown. An analysis of publicly available RNA-sequencing data and H3K27ac ChIP-sequencing data from oligodendrocyte progenitor cells (OPCs) and mature oligodendrocytes (MOs) revealed significant upregulation of THAP9 during oligodendrocyte maturation. Co-expression analysis demonstrated a strong correlation with established markers of oligodendrocyte development, including myelin-associated genes (MOG, MBP) and key transcriptional regulators (PDGFRA, SOX5, SOX6, SOX11). THAP9 lacks homologues in mice, highlighting potential human-specific mechanisms in oligodendrocyte development and emphasising the importance of studying species-specific factors in neurodevelopment.. Our findings suggest that THAP9 is a novel human-specific regulator of oligodendrocyte maturation and opens new avenues for studying myelination disorders.

Introduction:

Oligodendrocytes are glial cells in the central nervous system (CNS), responsible for forming the myelin sheath, a protective insulating layer surrounding and supporting axonal nerve fibers[1]. The mammalian CNS comprises approximately 70% myelin (by dry weight) [4], which acts as the

metabolic, physiological and structural support of axons[5]. Degeneration or dysfunction of

oligodendrocytes results in demyelination or dysmyelination, resulting in neurodegenerative

diseases like multiple sclerosis[6].

Understanding the maturation and development of oligodendrocytes is crucial for

uncovering the regulatory mechanisms that govern these processes, providing valuable insights

into neurodegenerative conditions. Neural stem cells give rise to OPCs during embryonic

development[1]. Depending upon the brain region and the CNS developmental stage,

oligodendrocyte precursor cells (OPCs), also called NG2 cells, differentiate into oligodendrocytes,

astrocytes and neurons[2]. However, the fate of OPCs is to develop into oligodendrocytes[3]. The

formation of MOs from OPCs occurs via many regulatory steps: OPCs differentiate into immature

oligodendrocytes, which undergo various morphological changes to form mature and myelinating

oligodendrocytes (MOs). This process is regulated by extrinsic factors, such as PDGF-α, IGF1,

and Neuregulin-1, and intrinsic factors involving transcription factors, such as SOX10, OLIG1,

and OLIG2 [9].

Transposons, or mobile genetic elements, comprise approximately 50% of the human

genome[7]. Transposable elements (TEs) play important roles in neural stem cell differentiation into

various neural lineages[10]. TEs act as sources of cis-regulatory elements, influencing gene

regulatory networks, altering chromatin accessibility, and facilitating transcriptional regulation

essential for the differentiation and function of these critical neural cells[8]. TEs can also integrate

into regulatory networks that control myelination by providing additional regulatory elements that

can enhance or suppress the expression of myelin-related genes, such as Myelin Basic Protein

(MBP) and Proteolipid Protein (PLP)[11]. TEs have also been associated with various

neurodegenerative conditions such as multiple sclerosis and dementia[12]; however, the molecular

mechanisms through which TEs contribute to these conditions are not fully understood [12].

This study focuses on a domesticated DNA transposable element-derived gene called

THAP9, a member of the THAP (THanatose-Associated Protein) family of proteins, which

consists of 12 proteins (THAP1-THAP12)[13] in humans. THAP9 gene maps to chromosome

4q21.22 and consists of 7 exons, though the exact numbers can vary depending on splice variants

[18]. The THAP9 protein is a *Drosophila* P-element transposase (DmTNP) homologue[19]. The

members of the THAP family are involved in functions including apoptosis[14], cell proliferation[14],

pluripotency[15,16] and differentiation of embryonic stem cells (ESCs)[17]. According to the Human

Protein Atlas [20], THAP9 is expressed in the brain, and single-cell studies suggest its enriched

expression in oligodendrocytes. However, its specific role in oligodendrocytes remains unknown,

Most studies on oligodendrocyte development have been performed in model organisms

such as mouse[21]. However, certain differences exist in the biology of oligodendrocytes of humans

and rodents [21], resulting in a lacuna of knowledge about human-specific mechanisms that result in

oligodendrocyte maturation and myelination. We hypothesised that since THAP9, which is highly

expressed in oligodendrocytes, has no homologues in mice; it could be potentially important for

human-specific oligodendrocyte functions or maturation.

In this study, we investigated the role of THAP9 gene in oligodendrocyte biology by

studying its regulation during the maturation and differentiation of OPCs into MOs. Our results

suggest that THAP9 could potentially be involved in oligodendrocyte lineage-specific

development as well as myelination.

Results:

THAP9 Shows Consistent Upregulation During Human Oligodendrocyte Maturation Across
Developmental Stages

The process of oligodendrocyte maturation is critical for proper myelination in the human brain;

despite the importance of this vital process, human-specific aspects of this process remain poorly

understood. We analysed RNAseq data (submitted in GEO [22,23]) of oligodendrocyte precursor cells

(OPCs) and mature oligodendrocytes (MOs) isolated from the gray matter of the dorsolateral

prefrontal cortex of human donors (10 infants samples aged 0-2 years; Figure 1:Ia) and 12 adults

samples aged 30-40 years, Figure 1:IIa). Each cell type forms a distinct cluster indicating different

expression profiles (Figure 1:Ia, IIa). Differential expression analysis revealed extensive

transcriptional changes during oligodendrocyte maturation in both age groups. In infant samples

(Figure 1:Ib), we identified 17,197 differentially expressed genes (9,047 downregulated, 8,150

upregulated). Adult samples (Figure 1:IIb) showed similar numbers with 19,430 differentially

expressed genes (10,479 downregulated, 8,951 upregulated). THAP9 showed consistent and

substantial upregulation in MOs compared to OPCs in both age groups [Figure 1: Ic (Infants: log2

fold change of 1.46 (MOs vs OPCs) with an adjusted p-value of 7.117729e-12) and & IIc (Adults:

log2 fold change of 1.34 (MOs vs OPCs) with an adjusted p-value of 1.17208e-47)].

Thus, through transcriptomic analysis of human dorsolateral prefrontal cortex samples, we
identified THAP9 as a consistently upregulated gene during oligodendrocyte maturation in infant
and adult brain tissue (Suppl. Fig. 1). It is to be noted that most studies about oligodendrocyte
maturation and myelination are performed in mice models. Interestingly, THAP9 lacks a homolog
in rodents like mice and rats, suggesting its potential human-specific role in myelination.
96

Figure 1: Transcriptional Analysis of Human Oligodendrocyte Maturation in Infant and
Adult Samples:
(a) *PCA plot* of I.Infant and II. Adult samples. MOs (red dots), OPCs (turquoise dots).
(b) *Venn diagram showing differential expression analysis comparing MOs and OPCs*, I. In infant
samples; Downregulated genes (salmon-coloured circle;12277), Upregulated genes (green-
coloured circle; 4920 genes, including THAP9) II. In Adult samples; Downregulated genes
(14530); Upregulated genes (4900 genes, including THAP9) [Statistical criteria: log2FoldChange
> 1, Adjusted p-value (p-adj) < 0.1 (indicating statistical significance with FDR correction)].

(c) *Boxplot representing THAP9 gene expression levels* in MOs (blue box) and OPCs (red box) in

I. infant II. adult samples. Expression levels are shown on a log2 scale (y-axis). Statistical analysis

was performed using DESeq2.

Epigenetic signatures suggest that THAP9 is transcriptionally more active in MOs compared

to OPCs.

We next decided to investigate if the higher expression of THAP9 in MOs than in OPCs could be

explained by its altered epigenetic regulation in these cell types. H3K27ac (histone H3 lysine 27

acetylation) is a histone modification associated with active enhancers and promoters and is

associated with positive transcriptional activity of a downstream gene. The THAP9 gene is located

on the forward strand of chromosome 4, at position 82,900,684-82,919,969. H3K27ac ChIP-seq

data revealed a conserved epigenetic signature at this locus, wherein enriched peaks (Figure 2c)

were observed in both infant and adult MOs, suggesting that THAP9 is transcriptionally more

active in mature oligodendrocytes than in oligodendrocyte precursor cells (Figure 2 a,b). Although

there is a slight increase in the H3K27ac signal in OPCs from infant to adult (Figure 2 a,b),

indicating some regulatory activity in these cells, the enrichment of H3K27ac in MOs indicates a

cell-type-specific regulation of THAP9. The stable H3K27ac signal in MOs across developmental

stages implies that THAP9 plays a consistent regulatory role in mature oligodendrocytes, while its

activity in OPCs is comparatively lower. These results are consistent with our observation that

THAP9 is upregulated (in MOs compared to OPCs) and suggest that the gene plays a role during

oligodendrocyte differentiation.

Figure 2: Histone H3K27ac enrichment at the THAP9 gene locus in OPCs and MOs from adult and infant brain samples. H3K27ac ChIP-seq track plots across the THAP9 gene for (a) infant samples for OPC (lavender) and MO (purple). (b) adult samples for OPCs (dark green) and MOs (light green) (c) Combined H3K27ac ChIP-seq track plots across the THAP9 gene for infant MO (purple peaks), adult MO (light green peaks), adult OPC (green peaks), infant OPC (purple peaks). Individual H3K27ac ChIP-seq tracks in THAP9 locus

Transcriptional Regulation, not alternative splicing, causes changes in THAP9 expression in MOs versus OPCs

To investigate the possible regulatory mechanisms that could dictate THAP9 expression during cellular development and maturation, we examined the expression of its alternate isoforms. Isoform switching or the differential usage of alternatively spliced isoforms of a gene can control its function, localisation or stability of the protein it encodes. Both expression and splicing of a gene can change in a spatiotemporal fashion and often vary during stages of development.

Thus, the isoform switching of the THAP9 gene was investigated in both mature oligodendrocytes (MO) and Oligodendrocyte progenitor cells (OPC) across infant and adult samples to identify if the gene is regulated during developmental transitions in these cell types.

Figure 3: Isoform switch analysis of THAP9 in Mature Oligodendrocytes (MO) and Oligodendrocyte Progenitor Cells (OPC). *Top panel*. Five different THAP9 isoforms with ENST numbers, THAP domain (blue) and Tnp_P_element domain (green) **(a)** In MO_Infants: Gene Expression (left panel) represents overall THAP9 gene expression; Isoform Expression (middle panel), represents the amount of specific isoform expressed; Isoform Usage (right panel) represents relative expression of an isoform **(b)** In MO_Adults: Gene Expression (left panel); Isoform Expression (middle panel); Isoform Usage (right panel).

The results obtained included information about gene expression, isoform expression and isoform usage. Gene expression represents the measure of total transcriptional output of the whole THAP9 gene, which is calculated by the sum of expression of all its isoforms (in TPM, Transcript Per

Million); Isoform expression measures the quantity of a specific isoform individually by

calculating the normalised read counts for each THAP9 transcript variant (labelled with unique

ENST number); Isoform usage (isoform fraction - IF) measures the relative amount of each

isoform compared to the total gene expression. Isoform switching refers to a condition when there

is a statistically significant shift in preferential usage of a specific isoform.

The isoform switch analysis demonstrated that the THAP9 gene did not undergo significant

change in isoform usage or switching in MOs and OPCs in samples obtained from adults and

infants (Figure 3). Moreover, the gene shows consistently higher expression of all its isoforms in

MOs (Figure 3 a, b bottom left and middle panels) as compared to OPCs. Thus, THAP9 regulation

appears to be primarily driven by transcriptional control rather than alternative splicing with

conserved expression patterns of the gene in adults as well as infant cells. Among all the isoforms,

ENST00000506208 (highest in MO_Infant) and ENST00000514440 (highest in MO_Adult) show

highest expression levels in both adults and infants. In infants (Figure 3a), ENST00000514440

appears to have higher usage (~ 0.3 or 30%, right panel) in OPCs; however statistical analysis

suggests that this difference is insignificant and cannot be considered an "isoform switch".

Similarly, in adult samples (Figure 3b), although ENST00000514244 has lower expression in

OPCs (middle panel), no significant switch between MOs and OPCs (right panel) was observed.

Thus, we concluded that in both infants and adults, there was no significant change in THAP9

isoform usage in either cell type. These results show that THAP9 maintains its cell specificity with

conserved regulatory mechanisms in both stages of life.

Gene co-expression analysis of THAP9 suggests its potential role in oligodendrocyte

development

To understand the function and regulatory mechanisms of THAP9 in OG cell types, we performed

weighted gene co-expression analysis (WGCNA). WGCNA helps in the identification of a co-

expression module in which a particular gene of interest lies, thus providing insights into its

associations with different cellular states (e.g., OPCs or MOs, across developmental stages). By

analysing these modules, WGCNA allows us to understand the possible regulatory and biological

networks in which a gene functions. In this study, WGCNA was performed to assess THAP9 co-

expression patterns between OPCs and MOs, elucidating the gene's system-dependent roles in

cellular differentiation and maturation processes.

We performed WGCNA separately for infant and adult samples of MOs and OPCs for network

construction with two datasets- one containing only protein-coding genes (Suppl. Fig. 2 for infant

samples, Suppl. Fig. 3 for adult samples) and one containing all genes (comprehensive analysis,

Figure 4).

In comprehensive (containing all genes, Figure 4 **I(b)+II(b)**) and coding gene-specific

analysis (Suppl. Fig 2e, 3e) across adult and infant samples, the THAP9 gene was consistently

found in the turquoise module (represented as MEturquoise) showing significant Module trait

relationship of 0.98 (infants, Suppl. Fig.3, Fig 4) and 1.0 (adults, Suppl. Fig.3, Fig 4)] but opposing

correlations with the two cell types (MOs and OPCs), suggesting the cell-type-specific expression

pattern of the genes in this module. Turquoise module genes are predicted to be highly

interconnected in a biologically important gene network in oligodendrocyte cells since they show

a strong positive correlation between module membership and gene significance (Figure 4:

I+II(c,d); Suppl. Fig 3f+g, 4f+g).

I. Co-expression analysis of oligodendrocyte specific genes in Infant samples

II. Co-expression analysis of oligodendrocyte specific genes in Adult samples

Figure 4: Co-expression network analysis of THAP9 and associated genes obtained by
comprehensive analysis using WGCNA

I(a)+II(a) *Scale mean independence and mean connectivity graphs*. The top graph represents
Scale-free topology model fit (R², Y-axis) vs soft-threshold power (X-axis). The red line indicates
the R² = 0.8 threshold, and the numbers indicate the different powers tested. The lower graph
represents the mean connectivity (Y-axis) vs soft-thresholding power (X-axis) and shows a
decrease in connectivity as the power increases. Selected soft-thresholding power (ß, circled in
green) = 10 (infants) and 8 (Adults). **I(b)+II(b)** *Module-Trait Correlation Heatmap* displays the
correlation between module eigengenes (ME) and two cell types (MOs and OPCs) in Infant (I) and
adult (II) samples. The turquoise module (represented as MEturquoise) contained the THAP9
gene. Positive correlation (red), negative correlation (blue), corresponding correlation values and
p-values in parentheses. **I+II(c,d)** *Module Membership vs Gene Significance* (OPCs and MOs)
shows a strong positive correlation in the infant (I: cor=0.98, p<1e-200) and adult (II:cor=1, p<1e-
200) samples, between module membership (MM) and gene significance (GS) for the Turquoise
module. The scatter plots reveal a strong linear relationship, with genes showing higher MM with

greater GS. The dense clustering of points along the diagonal indicates highly coordinated gene
expression within the Turquoise module. THAP9 (marked in red) appears at the right corner in
both plots, showing high MM (~0.9) and GS (~0.9) in both MOs and OPCs, indicating that THAP9
is a highly connected "hub gene" within the turquoise module in both infant (I) and adult (II)
samples. **I(e)+ II(e)** *Co-expression network* derived from WGCNA demonstrates consistent co-
expression patterns between THAP9 and oligodendrocyte regulatory genes (SOX4, SOX5, SOX6,
SOX11) and marker genes (MBP, MOG and PDGFRA) in both MOs and OPCs in infant (I) and
adult (II) samples. Each node represents a gene, and edges indicate a co-expression relationship.
230
In both MOs and OPCs, THAP9 co-expressed (Figure 4: Ie+ IIe; Suppl. Fig 3h, 4h) with well-
known oligodendrocyte lineage markers like MBP (Myelin Basic Protein, important myelin
structural protein in mature oligodendrocytes[24]), MOG (Myelin Oligodendrocyte Glycoprotein,
key maturation marker and competent of myelin[25]), PDGFRA (Platelet Derived Growth Factor
Receptor Alpha, important for OPC proliferation and maturation, important marker for
oligodendrocyte precursor cells[26]), SOX5 and SOX6 (essential transcription factors for
oligodendrocyte development[27]) and SOX4 and SOX11 (involved in early OPC specification[28]).
All these genes had high weight connections (weight>0.4, Supplementary file 2a + 2b)in coding
genes. THAP9 co-expression patterns with both precursor, PDGFRA and mature, MBP and MOG
markers give us a possible insight into the THAP9 gene's involvement in the multiple stages of
oligodendrocyte differentiation. The co-expression with the SOX genes indicates that THAP9 may
play a role in the transition from OPCs to MOs through its role in differentiation and myelination
processes. It is to be noted that THAP9 has a similar co-expression network architecture in adult
and infant samples. Thus, THAP9 could be a part of an unrecognised conserved regulatory network
controlling oligodendrocyte lineage progression and maintenance of myelination in human-
specific oligodendrocytes.
Interestingly, while the important oligodendrocyte-related genes (MOG, MBP, SOX4,
SOX5, SOX6 and PDGFRA) were all present within the turquoise module in both age groups, we
observed some differences in their strength of correlations (Supplementary file 2c + 2d). In adult
samples, MOG, MBP, SOX4, SOX5, SOX6 and PGDFRA maintained a strong correlation (weight
>0.4, Supplementary file 2d) with THAP9. However, in infant samples, MOG and MBP had lower
correlation values (weight ~0.2, Supplementary file 2c) with THAP9, while SOX4, SOX5,

`SOX6 and PDGFRA retained high correlation values (weight ≥0.4,`
Supplementary file 2c).

To further refine our network, we extracted the THAP9 gene from the edge file in the
turquoise module. We performed filtering to include only genes of our interest (i.e. the above-
mentioned oligodendrocyte regulatory and marker genes), thereby constructing a focused gene
interaction network (Figure 4: I(e)+ II(e)). We performed hierarchical clustering analysis on infant
and adult samples, showing distinct clustering patterns between the two groups. Infant samples
were categorised as MOs (green) and OPCs (orange) (Suppl. Figure 4a). Similarly, adult samples
were grouped into OPCs (green) and MOs (orange) (Suppl. Figure 4d) and block dendrograms
displaying block sizes, with infant samples containing 8893 genes in block 1 and 5597 genes in
block 2 (Suppl. Fig. 4:b,c) and adult samples containing 8558 genes in block 1 and 5875 genes in
block 2 (Suppl. Fig. 4:e,f), which included both coding and non-coding genes.

This comprehensive analysis of both coding and non-coding genes resulted in a more
robust understanding of the transcriptional landscape. We identified 6,455 genes in the turquoise
module of infant samples, with 20,830,285 edges, and 7,071 genes in adult samples, with
24,995,985 edges (Supplementary file 2c + 2d). In contrast, when restricting the analysis to only
coding genes, the turquoise module in infant samples contained 5,782 genes with 16,712,871
edges, while in adult samples, it contained 7,296 genes with 26,612,160 edges (Supplementary
file 2a + 2b). This difference demonstrates that including non-coding genes increases the number
of genes and the number of connections (edges) within the turquoise module. Non-coding genes
contribute additional layers of regulatory complexity and network connectivity.

Our detailed WGCNA analysis provides novel insights into the possible involvement of
THAP9 in oligodendrocyte development. This is exemplified by THAP9's consistent module
placement and high trait relationship (Suppl. Fig. 3,4; Figure 4). However, the variations observed
between the comprehensive and coding gene analysis highlight the complex molecular landscape
of oligodendrocyte development biology. For instance, the comprehensive analysis gives a more
detailed network pattern that includes all regulatory factors and associated genes (Supplementary
file 2c + 2d). On the other hand, the coding-gene-specific analysis demonstrates a network wherein
THAP9 shows more conserved co-expression with developmental genes in infant samples (Figure
4: Ie, Suppl. Fig. 2h), providing a stronger and more specific maturation process. Conversely, adult
samples display more dynamic molecular interaction (Figure 4: IIe, Suppl. Fig. 3h), with variations

in the gene co-expression relationships (i.e., topological overlap measures between genes) and

absence of some genes such as SOX11 (weight < 0.2) (<u>Supplementary file 2d</u>). These patterns

highlight the characteristics of ongoing developmental processes and early glial and neural

maturation. The observed differences in the co-expression network (Fig. I(e)+ II(e)) between the

adult and infant samples also display the dynamic nature of oligodendrocyte development. The

high module trait relationship (Fig I+II c,d) suggests that while the fundamental molecular

mechanism remains the same, some subtle regulatory changes exist across the developmental

stages. Despite these variations, the overall network structure and the strength of gene interactions

remained consistent across conditions, whether analysing all genes or only coding genes. This

consistency emphasises the robustness of the transcriptional networks governing oligodendrocyte

development. The high module-trait relationship (Fig. I+II c, d) suggests that while the

fundamental molecular mechanisms remain conserved, subtle regulatory shifts manage the

transition between developmental stages.

THAP9 expression during direct reprogramming of human fibroblasts to Oligodendrocyte

lineage cells

To understand the role of THAP9 in oligodendrocyte lineage cells, we further studied its

expression in mature cells that have been reprogrammed to oligodendrocyte lineage.

Reprogramming a mature cell into an oligodendrocyte lineage requires re-establishing all the

epigenetic states and cell identity; this can provide insights into a gene's role in lineage-specific

transcription or the pathways essential for transition and oligodendrocyte lineage specification.

Detailed expression analysis can help us understand whether a gene is required in earlier cell

commitment or for terminal differentiation and myelination.

308   Towards this, we analysed RNAseq data[29] from a study in which oligodendrocyte lineage

cells were obtained by reprogramming human fibroblasts using a small molecule cocktail (A83-

01, thiazovivin, purmorphamine, VPA, and forskolin) and OCT4 as induction factor. The study

used two reporter systems as indicators for reprogramming success to track and isolate cells at

specific developmental stages: OLIG2-positive (OLIG2::eGFP) cells marking early stages and

SOX10-positive cells (SOX10::eGFP) to mark the progression to more mature stages. Different

stages of the cells were made using specific markers: OLIG2, SOX10, O4+ and MBP. The study

 concluded that SOX10::eGFP cells show better maturation capacity, i.e., higher MBP+ cells and
 more mature morphology than OLIG2::eGFP cells[29].

Figure 5: Gene expression analysis in fibroblasts reprogrammed into maturing oligodendrocytes shows different transcriptomics profile

(**a**) *Heatmap of gene expression in logTPM*. High expression (red), low expression (blue), intermediate (white) expression. Rows represent different genes; columns represent different samples with colour-coded legends (on the right). The top of the dendrogram shows the hierarchical clustering of the samples. (**b**) *PCA (Principle Component Analysis) plot* shows the relationship between the samples in two dimensions, PC1 (x-axis) and PC2 (y-axis), with each coloured dot representing a sample. The samples clustered together show similar gene expression patterns, and the samples with different expressions are in separate groups. (**c**) THAP9 gene expression in induced cells. (**d**) MBP gene expression in induced cells. In c,d; Y-axis [mean TPM (Transcript per Million)], each bar with error bars (SD) representing a different sample

THAP9 displays distinct expression patterns in native (Figure 1: Ic+IIc) versus reprogrammed cells, suggesting its role may differ between natural development and induced reprogramming.

Figure 5c shows that THAP9 is highly expressed in early reprogramming stages of induced

fibroblasts [A2B5(DF), A2B5(BJ)] but decreases in later stages of induced OPCs[O4(D40),

O4(D60)], indicating a potential role in initiating lineage conversion.OThe high expression of

THAP9 in OLIG2_labelled reprogrammed cells across datasets suggests a functional relationship

between THAP9 and OLIG2 (Figure 5, 6), a key regulator of oligodendrocyte differentiation. On

the other hand, in normal human cells (Figure 1: Ic+IIc), THAP9 is upregulated in

oligodendrocytes when compared to OPCs. The expression patterns of THAP9 (upregulated in

native cells and downregulated in reprogrammed cells) are thus different. This analysis suggests

that the oligodendrocytes obtained by reprogramming human fibroblasts do not completely mimic

their natural developmental trajectory since there are marked differences in their transcriptomes as

well as the time durations and morphologies required to express relevant markers.

Here, THAP9 exhibits high expression (Figure 5c) in OPCs derived from induced

pluripotent stem cells (iPSCs), suggesting a role in the initial stages of oligodendrocyte

development. As OPCs mature, THAP9 expression decreases, but it remains elevated in cells

expressing key transcription factors involved in oligodendrocyte differentiation, such as OLIG2

and SOX10. This expression pattern suggests that THAP9 may interact or be regulated by these

transcriptional networks during oligodendrocyte maturation.

Figure 6: Specific marker proteins associated with each stage of oligodendrocyte

development. Markers for the progression from immature oligodendrocyte precursor cells

(iOPCs) to late OPCs, immature oligodendrocytes, and mature oligodendrocytes.

Marker gene	Developmental Stage	References
A2B5 (Antibody)	OPCs	30–34
OLIG2 (Transcription factor)	OPCs, Premyelinating Oligodendrocytes, Myelinating Oligo	30–36
SOX10 (Transcription factor)	OPCs, Premyelinating Oligodendrocytes, Myelinating Oligodendrocytes	30–33,35,36
O4 (Antibody)	Premyelinating Oligodendrocytes, Myelinating Oligodendrocytes	30–32,34–36
MBP (Myelin Component)	Myelinating Oligodendrocytes	30–32,34–36

A similar expression pattern was observed for the MBP (Myelin Basic Protein) gene. MBP is
essential for the formation and functions of mature oligodendrocytes[37]. Both THAP9 and MBP
have higher expression in the early stages of oligodendrocyte development (A2B5+, OLIG2+
cells), which decreases through maturation; this could signify that both genes are co-regulated
during reprogramming. Moreover, both genes show similar expression patterns (Figure 5c+d) to
OCT4-mediated reprogramming. MBP usually has high expression in mature cells, suggesting
reprogramming may need more optimisation to replicate natural oligodendrocyte development. It
also signifies that O4(D60) cells have undergone incomplete maturation (Figure 5d) or alternative
developmental pathways or there are issues with maintaining oligodendrocyte identity. THAP9
expression is specific to the oligodendrocyte lineage and may not be as highly expressed in other
cell types, such as fibroblasts. Overall, the observed expression patterns suggest that THAP9 plays
prominent roles in both the early stages of oligodendrocyte progenitor cell development and in
regulating oligodendrocyte development and differentiation. Further functional studies would be
necessary to elucidate the precise mechanisms by which THAP9 contributes to these processes.

Discussion:

Oligodendrocytes are cells responsible for myelination. Oligodendrocyte progenitor cells mature
and form mature myelinating oligodendrocytes. Oligodendrocytes from different developmental
stages exhibit different myelination abilities[39]. The proliferation, maturation and maintenance of

oligodendrocytes is important since dysregulation in these processes results in conditions like

neurodegenerative diseases or delayed repair ability after injury[38].

In this comprehensive bioinformatics investigation, we studied the role of THAP9 in

oligodendrocyte development. THAP9, encoded by a DNA transposable element-derived gene, is

homologous to the P-element transposase of *Drosophila melanogaster* and is a member of the

human THAP protein family of 12 members [40,41]. Our analysis revealed a pattern of THAP9

upregulation during oligodendrocyte maturation, consistently observed across infant and adult

tissue samples (Figure 1: Ic+IIc in infant and adult samples; 1.46 and 1.34 log2FoldChange,

respectively). The analysis of histone modification H3K27ac across the THAP9 gene locus in MOs

and OPCs provided us with further confirmation that THAP9 indeed shows more transcriptional

activity in the MOs than OPCs. This persistent and stage-specific expression profile highlights that

the THAP9 gene may play an important role in oligodendrocyte differentiation throughout human

development, potentially influencing crucial processes such as cellular maturation and myelination

in the central nervous system.

During the developmental trajectory of OPCs into MOs, transcription factors like SOX4,

SOX5, SOX6, and SOX11 are active in the OPCs and downregulate during maturation. Co-

expression network analysis in both infant and adult human samples (weight >0.4, Supplementary

file 2) show that THAP9 demonstrates an inverse expression pattern to these transcription factors.

On the other hand, THAP9 is upregulated in MOs, along with important myelination markers MBP

and MOG, which are expressed specifically in late stages of oligodendrocyte differentiation. It is

interesting to note that the upregulation of THAP9 along with MOG and MBP (Supplementary file

3) coincides with the downregulation of SOX4, SOX5, SOX6, and SOX11, as well as early

specification factors and PDGFRA. This indicates that THAP9 might be involved in the regulatory

network governing the maturation of oligodendrocytes or in the differentiation, maintenance and

myelination process in MOs.

Most studies in oligodendrocyte biology have been carried out in mouse model systems.

However, significant differences exist between human and mouse oligodendrocyte biology due to

the evolutionary divergence between humans and mouse[42]. These variations due to interspecies

differences have resulted in gaps in the information, making it important to study more human-

specific genes. Human and mouse oligodendrocyte biology varies in various ways, which include

differences in myelination timing[43], cell turnover rates, and molecular signatures[44]. Around two

hundred genes are expressed in human oligodendrocytes but are not expressed in mouse[44]. The

absence of the THAP9 gene in several rodents, including mice and rats, indicates that it may be

involved in human-specific pathways related to oligodendrocyte differentiation and maturation

and contribute to some of the unique features of human myelination compared to mouse models.

Our studies show that THAP9 is a promising candidate gene linked to human

oligodendrocyte maturation. We also highlight the importance of species-specific genes for

oligodendrocyte maturation and emphasise the importance of human-specific genes in this process.

Thus, we also highlight the fundamental challenges in neurogenetic research due to the limitations

of model organisms to completely understand the molecular mechanisms involved in human

neurological development.

Methodology:

Downloading of Datasets and Preprocessing:

GSE239662, GSE130063 (RNAseq) and GSE239660 (ChIP-seq) datasets from the GEO (Gene

Expression Omnibus) database[45] were downloaded. All the raw files were downloaded from the

NCBI SRA (Sequence Read Archive) using SRA-toolkit [46] and the prefetch (3.1.1) function. All

the .sra files were converted to .fastq. FastQC (version 0.12.1) [47] was used to control the quality

of all the sequencing files. The trimming and removal of low-quality sequences of sequencing files

were done using Fastp (version 0.23.4) [48].

The GSE239662 dataset consists of RNAseq samples obtained from the gray matter of the

dorsolateral prefrontal cortex of the human brain, comprising two age groups: Adults aged 30-40

years (n=12 individuals) and Infants ages 0-2 years (n=10 individuals). Two distinct

oligodendrocyte lineage populations were isolated from each individual and sequenced:

Oligodendrocyte precursor cells (OPCs) and Mature oligodendrocytes (MOs). Consisting of a total

of 39 RNA-sequencing files.

The GSE130063 dataset consists of RNA sequencing profiles of 24 samples (8 samples in different

stages in triplicates) of induced oligodendrocyte progenitor cells (iOPCs) generated from human

somatic cells. The different cell populations included (A2B5+ iOPCs, O4+iOPCs, OLIG2+iOPCs,

SOX10+iOPCs and Fibroblasts) were analysed in triplicates. A2B5(BJ): A2B5+ makers cells

derived from human fibroblast (BJ) induction representing early-stage iOPCs; A2B5(DF): A2B5+

makers cells derived from human dermal fibroblasts induction representing early-stage iOPCs;

OLIG2_labelled (cells with OLIG2+ maker derived from OLIG2::eGFP fibroblasts, representing

completely developed OPCs), SOX10_labelled (cells with SOX10+ maker derived from

SOX10::eGFP fibroblasts, representing a completely developed OPCs), O4(D40) and O4(D60),

iOPCs derived from BJ on day 40 and 60; OLIG2_ESC, OLIG2::eGFP fibroblasts and DF, dermal

fibroblasts. GFP+ cells had bipolar morphology, as expected in iOPCs. A total of 12 triplicate

samples were used in the study.

The GSE239660 dataset consists of 24 ChIP-seq samples analysing H3K27ac histone modification

marks in oligodendrocyte lineage populations isolated from the gray matter of the human

dorsolateral prefrontal cortex. Samples were obtained from both infant and adult donors.

Immunoprecipitation was performed using rabbit polyclonal anti-H3K27ac antibody to analyse

the distribution of active enhancer marks. The dataset includes corresponding input controls for

each experimental condition. The dataset includes samples from infants (5 MOs and 5 OPCs) and

adults (5 MOs and 5 OPCs)

All the downstream analysis was carried out on R (version 4.3.1)

RNAseq analysis: For RNAseq analysis, transcript quantification was done using Kallisto

(version 0.44.0)[49]. The abundance files were then processed in R (version 4.3.1) using

tximport(version 1.28.0). Gene-level counts were obtained by summarising transcript-level

abundances using "length-scaled TPM" values. Transcript-to-gene mapping was done using

human genome annotations from EnsDb.Hsapiens.v86. The differential expression analysis was

performed using DESeq2 (version 1.40.2) [50]. A DESeqDataSet was constructed using raw count

data with cell type as an experimental design factor. Differential expression between MOs and

OPCs cell types was assessed using DESeq2 implementation of the Wald test. Genes with an

adjusted p-value < 0.05 (Benjamini-Hochberg correction) were considered significantly

differentially expressed.

Isoform analysis: The RNAseq data was processed using R version 4.3.1, and an isoform analysis

was done using the IsoformSwitchAnalzeR package[51] in R. Transcript-level quantification data

was imported and analysed. The analysis was performed on the human genome assembly GRCh38

(Ensembl release 112) with corresponding transcript annotation. Differential exon usage was

assessed using DEXSeq through the IsoformSwitchAnalyzeR framework.

Multiple computational tools were employed to characterise the functional potential of

transcripts: Coding potential was evaluated using CPC2 (Coding Potential Calculator 2)[52] to

predict which isoforms were coding and non-coding. This tool supports Vector machine-based

classifiers and uses six biologically important sequence features to make predictions. Protein

domains were identified using PFam[53], and signal peptides were predicted using SignalP-5.0 [54];

Intrinsically disordered regions (IDRs) were identified using IUPred2A[55], which makes sequence-

based predictions of unstructured regions. Subcellular localisation prediction was performed using

the DeepLoc2 [56] tool, which uses deep learning methods to predict cellular compartment

localisation and map the cellular distribution of isoforms. Transmembrane topology was analysed

using the DeepTMHMM [57] tool that uses deep learning architecture.

Gene Co-expression analysis: Weighted Gene Co-expression Network Analysis (WGCNA)[58]

(version 1.72-5) was used to characterise the transcriptional networks and identify genes showing

coordinated expression patterns. The count's matrix was normalised using Variance Stabilizing

Transformation (vst). Outliers were checked using Hierarchical clustering and PCA. Soft-

thresholding power (ß) was calculated to get the scale-free network. To get the appropriate power,

the cut-off value for R^2 was set to 0.8. An adjacency matrix was formed, and a Topological Overlap

Matrix (TOM) was calculated using the calculated power. BlockwiseModules function was used

to detect the modules, where the minimum module size was set to 30 and the maximum block size

to 9000. To merge the module based on dissimilarity, the cut height was set to 0.25. Module

eigengenes (MEs) were obtained and used to calculate module trait (cell type) using Pearson

correlation and were displayed in the form of a heatmap. Module membership (MM) and gene

significance (GS) were calculated to get more information on individual genes' correlation with

the module eigengene and traits belonging to the Turquoise module (module with the genes of

interest). Genes with absolute MM and GS higher than 0.8 were selected and filtered to get the

edge file. Edges with a weight of less than 0.25 were removed to get a refined network. A

clusterProfiler (version 4.10.1) package was used for the functional enrichment analysis.

Molecular Function (MF), Biological Process (BP), and Cellular Component (CC) were obtained

for the desired module; the p-value and q-value cut-off were set to 0.05 and 0.1, respectively.

ChIP-seq analysis and visualisation: After trimming and removing low-quality reads, all the

sequence files were mapped to the human genome assembly GRCh38/hg38 using Bowtie2

(version 2.5.4)[59]. All the multi-mapping reads and low-quality alignments were removed using

SAMtools (version 1.20)[60]. Read mapping to ENCODE-blacklisted genomic regions was removed

using BEDTools (version 2.31.1)[61]. MACS2 (version 2.2.9.1)[62] peak calling was used to identify

H3K27-ac regions, with input controls for both cell types (OPCs and MOs) of both age groups

(infants and adults). All the signal tracks were normalised to 1 million reads and converted to

bigWig format (.bw files) for visualisation. The Integrative Genomics Viewer (IGV) (version

2.17.0)[63] was used to visualise H3K27ac peaks and signal distribution across the genome to get a

detailed view of chromatin modifications at specific genomic loci.

Author Contributions: TB performed all data analyses, prepared the figures, and drafted the

manuscript. DP conducted the WGCNA analysis and contributed to the interpretation of the

results. As the corresponding author, SM supervised the study, provided critical revisions, and

edited the manuscript. All authors reviewed and approved the final version of the manuscript.

Code and data: No new codes were developed for this study. All tools and scripts used in the

analysis have been documented and made publicly available in the GitHub repository.

Files attached:

Supplementary file 1

Supplementary file 2

Supplementary file 3

Conflict of interests: None

REFERENCES:

1.   Bradl, M. & Lassmann, H. Oligodendrocytes: biology and pathology. *Acta Neuropathol.*

119, 37 (2010).

2.   Suzuki, N. *et al.* Differentiation of Oligodendrocyte Precursor Cells from Sox10-Venus

Mice to Oligodendrocytes and Astrocytes. *Sci. Rep.* **7**, 1–11 (2017).

3.   Nishiyama, A., Suzuki, R. & Zhu, X. NG2 cells (polydendrocytes) in brain physiology and

repair. *Front. Neurosci.* **8**, 133 (2014).

4.   Morell, P. & Quarles, R. H. Characteristic Composition of Myelin. in *Basic*

Neurochemistry: Molecular, Cellular and Medical Aspects. 6th edition (Lippincott-Raven,

1999).

5.   Hughes, E. G. & Appel, B. The cell biology of CNS myelination. *Curr. Opin. Neurobiol.*

39, 93 (2016).

6.   Multiple sclerosis and demyelination. *Curr. Diagn. Pathol.* **13**, 193–202 (2007).

7.   Transposons. https://www.nature.com/scitable/topicpage/transposons-the-jumping-genes-

518/.

8.   Sekine, K., Onoguchi, M. & Hamada, M. Transposons contribute to the acquisition of cell

type-specific cis-elements in the brain. *Communications Biology* **6**, 1–16 (2023).

9.   Emery, B. & Richard Lu, Q. Transcriptional and Epigenetic Regulation of Oligodendrocyte

Development and Myelination in the Central Nervous System. *Cold Spring Harb. Perspect.*

Biol. **7**, (2015).

10.   Schumann, G. G. *et al.* The impact of transposable element activity on therapeutically

relevant human stem cells. *Mob. DNA* **10**, 1–23 (2019).

11.   Ghosh, T. *et al.* A retroviral link to vertebrate myelination through retrotransposon-RNA-

mediated control of myelin gene expression. *Cell* **187**, 814–830.e23 (2024).

12.   Ravel-Godreuil, C., Znaidi, R., Bonnifet, T., Joshi, R. L. & Fuchs, J. Transposable elements

as new players in neurodegenerative diseases. *FEBS Lett.* **595**, 2733–2755 (2021).

13. Roussigne, M. *et al.* The THAP domain: a novel protein motif with similarity to the DNA-

binding domain of P element transposase. *Trends Biochem. Sci.* **28**, 66–69 (2003).

14. Clouaire, T. *et al.* The THAP domain of THAP1 is a large C2CH module with zinc-

dependent sequence-specific DNA-binding activity. *Proc. Natl. Acad. Sci. U. S. A.* **102**,

6907–6912 (2005).

15. Baker, M. The latest pluripotency factor is a lone warrior. *Nat. Rep. Stem Cells* 1–1 (2008).

16. Ziaei, S., Rezaei-Tavirani, M., Ardeshirylajimi, A., Arefian, E. & Soleimani, M. Induced

Overexpression of THAP11 in Human Fibroblast Cells Enhances Expression of Key

Pluripotency Genes. *Galen Med J* **8**, e1308 (2019).

17. Aguilo, F. *et al.* THAP1: Role in Mouse Embryonic Stem Cell Survival and Differentiation.

Stem Cell Reports **9**, 92–107 (2017).

18. Home - Gene - NCBI. https://www.ncbi.nlm.nih.gov/gene.

19. Majumdar, S., Singh, A. & Rio, D. C. The human THAP9 gene encodes an active P-

element DNA transposase. *Science* **339**, 446–448 (2013).

20. Thul, P. J. & Lindskog, C. The human protein atlas: A spatial map of the human proteome.

Protein Science : A Publication of the Protein Society **27**, 233 (2017).

21. Dietz, K. C., Polanco, J. J., Pol, S. U. & Sim, F. J. Targeting human oligodendrocyte

progenitors for myelin repair. *Exp. Neurol.* **283**, 489 (2016).

22. Kozlenkov, A. *et al.* Novel method of isolating nuclei of human oligodendrocyte precursor

cells reveals substantial developmental changes in gene expression and H3K27ac histone

modification. *Glia* **72**, 69–89 (2024).

23. Barrett, T. *et al.* NCBI GEO: archive for functional genomics data sets--update. *Nucleic*

Acids Res **41**, D991–5 (2013).

24.   Galiano, M. R. *et al.* Myelin basic protein functions as a microtubule stabilizing protein in

differentiated oligodendrocytes. *Journal of neuroscience research* **84**, (2006).

25.   Peschl, P., Bradl, M., Höftberger, R., Berger, T. & Reindl, M. Myelin Oligodendrocyte

Glycoprotein: Deciphering a Target in Inflammatory Demyelinating Diseases. *Front.*

Immunol. **8**, 248125 (2017).

26.   Rivers, L. E. *et al.* PDGFRA/NG2 glia generate myelinating oligodendrocytes and piriform

projection neurons in adult mice. *Nature Neuroscience* **11**, 1392–1401 (2008).

27.   Ittner, E. *et al.* SoxD transcription factor deficiency in Schwann cells delays myelination in

the developing peripheral nervous system. *Scientific Reports* **11**, 1–11 (2021).

28.   Potzner, M. R. *et al.* Prolonged Sox4 Expression in Oligodendrocytes Interferes with

Normal Myelination in the Central Nervous System. *Molecular and Cellular Biology* **27**,

5316 (2007).

29.   Yun, W. *et al.* OCT4-induced oligodendrocyte progenitor cells promote remyelination and

ameliorate disease. *npj Regenerative Medicine* **7**, 1–15 (2022).

30.   Barateiro, A. & Fernandes, A. Temporal oligodendrocyte lineage progression: in vitro

models of proliferation, differentiation and myelination. *Biochim Biophys Acta* **1843**, 1917–

1929 (2014).

31.   Goldman, S. A. & Kuypers, N. J. How to make an oligodendrocyte. *Development* **142**,

3983–3995 (2015).

32.   Foerster, S., Hill, M. F. E. & Franklin, R. J. M. Diversity in the oligodendrocyte lineage:

Plasticity or heterogeneity? *Glia* **67**, 1797–1805 (2019).

33.   Antel, J. P. *et al.* Immunology of oligodendrocyte precursor cells in vivo and in vitro. *J*

Neuroimmunol **331**, 28–35 (2019).

34. Akay, L. A., Effenberger, A. H. & Tsai, L.-H. Cell of all trades: oligodendrocyte precursor

cells in synaptic, vascular, and immune function. *Genes Dev* **35**, 180–198 (2021).

35. Huang, H., He, W., Tang, T. & Qiu, M. Immunological Markers for Central Nervous

System Glia. *Neurosci Bull* **39**, 379–392 (2023).

36. Kamen, Y., Pivonkova, H., Evans, K. A. & Káradóttir, R. T. A Matter of State: Diversity in

Oligodendrocyte Lineage Cells. *Neuroscientist* **28**, 144–162 (2022).

37. Stage-specific Expression of Myelin Basic Protein in Oligodendrocytes Involves Nkx2.2-

mediated Repression That Is Relieved by the Sp1 Transcription Factor. *Journal of*

Biological Chemistry **280**, 16284–16294 (2005).

38. Disease-associated oligodendrocyte responses across neurodegenerative diseases. *Cell*

Reports **40**, 111189 (2022).

39. Cui, Q. L. *et al.* Human fetal oligodendrocyte progenitor cells from different gestational

stages exhibit substantially different potential to myelinate. *Stem cells and development* **21**,

(2012).

40. Roussigne, M. *et al.* The THAP domain: a novel protein motif with similarity to the DNA-

binding domain of P element transposase. *Trends in biochemical sciences* **28**, (2003).

41. Majumdar, S., Singh, A. & Rio, D. C. The human THAP9 gene encodes an active P-

element DNA transposase. *Science (New York, N.Y.)* **339**, (2013).

42. Semple, B. D., Blomgren, K., Gimlin, K., Ferriero, D. M. & Noble-Haeusslein, L. J. Brain

development in rodents and humans: Identifying benchmarks of maturation and

vulnerability to injury across species. *Progress in neurobiology* **0**, 1 (2013).

43. Gargareta, V.-I. *et al.* Conservation and divergence of myelin proteome and

621   oligodendrocyte transcriptome profiles between humans and mice. (2022)

622   doi:10.7554/eLife.77019.

623  44. Chamling, X. *et al.* Single-cell transcriptomic reveals molecular diversity and

624   developmental heterogeneity of human stem cell-derived oligodendrocyte lineage cells.

625   *Nature Communications* **12**, 1–20 (2021).

626  45. Home - GEO - NCBI. https://www.ncbi.nlm.nih.gov/geo/.

627  46. 01. Downloading SRA Toolkit. *GitHub* https://github.com/ncbi/sra-tools/wiki/01.-

628   Downloading-SRA-Toolkit.

629  47. Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput Sequence

630   Data. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

631  48. Chen, S. Ultrafast one-pass FASTQ data preprocessing, quality control, and deduplication

632   using fastp. *iMeta* **2**, e107 (2023).

633  49. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq

634   quantification. *Nature Biotechnology* **34**, 525–527 (2016).

635  50. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

636   for RNA-seq data with DESeq2. *Genome Biology* **15**, 1–21 (2014).

637  51. Vitting-Seerup, K. & Sandelin, A. IsoformSwitchAnalyzeR: analysis of changes in genome-

638   wide patterns of alternative splicing and its functional consequences. *Bioinformatics*

639   *(Oxford, England)* **35**, (2019).

640  52. Kang, Y.-J. *et al.* CPC2: a fast and accurate coding potential calculator based on sequence

641   intrinsic features. *Nucleic Acids Res* **45**, W12–W16 (2017).

642  53. Mistry, J. *et al.* Pfam: The protein families database in 2021. *Nucleic Acids Res* **49**, D412–

643   D419 (2021).

54. Almagro Armenteros, J. J. *et al.* SignalP 5.0 improves signal peptide predictions using deep

neural networks. *Nature Biotechnology* **37**, 420–423 (2019).

55. Mészáros, B., Erdos, G. & Dosztányi, Z. IUPred2A: context-dependent prediction of protein

disorder as a function of redox state and protein binding. *Nucleic Acids Res* **46**, W329–

W337 (2018).

56. Thumuluri, V., Almagro Armenteros, J. J., Johansen, A. R., Nielsen, H. & Winther, O.

DeepLoc 2.0: multi-label subcellular localization prediction using protein language models.

Nucleic Acids Res **50**, W228–W234 (2022).

57. Hallgren, J. *et al.* DeepTMHMM predicts alpha and beta transmembrane proteins using

deep neural networks. (2022) doi:10.1101/2022.04.08.487609.

58. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network

analysis. *BMC Bioinformatics* **9**, 1–13 (2008).

59. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nature*

Methods **9**, 357–359 (2012).

60. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, (2021).

61. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* **26**, 841–842 (2010).

62. Gaspar, J. M. Improved peak-calling with MACS2. (2018) doi:10.1101/496521.

63. Thorvaldsdóttir, H., Robinson, J. T. & Mesirov, J. P. Integrative Genomics Viewer (IGV):

high-performance genomics data visualization and exploration. *Briefings in Bioinformatics*

14, 178 (2012).

665