1

Main Manuscript for

When neighbours play a role: a systems-level analysis of protein interactions conditioning cancer driver mutation effects

João AI Miranda [§1,2], Márcia FS Vital[§1], Margarida Carrolo[§3], António Quintela[3], Francisco R Pinto[1]

[1] BioISI, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisbon, Portugal

[2] INESC-ID, Instituto Superior Técnico, Universidade de Lisboa, 1000-029, Lisbon, Portugal

[3] Centro Hospitalar Universitário Lisboa Norte, Hospital de Santa Maria, Lisboa, Portugal

[§] These authors have contributed equally to this work

*Francisco R Pinto

Email: frpinto@ciencias.ulisboa.pt

Author Contributions: FRP designed the study. JAIM, MFSV and FRP conducted computational work and data analysis. MC and AQ helped in cancer data interpretation. FRP wrote the first version of the manuscript. All authors revised the manuscript and contributed to the final version of the manuscript.

Competing Interest Statement: The authors have no competing interests to declare.

Classification: Biological Sciences / Systems Biology

Keywords: Cancer; Protein Physical Interactions; Driver Mutations.

This PDF file includes:

  Main Text
  Figures 1 to 4
  Tables 1 to 2

1
Abstract

Background: Cancer is driven by the accumulation of somatic mutations, including driver
mutations that confer a selective advantage to cancer cells. Driver proteins operate within
complex interaction networks, and their activity is conditioned by neighbour proteins.
Understanding the interplay between driver mutations and the expression of their neighbour
proteins can provide insights into cancer biology and potential therapeutic targets.

Methods: We assessed associations between expression of neighbour proteins and driver
mutation status, comparing both between and within cancer types. We further evaluated if
neighbours were enriched in significant associations with multiple drivers and characterised the
impact of neighbour expression on overall survival for all cancer types.

Results: We found a significant correlation between the number of driver associations a
neighbour gene has and the number of sign-coherent survival associations, particularly for
neighbours enriched in positive associations, where high neighbour expression correlated with
increased driver mutations and poorer survival. We identified 119 neighbours enriched in positive
driver associations with at least two unfavourable survival associations and 25 neighbours
enriched in negative driver associations with at least two favourable survival associations.

Conclusions: Our study systematically identified neighbours associated with driver mutation
status. Complementary evidence from survival analysis and the literature suggests that
neighbours enriched in driver associations can be further explored as drug target candidates.

Significance Statement
Cancers are caused by mutations in driver genes. The impact of those mutations in the cell can
be influenced by other proteins in the cell that physically interact with the mutated protein. In this
work, we analysed cancer patient data to uncover such neighbour proteins that may influence the
outcome of driver mutations. We discovered a subset of neighbour proteins that are associated
with mutations in multiple cancer drivers and, simultaneously, are associated with changes in
survival times of patients for multiple cancer types. These neighbour proteins may help explain
why some driver mutations are more common in certain cancer types. We also propose that
these neighbour proteins should be explored as candidate drug targets for cancer therapy.
30
31 **Main Text**
32
33 **Introduction**
34
Over the last decades, our understanding of oncogenesis has significantly expanded. The
discovery of cancer driver genes in the 1970s has contributed to the advancement of targeted
anticancer therapies and, more broadly, to the identification of genomic biomarkers for predicting
prognosis and treatment response (1). Likewise, the arrival of genomic sequencing of tumours has
led to a remarkable collection of disease-driving mutations, offering valuable insights into cancer
genomics (2).

Although 91% of tumours carry at least one gene driver mutation (3), these mutations alone are
insufficient to fully explain the process of cell transformation into tumour cells (4). Furthermore,
research in healthy individuals has shown that some cancer driver mutations are already present
in non-transformed cells across different somatic tissues, supporting the belief that mutations in
cancer driver genes *per se* are not enough to promote tumorigenesis (5, 6).

The cellular context in which a driver mutation occurs, namely the physical interactors that are
present, can condition the impact of the mutation on the cell phenotype (7). Protein-protein

interactions (PPIs) play a crucial role in both physiological and pathological processes, such as
cell proliferation, signal transduction and apoptosis (8, 9). Consequently, mapping of cancer-related
PPIs is an ongoing effort to better understand disease and contribute to the development of
precision medicine approaches (10–14).

In the last decades, PPIs have gained growing attention and emerged as therapeutic targets for
both haematological and solid tumours (15, 16). Examples of therapeutic approaches targeting
cancer driver PPIs include inhibitors of MDM2/p53, KRAS/PDEδ and c-Myc/Max interactions (15).

Despite these advances, the systematic investigation of the relationship between driver mutations,
neighbour gene expression, and patient survival remains largely unexplored. This study aims to
address this gap by analysing the association between driver mutations and the expression of their
neighbour genes in a pan-cancer cohort. We hypothesize that the expression of neighbour genes
can modulate the phenotypic effects of driver mutations and influence patient survival. If, for
example, the driver protein is a protein kinase, the effects of the mutation will be different depending
on the set of neighbour phosphorylation targets that are expressed in the cell type where the
mutation occurred. Alternatively, neighbours can influence the stability of the driver, and
consequently its steady-state abundance, as is the case of the PPIA stabilising effect on the
oncogene NRF2 (17). Moreover, we hypothesize that when such neighbour effects exist, we can
identify them through correlations between neighbour expression and driver mutation status in
cohorts of cancer patients. By integrating genomic, transcriptomic and interactomic data, we aim
to identify novel therapeutic targets and gain a deeper understanding of the complex interplay
between driver mutations and their protein interaction context in cancer.

69
70
71   **Results**

72
Rationale and computational workflow

In this study, we sought to find physical interactors (hereafter called "neighbours") of cancer driver
proteins that influence how driver mutations cause cancer. We reasoned that if a neighbour affects
the phenotypic consequences of driver mutations in a cancer cell, it should be possible to identify
a correlation between the neighbour gene expression and the presence of driver mutations in
cancer patient data. Pharmacologically targeting these neighbours might lead to an attenuation or
even reversion of the driver mutation effects. With this in mind, we designed a computational
workflow (Figure 1) to analyse cancer patient genomic and transcriptomic data (from the TCGA
Pan-Cancer cohort (3, 18) and evaluate Cancer Driver PPIs (known cancer driver genes extracted
from NCG (19) and PPIs from Biogrid (20), String (21), HuRI (22), Omnipath (23) and APID (24),
global network properties available in supplementary table S1).

The expression of neighbours can vary between tissues and between individuals. Therefore, we
followed two approaches to detect neighbours associated with driver mutation status: one
comparing neighbour expression between cancer types (BCT) and the other comparing neighbour
expression within cancer types (WCT) (Figure 1 A). Both methods considered the possible effect
of the tumour mutational burden (TMB) in each individual or cancer type. But the detection of these
associations does not imply a causal relationship. Three types of causal relationships can originate
these associations: 1) the driver mutation induces changes in neighbour expression; 2) the
neighbour expression influences the phenotypic outcome of the driver mutation; or 3) a third
variable influences simultaneously the occurrence of mutations in the driver and the neighbour
expression. We aim to find candidate neighbours with significant associations with cancer drivers
due to the second type of causal relationship. The third type of causal relationship can only be
excluded with experimental tests, but the first type can be excluded with 1) available cancer patient
data and 2) known signalling pathways and transcriptional regulations (Figure 1 B). First, using a

cohort of patients in which gene expression was measured in both tumour and adjacent normal
tissue, we could infer the effect of the driver mutation by quantifying the difference in the neighbour
expression between tumour and normal tissues (tumour/normal tissue comparison). If the
difference is the same in patients with and without driver mutations, the mutation does not have a
significant impact on the neighbour expression. Additionally, we identified short directed pathways
connecting driver to neighbour genes. These directed pathways were extracted from known
signalling networks (available in Omnipath (23)). The final step of each directed pathway was
always a transcriptional regulatory interaction targeting the neighbour gene (extracted from the
DoRothEA resource (25)). The existence of these short, directed pathways suggests that the driver
can influence the neighbour expression. Indeed, we determined that driver neighbour pairs
connected by directed pathways of length 1 or 2 were significantly more likely to exhibit a significant
difference in the tumour/normal tissue comparison (Supplementary table S2). Neighbour-driver
PPIs that 1) were not significant in the tumour/normal tissue comparison and 2) were not connected
by directed pathways of length 1 or 2, were kept for further analysis. Due to the very large number
of neighbour-driver PPIs under evaluation (401402 in the WCT analysis and 320055 in the BCT
analysis) it is very difficult to apply a statistical cut-off that ensures a low fraction of false positives
without drastically losing statistical power (meaning that many false negatives are produced).
Therefore, we focused on identifying neighbours that were involved in a larger number of significant
associations with drivers than expected by chance (Figure 1 C). Lastly, we compared our results
with a previously available survival analysis (26, 27) to determine whether the identified neighbours
might have an impact on disease progression and severity (Figure 1 D).

Neighbour-Driver associations are detectable between and within cancer types

We detected BCT associations by computing the Spearman rank correlation coefficient between
the fraction of individuals with mutations in a driver (normalised by the average TMB of each cancer
type) and the mean expression of the driver neighbour for each cancer type. Figure 2A shows one
example of a positive BCT association between neighbour expression and driver mutation. This
correlation and the associated statistical test were computed for all driver-neighbour PPIs. To
detect WCT associations, we applied linear models to predict neighbour expression using driver
mutation status, TMB and cancer types as predictor variables. Figure 2B shows one example of a
positive WCT association between neighbour expression and driver mutation. When model
coefficients for driver mutation status are significantly different from zero, we can infer that
neighbour expression is consistently distinct between individuals with and without driver mutation
for multiple cancer types. The obtained distribution of p-values in both analyses showed an
enrichment for lower p-values, contrasting with the uniform distribution obtained when the same
tests were applied with permuted expression datasets (Figure 2C and 2D). This departure from the
uniform distribution confirms the existence of driver-neighbour associations between neighbour
expression and driver mutation status, both across and within cancer types. Considering significant
associations with a p<0.05, we obtained 30762 BCT and 30361 WCT associations.

Although both analyses took into account TMB, results could be biased by complex differences in
mutational signatures across subgroups of samples. To determine whether our findings were being
influenced by such effects, we employed the Coselens method (28) to analyse 10 WCT
associations (chosen from the top-ranked associations according to statistical significance,
avoiding associations with repeated drivers or neighbours). This method compares the mutation
frequency of a gene across two sample groups, taking into account a fitted model of the mutational
profiles in each sample group. In our case, the two sample groups were defined by their level of
neighbour gene expression. For all 10 associations, we were able to detect evidence of significant
conditional selection of driver mutations by neighbour expression in at least one cancer type
(Supplementary data file 1). Moreover, the promotion or inhibition of driver mutations was
concordant with the sign of the WCT associations.

On the other hand, BCT associations could be due to a match between the neighbour expression
tissue specificity and the cancer types with higher (or lower) driver mutation frequency. In that

scenario, the neighbour could be associated with the driver mutation frequency, but not necessarily
involved in the tissue specific factors that condition the effects of driver mutations. These tissue
specificity induced BCT associations would be due to a small number of cancer types with high (or
low) driver mutation frequency and very high (or very low) neighbour expression. To rule out the
possibility that these phenomena were prevalent in our dataset, we partitioned the cancer types for
each driver-neighbour pair into two subsets based on driver mutation frequency, and calculated the
BCT correlations for each subset separately. For pairs with a significant BCT association, the signs
of the low-mutation (or high-mutation) BCT correlation were in agreement with the correlation using
all cancer types in more than 80% of the associations (Supplementary table S3). This sign
concordance was lower for non-significant associations (around 60%). This suggests that
significant BCT associations are supported by robust trends that are not only due to a few cancer
types that happen to coincide with the tissue specific expression of neighbour genes. Additionally,
we checked the tissue specificity of neighbour genes (Supplementary figure S1) and found that
neighbours enriched in BCT associations tend to have a lower tissue specificity index, meaning
that they are more broadly expressed across numerous tissues. This observation further supports
the notion that our of significant BCT associations are not due to a spurious neighbour tissue
specificity effect.

The list of driver-neighbour PPIs evaluated included 13748 (for BCT) and 14236 (for WCT)
neighbours, but only around 55% of them were involved in significant BCT associations and 60%
in WCT associations (Table 1). We compared these percentages with what would be expected by
chance if the same number of significant associations were attributed randomly among all driver-
neighbour PPIs. While the number of neighbours with significant WCT associations was close to
what would be expected by chance, the number with significant BCT associations was significantly
lower than expected. Regarding driver participation in significant associations, the fraction was
higher than for neighbours (above 80% for both BCT and WCT) but significantly lower than
expected by chance for both types of association (Table 2). These deviations from randomness
suggest that some drivers (and some neighbours in the case of BCT associations) are more likely
to be involved in significant driver-neighbour associations.

We then evaluated if some neighbours or drivers were involved in more associations than expected
by chance. For that, we repeated all analyses with 100 datasets where expression profiles were
randomly permuted between all patients. In this way, the eventual associations between neighbour
expression and driver mutation status were abolished, and the resulting distribution of the number
of associations per neighbour (or driver) was used to evaluate a null model random expectation.
For each neighbour and driver, the number of significant associations was transformed into a z-
score. Similar z-scores were computed for the permuted datasets, allowing us to set a z-score
threshold above which a false discovery rate lower than 0.1 was ensured. This resulted in the
identification of 510 and 257 neighbours enriched in BCT and WCT driver associations (DA),
respectively (Table 1). For drivers, 403 and 42 were enriched in BCT and WCT neighbour
associations (NA), respectively (Table 2). Among the cancer drivers included in our study, 591 are
considered well known canonical drivers, while the remaining 2456 drivers have been identified in
genomic studies of cancer patient cohorts (collected and annotated in the NCG resource (19)) but
the experimental evidence of their driver role is less well established. The subsets of drivers
enriched in BCT and WCT NA have significantly higher proportions of canonical drivers than
expected (Table 2). This suggests that these driver-neighbour associations are characteristic of
true cancer drivers. Some of the canonical drivers are labelled as oncogenes or tumour suppressor
genes. Oncogenes become overly active due to driver mutations, while mutations in tumour
suppressor genes result in the loss of their function. For drivers enriched in WCT NA, both
oncogenes and tumour suppressor genes tend to have higher z-scores than drivers with unknown
role, while for BCT NA, only tumour suppressor genes tend to have higher z-scores (Supplementary
figure S2).

Neighbours enriched in BCT associations tend to have positive sign-coherent effects

We classified enriched neighbours as "positive" (when at least 80% of their driver associations are positive), "negative" (when at least 80% of their driver associations are negative) or "mixed" (remaining cases). A higher expression of positive neighbours is associated with the tumorigenic effect of driver mutations. So, we can infer that these neighbours may have a cellular function that is intrinsically oncogenic. Conversely, negative neighbours should have tumour suppressor roles. Mixed neighbours have a context-dependent role according to which drivers are mutated. Neighbours enriched in BCT DA show a different pattern of association signs when compared with neighbours enriched in WCT DA (Figures 3A and 3B). A high proportion of neighbours enriched in BCT DA had the majority of their associations with a coherent sign, either positive or negative, while neighbours enriched in WCT DA tend to have a mixture of positive and negative associations.

Part of this difference in association sign pattern is probably due to differences in the structure of correlations between driver mutation patterns. Two drivers that tend to be simultaneously mutated in the same individuals will also be positively correlated in their cancer type mutation frequency profile. On the other hand, drivers with a positive correlation of cancer type mutation frequency are not necessarily positively correlated at the individual level. This is expected for drivers that are part of the same pathway and have a mutual-exclusivity mutation pattern, which should be positively correlated at the cancer type level but negatively correlated at the individual level. The impact of this different correlation structure is apparent in the distribution of the fraction of positive associations obtained in the analysis of randomly permuted expression datasets (Figures 3A and 3B). Random distributions were matched with observed data by the number of significant associations per neighbour. Compared with these random distributions, neighbours enriched in BCT DA have more positive associations and fewer negative or mixed patterns than expected. Neighbours enriched in WCT DA have a sign distribution that is similar to the random expectation. The distribution of the fraction of positive associations was also performed for drivers enriched in BCT and WCT NA (Figures 3C and 3D). For both types of analysis, drivers enriched in NA tend to have a mixed pattern of associations. Drivers enriched in BCT NA have a shift towards higher fractions of positive associations when compared with the random expectation (Figure 3C). This is in agreement with the predominance of neighbours enriched in positive BCT DA. Drivers enriched in WCT NA tend to have a mixed signal pattern, with slightly more negative than positive associations (Figure 3D). Drivers enriched in WCT NA with coherently positive or negative associations are less frequent than randomly expected.

Neighbours enriched with driver associations tend to have cancer survival impacts that are coherent with the association signs

In order to verify whether associations between neighbour expression and driver mutation status had a significant impact on the clinical outcome, we assessed the impact of neighbour expression on overall survival across 21 cancer types according to a Human Pathology Atlas resource (26, 27). First, we observed that neighbours enriched BCT and WCT DA had a significantly higher proportion of neighbours with at least two prognostic survival associations (SA) in two different cancer types (Table 1). Next, we found that, in both BCT and WCT analyses, neighbours with an unfavourable prognostic SA in more cancer types tended to have more positive driver associations (Figure 4A). This is a sign-coherent trend, as positive neighbour-driver associations indicate that high neighbour expression promotes the tumorigenic effects of driver mutations and, consequently, poorer survival outcomes. Conversely, neighbours with more favourable prognostic SA also tended to have more negative BCT DA (Figure 4B). However, a similar trend was not observed for negative WCT DA. Finally, we classified neighbours into four classes according to their SA: 1) neighbours that do not have any SA, 2) neighbours that only have favourable SA, 3) neighbours that only have unfavourable SA and 4) neighbours that have both favourable and unfavourable SA. We then tested if these four classes were more or less frequently found in sets of neighbours enriched in positive, negative or mixed DA, both for BCT (Figure 4C) and WCT analyses. Neighbours enriched in positive BCT DA were significantly enriched in unfavourable SA, and showed lower than expected proportions of neighbours with favourable or no SA (Figure 4C). Coherently, neighbours

enriched in negative BCT DA were significantly enriched in favourable SA and had lower than
expected proportions of neighbours with unfavourable or no SA (Figure 4C). For the WCT type of
associations, only neighbours enriched in mixed WCT DA were enriched in neighbours with both
favourable and unfavourable SA (Figure 4D). Globally, these results validate the impact of driver-
neighbour associations in disease outcome. Although neighbours with mixed WCT DA can have a
relevant role in conditioning the effect of several driver mutations, their context dependent effects
turn them less attractive as candidate drug targets. Therefore, we selected two lists of neighbours
for further exploration as potential drug targets: 1) 119 neighbours enriched in positive BCT DA
with at least two unfavourable SA (Supplementary file 2) and 2) 25 neighbours enriched in negative
BCT DA with at least two favourable SA (Supplementary file 3).

Most neighbours simultaneously enriched in driver and survival associations have evidence
of causing effects in cancer cells that are coherent with driver association sign

The candidate neighbour sets were selected through significant statistical associations with driver
mutation status and overall survival. Both associations were evaluated with observational data from
a large cancer patient cohort. Therefore, these associations alone are not enough to prove that
these neighbours have any causal effect on cancer cells. A fraction of these neighbours are
themselves cancer drivers (35 out of 119 positively associated neighbours and 6 out of 25
negatively associated neighbours). Being cancer drivers implies that changes in their activities
induce causal effects on cancer cells. Surprisingly, three tumour suppressor genes (*BUB1B*,
FANCA and *POLD1*) are among the neighbours enriched in positive associations with drivers and
decreased survival in multiple cancer types. The pro-tumorigenic effect of high expression of some
of these genes has been previously reported (29, 30). These genes are involved in DNA repair
pathways. Loss of function mutations in DNA repair genes may lead to accumulation of mutations
and chromosomal instability, promoting cancer development and explaining their tumour
suppressor roles. But when they are not mutated, higher expression may be associated with more
aggressive tumours, where these pathways may be activated to prevent excessive chromosomal
instability that could lead to cell death (31). Higher DNA repair capabilities may also increase the
resistance to some therapies, which may cause decreased survival times (32).

For neighbours not previously identified as cancer drivers, we conducted a bibliographic search to
find experimental evidence (functional tests using overexpression, knockdown, or knockout in
cancer cell lines) of causal effects that were coherent with the dominant sign of driver and survival
associations. We successfully identified such experimental evidence for 88% (74 out of 84,
Supplementary file 2) of positively associated neighbours and for 37% (7 out of 19, Supplementary
file 3) of negatively associated neighbours.

We note that, for some neighbours, we were able to identify evidence with opposing effects. As an
example, *PTGR2*, a neighbour enriched in negative driver associations, has been shown to
promote gastric cancer (33) while having favourable SA in lung and renal cancer. These dual roles
could be justified by a mixture of positive and negative driver associations. This is not the case for
PTGR2, for which only negative driver associations were detected. This suggests that *PTGR2* may
have other PPIs with cancer-related proteins that explain its pro-tumorigenic role in gastric cancer.

Finally, we searched the Open Target platform (34) and found that 28 of our selected neighbours
(25 neighbours enriched in positive driver and survival associations and 3 enriched in negative
driver and survival associations, Supplementary files 2 and 3) are known drug targets. 19 out of
these 28 were not previously recognized cancer drivers. 18 are targeted by drugs developed for
cancer treatment, confirming in this way the therapeutic potential of neighbours enriched in driver
associations. Interestingly, 10 neighbours (*CX3CL1, E2F1, LAMA3, MIF, RPL28, RPL35, RPL7A,*
RPLP0, RPS15A, RPS21) were targeted only by drugs for non-cancer related diseases, suggesting
an opportunity for drug repurposing.

Discussion

This study investigated the relationship between driver mutations, neighbour gene expression, and patient survival in cancer. We were able to identify significant associations between neighbour expression and driver mutation status that were not explained by a direct mutation effect. These associations were detectable both in comparisons between and within cancer types. Moreover, we could identify neighbours that had more driver associations than expected by chance. Many of these enriched neighbours tended to have sign-coherent associations with driver mutation status, either mostly positive or mostly negative associations. One key finding was the significant correlation between the number of driver associations and the number of sign-coherent survival associations for each neighbour. This correlation is particularly strong for neighbours enriched in BCT driver associations. Neighbours enriched in driver associations were significantly more likely to have multiple survival associations, particularly for neighbours with positive associations, where high neighbour expression correlated with both increased driver mutations and poorer survival.

The fact that direct neighbours of cancer drivers are associated with patient overall survival is not surprising, as it has been observed that physical interactors of cancer drivers have an increased probability of being cancer drivers themselves (13). There are also previously known examples of neighbours that condition the effects of driver mutations, such as EGFR and c-CBL (35) or TP53 and MDM2 (36). In both examples, the neighbour proteins are E3 ubiquitin ligases that control the degradation rate of the driver proteins, therefore regulating their abundance and activity. The novelty in this work resides in the finding that such neighbour effects can leave a statistical signature in genomic and transcriptomic datasets of cancer patient cohorts. This signature allows for the systematic discovery of neighbours with significant driver associations that can be relevant prognostic biomarkers or therapeutic targets.

Not all neighbours that condition driver mutations effects are discoverable in this way. For the statistical driver association to be detectable, it is necessary that the neighbour has a natural expression variability, either between tissues or between individuals. On the other hand, similar statistical associations with driver mutation status and with patient overall survival can also exist for other proteins that are not neighbours of cancer drivers. The advantage of detecting an association for a neighbour protein is that the PPI supports the existence of a molecular mechanism explaining the statistical association.

Our study identified 119 neighbours enriched in positive driver associations with multiple unfavourable survival associations and 25 neighbours enriched in negative driver associations with multiple favourable survival associations. These neighbours represent promising candidates for further investigation as potential drug targets. In fact, 18 of them are already known drug targets for cancer therapy. Additionally, a literature search revealed experimental evidence supporting the causal roles of a substantial proportion of these neighbours.

We opted to focus on neighbours that have most of their driver associations with the same sign. Their role in cancer development is less context-dependent, facilitating the detection of significant survival associations. Neighbours with a mixture of both positive and negative driver associations are not less valid. In fact, they can be fundamental to fully understand the role of drivers that behave both as tumour suppressors or as oncogenes, depending on the cellular context. But to untangle these context-dependent effects, either more data or experimental approaches are needed.

While the statistical associations observed in this study provide compelling evidence for the role of proteins that interact directly with cancer driver proteins, further experimental validation is essential to establish causality. Future research should focus on characterising the functional roles of the identified neighbours and exploring their potential as therapeutic targets. This could involve in vitro and in vivo studies to assess the impact of modulating neighbour gene expression on cancer cell growth, survival, and response to therapy. The impact of neighbour modulation should be stronger

when the cancer cells harbour mutations in drivers interacting with that neighbour. The confirmation of these neighbours as drug targets, especially effective for subsets of patients carrying particular driver mutations, would expand the toolkit of targeted therapies.

It is important to acknowledge certain limitations of this study. The analysis relied on observational data from the TCGA pan-cancer cohort. The main advantage of using observational data is that measurements are made in real patient tissues instead of animal or cell models. However, it is not possible to control all confounding variables in observational studies. This limitation implies that we cannot control for a situation in which some individuals were exposed to unknown factors that could affect simultaneously driver mutations and neighbour expression and, therefore, generate an artificial association between driver mutation and neighbour expression.

Using observational data also implies that the number of cases with a particular combination of attributes is not controlled by the researchers. This limits the application of more complex statistical models. In the linear models used to detect associations between driver mutation status and neighbour expression, it would be interesting to include interaction terms between mutation status and cancer type. This would allow us to detect more driver-neighbour associations that would be conditional on the cancer type. But given that for most drivers there is only a reduced number of mutated individuals among the less prevalent cancer types, the statistical power is not sufficient for a precise estimation of the referred interaction coefficients.

RNA-seq data was used to estimate the abundance of neighbour proteins in cancer cells. Correlation between mRNA and protein levels can be rather low, especially when comparing values of different genes in the same tissue or cell type (37). However, when comparing values for the same gene across different samples, mRNA-protein correlations are much higher (38). In the present study, mRNA expression values being compared are always for the same neighbour gene across different individuals and cancer types. Therefore, we expect that most of these changes in gene expression reflect similar changes in protein levels.

Other limitations arise due to tumour heterogeneity and the diverse composition of the tumour microenvironment. RNA-seq was performed on the bulk of cells collected from the tumour tissue. We cannot be sure that all tumour cells carry the same mutations and have similar gene expression profiles. And part of the expression signal may come from non-tumour cells that are infiltrated in the tumour tissue. This issue could be solved by utilizing single-cell genomic sequencing and mRNA sequencing for each tumour sample. In this way, we could identify neighbour gene expression in the same cells that carried a mutation in the interacting driver. Besides the additional layer of complexity in the data analysis, we are not aware of the availability of this type of data for large cohorts of cancer patients.

The PPI datasets used might have their own limitations. It is possible that interaction data is incomplete, meaning that drivers may have unknown interactions with other proteins. Additionally, some interactions are only established with mutated drivers (39) and are not included in the interaction databases used here. Although our focus was on protein physical interactions, relevant interactions may be established with other types of molecules, such as non-coding RNAs. Finally, we only considered binary interactions. Some context-dependent effects of driver mutations may be explained only by higher-order interactions, involving more than one interactor simultaneously.

Despite these limitations, the findings of this study provide valuable insights into the complex interplay between drivers and their neighbours. It also demonstrates a new approach to analyse genomic and transcriptomic data from cancer patient cohorts that uncovers neighbours enriched in driver associations as potential therapeutic targets and prognostic biomarkers.

Materials and Methods

Cancer drivers

A list of canonical and candidate cancer driver genes was retrieved from the Network of Cancer Genes & Healthy Drivers 7.1 (19). This list was used to search for protein physical interactors and to detect driver-neighbour associations (meaning that neighbour gene expression was statistically associated with driver mutation status). To evaluate if neighbours enriched in driver associations had any known cancer driver role, we retrieved the lists of cancer driver genes available in COSMIC Canger Gene Census (40), Intogen (2) and OncoKB (41, 42). Additionally, for the final lists of neighbours enriched in driver associations and with multiple survival associations, we searched the Disgenet database (43) for evidence of association with neoplastic diseases with association type "CausalMutation".

Protein physical interaction data

Human PPI were retrieved from five sources: String (only PPI with a confidence score equal or greater than 0.4) (21), Biogrid (20), HuRI (22), Omnipath (23) and APID (24). Interactions involving complexes were expanded to include all possible pairwise interactions. Protein physical interactors (neighbours) of each cancer driver were extracted from all five interactome sources. General characteristics if these networks are shown in Supplementary table 1.

Cancer patient data

Cancer patient data from the TCGA Pan-Cancer cohort was accessed through the UCSC Xena online tool (https://pancanatlas.xenahubs.net) (44). In particular, gene-level non-silent somatic mutation data (18), batch effects normalised mRNA data, sample phenotype data and patient clinical data (45) were retrieved. Patients with more than 72 mutated cancer driver genes were discarded as they were considered mild outliers using the Tukey's fence method (46). Missing values in the gene expression data were imputed with the minimum expression value observed for the respective gene across all patients.

Tumour mutational burden

Tumour mutational burden (TMB) is a measure of the total number of somatic mutations found in a tumour specimen. It is commonly used as a biomarker useful to predict the outcome of certain immunotherapies. TMB can be characterized experimentally using different methodologies that look for nonsynonymous mutations in coding regions and oncogenic drivers, including whole exome sequencing (WES) (47). In our work, we calculated the TMB for each tumour specimen (patient) using a dataset which includes all somatic mutation calls (SNPs and INDELs). First, this dataset was filtered to remove silent mutations—we considered as silent mutations mutations in noncoding regions—and then we counted all nonsynonymous mutations for each sample. For the BCT analysis we computed the mean mutational burden per cancer type. TMB values, both for individual patients and cancer type averages, were converted with a log_{10} transformation.

Comparison of paired tumour and normal samples

596 patients in the pan-cancer cohort had paired tumour and normal tissue samples. For each of these patients, we computed the difference in gene expression between the tumour sample and the matching normal tissue. For each driver-neighbour PPI, we fitted a linear regression model with fixed effects (Equation 1).

$$TNdiff_i = \beta_1 Mut_i + \beta_2 CT_i \qquad \text{(Equation 1)}$$

The dependent variable was the tumour-normal neighbour gene expression difference ($TNdiff_i$). The independent variables were 1) the mutation status of the driver (Mut_i) and 2) the cancer type (CT_i) of each patient i. Only data from cancer types that had at least one patient with a positive

440 mutation status in at least 2 cancer types were included in the model. A significant (p<0.05)
441 regression coefficient of the driver mutation status (β_i) was considered as evidence that the driver
442 mutation induced a change in the expression of the neighbour. These driver-neighbour PPIs were
443 excluded from further analyses.

Pathway analysis

Driver mutations might have an impact on neighbour gene expression if the driver protein directly
regulates signalling pathways and transcription factors (TF) that target the neighbour gene. To
exclude this type of association, we built a regulatory network by combining two directed networks:
a network of signalling pathways from Omnipath (23) with a TF-target gene network from
DoRothEA (25) (Supplementary table S1). For each driver-neighbour physical PPI, we computed
the shortest path distance between the driver and the neighbour so that the final interaction was
between a TF and the neighbour gene. A distance of 1 means the driver is a direct TF of the
neighbour. Driver-neighbour pairs with a shortest distance of 1 or 2 were more likely to have a
significant difference in the tumour/normal tissue comparison, and thus were excluded from the
analysis.

Statistical associations between neighbour expression and driver mutation status

We used data from 7317 patients (which passed the driver mutation outlier filter and had mutation
and gene expression data for tumour samples) to compute, for each driver-neighbour PPI, the
Spearman correlation coefficient between 1) the average neighbour expression per cancer type
and 2) the fraction of patients with mutated driver divided by the average TMB per cancer type. In
this way we can avoid detecting false correlations where the higher driver mutation frequency in
some cancer types is just due to a higher TMB of the individuals with those cancer types. A
statistical test was applied to each driver-neighbour PPI to evaluate if the computed coefficient was
significantly different from 0. Previous to the analysis, we conducted a power analysis with
simulated data and concluded that to achieve a statistical power of 0.7, the correlations should only
be computed when there are at least 10 cancer types with a driver mutation frequency greater than
0. Driver-neighbour PPIs with a significant correlation (p<0.05) were considered statistically
associated in a between cancer type comparison (BCT).

Using the same data, we fitted, for each driver-neighbour PPI, a linear regression model with fixed
effects (Equation 2).

$NeighbourExp_i = \beta_1 Mut_i + \beta_2 CT_i + \beta_3 TMB_i$ (Equation 2)

The dependent variable was the neighbour gene expression ($NeighbourExp_i$) and the independent
variables were 1) the driver mutation status (Mut_i), 2) the cancer type (CT_i) and 3) the tumour
mutation burden (TMB_i) for each patient i. Previous to the analysis, we conducted a power analysis
with simulated data and concluded that to achieve a statistical power of 0.7, the linear model should
only be fitted when there are at least 3 cancer types with at least one patient with a driver mutation.
A significant (p<0.05) regression coefficient of the driver mutation status (β_i) was considered as
evidence that the driver-neighbour PPI was statistically associated in a comparison within cancer
types (WCT).

Significant neighbour identification through random permutations

To evaluate the statistical significance of the number of driver-neighbour associations per
neighbour protein, the computation of BCT and WCT associations was repeated 100 times with
randomly permuted gene expression datasets. Gene expression values were organized in a matrix
where each row corresponded to an individual and each column corresponded to a gene.
Randomly permuted matrices were produced by randomly shuffling the row order without changing
the column orders. In each shuffle, the number of significant driver associations (x_i) of each type

(BCT or WCT) for each neighbour *i* was recorded, allowing the computation of the average ($\langle x_i^r \rangle$)
and standard deviation ($sd(x_i^r)$) of this number according to a random null model. These
parameters were used to compute z-scores (Equation 3) for the original observations and for each
of the 100 analyses with randomly permuted data.

$z_i = \frac{x_i - \langle x_i^r \rangle}{sd(x_i^r)}$ (Equation 3)

For any given z-score threshold (t) we computed the False Discovery Rate (FDR, Equation 4) using
the average number of neighbours with a z-score greater than the threshold in the 100 analyses
with the randomly permuted data ($\langle \#(z_i^r > t) \rangle$) as an estimate of the expected false positives.

$FDR(t) = \frac{\langle \#(z_i^r > t) \rangle}{\langle \#(z_i > t) \rangle}$ (Equation 4)

Equation 4 estimates FDR by excess, as the estimated number of false positives is done with the
total number of neighbours instead of using the unknown proportion of neighbours that follows the
null hypothesis (H_0: the neighbour does not have more driver associations than expected by
chance). For each type of driver association (BCT or WCT), we determined the lowest value of t
for which $FDR(t) < 0.1$.

Survival analysis

Cancer prognostic data was downloaded from the Human Protein Atlas, containing Log-rank p
values for Kaplan-Meier analysis of correlation between mRNA expression level and patient
survival in prognostic cohorts based on 21 cancers from TCGA and validation cohorts for 10 of the
cancers. Details of the analysis are described in previous publications (26, 27). According to the
authors, genes were designated as prognostic genes if they had log-rank p values less than 0.001.
Benjamini and Hochberg procedure (48) was applied to the 219519 p-values available in the
dataset. The p-value threshold of 0.001 corresponded to an FDR of 0.013. 22640 gene-cancer type
prognostic associations were considered using this threshold.

Tissue specificity

We used the Tau Tissue Specificity Index (49) to characterize the tissue specificity of genes across
different tissues. The tau index (τ), is defined as (Equation 5)

$\tau = \frac{\sum_{i=1}^{n}(1 - \hat{x}_i)}{n-1}; \hat{x}_i = \frac{x_i}{max(x_i)}, 1 \leq i \leq n$ (Equation 5)

Where n is the number of tissues and x_i is the expression of the gene in tissue i. We applied the
tau index to the Human Protein Atlas (HPA) Consensus RNA dataset. This dataset summarizes
HPA (50) and GTEx (51) transcriptomics data across 50 tissues using a normalized TPM value
(nTPM). To eliminate similar tissues that could impact the tissue specificity index calculation (e.g.
the brain is represented with 8 different regions), we processed the dataset into 41 distinct organs
by taking the median nTPM of tissues in the same organ. Finally, nTPM values were log_2
transformed, and an expression cutoff was applied: all genes with expression below 1 nTPM were
removed.

Code availability

523 All the code used to perform the analyses presented in this manuscript is available at
524 https://github.com/GamaPintoLab/driver_neighbours. All the data used in this work was obtained

from public sources. For reproducibility purposes, raw data files used are available at
https://doi.org/10.5281/zenodo.14284407.

Acknowledgments

Work supported by UID/04046/2025 – Instituto de Biosistemas & Ciências Integrativas Centre grant from FCT, Portugal.

References

1. D. Stehelin, H. E. Varmus, J. M. Bishop, P. K. Vogt, DNA related to the transforming gene(s) of avian sarcoma viruses is present in normal avian DNA. *Nature* **260**, 170–173 (1976).

2. F. Martínez-Jiménez, *et al.*, A compendium of mutational cancer driver genes. *Nat Rev Cancer* 1–18 (2020). https://doi.org/10.1038/s41568-020-0290-x.

3. T. I. P.-C. A. of W. G. Consortium, et al., Pan-cancer analysis of whole genomes. *Nature* **578**, 82–93 (2018).

4. B. Stransky, S. J. de Souza, Modeling tumor evolutionary dynamics. *Front. Physiol.* **3**, 480 (2013).

5. H. Lee-Six, et al., The landscape of somatic mutation in normal colorectal epithelial cells. Nature **574**, 532–537 (2019).

6. I. Martincorena, et al., Tumor evolution. High burden and pervasive positive selection of somatic mutations in normal human skin. *Sci. (N. York, NY)* **348**, 880–6 (2015).

7. G. Schneider, M. Schmidt-Supprian, R. Rad, D. Saur, Tissue-specific tumorigenesis: context matters. Nat Rev Cancer **17**, 239–253 (2017).

8. M. Vidal, M. E. Cusick, A.-L. Barabási, Interactome Networks and Human Disease. *Cell* **144**, 986–998 (2011).

9. M. E. Cusick, N. Klitgord, M. Vidal, D. E. Hill, Interactome: gateway into systems biology. *Hum. Mol. Genet.* **14**, R171–R181 (2005).

10. M. Kim, et al., A protein interaction landscape of breast cancer. *Science* **374**, eabf3066 (2021).

11. D. L. Swaney, et al., A protein network map of head and neck cancer reveals PIK3CA mutant drug sensitivity. Science 374, eabf2911 (2021).

12. M. S. Tabar, H. Francis, D. Yeo, C. G. Bailey, J. E. J. Rasko, Mapping oncogenic protein interactions for precision medicine. *Int. J. Cancer* **151**, 7–19 (2022).

13. M. Bouhaddou, et al., Mapping the protein–protein and genetic interactions of cancer to guide precision medicine. *Curr. Opin. Genet. Dev.* **54**, 110–117 (2019).

14. D. Módos, et al., Neighbours of cancer-related proteins have key influence on pathogenesis and could increase the drug target space for anticancer therapies. *npj Systems Biology and Applications* 1–13 (2017). https://doi.org/10.1038/s41540-017-0003-6.

15. H. Lu, et al., Recent advances in the development of protein–protein interactions modulators: mechanisms and clinical trials. *Signal Transduct. Target. Ther.* **5**, 213 (2020).

16. A. W. White, A. D. Westwell, G. Brahemi, Protein–protein interactions as targets for small-molecule therapeutics in cancer. *Expert Rev. Mol. Med.* **10**, e8 (2008).

17. W. Lu, et al., PPIA dictates NRF2 stability to promote lung cancer progression. *Nat.*

Commun. **15**, 4703 (2024).

18. K. Ellrott, et al., Scalable Open Science Approach for Mutation Calling of Tumor Exomes
Using Multiple Genomic Pipelines. *Cell Syst.* **6**, 271-281.e7 (2018).

19. L. Dressler, et al., Comparative assessment of genes driving cancer and somatic evolution in
non-cancer tissues: an update of the Network of Cancer Genes (NCG) resource. *Genome Biol*
23, 35 (2022).

20. R. Oughtred, et al., The BioGRID database: A comprehensive biomedical resource of
curated protein, genetic, and chemical interactions. *Protein Sci. : A Publ. Protein Soc.* **30**, 187–
200 (2021).

21. D. Szklarczyk, et al., The STRING database in 2023: protein–protein association networks
and functional enrichment analyses for any sequenced genome of interest. *Nucleic Acids Res.*
51, D638–D646 (2023).

22. K. Luck, et al., A reference map of the human binary protein interactome. Nature **580**, 402–
408 (2020).

23. D. Türei, et al., Integrated intra- and intercellular signaling knowledge for multicellular omics
analysis. *Mol. Syst. Biol.* **17**, MSB20209923 (2021).

24. D. Alonso-López, et al., APID database: redefining protein–protein interaction experimental
evidences and binary interactomes. *Database: J. Biol. Databases Curation* **2019**, baz005 (2019).

25. L. Garcia-Alonso, C. H. Holland, M. M. Ibrahim, D. Turei, J. Saez-Rodriguez, Benchmark and
integration of resources for the estimation of human transcription factor activities. *Genome Res*
29, 1363–1375 (2019).

26. M. Yuan, et al., The Human Pathology Atlas for deciphering the prognostic features of human
cancers. *eBioMedicine* **111**, 105495 (2024).

27. M. Uhlen, et al., A pathology atlas of the human cancer transcriptome. Science **357** (2017).

28. J. Iranzo, G. Gruenhagen, J. Calle-Espinosa, E. V. Koonin, Pervasive conditional selection of
driver mutations and modular epistasis networks in cancer. *Cell Rep.* **40**, 111272 (2022).

29. X. Fu, et al., Overexpression of BUB1B contributes to progression of prostate cancer and
predicts poor outcome in patients with prostate cancer. *OncoTargets Ther.* **9**, 2211–2220 (2016).

30. S. Bravo-Navas, L. Yáñez, Í. Romón, A. C. Pipaón, Elevated FANCA expression determines
a worse prognosis in chronic lymphocytic leukemia and interferes with p53 function. *FASEB J.*
33, 10477–10489 (2019).

31. M. J. O'Connor, Targeting the DNA Damage Response in Cancer. *Mol. Cell* **60**, 547–560
(2015).

32. K. Kiwerska, K. Szyfter, DNA repair in cancer initiation, progression, and therapy—a double-
edged sword. *J. Appl. Genet.* **60**, 329–334 (2019).

33. E. Y.-C. Chang, et al., Prostaglandin Reductase 2 Modulates ROS-Mediated Cell Death and
Tumor Transformation of Gastric *Cancer Cell*s and Is Associated with Higher Mortality in Gastric
Cancer Patients. *Am. J. Pathol.* **181**, 1316–1326 (2012).

34. D. Ochoa, et al., The next-generation Open Targets Platform: reimagined, redesigned,
rebuilt. Nucleic Acids Res. 51, D1353–D1359 (2023).

35. B. L. M. Crotchett, B. P. Ceresa, Knockout of c-Cbl slows EGFR endocytic trafficking and
enhances EGFR signaling despite incompletely blocking receptor ubiquitylation. *Pharmacol. Res.*
Perspect. **9**, e00756 (2021).

36. L. T. Vassilev, Small-molecule antagonists of p53-MDM2 binding: research tools and
potential therapeutics. *Cell cycle (Georget., Tex)* **3**, 419–21 (2004).

37. T. Maier, M. Güell, L. Serrano, Correlation of mRNA and protein in complex biological
samples. *FEBS Letters* **583**, 3966–3973 (2009).

38. F. Edfors, et al., Gene-specific correlation of RNA and protein levels in human cells and
tissues. *Mol Syst Biol* **12**, 883 (2016).
39. X. Mo, et al., Systematic discovery of mutation-directed neo-protein-protein interactions in
cancer. Cell **185**, 1974-1985.e12 (2022).
40. J. G. Tate, et al., COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic Acids*
Research **47**, D941–D947 (2018).
41. S. P. Suehnholz, et al., Quantifying the Expanding Landscape of Clinical Actionability for
Patients with Cancer. *Cancer Discov.* **14**, 49–65 (2023).
42. D. Chakravarty, et al., OncoKB: A Precision Oncology Knowledge Base. *JCO Precis. Oncol.*
2017, 1–16 (2017).
43. J. Piñero, et al., The DisGeNET knowledge platform for disease genomics: 2019 update.
Nucleic Acids Res. 48, D845–D855 (2019).
44. M. J. Goldman, et al., Visualizing and interpreting cancer genomics data via the Xena
platform. *Nat. Biotechnol.* **38**, 675–678 (2020).
45. J. Liu, et al., An Integrated TCGA Pan-Cancer Clinical Data Resource to Drive High-Quality
Survival Outcome Analytics. Cell **173**, 400-416.e11 (2018).
46. J. W. Tukey, *Exploratory Data Analysis.*, A.-W. P. Company, Ed. (1977).
47. D. L. Jardim, A. Goodman, D. de M. Gagliato, R. Kurzrock, The Challenges of Tumor
Mutational Burden as an Immunotherapy Biomarker. Cancer Cell 39, 154–173 (2021).
48. Y. Benjamini, Y. Hochberg, Controlling the false discovery rate—a practical and powerful
approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological)*
57, 289–300 (1995).
49. I. Yanai, et al., Genome-wide midrange transcription profiles reveal expression level
relationships in human tissue specification. *Bioinformatics* **21**, 650–659 (2004).
50. M. Uhlén, et al., Tissue-based map of the human proteome. Science **347**, 1260419–1260419
(2015).
51. T. Gte. Consortium, The GTEx Consortium atlas of genetic regulatory effects across human
tissues. Science **369**, 1318–1330 (2020).

Figures and Tables
2

3
4
Figure 1. Computational analysis workflow. (A) Using cancer patient mutation data, PPI between
cancer drivers and neighbour proteins, and gene expression data from the same patients, we
evaluated two types of statistical association between neighbour expression and driver mutation
status, one comparing between cancer types and the other comparing within cancer types. (B) To
avoid statistical associations where change in gene expression is caused by driver mutations, we
also evaluated associations between the difference in neighbour expression between tumour and
normal matching tissues and driver mutation status. Statistically significant associations found
with paired tumour-normal samples were excluded from further analyses. Additionally, we
searched signalling and transcriptional regulatory networks for directed pathways linking the
driver and the neighbour genes. If those pathways had a length of 2 or fewer interactions, the
driver-neighbour pair was also excluded from further analyses. (C) Next, we selected neighbours
that had more driver associations than expected by chance, using analyses with randomly
permuted expression datasets to estimate the random expectation. (D) Finally, we checked a
previously available survival analysis study for neighbours with a significant impact on patient
overall survival for several cancer types. These results allowed us to select neighbours that were
simultaneously enriched in driver association and had survival associations with coherent effect
signs (positive or negative association with driver mutations concurrently with a decrease or an
increase in overall survival times, respectively). Created in BioRender. Gama, M. (2025)
https://BioRender.com/o45x904.
24
25

Figure 2. Driver-neighbour associations. (A) Example of a significant BCT statistical association between neighbour (*PPIA* in the example) expression and the frequency of driver mutation (*DDX3X* in the example). Each dot represents data from a different cancer type. (B) Example of a significant WCT statistical association between neighbour (*CDK1* in the example) expression and driver (*WWTR1* in the example) mutation status. Neighbour expression values are represented as box plots according to cancer type and driver mutation status. In (A) and (B) expression values are log2(normalised TPM + 1). All significant BCT and WCT associations can be explored at https://joaoaimiranda.shinyapps.io/driver_neighbours_dashboard/, including correlation and regression coefficients and p-values. (C) Probability density distributions of Spearman correlation test p-values obtained using observed data or randomly permuted expression datasets. (D) Probability density distributions of mutation status linear model coefficient p-values obtained using observed data or randomly permuted expression datasets

Figure 3. Probability density distributions of the fraction of positive associations. (A) Probability density distributions of the fraction of positive associations for neighbours enriched in between cancer type (BCT) driver associations (DA) and comparison with the corresponding distribution in analyses using permuted expression datasets. (B) Probability density distributions of the fraction of positive associations for neighbours enriched in within cancer type (WCT) DA and comparison with the corresponding distribution in analyses using permuted expression datasets. (C) Probability density distributions of the fraction of positive associations for drivers enriched in BCT neighbour associations (NA) and comparison with the corresponding distribution in analyses using permuted expression datasets. (D) Probability density distributions of the fraction of positive associations for drivers enriched in WCT DA and comparison with the corresponding distribution in analyses using permuted expression datasets. All distributions obtained with permuted datasets were matched with observed distributions by the number of significant associations per neighbour or driver.

57
Figure 4. Association between driver and survival associations. (A) Boxplots representing the
distribution of significant driver associations (DA) with a positive sign as a function of the number
of cancer types for which a significant unfavourable survival association (SA) was detected, for
both between cancer types (BCT) and within cancer types (WCT) analyses. (B) Boxplots
representing the distribution of significant DA with a negative sign as a function of the number of
cancer types for which a significant favourable SA was detected, for both BCT and WCT
analyses. (C) Odds ratios for the association between neigbours with a particular SA pattern (only
unfavourable SA, only favourable SA, both favourable and unfavourable SA and no SA) and
neighbours enriched in BCT DA with a particular sign pattern (positive – 80% or more of positive
DA, negative – 20% or less of negative DA, mixed – between 20% and 80% of positive DA). (D)
Odds ratios for the association between neigbours with a particular SA pattern (only unfavourable
SA, only favourable SA, both favourable and unfavourable SA and no SA) and neighbours
enriched in WCT DA with a particular sign pattern (positive – 80% or more of positive DA,
negative – 20% or less of negative DA, mixed – between 20% and 80% of positive DA). In (C)
and (D), error bars represent 95% confidence intervals for the odds ratios. * represents p<0.05
and *** represents p<0.0001 for hypergeometric tests (Null hypothesis: Odds ratios are equal to
1). Odd ratios significantly greater than 1 mean that the proportion of neighbours with a particular
SA pattern within neighbours enriched in DA with a particular sign pattern is greater than
expected by chance.
77
78

79   **Table 1.** Characterisation of neighbours with significant driver associations.

Neighbour Analysis	Fraction of neighbours with significant DAs			Enrichment in DAs (FDR<0.1)		Multiple Survival Associations (SA>1)	
	Observed	Random	Z-score			Fraction	Odds Ratio (95% CI)
BCT	0.550	0.605	-20.50	**Yes**	510	0.533	2.00 (1.67-2.40)
				No	13238	0.363	
WCT	0.602	0.597	1.95	**Yes**	257	0.424	1.28 (0.99-1.66)
				No	13979	0.365	

80
81
82

83 **Table 2.** Characterisation of drivers with significant neighbour associations.
84

Driver Analysis	Fraction of drivers with significant NAs			Enrichment in NAs (FDR<0.1)		Canonical drivers	
	Observed	**Random**	**Z-score**			**Fraction**	**Odds Ratio (95% CI)**
BCT	0.840	0.899	-13.90	**Yes**	403	0.241	1.43 (1.10-1.86)
				No	2000	0.182	
WCT	0.821	0.879	-13.78	**Yes**	42	0.810	19.50 (8.80-49.09)
				No	3005	0.179	