SCAN-ACT: ADOPTIVE T CELL THERAPY TARGET DISCOVERY THROUGH SINGLE-CELL TRANSCRIPTOMICS

Stefano Testa[1,2], Aastha Pal[3], Ajay Subramanian[3], Sushama Varma[4], Jack Pengfei Tang[5],
Danielle Graham[6], Sara Arfan[7], Minggui Pan[8], Nam Q. Bui[8], Kristen N. Ganjoo[8], Sarah Dry[9],
Paul Huang[7], Matt van de Rijn[4], Wei Jiang[10], Anusha Kalbasi[3,11*], Everett J. Moding[3,11*]

[1]Department of Medicine, Stanford University, Stanford, CA, USA

[2]Current affiliation: Division of Hematology and Oncology, Beth Israel Deaconess Medical
Center, Harvard Medical School, Boston, MA, USA

[3]Department of Radiation Oncology, Stanford University, Stanford, CA, USA

[4]Department of Pathology, Stanford University, Stanford, CA, USA

[5]Department of Pediatrics, Boston Children's Hospital, Boston, MA, USA

[6]Department of Surgery, Cedars-Sinai Medical Center, Los Angeles, CA, USA

[7]Division of Molecular Pathology, The Institute of Cancer Research, London, UK

[8]Division of Oncology, Department of Medicine, Stanford University, Stanford, CA, USA

[9]Department of Pathology, University of California, Los Angeles, CA, USA

[10]JWE Science Technology, Inc., Bejing, China

[11]Stanford Cancer Institute, Stanford University, Stanford, CA, USA

*Corresponding Authors

Contact Information

Everett J. Moding, M.D., Ph.D., Assistant Professor, Department of Radiation Oncology,
Stanford Cancer Institute, 875 Blake Wilbur Drive, Stanford, CA 94305-5847. Email:
emoding@stanford.edu. Tel: 650-498-1625.
and

Anusha Kalbasi, M.D., Associate Professor, Department of Radiation Oncology, Stanford

Cancer Institute, 875 Blake Wilbur Drive, Stanford, CA. Email: akalbasi@stanford.edu. Tel: 650-

498-4796.

Summary

The FDA approval of T cell receptor-engineered T cells (TCR-T) for synovial sarcoma

demonstrates the potential for adoptive T cell therapies (ACTs) in solid tumors. However, the

paucity of tumor-specific targets without expression in normal tissues remains a major

bottleneck, especially in rare cancer subtypes. Here, we present a comprehensive

computational pipeline called SCAN-ACT that leverages single cell RNA sequencing and multi-

omics data from tumor and normal tissues to nominate and prioritize targets for both chimeric

antigen receptor (CAR)- and TCR-T cells. For surface membrane targets, SCAN-ACT proposes

target pairs for bispecific Boolean logic-gated CAR T cells. For peptide-MHC targets, SCAN-

ACT proposes intracellular peptides bound to a diverse set of human leukocyte antigens. We

applied the SCAN-ACT pipeline to soft tissue sarcoma (STS), analyzing 986,749 single cells to

identify and prioritize 395 monospecific CAR-T targets, 14,192 bispecific CAR-T targets, and

5,020 peptide-MHC targets for TCR-T cells. Selected targets were validated experimentally by

protein expression and for peptide-MHC binding. Proposed targets and target pairs reflected the

mesenchymal, neuronal, and hematopoietic ontogeny of STS. This work provides a robust data

repository along with a web-based and user-friendly set of analysis tools to accelerate ACT

development for solid tumors.

Keywords

Cellular therapy, CAR-T cells, TCR-T cells, Peptide-centric CAR-T, logic-gated CAR-T, Single

cell RNA sequencing, Soft tissue sarcoma, Immunotherapy

Introduction

Despite profound success in the treatment of hematologic malignancies, engineered adoptive T

cell therapies (ACT) have had limited success in solid tumors[1–3]. While novel engineering

strategies can address the major hurdles related to T cell function[4], including poor tumor

infiltration[5], terminal differentiation, effector dysfunction, and persistence or memory formation[6–

9], a lack of tumor-specific targets with uniformly high expression and low risk of on-target off-

tumor toxicity remains a challenge[10,11]. Advances in bispecific antigen targeting, synthetic gene

circuits, and Boolean logic-gating in engineered T cells have opened new opportunities to

improve antigen-discrimination between normal and tumor cells and increase the targetable

antigen pool[4,12,13]. However, there are no repositories of suitable target pairs for Boolean logic

CAR-T cells in solid tumors.

Prior ACT target discovery has focused on bulk RNA-sequencing and protein expression from

tumor and normal samples[3,14,15], which is confounded by non-malignant cells in the tumor

microenvironment. Similarly, analysis of cancer cell lines[16] may not reflect gene expression in

vivo or the heterogeneity of expression between individual cancer cells and across patients.

Furthermore, robust bulk transcriptomic and proteomic datasets are not available for many less

common human cancers. Compared to bulk RNA-seq and proteomics, scRNA-seq more

accurately represents the transcriptomic profile and heterogeneity of individual tumor cells

enabling the precise identification of tumor-specific genes[17,18].

Soft tissue sarcomas (STS) are rare and heterogeneous mesenchymal cancers with high rates

of metastatic disease and limited therapeutic options[19]. Although chemotherapy remains the

primary treatment for metastatic STS, most patients do not respond and the median overall

survival is less than two years[20,21]. Immune checkpoint blockers (ICBs) targeting programmed

cell death protein 1 (PD-1) and cytotoxic T-lymphocyte antigen-4 (CTLA-4) have shown promise

in specific STS histotypes[22–24], but are ineffective for most patients[25,26]. Recent studies have

shown promising efficacy for TCR-T cells in STS. Afamitresgene autoleucel (afami-cel) targeting

the cancer-testis antigen MAGE-A4 in patients with an HLA*02:01 allele achieved a 37% overall

response rate in heavily treated patients with synovial sarcoma (SS) and myxoid round cell

liposarcoma (MRCLS)[27,28], leading to the first FDA approval of a genetically engineered ACT in

solid tumors. Likewise, the most promising results for TCR-T cell products targeting NY-ESO-

1[29,30] have been reported in patients with STS[31]. However, additional targets will be critical to

expand the availability of ACTs to patients with STS that do not express MAGE-A4 and NY-

ESO-1. Furthermore, most patients lack the HLA-A*02:01 allele, which is most frequently

expressed in Europeans, raising concerns about the equity of current ACTs[32]. Taken together,

STS represent a challenging disease model with known effective ACT targets that can be used

to benchmark target identification pipelines.

Here, we leveraged single cell RNA sequencing (scRNA-seq) from patient tumors and collated

normal tissue and tumor genetic, transcriptomic, and proteomic data across public databases to

develop a comprehensive computational pipeline for discovery of ACT targets in solid tumors,

called SCAN-ACT (Single Cell Analysis Network for Adoptive Cellular therapy Targets). We

applied SCAN-ACT in STS to build CellSARCTx (Cellular SARComa Therapy), a database

describing and prioritizing over 20,000 TCR-T and CAR-T cell targets, including those for

Boolean logic-gated and peptide-centric CAR-T cells (PC-CAR-T). Proposed targets captured

the shared mesenchymal, neuronal and hematopoietic gene expression signatures of STS,

reflecting the intertwined developmental ontogeny of these three lineages from the mesoderm

and neural crest. We validated protein expression of select targets by IHC and binding of

peptides derived from intracellular proteins to HLA molecules with a novel yeast display

technology. We present two interactive web interfaces to explore and evaluate the identified

STS ACT targets (CellSARCTx) and to apply SCAN-ACT to new scRNA-Seq samples (SCAN-

ACT Custom). SCAN-ACT can be applied to individual patient samples to aid in therapy

selection or broader panels of patients to identify conserved targets and accelerate ACT

development. SCAN-ACT is freely accessible for academic use at scanact.stanford.edu.

Results

Overview of the SCAN-ACT pipeline

SCAN-ACT identifies tumor up-regulated and down-regulated genes through differential gene

expression (DGE) analysis using scRNA-seq (**Figure 1A**). Tumor up-regulated genes are then

compared against clinically tested CAR-T targets to identify novel targets with predicted low risk

of on-target off-tumor toxicity. SCAN-ACT then defines Boolean logic-gated CAR-T target pairs

by measuring tumor and normal cell/tissue co-expression patterns. Tumor-specific MHC-

restricted peptides ($pMHCs$) targetable by TCR-T or PC-CAR-T cells are identified by integrating

peptide immunogenicity and HLA presentation predictions with $pMHC$ cross-reactivity

assessment. Lastly, SCAN-ACT prioritizes targets by integrating target homogeneity with normal

tissue/cell expression, and functional, genomic, and proteomics tumor data.

scRNA-seq for cellular therapy target identification in STS

To identify ACT targets across STS histologies, we applied SCAN-ACT to newly sequenced and

publicly available scRNA-seq data[17,33] from both translocation-driven and non-translocation-

driven STS (**Table S1**). Our scRNA-seq dataset included 32 samples from 25 patients, totaling

128,122 cells after quality control (**Figure 1B, Figure S1A**, **S1B**, **S1C**, **S1E, S1G, S1I**). We

included visceral (31%) and extremity/trunk (69%) tumor samples from male (72%) and female

(28%) patients aged 22 to 77 years old. Samples included both primary (75%) and metastatic

tumors (25%) collected before (84%) or after treatment (16%). Sarcoma cells were identified

among the CD45⁻ cells (PTPRC count = 0), using inferred copy number abnormalities[34,35] and

transcriptional similarity to bulk samples of the same histology from The Cancer Genome Atlas

(TCGA)[36] (Similarity Score)[33] (**Figure S1D, S1F**, **S1H**, **S1H**, **S1J**). Within the tumor

microenvironment (TME), endothelial cells (ECs), myeloid cells (MCs), lymphoid cells (LCs),

and cancer-associated fibroblasts (CAFs) were defined using canonical markers from the

literature (**Table S2**).

Identification of genes differentially expressed by malignant cells within the STS TME

To identify tumor-specific genes, SCAN-ACT performs DGE analysis between sarcoma cells

and non-tumoral cells from the TME using Limma-voom[37], a pseudo-bulk method, and distinct[38],

a differential distribution method (**Figure 1C**). We also compared sarcoma cells with normal cell

clusters from 30 scRNA-seq datasets from the Human Protein Atlas (HPA)[39,40], Tabula Sapiens

(TS)[41], and the Human Cell Atlas (HCA)[42] covering the major normal human tissues (858,627

cells and 164 samples, **Figure 1D**). Unsupervised hierarchical clustering demonstrated that

STS were most similar to CAFs and ECs within the TME (**Figure S2A**, **S2C**) and mesenchymal

cells in normal tissues (**Figure S2B**, **S2D**), consistent with their putative mesenchymal origin. In

total, we identified 29,322 up-regulated genes and 15,812 down-regulated genes in STS cells

(**Table S3, S4**). SS and LMS had the most unique up-regulated and down-regulated genes,

respectively, while UPS, LPS, and MFS shared the most up-regulated and down-regulated

genes consistent with their biological similarity[36] (**Figure 1E**, **Figure S2E**).

There was concordance between Limma-voom and distinct for DGE analysis in the TME

(**Figure 1F**), with distinct identifying more unique differentially expressed genes. By measuring

differences in gene expression distribution between cell types, distinct identifies subtle

differences in expression[38], which can be advantageous when comparing transcriptionally

similar cells like STS and other mesenchymal cells in the TME. This approach identified a pool

of sarcoma-specific genes, including many like *FAP* or *MMP14* in UPS which are up-regulated

in STS compared to almost all normal cell types (**Figure 1G**). Notably, *FAP* has been targeted in

solid tumors due to its TME expression[43], though we found that *FAP* has intrinsic expression in

STS cells, particularly LPS, MFS, and UPS, consistent with IHC data from UPS patient samples

(**Figure 1H**), where 31/57 (54%) patients had *FAP* expression in $\geq$ 75% of cells (**Figure S2F,**

S2G, S2H). Other targets classically associated with the stroma that were upregulated in

malignant cells within the STS TME include *MRC2*, *CD248*, and *PDGFRB*.

Monospecific CAR-T cell targets in STS

The ideal monospecific CAR-T cell target would have homogeneously high expression on

cancer cells with low to no expression on normal cells to avoid on-target off-tumor toxicity[44].

However, development of CAR-T cells for solid tumors has been hampered by the scarcity of

antigens that meet these conditions with few targets tested in humans[10]. To address this, we

filtered the STS up-regulated genes to eliminate genes frequently expressed more highly in

normal cells than STS cells. We then selected genes coding for surface proteins with high

expression in STS by cross-referencing our scRNA-seq data with the TCGA STS bulk RNA-seq

data (TCGA SARC) and the HPA transcriptomics data in STS cell lines. Finally, we removed

targets with high RNA and protein expression in normal cells and tissues using CAR-T targets

tested in humans as references for maximum tolerable on-target off-tumor toxicity risk (**Figure**

2A, **Table S4**, **Table S5**, **Table S6**, **Table S7**)[45].

We identified targets with RNA and protein expression profiles in normal tissues that mimic the

normal tissue profiles of clinically tested CAR-T targets (**Figure 2B, Figure S3A**). Our approach

reassuringly identified several targets currently in clinical trials for sarcomas, including *IGF1R*[46]*,*

CD163, *EPHB4*, *FGFR1, KDR*, *NCAM1*, *SMO*, *PSEN1* and *PSEN2*[47], and *NT5E*/CD73[48].

Overall, we found 395 unique targets, with 87 common to all sarcomas and 41 common to LPS,

MFS, and UPS (**Figure S3B**, **Table S8, Figure S3C**, **S3D**). LPS and SS were the histotypes

with the highest number of unique targets with 30 and 21, respectively.

In each sarcoma, transmembrane enzymes were the most common class (**Figure 2C**, **Table**

S9), including metalloproteases of the ADAM family[49] (*ADAM10*, *ADAM33*), metallopeptidases

(*MME*), serine proteases (*PRSS21*, *FAP*), and protein tyrosine phosphatases (*PTPRJ*, *PTPRB*).

Other common target classes were glypicans[50] (*GPC3*, *GPC6*), immune checkpoint molecules[51]

(*CD276*, *NECTIN2*, *BTN2A2*), protein tyrosine kinases[52] (*FGFR1*, *ERBB4*, *EFNA4*), NOTCH

family receptors[53] (*NOTCH2*, *NOTCH4*), G-protein coupled receptors – GPCRs[54] (*ADGRL2*,

GPR107), cell adhesion molecules of the Immunoglobulin superfamily – Ig-like CAMs (*CLMP*,

NEO1), semaphorins[55] (*SEMA4A*, *SEMA6D*), cytokine receptors (*IL1R1*, *IFNAR2*), c-type

lectins[56] (*MRC1*, *MRC2*), and solute transporters (*ABCA1*, *MMGT1*). Gene set over-

representation analysis (ORA) of the 395 targets revealed enrichment in gene sets associated

with cell–matrix interaction, axon and neural projections growth, cell mobility and migration, and

solute transport across the plasma membrane (**Figure 2D**, **Figure S4A–E, Table S10, S11**). We

observed moderate correlation between bulk RNA-seq expression of CAR-T targets in TCGA

SARC and scRNA-seq expression (**Figure S3E**). For many targets like *CLMP, PDGFRB,*

NOTCH2, and *GPC6*, TCGA SARC samples had the highest median expression compared with

other cancer types in the TCGA Pan-Cancer Atlas (**Figure 2E**, **Figure S4F**).

To prioritize targets, we identified those essential for sarcoma cell line growth in vitro by

analyzing publicly available CRISPR KO screening data from DepMap[57] (**Figure 2F**, **Table**

S12). Twenty-two genes were essential for at least one sarcoma cell line, with some being more

selective for specific STS, like *TSPAN31* for LPS. Other genes were essential for multiple

sarcoma cell lines, like *PTK2*, *VPS28*, and *UBE2D3*. We then studied the frequency of CNAs,

gene fusions, and SNVs and small insertions/deletions (indels) involving each target, using

publicly available DNA sequencing data (TCGA SARC[36] and MSK-IMPACT, **Table S13, S14,**

S15, S16)[58]. We found targets that were frequently amplified in specific STS histotypes in TCGA

with high correlation between their CNA level and RNA expression, indicating that these events

could have a driver effect[59,60] (**Figure 2G**). For example, *TSPAN31* ($r = 0.89$) was amplified in

72.4% of DDLPS, and *FGFR1* ($r = 0.68$) was amplified in 12.5% of UPS. Integrating data from

TCGA and MSK-IMPACT with the results of a search in the public Mitelman Database[61],

encompassing 52 STS types and 3283 unique fusions (**Figure 2H**, **Table S15**), we found 213

fusions across 20 STS histologies that involved at least one target. The CAR-T targets most

altered by SNVs or indels included *MACF1* and *AXL* in UPS/MFH (10.2% and 6.1% in TCGA,

Figure 2I, **Table S16**).

Bispecific CAR-T targets in STS: 'AND' and 'AND–NOT' logic-gated CARs

To increase the surface antigen pool targetable by CAR-T cells in solid tumors, new bispecific

CAR-T designs have been developed. Specifically, Boolean-logic 'AND'-gated CAR-T cells[62] like

the logic-gated intracellular network (LINK) CAR-T[12], and the synthetic NOTCH receptor

(synNOTCH) CAR-T[63] cause T-cell activation only when two separate antigens are recognized

on target cells. Conversely, inhibitory CAR-T cells (iCARs)[64] are an example of Boolean-logic

'AND-NOT'-gated CAR-T cells, where an inhibitory domain prevents T-cell activation upon

recognition of an antigen selectively expressed on normal but not on tumoral cells.

To discover targets for 'AND' CAR-T cells in STS, we sought to identify genes with high co-

expression in STS cells and low to no co-expression in normal cells and tissues. (**Figure 3A,**

Figure S5A). To this end, we measured co-expression of STS up-regulated genes in tumor cells

at the single cell level using CS-CORE[65] (**Figure S5B**), and in normal cells and tissues using

the Genotype-Tissue Expression (GTEx)[66] bulk RNA-seq and proteomics datasets, the Human

Proteome Map (HMP)[67] proteomics dataset, and the HPA scRNA-seq, bulk RNA-seq, and IHC

datasets. We then compared tumor co-expression to normal tissue co-expression and

calculated the $TS_{\text{'AND'}}$ score as a measure of targetability for each 'AND' logic-gated CAR-T

gene pair, where pairs with $TS_{\text{'AND'}} \geq 1$ were those with better targetability. Although we found

3608 unique 'AND' CAR-T pairs (**Figure 3B, Table S17**), only 4 (0.1%) were common to all

sarcoma histotypes, highlighting the histotype specificity of gene pairs as well as the challenge

of identifying broadly applicable target pairs for 'AND' logic-gated CAR T cells.

Most 'AND' CAR-T pairs consisted of genes with restricted expression in orthogonal normal cell

types but shared expression in STS. Consistent with the mesenchymal cell origin of STS[68],

gene pairs from "Blood & Immune Cells" and "Neuronal Cells" were the most frequent in all STS

histotypes reflecting the shared development of mesenchymal, hematopoietic, and neuronal

cells from the embryonic mesoderm and neural crest[69] (**Figure 3C, Figure S5C–F**). We

previously observed expression of epithelial markers in malignant STS cells[17], and gene pairs

from both epithelial and mesenchymal cells were identified as targets for 'AND' CAR-T cells.

These results highlight the potential for bispecific CAR-T cells to leverage the dysregulated

differentiation of malignancies as a therapeutic target[70].

For 'AND-NOT' CAR-T target pairs, the ideal combination would include an activator highly

expressed on cancer cells, and an inhibitor with no expression on cancer cells but always co-

expressed with the activator on normal cells (**Figure 3D, Figure S5A**). To find activator-inhibitor

target pairs for 'AND-NOT' CAR-T cells, we used highly expressed STS up-regulated genes as

activators and STS down-regulated genes with average scRNA-seq tumor expression $\leq 50^{th}$

percentile as inhibitors and measured their expression in normal cells and tissues. We

combined the results of the co-expression analysis in each dataset to obtain a summative

targetability measure, the $TS_{\text{'AND-NOT'}}$ score, with values raging from -1 to 1, where higher values

indicate higher co-expression in normal tissues/cells. Overall, we found a total of 10,584 'AND-

NOT' CAR-T pairs across all sarcomas (**Figure 3E**, **Table S18**). Of these, 79 (0.7%) pairs were

shared across all five STS, including *TSPAN6–CLDN17*, *PDGFC–GPR33*, and *IL17RD–ZNRF4.*

Pairs with genes belonging to the same HPA single cell gene signatures were the most frequent,

reflecting similar patterns of expression in normal cells (**Figure 3F**).

Considering the 'AND' and 'AND-NOT' CAR-T target pairs with high *TS* scores, again we found

that most 'AND' pairs tend to have similar expression profiles in the same cells and tissues,

while 'AND-NOT' pairs show opposite expression in the same tissues and cells, which was true

both at the protein and RNA level (**Figure 3G-H**). Examples of 'AND' pairs include *NCSTN–*

BCAN ("Blood & Immune Cells" – "Glial Cells"), *TM9SF1–CD81* ("Blood & Immune Cells" –

"Glandular Epithelial Cells"), and *NTRK2–CD99* ("Neuronal Cells" – "Mesenchymal Cells").

Examples of 'AND–NOT' CAR-T target pairs include *PTPRC–ITGAM* ("Blood & Immune Cells"),

PTPRZ1–NRXN3 ("Glial Cells" – "Neuronal Cells"), and *ADAM2–LY6K* ("Germ Cells"). Many of

the bispecific CAR-T targets were essential for STS cell line growth (**Figure S6A**) and/or were

genetically altered in human STS (**Figure S6B-H**, **Table S19-S21**), suggesting their importance

in STS oncogenesis.

Targeting intracellular antigens in STS: TCR-T and PC-CAR-T cells

CAR-T cells target cell surface antigens but cannot directly recognize intra-cellular proteins,

which constitutes a large fraction of cancer antigens[71]. To address this issue, TCR-T and

peptide-centric CAR-T (PC-CAR-T)[72] cells have been developed targeting peptides presented

by the major histocompatibility complex (MHC) on tumor cells. Most peptides targeted by TCR-T

cells tested in humans so far derive from cancer germline antigens (CGAs)[73], which are highly

expressed in cancer and reproductive organs, such as *MAGEA4* or *CTAG1B,* coding for the NY-

ESO-1 protein[27,30]. SCAN-ACT uses DGE analysis to identify tumor cell up-regulated genes

coding for either intra- or extracellular proteins, with normal tissue expression similar to CGAs

targeted by TCR-T cells tested in humans to find tumor-specific genes with expression restricted

to immune-privileged sites thus minimizing on-target off-tumor toxicity (**Figure 4A**, **Table S22**).

SCAN-ACT identified 76 genes across five STS, with some showing uniformly high expression,

and others more specific to one or few histologies (**Figure 4B**, **Table S23**). UPS had the highest

number of targets (45), followed by LMS (35), LPS (28), SS and MFS (26), with 7 shared across

all histologies (*ACTRT3*, *ADAM12*, *ADAM32*, *CXCL16*, *MAP10*, *MSX1*, and *SUV39H2*, **Figure**

S7A). Overall, 34/76 antigens (44.7%) were known CGAs[73], and gene set ORA identified a roles

of the targets in gametogenesis and gonadal cells function (**Figure 4C**). Few of the targets were

essential for STS cell line growth in vitro (**Figure S7B**), amplified in human STS (**Figure S7C**,

Table S24), or involved in SNVs/Indels (**Figure S7E**, **Table S25**). However, 20 gene fusions in

STS involved at least one target (**Figure S7D**).

Next, we identified a pool of $pMHCs$ suitable as TCR-T or PC-CAR-T targets for each antigen.

Specifically, we assessed the HLA presentation and immunogenicity potential of all the 9-mer

and 10-mer peptides derived from each antigen, using HLAthena[74] and DeepImmuno[75],

respectively, focusing on 9-mers and 10-mers as these account for more than 97.5% of HLA

class I-bound peptides[76]. Also, we chose the top 10 most frequent HLA alleles in each major

geographical area as reported in the Allele Frequency Net Database (AFND)[77], accounting for

118 HLA alleles (**Table S26**). A major limitation of TCR-T cells is their ability to recognize

peptides presented on one or rarely two HLA alleles, making them accessible to a small fraction

of the population. To address this, we combined the HLAthena and DeepImmuno predictions

with the average global HLA allele frequency of each $pMHC$ to obtain a combined TCR-T/PC-

CAR-T targetability score, $TS_{TCR/PC}$. We retained the top 5,020 $pMHCs$ formed by 348 peptides

and 115 HLA alleles with 322 $pMHCs$ shared between $\geq$ 1 antigen (**Table S27**). *ADAM32* had

the highest number of targetable $pMHCs$ (n =136), covering 64 HLA alleles (**Figure S7F**). We

found a higher number of targetable $pMHCs$ for HLA-C (n = 2,041), compared to HLA-B (n =

1,540) and HLA-A (n = 1,439). The top three alleles by number of targetable $pMHCs$ were

C*07:01 (n =168), C*06:02 (n =145), and C*07:02 (n =131), while C*04:01 had $pMHCs$ with the

highest median $TS_{TCR/PC}$ (2.1) (**Figure S7G**).

Because TCR degeneracy allows T-cells to recognize different peptides presented on the same

MHC molecule[78], we designed a $pMHC$ cross-reactivity prediction algorithm called PS-TRACT

(<u>P</u>eptide <u>S</u>imilarity and <u>T</u>CR <u>R</u>eactivity <u>A</u>nalysis for <u>C</u>ellular <u>T</u>herapies). PS-TRACT measures

the similarity of each tumoral peptide to peptides of the same length in the normal MHC

ligandome (**Supplementary Data**) while considering the tissue-specific expression and

likelihood of presentation of each cross-reactive peptide. Peptide similarity was measured in

terms of amino acid physicochemical[79] and HLA binding energy covariance similarity[80], given

that peptide sequence identity alone cannot fully predict T-cell cross-reactivity[81–83]. Also,

because not all peptide residues bear the same importance in mediating TCR recognition[84], we

applied an entropy-based weighting, postulating that residues with higher entropy could be more

relevant for $pMHC$ TCR recognition[74].

We validated PS-TRACT on experimentally observed cross-reactive peptides, including tumoral

and microbial epitopes[85], and the *TTN* peptide ESDPIVAQY which was found to be the off-tumor

target in cases of fatal myocarditis in patients receiving TCR-T cells directed at the *MAGEA3*

epitope EVDPIGHLY[86] (**Figure S8A**). In each known cross-reactive pair, the target peptide was

scored for similarity both against its known cross-reactive peptide and against all the normal

peptides of same length binding to the same MHC molecule, used as controls. As expected,

known cross-reactive peptides had peptide similarity scores (S_{sim}) $\geq$ 99.9[th] percentile

(**Supplementary and Source Data**). We applied PS-TRACT to each STS $pMHC$ and quantified

cross-reactivity scores, S_{CR}, for each tissue in the HPA/GTEx consensus database

(**Supplementary Data**).

To test if PS-TRACT can predict tissue-specific TCR cross-reactivity, we applied it to $pMHC$

targets for which TCR-T cells have been tested in clinical trials with safety outcome data

available[11] (**Figure S8B**, **Table S28**). We selected the S_{CR} score for the $pMHC$ target of afami-

cel, GVYDGREHTV–HLA-A*02:01, as the upper limit for our analyses given afami-cel's

promising safety profile in clinical trials[27,28]. We found that $pMHCs$ for which cross-reactivity was

observed in clinical trials[11] had higher tissue-specific S_{CR} scores compared to GVYDGREHTV–

HLA-A*02:01, as well as higher overall S_{CR} profile difference ($S_{CR}Diff$) with GVYDGREHTV–

HLA-A*02:01 when considering all tissues (**Figure S8C**). For example, the *MAGEA3*-derived

$pMHC$ KVAELVHFL–HLA-A*02:01, targeted by TCR-T cells causing two cases of fatal

necrotizing leukoencephalopathy[87], had a S_{CR} in the cerebral cortex which was 2.4 standard

deviations above the S_{CR} of GVYDGREHTV–HLA-A*02:01 with the cross-reactive peptide

identified as AVADLVHIV from *MCHR2,* which is highly expressed on neurons. Similarly, we

observed a high cardiac S_{CR} score for EVDPIGHLY–HLA-A*01:01 driven by the known cross-

reactive *TTN* epitope ESDPIVAQY[86].

Using PS-TRACT, we found 1113 STS $pMHCs$ with $S_{CR}Diff \leq 0$ across 182 peptides, 69

antigens, and 113 HLA alleles, and of these 74 were shared between ≥ 2 antigens (**Table S27**).

We found several peptides predicted to form $pMHCs$ with $S_{CR}Diff \leq 0$ with ≥ 10 different HLA

alleles for either the class A, class B, or class C locus (**Figure 4D**). Such MHC-promiscuous

peptides could be targetable with a PC-CAR-T recognizing the same peptide presented on

different MHC molecules as long as the molecular surface of the resulting $pMHCs$ is similar. For

example, YVVDGQIIIY derived from both *PSG8* and *PSG11* bound to 17 class C (total allele

frequency: 70%), 10 class B (total allele frequency: 8%), and 4 class A alleles (total allele

frequency: 6%). Lastly, we found peptides forming $pMHCs$ with very low risk of on-target off-

tumor toxicity ($S_{CR}Diff \leq 10^{th}$ percentile) with highly prevalent HLA alleles (**Figure 4E**). These

peptides could be ideal TCR-T targets, given the increased TCR dependency on HLA restriction,

and could be targeted in a larger fraction of the population due to the high prevalence of their

HLA alleles. For example, YFKDKRVHHF from *DAZL* and QFYDNKGITY from *SUV39H2* were

both targetable on C*07:01 (allele frequency: 14%).

Validation of peptide-MHC binding for high-priority targets

Most $pMHC$ binding prediction tools rely on datasets from mass spectrometric measurements of

eluted peptide ligands (EL method), which require further verifications using the actual binding

measurements (BA method) to eliminate false identifications[88]. Here, we adopted a recently

developed empirical system, IMMUNE (Identifying Massive MHC Utilized Novel Epitopes), to

validate the peptide-MHC binding of targets identified with PS-TRACT. IMMUNE employs a

yeast display system enabling fast cloning as well as mammal-like post-translational

modifications for the ex-vivo expression of complex HLA heterodimers[89,90]. Through anchorage

to agglutinin (formed by Aga1p and Aga2p), a peptide-receptive format of HLA-A, B, or C non-

covalent heterodimer can be assembled on the cell surface for antibody detection and peptide

accommodation (**Figure 5A**). Using A*02:01, A*11:01, B*35:01, and C*12:03 as examples, we

observed the entire heavy chain fusion protein represented by its C-terminal HA staining, the

soluble form of β2m in association with the heavy chain, and the correctly folded heavy

chain/β2m heterodimer represented by the HLA-A, B, C conformation-specific antibody staining,

on the surface of correspondingly transformed yeast strains (**Figure S9A**). The successful

detection of each component of the separately expressed heavy chain and β2m without any

covalent linkers allowed further development of on-yeast peptide binding assays. Without the

necessity to purify HLA proteins, the IMMUNE system substantially expedited BA evaluation and

synergized with our computational prediction for rapid TCR-T or PC-CAR-T target discovery.

By comprehensively analyzing the epitope repertoire of each HLA in the immune epitope

database (IEDB), we designed indicator peptides that could be biotinylated without affecting

HLA binding. Co-staining of HLA molecules and the exogenously loaded biotinylated indicators

confirmed specific binding on the surface of each yeast strain (**Figure S9B**). The capability of

IMMUNE to display peptide-receptive HLA mono-alleles represents an ideal system for

competitive binding analysis against the indicator. Based on overall S_{CR} profile and $TS_{TCR/PC}$, we

selected 9 peptides forming 14 $pMHC$ targets for testing. All test peptides bound A*02:01,

including RLHRIIPKI from SSX (**Figure 5B**, **5C**). However, other than AVLDGLDVLL and

VLYPVPLESY from PRAME, most peptides showed low to moderately competitive binding for

B*35:01 and C*12:03. These results suggest that IMMUNE can be utilized to prioritize pMHCs

for targeting.

Target expression in STS Proteomics Datasets

Next, we analyzed target expression in STS clinical proteomics databases[91,92], sarcoma cell

lines[93,94], and STS tissue micro-arrays (TMAs). Since none of the TCR-T/PC-CAR-T target

antigens were measured in the clinical STS proteomics datasets, we analyzed their expression

in STS cell lines, as reported in the Cancer Cell Line Encyclopedia (CCLE)[93] and in the ProCan-

DepMapSanger[94] repositories. Overall, 33/76 (43.4%) targets were expressed in 21 cell lines

spanning 13 STS histotypes (**Figure 5D, Figure S9C**). Of all the targets assessable for

expression, *SSX2/SSX2B, LIN28B*, and *MAGEA4* had the most cell lines with medium-high

expression in ProCan-DepMapSanger at 32%, 18% and 14%, respectively. In the Tang et al.

STS clinical proteomics dataset[91], we found overall good correlation between scRNA-seq and

protein expression for the STS up-regulated surfaceome genes (**Figure S9D**, **S9E**). This

correlation was consistent across histotypes when considering the monospecific CAR-T targets,

particularly only those with protein expression $\geq 50^{th}$ percentile (**Figure 5E**, **Figure S9F**). We

found that many targets had high expression in a large fraction of samples (**Figure 5F, Figure**

S10A, S10C). Similar results were observed in the Burns et al. dataset (**Figure S10B**).

For the 'AND' CAR-T targets with measurable expression of both genes in the Tang et al.

dataset, we measured co-expression across all the STS samples (**Table S29**). Of the 1207

'AND' pairs, 274 showed positive correlation in protein expression between the two targets

across all STS samples (**Figure 5G**). Of the 438 'AND' CAR-T pairs found in the Burns et al.

dataset, 79 had positive correlation in expression across all samples (**Table S29**). Notably, we

were unable to distinguish tumor and normal cell protein expression from the bulk analysis,

which may confound the correlation.

Lastly, we measured expression of selected targets using in-house TMAs of LMS, SS, MFS,

UPS, and MPNST samples (**Figure 5H**). *GPC6* was expressed in 100% (*n* = 10) of SS, 81.4%

(*n* = 59) of UPS, and 95% (*n* = 537) of LMS samples, similarly to *MRC2* which was expressed in

all SS samples, 84.6% LMS, 79.7% UPS, and 76.1% LPS (*n* = 21) samples. LMS had the

highest proportion of positive samples for *KITLG* (91.0%), while SS had the highest number of

samples positive for *CLMP* (90.0%), followed by LMS (89.4%). These antigens were selected

for IHC assessment due to their overall high STS expression and low normal expression profile.

Target Prioritization

We calculated a Target Priority Index (TPI) for each cell therapy target class, integrating normal

cell/tissue expression with functional, genomics, and proteomics STS data, while considering

single-cell tumor expression homogeneity and existing immunotherapeutic agents (**Table S30-**

S33). We defined scRNA-seq expression homogeneity as the product of average expression

and fraction of cells with non-zero expression (H_{RNA} score). For monospecific and bispecific

CAR-T targets, we used the average H_{RNA} score of *CD276* across all histotypes as the

homogeneity reference, and the average fraction of samples in the Tang et al. proteomics

dataset with *CD276* z-score ≥ 0 as the protein expression reference because *CD276* has been

tested as a CAR-T target in STS[95]. Using LPS as an example (**Figure 6A**, **6B**), notable

monospecific CAR-T targets in the top 100 TPI_{CAR} included *GPC6* (high protein expression and

H_{RNA}), *FGFR1* (essential in LPS cell lines and with ADCs in clinical development[96]), and

TSPAN31 (essential for LPS cell lines and involved by gene fusions and amplifications in LPS).

For bispecific CAR-T targets, we integrated the average $TS_{,AND,}$ and $TS_{,AND-NOT,}$ for all the

targetable pairs in which each gene appears into the TPI, prioritizing genes suitable both as

'AND' CAR-T targets and 'AND-NOT' CAR-T activators. In LMS as an example (**Figure 6C**, **6D**,

6E), notable targets with antibodies developed clinically or pre-clinically include *GPNMB*[97],

RTN4[98] and *CD46*[99]. Considering all the monospecific and bi-specific CAR-T targets identified,

69 had a clinically developed monoclonal antibody (mAb), antibody-drug conjugate (ADC),

single-chain variable fragments (scFv), fusion protein, nanobody, or fragment antigen binding

region (Fab), while 36 had at least a mAb or ADC in pre-clinical development. Also, of all the

CAR-T targets described, 26 and 32 have been previously tested as CAR-T targets either

clinically or pre-clinically, respectively.

Finally, we calculated a TPI for TCR-T and PC-CAR-T target antigens, using *MAGEA4* H_{RNA} in

SS as the homogeneity reference, given afami-cel's efficacy in SS. In addition, we considered

protein expression in STS cell lines, the total number of target $pMHCs$, and the HLA allele

frequency per antigen (**Table S33**). Top targets included *PRAME* and *LY6K*, which have more

homogenous expression than *MAGEA4* in SS and a higher total HLA allele frequency. In

addition, *SSX1* is recurrently involved in pathogenic fusions in SS and has a higher predicted

number of targetable $pMHCs$ than *MAGEA4* (**Figure 6F**, **6G**). Given the large number of targets

predicted by SCAN-ACT, the TPI will be helpful to identify the most promising tumor-specific

antigens for further testing.

Discussion

The major bottleneck for development of ACTs in solid tumors continues to be the lack of tumor-

specific antigens with low expression on normal cells[100,101]. Differently from hematological

malignancies, solid tumors lack lineage-specific antigens with homogeneous high-density

expression. Also, it remains challenging to ascertain an acceptable level of target expression in

normal tissues. To address this, we developed SCAN-ACT, a computational pipeline leveraging

scRNA-seq to discover targets for TCR-T cells, and monospecific, bispecific, and peptide-centric

CAR-T cells, targeting both intracellular and surface proteins.

We utilized SCAN-ACT to screen for targets in STS, a disease with limited therapeutic options

where ACT has proven potential. The characteristic heterogeneity of STS is also an ideal small-

scale representation of solid tumor heterogeneity, thus providing a high-bar test case for SCAN-

ACT. We screened sarcoma-specific antigens using single-cell DGE analysis between STS

cells, non-tumoral cells in the TME, and normal cells across 30 human tissues. scRNA-seq can

identify genes that are more selectively expressed in tumoral cells compared to non-tumoral

cells in the TME, as opposed to bulk transcriptomics and proteomics[17,18]. Additionally, scRNA-

seq allows measurement of target expression heterogeneity, a crucial barrier to ACT efficacy in

solid tumors[1], and the availability of normal tissue scRNA-seq data allows for better estimation

of the on-target off-tumor toxicity risk[41,42]. Previous studies demonstrated that target expression

even on rare cellular populations in vital organs can mediate CAR-T cell toxicity[102], and scRNA-

seq may better predict such off-tumor effects than bulk RNA-seq[103].

We extensively validated the off-tumor toxicity risk of the identified targets using publicly

available multiomics data in normal cells and tissues, including scRNA-seq, bulk RNA-seq, IHC,

and proteomics, with targets already tested in humans as reference. Through this approach, we

identified targets with lower predicted risk of on-target off-tumor toxicity than those currently

available. We assessed the relevance of the identified targets to STS biology at the

transcriptomic, genomic, and proteomic level, as well as functionally, through CRISPR KO

screens. This highlighted *TSPAN31* on chromosome 12, which is frequently involved by CNAs in

DDLPS and co-amplified with *MDM2* in both WDLPS and DDLPS[104], indicating a potential

oncogenic driver role in LPS.

We found targets not previously studied in STS like *GPC6,* a member of the glypican family

which is pathogenic in sarcomas[105], or *CLMP,* described as tumor suppressor in colon cancer[106].

We show targetable pathways and protein families in STS, including semaphorins and plexins

that are physiologically involved in neural development[107] and poorly studied in STS[108]; the

NOTCH pathway, previously targeted in STS[109,110]; the EPH/Ephrin system, described as

pathogenic in sarcomas[111] and physiologically involved in neuron migration, axon guidance, and

vascular remodeling[112]; and the WNT/β-Catenin pathway, implicated in sarcomagenesis and

previously targeted in STS[113]. Also, we identify stromal antigens, previously targeted to

manipulate the TME, as tumor cell intrinsic in STS like *FAP*[43], *FOLR2*[114], or *CD248*[115].

Here we also provide a tool for screening safe and effective $pMHC$ targets for ACT in solid

tumors. We expand the pool of known CGAs targetable in STS[116], identifying antigens not

previously described such as *SUV39H2* or *MEIOC*, while confirming the role of known CGAs

like *PRAME*, *SSX1*, and those of the MAGE and GAGE families[116]. Targetable $pMHCs$ were

identified using computational tools measuring peptide HLA presentation and immunogenicity.

Despite being an in silico analysis, this approach prioritizes peptides likely to trigger cytotoxic T-

cell responses while maximizing the number of patients who could benefit from targeting a

specific $pMHC$. Almost all TCR-T cells tested in humans so far recognize peptides presented on

HLA-A*02:01[11], which is highly prevalent in Caucasians, and less so in African Americans and

Asians and Pacific Islanders[77]. This creates equity and generalizability issues for HLA-restricted

ACTs[32]. To address this, we integrated allele frequency into $pMHC$ target prioritization, including

the most frequent HLA alleles from each major geographical area[77]. We curated a normal MHC

ligandome database of $46x10^6$ $pMHCs$, including 630,022 experimentally validated peptides,

and developed PS-TRACT, a tool for $pMHC$ T-cell cross-reactivity assessment. PS-TRACT

overcomes limitations of current methods[72,117], allowing predictions for 9-mer and 10-mer

peptides over 223 HLA alleles and measuring peptide similarity through biophysical properties

while applying an entropy-based weighting that prioritizes amino acids critical for TCR

recognition. By using a large pool of experimentally observed and computationally predicted

normal $pMHCs$, PS-TRACT provides a multi-tissue cross-reactivity risk estimation for each

tumoral $pMHC$. Finally, we experimentally measured HLA binding of selected target peptides

using a novel yeast display technology, IMMUNE, especially critical for validation of peptide-

binding to less common HLA alleles where in silico predictions may be inaccurate.

Compared to prior CAR-T target discovery efforts that either performed DGE analysis between

tumoral and normal bulk RNA-seq samples[14,15] or analyzed patterns of protein expression in

tumor samples[45] or cancer cell lines[16], we leveraged single-cell DGE analysis with a large cohort

of non-tumoral reference cells, allowing a more precise screening for tumor-specific genes.

Unlike prior studies on bispecific CAR-T targets[14,16,45], we defined tumoral target co-expression

at the single cell level and directly measured co-expression in normal cells and tissues using

correlation analyses, including normal tissue scRNA-seq and proteomics data. Lastly, compared

to prior scRNA-seq studies for CAR-T target discovery[18], we analyzed a larger number of tumor

samples, included more normal reference cells, applied robust single-cell DGE analyses less

prone to false discovery[118], validated the identified targets with genomics, proteomics,

transcriptomics, and functional assays in STS, and identified TCR-T/PC-CAR-T and Boolean-

logic gated CAR-T cell targets.

In summary, we developed a comprehensive tool for the identification of ACT targets in solid

tumors enabling the targeting of both intracellular and membrane proteins. By applying the tool

for STS, we nominate and prioritize novel putative targets to expand and accelerate the early

success of ACT in this disease. Our pipeline and sarcoma ACT target database are freely

accessible for academic use at scanact.stanford.edu.

Acknowledgments

We thank the patients and families who participated in this study. This work was supported by

the My Blue Dots organization (E.J.M.) and the Tad and Diane Taube Family Foundation

(M.v.R., E.J.M.). This work used the UCLA Technology Center for Genomics and Bioinformatics

and the UCLA Translational Pathology Core Lab. The schematics in Figure 1, 2, 3, 4, and Figure

S5 were created with BioRender.com.

Author contributions

S.T., A.K., and E.J.M. were responsible for conceptualization. S.T., A.K., and E.J.M. were

responsible for the methodology. S.T. and E.J.M. were responsible for software. S.T., A.P., A.K.,

and E.J.M were responsible for formal analysis. S.T., A.S., A.K., and E.J.M. were responsible for

data curation. S.T., A.K., and E.J.M. were responsible for writing the original draft. W.J. and A.K.

were responsible for the IMMUNE validation and related methods and results. S.T., A.P., A.S.,

S.V., M.P., N.Q.B, K.N.G., M.v.R., P.H., D.G., S.D., W.J., A.K., and E.J.M. were responsible for

review and editing. S.T. A.P., S.A., P.T., S.A., D.G. and E.J.M. were responsible for visualization.

P.H., M.v.R., A.K., and E.J.M. were responsible for supervision. A.K. and E.J.M. were

responsible for funding acquisition.

Declaration of Interests

E.J.M. has served as a consultant for GLG and Guidepoint. A.K. serves on the advisory board

and holds stock for Dispatch Therapeutics, and Certis Oncology and consults for Sastra Cell

Therapy. W.J. serves as CSO for JWE Science Technology, Inc. who owns the intellectual

property of the IMMUNE technology. The other authors declare no financial interests.

Data and Code Availability

The processed scRNA-seq data new to this paper have been deposited to GEO with accession

code GSE279852 (reviewer token: qbejwckcjnkvbgv). The raw scRNA-seq data new to this

project is available on SRA with project code PRJNA117421. The previously published scRNA-

seq datasets reanalyzed in this project are available in GEO with accession codes GSE212527

and GSE131309. The SCAN-ACT pipeline and code used to create the images in the

manuscript are available on Github at https://github.com/StefTesta/SCAN_ACT and as

supplementary material. The SCAN-ACT pipeline and the CellSARCTx interactive interface are

fully accessible for academic use at https://scanact.stanford.edu.

Supplemental Information

Document S1. Figures S1-S10.

Document S2. Excel File containing Supplementary Tables S1-S34.

Methods

Human subjects

All samples analyzed in this study were collected with informed consent from subjects enrolled

on protocols approved by the Institutional Review Boards at Stanford University and University

of California Los Angeles.

Single cell RNA sequencing of STS samples

STS belonging to the following STS histotypes were included in the study: LMS, MFS, UPS,

DDLPS, WDLPS, and SS. Pathological diagnosis was confirmed through intuitional pathologist

review for the samples analyzed at Stanford University and University of California Los Angeles,

and as reported in the original study for the SS samples[33]. For the MFS2 sample, initial tumor

biopsy was diagnosed as MFS, while final surgical specimen was classified as UPS. Due to the

ambiguity of diagnosis, and the lower representation of MFS samples in our cohort, we decided

to analyze this sample as MFS. Also, the UPS6 sample was classified as Spindle Cell Sarcoma

versus UPS and was hence analyzed with the other UPS samples in the study.

For samples processed at Stanford University, fresh tumors were minced and enzymatically

dissociated using the Human Tumor Dissociation Kit (Miltenyi Biotec) using a GentleMACS Octo

Dissociator with heaters (Miltenyi Biotec) following the manufacturer's instructions. The

dissociated cell suspension was then applied to a 70 um MACS SmartStrainer (130-098-462;

Miltenyi Biotec). The quality of the suspension was checked by generating a hematoxylin and

eosin (H&E) smear and analyzed using a TC10 automated cell counter (BioRad). Samples were

further processed by the Stanford Genomics Shared Resource or the using the Chromium 3'

Gene Expression Solution v3.1 (10x Genomics) per the manufacturer's instructions and

sequenced on a NovaSeq 6000 (Illumina). For samples processed at the University of California

Los Angeles (MFS1/BO112_sarc04, UPS1/BO112_sarc10, UPS2/BO112_sarc11,

UPS3/BO112_sarc13, UPS4/BO112_sarc15, UPS5/BO112_sarc17, LPS1/BO112_sarc07,

LPS2/BO112_sarc09, LPS2sx/BO112_sarc09_sx, LPS3sx/BO112_sarc16_sx), these were

dissociated by treatment with collagenase D (100mg/mL) and DNase I (100mg/mL) using a 100

μm cell strainer, following the manufacturer's instructions. After digestion, RBC lysis was

performed by adding 1mL of BD lysing buffer (BD Biosciences) and incubating for 5 mins. Then,

dead cells were eliminated using a dead cell removal kit (Miltenyi Biotec). Cells were counted

using a Countess II cell counter (Thermo Scientific). Samples were then further processed by

the UCLA Technology Center for Genomics & Bioinformatics (TCGB) using the Chromium 3'

Gene Expression Solution v3.1 (10x Genomics) per the manufacturer's instructions and

sequenced on a NexSeq500 High Output instrument (Illumina).

Pre-processing of scRNA-seq STS samples

Sample demultiplexing, read alignment to the GRCh38 human reference genome, and

generation of feature-barcode matrices were performed using CellRanger v6.0.0 (10x

Genomics). Raw data was pre-processed in R (version 4.2.2.)[119] using the Seurat package

(version 4.3.0)[120]. Quality control was performed separately for each sample and consisted of

excluding cells with number of expressed genes < 200 and percentage of mitochondrial genes >

15%. We then identified and excluded doublets using the R package DoubletFinder[121] (version

2.0.3) applying the standard workflow as described by the developer.

Sample Integration and cell annotation

Log-normalization was performed independently for each sample using the Seurat function

NormalizeData. After this step, the Seurat objects for each independent sample were merged

preserving the individual normalization of the data. Sample integration and clustering were

performed separately for each histotype, accounting for batch effects and differences in

sequencing technologies. The top 2000 highly variable genes of the merged and normalized

expression matrices were identified using the function FindVariableFeatures, and then centered

and scaled using the ScaleData function. Principal component analysis was then run using the

top 2000 highly variable genes previously identified. To integrate multiple samples, the harmony

R package[122] (version 0.1.0) was employed for batch correction. Harmony was ran using both

the sample of origin and the sequencing technology variables as arguments of "group.by.vars",

and based on the first 25 PCA dimensions previously identified. UMAP embeddings were then

obtained using the first 25 Harmony dimensions, and clusters were obtained by calculating the

k-nearest neighbors (k-NN) and a shared nearest-neighbor graph using the Louvain algorithm

implemented in the FindClusters function of the Seurat package with a resolution of 1.0. For the

LPS, MFS, UPS, and LMS samples, endothelial cells, myeloid cells, lymphoid cells, and CAFs

were annotated using the differentially expressed genes discovered with the FindMarkers

function in Seurat, which was run with the default parameters except for only.pos = T and

min.pct = 0.25. Canonical cell identity markers obtained from the literature were used for cell

type annotations. A list of the marker genes for each cell type in our samples is provided in

Table S2. A small number of cells could not be uniquely assigned to a specific cell identity and

were removed from further downstream analysis. UMAP plots were generated with Seurat and

scCustomize[123]. Density plots were generated with the R package Nebulosa[124].

Public scRNA-seq Datasets

Synovial Sarcoma Samples

Previously published synovial sarcoma 10x Genomics and SMART-seq2 scRNA-seq data from

Jerby-Arnon et al.[33] were downloaded from GEO (GSE131309). After applying quality control

filtering as described above, the Seurat objects for each sample were integrated using the

harmony[122] pipeline as previously described. Author-provided annotations were used to classify

cells as sarcoma cells, endothelial cells, lymphoid cells, myeloid cells, and CAFs.

Normal Tissue scRNA-seq Datasets

A list of normal tissue scRNA-seq samples of reference used for differential gene expression

was obtained from the HPA website

(https://www.proteinatlas.org/about/assays+annotation#singlecell_rna). If for any of the

reference datasets listed on the HPA dataset, only one sample was available, additional data

was obtained from either Tabula Sapiens[41] or from the Human Cell Atlas[42]. For each dataset, the

raw count matrices were downloaded from the publication-specific repositories as follows:

Adipose Tissue (GEO: accession code GSE155960); Bone Marrow (GEO: accession codes

GSE159929-GSM4850584, and Tabula Sapiens: https://cellxgene.cziscience.com/e/4f1555bc-

4664-46c3-a606-78d34dd10d92.cxg/); Brain (The Allen Brain Map: https://portal.brain-

map.org/atlases-and-data/rnaseq/human-m1-10x); Breast (GEO: accession code GSE164898,

except for D6-D10 samples); bronchus (figshare: accession code fig11981034); Colon (GEO:

accession code GSE116222); Endometrium (GEO: accession code GSE111976); Esophagus

(GEO: accession codes GSE159929-GSM4850580 and HCA:

https://cellxgene.cziscience.com/e/b07fb54c-d7ad-4995-8bb0-8f3d8611cabe.cxg/); Eye (GEO:

accession code GSE137537); Fallopian Tube (GEO: accession code GSE178101, excluding

hydrosalpinx samples); Heart Muscle (GEO: accession code GSE109816 and Tabula Sapiens:

https://cellxgene.cziscience.com/e/e6a11140-2545-46bc-929e-da243eed2cae.cxg/); Kidney

(GEO: accession code GSE131685); Liver (GEO: accession code GSE115469); Lung (Tabula

Sapiens: https://cellxgene.cziscience.com/e/0d2ee4ac-05ee-40b2-afb6-ebb584caa867.cxg/);

Lymph Node (GEO: accession codes GSE159929-GSM4850583 and Tabula Sapiens:

https://cellxgene.cziscience.com/e/18eb630b-a754-4111-8cd4-c24ec80aa5ec.cxg/); Ovary

(EMBL-EBI: accession code E-MTAB-8381); Pancreas (GEO: accession code GSE131886);

PBMC (GEO: accession code GSE112845 and Tabula Sapiens:

https://cellxgene.cziscience.com/e/983d5ec9-40e8-4512-9e65-a572a9c486cb.cxg/); Placenta

(EMBL-EBI: accession code E-MTAB-6701); Prostate (Tabula Sapiens:

https://cellxgene.cziscience.com/e/d77ec7d6-ef2e-49d6-9e79-05b7f8881484.cxg/); Rectum

(GEO: accession code GSE125970); Salivary Gland (Tabula Sapiens:

https://cellxgene.cziscience.com/e/f01bdd17-4902-40f5-86e3-240d66dd2587.cxg/); Skeletal

Muscle (GEO: accession code GSE143704); Skin (GEO: accession code GSE130973); Small

Intestine (GEO: accession code GSE125970); Spleen (GEO: accession codes GSE159929-

GSM4850589 and Tabula Sapiens: https://cellxgene.cziscience.com/e/cee11228-9f0b-4e57-

afe2-cfe15ee56312.cxg/); Thymus (Tabula Sapiens:

https://cellxgene.cziscience.com/e/0ced5e76-6040-47ff-8a72-93847965afc0.cxg/); Testis (GEO:

accession code GSE120508); Tongue (Tabula Sapiens:

https://cellxgene.cziscience.com/e/55cf0ea3-9d2b-4294-871e-bb4b49a79fc7.cxg/); Vascular

(Tabula Sapiens: https://cellxgene.cziscience.com/e/a2d4d33e-4c62-4361-b80a-

9be53d2e50e8.cxg/). After download, each dataset was pre-processed in a similar fashion as

our own STS samples. The pre-processed samples for a given tissue were then integrated with

the harmony pipeline[122] using the sample variable, and for the SS samples the sequence

technology variable as well, for batch correction, as previously described. Then, unsupervised

clustering using the Louvain algorithm with a resolution of 1 was applied for all datasets.

Exceptions to this workflow include the brain dataset where the dimensionality reduction and

clustering from HPA were used, as well as the endometrium and the bronchus datasets where

the author-provided clusters and annotations were used. Author-provided annotations were

used whenever possible, and if these were not available, the package SingleR[125] and common

gene markers were used for cluster cell-type annotation, where each cluster was assigned to its

most frequent cell type. In SingleR[125], the Encode[126] and Blueprint epigenomics[127] datasets

were used for annotation of all datasets except for the pancreas for which the scRNA-seq

reference data from Baron et al.[128] was used instead.

Identification of tumor cells based on InferCNV

CNVs were calculated using the R package inferCNV[35]. Each sample was independently

analyzed using as reference cells the previously annotated CAFs, endothelial cells, myeloid

cells, and lymphoid cells from the TME. These were identified based on expression of known

marker genes available from the literature as described above. The following parameters were

used to run the inferCNV function: analysis_mode="subclusters";

tumor_subcluster_partition_method="qnorm"; cluster_by_groups= FALSE; k_obs_groups = 3,

denoise=T, HMM=TRUE. All the other parameters were set as default. Only for UPS

k_obs_groups was set to 2 since this resulted in better overlap with the classification based on

comparison to bulk sarcoma tumors (see below). To define a cell as potentially malignant, we

used the CNA states across all genes calculated using a hidden Markov model (HMM) as

implemented in the inferCNV package. To reduce the number of false positive CNVs identified,

a Bayesian latent mixture model was applied with a default p-value threshold of 0.1 for all

samples. For each gene, inferCNV reports a CNA state as one of the following: state 1

corresponding to a complete allele loss, state 2 corresponding to loss of one allele, state 3

corresponding to the neutral state with both allelic copies intact, state 4 corresponding to gain of

one allele, state 5 corresponding to gain of 2 alleles, and state 6 corresponding to gain of more

than two alleles. For each sample, CNA states of 6 in the output data matrix obtained in the 19th

step of the inferCNV workflow were first converted to a state of 5, and then the CNA states were

normalized to values ranging from -1 to +1 and centered on 0 using the following formula:

$(\text{CNA State} - 3)/2$. The matrix values were then squared to give the same weight in the

analysis to copy gains and copy losses. Then we calculated the overall CNA level of each cell.

For each sample, we identified, the CD45^{-} cells with the top 10% CNA level. Cells with

normalized PTPRC RNA count = 0 were defined as CD45^{-}. The average CNA level for these

cells was defined as the CNA profile of the tumor. A spearman correlation coefficient was then

calculated for each cell between its overall CNA level and the CNA profile of the tumor. Cells

whose spearman correlation coefficient was above the 25th percentile of all the cells in each

sample were considered to have detected CNAs and labeled as potentially malignant. The

output of the inferCNV analysis used to identify tumor cells is provided in the supplementary

data.

Bulk sarcoma similarity score

Bulk RNA-seq of STS samples from TCGA were downloaded (Tatlow and Piccolo, 2016) and

scaled to transcripts per million (TPM) as previously described and used as the bulk tumor

references. Cells from scRNA-seq data available from Jerby-Arnon et al.[33] that were annotated

as normal fibroblasts, macrophages, and endothelial cells were used as the normal reference

cells. First, the sarcoma histotypes of reference were extracted from TCGA, then the expression

matrices for both the bulk RNA-seq sarcoma samples (TPM) and the scRNA-seq samples (raw

UMI counts) were subsampled to include only genes in common between each dataset to

compare. Then, two spearman correlation coefficients were calculated for a given cell of a

scRNA-seq sample, respectively comparing its gene expression profile to the average

expression profile of all the bulk RNA-seq tumor samples of reference (ρ Tumor) and to the

average gene expression profile of all the normal cells in the scRNA-seq dataset used as

normal control (ρ Normal). Based on these, we defined the similarity score as

ρ Tumor – ρ Normal. Cells with similarity score > 0 were classified as potentially malignant. The

InferCNV and similarity to bulk analyses were combined so that a cell was labeled as malignant

if it was CD45⁻ (PTPRC RNA count = 0) and had either detected CNV or a similarity score > 0.

Cells that could not be classified into a specific cell type based on any of the above methods

were labeled as "unknown" and excluded from further downstream analysis.

Differential Expression Analysis

To limit false discovery of differentially expressed genes in STS using scRNA-seq[118], we used a

pseudobulk DGE method, limma-voom[37], which has been shown to be statistically more robust

and less prone to false discovery than other methods when applied to single cell

transcriptomics. Also, for the STS cells versus non-tumoral cells in TME, we complemented the

limma-voom results with a new differential distribution testing method, distinct[38], which was

specifically designed for DGE using scRNA-seq.

Pseudobulk Limma-Voom

Sarcoma cells and normal cells were identified as above. For each STS histotype, the tumor

cells were compared to endothelial cells, myeloid cells, lymphoid cells, and CAFs in the TME, as

well as with each normal cell cluster of the normal tissues scRNA-seq samples of reference. For

each comparison, first a single integrated Seurat object was created and from this, the raw non-

normalized mRNA count matrix was extracted. To minimize variability in DE analysis and reduce

false discoveries, the synovial sarcoma samples analyzed with plate sequencing technology

and those analyzed with droplet-based technology were tested separately. Also, when

comparing tumor samples to normal samples from HPA-TS-HCA, only genes that were present

in both gene expression matrices were included in the differential expression test, to avoid

identification of falsely up-regulated or down-regulated genes. Based on the information in the

Seurat object metadata, the total mRNA counts for each gene were calculated by summing

across all the cells of a certain cell type belonging to a specific sample, obtaining a unique

pseudobulk for each cell type belonging to a given sample in the analysis. The pseudobulk gene

expression matrix was then converted to a DGEList object using the DGEList function in the

edgeR[129] package (version 3.99.4), and subsequently normalized using the calcNormFactors

function. Before voom transformation using the voom function from the R package limma[37]

(version 3.57.11), lowly expressed genes were filtered from the expression matrix, using a cutoff

of 1 CPM. Counts data was then voom-transformed and a limma workflow was implemented

using the sample variable as the blocking variable to control for batch effect. The limma model

was then fitted using the patient sex as a covariate. Contrasts were then compared between

each normal cell cluster and each sarcoma cluster separately (UPS, MFS, LMS, LPS, and SS).

Genes with a Benjamini-Hochberg (BH) corrected p-value < 0.25 and $\log_2(FC) \neq 0$ were

considered as significantly differentially expressed. We used a more lenient BH-corrected p-

value cutoff given the screening and exploratory nature of this part of our analysis.

Differential Distribution Testing with distinct

The R package distinct[38] (version 1.10.2) was used to perform differential gene distribution

testing between sarcoma cells and non-tumoral cells of the TME. Since distinct compares

empirical cumulative distribution functions for a given gene across cell types, it allows

identification of subtle differences in expression that are not captured by other methods which

might consider only measures of central tendency in the expression distribution. We

hypothesized that this could be particularly useful in comparing transcriptionally similar cells, as

STS and non-tumoral mesenchymal cells in TME. Due to its computational load, it resulted

unfeasible to use distinct for the HPA-TS-HCA normal cells comparisons. The raw gene

expression matrices of sarcoma cells and non-tumoral cells were first processed in Seurat as

described above to create a merged Seurat object. Afterwards, a single cell experiment object

specific to each tumor versus normal comparison was created using the SingleCellExperiment

R package[130] (version 1.20.1). The expression data was then log normalized using the

logNormCounts function from the R package Scuttle[131] (version 1.8.4). A "batch" variable

indicating the sex of the patient was included in the design matrix as a covariate, unless patients

were of the same sex. The distinct_test function was then used to run differential distribution

testing using default parameters. To obtain the $\log_2$ FC of the gene expression between

sarcoma and normal cells, counts were first converted to CPM using the calculateCPM function

from Scuttle[131], and these were used as input for the log2_FC function in distinct[38]. Then the

$\log_2$FC was then calculated as $\log_2 \frac{(\text{average sarcoma cells CPM} + 0.01)}{(\text{average normal cells CPM} + 0.01)}$, where the average RNA

expression in the normal and sarcoma cells were calculated through the distinct function

log2_FC. Genes with a $\log_2 FC \neq 0$ and a "p_adj.glb" value < 0.25 were considered as

significantly differentially expressed between sarcoma and normal cells.

Identification of differentially expressed genes in STS

For each STS histotype, we defined a gene as up-regulated, if it resulted as differentially up-

regulated ($\log_2 FC > 0$, FDR < 0.25) in sarcoma cells in at least one of the comparisons made,

either against reference cells in the TME (limma-voom or distinct), or in the HPA-TS-HCA

datasets (limma-voom). STS down-regulated genes were instead defined as those that resulted

as down-regulated in sarcoma cells ($\log_2 FC < 0$, FDR < 0.25) in at least one of the comparisons

made, and that were never up-regulated in sarcoma cells, separately considering the TME and

the HPA-TS-HCA tests. For genes with more than one alternative transcript, these were tested

separately.

Hierarchical Clustering of STS versus non-tumoral cells in the TME and normal tissues

To perform hierarchical clustering between STS and non-tumoral cells in the TME as well as

normal cells in the HPA-TS-HCA datasets, we used the R package factoextra (version 1.0.7)

and base R functions. First, we filtered the CPM normalized scRNA count matrices to retain only

genes defined as differentially expressed in STS (either up- or down-regulated), considering the

TME and HPA-TS-HCA comparisons separately. Then we merged the count matrices from

different samples/tissues, averaged gene expression per cell type and $\log_2(+1)$ transformed the

expression values. Pearson correlation was then used to obtain pairwise distances between cell

types, separately analyzing STS cells versus non-tumoral cells in the TME, and STS cells

versus normal cells in HPA-TS-HCA (supplementary data). The Pearson distance matrices were

used to create a dendrogram with the hclust base R function, setting the agglomeration method

to "ward.D2", and this was then visualized using the fviz_dend function from factoextra. The

optimal number of clusters, k, was obtained by calculating the silhouette score to estimate how

similar cells within the same cluster are to each other and how different they are from cells in

other clusters, for $k \in \{2, ..., n-1\}$, where n is the number of cells considered. The silhouette

score peaked at $k = 3$ for the TME comparisons, and around $k = 5$ and $k = 6$, for the HPA-TS-

HCA comparisons. We chose $k = 6$ for the latter as it provided a more biologically relevant

separation of the different cell types.

List of Surface Proteins

To obtain a comprehensive list of genes coding for proteins that are known to or predicted to be

expressed on the plasma membrane, we combined The Cancer Surfaceome Atlas[14] with the in

silico human surfaceome (SURFY)[132] after conversion of gene names to standard HGNC

symbols (**Table S4**).

Normal Tissue Expression Datasets

To assess the suitability of a given gene as a cellular therapy target, we used several normal

tissue and cell datasets to measure RNA and protein expression.

Normal tissue RNA expression was assessed using the bulk RNA-seq dataset from the Human

Protein Atlas[39] (version 22.0, Ensembl version 103.38) accessed at

https://www.proteinatlas.org/about/download (file name: rna_tissue_hpa.tsv.zip), and the bulk

RNA-seq dataset from the GTEx[66] accessed at https://gtexportal.org/home/datasets

(expression file: GTEx_Analysis_2017-06-05_v8_RNASeQCv1.1.9_gene_tpm.gct; annotation

file: GTEx_Analysis_v8_Annotations_SampleAttributesDS.txt).

RNA expression in normal cells was evaluated through the HPA single cell RNA-seq dataset

(version 22.0, Ensembl version 103.38), accessed at

https://www.proteinatlas.org/about/download (file name: rna_single_cell_type.tsv.zip).

Protein expression in normal tissues was evaluated using three separate datasets:

1. The HPA dataset (version 22.0, Ensembl version 103.38) obtained at (file name:

normal_tissue.tsv.zip), which contains protein expression data across 44 normal human

tissues obtained through immunohistochemistry.

2. The Human Proteome Map[67] (HPM) dataset obtained at

http://www.humanproteomemap.org/download.php , which reports protein expression

data in 30 normal human tissues obtained through LC-MS/MS proteomics.

3. The GTEx Proteomics dataset[133] which reports protein expression data in 32 normal

human tissues, obtained through tandem mass tag (TMT) proteomics. Data was

obtained from the Supplemental data (Table S2D) of the corresponding manuscript.

Specifically, we used the $\log_2$ transformed protein abundances relative to the average of

the reference in the same run.

Dataset pre-processing

Gene names across datasets were standardized using the R package mygene[134] (version

1.38.0) by converting the gene name aliases to the corresponding standardized HGNC symbol.

For the bulk RNA-seq datasets (GTEx and HPA), TPM expression values were converted to

$\log_2$(TPM +1). Then, the expression data was processed differently based on the target

discovery pipeline considered. For the permissive on-target off-tumor toxicity filter of the

monospecific CAR target discovery pipeline, RNA expression from the GTEx dataset was first

averaged across all samples belonging to a given tissue type. Then tissues were organized in

tissue groups in a similar fashion as done by MacKay et al.[45], and the average gene expression

per tissue group was calculated (**Tables S5**, **S6**). Similarly, for the HPA dataset, the RNA

expression was averaged across tissues belonging to same tissue group. Then, tissue

expression in both the GTEx and HPA dataset was categorized as "Medium–High" (M-H) for

$\log_2$(TPM+1) $\geq$ 4, and "Not Detected–Low" (ND-L) for $\log_2$(TPM+1) $<$ 4. In all the other

pipelines, the expression matrices of the bulk RNA-seq GTEx and HPA dataset were median-

centered across tissues (HPA) or samples (GTEx), and then z-scored within genes, before use

for downstream analysis.

In the HPA IHC protein expression dataset, the protein expression is reported for each cell type

in each tissue as either "Not Detected", "Low", "Medium", or "High". For use in the permissive

on-target off-tumor toxicity filter of the monospecific CAR target discovery pipeline, protein

expression for each tissue was classified as follows: "ND-L" if all the cells in a tissue had only

"Not Detected" or "Low" expression; "M-H" if all the cells in a tissue had "Medium" or "High"

expression, or only "High" expression; "Medium" if all the cells in a tissue had "Medium"

expression; "Variable" if the cells in a tissue had either "Not Detected", "Low", "Medium" or

"High" expression. Tissue group expression was then calculated as the highest tissue

expression within a tissue group using the following hierarchy: M-H > Variable > Medium > ND-

L. Again, tissue groups were defined as above. For use in all the other pipelines, we first

converted the cell type protein expression level from a categorical to a numerical score, where

"Not Detected" = 1, "Low" = 2, "Medium" = 3, and "High" = 4. Then, we calculated the average

expression value per tissue group for each gene, and then obtained a gene per tissue group

expression matrix that was median-centered across tissue groups and z-scored within genes for

downstream analysis.

In the HPA single cell RNA-seq dataset, gene expression is reported as normalized TPM

(nTPM) for each normal cell cluster in 31 tissues. We first, converted the nTPM values to

$\log_2(\text{nTPM}+1)$ and then obtained a gene per cell cluster matrix, where columns represented cell

clusters in each normal tissue in the dataset. The $\log_2(\text{nTPM}+1)$ expression values were then

median-centered across cell clusters and z-scored within genes, and this matrix was used for

downstream analysis.

The LC-MS/MS proteomics expression data from the HPM was $\log_2(+1)$ transformed and then

median-centered across tissues and z-scored within genes, before further analysis. Lastly, in

the GTEx TMT proteomics dataset the relative $\log_2$ transformed protein abundance data

provided by the authors was median-centered across tissue samples and z-scored within

genes, similarly to the other datasets. All the processed normal tissue and cell expression

datasets used in the study are provided in the supplementary data.

Sarcoma Cell Lines RNA Expression

The HPA cancer cell line expression was downloaded from HPA (version 22.0, Ensembl version

103.38) at https://www.proteinatlas.org/about/download (file: rna_celline_cancer.tsv.zip). The

dataset was subsetted to retain only data corresponding to sarcoma cell lines including: GCT,

HT-1080, MES-SA, RD, Rh18, Rh30, Rh41, RKN, S-117, SK-LMS-1, SK-UT-1, TE 441.T, TE

617.T, and U-2197. For each gene, HPA reports the average expression as nTPM for all the cell

lines of a given cancer type. The nTPM values were averaged for those genes with alternative

transcripts, and then converted to $\log_2(nTPM+1)$ before use for downstream analyses. The

processed data is provided in the supplementary data.

Monospecific CAR-T Cell Discovery Pipeline

To define genes suitable for a monospecific CAR-T cell design, first we removed from our initial

pool of STS up-regulated genes any genes that were up-regulated more often in normal cells

than in sarcoma cells, considering all the comparisons in which that gene appeared. Then this

gene pool was filtered to retain only those present in the surfaceome. HLA class I and II genes

were also removed from the pool.

To increase tumor-specificity of our candidate targets, we filtered the gene pool to retain only

those with high tumor expression in both our scRNA-seq data, in TCGA SARC, and in the HPA

sarcoma cell lines (bulk RNA-seq). Tumor expression in our scRNA-seq data was calculated as

the average $\log_2$(CPM+1) across the sarcoma cells of a given STS histotype, where CPM

indicates RNA expression in counts per million, while RNA expression in TCGA SARC was

calculated as the average $\log_2$(TPM+1) across all the samples of a given STS histotype. Then,

we filtered our candidate pool to retain any gene that had above-median expression in both the

SARC TCGA data and in the scRNA-seq data, or above-median expression in the scRNA-seq

data and in the HPA sarcoma cell lines. Genes without expression data in SARC TCGA or

sarcoma HPA cell lines but with scRNA-seq expression $\geq$ the 50th percentile were also retained.

Then we assessed the potential for on-target off-tumor toxicity of each candidate target by

measuring its RNA and protein expression in normal tissues and cells and comparing it to that of

targets for which CAR-T cells are currently available for clinical use or in clinical development.

The list of tested CAR-T targets was obtained from the previous work from MacKay et al[45]

(**Table S7**). Specifically, we identified genes at risk for on-target off-tumor toxicity by applying

both a *permissive* and a *stringent filter*.

For our *permissive filter*, we used the HPA IHC protein database, as well as the GTEx and HPA

bulk RNA-seq databases. Specifically, protein and RNA expression were categorized as

described above, and then the number of tissue groups (TGs) with medium-high (M-H)

expression was measured for the tested CAR-T targets. We used the 90th percentile of the

number of TGs with M-H expression for the tested CAR-T targets in each of the three datasets

as cutoffs. A gene was considered to have passed the filter in a dataset if it had number of TGs

with M-H expression below the respective cutoff.

For our *stringent filter*, we assessed the risk of on-target off-tumor toxicity both at the single cell

level, using the HPA scRNA-seq dataset, and by measuring protein expression at the tissue

level, using the HPA, HPM, and the GTEx protein expression datasets. For this analysis, we

used the HPA protein numerical scoring classification obtained as described above. Also, for the

HPA scRNA-seq dataset we used the average cell type expression across all tissues. Again, we

used the normal cell/tissue expression of tested CAR-T targets as a reference to score our

candidate genes against.

For a gene to have an acceptable on-target off-tumor toxicity risk based on its expression in the

HPA scRNA-seq dataset, it had to meet two conditions. First, we categorized gene expression

as "high" for z-score> 1, and "low" for z-score ≤ 1. Based on this, genes from our pool that had

"high" expression in $\leq$14 cell types, which corresponds to the 75[th] percentile of the number of

cell types with "high" expression for the tested CAR-T targets, were initially retained. Then, we

calculated the maximum expression value per cell type for the tested CAR-T targets and

retained only those genes that had RNA expression below this threshold in each cell type. The

following cell types were not considered: lymphoid immune cells (T-cells, NK-cells, B-cells,

plasma cells), due to the potentially higher tolerability of their depletion; fibroblasts, adipocytes,

and endometrial stromal cells due to potential high transcriptional similarity to the STS in the

study; and cells in the gonads and placenta (ovarian stromal cells, granulosa cells, oocytes,

peritubular cells (testis), cytotrophoblasts, extravillous trophoblasts, Hofbauer cells,

syncytiotrophoblasts, Leydig cells, spermatocytes, late spermatids, spermatogonia, early

spermatids, and Sertoli cells) being sites of immune privilege. Only those genes that met both

criteria were retained based on their single cell RNA expression in normal cells.

1000

For each of the 3 protein expression datasets, we calculated the maximum expression values

per tissue (GTEx and HPM) or tissue group (HPA) for the tested CAR-T targets and used these

as thresholds so that only the candidate targets with protein expression below the threshold in

each of the tissues or tissue groups studied were retained. The following tissues and tissue

groups were not considered: fetal tissues, male or female reproductive tissues, placenta, CD4

T-cells, CD8 T-cells, NK-cells, B-cells, spleen, immune cells and immune tissue. Notably, for the

GTEx proteomics dataset, we used the average z-scored protein expression across all samples

belonging to a given tissue.

1009

1010   For each candidate gene (g), we then combined the results of both filters by calculating a

1011   monospecific CAR Targetability Score, TS_{CAR}, as follows:

1012
$$TS_{CAR}(g) \; = \; \sum_{i=1}^{2} m_i \; + \; \sum_{j=1}^{4} m_j \; + \; m_k$$

Where:

• m_i represents the score from the 2 bulk RNA-seq datasets (HPA and GTEx categorical)

and is calculated as +1 if the gene passes the filter, −1 if it does not, and 0 if the gene is

not present in the dataset.

• m_j represents the score from any of the 4 protein expression datasets (HPA categorical,

GTEx proteomics, HPM, and HPA numeric) and is calculated as +2 if the gene passes

the filter, −2 if it does not, and 0 if it is not present in the dataset.

• m_k represents the score from the HPA scRNA-seq dataset and is calculated as +1.5 if

the gene passes the filter, −1.5 if it does not, and 0 if it is not present in the dataset.

Any gene with $TS_{CAR} \geq 5$ which corresponds to the 25th percentile of the TS_{CAR} for the tested

CAR-T targets was considered as a suitable target for a monospecific CAR-T cell. The results of

the pipeline are reported in **Table S8**.

1025

Gene set over-representation analyses

Gene set over-representation analysis (ORA) was performed using the R packages,

clusterProfiler[135] (version 4.10.0), ReactomePA[136,137] (version 1.46.0), and org.Hs.eg.db[138]

(version 3.18.0). Before analysis, the gene symbols were converted to Entrez gene IDs. Then,

we analyzed all the candidate targets for monospecific CAR-T cells for each STS, using the

function enrichGO for Gene Ontology (GO) term enrichment, the function enrichKEGG for Kyoto

Encyclopedia of Genes and Genomes (KEGG) pathways enrichment, the function

enrichPathway for enrichment of Reactome pathways, and the function enrichWP for

enrichment of WikiPathways terms. The p-value cutoff for significance was set to 0.05 and the

Bonferroni-Hochberg method was used for FDR correction, while minimum and maximum gene

sets sizes were set respectively to 10 and 500. We then performed a clustered ORA on the

monospecific CAR-T targets by STS histotype. In this case we used the compareCluster

function from clusterProfiler and within this we used the same enrichment functions as

described above, depending on the reference database used, again setting FDR cutoff to 0.05

and minimum and maximum gene sets sizes to 10 and 500. When performing the cluster ORA

analysis on the TCR-T and PC-CAR targets, the FDR cutoff was instead set to 0.1 due to

smaller gene pool to test, while the minimum and maximum gene sets sizes were set to 2 and

500, respectively.

1044

TCGA Pan-Cancer Atlas

RNA expression data for the Pan-Cancer Atlas TCGA[139] samples (batch normalized RSEM

values) was obtained from cBioPortal[140,141] for the genes of interest. The downloaded data is

available in the Source Data. Data for the following cancer types was analyzed: KICH: Kidney

Chromophobe; ACC: Adrenocortical carcinoma; LIHC: Liver hepatocellular carcinoma; UVM:

Uveal Melanoma; PCPG: Pheochromocytoma and Paraganglioma; KIRP: Kidney renal papillary

cell carcinoma; LAML: Acute Myeloid Leukemia; THYM: Thymoma; SKCM: Skin Cutaneous

Melanoma; KIRC: Kidney renal clear cell carcinoma; READ: Rectum Adenocarcinoma; LGG:

Brain Lower Grade Glioma; CHOL: Cholangiocarcinoma; COAD: Colon Adenocarcinoma;

PRAD: Prostate Adenocarcinoma; STAD: Stomach Adenocarcinoma; UCEC: Uterine Corpus

Endometrial Carcinoma; CESC: Cervical squamous cell carcinoma and endocervical

adenocarcinoma; BLCA: Bladder Urothelial Carcinoma; ESCA: Esophageal Carcinoma; DLBC:

Diffuse Large B-Cell Lymphoma; TGCT: Testicular Germ Cell Tumors; OV: Ovarian serous

cystadenocarcinoma; LUSC: Lung Squamous Cell Carcinoma; LUAD: Lung Adenocarcinoma;

HNSC: Head and Neck Squamous Cell Carcinoma; BRCA: Breast Invasive Carcinoma; PAAD:

Pancreatic Adenocarcinoma; GBM: Glioblastoma Multiforme; THCA: Thyroid Carcinoma; UCS:

Uterine Carcinosarcoma; SARC: Sarcoma; MESO: Mesothelioma.

1062

Cell Line Gene Dependency

The DepMap project[57,71] reports the results of CRISPR KO experiments on 1100 human cell

lines. To assess the essentiality of our candidate targets for sarcoma cell lines (including both

soft tissue sarcomas as well as chondrosarcoma, osteosarcoma and Ewing's sarcoma), we

used the "CRISPRGeneEffect.csv" file obtained from the DepMap website, containing data

regarding the effect of the KO of each gene in each cell line. Gene effects less than –1 indicate

a significant reduction in cell growth and proliferation, as this was the median gene effect

estimate of genes known to be essential for cell growth and survival.

1071

Soft Tissue Sarcoma Genomics Data

Data on copy number alterations (CNAs) and mutations in soft tissue sarcomas were obtained

from the TCGA SARC firehose database[36] and from the MSK-IMPACT[58] sarcoma study. Both

datasets were downloaded from cBioPortal[140,141]. The file named "data_linear_cna.txt" was used

to analyze linear CNA data from TCGA firehose, while the file named "data_cna.txt" was used to

determine amplification status of a given gene in each sample for both the TCGA and MSK-

IMPACT databases. For the Pearson correlation analysis between linear CNA and RNA

expression, the TCGA SARC bulk RNA-seq expression data obtained as above, was used.

Correction for multiple hypothesis was obtained using the BH method setting the threshold for

significance to 0.01.

1082

Data regarding single nucleotide variants (SNVs) and small insertions/deletions (Indels) was

retrieved from the "data_mutations.txt" files for both the TCGA and the MSK-IMPACT cohorts.

Lastly, data on gene fusions in soft tissue sarcomas was obtained by integrating data in the

"data_sv.txt" file obtained from cBioPortal for the MSK-IMPACT cohort, data from the study from

Gao et al.[142] on recurrent gene fusions in TCGA tumor samples, and from a search through the

Mitelman Database[61,143] (https://mitelmandatabase.isb-cgc.org/mb_search) that included the

following sarcoma types: Ewing sarcoma; Atypical lipomatous tumor/atypical lipoma/well-

differentiated liposarcoma; Alveolar rhabdomyosarcoma; Myxoinflammatory fibroblastic

sarcoma; Undifferentiated pleomorphic sarcoma; Liposarcoma, dedifferentiated; Inflammatory

myofibroblastic tumor/myofibroblastic sarcoma; Soft tissue tumor, NOS; Clear cell sarcoma;

Undifferentiated round cell sarcoma; Undifferentiated sarcoma; Mesenchymal tumor, NOS;

Fibrosarcoma; Embryonal rhabdomyosarcoma; Spindle cell/sclerosing rhabdomyosarcoma;

Myxofibrosarcoma; Endometrial stromal sarcoma; Pleomorphic rhabdomyosarcoma;

Liposarcoma, NOS; Alveolar soft part sarcoma; Leiomyosarcoma; Sclerosing epithelioid

fibrosarcoma; Malignant soft tissue tumor, NOS; Angiosarcoma; Rhabdomyosarcoma, NOS;

Epithelioid sarcoma; Dermatofibrosarcoma protuberans/Bednar tumor/giant cell fibroblastoma;

Low-grade fibromyxoid sarcoma; Synovial sarcoma; Histiocytic sarcoma/malignant histiocytosis;

Liposarcoma, myxoid/round cell; Liposarcoma, pleomorphic. The combined fusion data in STS

is available from **Table S15**.

1102

Logic-gated 'AND' CAR Targets Discovery Pipeline

To discover gene pairs suitable for a bispecific logic-gated 'AND' CAR design, first we selected

genes up-regulated in sarcoma cells, separately for each STS, as described above. Next, we

removed from this pool of STS up-regulated genes any genes that were up-regulated more

often in normal cells than in sarcoma cells, considering all the comparisons in which that gene

appeared. Then, we filtered this gene pool to include only highly expressed genes, as defined in

the monospecific CAR-T cell target discovery pipeline. Then, for the selected genes we

computed all possible pairs and measured gene co-expression at the single cell level by running

the CS-CORE algorithm from the R package CSCORE[65] (version 0.0.0.9). Specifically, we ran

CS-CORE on all the tumor cells of a given sarcoma histotype and corrected p-values for

multiple hypothesis testing using the BH method. Any non-significant correlations were

converted to a score of 0, using an adjusted p-value significance threshold of 0.05. For a given

gene pair (g_i, g_j), CS-CORE computes a correlation score, which we will refer to as

$\rho_{Tumor}(g_i, g_j)$, and which can assume values between –1 (lowest likelihood of co-expression)

and +1 (highest likelihood of co-expression).

1118

Then, for each gene pair we assessed co-expression in normal cells and tissues, with the

caveat that genes for an ideal target pair of an 'AND' logic-gated CAR would have mutually

exclusive expression across all normal cells and tissues. To assess this, we used, the GTEx and

HPA bulk RNA-seq databases, the HPA scRNA-seq database, the HPA IHC protein expression

database, as well as the TMT GTEx proteomics and the HPM LC-MS/MS proteomics datasets.

For the bulk RNA-seq datasets (GTEx and HPA), the median-centered, z-scored numerical

gene expression matrices were used, where expression per each tissue (HPA), or tissue sample

(GTEx) was measured. In this case, we used Pearson correlation to assess gene co-expression

across normal tissues. Correction for multiple hypothesis testing was performed using the BH

method, and for pairs with a corrected p-value ≥ 0.05 the correlation coefficient ρ was set to 0.

Similarly, for the scRNA-seq HPA database, we used Pearson correlation to measure gene co-

expression across normal cell clusters, with non-significant correlations set to $\rho = 0$, using a

BH-corrected p-value threshold of 0.05. For the GTEx TMT proteomics database, we first

removed any genes with more than 25% missing values across all tissue samples analyzed.

Then, we applied the Pearson correlation as above to assess protein co-expression across

normal tissue samples, again using a BH-corrected p-value threshold of 0.05 for significance

and setting $\rho = 0$ for any non-significant correlations.

1136

1137 For the HPA IHC database (numerical score data) and for the HPM proteomics database, given

1138 the low number of comparisons, we did not use Pearson correlation but instead calculated a

1139 modified odds ratio (OR) for each gene pair (g_i, g_j) as follows:

1140
$$OR_{mod(g_i,g_j)} = \frac{d}{b \times c}$$

1141 Where:

1142 • d is the number of tissues/cells where both g_i and g_j have z-score > 1

1143 • b is the number of tissues/cells where g_i has z-score > 1 and g_j has z-score ≤ 1

1144 • c is the number of tissues/cells where g_j has z-score > 1 and g_i has z-score ≤ 1

1145

1146 In this case, pairs with mutually exclusive protein expression were considered those where

1147 $OR_{mod} = 0$. Of note, if both d and $b \times c$ were 0, the OR was set to 0.

1148

1149 Then, for each gene pair (g_i, g_j) we combined results from each normal tissue/cell database to

1150 obtain a score that estimates gene co-expression across normal tissues and cells, $N(g_i, g_j)$, as

1151 follows:

1152
$$N(g_i, g_j) = \sum_{k=1}^{2} r_{g_i,g_j}^{k} + \sum_{l=1}^{2} r_{g_i,g_j}^{l} + r_{g_i,g_j}^{m} + r_{g_i,g_j}^{n}$$

1153 Where:

1154 • r_{g_i,g_j}^k represents the score from the 2 bulk RNA-seq datasets and is calculated as:

1155
$$\begin{cases} +1 \ if \ \rho_{g_i,g_j} < 0 \ \text{and} \ p_{g_i,g_j} < 0.05 \\ -1 \times if \ \rho_{g_i,g_j} > 0 \ \text{and} \ p_{g_i,g_j} < 0.05 \\ 0 \ if \ p_{g_i,g_j} \geq 0.05 \ or \ if \ \text{either} \ g_i \ or \ g_j \ \text{is not in the dataset or if} \ g_i \ or \ g_j \ SD = 0 \end{cases}$$

1156 • r_{g_i,g_j}^l represents the score from the HPA IHC and the HPM proteomics datasets and is

1157 calculated as:

1158
$$\begin{cases} +2 \ if \ OR_{mod(g_i,g_j)} = 0 \\ -2 \times if \ OR_{mod(g_i,g_j)} \neq 0 \\ 0 \ if \ \text{either} \ g_i \ or \ g_j \ \text{is not in the dataset} \end{cases}$$

1159 • r_{g_i,g_j}^m represents the score from the HPA scRNA-seq dataset and is calculated as:

1160
$$\begin{cases} +1.5 \ if \ \rho_{g_i,g_j} < 0 \ \text{and} \ p_{g_i,g_j} < 0.05 \\ -1.5 \times if \ \rho_{g_i,g_j} > 0 \ \text{and} \ p_{g_i,g_j} < 0.05 \\ 0 \ if \ p_{g_i,g_j} \geq 0.05 \ or \ if \ \text{either} \ g_i \ or \ g_j \ \text{is not in the dataset or if} \ g_i \ or \ g_j \ SD = 0 \end{cases}$$

1161 • r_{g_i,g_j}^n represents the score from the GTEx proteomics dataset and is calculated as:

1162
$$\begin{cases} +2 \ if \ \rho_{g_i,g_j} < 0 \ \text{and} \ p_{g_i,g_j} < 0.05 \\ -2 \times if \ \rho_{g_i,g_j} > 0 \ \text{and} \ p_{g_i,g_j} < 0.05 \\ 0 \ if \ p_{g_i,g_j} \geq 0.05 \ or \ if \ \text{either} \ g_i \ or \ g_j \ \text{is not in the dataset or if} \ g_i \ or \ g_j \ SD = 0 \end{cases}$$

1163 • ρ_{g_i,g_j} is the Pearson correlation coefficient in a dataset between the two genes in a pair

1164 and p_{g_i,g_j} is the respective BH-corrected p-value.

1165

1166 The raw score $N(g_i, g_j)$ was then normalized and scaled as follows:

1167

1168
$$N_{norm}(g_i, g_j) = -\left\{ \left(\frac{N(g_i, g_j) - min(N)}{max(N) - min(N)} \times 2 \right) - 1 \right\}$$

1169 Where:

1170 • $N(g_i, g_j)$ is the raw score for a given gene pair (g_i, g_j)

1171 • min(N) and max(N) are the maximum and minimum raw scores across all gene pairs for

1172 a given STS histotype, respectively.

1173

1174 This normalization ensures that a score between 0 and 1 indicates higher likelihood of gene co-

1175 expression across normal tissues/cells and conversely a score between -1 and 0 indicates a

1176 lower likelihood of co-expression.

1177

1178 Finally, for each gene pair (g_i, g_j) we calculated an 'AND' logic-gated Targetability Score, called

1179 $TS_{,AND,}$ score, as follows:

1180
$$TS_{,AND,}(g_i, g_j) = \rho_{Tumor}(g_i, g_j) - N_{norm}(g_i, g_j)$$

1181 Gene pairs with $TS_{,AND,}$ score ≥ 1 were considered to have higher fitness as targets for an

1182 'AND'-gated CAR design. The results of the pipeline by STS histotype are reported in **Table**

1183 **S17**.

1184

Logic-gated 'AND–NOT' CAR Targets Discovery Pipeline

To discover gene pairs for an 'AND–NOT' bispecific CAR design, we applied a similar logic to

that used for discovery of gene pairs for an 'AND'-gated CAR. In this case, we looked for genes

with high expression in sarcoma cells that would function as activators, and for genes with low

to no expression in sarcoma cells but high co-expression with the paired activator in normal

cells/tissues that would function as inhibitors, thus preventing on-target off-tumor toxicity.

1191

To do this, first we obtained a list of highly expressed sarcoma up-regulated genes for each STS

histotype, removing any genes up-regulated in normal cells more often than STS cells, as

described above, and considered these as candidate activators. Then, to define a list of

possible inhibitors with little to no expression in sarcoma cells, for each STS histotype we

selected surfaceome genes with scRNA-seq expression below the 50th percentile among the

sarcoma down-regulated genes as defined in our differential gene expression pipeline. Single

cell RNA expression was again measured as the average log$_2$(CPM+1) across all the tumor

cells in a sarcoma histotype.

1200

We then calculated all the possible unique pairs between the candidate activators and inhibitors

and measured their co-expression in normal cells/tissues using the same datasets used for the

'AND'-gated CAR targets discovery pipeline. Specifically, for the GTEx and HPA bulk RNA-seq

datasets, as well as for the scRNA-seq HPA and the GTEx TMT proteomics datasets we used

Pearson correlation to measure co-expression of the two genes in each pair, and applied BH

multiple hypothesis testing correction setting the threshold for significance to a p-value < 0.05.

For non-significant correlations, we then set $\rho = 0$. For the GTEx TMT proteomics dataset,

again we first removed any genes with more than 25% missing values across all tissue samples

analyzed before performing correlation testing.

1210

1211 For the HPA IHC and the HPM proteomics databases, instead of performing Pearson correlation

1212 analysis we calculated a modified odds ratio (OR) for each gene pair (g_a, g_i) between an

1213 activator, g_a, and an inhibitor, g_i as follows:

1214
$$OR_{mod(g_a,g_i)} = \frac{d}{b}$$

1215 Where:

1216 • d is the number of tissues where both the activator g_a and the inhibitor g_i have z-score

1217 > 1

1218 • b is the number of tissues where the activator g_a has z-score > 1 and the inhibitor g_i has

1219 z-score ≤ 1

1220

1221 Pairs where $OR_{mod} = Inf$ were considered as suitable for an 'AND–NOT' CAR design. In cases

1222 where both d and b were 0, we set OR_{mod} to Inf.

1223

1224 Then, for each gene pair (g_a, g_i) we combined results from each normal tissue/cell database to

1225 calculate an 'AND–NOT' logic-gated Targetability Score, $TS_{,AND-NOT,}$ as follows:

1226
$$TS_{,AND-NOT,}(g_a, g_i) = \sum_{k=1}^{2} S^k_{g_a,g_i} + \sum_{l=1}^{2} S^l_{g_a,g_i} + S^m_{g_a,g_i} + S^n_{g_a,g_i}$$

1227 Where:

1228 • $S^k_{g_a,g_i}$ represents the score from the 2 bulk RNA-seq datasets and is calculated as:

1229
$$\begin{cases} +1 \times \left(1 + I_{g_a,g_i}\right) if \ \rho_{g_a,g_i} > 0 \ and \ p_{g_a,g_i} < 0.05 \\ -1 \times \left(1 + A_{g_a,g_i}\right) if \ \rho_{g_a,g_i} < 0 \ and \ p_{g_a,g_i} < 0.05 \\ 0 \ if \ p_{g_a,g_i} \geq 0.05 \ or \ if \ either \ g_a \ or \ g_i \ is \ not \ in \ the \ dataset \ or \ if \ g_a \ or \ g_i \ SD = 0 \end{cases}$$

1230 • $S^l_{g_a,g_i}$ represents the score from the HPA IHC and the HPM proteomics datasets and is

1231 calculated as:

1232
$$\begin{cases} +2 \times \left(1 + I_{g_a,g_i}\right) if \ OR_{mod(g_a,g_i)} = Inf \\ -2 \times \left(1 + A_{g_a,g_i}\right) if \ OR_{mod(g_a,g_i)} \neq Inf \\ 0 \ if \ either \ g_a \ or \ g_i \ is \ not \ in \ the \ dataset \end{cases}$$

1233 • $S^m_{g_a,g_i}$ represents the score from the HPA scRNA-seq dataset and is calculated as:

1234
$$\begin{cases} +1.5 \times \left(1 + I_{g_a,g_i}\right) if \ \rho_{g_a,g_i} > 0 \ and \ p_{g_a,g_i} < 0.05 \\ -1.5 \times \left(1 + A_{g_a,g_i}\right) if \ \rho_{g_a,g_i} < 0 \ and \ p_{g_a,g_i} < 0.05 \\ 0 \ if \ p_{g_a,g_i} \geq 0.05 \ or \ if \ either \ g_a \ or \ g_i \ is \ not \ in \ the \ dataset \ or \ if \ g_a \ or \ g_i \ SD = 0 \end{cases}$$

1235 • $S^n_{g_a,g_i}$ represents the score from the GTEx proteomics dataset and is calculated as:

1236
$$\begin{cases} +2 \times \left(1 + I_{g_a,g_i}\right) if \ \rho_{g_a,g_i} > 0 \ and \ p_{g_a,g_i} < 0.05 \\ -2 \times \left(1 + A_{g_a,g_i}\right) if \ \rho_{g_a,g_i} < 0 \ and \ p_{g_a,g_i} < 0.05 \\ 0 \ if \ p_{g_a,g_i} \geq 0.05 \ or \ if \ either \ g_a \ or \ g_i \ is \ not \ in \ the \ dataset \ or \ if \ g_a \ or \ g_i \ SD = 0 \end{cases}$$

• I_{g_a,g_i} is the proportion of comparisons in a dataset where the expression of the inhibitor

is equal or above that of the activator ($z - score_{g_i} \geq z - score_{g_a}$)

• A_{g_a, g_i} is the proportion of comparisons in a dataset where the expression of the activator

is above that of the activator ($z - score_{g_a} > z - score_{g_i}$)

• ρ_{g_a, g_i} is the Pearson correlation coefficient in a dataset between the activator and the

inhibitor and p_{g_a, g_i} is the respective BH-corrected p-value.

1243

1244 The $TS_{\text{'AND-NOT'}}$ scores were then normalized as follows:

1245
$$TS_{\text{'AND-NOT'}}(g_a, g_i) = \left\{ \left(\frac{TS_{\text{'AND-NOT'}}(g_a, g_i) - min(TS_{\text{'AND-NOT'}})}{max(TS_{\text{'AND-NOT'}}) - min(TS_{\text{'AND-NOT'}})} \times 2 \right) - 1 \right\}$$

Where:

• $TS_{\text{'AND-NOT'}}(g_a, g_i)$ is the raw score for a given gene pair (g_a, g_i)

• $min(TS_{\text{'AND-NOT'}})$ and $max(TS_{\text{'AND-NOT'}})$ are the maximum and minimum raw scores

across all gene pairs for a given STS histotype, respectively.

In each sarcoma histotype, gene pairs with values above the 99[th] percentile of the normalized

$TS_{\text{'AND-NOT'}}$ score were considered as those suitable for an 'AND-NOT'-gated CAR design. The

results of the pipeline by STS histotype are reported in **Table S18**.

1253

HPA Single Cell Gene Clusters and Signatures

HPA categorizes genes based on their expression profile across different normal cell types in

their scRNA-seq dataset[144]. Specifically, genes are categorized into 80 clusters, with one or

more clusters corresponding to a specific cell type. We downloaded the gene cluster

assignments directly from the HPA website and grouped gene clusters specific to a given cell

type together to obtain cell type-specific gene signatures that were used to annotate genes in

each bispecific CAR target pair. Also, the expression of these gene signatures in our STS

scRNA-seq data was measured using the Seurat function AddModuleScore and shown in

Figure S7. The HPA single cell gene clusters and signatures are provided in the Supplementary

Data.

1264

T cell receptor-engineered T cell and Peptide-Centric CAR Target Discovery Pipeline

To identify targets for TCR-T or PC-CAR-T cells in STS, we obtained a list of up-regulated

genes for each of the STS histotype studied. This was defined as any gene, either coding for

intracellular or surface proteins, that was up-regulated in at least one of the DET comparisons

made. Within this gene pool we then identified genes with high STS expression, defined in this

case as those with expression $\geq 25^{th}$ percentile in both our scRNA-seq data and in the SARC

TCGA samples, or those genes with expression $\geq 25^{th}$ percentile in both the sarcoma HPA cell

lines and our scRNA-seq data. Genes without expression data in SARC TCGA or sarcoma HPA

cell lines but with scRNA-seq expression $\geq$ the 25^{th} percentile were also considered to have

high STS expression.

1275

Then we compared the expression in normal cells and tissues of each of the up-regulated

genes to that of known TCR-T cell therapy targets that are currently in clinical development

identified through a search on clinicaltrials.gov with the term "solid tumor" for the

Condition/disease field, and "TCR-T cells" or "cellular therapy" for the Intervention/treatment

field (**Table S22**). Specifically, only genes belonging to the cancer germline antigen (CGA) class

were chosen as reference due to their preferential expression in immune-privileged sites like the

gonads.

1283

1284 Normal cell and tissue expression was measured again using the HPA scRNA-seq dataset

1285 (average cell type expression), the HPA and GTEx bulk RNA-seq datasets (average tissue

1286 group expression), the GTEx TMT proteomics dataset (average tissue expression), the HPM

1287 LC-MS/MS proteomics dataset, or the HPM IHC data, obtained as previously described.

1288

1289 For each gene (g) we calculated a summative score $N(g)$, representing its normal cell/tissue

1290 RNA and protein expression as compared to that of the CGAs used as TCR-T cell targets, and

1291 calculated as follows:

$$N(g) = \sum_{i=1}^{2} t_i + \sum_{j=1}^{3} t_j + t_k$$

1292

Where:

• t_i is the score calculated by comparing the candidate target expression to that of the

reference targets in the HPA and GTEx bulk RNA-seq datasets and is calculated as $+1$ if

the candidate has expression below the maximum expression value for the reference

targets in each tissue group, 0 if the candidate has expression above the maximum

value for the reference targets in at least one of the tissue groups or if the gene is not

present in the parent dataset;

• t_j is the score calculated by comparing the candidate target expression to that of the

reference targets in the HPA IHC, GTEx, and HPM proteomics datasets and is

calculated as $+2$ if the candidate has expression below the maximum expression value

for the reference targets in each tissue or tissue group, 0 if the candidate has expression

above the maximum value for the reference targets in at least one of the tissues or

tissue groups or if the gene is not present in the parent dataset;

• t_k is the score calculated by comparing the candidate target expression to that of the

reference targets in the HPA scRNA-seq dataset and is calculated as $+1.5$ if the

candidate has expression below the maximum expression value for the reference targets

in each cell type and 0 if the candidate has expression above the maximum value for the

reference targets in at least one of the cell types or if the gene is not present in the

parent dataset.

1312

Genes with a $N(g)$ score $\geq$ 3.5 and with high STS expression as defined above were retained

for further analysis. Due to the low number of genes with $N(g) \geq$ 3.5 meeting the combined high

expression criteria, possibly related to RNA expression dilution in bulk tissue analysis (TCGA

SARC) and differences between in vivo and in vitro analysis (HPA cell lines), genes that had

$N(g)$ score $\geq$ 3.5 and scRNA-seq expression $\geq$ the 25[th] percentile, regardless of TCGA SARC

or HPA cell line expression, were also considered for further analysis as below (**Table S23**).

1319

Global HLA Allele Frequency Estimation

To estimate the global HLA allele frequencies at the HLA-B, HLA-C, and HLA-A loci, we

downloaded allele frequencies for all the population studies available in the Allele Frequency

Net Database (AFND)[77]. Then, we used the python package HLAfreq[145] to combine allele

frequencies across populations, first within countries and then across countries to obtain an

average world allele frequency. Analyses were ran using default parameters. The population

size of each country was use as weight in the global allele frequency calculation, as described

by the developer. Population sizes by country updated to 2022 were downloaded from the World

Bank (https://data.worldbank.org/indicator/SP.POP.TOTL?most_recent_year_desc=false), or

from the United Nations Populations Division (Martinique, Turkey, Taiwan), or from the European

Union Commission (Madeira, Azores).

1331

Peptide-MHC targetability assessment

Next for each candidate target antigen, we used the MSiC model from HLAthena[74] to infer

presentation and binding to HLA class I molecules of all the possible 9-mer and 10-mer peptides

derived from each target. The amino acid sequence of each candidate target in FASTA format

was obtained from UniProt[146]. For our analysis, we chose the top 10 most frequent HLA class I

alleles by geographical region worldwide as reported in the Allele Frequency Net Database

(AFND)[77] https://www.allelefrequencies.net/top10freqs.asp, and then applied DeepImmuno[75] to

measure the predicted immunogenicity, the ability to elicit a CD8[+] immune response, of each

peptide-MHC pair ($pMHC$). Then, for each HLA allele we calculated the average worldwide

allele frequency across all the populations in which that allele was detected in the AFND (**Table**

S26). Finally, for each $pMHC$ of each candidate target gene *g*, a targetability score, $TS_{TCR/PC}$,

was calculated to assess its suitability as a target for either TCR-T or PC-CAR-T cells as

follows:

1345

1346 $$TS_{TCR/PC}(pMHC_i) = \left(S_{HLAthena} + S_{DeepImmuno}\right) \times \left(1 + F_{allele}\right)$$

1347

Where $pMHC_i$ represents the i^{th} $pMHC$ pair of a target gene *g*, $S_{HLAthena}$ is the score from the

MSiC HLAthena model for $pMHC_i$, $S_{DeepImmuno}$ is the immunogenicity score assigned by

DeepImmuno for $pMHC_i$, and F_{allele} is the average worldwide frequency for the HLA allele in

$pMHC_i$. For each target antigen, we then ranked peptides based on their cumulative HLA allele

frequency calculated as the sum of the allele frequencies of the HLA alleles in all of the $pMHCs$

formed by each peptide, and then retained for cross-reactivity assessment only those $pMHCs$

originating from peptides with the top five highest cumulative HLA allele frequency, and with

$TS_{TCR/PC} \geq$ the 99[th] percentile, and $S_{HLAthena} \geq 0.85$. The results for HLAthena and

DeepImmuno for all the target antigens, as well as the $TS_{TCR/PC}$ scores, are available in the

supplementary data.

1358

Peptide-MHC cross-reactivity assessment

For the $pMHCs$ of each target antigen selected as described above, we assessed the likelihood

of cross-reactivity with $pMHCs$ derived from other proteins expressed in normal cells and

tissues. To do this, we developed an algorithm called PS-TRACT (<u>P</u>eptide <u>S</u>imilarity and <u>T</u>CR

R̲eactivity A̲nalysis for C̲ellular T̲herapies), that considers the similarity of each tumoral $pMHC$ to

a pool of normal $pMHCs$ of reference while accounting for the likelihood of presentation of the

cross-reactive $pMHCs$ in normal tissues. Specifically, first we obtained a list of $pMHCs$

experimentally detected in normal tissues/cells (*observed normal ligandome*) by integrating data

from IEDB[147], systeMHC[148], and the HLA Ligand Atlas[149]. For IEDB, the full dataset was

downloaded from https://iedb.org/database_export_v3.php (file: "mhc_ligand_full.csv.csv"), and

epitopes were filtered to retain only linear peptides derived from human proteins and binding to

MHC class I, with any qualitative binding measurement other than "Negative", and of either 9 or

10 amino acids in length. Epitopes where MHC binding was measured with any of the following

methods were also removed: half maximal effective concentration (EC50), dissociation constant

KD (~EC50), and dissociation constant KD (~IC50). Lastly, only epitope binding data for alleles

in standard four-digit typing HLA allele nomenclature were retained. For the HLA Ligand Atlas,

the 2020.12 data release was obtained from https://hla-ligand-atlas.org/downloads. Epitope data

was filtered to retain only 9-mer and 10-mer peptides binding to HLA class I or HLA class I and

II, categorized as either strong or weak binders, and only those binding to HLA alleles reported

in standard four-digit typing nomenclature. For systeMHC, data for the following SYSMHCIDs

were obtained from https://systemhc.sjtu.edu.cn/datasets: SYSMHC00002, SYSMHC00003,

SYSMHC00004, SYSMHC00007, SYSMHC00026, SYSMHC00028, SYSMHC00029,

SYSMHC00031, SYSMHC00037, SYSMHC00039, SYSMHC00040, SYSMHC00044,

SYSMHC00047, SYSMHC00052, SYSMHC00053, SYSMHC00055, SYSMHC00057,

SYSMHC00064, SYSMHC00073, SYSMHC00088, and SYSMHC00089. In this case, only 9-

mer and 10-mer peptides binding HLA class I and derived from human proteins were retained.

To expand the pool of $pMHCs$ presented in normal tissues, we ran MHCflurry[150] for all the

possible 9-mer and 10-mer peptides derived from the whole human proteome (*predicted normal*

ligandome), using all the HLA alleles tested in our $pMHC$ targetability pipeline as described

above. Protein FASTA sequences for 20,409 proteins total were obtained from UniProt and

mhcflurry-predict-scan was used to obtain binding, processing, and presentation predictions,

with results filtered for affinity percentile < 2.0. Data from the observed and predicted normal

ligandomes were combined to obtain a unified non-redundant reference dataset of unique

$pMHC$/protein combinations (*unified normal ligandome*, supplementary data). Those $pMHCs$

with MHCflurry presentation score ≤ 0.5 were removed from the unified normal ligandome

unless they were listed in the observed ligandome. Then for each selected tumoral peptide we

measured its similarity with normal peptides accounting for both their physicochemical and HLA

binding properties. Specifically, given a tumoral peptide p_t and a normal reference peptide p_n

both binding to an HLA allele A, peptide similarity S_{sim} was calculated as:

1398

1399 $$S_{sim}(A, p_t, p_n) = \frac{simPMBEC - min(simPMBEC)}{max(simPMBEC) - min(simPMBEC)} + \frac{simAAindex - min(simAAindex)}{max(simAAindex) - min(simAAindex)}$$

1400

1401   Where $simPMBEC$ is a similarity score based on the peptide:MHC binding energy covariance

1402   (PMBEC) matrix[80] which measures amino acid similarity based on their effect on peptide HLA

1403   binding. Specifically, $simPMBEC$ was calculated as:

1404

$$\frac{\sum_{i=1}^{L}\left(\frac{1}{1 + distPMBEC(p_{ti}, p_{ni})} \times (1 + H_{Ai})\right)}{L}$$

Where $distPMBEC = max(PMBEC) - PMBEC$, $PMBEC$ is the value in the PMBEC matrix for the

pair of amino acids at the i^{th} position of p_t and p_n, L is the peptide length (9 or 10 amino acids),

and H_{Ai} is the entropy at the i^{th} peptide residue calculated for all the peptides of same length as

p_t binding to the HLA allele A in both the observed and predicted ligandome. This was obtained

using the MolecularEntropy function from the R package HDMD[151] (version 1.2), with values

ranging from 0 to 1, indicating respectively no amino acid sequence variability and maximum

sequence variability at the i^{th} peptide residue.

1413

1414 On the other end, $simAAindex$ was calculated as follows:

1415
$$\frac{\sum_{i=1}^{L}\left(\frac{1}{1+distAAindex(p_{ti},p_{ni})}\times(1+H_{Ai})\right)}{L}$$

Where $distAAindex$ is the Euclidean distance calculated between the pair of amino acids at the

i^{th} position of p_t and p_n, considering the first nine principal components of a PCA dimensionality

reduction of 554 physicochemical amino acid properties available from AAindex[79] and obtained

through the R package ftrCOOL[152] (version 2.0.0), L is the peptide length, and H_{Ai} is the residue

entropy calculated as above. The results of the peptide similarity analysis for the candidate

$pMHCs$ for each target antigen are available in the supplementary data.

1422

To define a significant threshold for S_{sim}, we tested PS-TRACT on known cross-reactive peptide

pairs[85] and found that a similarity score of 0.95 was the lowest observed for cross-reactive

peptides. Also, we ran PS-TRACT between the *MAGEA3*-derived peptide EVDPIGHLY and the

TTN-derived peptide ESDPIVAQY, which was found to be the off-tumor target leading to cases

of fatal myocarditis in patients that received EVDPIGHLY-targeted TCR-T cells[86], obtaining a

peptide similarity score of 1.19. Based on these results, we chose a threshold of $S_{sim} > 1.15$ as

this provided the highest specificity in identifying known cross-reactive peptides (**Figure S11A**).

Thus, after running PS-TRACT on each candidate tumoral $pMHC$, we selected only the

corresponding normal $pMHCs$ with $S_{sim} > 1.15$ for further cross-reactivity assessment.

Specifically, for each of these highly similar normal $pMHCs$ we calculated a tissue-specific

cross-reactivity score, S_{CR}, that considers the tissue abundance of the antigen from which the

cross-reactive peptide originates, the likelihood of presentation based on MHCflurry, and the

presence of the peptide in the observed ligandome:

1436

1437 $$S_{CR}(pMHC_y, pMHC_i)$$

1438
$$= (P_{MHCflurry} + 0.5) \times (E_{HPA/GTEx}) \times \begin{cases} 1.25 \ if \ pMHC_y \in observed \ ligandome \\ 1 \ if \ pMHC_y \notin observed \ ligandome \end{cases}$$

1439

Where $pMHC_y$ is the y^{th} cross-reactive $pMHC$ for a given tumoral $pMHC_i$, $P_{MHCflurry}$ is the

MHCflurry presentation score for $pMHC_y$, and $E_{HPA/GTEx}$ is the median-centered, z-scored RNA

expression for the parent antigen of the $pMHC_y$ peptide in the HPA and GTEx bulk RNA-seq

consensus datasets (source: https://www.proteinatlas.org/about/download, file:

rna_tissue_consensus.tsv.zip). Then for each tumoral $pMHC$ in each of the 50 normal tissues,

we selected the maximum S_{CR} score among all the filtered cross-reactive normal $pMHCs$, and

this vector of S_{CR} scores was used to define the cross-reactivity profile of the tumoral target

$pMHC$ considered.

1448

Finally, we benchmarked the cross-reactivity profile of each candidate tumoral $pMHC$ pair

against that of the $pMHC$ targeted by afami-cel (GVYDGREHTV–HLA-A*02:01) which was used

as reference given afami-cel's promising safety and tolerability profile observed in clinical

trials[27,28]. Specifically, for each candidate tumoral peptide-MHC pair $pMHC_i$, we calculated the

difference $S_{CR}Diff(i)$ between its S_{CR} cross-reactivity profile and that of GVYDGREHTV–HLA-

A*02:01, as follows:

1455

1456
$$S_{CR}Diff(i) = \frac{1}{n} \sum_{j=1}^{n} \begin{cases} \left((S_{CR}^{ij} - S_{CR}^{refj}) / \sigma_j \right) \times \alpha \ if \ S_{CR}^{ij} > S_{CR}^{refj} \\ (S_{CR}^{ij} - S_{CR}^{refj}) / \sigma_j \qquad otherwise \end{cases}$$

1457

1458 Where S_{CR}^{ij} is the S_{CR} score in the j^{th} tissue of $pMHC_i$, while S_{CR}^{refj} is the S_{CR} score in the j^{th}

1459 tissue of the reference $pMHC$ (GVYDGREHTV–HLA-A*02:01), n is the number of tissues

1460 considered, σ_j is the standard deviation of the S_{CR} scores for all the $pMHC$ pairs in the j^{th} tissue,

1461 and $\alpha = 2$. Target tumoral $pMHCs$ with differential S_{CR} profile compared to GVYDGREHTV–HLA-

1462 A*02:01 ($S_{CR}Diff$) ≤ 0 were considered to have acceptable risk of cross-reactivity.

1463

Clinical STS proteomics datasets

Two clinical STS MS proteomics datasets were used to validate the expression of the identified

cell therapy targets at the protein level, one from Tang et al.[91] and one from Burns et al.[92] The

dataset from Tang et al. was obtained from the supplementary material of the relative paper

("Proteomic_matrix(tumor)" and "Clinical_information" tables). The relative fraction of total (Fot)

protein intensities for the Tang et al. dataset were $\log_2(+1)$ transformed and then median

centered across samples and z-scored across all proteins for standardization. Protein symbols

were standardized to the current HGNC symbols before analysis. The proteomics dataset from

Burns et al. was obtained from the supplementary material (Supplementary Data 2). The

expression matrix in this case was used as is given that protein expression values were already

log-transformed, and median centered across samples and z-scored across proteins. Protein

symbols were standardized to the current HGNC symbols before analysis. For SCAN-ACT

implementation in other cancers, we obtained pre-processed mass spectrometry-based

proteomics data from either LinkedOmicsKB[153], the Proteomic Data Commons (PDC)[154], or from

a literature search (**Table S34**). Most of the data was derived from studies conducted by Clinical

Proteomic Tumor Analysis Consortium (CPTAC) or the International Cancer Proteogenome

Consortium (ICPC). For data obtained from LinkedOmicsKB, which was provided as $\log_2$

reference-normalized protein abundance, we applied median-centering across samples and z-

scoring across proteins. In contrast, for data from the PDC, where the $\log_2$ reference-normalized

protein abundance was already standardized across proteins, we performed only median-

centering across samples. For the gastric adenocarcinoma dataset, expression data from

different technical replicates of the same patient was averaged before analysis. For other

cancers, expression data provided in the supplementary materials of the related publications

was used[155–158]. For the prostate cancer dataset, which was provided as log2 reference-

normalized protein abundance, we applied median-centering across samples and z-scoring

across proteins. For the melanoma and esophageal cancer datasets, only median-centering

across samples was necessary, as the data was already standardized across proteins. The

bladder cancer expression data was used as provided by the authors, as it was already

standardized across both samples and proteins. Prior to analysis, gene symbols across all

datasets were updated to the current HGNC version. Proteins with missing values in 50% or

more samples were removed, and the remaining missing values were imputed with a value of

1e-6. For genes with multiple protein isoforms, the isoform with the highest expression was

used.

1497

STS cell line proteomics data

Protein expression data in soft tissue sarcoma cell lines was obtained from the Cancer Cell Line

Encyclopedia[93] (CCLE) and the ProCan-DepMapSanger repository. For CCLE, the normalized

$\log_2$ transformed relative protein intensities were obtained from

https://gygi.hms.harvard.edu/publica Shtions/ccle.html (file:

protein_quant_current_normalized.csv). Gene symbols were standardized to the current HGNC

symbols, and for different protein isoforms for a given gene symbol we retained the isoform with

maximum expression (supplementary data). Then, data was median-centered across all cell

lines and z-scored within genes before use for downstream analysis. The following "soft tissue"

cell lines were considered in the CCLE dataset for further analysis: "A-204", "RD", "RH-41", "HT-

1080", "RH-30", and "KYM-1". The ProCan-DepMapSanger dataset was obtained from the

figshare repository associated with the corresponding publication:

https://doi.org/10.6084/m9.figshare.19345397 (file: ProCan-

DepMapSanger_protein_matrix_8498_averaged.txt). Similarly to the CCLE dataset, first we

standardized gene symbols to the current HGNC nomenclature, then for genes with more than

one protein isoform we retained the isoform with highest expression (supplementary data).

Since the data was already provided as log_2 transformed relative protein intensities, the protein

per cell line expression matrix was median-centered across cell lines and z-scored within genes

before further analysis. The following soft tissue sarcoma cell lines were considered for

downstream analysis: "A204" , "ESS-1", "G-402", "GCT", "Hs-633T", "HT-1080" , "KYM-1",

"MES-SA", "MFH-ino", "RD", "RH-18", "RH-41", "RKN", "SJRH30", "SK-LMS-1", "SK-UT-1" ,

"SKN", "SW684", "SW872", "SW982", "TE-441-T", and "VA-ES-BJ".

1520

IHC of Selected CAR-T Targets

IHC was performed by the Stanford Human Pathology/Histology Service Center for ACVR1,

AXL, CLMP, GPC6, KITLG, and MRC2. Previously obtained tissue microarrays constructed

using one to seven 1-mm FFPE cores per patient were used for the analysis, including UPS,

LPS, LMS, SS, and MPNST samples. The following antibodies were used for the analysis:

ACVR1 Rabbit IgG (cat. HPA007505, Sigma Aldrich, MO, USA), AXL Rabbitt IgG (cat.

HPA037422, Sigma Aldrich, MO, USA), GPC6 (cat. HPA017671, Sigma Aldrich, MO, USA),

KITLG (cat. HPA070395, HPA017671), CLMP (cat. 16127-1-AP, Thermo Fisher Scientific, MA,

USA), and MRC2 (cat. HPA041991, Sigma Aldrich, MO, USA). All the antibodies were used at a

dilution titer of 1:50 except for the CLMP antibody (1:400), and the KITLG antibody which was

used at a 1:100, 1:200, or 1:50 dilution depending on the TMA analyzed. Tumor cell staining

intensity was scored using a three-point scale (0, negative; 1, weakly positive; and 2, strongly

positive). Staining for FAP was performed at the UCLA pathology department on a tissue

microarray of 60 undifferentiated pleomorphic sarcomas in triplicate (no previous treatment), of

which 57 samples were evaluable. FAP immunostaining was performed on Leica Bond platform

using a polyclonal rabbit anti-FAP alpha antibody (Abcam, ab207178). All slides were scanned

at 20X magnification. The percentage of FAP expressing cells was quantified using the HALO®

image analysis platform based on automatic cell segmentation (Indica Labs). Missing or folded

tissue cores were excluded. Replicate cores were averaged.

1540

Measurement of peptide-MHC binding through IMMUNE

Plasmid construction

To construct plasmids driving the expression of non-covalent HLA-I heterodimers, the

commercially available backbone vector pRS315 was modified[89]. Briefly, the entire expression

cassette containing genes encoding the Aga2 leader sequence, the ectodomain of a particular

HLA-I heavy chain allele (A*02:01 or A*11:01 or B*35:01 or C*12:03), the Aga2p yeast native

protein subunit, an HA epitope tag, and the MFα terminator was cloned to the downstream of

the GAL10 promoter in pRS315, so that the expressed HLA-I heavy chain ectodomain was

fused to the N-terminus of Aga2p for yeast surface display. Subsequently, a second expression

cassette containing genes encoding the Aga2 leader sequence, β2m, a C-myc epitope tag and

the MFα terminator was cloned to the downstream of the GAL1 promoter in the same pRS315-

based vector, allowing the secretion of the β2m domain for HLA heavy chain/β2m dimerization

and surface detection. In addition, the Aga2p subunit was further removed from the above

plasmids, which resulted in control plasmids yielding no HLA detection on the surface of yeast.

Gene synthesis and subcloning services were provided by GenScript (Jiangsu, China).

1556

Electrotransformation of yeast cells and expression of HLA

The *Saccharomyces cerevisiae* yeast parent strain, EBY100[159] was used for generation of

daughter strains displaying each of the aforementioned HLA alleles. The constructed plasmids

were transformed into EBY100 one at a time by electroporation, following the MicroPulser

electroporation mannual (BioRad). Transformed cells that received the corresponding plasmid

were recovered on selected dextrose-based (SD) medium agar plates minus the nutrient

corresponding to the selection marker gene within the plasmid. Single colonies on SD agar

plates appeared after 2-3 days of incubation at 30°C. To induce yeast expression of target

proteins, a single yeast colony was inoculated with liquid SD medium and cultured in a shaking

tube at 30°C until it reached the exponential growth phase. A sufficient amount of yeast cells

was then transferred from the SD medium to selected galactose-based (SG) medium for protein

expression under the promotion of GAL1 or GAL10. Typically, an overnight shaking and

induction at 30°C can promote effective expression of the protein of interest in yeast for surface

detection by flow cytometry.

1571

Immunofluorescent labeling and flow cytometric analysis

A sufficient amount of galactose-induced yeast cells was collected by centrifugation and labeled

by primary antibodies in PBS + 0.5% bovine serum albumin (BSA) for 1 h on ice. The primary

antibody was rabbit anti-HA tag monoclonal antibody (mAb clone C29F4, Cell signaling), mouse

anti-human β2m mAb (clone 2M2, Biolegend), or mouse anti-HLA-A, B, C conformation-specific

mAb (clone W6/32, Biolegend). After the primary staining, the cells were washed at least twice

with ice cold PBS + 0.5% BSA and labeled by secondary antibodies on ice for another 1 h. The

secondary antibody was Alexa Fluor 488 conjugated goat anti-rabbit IgG (H+L) highly cross-

adsorbed secondary antibody, Alexa Fluor 488 or Alexa Fluor 647 conjugated goat anti-mouse

IgG (H+L) highly cross-adsorbed secondary antibody (Thermo Fisher Scientific). After the

secondary staining, cells were washed with ice cold PBS + 0.5% BSA and resuspended in PBS

+ 0.5% BSA for flow cytometric analysis. A FACSCalibur flow cytometer (BD biosciences) was

used to acquire flow data. At least 10000 cell events were collected for each sample and gated

through forward and side scattering. FL1 and FL4 fluorescence channels were selected for

Alexa Fluor 488 and Alexa Fluor 647, respectively, according to the fluorophore selected at the

secondary labeling step as mentioned above. Flowjo (BD biosciences) was used for data

analysis.

1589

Peptide design and synthesis

To design indicator peptides appropriate for evaluation of HLA peptide binding on the surface of

yeast cells and for downstream competitive assay development, the reported epitope repertoire

for each aforementioned HLA allele from the Immune Epitope Database (https://www.IEDB.org)

was carefully analyzed. First, an epitope peptide was considered only if it contained a cysteine

residue at P4 or P5 or P6 anchor position available for site-specific modification, i.e.,

biotinylation or fluorophore conjugation. After that, the chosen peptide was further analyzed and

validated 1) based on its predicted %Rank_EL (<2) and %Rank_BA scores (<2) from the

NetMHCpan-4.1 webserver, and 2) based on its sequence similarity to the peptides with

available X-ray structure in complex with the given HLA. For example, Bio-KLT (amino acids:

KLTPL{Cys-Biotin}VTL), Bio-AIF (aa: AIFQ{Cys-Biotin}SMTK), Bio-RYP (aa: RYPV{Cys-

Biotin}KFPSL), and Bio-FAF (aa: FAF{Cys-Biotin}RITSF) were designed as the indicator peptide

for A*02:01, A*11:01, B*35:01, and C*12:03, respectively. All peptides were synthesized

(GenScript) with over 95% purify.

On yeast peptide binding

About half a million yeast cells displaying a peptide-receptive HLA were collected by

centrifugation, resuspended in 40 µl of PBS, and incubated with 0.5 uM biotinylated indicator

peptides ([peptide]>>[HLA]) overnight prior to fluorescent labeling for flow cytometry. After

peptide binding reached equilibrium, yeast cells were washed twice with ice-cold PBS + 0.5%

BSA and then simultaneously stained for bound biotinylated indicator peptides and HLA

molecules on the surface using Alexa Fluor 647 conjugated streptavidin and HLA conformation-

specific mAb W6/32 followed by Alexa Fluor 488 conjugated goat anti-mouse IgG (H+L) highly

cross-adsorbed secondary antibody (Thermo Fisher Scientific). Double stained cells were then

washed and resuspended in ice-cold PBS + 0.5% BSA for analysis on a BD FACSCalibur. Flow

cytometric data was analyzed using FlowJo.

1615

On yeast peptide competition

To evaluate the competitive binding of a test peptide against the indicator peptide to HLA on the

surface of yeast cells, about half a million yeast cells were collected and incubated with the

biotinylated peptide as mentioned above, but with addition of a 20-fold more concentrated test

peptide in its non-biotinylated format. The peptide competition assay was also performed under

a similar condition as mentioned above, allowing reaction to reach equilibrium. Yeast cells were

then collected, washed, and co-stained for bound biotinylated indicator peptides and HLA

molecules using the aforementioned fluorescent labeling method for flow cytometric analysis.

The %Binding in the presence of competitors was quantified as $[(MFI_{with\ competitor}-BG)/(MFI_{no}$

$_{competitor}-BG)]\times 100\%$, where $MFI_{with\ competitor}$ represents the average fluorescence intensity in the

presence of test peptide (a competitor), $MFI_{no\ competitor}$ represents the average fluorescence

intensity in the absence of competitors, and BG represents the background fluorescence

intensity. The %Competition was calculated as 100%−%Binding to reveal binding of the test

peptide competitor to HLA molecules displayed on yeast. MFI was determined using

appropriately gated cell populations.

1631

Target Priority Index

For each target discovery pipeline, we calculated a final Target Priority Index (TPI) that

integrates normal tissues/cells target expression, scRNA-seq tumor expression and

homogeneity, STS protein expression, DepMap sarcoma cell line essentiality, incidence of

SNVs/Indels, gene fusions and CNAs in STS involving the target considered, and the presence

of clinically or pre-clinically developed immunotherapeutic agents against each target (for

monospecific and bispecific CAR-T cells targets only).

1639

1640 Homogeneity of expression at the single-cell level in each STS was calculated as the product of

1641 the average tumor expression in $\log_2$(CPM+1) and the percentage of tumor cells with

1642 $\log_2$(CPM+1) > 0, which we defined as H_{RNA} score:

1643
$$H_{RNA} = \overline{(log_2(CPM + 1))} \times \left(\frac{n\ cells\ log_2(CPM + 1) > 0}{total\ cells} \times 100 \right)$$

To assess if a gene has been previously studied as an immunotherapy target, we focused on

therapeutic monoclonal antibodies (mAb), ADCs, single-chain variable fragments (scFv), fusion

proteins, nanobodies, and fragment antigen binding regions (Fab), as well as CAR-T cells.

Specifically, we downloaded a list of clinically developed mAbs, ADCs, scFvs, Fabs, nanobodies

and fusion proteins from Thera-SAbDab[160] (https://opig.stats.ox.ac.uk/webapps/sabdab-

sabpred/therasabdab, accessed on 07/16/2024, Supplementary Data), retaining only agents

developed for human use, and removing all those agents with clinical development phase

reported as "TBC", "Preregistration", or "Preregistration (w)". We complemented Thera-SAbDab

with a PubMed search for pre-clinically developed therapeutic antibodies or ADCs against any

genes in the surfaceome, using the following search query: "gene AND antibody, OR gene AND

antibody drug conjugate, OR gene AND ADC OR gene AND antibody-drug conjugate" (**Table**

S30). Targets with clinical development phase reported as "TBC", "Preregistration", or

"Preregistration (w)" in Thera-SAbDab were included as pre-clinical targets, unless they were

described as clinical targets. As far CAR-T cells, for clinically developed targets we used the

previously defined list of targets as per MacKay et al[45], while for pre-clinically developed targets

we performed a PubMed search for all the genes in the surfaceome using the following search

query: "gene AND CAR-T OR gene AND chimeric antigen receptor T-cells OR gene AND

chimeric antigen receptor", **Table S30**).

1662

1663 *Monospecific CAR-T Cell Target TPI Calculation*

1664   To obtain the TPI for monospecific CAR-T targets, we used the list of targets defined based on

1665   the TS_{CAR} as described above. Then, for these targets we normalized the TS_{CAR} to assume

1666   values between 0 and 1 as follows:

1667

$$TS_{CAR}(g)(norm) = \frac{TS_{CAR}(g) - \min(TS_{CAR})}{\max(TS_{CAR}) - \min(TS_{CAR})}$$

1668   Where $TS_{CAR}(g)$ is the TS_{CAR} for a target g, and $\min(TS_{CAR})$ and $\max(TS_{CAR})$ represents the

1669   minimum and maximum TS_{CAR} values, respectively, for all the final targets being considered for

1670   the STS histotype analyzed.

1671

1672   Then, for each target g, we obtained a summative validation score, V_{CAR}, calculated as the sum

1673   of different scores assigned for each validation feature, defined as follows:

1674   •   If the H_{RNA} score for g is equal or greater than the average H_{RNA} for *CD276* across the

1675   five STS studied, g is assigned a score of 2, otherwise g is assigned a score of 0.

1676   *CD276* was used as reference as it has been clinically (NCT04433221) and pre-

1677   clinically[95] studied as a CAR-T target in STS.

1678   •   For each gene g and STS histotype we measured the percentage of samples in the STS

1679   proteomics Tang et al. dataset with medium-high (M-H) expression (z-score ≥ 0).

1680   DDLPS and WDLPS samples were considered as one group (LPS) for this analysis. If

1681   the percentage of samples of the STS analyzed with M-H expression for g was greater

1682   than or equal to the average percentage of samples with M-H expression for CD276

1683   across SS, MFS, UPS, LMS and LPS, then we assigned a score of 2, otherwise we

1684   assigned a score of 0.

1685   •   If g was reported as target for clinically developed mAbs, ADCs, scFv, Fabs, nanobodies

1686   or fusion proteins, we assigned a score of 1. If instead g was only studied preclinically as

1687   target for therapeutic antibodies or ADCs, we assigned a score of 0.5. Otherwise, we

1688   assigned a score of 0.

• If g was reported as target for clinically developed CAR-T cells, we assigned a score of

1, while we assigned a score of 0.5 if g was only explored as a CAR-T target

preclinically, and a score of 0 if it was never tested as a CAR-T target.

• If g resulted as essential (Chronos DepMap score ≤ -1) in sarcoma cell lines derived

from the STS histotype studied, we assigned a score of 1, otherwise we assigned a

score of 0.5 if g was essential in any sarcoma cell line, or a score of 0 if g was not

essential in any sarcoma cell lines.

• If g was involved in at least one gene fusion in the STS histotype studied we assigned a

score of 1, otherwise we assigned a score of 0.5 if g was involved in at least one gene

fusion in any STS, or a score of 0 if g was never involved in gene fusions in STS.

• If g was involved by at least one SNV or Indel in the STS histotype studied we assigned

a score of 1, otherwise we assigned a score of 0.5 if g was involved by at least one

SNV/Indel in any STS histotype, or a score of 0 if g was never involved by SNVs/Indels.

• If g was involved by CNAs in the STS histotype studied and there was a positive

correlation between its linear CNV level and RNA expression in TCGA, we assigned a

score of 1, otherwise if it was involved by CNAs either in TCGA or MSK-IMPACT in the

same STS histotype studied we assigned a score of 0.5. Instead, if g was never involved

by CNAs in STS we assigned a score of 0.

We then normalized V_{CAR} to assume values ranging from 0 to 1 as follows:

1708
$$V_{CAR}(g)(norm) = \frac{V_{CAR}(g) - \min(V_{CAR})}{\max(V_{CAR}) - \min(V_{CAR})}$$

1709 Finally, we calculated the TPI for each gene g as follows:

1710
$$TPI_{CAR} = TS_{CAR}(g)(norm) + V_{CAR}(g)(norm)$$

1711 The TPI for the monospecific CAR-T targets is reported in **Table S31**.

1712

Bispecific CAR-T Cells Targets TPI Calculation

For bispecific CAR-T cells targets we obtained a TPI for logic-gated 'AND' CAR-T cells

$(TPI_{,AND,})$, and a separate TPI for 'AND-NOT' CAR-T cells $(TPI_{,AND-NOT,})$. Specifically, we

defined suitable target pairs for both 'AND' and 'AND-NOT' CAR-T cells as described above,

and then for each 'AND' CAR-T target and 'AND-NOT' CAR-T activator we calculated the

average $TS_{,AND,}$ and $TS_{,AND-NOT,}$ scores considering all the target pairs in which that gene

appears. We then normalized the average $TS_{,AND,}$ and $TS_{,AND-NOT,}$ for each gene g to assume

values ranging from 0 to 1 as follows:

1721
$$TS_{,AND,}(g)(norm) = \frac{\overline{TS_{,AND,}}(g) - \min(\overline{TS_{,AND,}})}{\max(\overline{TS_{,AND,}}) - \min(\overline{TS_{,AND,}})}$$

1722

1723
$$TS_{,AND-NOT,}(g)(norm) = \frac{\overline{TS_{,AND-NOT,}}(g) - \min(\overline{TS_{,AND-NOT,}})}{\max(\overline{TS_{,AND-NOT,}}) - \min(\overline{TS_{,AND-NOT,}})}$$

1724 Then, for each target g we calculated $V_{CAR}(g)$ and $V_{CAR}(g)(norm)$ as described above, and

1725 finally obtained the TPIs as follows:

1726
$$TPI_{,AND,} = TS_{,AND,}(g)(norm) + V_{CAR}(g)(norm)$$

1727
$$TPI_{,AND-NOT,} = TS_{,AND-NOT,}(g)(norm) + V_{CAR}(g)(norm)$$

1728 The TPIs for the bispecific CAR-T targets are reported in **Table S32**.

1729

1730 *TCR and PC-CAR-T Cells Targets TPI Calculation*

1731 For each of the defined target antigens g for TCR-T and PC-CAR-T cells in each STS, we first

1732 normalized the $N(g)$ score, indicative of the normal tissue and cell expression of g as previously

1733 defined, to assume values between 0 and 1:

1734
$$N(g)_{norm} = \frac{N(g) - \min(N)}{\max(N) - \min(N)}$$

1735 Then, for each target g we obtained a summative validation score, $V_{TCR/PC}$, calculated as the

1736 sum of the following scores:

- 1737   •   If the H_{RNA} score for g is equal or greater than the H_{RNA} for *MAGEA4* in SS, g is
- 1738   assigned a score of 2, otherwise is assigned a score of 0. *MAGEA4* was used as
- 1739   reference in this case since it is the source antigen for the target of afami-cel, the first
- 1740   TCR-T cell therapy to receive FDA approval in soft tissue sarcomas.
- 1741   •   If g had a z-score ≥ 0 in at least two separate STS cell lines in either CCLE or in the
- 1742   ProCan-DepMapSanger datasets, we assigned a score of 1, otherwise we assigned a
- 1743   score of 0.
- 1744   •   If g resulted as essential (Chronos DepMap score ≤ -1) in sarcoma cell lines derived
- 1745   from the STS histotype studied, we assigned a score of 1, otherwise we assigned a
- 1746   score of 0.5 if g was essential in any sarcoma cell line, or a score of 0 if g was not
- 1747   essential in any sarcoma cell lines.
- 1748   •   If g was involved in at least one gene fusion in the STS histotype studied we assigned a
- 1749   score of 1, otherwise we assigned a score of 0.5 if g was involved in at least one gene
- 1750   fusion in any STS, or a score of 0 if g was never involved in gene fusions in STS.
- 1751   •   If g was involved by at least one SNV or Indel in the STS histotype studied we assigned
- 1752   a score of 1, otherwise we assigned a score of 0.5 if g was involved by at least one
- 1753   SNV/Indel in any STS histotype, or a score of 0 if g was never involved by SNVs/Indels.
- 1754   •   If g was involved by CNAs in the STS histotype studied and there was a positive
- 1755   correlation between its linear CNV level and RNA expression in TCGA, we assigned a
- 1756   score of 1, otherwise if it was involved by CNAs either in TCGA or MSK-IMPACT in the
- 1757   same STS histotype studied we assigned a score of 0.5. Instead, if g was never involved
- 1758   by CNAs in STS we assigned a score of 0.
- 1759   We then normalized $V_{TCR/PC}$ to assume values ranging from 0 to 1 as follows:

- 1760

$$V_{TCR/PC}(g)(norm) = \frac{V_{TCR/PC}(g) - \min(V_{TCR/PC})}{\max(V_{TCR/PC}) - \min(V_{TCR/PC})}$$

1761   Next, for each target antigen we considered the target $pMHCs$ defined based on their $TS_{TCR/PC}$

1762   as reported in **Table S27**. Specifically, for each target antigen g we calculated the average HLA

1763   allele frequency at each locus, by considering all the peptides derived from each antigen binding

1764   to HLA-A, HLA-B, or HLA-C class alleles, as follows:

1765
$$HLA_A(f_{TOT}) = \frac{\sum_{i=1}^{pA} HLA_A(f)}{pA}$$

1766
$$HLA_B(f_{TOT}) = \frac{\sum_{i=1}^{pB} HLA_B(f)}{pB}$$

1767
$$HLA_C(f_{TOT}) = \frac{\sum_{i=1}^{pC} HLA_C(f)}{pC}$$

1768   Where pA, pB, pC are respectively the number of $pMHC$ pairs including class A, class B and

1769   class C alleles for the target g, while $HLA_A(f), HLA_B(f)$, and $HLA_C(f)$ represent the HLA allele

1770   frequencies of the $i^{th} pMHC$ considered. We then obtained the total HLA allele frequency,

1771 $HLA_{TOT}(g)$ for each target g by adding $HLA_A(f_{TOT})$, $HLA_B(f_{TOT})$, and $HLA_C(f_{TOT})$, and

1772   normalized this to obtain values ranging from 0 to 1, as follows:

1773
$$HLA_{TOT}(g)(norm) = \frac{HLA_{TOT}(g) - \min(HLA_{TOT})}{\max(HLA_{TOT}) - \min(HLA_{TOT})}$$

1774   Next, for each target g we calculated the total number of $pMHC$ targets ($pMHC_{TOT}$) and the

1775   number of $pMHC$ targets with $S_{CR}Diff \leq 0$ ($pMHC_{SCR}$). We divided these by the target antigen

1776   total number of amino acids (L_{AA}) to account for differences in protein size, and then normalized

1777   them to obtain values ranging from 0 to 1, as follows:

1778
$$pMHC_{TOT} = \frac{n(pMHC_{TOT})}{L_{AA}}$$

1779
$$pMHC_{TOT}(g)(norm) = \frac{pMHC_{TOT}(g) - \min(pMHC_{TOT})}{\max(pMHC_{TOT}) - \min(pMHC_{TOT})}$$

1780

1781
$$pMHC_{SCR} = \frac{n(pMHC_{SCR})}{L_{AA}}$$

1782
$$pMHC_{SCR}(g)(norm) = \frac{pMHC_{SCR}(g) - \min(pMHC_{SCR})}{\max(pMHC_{SCR}) - \min(pMHC_{SCR})}$$

1783 Finally, we calculated the TPI for each target g as follows:

1784
$$TPI_{TCR/PC} = N(g)_{norm} + V_{TCR/PC}(g)(norm) + HLA_{TOT}(g)(norm) + pMHC_{TOT}(g)(norm)$$

1785
$$+ pMHC_{SCR}(g)(norm)$$

1786 The TPIs for the TCR-T and PC-CAR-T cells targets are reported in **Table S33**.

1787

References

1. Wagner, J., Wickman, E., DeRenzo, C., and Gottschalk, S. (2020). CAR T Cell Therapy for

Solid Tumors: Bright Future or Dark Reality? Mol. Ther. J. Am. Soc. Gene Ther. *28*, 2320–

2339. https://doi.org/10.1016/j.ymthe.2020.09.015.

2. Neelapu, S.S., Locke, F.L., Bartlett, N.L., Lekakis, L.J., Miklos, D.B., Jacobson, C.A.,

Braunschweig, I., Oluwole, O.O., Siddiqi, T., Lin, Y., et al. (2017). Axicabtagene Ciloleucel

CAR T-Cell Therapy in Refractory Large B-Cell Lymphoma. N. Engl. J. Med. *377*, 2531–

2544. https://doi.org/10.1056/NEJMoa1707447.

3. Maude, S.L., Laetsch, T.W., Buechner, J., Rives, S., Boyer, M., Bittencourt, H., Bader, P.,

Verneris, M.R., Stefanski, H.E., Myers, G.D., et al. (2018). Tisagenlecleucel in Children

and Young Adults with B-Cell Lymphoblastic Leukemia. N. Engl. J. Med. *378*, 439–448.

https://doi.org/10.1056/NEJMoa1709866.

4. Labanieh, L., and Mackall, C.L. (2023). CAR immune cells: design principles, resistance

and the next generation. Nature *614*, 635–648. https://doi.org/10.1038/s41586-023-

05707-3.

5. Adachi, K., Kano, Y., Nagai, T., Okuyama, N., Sakoda, Y., and Tamada, K. (2018). IL-7 and

CCL19 expression in CAR-T cells improves immune cell infiltration and CAR-T cell

survival in the tumor. Nat. Biotechnol. *36*, 346–351. https://doi.org/10.1038/nbt.4086.

6. Lynn, R.C., Weber, E.W., Sotillo, E., Gennert, D., Xu, P., Good, Z., Anbunathan, H., Lattin,

J., Jones, R., Tieu, V., et al. (2019). c-Jun overexpression in CAR T cells induces

exhaustion resistance. Nature *576*, 293–300. https://doi.org/10.1038/s41586-019-1805-

z.

7. Rafiq, S., Yeku, O.O., Jackson, H.J., Purdon, T.J., van Leeuwen, D.G., Drakes, D.J., Song,

M., Miele, M.M., Li, Z., Wang, P., et al. (2018). Targeted delivery of a PD-1-blocking scFv by

CAR-T cells enhances anti-tumor efficacy in vivo. Nat. Biotechnol. *36*, 847–856.

https://doi.org/10.1038/nbt.4195.

8. Narayan, V., Barber-Rotenberg, J.S., Jung, I.-Y., Lacey, S.F., Rech, A.J., Davis, M.M.,

Hwang, W.-T., Lal, P., Carpenter, E.L., Maude, S.L., et al. (2022). PSMA-targeting TGFβ-

insensitive armored CAR T cells in metastatic castration-resistant prostate cancer: a

phase 1 trial. Nat. Med. *28*, 724–734. https://doi.org/10.1038/s41591-022-01726-1.

9. Doan, A.E., Mueller, K.P., Chen, A.Y., Rouin, G.T., Chen, Y., Daniel, B., Lattin, J.,

Markovska, M., Mozarsky, B., Arias-Umana, J., et al. (2024). FOXO1 is a master regulator

of memory programming in CAR T cells. Nature *629*, 211–218.

https://doi.org/10.1038/s41586-024-07300-8.

10.   Larson, R.C., and Maus, M.V. (2021). Recent advances and discoveries in the

mechanisms and functions of CAR T cells. Nat. Rev. Cancer *21*, 145–161.

https://doi.org/10.1038/s41568-020-00323-z.

11.   Baulu, E., Gardet, C., Chuvin, N., and Depil, S. (2023). TCR-engineered T cell therapy

in solid tumors: State of the art and perspectives. Sci. Adv. *9*, eadf3700.

https://doi.org/10.1126/sciadv.adf3700.

12.   Tousley, A.M., Rotiroti, M.C., Labanieh, L., Rysavy, L.W., Kim, W.-J., Lareau, C.,

Sotillo, E., Weber, E.W., Rietberg, S.P., Dalton, G.N., et al. (2023). Co-opting signalling

molecules enables logic-gated control of CAR T cells. Nature *615*, 507–516.

https://doi.org/10.1038/s41586-023-05778-2.

13.   Choe, J.H., Watchmaker, P.B., Simic, M.S., Gilbert, R.D., Li, A.W., Krasnow, N.A.,

Downey, K.M., Yu, W., Carrera, D.A., Celli, A., et al. (2021). SynNotch-CAR T cells

overcome challenges of specificity, heterogeneity, and persistence in treating

glioblastoma. Sci. Transl. Med. *13*, eabe7378.

https://doi.org/10.1126/scitranslmed.abe7378.

14.   Hu, Z., Yuan, J., Long, M., Jiang, J., Zhang, Y., Zhang, T., Xu, M., Fan, Y., Tanyi, J.L.,

Montone, K.T., et al. (2021). The Cancer Surfaceome Atlas integrates genomic, functional

and drug response data to identify actionable targets. Nat. Cancer *2*, 1406–1422.

https://doi.org/10.1038/s43018-021-00282-w.

15.   Shaw, T.I., Wagner, J., Tian, L., Wickman, E., Poudel, S., Wang, J., Paul, R., Koo, S.C.,

Lu, M., Sheppard, H., et al. (2024). Discovery of immunotherapy targets for pediatric

solid and brain tumors by exon-level expression. Nat. Commun. *15*, 3732.

https://doi.org/10.1038/s41467-024-47649-y.

16.   Perna, F., Berman, S.H., Soni, R.K., Mansilla-Soto, J., Eyquem, J., Hamieh, M.,

Hendrickson, R.C., Brennan, C.W., and Sadelain, M. (2017). Integrating Proteomics and

Transcriptomics for Systematic Combinatorial Chimeric Antigen Receptor Therapy of

AML. Cancer Cell *32*, 506-519.e5. https://doi.org/10.1016/j.ccell.2017.09.004.

17.   Subramanian, A., Nemat-Gorgani, N., Ellis-Caleo, T.J., Van IJzendoorn, D.G.P., Sears,

T.J., Somani, A., Luca, B.A., Zhou, M.Y., Bradic, M., Torres, I.A., et al. (2024). Sarcoma

microenvironment cell states and ecosystems are associated with prognosis and predict

response to immunotherapy. Nat. Cancer *5*, 642–658. https://doi.org/10.1038/s43018-

024-00743-y.

18.   Gottschlich, A., Thomas, M., Grünmeier, R., Lesch, S., Rohrbacher, L., Igl, V.,

Briukhovetska, D., Benmebarek, M.-R., Vick, B., Dede, S., et al. (2023). Single-cell

transcriptomic atlas-guided development of CAR-T cells for the treatment of acute

myeloid leukemia. Nat. Biotechnol. *41*, 1618–1632. https://doi.org/10.1038/s41587-023-

01684-0.

19.   Lochner, J., Menge, F., Vassos, N., Hohenberger, P., and Kasper, B. (2020). Prognosis

of Patients with Metastatic Soft Tissue Sarcoma: Advances in Recent Years. Oncol. Res.

Treat. *43*, 613–619. https://doi.org/10.1159/000509519.

20.   Judson, I., Verweij, J., Gelderblom, H., Hartmann, J.T., Schöffski, P., Blay, J.-Y., Kerst,

J.M., Sufliarsky, J., Whelan, J., Hohenberger, P., et al. (2014). Doxorubicin alone versus

intensified doxorubicin plus ifosfamide for first-line treatment of advanced or metastatic

soft-tissue sarcoma: a randomised controlled phase 3 trial. Lancet Oncol. *15*, 415–423.

https://doi.org/10.1016/S1470-2045(14)70063-4.

21.   Tap, W.D., Papai, Z., Van Tine, B.A., Attia, S., Ganjoo, K.N., Jones, R.L., Schuetze, S.,

Reed, D., Chawla, S.P., Riedel, R.F., et al. (2017). Doxorubicin plus evofosfamide versus

1869 doxorubicin alone in locally advanced, unresectable or metastatic soft-tissue sarcoma

1870 (TH CR-406/SARC021): an international, multicentre, open-label, randomised phase 3

1871 trial. Lancet Oncol. *18*, 1089–1103. https://doi.org/10.1016/S1470-2045(17)30381-9.

1872 22. Tawbi, H.A., Burgess, M., Bolejack, V., Van Tine, B.A., Schuetze, S.M., Hu, J.,

1873 D'Angelo, S., Attia, S., Riedel, R.F., Priebat, D.A., et al. (2017). Pembrolizumab in

1874 advanced soft-tissue sarcoma and bone sarcoma (SARC028): a multicentre, two-cohort,

1875 single-arm, open-label, phase 2 trial. Lancet Oncol. *18*, 1493–1501.

1876 https://doi.org/10.1016/S1470-2045(17)30624-1.

1877 23. Burgess, M.A., Bolejack, V., Schuetze, S., Van Tine, B.A., Attia, S., Riedel, R.F., Hu,

1878 J.S., Davis, L.E., Okuno, S.H., Priebat, D.A., et al. (2019). Clinical activity of

1879 pembrolizumab (P) in undifferentiated pleomorphic sarcoma (UPS) and

1880 dedifferentiated/pleomorphic liposarcoma (LPS): Final results of SARC028 expansion

1881 cohorts. J. Clin. Oncol. *37*, 11015–11015.

1882 https://doi.org/10.1200/JCO.2019.37.15_suppl.11015.

1883 24. Somaiah, N., Conley, A.P., Parra, E.R., Lin, H., Amini, B., Solis Soto, L., Salazar, R.,

1884 Barreto, C., Chen, H., Gite, S., et al. (2022). Durvalumab plus tremelimumab in

1885 advanced or metastatic soft tissue and bone sarcomas: a single-centre phase 2 trial.

1886 Lancet Oncol. *23*, 1156–1166. https://doi.org/10.1016/S1470-2045(22)00392-8.

1887 25. Keung, E.Z., Burgess, M., Salazar, R., Parra, E.R., Rodrigues-Canales, J., Bolejack, V.,

1888 Van Tine, B.A., Schuetze, S.M., Attia, S., Riedel, R.F., et al. (2020). Correlative Analyses of

1889 the SARC028 Trial Reveal an Association Between Sarcoma-Associated Immune

Infiltrate and Response to Pembrolizumab. Clin. Cancer Res. Off. J. Am. Assoc. Cancer

Res. *26*, 1258–1266. https://doi.org/10.1158/1078-0432.CCR-19-1824.

26.   Petitprez, F., De Reyniès, A., Keung, E.Z., Chen, T.W.-W., Sun, C.-M., Calderaro, J.,

Jeng, Y.-M., Hsiao, L.-P., Lacroix, L., Bougoüin, A., et al. (2020). B cells are associated

with survival and immunotherapy response in sarcoma. Nature *577*, 556–560.

https://doi.org/10.1038/s41586-019-1906-8.

27.   D'Angelo, S.P., Araujo, D.M., Abdul Razak, A.R., Agulnik, M., Attia, S., Blay, J.-Y.,

Carrasco Garcia, I., Charlson, J.A., Choy, E., Demetri, G.D., et al. (2024). Afamitresgene

autoleucel for advanced synovial sarcoma and myxoid round cell liposarcoma

(SPEARHEAD-1): an international, open-label, phase 2 trial. The Lancet *403*, 1460–1471.

https://doi.org/10.1016/S0140-6736(24)00319-2.

28.   Hong, D.S., Van Tine, B.A., Biswas, S., McAlpine, C., Johnson, M.L., Olszanski, A.J.,

Clarke, J.M., Araujo, D., Blumenschein, G.R., Kebriaei, P., et al. (2023). Autologous T cell

therapy for MAGE-A4+ solid cancers in HLA-A*02+ patients: a phase 1 trial. Nat. Med. *29*,

104–114. https://doi.org/10.1038/s41591-022-02128-z.

29.   D'Angelo, S.P., Melchiori, L., Merchant, M.S., Bernstein, D., Glod, J., Kaplan, R.,

Grupp, S., Tap, W.D., Chagin, K., Binder, G.K., et al. (2018). Antitumor Activity Associated

with Prolonged Persistence of Adoptively Transferred NY-ESO-1 c259T Cells in Synovial

Sarcoma. Cancer Discov. *8*, 944–957. https://doi.org/10.1158/2159-8290.CD-17-1417.

30.   D'Angelo, S.P., Druta, M., Van Tine, B.A., Liebner, D.A., Schuetze, S., Hasan, A.N.,

Holmes, A.P., Huff, A., Kapoor, G.S., Zajic, S., et al. (2021). Safety and efficacy of

letetresgene autoleucel (lete-cel; GSK3377794) in advanced myxoid/round cell

liposarcoma (MRCLS) following high lymphodepletion (Cohort 2): Interim analysis. J.

Clin. Oncol. *39*, 11521–11521. https://doi.org/10.1200/JCO.2021.39.15_suppl.11521.

31.   Robbins, P.F., Morgan, R.A., Feldman, S.A., Yang, J.C., Sherry, R.M., Dudley, M.E.,

Wunderlich, J.R., Nahvi, A.V., Helman, L.J., Mackall, C.L., et al. (2011). Tumor regression

in patients with metastatic synovial cell sarcoma and melanoma using genetically

engineered lymphocytes reactive with NY-ESO-1. J. Clin. Oncol. Off. J. Am. Soc. Clin.

Oncol. *29*, 917–924. https://doi.org/10.1200/JCO.2010.32.2537.

32.   Smithy, J.W., Blouin, A., Diamond, L.C., and Postow, M. (2022). Ensuring equity in the

era of HLA-restricted cancer therapeutics. J. Immunother. Cancer *10*, e005600.

https://doi.org/10.1136/jitc-2022-005600.

33.   Jerby-Arnon, L., Neftel, C., Shore, M.E., Weisman, H.R., Mathewson, N.D., McBride,

M.J., Haas, B., Izar, B., Volorio, A., Boulay, G., et al. (2021). Opposing immune and

genetic mechanisms shape oncogenic programs in synovial sarcoma. Nat. Med. *27*,

289–300. https://doi.org/10.1038/s41591-020-01212-6.

34.   Patel, A.P., Tirosh, I., Trombetta, J.J., Shalek, A.K., Gillespie, S.M., Wakimoto, H.,

Cahill, D.P., Nahed, B.V., Curry, W.T., Martuza, R.L., et al. (2014). Single-cell RNA-seq

highlights intratumoral heterogeneity in primary glioblastoma. Science *344*, 1396–1401.

https://doi.org/10.1126/science.1254257.

35.   Timothy Tickle, Itay Tirosh, Christophe Georgescu, Maxwell Brown, Brian Haas

inferCNV of the Trinity CTAT Project.. Klarman Cell Observatory, Broad Institute of MIT

and Harvard, Cambridge, MA, USA. (Bioconductor).

https://doi.org/10.18129/B9.BIOC.INFERCNV

https://doi.org/10.18129/B9.BIOC.INFERCNV.

36.   Abeshouse, A., Adebamowo, C., Adebamowo, S.N., Akbani, R., Akeredolu, T., Ally,

A., Anderson, M.L., Anur, P., Appelbaum, E.L., Armenia, J., et al. (2017). Comprehensive

and Integrated Genomic Characterization of Adult Soft Tissue Sarcomas. Cell *171*, 950-

965.e28. https://doi.org/10.1016/j.cell.2017.10.014.

37.   Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015).

limma powers differential expression analyses for RNA-sequencing and microarray

studies. Nucleic Acids Res. *43*, e47. https://doi.org/10.1093/nar/gkv007.

38.   Tiberi, S., Crowell, H.L., Samartsidis, P., Weber, L.M., and Robinson, M.D. distinct: a

novel approach to differential distribution analyses.

39.   Uhlén, M., Fagerberg, L., Hallström, B.M., Lindskog, C., Oksvold, P., Mardinoglu, A.,

Sivertsson, Å., Kampf, C., Sjöstedt, E., Asplund, A., et al. (2015). Proteomics. Tissue-

based map of the human proteome. Science *347*, 1260419.

https://doi.org/10.1126/science.1260419.

40.   Pontén, F., Gry, M., Fagerberg, L., Lundberg, E., Asplund, A., Berglund, L., Oksvold,

P., Björling, E., Hober, S., Kampf, C., et al. (2009). A global view of protein expression in

human cells, tissues, and organs. Mol. Syst. Biol. *5*, 337.

https://doi.org/10.1038/msb.2009.93.

41.   The Tabula Sapiens Consortium*, Jones, R.C., Karkanias, J., Krasnow, M.A., Pisco,

A.O., Quake, S.R., Salzman, J., Yosef, N., Bulthaup, B., Brown, P., et al. (2022). The Tabula

Sapiens: A multiple-organ, single-cell transcriptomic atlas of humans. Science *376*,

eabl4896. https://doi.org/10.1126/science.abl4896.

42.   Regev, A., Teichmann, S.A., Lander, E.S., Amit, I., Benoist, C., Birney, E.,

Bodenmiller, B., Campbell, P., Carninci, P., Clatworthy, M., et al. (2017). The Human Cell

Atlas. eLife *6*, e27041. https://doi.org/10.7554/eLife.27041.

43.   Bughda, R., Dimou, P., D'Souza, R.R., and Klampatsa, A. (2021). Fibroblast

Activation Protein (FAP)-Targeted CAR-T Cells: Launching an Attack on Tumor Stroma.

ImmunoTargets Ther. *Volume 10*, 313–323. https://doi.org/10.2147/ITT.S291767.

44.   June, C.H., and Sadelain, M. (2018). Chimeric Antigen Receptor Therapy. N. Engl. J.

Med. *379*, 64–73. https://doi.org/10.1056/NEJMra1706169.

45.   MacKay, M., Afshinnekoo, E., Rub, J., Hassan, C., Khunte, M., Baskaran, N., Owens,

B., Liu, L., Roboz, G.J., Guzman, M.L., et al. (2020). The therapeutic landscape for cells

engineered with chimeric antigen receptors. Nat. Biotechnol. *38*, 233–244.

https://doi.org/10.1038/s41587-019-0329-2.

46.   DuBois, S.G., Krailo, M.D., Glade-Bender, J., Buxton, A., Laack, N., Randall, R.L.,

Chen, H.X., Seibel, N.L., Boron, M., Terezakis, S., et al. (2023). Randomized Phase III Trial

of Ganitumab With Interval-Compressed Chemotherapy for Patients With Newly

Diagnosed Metastatic Ewing Sarcoma: A Report From the Children's Oncology Group. J.

Clin. Oncol. Off. J. Am. Soc. Clin. Oncol. *41*, 2098–2107.

https://doi.org/10.1200/JCO.22.01815.

47.   Gounder, M.M., Rosenbaum, E., Wu, N., Dickson, M.A., Sheikh, T.N., D'Angelo, S.P.,

Chi, P., Keohan, M.L., Erinjeri, J.P., Antonescu, C.R., et al. (2022). A Phase Ib/II

Randomized Study of RO4929097, a Gamma-Secretase or Notch Inhibitor with or

without Vismodegib, a Hedgehog Inhibitor, in Advanced Sarcoma. Clin. Cancer Res. *28*,

1586–1594. https://doi.org/10.1158/1078-0432.CCR-21-3874.

48.   Luke, J.J., Powderly, J.D., Merchan, J.R., Barve, M.A., Hotson, A.N., Mobasher, M.,

Kwei, L., Luciano, G., Buggy, J.J., Piccione, E., et al. (2019). Immunobiology, preliminary

safety, and efficacy of CPI-006, an anti-CD73 antibody with immune modulating activity,

in a phase 1 trial in advanced cancers. J. Clin. Oncol. *37*, 2505–2505.

https://doi.org/10.1200/JCO.2019.37.15_suppl.2505.

49.   Edwards, D., Handsley, M., and Pennington, C. (2008). The ADAM

metalloproteinases. Mol. Aspects Med. *29*, 258–289.

https://doi.org/10.1016/j.mam.2008.08.001.

50.   Li, N., Spetz, M.R., and Ho, M. (2020). The Role of Glypicans in Cancer Progression

and Therapy. J. Histochem. Cytochem. *68*, 841–862.

https://doi.org/10.1369/0022155420933709.

51.   Qin, S., Xu, L., Yi, M., Yu, S., Wu, K., and Luo, S. (2019). Novel immune checkpoint

targets: moving beyond PD-1 and CTLA-4. Mol. Cancer *18*, 155.

https://doi.org/10.1186/s12943-019-1091-2.

52.   Du, Z., and Lovly, C.M. (2018). Mechanisms of receptor tyrosine kinase activation in

cancer. Mol. Cancer *17*, 58. https://doi.org/10.1186/s12943-018-0782-4.

53.   Zhou, B., Lin, W., Long, Y., Yang, Y., Zhang, H., Wu, K., and Chu, Q. (2022). Notch

signaling pathway: architecture, disease, and therapeutics. Signal Transduct. Target.

Ther. *7*, 95. https://doi.org/10.1038/s41392-022-00934-y.

54.   Dorsam, R.T., and Gutkind, J.S. (2007). G-protein-coupled receptors and cancer.

Nat. Rev. Cancer *7*, 79–94. https://doi.org/10.1038/nrc2069.

55.   Neufeld, G., Mumblat, Y., Smolkin, T., Toledano, S., Nir-Zvi, I., Ziv, K., and Kessler, O.

(2016). The role of the semaphorins in cancer. Cell Adhes. Migr. *10*, 652–674.

https://doi.org/10.1080/19336918.2016.1197478.

56.   Yan, H., Kamiya, T., Suabjakyong, P., and Tsuji, N.M. (2015). Targeting C-Type Lectin

Receptors for Cancer Immunity. Front. Immunol. *6*.

https://doi.org/10.3389/fimmu.2015.00408.

57.   Pacini, C., Dempster, J.M., Boyle, I., Gonçalves, E., Najgebauer, H., Karakoc, E., Van

Der Meer, D., Barthorpe, A., Lightfoot, H., Jaaks, P., et al. (2021). Integrated cross-study

datasets of genetic dependencies in cancer. Nat. Commun. *12*, 1661.

https://doi.org/10.1038/s41467-021-21898-7.

58.   Nacev, B.A., Sanchez-Vega, F., Smith, S.A., Antonescu, C.R., Rosenbaum, E., Shi, H.,

Tang, C., Socci, N.D., Rana, S., Gularte-Mérida, R., et al. (2022). Clinical sequencing of

soft tissue and bone sarcomas delineates diverse genomic landscapes and potential

therapeutic targets. Nat. Commun. *13*, 3405. https://doi.org/10.1038/s41467-022-

30453-x.

59.   Woo, H.G., Park, E.S., Lee, J.-S., Lee, Y.-H., Ishikawa, T., Kim, Y.J., and Thorgeirsson,

S.S. (2009). Identification of Potential Driver Genes in Human Liver Carcinoma by

Genomewide Screening. Cancer Res. *69*, 4059–4066. https://doi.org/10.1158/0008-

5472.CAN-09-0164.

60.　　Pollack, J.R., Sørlie, T., Perou, C.M., Rees, C.A., Jeffrey, S.S., Lonning, P.E., Tibshirani,

R., Botstein, D., Børresen-Dale, A.-L., and Brown, P.O. (2002). Microarray analysis reveals

a major direct role of DNA copy number alteration in the transcriptional program of

human breast tumors. Proc. Natl. Acad. Sci. U. S. A. *99*, 12963–12968.

https://doi.org/10.1073/pnas.162471999.

61.　　Wang, J., Zheng, J., Lee, E.E., Aguilar, B., Phan, J., Abdilleh, K., Taylor, R.C.,

Longabaugh, W., Johansson, B., Mertens, F., et al. (2023). A cloud-based resource for

genome coordinate-based exploration and large-scale analysis of chromosome

aberrations and gene fusions in cancer. Genes. Chromosomes Cancer *62*, 441–448.

https://doi.org/10.1002/gcc.23128.

62.　　Kloss, C.C., Condomines, M., Cartellieri, M., Bachmann, M., and Sadelain, M.

(2013). Combinatorial antigen recognition with balanced signaling promotes selective

tumor eradication by engineered T cells. Nat. Biotechnol. *31*, 71–75.

https://doi.org/10.1038/nbt.2459.

63.　　Roybal, K.T., Williams, J.Z., Morsut, L., Rupp, L.J., Kolinko, I., Choe, J.H., Walker, W.J.,

McNally, K.A., and Lim, W.A. (2016). Engineering T Cells with Customized Therapeutic

Response Programs Using Synthetic Notch Receptors. Cell *167*, 419-432.e16.

https://doi.org/10.1016/j.cell.2016.09.011.

64.　　Fedorov, V.D., Themeli, M., and Sadelain, M. (2013). PD-1- and CTLA-4-based

inhibitory chimeric antigen receptors (iCARs) divert off-target immunotherapy

responses. Sci. Transl. Med. *5*, 215ra172.

https://doi.org/10.1126/scitranslmed.3006597.

65.   Su, C., Xu, Z., Shan, X., Cai, B., Zhao, H., and Zhang, J. (2023). Cell-type-specific co-

expression inference from single cell RNA-sequencing data. Nat. Commun. *14*, 4846.

https://doi.org/10.1038/s41467-023-40503-7.

66.   GTEx Consortium, Laboratory, Data Analysis &Coordinating Center (LDACC)—

Analysis Working Group, Statistical Methods groups—Analysis Working Group,

Enhancing GTEx (eGTEx) groups, NIH Common Fund, NIH/NCI, NIH/NHGRI, NIH/NIMH,

NIH/NIDA, Biospecimen Collection Source Site—NDRI, et al. (2017). Genetic effects on

gene expression across human tissues. Nature *550*, 204–213.

https://doi.org/10.1038/nature24277.

67.   Kim, M.-S., Pinto, S.M., Getnet, D., Nirujogi, R.S., Manda, S.S., Chaerkady, R.,

Madugundu, A.K., Kelkar, D.S., Isserlin, R., Jain, S., et al. (2014). A draft map of the

human proteome. Nature *509*, 575–581. https://doi.org/10.1038/nature13302.

68.   Xiao, W., Mohseny, A.B., Hogendoorn, P.C.W., and Cleton-Jansen, A.-M. (2013).

Mesenchymal stem cell transformation and sarcoma genesis. Clin. Sarcoma Res. *3*, 10.

https://doi.org/10.1186/2045-3329-3-10.

69.   Ferretti, E., and Hadjantonakis, A.-K. (2019). Mesoderm specification and

diversification: from single cells to emergent tissues. Curr. Opin. Cell Biol. *61*, 110–116.

https://doi.org/10.1016/j.ceb.2019.07.012.

70.   Shekhani, M.T., Jayanthy, A.-S., Maddodi, N., and Setaluri, V. (2013). Cancer stem

cells and tumor transdifferentiation: implications for novel therapeutic strategies. Am. J.

Stem Cells *2*, 52–61.

71. Tsherniak, A., Vazquez, F., Montgomery, P.G., Weir, B.A., Kryukov, G., Cowley, G.S.,

Gill, S., Harrington, W.F., Pantel, S., Krill-Burger, J.M., et al. (2017). Defining a Cancer

Dependency Map. Cell *170*, 564-576.e16. https://doi.org/10.1016/j.cell.2017.06.010.

72. Yarmarkovich, M., Marshall, Q.F., Warrington, J.M., Premaratne, R., Farrel, A., Groff,

D., Li, W., Di Marco, M., Runbeck, E., Truong, H., et al. (2023). Targeting of intracellular

oncoproteins with peptide-centric CARs. Nature *623*, 820–827.

https://doi.org/10.1038/s41586-023-06706-0.

73. Gjerstorff, M.F., Andersen, M.H., and Ditzel, H.J. (2015). Oncogenic cancer/testis

antigens: prime candidates for immunotherapy. Oncotarget 6, 15772–15787.

https://doi.org/10.18632/oncotarget.4694.

74. Sarkizova, S., Klaeger, S., Le, P.M., Li, L.W., Oliveira, G., Keshishian, H., Hartigan,

C.R., Zhang, W., Braun, D.A., Ligon, K.L., et al. (2020). A large peptidome dataset

improves HLA class I epitope prediction across most of the human population. Nat.

Biotechnol. *38*, 199–209. https://doi.org/10.1038/s41587-019-0322-9.

75. Li, G., Iyer, B., Prasath, V.B.S., Ni, Y., and Salomonis, N. (2021). DeepImmuno: deep

learning-empowered prediction and generation of immunogenic peptides for T-cell

immunity. Brief. Bioinform. *22*, bbab160. https://doi.org/10.1093/bib/bbab160.

76. Nielsen, M., Lundegaard, C., Blicher, T., Lamberth, K., Harndahl, M., Justesen, S.,

Røder, G., Peters, B., Sette, A., Lund, O., et al. (2007). NetMHCpan, a method for

quantitative predictions of peptide binding to any HLA-A and -B locus protein of known

sequence. PloS One *2*, e796. https://doi.org/10.1371/journal.pone.0000796.

77.   Gonzalez-Galarza, F.F., McCabe, A., Santos, E.J.M.D., Jones, J., Takeshita, L., Ortega-

Rivera, N.D., Cid-Pavon, G.M.D., Ramsbottom, K., Ghattaoraya, G., Alfirevic, A., et al.

(2019). Allele frequency net database (AFND) 2020 update: gold-standard data

classification, open access genotype data and new query tools. Nucleic Acids Res.,

gkz1029. https://doi.org/10.1093/nar/gkz1029.

78.   Sewell, A.K. (2012). Why must T cells be cross-reactive? Nat. Rev. Immunol. *12*,

669–677. https://doi.org/10.1038/nri3279.

79.   Kawashima, S., Pokarowski, P., Pokarowska, M., Kolinski, A., Katayama, T., and

Kanehisa, M. (2007). AAindex: amino acid index database, progress report 2008. Nucleic

Acids Res. *36*, D202–D205. https://doi.org/10.1093/nar/gkm998.

80.   Kim, Y., Sidney, J., Pinilla, C., Sette, A., and Peters, B. (2009). Derivation of an amino

acid similarity matrix for peptide:MHC binding and its application as a Bayesian prior.

BMC Bioinformatics *10*, 394. https://doi.org/10.1186/1471-2105-10-394.

81.   Antunes, D.A., Rigo, M.M., Freitas, M.V., Mendes, M.F.A., Sinigaglia, M., Lizée, G.,

Kavraki, L.E., Selin, L.K., Cornberg, M., and Vieira, G.F. (2017). Interpreting T-Cell Cross-

reactivity through Structure: Implications for TCR-Based Cancer Immunotherapy. Front.

Immunol. *8*, 1210. https://doi.org/10.3389/fimmu.2017.01210.

82.   Bentzen, A.K., Such, L., Jensen, K.K., Marquard, A.M., Jessen, L.E., Miller, N.J.,

Church, C.D., Lyngaa, R., Koelle, D.M., Becker, J.C., et al. (2018). T cell receptor

fingerprinting enables in-depth characterization of the interactions governing recognition

of peptide-MHC complexes. Nat. Biotechnol. https://doi.org/10.1038/nbt.4303.

83. Birnbaum, M.E., Mendoza, J.L., Sethi, D.K., Dong, S., Glanville, J., Dobbins, J.,

Ozkan, E., Davis, M.M., Wucherpfennig, K.W., and Garcia, K.C. (2014). Deconstructing

the peptide-MHC specificity of T cell recognition. Cell *157,* 1073–1087.

https://doi.org/10.1016/j.cell.2014.03.047.

84. Singh, N.K., Riley, T.P., Baker, S.C.B., Borrman, T., Weng, Z., and Baker, B.M. (2017).

Emerging Concepts in TCR Specificity: Rationalizing and (Maybe) Predicting Outcomes.

J. Immunol. *199*, 2203–2213. https://doi.org/10.4049/jimmunol.1700744.

85. Tagliamonte, M., Cavalluzzo, B., Mauriello, A., Ragone, C., Buonaguro, F.M.,

Tornesello, M.L., and Buonaguro, L. (2023). Molecular mimicry and cancer vaccine

development. Mol. Cancer *22*, 75. https://doi.org/10.1186/s12943-023-01776-0.

86. Linette, G.P., Stadtmauer, E.A., Maus, M.V., Rapoport, A.P., Levine, B.L., Emery, L.,

Litzky, L., Bagg, A., Carreno, B.M., Cimino, P.J., et al. (2013). Cardiovascular toxicity and

titin cross-reactivity of affinity-enhanced T cells in myeloma and melanoma. Blood *122*,

863–871. https://doi.org/10.1182/blood-2013-03-490565.

87. Morgan, R.A., Chinnasamy, N., Abate-Daga, D., Gros, A., Robbins, P.F., Zheng, Z.,

Dudley, M.E., Feldman, S.A., Yang, J.C., Sherry, R.M., et al. (2013). Cancer regression and

neurological toxicity following anti-MAGE-A3 TCR gene therapy. J. Immunother.

Hagerstown Md 1997 *36*, 133–151. https://doi.org/10.1097/CJI.0b013e3182829903.

88. Xie, N., Shen, G., Gao, W., Huang, Z., Huang, C., and Fu, L. (2023). Neoantigens:

promising targets for cancer therapy. Signal Transduct. Target. Ther. *8*, 9.

https://doi.org/10.1038/s41392-022-01270-x.

89. Jiang, W., and Boder, E.T. (2010). High-throughput engineering and analysis of

peptide binding to class II MHC. Proc. Natl. Acad. Sci. U. S. A. *107*, 13258–13263.

https://doi.org/10.1073/pnas.1006344107.

90. Liu, R., Jiang, W., and Mellins, E.D. (2021). Yeast display of MHC-II enables rapid

identification of peptide ligands from protein antigens (RIPPA). Cell. Mol. Immunol. *18*,

1847–1860. https://doi.org/10.1038/s41423-021-00717-5.

91. Tang, S., Wang, Y., Luo, R., Fang, R., Liu, Y., Xiang, H., Ran, P., Tong, Y., Sun, M., Tan,

S., et al. (2024). Proteomic characterization identifies clinically relevant subgroups of

soft tissue sarcoma. Nat. Commun. *15*, 1381. https://doi.org/10.1038/s41467-024-

45306-y.

92. Burns, J., Wilding, C.P., Krasny, L., Zhu, X., Chadha, M., Tam, Y.B., Ps, H.,

Mahalingam, A.H., Lee, A.T.J., Arthur, A., et al. (2023). The proteomic landscape of soft

tissue sarcomas. Nat. Commun. *14*, 3834. https://doi.org/10.1038/s41467-023-39486-2.

93. Barretina, J., Caponigro, G., Stransky, N., Venkatesan, K., Margolin, A.A., Kim, S.,

Wilson, C.J., Lehár, J., Kryukov, G.V., Sonkin, D., et al. (2012). The Cancer Cell Line

Encyclopedia enables predictive modelling of anticancer drug sensitivity. Nature *483*,

603–607. https://doi.org/10.1038/nature11003.

94. Gonçalves, E., Poulos, R.C., Cai, Z., Barthorpe, S., Manda, S.S., Lucas, N., Beck, A.,

Bucio-Noble, D., Dausmann, M., Hall, C., et al. (2022). Pan-cancer proteomic map of

949 human cell lines. Cancer Cell *40*, 835-849.e8.

https://doi.org/10.1016/j.ccell.2022.06.010.

95.   Majzner, R.G., Theruvath, J.L., Nellan, A., Heitzeneder, S., Cui, Y., Mount, C.W.,

Rietberg, S.P., Linde, M.H., Xu, P., Rota, C., et al. (2019). CAR T Cells Targeting B7-H3, a

Pan-Cancer Antigen, Demonstrate Potent Preclinical Activity Against Pediatric Solid

Tumors and Brain Tumors. Clin. Cancer Res. *25*, 2560–2574.

https://doi.org/10.1158/1078-0432.CCR-18-0432.

96.   Yoshida, K., Poon, V., Dash, A., Kunder, R., Chinn, L., and Kågedal, M. (2024).

Simulation-based evaluation of personalized dosing approaches for ANTI-FGFR / KLB

bispecific antibody fazpilodemab. CPT Pharmacomet. Syst. Pharmacol. *13*, 544–550.

https://doi.org/10.1002/psp4.13111.

97.   Vahdat, L.T., Schmid, P., Forero-Torres, A., Blackwell, K., Telli, M.L., Melisko, M.,

Möbus, V., Cortes, J., Montero, A.J., Ma, C., et al. (2021). Glembatumumab vedotin for

patients with metastatic, gpNMB overexpressing, triple-negative breast cancer

("METRIC"): a randomized multicenter study. Npj Breast Cancer *7*, 57.

https://doi.org/10.1038/s41523-021-00244-6.

98.   Meininger, V., Genge, A., Van Den Berg, L.H., Robberecht, W., Ludolph, A., Chio, A.,

Kim, S.H., Leigh, P.N., Kiernan, M.C., Shefner, J.M., et al. (2017). Safety and efficacy of

ozanezumab in patients with amyotrophic lateral sclerosis: a randomised, double-blind,

placebo-controlled, phase 2 trial. Lancet Neurol. *16*, 208–216.

https://doi.org/10.1016/S1474-4422(16)30399-4.

99.   Wang, X., Qi, L., Chen, M., Zhang, Y., Gao, X., and Cai, Y. (2024). Feasibility study of

ADCs targeting TROP-2, HER2, and CD46 in Ductal Adenocarcinoma and Intraductal

2167 Carcinoma of the prostate. World J. Urol. *42*, 404. https://doi.org/10.1007/s00345-024-

2168 05109-8.

2169 100. Rafiq, S., Hackett, C.S., and Brentjens, R.J. (2020). Engineering strategies to

2170 overcome the current roadblocks in CAR T cell therapy. Nat. Rev. Clin. Oncol. *17*, 147–

2171 167. https://doi.org/10.1038/s41571-019-0297-y.

2172 101. Rodriguez-Garcia, A., Palazon, A., Noguera-Ortega, E., Powell, D.J., and Guedan, S.

2173 (2020). CAR-T Cells Hit the Tumor Microenvironment: Strategies to Overcome Tumor

2174 Escape. Front. Immunol. *11*, 1109. https://doi.org/10.3389/fimmu.2020.01109.

2175 102. Parker, K.R., Migliorini, D., Perkey, E., Yost, K.E., Bhaduri, A., Bagga, P., Haris, M.,

2176 Wilson, N.E., Liu, F., Gabunia, K., et al. (2020). Single-Cell Analyses Identify Brain Mural

2177 Cells Expressing CD19 as Potential Off-Tumor Targets for CAR-T Immunotherapies. Cell

2178 *183*, 126-142.e17. https://doi.org/10.1016/j.cell.2020.08.022.

2179 103. Jing, Y., Liu, Y., Li, Q., Ye, Y., Diao, L., Huang, Y., Zhou, Y., Green, M.R., Mills, G.B., and

2180 Han, L. (2021). Expression of chimeric antigen receptor therapy targets detected by

2181 single-cell sequencing of normal cells may contribute to off-tumor toxicity. Cancer Cell

2182 *39*, 1558–1559. https://doi.org/10.1016/j.ccell.2021.09.016.

2183 104. Lee, A.T.J., Thway, K., Huang, P.H., and Jones, R.L. (2018). Clinical and Molecular

2184 Spectrum of Liposarcoma. J. Clin. Oncol. *36*, 151–159.

2185 https://doi.org/10.1200/JCO.2017.74.9598.

2186 105. Cassinelli, G., Zaffaroni, N., and Lanzi, C. (2016). The heparanase/heparan sulfate

2187 proteoglycan axis: A potential new therapeutic target in sarcomas. Cancer Lett. *382*,

2188 245–254. https://doi.org/10.1016/j.canlet.2016.09.004.

106.   Wu, Z., Zhang, X., An, Y., Ma, K., Xue, R., Ye, G., Du, J., Chen, Z., Zhu, Z., Shi, G., et al.

(2023). CLMP is a tumor suppressor that determines all-trans retinoic acid response in

colorectal cancer. Dev. Cell *58*, 2684-2699.e6.

https://doi.org/10.1016/j.devcel.2023.10.006.

107.   Limoni, G., and Niquille, M. (2021). Semaphorins and Plexins in central nervous

system patterning: the key to it all? Curr. Opin. Neurobiol. *66*, 224–232.

https://doi.org/10.1016/j.conb.2020.12.014.

108.   Worzfeld, T., and Offermanns, S. (2014). Semaphorins and plexins as therapeutic

targets. Nat. Rev. Drug Discov. *13*, 603–621. https://doi.org/10.1038/nrd4337.

109.   Gounder, M., Ratan, R., Alcindor, T., Schöffski, P., van der Graaf, W.T., Wilky, B.A.,

Riedel, R.F., Lim, A., Smith, L.M., Moody, S., et al. (2023). Nirogacestat, a γ-Secretase

Inhibitor for Desmoid Tumors. N. Engl. J. Med. *388*, 898–912.

https://doi.org/10.1056/NEJMoa2210140.

110.   Mir, O., Azaro, A., Merchan, J., Chugh, R., Trent, J., Rodon, J., Ohnmacht, U., Diener,

J.T., Smith, C., Yuen, E., et al. (2018). Notch pathway inhibition with LY3039478 in soft

tissue sarcoma and gastrointestinal stromal tumours. Eur. J. Cancer Oxf. Engl. 1990 *103*,

88–97. https://doi.org/10.1016/j.ejca.2018.08.012.

111.   Hadjimichael, A.C., Pergaris, A., Kaspiris, A., Foukas, A.F., Kokkali, S., Tsourouflis,

G., and Theocharis, S. (2022). The EPH/Ephrin System in Bone and Soft Tissue Sarcomas'

Pathogenesis and Therapy: New Advancements and a Literature Review. Int. J. Mol. Sci.

23, 5171. https://doi.org/10.3390/ijms23095171.

112.   Kania, A., and Klein, R. (2016). Mechanisms of ephrin–Eph signalling in

development, physiology and disease. Nat. Rev. Mol. Cell Biol. *17*, 240–256.

https://doi.org/10.1038/nrm.2015.16.

113.   Martinez-Font, E., Pérez-Capó, M., Vögler, O., Martín-Broto, J., Alemany, R., and

Obrador-Hevia, A. (2021). WNT/β-Catenin Pathway in Soft Tissue Sarcomas: New

Therapeutic Opportunities? Cancers *13*, 5521.

https://doi.org/10.3390/cancers13215521.

114.   Rodriguez-Garcia, A., Lynn, R.C., Poussin, M., Eiva, M.A., Shaw, L.C., O'Connor, R.S.,

Minutolo, N.G., Casado-Medrano, V., Lopez, G., Matsuyama, T., et al. (2021). CAR-T cell-

mediated depletion of immunosuppressive tumor-associated macrophages promotes

endogenous antitumor immunity and augments adoptive immunotherapy. Nat.

Commun. *12*, 877. https://doi.org/10.1038/s41467-021-20893-2.

115.   Ash, S.L., Orha, R., Mole, H., Dinesh-Kumar, M., Lee, S.P., Turrell, F.K., and Isacke,

C.M. (2024). Targeting the activated microenvironment with endosialin (CD248)-directed

CAR-T cells ablates perivascular cells to impair tumor growth and metastasis. J.

Immunother. Cancer *12*, e008608. https://doi.org/10.1136/jitc-2023-008608.

116.   Wei, R., Dean, D.C., Thanindratarn, P., Hornicek, F.J., Guo, W., and Duan, Z. (2020).

Cancer testis antigens in sarcoma: Expression, function and immunotherapeutic

application. Cancer Lett. *479*, 54–60. https://doi.org/10.1016/j.canlet.2019.10.024.

117.   Fonseca, A.F., and Antunes, D.A. (2023). CrossDome: an interactive R package to

predict cross-reactivity risk using immunopeptidomics databases. Front. Immunol. *14*,

1142573. https://doi.org/10.3389/fimmu.2023.1142573.

118. Squair, J.W., Gautier, M., Kathe, C., Anderson, M.A., James, N.D., Hutson, T.H.,

Hudelle, R., Qaiser, T., Matson, K.J.E., Barraud, Q., et al. (2021). Confronting false

discoveries in single-cell differential expression. Nat. Commun. *12*, 5692.

https://doi.org/10.1038/s41467-021-25960-2.

119. R Core Team (2023). R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna, Austria.

120. Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., Zheng, S., Butler, A., Lee, M.J.,

Wilk, A.J., Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-cell

data. Cell *184*, 3573-3587.e29. https://doi.org/10.1016/j.cell.2021.04.048.

121. McGinnis, C.S., Murrow, L.M., and Gartner, Z.J. (2019). DoubletFinder: Doublet

Detection in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. Cell

Syst. *8*, 329-337.e4. https://doi.org/10.1016/j.cels.2019.03.003.

122. Korsunsky, I., Millard, N., Fan, J., Slowikowski, K., Zhang, F., Wei, K., Baglaenko, Y.,

Brenner, M., Loh, P., and Raychaudhuri, S. (2019). Fast, sensitive and accurate

integration of single-cell data with Harmony. Nat. Methods *16*, 1289–1296.

https://doi.org/10.1038/s41592-019-0619-0.

123. Samuel Marsh, Maëlle Salmon, and Paul Hoffman (2024). samuel-

marsh/scCustomize: Version 2.1.2. Version v2.1.2 (Zenodo).

https://doi.org/10.5281/ZENODO.5706430 https://doi.org/10.5281/ZENODO.5706430.

124. Alquicira-Hernandez, J., and Powell, J.E. (2021). Nebulosa recovers single-cell gene

expression signals by kernel density estimation. Bioinforma. Oxf. Engl. *37*, 2485–2487.

https://doi.org/10.1093/bioinformatics/btab003.

125.   Aran, D., Looney, A.P., Liu, L., Wu, E., Fong, V., Hsu, A., Chak, S., Naikawadi, R.P.,

Wolters, P.J., Abate, A.R., et al. (2019). Reference-based analysis of lung single-cell

sequencing reveals a transitional profibrotic macrophage. Nat. Immunol. *20*, 163–172.

https://doi.org/10.1038/s41590-018-0276-y.

126.   The ENCODE Project Consortium (2012). An integrated encyclopedia of DNA

elements in the human genome. Nature *489*, 57–74.

https://doi.org/10.1038/nature11247.

127.   Stunnenberg, H.G., International Human Epigenome Consortium, and Hirst, M.

(2016). The International Human Epigenome Consortium: A Blueprint for Scientific

Collaboration and Discovery. Cell *167*, 1145–1149.

https://doi.org/10.1016/j.cell.2016.11.007.

128.   Baron, M., Veres, A., Wolock, S.L., Faust, A.L., Gaujoux, R., Vetere, A., Ryu, J.H.,

Wagner, B.K., Shen-Orr, S.S., Klein, A.M., et al. (2016). A Single-Cell Transcriptomic Map

of the Human and Mouse Pancreas Reveals Inter- and Intra-cell Population Structure.

Cell Syst. *3*, 346-360.e4. https://doi.org/10.1016/j.cels.2016.08.011.

129.   Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR : a Bioconductor

package for differential expression analysis of digital gene expression data.

Bioinformatics *26*, 139–140. https://doi.org/10.1093/bioinformatics/btp616.

130.   Amezquita, R.A., Lun, A.T.L., Becht, E., Carey, V.J., Carpp, L.N., Geistlinger, L.,

Marini, F., Rue-Albrecht, K., Risso, D., Soneson, C., et al. (2020). Orchestrating single-

cell analysis with Bioconductor. Nat. Methods *17*, 137–145.

https://doi.org/10.1038/s41592-019-0654-x.

131.   McCarthy, D.J., Campbell, K.R., Lun, A.T.L., and Wills, Q.F. (2017). Scater: pre-

processing, quality control, normalization and visualization of single-cell RNA-seq data

in R. Bioinformatics *33*, 1179–1186. https://doi.org/10.1093/bioinformatics/btw777.

132.   Bausch-Fluck, D., Goldmann, U., Müller, S., Van Oostrum, M., Müller, M., Schubert,

O.T., and Wollscheid, B. (2018). The in silico human surfaceome. Proc. Natl. Acad. Sci.

115. https://doi.org/10.1073/pnas.1808790115.

133.   Jiang, L., Wang, M., Lin, S., Jian, R., Li, X., Chan, J., Dong, G., Fang, H., Robinson,

A.E., Snyder, M.P., et al. (2020). A Quantitative Proteome Map of the Human Body. Cell

183, 269-283.e19. https://doi.org/10.1016/j.cell.2020.08.036.

134.   Adam Mark, R.T. (2017). mygene. (Bioconductor).

https://doi.org/10.18129/B9.BIOC.MYGENE https://doi.org/10.18129/B9.BIOC.MYGENE.

135.   Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L.,

et al. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.

Innov. Camb. Mass *2*, 100141. https://doi.org/10.1016/j.xinn.2021.100141.

136.   Yu, G., and He, Q.-Y. (2016). ReactomePA: an R/Bioconductor package for reactome

pathway analysis and visualization. Mol. Biosyst. *12*, 477–479.

https://doi.org/10.1039/C5MB00663E.

137.   Yu, G., Wang, L.-G., Han, Y., and He, Q.-Y. (2012). clusterProfiler: an R Package for

Comparing Biological Themes Among Gene Clusters. OMICS J. Integr. Biol. *16*, 284–287.

https://doi.org/10.1089/omi.2011.0118.

138.   Carlson M (2019). org.Hs.eg.db: Genome wide annotation for Human. R package

version 3.8.2.

139.   The Cancer Genome Atlas Research Network, Weinstein, J.N., Collisson, E.A., Mills,

G.B., Shaw, K.R.M., Ozenberger, B.A., Ellrott, K., Shmulevich, I., Sander, C., and Stuart,

J.M. (2013). The Cancer Genome Atlas Pan-Cancer analysis project. Nat. Genet. *45*,

1113–1120. https://doi.org/10.1038/ng.2764.

140.   Cerami, E., Gao, J., Dogrusoz, U., Gross, B.E., Sumer, S.O., Aksoy, B.A., Jacobsen, A.,

Byrne, C.J., Heuer, M.L., Larsson, E., et al. (2012). The cBio cancer genomics portal: an

open platform for exploring multidimensional cancer genomics data. Cancer Discov. *2*,

401–404. https://doi.org/10.1158/2159-8290.CD-12-0095.

141.   Gao, J., Aksoy, B.A., Dogrusoz, U., Dresdner, G., Gross, B., Sumer, S.O., Sun, Y.,

Jacobsen, A., Sinha, R., Larsson, E., et al. (2013). Integrative analysis of complex cancer

genomics and clinical profiles using the cBioPortal. Sci. Signal. *6*, pl1.

https://doi.org/10.1126/scisignal.2004088.

142.   Gao, Q., Liang, W.-W., Foltz, S.M., Mutharasu, G., Jayasinghe, R.G., Cao, S., Liao, W.-

W., Reynolds, S.M., Wyczalkowski, M.A., Yao, L., et al. (2018). Driver Fusions and Their

Implications in the Development and Treatment of Human Cancers. Cell Rep. *23*, 227-

238.e3. https://doi.org/10.1016/j.celrep.2018.03.050.

143.   Mertens F, Mitelman Database of Chromosome Aberrations and Gene Fusions in

Cancer (2024). Mitelman F, Johansson B and Mertens F (Eds.),

https://mitelmandatabase.isb-cgc.org.

144.   Karlsson, M., Zhang, C., Méar, L., Zhong, W., Digre, A., Katona, B., Sjöstedt, E.,

Butler, L., Odeberg, J., Dusart, P., et al. (2021). A single–cell type transcriptomics map of

human tissues. Sci. Adv. *7*, eabh2169. https://doi.org/10.1126/sciadv.abh2169.

145. Wells, D.A., and McAuley, M. (2023). HLAfreq: Download and combine HLA allele

frequency data. Preprint, https://doi.org/10.1101/2023.09.15.557761

https://doi.org/10.1101/2023.09.15.557761.

146. The UniProt Consortium, Bateman, A., Martin, M.-J., Orchard, S., Magrane, M.,

Ahmad, S., Alpi, E., Bowler-Barnett, E.H., Britto, R., Bye-A-Jee, H., et al. (2023). UniProt:

the Universal Protein Knowledgebase in 2023. Nucleic Acids Res. *51*, D523–D531.

https://doi.org/10.1093/nar/gkac1052.

147. Vita, R., Mahajan, S., Overton, J.A., Dhanda, S.K., Martini, S., Cantrell, J.R., Wheeler,

D.K., Sette, A., and Peters, B. (2019). The Immune Epitope Database (IEDB): 2018

update. Nucleic Acids Res. *47*, D339–D343. https://doi.org/10.1093/nar/gky1006.

148. Huang, X., Gan, Z., Cui, H., Lan, T., Liu, Y., Caron, E., and Shao, W. (2024). The

SysteMHC Atlas v2.0, an updated resource for mass spectrometry-based

immunopeptidomics. Nucleic Acids Res. *52*, D1062–D1071.

https://doi.org/10.1093/nar/gkad1068.

149. Marcu, A., Bichmann, L., Kuchenbecker, L., Kowalewski, D.J., Freudenmann, L.K.,

Backert, L., Mühlenbruch, L., Szolek, A., Lübke, M., Wagner, P., et al. (2021). HLA Ligand

Atlas: a benign reference of HLA-presented peptides to improve T-cell-based cancer

immunotherapy. J. Immunother. Cancer 9, e002071. https://doi.org/10.1136/jitc-2020-

002071.

150. O'Donnell, T.J., Rubinsteyn, A., and Laserson, U. (2020). MHCflurry 2.0: Improved

Pan-Allele Prediction of MHC Class I-Presented Peptides by Incorporating Antigen

Processing. Cell Syst. *11*, 42-48.e7. https://doi.org/10.1016/j.cels.2020.06.010.

151.   McFerrin L (2013). HDMD: Statistical Analysis Tools for High Dimension Molecular

Data (HDMD).

152.   Amerifar S (2021). ftrCOOL: Feature Extraction from Biological Sequences. Version

version 2.0.0.

153.   Liao, Y., Savage, S.R., Dou, Y., Shi, Z., Yi, X., Jiang, W., Lei, J.T., and Zhang, B. (2023). A

proteogenomics data-driven knowledge base of human cancer. Cell Syst. *14*, 777-

787.e5. https://doi.org/10.1016/j.cels.2023.07.007.

154.   Thangudu, R.R., Rudnick, P.A., Holck, M., Singhal, D., MacCoss, M.J., Edwards, N.J.,

Ketchum, K.A., Kinsinger, C.R., Kim, E., and Basu, A. (2020). Abstract LB-242: Proteomic

Data Commons: A resource for proteogenomic analysis. Cancer Res. *80*, LB-242-LB-

242. https://doi.org/10.1158/1538-7445.AM2020-LB-242.

155.   Sinha, A., Huang, V., Livingstone, J., Wang, J., Fox, N.S., Kurganovs, N., Ignatchenko,

V., Fritsch, K., Donmez, N., Heisler, L.E., et al. (2019). The Proteogenomic Landscape of

Curable Prostate Cancer. Cancer Cell *35*, 414-427.e6.

https://doi.org/10.1016/j.ccell.2019.02.005.

156.   Liu, W., Xie, L., He, Y.-H., Wu, Z.-Y., Liu, L.-X., Bai, X.-F., Deng, D.-X., Xu, X.-E., Liao,

L.-D., Lin, W., et al. (2021). Large-scale and high-resolution mass spectrometry-based

proteomics profiling defines molecular subtypes of esophageal cancer for therapeutic

targeting. Nat. Commun. *12*, 4961. https://doi.org/10.1038/s41467-021-25202-5.

157.   Beck, L., Harel, M., Yu, S., Markovits, E., Boursi, B., Markel, G., and Geiger, T. (2021).

Clinical Proteomics of Metastatic Melanoma Reveals Profiles of Organ Specificity and

Treatment Resistance. Clin. Cancer Res. *27*, 2074–2086. https://doi.org/10.1158/1078-

0432.CCR-20-3752.

158.   Groeneveld, C.S., Sanchez-Quiles, V., Dufour, F., Shi, M., Dingli, F., Nicolle, R.,

Chapeaublanc, E., Poullet, P., Jeffery, D., Krucker, C., et al. (2024). Proteogenomic

Characterization of Bladder Cancer Reveals Sensitivity to Apoptosis Induced by Tumor

Necrosis Factor–related Apoptosis-inducing Ligand in FGFR3-mutated Tumors. Eur. Urol.

85, 483–494. https://doi.org/10.1016/j.eururo.2023.05.037.

159.   Boder, E.T., and Wittrup, K.D. (1997). Yeast surface display for screening

combinatorial polypeptide libraries. Nat. Biotechnol. *15*, 553–557.

https://doi.org/10.1038/nbt0697-553.

160.   Raybould, M.I.J., Marks, C., Lewis, A.P., Shi, J., Bujotzek, A., Taddese, B., and Deane,

C.M. (2020). Thera-SAbDab: the Therapeutic Structural Antibody Database. Nucleic

Acids Res. *48*, D383–D388. https://doi.org/10.1093/nar/gkz827.

2376

2377

2378

2379

2380

2381

2382

2383

2384

2385 **Figures**

Figure 1. scRNA-seq for cellular therapy target identification.

(A). Flow diagram showing the full SCAN-ACT pipeline. Stacked bar charts showing the fraction of cells identified as tumoral, endothelial, myeloid, lymphoid, or CAFs, sample site of origin, prior treatment, and metastatic or primary tumors. (C, D) Heatmaps showing average RNA expression in STS versus non-tumoral cells in the TME (C) and normal cells in the HPA, TS, and HCA (D). RNA expression (log2(CPM+1)) was averaged and median-centered across cell types, and z-scored across genes. The number of cells analyzed is shown in the legend. Only genes differentially expressed in each STS and measured in all samples are shown. (E) Venn diagrams showing the number of STS up-regulated and down-regulated genes shared between histotypes. (F) Stacked bar chart showing the fraction of DEGs in STS versus non-tumoral cells in the TME, identified with Limma-voom, distinct, or both methods, considering surfaceome and total genes separately. (G) Lollipop plots showing the log2FC in expression for MMP14 and FAP between UPS cells and the normal cell clusters in the HPA, TS, and HCA using Limma-voom. Only statistically significant comparisons are shown (FDR < 0.25). (H) FAP IHC expression on tissue microarrays of UPS samples. CAFs: cancer associated fibroblasts; ECs: endothelial cells; LCs: lymphoid cells; MCs: myeloid cells; Mφ: Macrophages.

2386

2387

2388

2389

2390

2391

2392

2393

2394

Figure 2. Monospecific CAR-T cell targets in STS.

(A) Diagram showing discovery pipeline for monospecific STS CAR-T targets. (B) Heatmap showing the protein expression in normal tissues and cells or tissue groups for CAR-T targets tested in humans (left panel) and for the STS targets (right panel) in the HPM dataset. (C) Bar chart showing the proportion of STS CAR-T targets for each protein class/family. (D) Scatter plot showing gene set ORA on the 449 CAR-T targets, using the gene ontology (GO) molecular function (MF) and biological process (BP), as well as the KEGG, Reactome, and WikiPathways terms. The x-axis shows the –log10(FDR) while the dots size indicates the number of targets per gene set. The horizontal line intersects at the significance threshold (FDR = 0.05). (E) Ridge plots showing RNA expression (log2(TPM+1)) in the Pan-Cancer TCGA Atlas for STS CAR-T targets. STS samples (SARC) are highlighted in bold red. (F) Violin plots showing the effect of CAR-T targets KO on sarcoma cell line growth from DepMap, quantified using the Chronos algorithm. Negative values indicate genes essential for cell growth. The vertical line indicates the median Chronos value for known essential genes (–1). Dots are colored based on the cell line cancer of origin. (G) Pearson correlation between linear CNV level and RNA expression in TCGA SARC for monospecific CAR-T targets. P-values were adjusted using the BH method setting significance to FDR < 0.01. (H) Circos plot showing gene fusions involving a CAR-T target in STS from the Mitelman database, MSK-IMPACT, and TCGA SARC. Colored links show fusions for the top 30 most recurrently fused genes, with link colors indicating the STS histotype. (I) Oncoplots showing recurring SNVs/Indels for monospecific CAR-T targets in MSK-IMPACT (left) and TCGA SARC (right). ALT: Atypical Lipomatous Tumor; GIST: Gastrointestinal Stromal Tumor; MPNST: Malignant Peripheral Nerve Sheath Tumor; IMT: Inflammatory Myofibroblastic Tumor; MFH: Malignant Fibrous Histiocytoma; SCS: High-grade Spindle Cell Sarcoma.

2395

Figure 3. Bispecific CAR-T targets in STS: 'AND' and 'AND–NOT' logic-gated CARs.

(A) Illustration of an 'AND' CAR-T cell activated by two different tumor antigens. (B) Venn diagram showing the number of 'AND' pairs in each STS. (C) Bar chart showing the top 5 most frequent 'AND' pairs by STS. Genes are classified based on their HPA single cell type signature. (D) Illustration of an 'AND-NOT' CAR-T cell activated by a tumor antigen, and inhibited by a normal cell antigen. (E) Venn diagram showing the number of shared and unique 'AND-NOT' pairs in each STS. (F) Bar chart showing the top 5 most frequent 'AND-NOT' pairs by STS. Genes are classified based on their HPA single cell signature, with the first indicating the activator and the second the inhibitor. (G, H) Heatmaps showing selected 'AND' and 'AND-NOT' pairs expression in the HPA scRNA-seq (G), and the HPM (H) datasets. Genes are annotated based on the HPA single-cell signature they belong to (top annotation), and their functional role as activators or inhibitors (bottom annotation, 'AND-NOT' pairs).

2396

Figure 4. Targeting intracellular antigens in STS: TCR-T and PC-CAR-T cells.

(A) TCR-T and PC-CAR-T pMHC target discovery pipeline. (B) Heatmap showing bulk RNA-seq GTEx expression for the 76 target antigens. Top annotation indicates predicted antigen in each STS, with average expression in STS scRNA-seq data and HPA STS cell lines. (C) Gene set ORA using GO BP terms for PC-CAR-T and TCR-T target antigens (FDR < 0.1, minimum and maximum gene set sizes of 2 and 500). (D, E) Heatmaps showing HPA/GTEx tissue-specific S_CR scores for target pMHCs scaled to the S_CR scores of afami-cel's target, GVYDGREHTV–HLA-A*02:01, obtained as follows: $(((S\_CR^j (target)-S\_CR^j (reference)))/\sigma\_j) \times 2$ if $S\_CR^j (target)>S\_CR^j (reference)$, where σ_j is the standard deviation of the S_CR values for the 5,020 target pMHCs in the jth tissue. In (D), pMHCs with S_CR Diff ≤0, and formed by peptides that form pMHCs with ≥10 different HLA alleles in either locus are shown. In (E), only pMHCs with S_CR Diff ≤ 10th percentile and formed by one of the top 10 most prevalent HLA alleles at either locus are shown.

2397

Figure 5. Target validation.

(A) Vignette demonstrating the IMMUNE workflow for peptide-MHC binding assessment. (B) IMMUNE results for 14 pMHC targets identified through PS-TRACT, including KAFDDIATYF and RLHRIIPKI from SSXI, AVLDGLDVLL and VLYPVPLESY from PRAME, NVFPPPSQTW and YLQDGTDVSL from ADAM12, and SLLPFSVEAL, KLPEDALVKA and AALPVAPVGL from MSX1. The scatterplots report the mean fluorescence intensity (MFI) for the peptide-HLA binding indicator (x-axis) and for the HLA molecule tested (y-axis). Each peptide is identified by its first three amino acids. Strong binders are indicated in red, while moderate binders are shown in yellow, and weak binders in light blue. The KLT, AIF, RYP, and FAF peptides are used as positive controls for known strong binders for the HLA considered. (C) Bar chart showing the % Competition as measured with IMMUNE, for each tested pMHC target. Strong binders are indicated in red, moderate binders in yellow, and weak binders in light blue. (D) Dot plot showing protein expression in STS cell lines in ProCan-DepMapSanger. (E) Spearman correlation of scRNA-seq expression and protein abundance in the Tang et al. dataset. Only monospecific CAR-T targets for each histotype are plotted. LPS scRNA-seq samples were compared to DDLPS and WDLPS proteomics samples separately. (F) Bar charts showing the fraction of samples in each STS with high, medium, and low expression of selected monospecific CAR-T targets in the Tang et al. dataset. (G) Spearman correlation of protein expression in the Tang et al. dataset for 'AND' target pairs. (H) Protein expression by IHC in STS tissue micro-arrays. The bar chart shows the fraction of samples in each STS with negative, weakly positive, and strongly positive expression. Images show tissue sections of LMS samples scored as strongly positive for GPC6, MRC2, CLMP, and KITLG. MPNST: malignant peripheral nerve sheath tumor; MLPS: mucinous liposarcoma; others: other fibrosarcoma; ES: Ewing sarcoma; RMS: rhabdomyosarcoma; AS: angiosarcoma.

2398

Figure 6. Integrated target prioritization.

(A, C, D) Circular heatmaps showing the top 100 CAR-T targets ranked by TPI for LPS (A, monospecific CAR-T targets), and LMS (C, D, bispecific CAR-T targets). (B, E, G) UMAP plots showing expression density of selected monospecific CAR-T targets in LPS (B), bispecific CAR-T targets in LMS (E), and TCR-T/PC-CAR-T target antigens in SS (G). In one UMAP plot per panel, cells are colored based on identity. (F) Circular heatmap showing TPI scoring for TCR-T and PC-CAR-T target antigens in SS.

2399

2400