Pyruvate and Related Energetic Metabolites Modulate Resilience Against High Genetic

Risk for Glaucoma

Keva Li[1,*], Nicholas Tolman[2,*], Ayellet V. Segrè[3,4], Kelsey V. Stuart[5], Oana A. Zeleznik[6], Neeru

A. Vallabh[7,8], Kuang Hu[5], Nazlee Zebardast[3], Akiko Hanyuda[9,10], Yoshihiko Raita[11], Christa

Montgomery[2], Chi Zhang[2], Pirro G. Hysi[12,13], Ron Do[14], Anthony P. Khawaja[5], Janey L.

Wiggs[3,4], Jae H. Kang[6,‡], Simon WM John[2,15,‡], Louis R. Pasquale[1,‡] & UK Biobank Eye and

Vision Consortium

[1]Department of Ophthalmology, Icahn School of Medicine at Mount Sinai, New York, NY

10029, USA

[2]Department of Ophthalmology, Columbia University Irving Medical Center, New York, NY

10027, USA

[3]Department of Ophthalmology, Massachusetts Eye and Ear, Harvard Medical School, Boston,

MA 02142, USA

[4]Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA

[5]NIHR Biomedical Research Centre, Moorfields Eye Hospital NHS Foundation Trust, and

University College London Institute of Ophthalmology, London, EC1V 2PD, United Kingdom

[6]Channing Division of Network Medicine, Department of Medicine, Harvard Medical School

and Brigham and Women's Hospital, 181 Longwood Avenue, Boston, MA 02115, USA.

[7]Department of Eye and Vision Science, Institute of Life Course and Medical Sciences,

University of Liverpool, Liverpool, L69 3BX, United Kingdom

[8]St. Paul's Eye Unit, Liverpool University Hospital NHS Foundation Trust, Liverpool, L7 8XP,

United Kingdom

[9]Department of Ophthalmology, Keio University School of Medicine, 35 Shinanomachi,

Shinjuku-ku, Tokyo 160-8582, Japan

[10]Epidemiology and Prevention Group, Center for Public Health Sciences, National Cancer

Center, Tokyo 104-0045, Japan

[11]Okinawa Kenritsu, Chubu Byoin, Uruma, Okinawa 904-2293, Japan

[12]Department of Ophthalmology, St Thomas' Hospital, King's College London, London, SE1

7EH, United Kingdom.

[13]Department of Twin Research & Genetic Epidemiology, St Thomas' Hospital, King's College

London, London, SE1 7EH, United Kingdom.

[14]Department of Genetics and Genomics Science, Icahn School of Medicine at Mount Sinai,

New York, NY 10029, USA.

[15]Zuckerman Mind Brain Behavior Institute, Columbia University, New York, NY 10032, USA

[*] Co-first authors

[‡] Co-senior authors

Running Head: Pyruvate as a Modulator of Resilience Against High Genetic Risk for Glaucoma

Corresponding authors:

Louis R. Pasquale, MD, FARVO

Department of Ophthalmology, Icahn School of Medicine at Mount Sinai, New York, NY, USA

Louis.Pasquale@mssm.edu

Simon WM John, PhD

Department of Ophthalmology, Columbia University Irving Medical Center, New York, NY

10032, USA

Keywords: metabolite risk score; metabolites; genetic risk score; genetics; glaucoma; glaucoma

resilience; glycolysis; pyruvate; tricarboxylic acid cycle

Abbreviations and Acronyms:

AUC, area under the curve; BMI, body mass index; CI, confidence interval; HDL, high-density

lipoprotein; ICD, International Classification of Diseases; IOP, intraocular pressure; mRNFL,

macular retinal nerve fiber layer; MET, metabolic equivalent of task; MRS, metabolic risk score;

MTAG, Multi-Trait Analysis of Genome-wide association study data; NEF, number of effective;

NMR, nuclear magnetic resonance; OR, odds ratio; POAG, primary open-angle glaucoma; PRS,

polygenic risk score; RGC, retinal ganglion cell; ROC, receiver operating characteristic; SD,

standard deviation; SNP, single nucleotide polymorphism; TCA cycle, Tricarboxylic acid cycle;

VCDR, vertical cup-disc ratio; VLDL, very low-density lipoprotein.

Abstract

Glaucoma polygenic risk scores (PRS) effectively identify disease risk, but some individuals with high PRS do not develop glaucoma. Factors contributing to this resilience remain unclear. Using 4,658 glaucoma cases and 113,040 controls in a cross-sectional study in the UK Biobank, we investigated whether plasma metabolites enhanced glaucoma prediction and if a metabolomic signature of resilience in high-genetic risk individuals existed. Logistic regression models incorporating 168 NMR-based metabolites into PRS-based glaucoma assessments were developed, with multiple comparison corrections applied. While metabolites weakly predicted glaucoma (Area Under the Curve=0.579), they offered modest prediction improvement in PRS-only-based models (P=0.004). We identified a metabolomic signature associated with resilience in the top PRS decile, with elevated glycolysis-related metabolites—lactate (P=8.8E-12), pyruvate (P=1.9E-10), and citrate (P=0.02)—linked to reduced glaucoma prevalence. These metabolites combined significantly modified the PRS-glaucoma relationship ($P_{interaction}$=0.011). Higher total resilience metabolite levels within the highest PRS quartile corresponded to lower glaucoma prevalence (Odds Ratio$_{highest\ vs.\ lowest\ total\ resilience\ metabolite\ quartile}$=0.71, 95% Confidence Interval [CI]=0.64–0.80). As pyruvate is a foundational metabolite linking glycolysis to tricarboxylic acid cycle metabolism and ATP generation, we pursued experimental validation for this putative resilience biomarker in a human-relevant Mus musculus glaucoma model. Dietary pyruvate mitigated elevated intraocular pressure (P=0.002) and optic nerve damage (P<0.0003) in $Lmx1b^{V265D}$ mice. These findings highlight the protective role of pyruvate-related metabolism against glaucoma and suggest potential avenues for therapeutic intervention.

Introduction

Glaucoma is a polygenic, progressive neurodegenerative disease and a leading cause of irreversible blindness.[1] The disease is typically asymptomatic until advanced visual field loss occurs, and around 50 to 70% of people affected remain undiagnosed.[2,3] Early detection and intervention are essential to stop disease progression and prevent visual impairment in glaucoma patients, as there is no cure available. However, population-based glaucoma screening is not cost-effective from a public health perspective.[4–6]

Glaucoma is well-suited for developing and applying polygenic risk scores (PRS) to facilitate disease identification and risk stratification as it is a condition with several endophenotypes, such as elevated intraocular pressure (IOP) and thinning of the retinal nerve fiber layer (RNFL), and significant heritability.[7–9] However, glaucoma is also significantly affected by environmental and lifestyle factors,[10–14] which are not accounted for in a PRS profile. Glaucoma was five times more likely in the UK Biobank (UKBB) among participants with the highest genetic risk decile versus the lowest decile (7.4% vs 1.3%).[9] While the relatively low glaucoma prevalence in the highest decile group could be explained by disease under-ascertainment or a PRS that incompletely reflects the glaucoma genetic architecture, it is possible that resilience biomarkers could explain this result, in addition to protective genetic or epigenetic factors.

Recent advancements in metabolomics have opened up avenues to explore metabolites as potential biomarkers for glaucoma.[15] Metabolites are intermediate and end products of cellular processes critical for driving cellular growth and tissue homeostasis.[16] These small molecules provide a holistic measure of physiological status, reflecting both genetic predispositions and environmental influences. Previous studies have indicated a potential role for plasma metabolites

to stratify glaucoma risk; however, these studies were limited by small sample sizes,[17–21] with

restricted coverage of metabolites. In addition, the potential benefits of the integration of

metabolomics with genetics to identify individuals at the highest risk of glaucoma remain

unexplored.

This study aims to evaluate the relevance of plasma metabolite data as risk factors for

glaucoma. First, we explored whether incorporating plasma metabolite data can improve the

predictive accuracy of a PRS for glaucoma risk based on available data in the UKBB, where

genetic and metabolomic data were available in 117,698 participants. Additionally, we evaluated

interactions between PRS and metabolomics for further stratifying individuals with high genetic

risk but without glaucoma. We undertook an agnostic approach to identify a metabolite signature

associated with resilience to high glaucoma genetic risk. Finally, we experimentally validated the

top resilience metabolite by assessing its ability to rescue the ocular phenotype in a human-

relevant, genetic mouse model of adult-onset glaucoma.

Results

UK Biobank Study Characteristics

A total of 117,698 participants (4,658 glaucoma cases and 113,040 non-cases) were

included in this study. UK Biobank participants were predominately of European ancestry

(85.8%) but were also of admixed American (0.22%), Asian (2.77%), and African (1.73%)

ancestry. The design of the human studies is depicted graphically in **Figures 1 and 2**. There were

significant differences seen in baseline demographic and clinical characteristics between the

glaucoma cases and the non-cases presented in **Table 1**. Notably, characteristics associated with

glaucoma included male gender, older age, genetic African ancestry (see ref #22 for details

regarding the derivation of genetic ancestry in the UK Biobank)[22], prior history of smoking,

slightly lower cholesterol levels, higher body mass index (BMI), higher hemoglobin A1C

(HbA1C), higher IOP, thinner macular retinal nerve fiber layer (mRNFL) thickness, higher

caffeine intake, higher alcohol intake, higher oral steroid use, diabetes, and coronary artery

disease.

Plasma metabolites modestly improve glaucoma risk prediction.

We first sought to determine whether the addition of metabolites could improve the

prediction of glaucoma from basic demographics, clinical variables, and genetic data. We created

four logistic regression models, each with increasing predictive variables considered, to evaluate

their added utility for predicting glaucoma (see Methods and **Supplemental Table 2** for model

construction details). We analyzed two sets of metabolite variables: a comprehensive set of 168

metabolites measured by nuclear magnetic resonance (NMR) spectroscopy included in the UK

Biobank dataset and a limited set of 27 metabolites approved by the European Union (EU) for *in*

vitro diagnostic use.

To evaluate the performance of each model, we plotted receiver operating characteristic

(ROC) curves and calculated the area under the curve (AUC) metrics. In model 1, which utilized

only metabolite data, the highest performance was achieved using the full panel of 168

metabolites, yielding an AUC value of 0.602 (95% CI=0.592-0.612) compared to 0.579 (95%

CI=0.569-0.589; P=0.0003) using the 27 metabolites (**Figure 3**). Model 2 containing

demographic information and Model 3, which also included clinical variables, performed

similarly before the addition of metabolite data. In model 2, the addition of the 168 metabolite

panel demonstrated the best performance, with an AUC value of 0.670 (95% CI=0.660-0.680)

compared to 0.664 (95% CI=0.654-0.674) for the model without metabolites (P=0.002) and

0.666 (95% CI=0.656-0.676) for the limited panel of 27 metabolites. This performance trend

continued with model 3 with the inclusion of 168 metabolites, producing an AUC of 0.680 (95%

CI=0.660-0.700), which represented in an increase in AUC (P=0.02) from model 3 with the

exclusion of metabolites (AUC 0.670; 95% CI=0.650-0.690). We subsequently examined

whether integrating metabolite data into a PRS-based model could improve glaucoma prediction

algorithms. In Model 4, the panel of 168 metabolites yielded the best performance with an AUC

value of 0.806 (95% CI=0.796-0.816), while the model without metabolites (AUC=0.801, 95%

CI=0.791-0.811; P=0.004) as well as the model with a limited panel of 27 metabolites

(AUC=0.802, 95% CI=0.792-0.812; P=0.006) both had lower AUC values. Thus, these findings

suggest that the addition of metabolite data modestly enhances the prediction of glaucoma

beyond the use of demographic and genetic data collected on an aggregated basis.

We then investigated whether the inclusion of the 168 metabolites could improve

predictive value for glaucoma by stratifying patients into specific subgroups and calculating

AUC curves for each stratum. Accordingly, we found that metabolites showed a modestly

improved predictive value for glaucoma among people of White ethnicity (P<0.001), those aged

55 and older (P=0.002), and males (P=0.002) (**Table 2**).

Metabolites associated with resilience to a high glaucoma polygenic risk score

As metabolite data only modestly augmented clinical or PRS predictions of glaucoma, we

hypothesized that metabolites might provide utility for differentiating patients within risk groups.

Specifically, we studied participants who possessed a high glaucoma PRS but did not have

glaucoma. For this study, we labeled them resilient while recognizing there are many reasons

they may not have glaucoma. Participants were stratified into PRS deciles, and metabolite

signatures were identified to differentiate patients with and without glaucoma within the top

decile (N=11,770) and in the bottom half (N=58,358) of the glaucoma PRS distribution (**Figure

4). Within the top decile of glaucoma PRS, compared to participants without glaucoma,

participants with glaucoma were more likely to be older, male, of White ethnicity, and were prior

smokers. Glaucoma participants with the highest PRS also had higher BMI, higher HbA1C,

higher spherical equivalent, consumed more caffeine and alcohol, and were more likely to have

diabetes and coronary artery disease (**Table 3**). Among participants in the bottom half of

glaucoma PRS, participants with glaucoma were also more likely to be older, male, of Black and

Asian ethnicity, prior smokers, had higher BMI and higher A1C, and were more likely to have

diabetes and coronary artery disease (**Table 4**). As expected, participants with glaucoma in both

bins of glaucoma genetic risk had higher IOP and thinner mRNFL thickness.

Our multivariable-adjusted analysis revealed that higher probit-transformed levels of

lactate (adjusted P_{NEF}=8.8E-12), pyruvate (adjusted P_{NEF}=2.9E-10), and citrate (adjusted

P_{NEF}=0.018) were independently associated with no glaucoma in the top decile of glaucoma PRS

(**Table 5**). In addition, lower levels of triglycerides and higher levels of selected HDL analytes

had an adjusted $P_{NEF} < 0.2$ and were associated with no glaucoma among participants with high

genetic risk. Among the bottom half of the PRS distribution, higher levels of albumin were

associated with no glaucoma (adjusted P_{NEF}=0.047), while higher levels of small HDL, omega-3

fatty acids, docosahexaenoic acid, lactate, and citrate were associated with no glaucoma, albeit

with an adjusted $P_{NEF} < 0.2$.

Interaction of metabolic and genetic biomarkers

The three metabolites (lactate, pyruvate, and citrate) for which higher levels were

associated with reduced glaucoma prevalence in individuals with high genetic susceptibility had

no statistically significant relationship with glaucoma for participants in the lower half of the

PRS distribution. Thus, we hypothesized that there may be an interaction between these

metabolites and genetic risk. Such an interaction would suggest that these

glycolysis/tricarboxylic acid cycle (TCA) metabolites are relevant primarily in the setting of high

PRS, establishing them as resilience factors against elevated genetic risk of glaucoma. Indeed,

we observed a significant interaction between elevated levels of total lactate, pyruvate, and

citrate with glaucoma PRS for predicting the risk of glaucoma ($P_{interaction}=0.011$), as shown in

Figure 5A.

To confirm and visualize this interaction, we calculated and plotted the glaucoma odds

ratio (OR) as a function of PRS quartile with each genetic predisposition bin stratified by the

sum of the resilience-metabolite probit score quartiles (**Figures 5A and 5B**). Notably, high

levels of resilience-associated metabolites were significantly associated with lower odds of

glaucoma, particularly within the higher PRS quartiles (Q3 and Q4). For example, for individuals

in PRS Q3 with total resilience-associated metabolite sum Q3, the glaucoma OR was 0.78 (95%

CI=0.65-0.93; P=0.0059), using metabolite sum Q1 as the reference. In the same PRS Q3 with a

higher quartile of total resilience-associated metabolite sum (Q4), the OR was 0.74 (95%

CI=0.62-0.88; P=0.0009). The lowest OR of glaucoma was observed in participants with both

the highest PRS (Q4) and highest resilience-associated metabolite sum (Q4), with an OR of 0.71

(95% CI=0.64-0.80; P<0.001). Although both age and PRS were significant predictors of

glaucoma prevalence (both are strongly associated with increased risk), there was no evidence of

a significant three-way interaction between the resilience-associated metabolite sum, PRS, and

age ($P_{3\text{-}way\text{-}interaction}=0.65$).

Next, we investigated whether a holistic metabolic risk score (MRS) incorporating

resilience metabolites and other measured metabolites (the total of 168 metabolites; see Methods

and **Supplemental Table 3** for betas related to MRS construction) can be used in conjunction

with a glaucoma PRS to predict risk. To quantify the degree of interaction of glaucoma PRS and

MRS, we plotted glaucoma OR at various quartiles of PRS and MRS compared to the first MRS

quartile within each PRS quartile. We found a significant interaction between glaucoma MRS

built from 168 metabolites and PRS ($P_{interaction}$=0.0012; see **Supplemental Figure 1A**).

In stratified analyses by quartile of PRS, compared to the first quartile of MRS, higher

MRS was associated with increased glaucoma prevalence across most PRS quartiles, with the

most pronounced effect observed in individuals in the highest PRS and MRS quartile with an OR

of 2.14 compared to those in the highest PRS and lowest MRS quartile (**Supplemental Figure**

1B). Among individuals in the highest MRS quartile (Q4), PRS Q1 had an OR of 1.66 (95%

CI=1.15-2.42; P=0.0086), PRS Q2 an OR of 2.08 (95% CI=1.62-2.67; P<0.001), PRS Q3 an OR

of 2.20 (95% CI=1.83-2.65; P<0.001), and PRS Q4 an OR of 2.14 (95% CI=1.91-2.40;

P<0.001). This suggests potential synergistic effects of genetic and metabolite risk factors on

glaucoma risk that transcend the impacts of the resilience metabolites.

Most importantly, using individuals in the lowest quartiles of PRS and MRS (PRS Q1,

MRS Q1) as the reference group for the entire population (n=117,698), we determined glaucoma

OR as a function of PRS quartile, with further stratification by holistic MRS in each PRS bin

(**Figure 6**). The OR for PRS Q1 with MRS Q4 relative to PRS Q1 with MRS Q1 was modest

(OR=1.66; 95% CI= 1.15–2.42; P=0.0086). However, those with PRS Q4 (highest genetic risk)

and MRS Q1 (lowest metabolic risk) had an OR of 11.7 (95% CI=8.72-16.0; P<0.001) compared

to PRS Q1 MRS Q1. This illustrates the importance of genetics in impacting glaucoma risk

($P_{interaction}$=0.019). Strikingly, the glaucoma OR further increased to 25.1 (95% CI=18.8-34.1;

P<0.001) for those with PRS Q4, MRS Q4 (combination of highest genetic and highest

metabolic risk). Altogether, this demonstrates the strong prognostic utility of combining both

PRS and MRS measurements for assessing glaucoma risk, particularly in individuals with a high

genetic predisposition.

Pyruvate supplementation lessens intraocular pressure and glaucoma

To functionally test the association between higher levels of pyruvate and resilience to

glaucoma, we experimentally tested whether treatment with pyruvate induces resilience to IOP

elevation and glaucoma in a human-relevant mouse model. Mutations in *LMX1B* contribute to a

spectrum of human glaucoma including the most common form, primary open-angle

glaucoma.[23–28] We have previously demonstrated that mice with a dominant mutation in *Lmx1b*

(*Lmx1b*$^{V265D/+}$) develop IOP elevation and glaucoma.[29,30] Depending on genetic background, this

Lmx1b^{V265D} mutation induces either early-onset or later glaucoma[30]. Mutant mice with a

C57BL/6J strain background develop severe, early-onset, IOP elevation and glaucoma. As

pyruvate and its metabolites were associated with no glaucoma despite strong genetic

predisposition in the UK Biobank cohort (highest decile of PRS), we tested the ability of dietary

pyruvate to induce resilience against the *Lmx1b*^{V265D}-induced glaucoma on this C57BL/6J

genetic background. Pyruvate supplementation through drinking water substantially protected

mice from IOP elevation and glaucoma. Pyruvate significantly protected against both anterior

chamber deepening (ACD), a consequence of IOP elevation in mouse eyes (**Figure 7A-B**), and

IOP elevation itself (**Figure 7C**). Importantly, pyruvate treatment protected against

glaucomatous optic nerve degeneration (**Figure 7D**). Together, our findings strongly support the

role of endogenous pyruvate in conferring resilience against glaucoma even countering strong

genetic predisposition. Further, they show that pyruvate can act as a potent resilience factor

against IOP elevation and glaucoma when delivered orally.

Discussion

In this large cohort study, we found that the inclusion of metabolite data from an NMR platform only modestly improved various glaucoma prediction algorithms **(Figure 3)**. Nonetheless, using an agnostic approach, we found that lactate, pyruvate, and citrate levels collectively were associated with a 29% reduced risk of glaucoma among those in the top quartile of glaucoma genetic predisposition **(Figure 5B)**. We validated pyruvate as a glaucoma resilience factor by demonstrating that pyruvate supplementation reduced glaucoma incidence in a human-relevant genetic mouse model (**Figure 7**). Furthermore, the interaction between glaucoma PRS and an MRS based on 168 metabolites provided synergistic predictive information regarding glaucoma risk (**Figure 6**). The statistical interaction we demonstrate between metabolite scores and PRS-derived glaucoma risk helps identify individuals who, despite high genetic risk, are less likely to develop glaucoma due to favorable metabolic profiles or on the contrary, are more likely to develop glaucoma and should be monitored at an earlier age. Overall, these findings indicate that an MRS may provide clinically useful measures of metabolic state and improve patient risk stratification in glaucoma.

While earlier studies[31–33] showed modest discriminatory powers for a glaucoma PRS, recent studies backed by larger genome-wide association studies have shown significantly improved risk prediction.[7,34–36] However, no studies have examined the utility of incorporating metabolite data into glaucoma prediction algorithms combined with PRS. Future work is needed to understand how non-genetic factors known to alter glaucoma risk or IOP levels like air pollution, psychological stress, physical activity, and dietary factors,[14] could change the MRS and alter high glaucoma genetic predisposition.

Previous studies have explored metabolomics to identify open-angle glaucoma (OAG)

biomarkers. A systematic review identified 13 studies with a total of 144 metabolites, of which

12 metabolites were identified to be associated with OAG.[21] In studies using plasma samples,

four metabolic pathways were significantly enriched: sphingolipid metabolism, arginine and

proline metabolism, and beta-alanine metabolism. In another study among three US cohorts and

the UKBB, higher levels of plasma diglycerides and triglycerides were found to be adversely

associated with glaucoma.[37] This is consistent with our observation that lower levels of

triglycerides are associated with glaucoma resilience, albeit at less significance compared to the

three glycolysis-related metabolites. The biological mechanism is suggested to be the association

of hypertriglyceridemia with increased blood viscosity and increased IOP.[38–41] A meta-analysis

showed that patients with glaucoma had higher mean triglyceride levels with an absolute

difference of 14.2 mg/dL compared to patients without glaucoma.[40]

While studies have identified biomarkers associated with glaucoma, to the best of our

knowledge, there has been no investigation on biomarkers that are associated with being resilient

to a high genetic predisposition to glaucoma. We identified elevated levels of lactate, pyruvate,

and citrate among individuals without glaucoma despite high genetic risk. The retina commonly

utilizes energy produced from glycolysis and oxidative phosphorylation. Therefore, glycolysis

end products (pyruvate and lactate) are important energy sources for retinal ganglion cells

(RGCs).[42] Lactate is critical for RGC survival during periods of glucose deprivation,[43,44] and

bioenergetic insufficiency contributes to RGC loss in glaucoma.[45–48] Finally, a study found that

baseline levels of lactate were significantly decreased in patients with normal-tension glaucoma

compared to controls without optic neuropathy.[49]

Pyruvate is central in energy metabolism linking glycolysis to the TCA and promoting

ATP production by the electron transport chain. Pyruvate is converted into acetyl-CoA, which

drives the TCA, and into lactate, a key molecule that regulates metabolism and serves as an

important energy source.[50] The TCA is precisely regulated to determine the appropriate balance

between biosynthesis (producing the metabolic precursors for lipids, carbohydrates, amino acids,

nucleic acids, and co-factors) and energy production depending on the real-time need to enhance

cell functions and survival.[51] Thus, pyruvate is pivotal in metabolic control and cellular health.

Pyruvate also scavenges reactive oxygen species (ROS), can induce expression of the antioxidant

response control gene *Nrf2*, and can promote beneficial autophagy.[52–54] Neuroprotective effects

of pyruvate on RGCs are demonstrated in cell culture and rodent eyes, including an induced

model of glaucoma in rats[55] with enhanced protection when nicotinamide and pyruvate were

both administered in mice.[56] Importantly, the current study extends this to strong protection

against glaucoma by pyruvate alone in a mouse model with a mutation in the ortholog of a

human POAG gene. Further, it extends the protective effects of pyruvate to the ocular drainage

tissues and the lessening of IOP elevation. A randomized Phase II clinical trial that tested

pyruvate and nicotinamide in POAG patients found improved visual function in glaucoma

patients compared to the placebo group.[57] Thus, both preclinical and clinical evidence agree with

our findings that pyruvate serves as a protective factor against glaucoma.

Citrate is a key intermediate in the TCA cycle and is important for aerobic energy

production. Elevated citrate levels might reflect enhanced energy supply and antioxidative

capacity of ocular tissues, potentially protecting RGCs against oxidative stress.[47,58–61] Decreased

levels of citrate may indicate increased usage or mitochondrial dysfunction.[62] Studies have found

lower levels of plasma citrate in glaucoma patients among adults and children.[63,64] A study

measuring plasma citrate with a cutoff of 110 μmol/L to detect glaucoma had a sensitivity of

66.7% and a specificity of 71.4%.[63] These findings corroborate the protective effects of citrate in

conferring resilience to glaucoma.

We also identified several lipoproteins associated with resilience, most notably

cholesteryl esters in small high-density lipoprotein (HDL) and medium HDL. This is consistent

with studies describing the neuroprotective effect of HDL in the pathophysiology of

glaucoma.[65,66] Cholesteryl esters and triglycerides make up the core of lipoproteins, which

transport lipids in the blood. Lipoproteins are characterized by their size, where HDL ranges

from 8-12nm in diameter, and low-density lipoprotein (LDL) ranges from 18-25nm in diameter.

The retinal pigment epithelium can take up the lipoproteins.[67,68] However, dysregulation of lipid

metabolism can result in the accumulation of lipid and oxidative stress in the retina, resulting in

RNFL thinning.[37,69–72] Thus, HDL has antioxidant properties to protect RGCs and facilitate

cholesterol efflux from accumulating in the retina.[73,74] Notably, genes implicated by large cross-

ancestry and European subset GWAS meta-analyses of POAG risk are enriched in processes

related to apolipoprotein binding, lipid transport, and decreased circulating high-density

lipoprotein cholesterol levels.[25,75]

This study had several strengths. First, the UKBB is a large cohort study with detailed

covariate information on demographics, clinical characteristics, and endophenotypes for

glaucoma, allowing for better glaucoma risk profiling. Second, we used a PRS calculated from

the large (~600k individuals) multi-trait analysis of genome-wide association studies on

glaucoma and endophenotypes, including optic nerve head structural features and IOP data.[7]

Finally, we supported our biobank analyses with functional tests in a disease-relevant mouse

model, identifying a protective effect of pyruvate supplementation in animals susceptible to

356 glaucoma. The data showing that pyruvate protected against IOP elevation (a key causative risk

357 factor for glaucoma) and glaucomatous nerve damage in a mouse model, as well as pyruvate's

358 biochemical relatedness to lactate and citrate regarding energy metabolism, mitigate against

359 concerns about false discovery, uncontrolled confounding, reverse causation and collinearity

360 between biomarkers.

361   A major limitation of our study is that it includes a predominately European population,

362 with an underrepresentation of African, Asian, and mixed American genetic ancestries, limiting

363 the generalizability of our findings in more heterogeneous populations. In addition, a subset of

364 UKBB participants have received glaucoma treatment, which could alter metabolite levels and

365 could lead to an underestimate of the protective effects of metabolites. Moreover, the UKBB

366 study includes a limited set of 168 metabolites which is not inclusive of the comprehensive set of

367 metabolites, known and unknown. This study also only focuses on plasma metabolites and does

368 not account for ocular metabolite profiles. Our study was cross-sectional, making it difficult to

369 determine the temporality; therefore, it is unclear whether certain metabolic changes observed

370 drive the onset of glaucoma or are a consequence of the disease. However, the mouse data

371 support a causal role for pyruvate as a factor that mitigates high genetic risk. The exploratory

372 nature of this study, in which we investigated metabolite associations in an agnostic, hypothesis-

373 free manner, highlights the need for replication of our findings, although we review considerable

374 evidence that provides support for our findings. Concerning the mouse studies, we performed ad-

375 lib pyruvate dosing, and we only tested one human-related mutation, while humans with

376 glaucoma genetic predisposition had a collective high dose of many common glaucoma-related

377 variants. Lastly, neither the mouse studies nor the human data accurately inform the exact dosing

378 of TCA biomarkers needed to effectively mitigate glaucoma risk. We find considerable overlap

between TCA biomarkers between those with and without glaucoma in the top decile of

glaucoma risk (**Supplementary Table 1** and **Supplementary Figure 2**).

Overall, we have shown that plasma metabolites modestly enhance the predictive power

of PRS and may have a role in identifying those who may be resilient to glaucoma despite high

genetic predisposition. Our study identified protective metabolites linked to glycolysis and

mitochondrial function, suggesting important pathophysiology in glaucoma. One of these

metabolites of significance is pyruvate. Oral pyruvate supplementation substantially protected

against IOP elevation and conferred resilience against glaucoma. Thus, these results open new

avenues for therapeutic strategies of pyruvate and its metabolites as resilience factors against

glaucoma in genetically predisposed individuals.

Methods

Study Design

Our initial human analysis was conducted in three steps. First, we assessed whether

plasma metabolite data alone can predict glaucoma risk with receiver operating characteristic

(ROC) curves in the UK Biobank. We then integrated the metabolite data into a polygenic risk

score (PRS)-based glaucoma risk assessment model to see if metabolites could enhance

glaucoma prediction. Second, we focused on identifying a metabolomic signature of resilience to

high glaucoma PRS by comparing plasma metabolites in glaucoma cases versus participants

without glaucoma (henceforth referred to as resilient participants) from the top 10% of the PRS.

Third, we explored the interactions between metabolic scores and PRS in modifying glaucoma

risk.

	Study population and data collection

The UKBB is a prospective cohort study of over half a million participants aged 37 to 73

years at recruitment across the United Kingdom from 2006 to 2010. Participants were recruited

through the National Health Service registers from 22 assessment centers, where they signed

electronic informed consent to participate. Participants then completed in-depth touchscreen

questionnaires and trained staff-led interviews, performed body measurements, and provided

biological samples.[76] Additional outcomes are available from data linkage to hospital episode

statistics, the death register, and primary data.[76,77] Biological samples collected from these

participants, including blood, urine, and saliva specimens, were used to generate genetic,

metabolomic, and proteomic data.[78] Details of the UKBB study design and population are

described online (https://www.ukbiobank.ac.uk). The UKBB study received approval from the

National Health Service North West Multicentre Research Ethics Committee (reference number

06/MRE08/65) and the National Information Governance Board for Health and Social Care. This

research, conducted under UK Biobank application number 36741, adhered to the tenets of the

Declaration of Helsinki.

The UKBB included 502,613 participants, of which 173,679 participants had self-

reported glaucoma data, ICD-coded, or previous glaucoma laser/surgical therapy (34.6%) at

baseline. A subset consisting of 117,698 participants with metabolomic data and genetic

profiling comprise the study population (**Figure 1**). The glaucoma cases were defined based on

those with self-reported glaucoma, in which participants selected "glaucoma" when they

completed a touchscreen questionnaire with the question, "Has a doctor told you that you have

any of the following problems with your eyes?" Glaucoma cases also included participants who

reported a history of glaucoma surgery or laser therapy on the questionnaire or if an International

Classification of Diseases (ICD) code for glaucoma (ICD 9[th] revision: 365.* [excluding 365.0];

ICD 10[th]s revision: H40.* [excluding H40.0] and H42.*) was carried in linked Hospital Episode

Statistics before the baseline assessment. The described approach to identifying glaucoma has

been strongly supported by prior research publications.[26,27,79]

Metabolite Profiling

In the UKBB, a high-throughput nuclear magnetic resonance (NMR)-based biomarker

platform (Nightingale Health Ltd; Helsinki, Finland) was used to measure the metabolomic

profile in the randomly selected non-fasting EDTA plasma samples from a subset of

participants.[80] In contrast to liquid chromatography/mass spectroscopy, NMR spectroscopy

produces distinctive spectral shapes for molecules containing hydrogen atoms.[81] The areas under

the curve (AUC) are proportional to the concentration for each molecule based on chemical

shifts and J coupling split patterns derived from quantum mechanics.[82,83] The NMR platform

contains data on 249 metabolic biomarkers (168 absolute levels and 81 ratio measures),

including a subset of 36 biomarkers (27 absolute levels and 9 ratio measures) certified for broad

diagnostic use by the European Union (EU).[80] In this study, we focused on the 168-metabolite

and the 27-metabolite sets for analysis (for a complete list of quantified metabolites, refer to the

Supplementary File). Individual metabolite values were transformed to probit scores to

standardize their range and reduce the impact of skewed distributions.

Construction of PRS for glaucoma

The glaucoma PRS used in this study was from a multi-trait analysis of GWAS (MTAG)

on glaucoma.[7] This PRS was derived based on GWAS data from glaucoma (7,947 cases and

119,318 controls) and its endophenotypes: optic nerve head structure using vertical cup-disc ratio

(VCDR) (including additional data from 67,040 UKBB and 23,899 International Glaucoma

Genetics Consortium, IGGC, participants), and intraocular pressure (IOP, including additional

data on 103,914 UKBB and 29,578 IGGC participants). The glaucoma PRS was constructed

using single nucleotide polymorphisms (SNPs) with MTAG P-values $\leq$ 0.001, resulting in 2,673

uncorrelated SNPs after LD-clumping at r^2 =0.1, and was found to have a predictive ability with

an area under the curve (AUC) of 0.68 (95% confidence interval, CI, [0.67-0.70]) and 0.80

accounting for age, sex, family history, and PRS tested in the Australian and New Zealand

Registry of Advanced Glaucoma cohort.

Model building/covariates

We first assessed whether the inclusion of probit-transformed metabolite data enhances

glaucoma prediction algorithms. We adjusted for factors involving major determinants of

variability in metabolites, POAG-established and suspected risk factors, and other comorbidities.

In model 1, we considered metabolites only. Model 2 incorporated additional demographic

covariates, including age (years), sex, genetic ancestry, season, time of day of specimen

collection (morning, afternoon, night), and fasting time (hours). Model 3 incorporated covariates

in model 2 and additional variables accounting for comorbidities potentially related to glaucoma

including smoking status (never, past, and current smoker), alcohol intake (g/week),[10] caffeine

intake (mg/day),[84] physical activity (metabolic equivalent of task [MET], hours/week), body

mass index (kg/m^2), average systolic blood pressure (mm Hg), history of diabetes, hemoglobin

A1C (HbA1c, mmol/mol), self-reported history of coronary artery disease, systemic beta-blocker

use, oral steroid use, and mean spherical equivalent refractive error (diopters) across both eyes.

Model 4 incorporated covariates in model 3 and a glaucoma PRS. We compared these models

with and without the metabolites to determine the predictive ability of the metabolites above and

beyond a glaucoma PRS and known risk factors. Missing values were imputed with medians for

numeric variables and modes for factor variables.

Statistical analysis

Metabolite-Based Predictive Modeling

Multivariable logistic regression models with regularization (implemented with *glmnet*)

were built to investigate the associations of metabolic biomarkers and the risk of glaucoma. We

used four sequential models, adjusting for covariates described above. For each model, we

examined three groupings: no metabolites, the 27 metabolites corresponding to EU-stamped

validated markers, and 168 metabolites corresponding to all the measured plasma metabolites

captured in the UKBB. We used regularization to address for collinearity, which reduced the set

of metabolites and other covariates considered in each model (reported in **Supplementary Table**

2). To examine model accuracy for predicting glaucoma prevalence, we utilized ROC curves

and associated AUC measurements as a metric for model performance. We performed five-fold

cross-validation and split data where 80% were used for training, and 20% was used for

evaluation. All metabolites' values were probit transformed and used as continuous variables

(per 1 standard deviation (SD) increase). Additionally, we stratified glaucoma by age, ethnicity,

and sex. The DeLong test was utilized to examine the statistical significance of AUC differences

between models, with a threshold of $P<0.05$ as significant.

Identifying resilience metabolites to high glaucoma polygenic risk scores

To identify individual metabolites associated with resilience to a high glaucoma PRS, we

first performed a logistic regression model to obtain metabolite residuals from probit-

transformed concentration data adjusting for the following variables: age, age-squared, time

since the last meal/drink (≤4, 5–8, and 9+ hours), sex, ethnicity (Asian, Black, White, and

mixed/other), season, time of day of specimen collection, smoking status (never, past, and

current smoker), alcohol intake, caffeine intake, physical activity (metabolic equivalent of task

[MET] hours/week), body mass index (kg/m^2), average systolic blood pressure (mm Hg), history

of diabetes, HbA1c (mmol/mol), history of coronary artery disease, systemic beta-blocker use,

oral steroid use, and spherical equivalent refractive error (diopters). Metabolites associated with

glaucoma were nominated by performing a t-test comparing residuals in participants with and

without glaucoma to identify metabolites significantly associated with glaucoma in both the top

10% and the bottom 50% of glaucoma PRS score. This stratification balanced case counts to

enable comparison of groups with adequate sample size (780 glaucoma cases in the bottom 50%,

and 1,693 cases in the top 10% of PRS score). To account for covariance in metabolite

abundance, we corrected for multiple comparisons using the number of effective tests (NEF)

method.[85] First, a correlation matrix was generated using pairwise complete observations to

address any missing values. Eigenvalue decomposition was performed on the correlation matrix

to capture the principal components. Principal components contributing more than 1% of the

total variance were calculated, yielding 9 significant components, capturing 91.6% of the

variance. Thus, we applied a Bonferroni adjustment based on the number of significant

components (n=9) to calculate the adjusted p-values accounting for multiple hypotheses. P-

values are considered statistically significant if the NEF-adjusted P-value was <0.05.[85] Given this

was an exploratory analysis, P-values considered worthy of additional analysis if the NEF-adjust

P-value <0.2.

Construction of Metabolite Risk Scores

We constructed metabolite risk scores (MRS), which were calculated as a weighted sum

of each metabolic biomarker from coefficients in the 168-metabolite base model. Specifically,

we utilized a logistic regression model (implemented with a *binomial glm*) using probit

transformed metabolite values to predict the prevalence of glaucoma using five-fold cross-

validation. The resulting beta values were extracted and are reported in **Supplementary Table 3**.

To determine if there is an interaction between glaucoma MRS and PRS, we utilized extracted

model coefficients and corresponding significance from a binomial generalized linear model. We

fit this model using the formula *Glaucoma ~ PRS * MRS + PRS + MRS + Covariates.* We

classified participants into 16 groups by quartiles of MRS and PRS, defined as quartiles. Within

each PRS quartile, we used MRS quartile 1 as the reference for comparison. Additionally, we

used the group in both PRS quartile 1 and MRS quartile 1 as the overall reference group for all

comparisons among the groups. A chi-squared test was used to assess statistical significance

between odds ratios (ORs) with 95% confidence intervals (CIs).

To investigate whether age modifies the associations of resilience metabolites with

glaucoma, we conducted a stratified analysis and assessed a three-way interaction term of age ($\geq$

or < age 58 years, based on median age), the sum of resilience metabolite levels, and PRS using

a Wald test for individuals in the top 10% PRS. All statistical analyses and plots were produced

using R version 4.2.1 (R Foundation for Statistical Computing, Vienna, Austria). All statistical

tests were two-sided.

Animal husbandry and ethics statement

Experimental mice had a C57BL/6J (Jackson Laboratory Stock #000664) genetic

background. The *Lmx1b*[V265D] (alias *Lmx1b*[Icst]) mutation is previously reported to cause high IOP

and glaucoma in mice with IOP becoming elevated in some eyes during the first months after

birth.[29,30,86] The mutation was backcrossed to the C57BL/6J strain for at least 30 generations. All

mice were treated in accordance with the Association for Research in Vision and

Ophthalmology's statement on the use of animals in ophthalmic research. The Institutional

Animal Care and Use Committee of Columbia University approved all experimental protocols

performed.

Genotyping mice

Lmx1b^{V265D} and *Lmx1b^{+}* genotypes were determined by direct Sanger sequencing of a

specific PCR product. Genomic DNA was PCR amplified with forward primer 5′-

CTTTGAGCCATCGGAGCTG-3′ and reverse primer 5′-ATCTCCGACCGCTTCCTGAT-3′

using the following program: (1) 94 °C for 3 min; (2) 94 °C for 30 s; (3) 57 °C for 30 s; (4) 72 °C

for 1 min; (5) repeat steps 2–4 35 times; and (6) 72 °C for 5 min. PCR products were purified and

sequenced by the Genewiz (Azenta Life Sciences).

Slit-lamp examination

Lmx1b^{V265D} and *Lmx1b^{+}* genotypes were determined by direct Sanger sequencing of a

specific PCR product. Genomic DNA was PCR amplified with forward primer 5′-

CTTTGAGCCATCGGAGCTG-3′ and reverse primer 5′-ATCTCCGACCGCTTCCTGAT-3′

using the following program: (1) 94 °C for 3 min; (2) 94 °C for 30 s; (3) 57 °C for 30 s; (4) 72 °C

for 1 min; (5) repeat steps 2–4 35 times; and (6) 72 °C for 5 min. PCR products were purified and

sequenced by the Genewiz (Azenta Life Sciences).

Intraocular pressure measurement

IOP was measured with the microneedle method as previously described in detail.[87,88]

Before cannulation, mice were acclimatized to the procedure room and anesthetized via an

intraperitoneal injection of a mixture of ketamine (99 mg/kg; Ketlar, Parke-Davis, Paramus, NJ,

USA) and xylazine (9 mg/kg; Rompun, Phoenix Pharmaceutical, St Joseph, MO, USA)

immediately prior to IOP assessment, a procedure that does not alter IOP in the experimental

window.[88] IOP was measured at both 5-6 weeks and 8-10 weeks of age in WT and $Lmx1b^{V265D/+}$

eyes. Balanced groups of males and females were examined. During each IOP measurement

period, the eyes of independent WT B6 mice were assessed in parallel with experimental mice as

a methodological control to ensure proper calibration and equipment function. *Lmx1b* mutant

groups were compared by ANOVA followed by Tukey's honestly significant difference. n > 30

eyes examined in each $Lmx1b^{V265D}$ mutant group and n > 20 eyes were examined in WT groups.

Pyruvate administration

Ethyl pyruvate (Sigma-Aldrich, St. Louis, Missouri) was dissolved in the standard

institutional drinking water to a dose of 2000 mg/kg/day for adult mice based on the average

volume mice consume. The mothers consumed this dose and delivered it to their suckling pups at

an unknown dose throughout the first 3 to 4 weeks of life. At 4 weeks of age, the mice were

weaned into separate cages. From 4 weeks on, the young mice consumed a dose of

approximately 2000 mg/kg/day based on their water consumption and body weight over weekly

intervals. Untreated groups received the same drinking water without pyruvate. The water was

changed once per week. Treatment was started at postnatal day 2. Births were checked daily

between 9am-12pm to determine the pup's age.

Optic nerve assessment

We analyzed $Lmx1b^{V265D/+}$ and WT controls optic nerve for glaucomatous damage at 5.5

months of age (5.3-5.9, sex balanced). Intracranial portions of optic nerves were dissected,

processed, and analyzed as previously described.[45,89,90] Briefly, optic nerve cross-sections were

stained with para-phenylenediamine (PPD) and examined for glaucomatous damage. PPD stains

all myelin sheaths, but differentially darkly stains the myelin sheaths and the axoplasm of

damaged axons. This allows for the sensitive detection and quantification of axon damage and

loss. Optic nerves were prepared for analysis with a 48-h fixation in 0.8% paraformaldehyde and

1.2% glutaraldehyde in 0.08 M phosphate buffer (pH 7.4) at 4°C followed by overnight post-fix

in osmium tetroxide at 4°C. Nerves were washed twice for 10 min in 0.1 M phosphate buffer,

once in 0.1 M sodium-acetate buffer, and dehydrated in graded ethanol concentrations. Tissues

were then embedded in Embed 812 resin (Electron Microscopy Sciences, Fort Washington, PA,

USA), and 1-µm-thick sections were stained in 1% PPD for ~40 minutes. Stained sections were

compared using a damage scale that is validated against axon counting.[89,91] Multiple sections of

each nerve were considered when determining damage level. Nerves were determined to have

one of 4 damage levels: 1) No glaucoma (NOE) – less than 5% axons damaged. This level of

damage is seen in age- and sex-matched non-glaucomatous mice and is not due to glaucoma. We

named this level no or early stage as some have early molecular changes when assessed

transcriptomically, but they cannot be distinguished from control by morphology;[47,92] 2)

Moderate damage (MOD) – average of 30% axon loss.; 3) Severe (SEV) – >50% axonal loss and

extensive axon damage; and 4) Very severe (V. SEV) – glial scar over the vast majority of nerve

with few remaining axons.

Acknowledgements

599   The authors would like to thank the participants who contributed their data to the UK Biobank

600   study and for their contributions to the study. The authors would also like to thank the members

601   the Simon John Laboratory for experimental and technical assistance and the Institute of

602   Comparative Medicine for animal and veterinary care.

Consortia

UK Biobank Eye and Vision Consortium

Pirro Hysi & Anthony Khawaja

CRediT authorship contribution statement

Keva Li: Writing – original draft, Writing – review & editing, Methodology, Investigation, Formal analysis, Conceptualization. Nicholas Tolman: original draft, Writing – review & editing, Methodology, Investigation, Formal analysis, Conceptualization. Ayellet V. Segrè: Writing – review & editing, Methodology, Investigation. Kelsey V. Stuart: Data Curation, Methodology, Resources, Writing – review & editing, Methodology, Investigation. Oana A. Zeleznik: Writing – review & editing, Investigation. Neeru A. Vallabh: Writing – review & editing, Investigation. Kuang Hu: Writing – review & editing, Investigation. Nazlee Zebardast: Writing – review & editing, Investigation. Akiko Hanyuda: Writing – review & editing, Investigation. Yoshihiko Raita: Writing – review & editing, Investigation. Christa Montgomery: Writing – review & editing, Investigation. Chi Zhang: Writing – review & editing, Investigation. Pirro G. Hysi: Writing – review & editing, Resources, Project administration, Methodology, Investigation, Funding acquisition. Ron Do: Writing – review & editing, Investigation. Anthony Khawaja: Writing – review & editing, Supervision, Resources, Project administration, Methodology, Investigation, Funding acquisition, Conceptualization. Janey L. Wiggs: Writing – review & editing, Investigation, Funding acquisition. Jae H. Kang: Writing – review & editing, Supervision, Resources, Methodology, Investigation, Funding acquisition. Simon WM. John: Writing – review & editing, Supervision, Resources, Project administration, Methodology, Investigation, Conceptualization. Louis R. Pasquale: Writing – review & editing, Supervision, Resources, Project administration, Methodology, Investigation, Funding acquisition,

626   Conceptualization. All authors were involved in the interpretation of the results, and the critical

627   review and approval of the manuscript.

Data Availability

629   Data from the UK Biobank cannot be shared due to our Material Transfer Agreement. To request

630   access to the UK Biobank data, please make requests directly to the UK Biobank via

631   https://www.ukbiobank.ac.uk/enable-your-research/apply-for-access.

References

1. Soh Z, Yu M, Betzler BK, et al. The Global Extent of Undetected Glaucoma in Adults: A
Systematic Review and Meta-analysis. *Ophthalmology*. 2021;128(10):1393-1404.
doi:10.1016/j.ophtha.2021.04.009

2. Khawaja AP, Viswanathan AC. Are we ready for genetic testing for primary open-angle
glaucoma? *Eye*. 2018;32(5):877-883. doi:10.1038/s41433-017-0011-1

3. Jan C, He M, Vingrys A, Zhu Z, Stafford RS. Diagnosing glaucoma in primary eye care and
the role of Artificial Intelligence applications for reducing the prevalence of undetected
glaucoma in Australia. *Eye*. 2024;38(11):2003-2013. doi:10.1038/s41433-024-03026-z

4. Moyer VA, US Preventive Service Task Force. Screening for Glaucoma: U.S. Preventive
Services Task Force Recommendation Statement. *Ann Intern Med*. 2013;159(7):484-489.
doi:10.7326/0003-4819-159-6-201309170-00686

5. US Preventive Services Task Force. Screening for Primary Open-Angle Glaucoma: US
Preventive Services Task Force Recommendation Statement. *JAMA*. 2022;327(20):1992-
1997. doi:10.1001/jama.2022.7013

6. Zukerman R, Harris A, Oddone F, Siesky B, Verticchio Vercellin A, Ciulla TA. Glaucoma
Heritability: Molecular Mechanisms of Disease. *Genes*. 2021;12(8):1135.
doi:10.3390/genes12081135

7. Craig JE, Han X, Qassim A, et al. Multitrait analysis of glaucoma identifies new risk loci and
enables polygenic prediction of disease susceptibility and progression. *Nat Genet*.
2020;52(2):160-166. doi:10.1038/s41588-019-0556-y

8. de Vries VA, Hanyuda A, Vergroesen JE, et al. The Clinical Usefulness of a Glaucoma
Polygenic Risk Score in 4 Population-Based European Ancestry Cohorts. *Ophthalmology*.
Published online August 14, 2024. doi:10.1016/j.ophtha.2024.08.005

9. Sekimitsu S, Xiang D, Smith SL, et al. Deep Ocular Phenotyping Across Primary Open-Angle
Glaucoma Genetic Burden. *JAMA Ophthalmol*. 2023;141(9):891-899.
doi:10.1001/jamaophthalmol.2023.3645

10. Stuart KV, Luben RN, Warwick AN, et al. The Association of Alcohol Consumption
with Glaucoma and Related Traits: Findings from the UK Biobank. *Ophthalmol Glaucoma*.
2022;6(4):366. doi:10.1016/j.ogla.2022.11.008

11. Kang JH, Willett WC, Rosner BA, Buys E, Wiggs JL, Pasquale LR. Association of
Dietary Nitrate Intake With Primary Open-Angle Glaucoma: A Prospective Analysis From the
Nurses' Health Study and Health Professionals Follow-up Study. *JAMA Ophthalmol*.
2016;134(3):294-303. doi:10.1001/jamaophthalmol.2015.5601

12. Kang JH, Ivey KL, Boumenna T, Rosner B, Wiggs JL, Pasquale LR. Prospective study of
flavonoid intake and risk of primary open-angle glaucoma. *Acta Ophthalmol (Copenh)*.
2018;96(6):e692. doi:10.1111/aos.13705

13. Kang JH, Wu J, Cho E, et al. Contribution of the Nurses' Health Study to the
Epidemiology of Cataract, Age-Related Macular Degeneration, and Glaucoma. *Am J Public
Health*. 2016;106(9):1684. doi:10.2105/AJPH.2016.303317

14. Stuart KV, Pasquale LR, Kang JH, Foster PJ, Khawaja AP. Towards modifying the
genetic predisposition for glaucoma: An overview of the contribution and interaction of
genetic and environmental factors. *Mol Aspects Med*. 2023;93:101203.
doi:10.1016/j.mam.2023.101203

15. Nazifova-Tasinova N, Radeva M, Galunska B, Grupcheva C. Metabolomic analysis in
ophthalmology. *Biomed Pap*. 2020;164(3):236-246. doi:10.5507/bp.2020.028

16. Johnson CH, Ivanisevic J, Siuzdak G. Metabolomics: beyond biomarkers and towards
mechanisms. *Nat Rev Mol Cell Biol*. 2016;17(7):451-459. doi:10.1038/nrm.2016.25

17. Javadiyan S, Burdon KP, Whiting MJ, et al. Elevation of Serum Asymmetrical and
Symmetrical Dimethylarginine in Patients with Advanced Glaucoma. *Invest Ophthalmol Vis
Sci*. 2012;53(4):1923-1927. doi:10.1167/iovs.11-8420

18. Burgess LG, Uppal K, Walker DI, et al. Metabolome-Wide Association Study of Primary
Open Angle Glaucoma. *Invest Ophthalmol Vis Sci*. 2015;56(8):5020-5028.
doi:10.1167/iovs.15-16702

19. Kouassi Nzoughet J, Guehlouz K, Leruez S, et al. A Data Mining Metabolomics
Exploration of Glaucoma. *Metabolites*. 2020;10(2):49. doi:10.3390/metabo10020049

20. Leruez S, Marill A, Bresson T, et al. A Metabolomics Profiling of Glaucoma Points to
Mitochondrial Dysfunction, Senescence, and Polyamines Deficiency. *Invest Ophthalmol Vis
Sci*. 2018;59(11):4355-4361. doi:10.1167/iovs.18-24938

21.   Wang Y, Hou XW, Liang G, Pan CW. Metabolomics in Glaucoma: A Systematic
Review. *Invest Ophthalmol Vis Sci*. 2021;62(6):9. doi:10.1167/iovs.62.6.9

22.   Privé F, Aschard H, Carmi S, et al. Portability of 245 polygenic scores when derived
from the UK Biobank and applied to 9 ancestry groups from the same cohort. *Am J Hum*
Genet. 2022;109(1):12-23. doi:10.1016/j.ajhg.2021.11.008

23.   Choquet H, Paylakhi S, Kneeland SC, et al. A multiethnic genome-wide association study
of primary open-angle glaucoma identifies novel risk loci. *Nat Commun*. 2018;9:2278.
doi:10.1038/s41467-018-04555-4

24.   Gao XR, Huang H, Nannini DR, Fan F, Kim H. Genome-wide association analyses
identify new loci influencing intraocular pressure. *Hum Mol Genet*. 2018;27(12):2205-2213.
doi:10.1093/hmg/ddy111

25.   Gharahkhani P, Jorgenson E, Hysi P, et al. Genome-wide meta-analysis identifies 127
open-angle glaucoma loci with consistent effect across ancestries. *Nat Commun*.
2021;12:1258. doi:10.1038/s41467-020-20851-4

26.   Khawaja AP, Cooke Bailey JN, Wareham NJ, et al. Genome-wide analyses identify 68
new loci associated with intraocular pressure and improve risk prediction for primary open-
angle glaucoma. *Nat Genet*. 2018;50(6):778-782. doi:10.1038/s41588-018-0126-8

27.   MacGregor S, Ong JS, An J, et al. Genome-wide association study of intraocular pressure
uncovers new pathways to glaucoma. *Nat Genet*. 2018;50(8):1067-1071. doi:10.1038/s41588-
018-0176-y

28.   Shiga Y, Akiyama M, Nishiguchi KM, et al. Genome-wide association study identifies
seven novel susceptibility loci for primary open-angle glaucoma. *Hum Mol Genet*.
2018;27(8):1486-1496. doi:10.1093/hmg/ddy053

29.   Cross SH, Macalinao DG, McKie L, et al. A Dominant-Negative Mutation of Mouse
Lmx1b Causes Glaucoma and Is Semi-lethal via LBD1-Mediated Dimerisation. *PLoS Genet*.
2014;10(5):e1004359. doi:10.1371/journal.pgen.1004359

30.   Tolman NG, Balasubramanian R, Macalinao DG, et al. Genetic background modifies
vulnerability to glaucoma-related phenotypes in Lmx1b mutant mice. *Dis Model Mech*.
2021;14(2):dmm046953. doi:10.1242/dmm.046953

31.   Mabuchi F, Mabuchi N, Sakurada Y, et al. Additive effects of genetic variants associated
with intraocular pressure in primary open-angle glaucoma. *PLOS ONE*. 2017;12(8):e0183709.
doi:10.1371/journal.pone.0183709

32.   Tham YC, Liao J, Vithana EN, et al. Aggregate Effects of Intraocular Pressure and Cup-
to-Disc Ratio Genetic Variants on Glaucoma in a Multiethnic Asian Population.
Ophthalmology. 2015;122(6):1149-1157. doi:10.1016/j.ophtha.2015.01.024

33. Zanon-Moreno V, Ortega-Azorin C, Asensio-Marquez EM, et al. A Multi-Locus Genetic
Risk Score for Primary Open-Angle Glaucoma (POAG) Variants Is Associated with POAG
Risk in a Mediterranean Population: Inverse Correlations with Plasma Vitamin C and E
Concentrations. *Int J Mol Sci*. 2017;18(11):2302. doi:10.3390/ijms18112302

34. Han X, Souzeau E, Ong JS, et al. Myocilin Gene Gln368Ter Variant Penetrance and
Association With Glaucoma in Population-Based and Registry-Based Studies. *JAMA
Ophthalmol*. 2019;137(1):28-35. doi:10.1001/jamaophthalmol.2018.4477

35. Gao XR, Huang H, Kim H. Polygenic Risk Score Is Associated With Intraocular Pressure
and Improves Glaucoma Prediction in the UK Biobank Cohort. *Transl Vis Sci Technol*.
2019;8(2):10. doi:10.1167/tvst.8.2.10

36. Nannini DR, Kim H, Fan F, Gao X. Genetic risk score is associated with vertical cup-disc
ratio and improves prediction of primary open angle glaucoma in Latinos. *Ophthalmology*.
2018;125(6):815-821. doi:10.1016/j.ophtha.2017.12.014

37. Zeleznik OA, Kang JH, Lasky-Su J, et al. Plasma metabolite profile for primary open-
angle glaucoma in three US cohorts and the UK Biobank. *Nat Commun*. 2023;14(1):2860.
doi:10.1038/s41467-023-38466-w

38. Madjedi KM, Stuart KV, Chua SY, et al. The Association between Serum Lipids and
Intraocular Pressure in 2 Large United Kingdom Cohorts. *Ophthalmology*. 2022;129(9):986.
doi:10.1016/j.ophtha.2022.04.023

39. Wang S, Bao X. Hyperlipidemia, Blood Lipid Level, and the Risk of Glaucoma: A Meta-
Analysis. *Invest Ophthalmol Vis Sci*. 2019;60(4):1028-1043. doi:10.1167/iovs.18-25845

40. Pertl L, Mossböck G, Wedrich A, et al. Triglycerides and Open Angle Glaucoma - A
Meta-analysis with meta-regression. *Sci Rep*. 2017;7(1):7829. doi:10.1038/s41598-017-
08295-1

41. Rasoulinejad SA, Kasiri A, Montazeri M, et al. The Association Between Primary Open
Angle Glaucoma and Clustered Components of Metabolic Syndrome. *Open Ophthalmol J*.
2015;9:149-155. doi:10.2174/1874364101509010149

42. Williams PA, Casson RJ. Glycolysis and glucose metabolism as a target for bioenergetic
and neuronal protection in glaucoma. *Neural Regen Res*. 2024;19(8):1637. doi:10.4103/1673-
5374.389638

43. Vohra R, Aldana BI, Bulli G, et al. Lactate-Mediated Protection of Retinal Ganglion
Cells. *J Mol Biol*. 2019;431(9):1878-1888. doi:10.1016/j.jmb.2019.03.005

44. Rajala RVS, Rajala A. Unlocking the role of lactate: metabolic pathways, signaling, and
gene regulation in postmitotic retinal cells. *Front Ophthalmol*. 2023;3.
doi:10.3389/fopht.2023.1296624

45. Williams PA, Harder JM, John SWM. Glaucoma as a metabolic optic neuropathy:
making the case for nicotinamide treatment in glaucoma. *J Glaucoma*. 2017;26(12):1161-
1168. doi:10.1097/IJG.0000000000000767

46. Inman DM, Harun-Or-Rashid M. Metabolic Vulnerability in the Neurodegenerative
Disease Glaucoma. *Front Neurosci*. 2017;11. doi:10.3389/fnins.2017.00146

47. Williams PA, Harder JM, Foxworth NE, et al. Vitamin B3 modulates mitochondrial
vulnerability and prevents glaucoma in aged mice. *Science*. 2017;355(6326):756-760.
doi:10.1126/science.aal0092

48. Chen S da, Wang L, Zhang X lan. Neuroprotection in glaucoma: present and future. *Chin*
Med J (Engl). 2013;126(8):1567-1577.

49. Vohra R, Dalgaard LM, Vibæk J, et al. Potential metabolic markers in glaucoma and their
regulation in response to hypoxia. *Acta Ophthalmol (Copenh)*. 2019;97(6):567-576.
doi:10.1111/aos.14021

50. Brooks GA. Lactate as a fulcrum of metabolism. *Redox Biol*. 2020;35:101454.
doi:10.1016/j.redox.2020.101454

51. Arnold PK, Finley LWS. Regulation and function of the mammalian tricarboxylic
acid cycle. *J Biol Chem*. 2022;299(2):102838. doi:10.1016/j.jbc.2022.102838

52. Jiménez-Moreno N, Kollareddy M, Stathakos P, et al. ATG8-dependent LMX1B-
autophagy crosstalk shapes human midbrain dopaminergic neuronal resilience. *J Cell Biol*.
2023;222(5):e201910133. doi:10.1083/jcb.201910133

53. Kim SW, Lee HK, Shin JH, Lee JK. Up-down Regulation of HO-1 and iNOS Gene
Expressions by Ethyl Pyruvate via Recruiting p300 to Nrf2 and Depriving It from p65. *Free*
Radic Biol Med. 2013;65:468-476. doi:10.1016/j.freeradbiomed.2013.07.028

54. Shin JH, Kim SW, Jin Y, Kim ID, Lee JK. Ethyl pyruvate-mediated Nrf2 activation and
hemeoxygenase 1 induction in astrocytes confer protective effects via autocrine and paracrine
mechanisms. *Neurochem Int*. 2012;61(1):89-99. doi:10.1016/j.neuint.2012.04.005

55. Harder JM, Guymer C, Wood JPM, et al. Disturbed glucose and pyruvate metabolism in
glaucoma with neuroprotection by pyruvate or rapamycin. *Proc Natl Acad Sci*.
2020;117(52):33619-33627. doi:10.1073/pnas.2014213117

56. Guymer C, Chidlow G, Wood JPM, Casson RJ. Oral pyruvate supplementation protects
against neurodegeneration in a rat model of glaucoma. *Invest Ophthalmol Vis Sci*.
2018;59(9):2618.

57. De Moraes CG, John SWM, Williams PA, Blumberg DM, Cioffi GA, Liebmann JM.
Nicotinamide and Pyruvate for Neuroenhancement in Open-Angle Glaucoma. *JAMA*
Ophthalmol. 2022;140(1):11-18. doi:10.1001/jamaophthalmol.2021.4576

58.   Wu X, Dai H, Liu L, et al. Citrate reduced oxidative damage in stem cells by regulating
cellular redox signaling pathways and represent a potential treatment for oxidative stress-
induced diseases. *Redox Biol*. 2019;21:101057. doi:10.1016/j.redox.2018.11.015

59.   Morganti C, Bonora M, Marchi S, et al. Citrate Mediates Crosstalk between
Mitochondria and the Nucleus to Promote Human Mesenchymal Stem Cell In Vitro
Osteogenesis. *Cells*. 2020;9(4):1034. doi:10.3390/cells9041034

60.   Catalina-Rodriguez O, Kolukula VK, Tomita Y, et al. The mitochondrial citrate
transporter, CIC, is essential for mitochondrial homeostasis. *Oncotarget*. 2012;3(10):1220-
1235. doi:10.18632/oncotarget.714

61.   Ohanele C, Peoples JN, Karlstaedt A, et al. The mitochondrial citrate carrier SLC25A1
regulates metabolic reprogramming and morphogenesis in the developing heart. *Commun
Biol*. 2024;7(1):1-16. doi:10.1038/s42003-024-07110-8

62.   Chhimpa N, Singh N, Puri N, Kayath HP. The Novel Role of Mitochondrial Citrate
Synthase and Citrate in the Pathophysiology of Alzheimer's Disease. *J Alzheimers Dis*.
94(Suppl 1):S453-S472. doi:10.3233/JAD-220514

63.   Fraenkl SA, Muser J, Groell R, et al. Plasma Citrate Levels as a Potential Biomarker for
Glaucoma. *J Ocul Pharmacol Ther*. 2011;27(6):577-580. doi:10.1089/jop.2011.0062

64.   Michalczuk M, Tadeusz P, Urban B, Anna W, Łazarczyk AB. Plasma citrate
concentration: a possible biomarker for glaucoma in children. *BMJ Paediatr Open*.
2017;1(1):e000023. doi:10.1136/bmjpo-2017-000023

65.   Nusinovici S, Li H, Thakur S, et al. High-Density Lipoprotein 3 Cholesterol and Primary
Open-Angle Glaucoma: Metabolomics and Mendelian Randomization Analyses.
Ophthalmology. 2022;129(3):285-294. doi:10.1016/j.ophtha.2021.09.013

66.   Masson EAY, Serrano J, Leger-Charnay E, Acar N. Cholesterol and oxysterols in retinal
neuron-glia interactions: relevance for glaucoma. *Front Ophthalmol*. 2024;3:1303649.
doi:10.3389/fopht.2023.1303649

67.   REvisiting Lipids in REtinal Diseases: A Focused Review on Age-related Macular
Degeneration and Diabetic Retinopathy - PMC. Accessed August 3, 2024.
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7521975/

68.   Increased High-Density Lipoprotein Levels Associated with Age-Related Macular
Degeneration: Evidence from the EYE-RISK and European Eye Epidemiology Consortia -
ScienceDirect. Accessed August 3, 2024.
https://www.sciencedirect.com/science/article/pii/S0161642018310911

69.   Pasquale LR, Khawaja AP, Wiggs JL, et al. Metabolite and Lipid Biomarkers Associated
With Intraocular Pressure and Inner Retinal Morphology: [1] H NMR Spectroscopy Results
From the UK Biobank. *Investig Opthalmology Vis Sci*. 2023;64(11):11.
doi:10.1167/iovs.64.11.11

70.   Ana R da, Gliszczyńska A, Sanchez-Lopez E, et al. Precision Medicines for Retinal Lipid
Metabolism-Related Pathologies. *J Pers Med*. 2023;13(4):635. doi:10.3390/jpm13040635

71.   Fu Z, Chen CT, Cagnone G, et al. Dyslipidemia in retinal metabolic disorders. *EMBO*
Mol Med. 2019;11(10):e10473. doi:10.15252/emmm.201910473

72.   Yuksel S, Ufret-Vincenty R, Butovich IA. The role of Lipe in lipid metabolism in the
retina. *Invest Ophthalmol Vis Sci*. 2023;64(8):5454.

73.   Kelly UL, Grigsby D, Cady MA, et al. High-density lipoproteins are a potential
therapeutic target for age-related macular degeneration. *J Biol Chem*. 2020;295(39):13601-
13616. doi:10.1074/jbc.RA119.012305

74.   Morvaridzadeh M, Zoubdane N, Heshmati J, Alami M, Berrougui H, Khalil A. High-
Density Lipoprotein Metabolism and Function in Cardiovascular Diseases: What about Aging
and Diet Effects? *Nutrients*. 2024;16(5):653. doi:10.3390/nu16050653

75.   Hamel AR, Yan W, Rouhana JM, et al. Integrating genetic regulation and single-cell
expression with GWAS prioritizes causal genes and cell types for glaucoma. *Nat Commun*.
2024;15:396. doi:10.1038/s41467-023-44380-y

76.   Sudlow C, Gallacher J, Allen N, et al. UK Biobank: An Open Access Resource for
Identifying the Causes of a Wide Range of Complex Diseases of Middle and Old Age. *PLoS*
Med. 2015;12(3):e1001779. doi:10.1371/journal.pmed.1001779

77.   Bycroft C, Freeman C, Petkova D, et al. The UK Biobank resource with deep
phenotyping and genomic data. *Nature*. 2018;562(7726):203-209. doi:10.1038/s41586-018-
0579-z

78.   Elliott P, Peakman TC, on behalf of UK Biobank. The UK Biobank sample handling and
storage protocol for the collection, processing and archiving of human blood and urine. *Int J*
Epidemiol. 2008;37(2):234-244. doi:10.1093/ije/dym276

79.   Zeleznik O, Kang JH, Lasky-Su J, et al. Plasma metabolomics of primary open-angle
glaucoma in three prospective US cohorts and the UK Biobank. Published online February 28,
2022. doi:10.1101/2022.02.24.22271483

80.   Julkunen H, Cichońska A, Slagboom PE, Würtz P, Nightingale Health UK Biobank
Initiative. Metabolic biomarker profiling for identification of susceptibility to severe
pneumonia and COVID-19 in the general population. Janus ED, Barton M, Karunajeewa H,
Janus ED, eds. *eLife*. 2021;10:e63033. doi:10.7554/eLife.63033

81.   Takis PG, Ghini V, Tenori L, Turano P, Luchinat C. Uniqueness of the NMR approach to
metabolomics. *TrAC Trends Anal Chem*. 2019;120:115300. doi:10.1016/j.trac.2018.10.036

82.   Soininen P, Kangas AJ, Würtz P, Suna T, Ala-Korpela M. Quantitative serum nuclear
magnetic resonance metabolomics in cardiovascular epidemiology and genetics. *Circ*
Cardiovasc Genet. 2015;8(1):192-206. doi:10.1161/CIRCGENETICS.114.000216

83.   Mihaleva VV, Korhonen SP, van Duynhoven J, Niemitz M, Vervoort J, Jacobs DM.
Automated quantum mechanical total line shape fitting model for quantitative NMR-based
profiling of human serum metabolites. *Anal Bioanal Chem*. 2014;406(13):3091-3102.
doi:10.1007/s00216-014-7752-5

84.   Kim J, Aschard H, Kang JH, et al. Intraocular Pressure, Glaucoma, and Dietary Caffeine
Consumption: A Gene—Diet Interaction Study from the UK Biobank. *Ophthalmology*.
2020;128(6):866. doi:10.1016/j.ophtha.2020.12.009

85.   Gao X, Starmer J, Martin ER. A multiple testing correction method for genetic
association studies using correlated single nucleotide polymorphisms. *Genet Epidemiol*.
2008;32(4):361-369. doi:10.1002/gepi.20310

86.   Thaung C, West K, Clark BJ, et al. Novel ENU-induced eye mutations in the mouse:
models for human eye disease. *Hum Mol Genet*. 2002;11(7):755-767.
doi:10.1093/hmg/11.7.755

87.   John SW, Hagaman JR, MacTaggart TE, Peng L, Smithes O. Intraocular pressure in
inbred mouse strains. *Invest Ophthalmol Vis Sci*. 1997;38(1):249-253.

88.   Savinova OV, Sugiyama F, Martin JE, et al. Intraocular pressure in genetically distinct
mice: an update and strain survey. *BMC Genet*. 2001;2:12. doi:10.1186/1471-2156-2-12

89.   Howell GR, Libby RT, Jakobs TC, et al. Axons of retinal ganglion cells are insulted in
the optic nerve early in DBA/2J glaucoma. *J Cell Biol*. 2007;179(7):1523-1537.
doi:10.1083/jcb.200706181

90.   Nair KS, Cosma M, Raghupathy N, et al. YBR/EiJ mice: a new model of glaucoma
caused by genes on chromosomes 4 and 17. *Dis Model Mech*. 2016;9(8):863-871.
doi:10.1242/dmm.024307

91.   Howell GR, Soto I, Zhu X, et al. Radiation treatment inhibits monocyte entry into the
optic nerve head and prevents neuronal damage in a mouse model of glaucoma. *J Clin Invest*.
2012;122(4):1246-1261. doi:10.1172/JCI61135

92.   Howell GR, Macalinao DG, Sousa GL, et al. Molecular clustering identifies complement
and endothelin induction as early events in a mouse model of glaucoma. *J Clin Invest*.
2011;121(4):1429-1444. doi:10.1172/JCI44646

Figures

899
900
901   **Figure 1. Participant flow chart describing inclusion and exclusion criteria from the UK**
902   **Biobank**
903
904

Figure 2. Study design from the UK Biobank. (a) 117,698 individuals had metabolomics data
available from the UK Biobank, which was divided into a training and test set to formulate a
metabolic risk score (MRS) model. (b) The inclusion of metabolites (either 168 metabolites on
the NMR platform or a subset of 27 metabolites with European Union (EU) certification) in
relation to prevalent glaucoma risk prediction was studied. (c) A histogram showing the
polygenic risk score (PRS) distribution is shown. Overall, 4,658 cases and 113,040 individuals
without glaucoma are available for analysis. The metabolomic signature of resilience to the top
10% of glaucoma PRS was assessed among 1,693 cases (14.4%) and 10,077 individuals without
glaucoma (85.6%). (d) Interactions of prevalent glaucoma with MRS and PRS quartiles were
examined.

Figure 3. Inclusion of metabolite data into glaucoma prediction algorithms. Model 1
includes metabolites only; Model 2 incorporates additional covariates including age, sex, genetic
ancestry, season, time of day of specimen collection, and fasting time; Model 3 incorporates
covariates in Model 2 and smoking status (never, past, and current smoker), alcohol intake
(g/week), caffeine intake (mg/day), physical activity (metabolic equivalent of task [MET],
hours/week), body mass index (kg/m^2), average systolic blood pressure (mm Hg), history of
diabetes, HbA1c (mmol/mol), history of coronary artery disease, systemic beta-blocker use, oral
steroid use, and spherical equivalent refractive error (diopters); Model 4 incorporates covariates
in model 3 and a glaucoma polygenic risk score (PRS). Each color represents a different panel of
metabolites (grey = no metabolites; light blue = 27 metabolites; and dark blue = 168
metabolites). The white text represents the AUC ± 95% confidence interval.
Abbreviations: ROC, receiver operator curve; AUC, area under the curve; EU, European Union.

Figure 4. The distribution of glaucoma cases and no glaucoma stratified by polygenic risk score (PRS) deciles. Participants were divided into the bottom 50% and top 10% based on their glaucoma PRS, where the prevalence of glaucoma cases from (A) the bottom 50% (n=58,358) was 1.3% and from the top 10% (n=11,770) was 14.4%. (B) Box plot illustrating the distribution of participants with glaucoma (top) and no glaucoma (bottom) as a function of PRS decile. The blue line denotes participants at the bottom 50% of glaucoma PRS, the red line highlights the participants at the top decile of glaucoma PRS, and the red box represents the participants resilient to glaucoma despite high PRS.

A.

Quartiles				
	Q1	**Q2**	**Q3**	**Q4**
PRS	-0.0090, -0.0075	-0.0075, -0.0073	-0.0073, -0.0070	-0.0070, -0.0055
Metabolite sum	-8.2, -1.5	-1.5, -0.2	-0.2, 1.3	1.3, 32.2

B.

	Glaucoma cases	Glaucoma Metabolite Reference	Odds Ratio (95% CI)	p-value
PRS Q1				
Metabolite sum Q2	66	55	1.18 (0.82, 1.69)	0.42
Metabolite sum Q3	61	55	1.1 (0.77, 1.60)	0.66
Metabolite sum Q4	57	55	1.0 (0.69, 1.46)	1.00
PRS Q2				
Metabolite sum Q2	138	161	0.87 (0.69, 1.10)	0.27
Metabolite sum Q3	128	161	0.79 (0.63, 1.00)	0.057
Metabolite sum Q4	127	161	0.80 (0.63, 1.01)	0.069
PRS Q3				
Metabolite sum Q2	266	297	0.90 (0.76, 1.06)	0.23
Metabolite sum Q3	228	297	0.78 (0.65, 0.93)	0.0059
Metabolite sum Q4	219	297	0.74 (0.62, 0.88)	0.0009
PRS Q4				
Metabolite sum Q2	750	793	0.93 (0.84, 1.03)	0.19
Metabolite sum Q3	733	793	0.90 (0.81, 1.00)	0.054
Metabolite sum Q4	579	793	0.71 (0.64, 0.80)	<0.0001

Figure 5. Interaction of three putative resilient metabolites (lactate, pyruvate, and citrate) and polygenic risk score (PRS) on glaucoma risk. (A) The bar chart shows the interaction of resilient probit-transformed metabolite sum with glaucoma genetic predisposition in each PRS quartile. In each glaucoma PRS quartile, the lowest metabolic sum quartile (Q1) is the metabolite reference group used to calculate the odds ratios. Each color represents resilient-metabolite sum quartiles (red = second quartile; blue = third quartile; and green = fourth quartile). Error bars show 95% confidence interval (CI). The table under the bar chart shows the ranges for the PRS and metabolite sum value quartiles. (B) The table shows odds ratios for glaucoma by PRS and putative resilient metabolite sum within various quartiles. The number of glaucoma cases within each resilient metabolite sum quartile and the number of glaucoma cases in the first quartile of resilient metabolite sum (Q1, labeled as glaucoma metabolite reference) are used to calculate the odd ratios. This analysis is adjusted for time since the last meal/drink (hours), age, age-squared, sex, ethnicity (Asian, Black, White, and other), season, time of day of specimen collection (morning, afternoon, night), smoking status (never, past, and current smoker), alcohol intake, caffeine intake, physical activity (metabolic equivalent of task [MET] hours/week), body mass index (kg/m^2), average systolic blood pressure (mm Hg), history of diabetes (yes or no), HbA1c (mmol/mol), history of coronary artery disease, systemic beta-blocker use, oral steroid use, and spherical equivalent refractive error (diopters).

A.

P-Interaction= 0.019
Sample size= 117,697

B.

	Glaucoma cases	Odds Ratio (95% CI)	p-value
PRS Q1			
MRS Q1	46	reference	1
MRS Q2	56	1.20 (0.81, 1.79)	0.40
MRS Q3	64	1.39 (0.95, 2.04)	0.11
MRS Q4	73	1.66 (1.15, 2.42)	0.0086
PRS Q2			
MRS Q1	94	2.06 (1.45, 2.96)	0.0001
MRS Q2	119	2.62 (1.87, 3.72)	<0.0001
MRS Q3	149	3.32 (2.40, 4.67)	<0.0001
MRS Q4	192	4.29 (3.12, 5.98)	<0.0001
PRS Q3			
MRS Q1	171	3.79 (2.75, 5.30)	<0.0001
MRS Q2	220	4.98 (3.65, 6.92)	<0.0001
MRS Q3	254	5.70 (4.19, 7.90)	<0.0001
MRS Q4	365	8.33 (6.17, 11.44)	<0.0001
PRS Q4			
MRS Q1	495	11.71 (8.72, 16.00)	<0.0001
MRS Q2	582	13.90 (10.36, 16.03)	<0.0001
MRS Q3	747	18.27 (13.66, 24.89)	<0.0001
MRS Q4	1031	25.08 (18.80, 34.12)	<0.0001

	Quartiles			
	Q1	Q2	Q3	Q4
PRS	-0.0090, -0.0075	-0.0075, -0.0073	-0.0073, -0.0070	-0.0070, -0.0055
MRS	0.0000, 0.0316	0.0316, 0.0384	0.0384, 0.0462	0.0462, 0.5185

Figure 6. Interaction of the holistic metabolite risk score (n=168 metabolites) and polygenic risk score (PRS) on glaucoma risk. (A) The bar chart plots the odds ratio of glaucoma as a function of holistic probit-transformed MRS quartile with further stratification by glaucoma PRS in each MRS bin. The lowest quartile of glaucoma PRS and MRS is the reference group (see dotted red line) for the entire population. Each color represents the MRS quartiles (red = first quartile; blue = second quartile; green = third quartile; and purple = fourth quartile). Error bars show 95% confidence interval (CI). The table under the bar chart shows the ranges for the PRS and MRS quartiles. (B) Table showing odds ratios for glaucoma by polygenic risk score (PRS) and MRS within various quartiles. The number of glaucoma cases within each MRS and the number of glaucoma cases in PRS Q1 and MRS Q1 are used to calculate the odds ratios. This analysis is adjusted for time since the last meal/drink (hours), age, age-squared, sex, ethnicity (Asian, Black, White, and other), season, time of day of specimen collection (morning, afternoon, night), smoking status (never, past, and current smoker), alcohol intake, caffeine intake, physical activity (metabolic equivalent of task [MET] hours/week), body mass index (kg/m^2), average systolic blood pressure (mm Hg), history of diabetes (yes or no), HbA1c (mmol/mol), history of coronary artery disease, systemic beta-blocker use, oral steroid use, and spherical equivalent refractive error (diopters).

977
Figure 7. Pyruvate treatment protects from IOP elevation and glaucoma. (A) Representative
photos of eyes from mice of the indicated genotypes and treatments (Unt = untreated, Pyr =
pyruvate treated). *Lmx1b* is expressed in the iris and cornea and so *Lmx1b*^{V265D} mutant eyes have
primary abnormalities of the iris and cornea. This includes corneal haze, which is present prior to
IOP elevation in many eyes and likely reflects a direct transcriptional role of LMX1B in collagen
gene expression. *Lmx1b*^{V265D} mutant eyes also develop anterior chamber deepening (ACD), a
sensitive indicator of IOP elevation in mice. The WT and pyruvate-treated mutant eyes have
shallow anterior chambers, while the untreated mutant eye has a deepened chamber
(arrowheads). (B) Distributions of ACD are based on a previously defined scoring system.[30]
Groups are compared by Fisher's exact test. n > 30 eyes were examined in each group. (C)
Boxplots of IOP (interquartile range and median line) in WT and mutant eyes. Pyruvate
treatment significantly lessens IOP elevation in mutants compared to untreated mutant controls.
Groups were compared by ANOVA followed by Tukey's honestly significant difference. n > 30
eyes were examined in each *Lmx1b*^{V265D} mutant group, and n > 20 eyes were examined in WT
groups. (D) Distributions of damage based on analysis of PPD-stained optic nerve cross sections
from 5.5-month-old mice (Methods). Pyruvate treatment lessened the incidence of glaucoma
(Fisher's exact test). No glaucoma was found in WT mice. Geno = genotype. n > 40 nerves
examined per group. NOE = no glaucoma, MOD = moderate. SEV = severe, and V. SEV = very
severe (see Methods).
997

998   Tables

999   **Table 1. Demographic and clinical characteristics of the UK Biobank study population**
1000 **assessed in 2006-2010**

1001

Characteristic	No Glaucoma	Glaucoma	P-value
Sample Size (%)	113,040 (96.0)	4,658 (4.0)	
Sex - Male (%)	52,497 (46.4)	2,493 (53.5)	<0.001
Age at recruitment (mean (SD))	56.7 (8.0)	60.9 (6.6)	<0.001
Ethnicity (%)			0.0010
White	105,912 (94)	4,331 (93)	
Asian	2,977 (2.6)	127 (2.7)	
Black	2,228 (2.0)	130 (2.8)	
Other	1,923 (1.7)	70 (1.5)	
Genetic Ancestry (%)			<0.001
African	1,911 (1.7)	123 (2.7)	
AMR	261 (0.2)	2 (0.0)	
Asian	3,125 (2.8)	134 (2.9)	
European	96,991 (86.6)	4,001 (86.7)	
Smoking Status (%)			<0.001
Never	63,240 (56)	2,339 (52)	
Prefer not to answer	430 (0.4)	16 (0.4)	
Previous	38,926 (34)	1,732 (39)	
Current	10,605 (9.4)	410 (9.1)	
Total Cholesterol, mmol/l (median [IQR])	4.6 [3.98, 5.22]	4.5 [3.88, 5.18]	<0.001
Physical Activity, MET-minutes per week (mean (SD))	2,465 (2430)	2,458 (2,438)	0.84
Body Mass Index, kg/m^2 (mean (SD))	27.4 (4.8)	27.7 (4.7)	<0.001
HbA1c, mmol/mol (mean (SD))	36.0 (5.8)	37.4 (6.8)	<0.001
Spherical Equivalent, diopter (mean (SD))	-0.1 (2.1)	-0.1 (1.9)	0.67
Intraocular pressure, mmHg (mean (SD))	15.9 (2.6)	17.8 (4.5)	<0.001
mRNFL thickness, μm (mean (SD))	28.7 (1.8)	28.5 (1.3)	<0.001
Beta blocker use (%)	8,398 (7.4)	447 (9.6)	<0.001
Caffeine intake, mg/day (mean (SD))	165 (67)	168 (59)	0.0020
Alcohol intake, g/week (median [IQR])	84 [40.3, 146.9]	84 [48.0, 154.6]	<0.001
Diabetes (%)	6,512 (5.8)	463 (9.9)	<0.001
Oral steroid use (%)	3,043 (2.7)	184 (4.0)	<0.001
Coronary Artery Disease (%)	5,153 (4.6)	377 (8.1)	<0.001

1002

1003 Abbreviations: SD, standard deviation; IQR, interquartile range; AMR, mixed American; MET
1004 metabolic equivalent of task mRNFL, macular retinal nerve fiber layer

Table 2. Stratification of glaucoma by ethnicity, age, and gender for predictive assessment with and without using metabolite data.

Stratification	n	Glaucoma cases	No Metabolites (AUC)	Metabolites (AUC)	P-value
Ethnicity					
White	110,243	4,331	0.675	0.686	<0.001
Asian	3,104	127	0.780	0.768	0.062
Black	2,358	130	0.704	0.706	0.52
Age					
<55 years	43,648	788	0.576	0.566	0.52
≥55 years	74,050	3,870	0.569	0.596	0.002
Gender					
Female	62,708	2,165	0.689	0.696	0.25
Male	54,990	2,493	0.659	0.673	0.002

The area under the curve (AUC) of the receiver operating characteristic (ROC) curve was calculated for each demographic stratification to evaluate the predictive performance of models both with and without metabolite data. Differences in model AUC were tested using the DeLong test, and p-values were reported. For models excluding metabolite data, the predictors include: age, sex, genetic ancestry, season, time of day of specimen collection, fasting time (hours), smoking status (never, past, and current smoker), alcohol intake (g/week), caffeine intake (mg/day), physical activity (metabolic equivalent of task [MET], hours/week), body mass index (kg/m^2), average systolic blood pressure (mm Hg), history of diabetes, HbA1c (mmol/mol), history of coronary artery disease, systemic beta-blocker use, oral steroid use, and spherical equivalent refractive error (diopters). Models including metabolite data incorporated the same predictors with the addition of the 168 metabolite measurements.

Table 3. Demographic and clinical characteristics in 2006-2010 of participants among the
top 10% of glaucoma polygenic risk score (PRS).

Characteristic	No Glaucoma	Glaucoma	P-value
Sample size (%)	10,077 (85.6)	1,693 (14.4)	
Sex - Male (%)	4,667 (46.3)	893 (52.7)	<0.001
Age at recruitment (mean (SD))	56.3 (8.0)	61.0 (6.3)	<0.001
Ethnicity (%)			<0.001
White	9,256 (91.9)	1,636 (96.6)	
Asian	279 (2.8)	24 (1.4)	
Black	345 (3.4)	20 (1.2)	
Other	197 (2.0)	13 (0.8)	
Smoking Status (%)			0.0030
Never	5,806 (57.6)	899 (53.1)	
Prefer not to answer	48 (0.5)	5 (0.3)	
Previous	3,402 (33.8)	640 (37.8)	
Current	821 (8.1)	149 (8.8)	
Physical Activity, MET-minutes per week (mean (SD))	2,439 (2,352)	2,514 (2,480)	0.23
Body mass index, kg/m^2 (mean (SD))	27.4 (4.7)	27.7 (4.8)	0.0080
HbA1c, mmol/mol (mean (SD))	36.00 (5.9)	37.0 (6.1)	<0.001
Spherical Equivalent, diopter (mean (SD))	-0.3 (2.2)	-0.2 (1.9)	0.0090
Intraocular pressure, mmHg (mean (SD))	17.2 (3.0)	18.3 (5.0)	<0.001
mRNFL thickness, μm (mean (SD))	28.7 (1.8)	28.5 (1.2)	<0.001
Beta blocker use (%)	719 (7.1)	179 (10.6)	<0.001
Caffeine intake, mg/day (mean (SD))	164 (65)	170 (60)	0.0010
Alcohol intake, g/week (median [IQR])	84 [38.6, 145.3]	84 [51.5, 155.3]	0.002
Diabetes mellitus (%)	607 (6.0)	151 (8.9)	<0.001
Oral steroid use (%)	292 (2.9)	59 (3.5)	0.22
Coronary Artery Disease (%)	443 (4.4)	148 (8.7)	<0.001
Total cholesterol, mmol/l (median [IQR])	4.61 [4.00, 5.25]	4.54 [3.90, 5.21]	0.006
Lactate, mmol/l (median [IQR])	3.96 [3.27, 4.78]	3.79 [3.16, 4.58]	<0.001
Pyruvate, mmol/l (median [IQR])	0.080 [0.06, 0.10]	0.077 [0.06, 0.09]	<0.001
Citrate, mmol/l (median [IQR])	0.065 [0.06, 0.07]	0.065 [0.06, 0.07]	0.87
Cholesteryl Esters in Small HDL, mmol/l (median [IQR])	0.33 [0.30, 0.36]	0.33 [0.30, 0.36]	0.005
Triglycerides in Very Large VLDL, mmol/l (median [IQR])	0.091 [0.05, 0.15]	0.10 [0.05, 0.17]	<0.001
Alanine, mmol/l (median [IQR])	0.29 [0.24, 0.35]	0.29 [0.24, 0.35]	0.58
Triglycerides in Chylomicrons and extremely Large VLDL, mmol/l (median [IQR])	0.088 [0.03, 0.18]	0.10 [0.04, 0.21]	<0.001
Acetoacetate, mmol/l (median [IQR])	0.010 [0.01, 0.02]	0.011 [0.01, 0.02]	<0.001
Cholesteryl Esters in Medium HDL, mmol/l (median [IQR])	0.41 [0.34, 0.48]	0.40 [0.34, 0.47]	0.003
Triglycerides in Large VLDL, mmol/l (median [IQR])	0.15 [0.09, 0.22]	0.16 [0.10, 0.23]	<0.001
Cholesterol in Medium HDL, mmol/l (median [IQR])	0.49 [0.42, 0.58]	0.48 [0.41, 0.57]	0.005

1041   Abbreviations: SD, standard deviation; IQR, interquartile range; MET, metabolic equivalent of
1042   task HbA1c, hemoglobin A1C; mRNFL, macular retinal nerve fiber layer; HDL, high-density
1043   lipoprotein; VLDL, very low-density lipoprotein
1044

Table 4. Demographic and clinical characteristics in 2006-2010 of participants among the bottom 50% of glaucoma polygenic risk score (PRS).

Characteristic	No Glaucoma	Glaucoma	P-value
Sample size (%)	57,578 (98.7)	780 (1.3)	
Sex - Male (%)	26,887 (46.7)	409 (52.4)	0.002
Age at recruitment (mean (SD))	57 (8.0)	60 (7.0)	<0.001
Ethnicity (%)			<0.001
White	55,034 (96)	690 (89)	
Asian	1,129 (2.0)	41 (5.3)	
Black	574 (1.0)	27 (3.5)	
Other	841 (1.5)	22 (2.8)	
Smoking Status (%)			0.67
Never	200 (0.3)	4 (0.5)	
Prefer not to answer	31,593 (54.9)	418 (53.6)	
Previous	20,276 (35.2)	287 (36.8)	
Current	5,509 (9.6)	71 (9.1)	
MET, minutes per week (mean (SD))	2,483 (2,450)	2,612 (2,562)	0.14
Body mass index kg/m^2 (mean (SD))	27.4 (4.8)	27.8 (4.7)	0.021
HbA1c, mmol/mol (mean (SD))	35.9 (5.7)	38.1 (7.8)	<0.001
Spherical Equivalent, diopter (mean (SD))	-0.03 (2.1)	0.01 (1.9)	0.63
Intraocular pressure, mmHg (mean (SD))	15.4 (2.4)	17.1 (3.9)	<0.001
mRNFL thickness, μm (mean (SD))	28.7 (1.7)	28.6 (1.3)	0.02
Beta blocker use (%)	4,322 (7.5)	71 (9.1)	0.11
Caffeine intake, mg/day (mean (SD))	1,66.3 (67.4)	166.6 (61.5)	0.90
Alcohol intake, g/week (median [IQR])	84 [41.7, 149.3]	84 [42.5, 145.9]	0.91
Diabetes (%)	3209 (5.6)	104 (13.3)	<0.001
Oral steroid use (%)	1,538 (2.7)	39 (5.0)	<0.001
Coronary Artery Disease (%)	2,656 (4.6)	64 (8.2)	<0.001
Total cholesterol, mmol/l (median [IQR])	4.59 [3.98, 5.22]	4.49 [3.76, 5.13]	0.001
Lactate, mmol/l (median [IQR])	3.95 [3.24, 4.75]	3.84 [3.21, 4.69]	0.16
Concentration of Small HDL Particles, mmol/l (median [IQR])	0.0097 [0.0089, 0.011]	0.0095 [0.0088, 0.010]	0.002
Cholesteryl esters in small HDL, mmol/l (median [IQR])	0.33 [0.30, 0.36]	0.32 [0.30, 0.36]	0.001
Albumin, mmol/l (median [IQR])	39.4 [37.3, 41.45]	38.9 [36.7, 40.8]	<0.001
Total lipids in small HDL, mmol/l (median [IQR])	1.16 [1.06, 1.26]	1.15 [1.05, 1.25]	0.018
Citrate, mmol/l (median [IQR])	0.065 [0.057, 0.074]	0.065 [0.057, 0.073]	0.52
Pyruvate, mmol/l (median [IQR])	0.080 [0.06, 0.10]	0.079 [0.061, 0.098]	0.14
Alanine, mmol/l (median [IQR])	0.29 [0.24, 0.35]	0.29 [0.24, 0.35]	0.95
Phospholipids in Small HDL, mmol/l (median [IQR])	0.66 [0.60, 0.72]	0.66 [0.60, 0.71]	0.033

Abbreviations: SD, standard deviation; MET, metabolic equivalents; HbA1c, hemoglobin A1C; mRNFL, macula region retinal nerve fiber layer; HDL, high density lipoprotein; VLDL, very low density lipoprotein

1053 **Table 5. Metabolites associated with glaucoma among participants in the top decile and the**
1054 **bottom half of glaucoma polygenic risk score.**
1055

Top 10% of glaucoma polygenic risk score			
Metabolites (Probit score)	Glaucoma	No Glaucoma	Adjusted P-value (NEF)
Lactate	-0.146	0.0239	8.8E-12
Pyruvate	-0.137	0.0117	2.9E-10
Citrate	-0.0693	0.0079	0.018
Triglycerides in Very Large VLDL	0.0606	-0.0043	0.10
Triglycerides in Chylomicrons and Extremely Large VLDL	0.0572	-0.0076	0.11
Acetoacetate	0.0744	0.0011	0.11
Cholesteryl Esters in Small HDL	-0.0239	0.0376	0.12
Cholesteryl Esters in Medium HDL	-0.0261	0.0276	0.13
Alanine	-0.0612	-0.0006	0.14
Triglycerides in Large VLDL	0.0629	0.0005	0.15

1056
1057

Bottom 50% of glaucoma polygenic risk score			
Metabolites (Probit score)	Glaucoma	No Glaucoma	Adjusted P-value (NEF)
Albumin	-0.110	-0.0099	0.047
Concentration of Small HDL Particles	-0.0943	-0.0062	0.071
Cholesteryl Esters in Small HDL	-0.0925	-0.0055	0.085
Cholesterol in Small HDL	-0.0919	-0.0063	0.088
Omega-3 Fatty Acids	-0.0896	-0.0052	0.092
Docosahexaenoic Acid	-0.0876	-0.0061	0.10
Total Lipids in Small HDL	-0.0844	-0.0051	0.12
Lactate	-0.0823	0.0031	0.12
Citrate	-0.0882	-0.0023	0.16
Alanine	-0.0766	0.0011	0.18

Potential confounders adjusted by regression include time since the last meal/drink (hours), age,
age-squared, sex, ethnicity (Asian, Black, White, and other), season, time of day of specimen
collection (morning, afternoon, night), smoking status (never, past, and current smoker), alcohol
intake, caffeine intake, physical activity (metabolic equivalent of task [MET] hours/week), body
mass index (kg/m^2), average systolic blood pressure (mm Hg), history of diabetes (yes or no),
HbA1c (mmol/mol), history of coronary artery disease, systemic beta-blocker use, oral steroid
use, and spherical equivalent refractive error (diopters).
1065

Financial Support:

This work is supported by NEI R01 015473 (LRP), NEI R01 032559 (LRP, JLW, AVS),

EY036460 (LRP, JHK), NEI P30 EY014104 (JLW, AVS), NEI R01EY031424 (AVS), NIGMS

R35-GM124836 (RD), NEI EY032507, EY032062, and EY018606 (SWMJ), an unrestricted

Challenge Grant from Research to Prevent Blindness (NYC), and The Glaucoma Foundation

(NYC). Further funding was provided by startup funds at Columbia University including the

Precision Medicine Initiative, and by the New York Fund for Innovation in Research and

Scientific Talent (NYFIRST; EMPIRE CU19-2660; SWMJ). This work is also supported by K23

(1K23EY032634) and the Research to Prevent Blindness Career Development Award (N.Z), a

Vision Core grant P30EY019007 (Columbia University) an unrestricted departmental award

from Research to Prevent Blindness (Columbia University). Supported by Fight for Sight (UK)

(1956A), and The Desmond Foundation (K.V.S.); UK Research and Innovation Future Leaders

Fellowship (MR/T040912/1), Alcon Research Institute Young Investigator Award and a Lister

Institute of Preventative Medicine Fellowship (A.P.K.); financial support from the UK

Department of Health through an award made by the National Institute for Health Research

(NIHR) to Moorfields Eye Hospital National Health Service (NHS) Foundation Trust and

University College London (UCL) Institute of Ophthalmology for a Biomedical Research Centre

(BRC) for Ophthalmology (A.P.K.). The sponsor or funding organizations had no role in the

design or conduct of this research. The content is solely the responsibility of the authors and does

not necessarily represent the official views of the National Institutes of Health.

1086

Conflict of Interest:

The authors declare the following financial interests/personal relationships that may be

considered potential competing interests: LRP reports consulting honoraria from Twenty

Twenty. JLW reports consulting fees from Editas and CRISPR Therapeutics. NZ reports

consulting fees from Character Biosciences. APK has acted as a paid consultant or lecturer to

Abbvie, Aerie, Allergan, Google Health, Heidelberg Engineering, Novartis, Reichert, Santen,

Thea and Topcon. NV has acted as a paid consultant or lecturer to Santen, Thea, NovaEye,

Glaukos, and Elios. RD reports being a scientific co-founder, consultant, and equity holder for

Pensieve Health (pending) and being a consultant for Variant Bio and Character Bio. SWMJ is

founder of Myco Advising LLC, a founder of Qura Inc and advises Starlight Bio Inc with no

overlap to the currently presented research. SWMJ and NT are inventors on patents/patent

applications held by The Jackson Laboratory and Columbia University on the use of pyruvate to

treat glaucoma. No other disclosures were reported.