Pyruvate and Related Energetic Metabolites Modulate Resilience Against High Genetic Risk for Glaucoma

Keva Li[1,*], Nicholas Tolman[2,*], Ayellet V. Segrè[3,4], Kelsey V. Stuart[5], Oana A. Zeleznik[6], Neeru A. Vallabh[7,8], Kuang Hu[5], Nazlee Zebardast[3], Akiko Hanyuda[9,10], Yoshihiko Raita[11], Christa Montgomery[2], Chi Zhang[2], Pirro G. Hysi[12,13], Ron Do[14], Anthony P. Khawaja[5], Janey L. Wiggs[3,4], Jae H. Kang[6,‡], Simon WM John[2,15,‡], Louis R. Pasquale[1,‡] & UK Biobank Eye and Vision Consortium

[1]Department of Ophthalmology, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA

[2]Department of Ophthalmology, Vagelos College of Physicians and Surgeons, Columbia University Irving Medical Center, New York, NY 10032, USA

[3]Department of Ophthalmology, Massachusetts Eye and Ear, Harvard Medical School, Boston, MA 02142, USA

[4]Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA

[5]NIHR Biomedical Research Centre, Moorfields Eye Hospital NHS Foundation Trust, and University College London Institute of Ophthalmology, London, EC1V 2PD, United Kingdom

[6]Channing Division of Network Medicine, Department of Medicine, Harvard Medical School and Brigham and Women's Hospital, 181 Longwood Avenue, Boston, MA 02115, USA.

[7]Department of Eye and Vision Science, Institute of Life Course and Medical Sciences, University of Liverpool, Liverpool, L69 3BX, United Kingdom

[8]St. Paul's Eye Unit, Liverpool University Hospital NHS Foundation Trust, Liverpool, L7 8XP,

United Kingdom

[9]Department of Ophthalmology, Keio University School of Medicine, 35 Shinanomachi,

Shinjuku-ku, Tokyo 160-8582, Japan

[10]Epidemiology and Prevention Group, Center for Public Health Sciences, National Cancer

Center, Tokyo 104-0045, Japan

[11]Okinawa Kenritsu, Chubu Byoin, Uruma, Okinawa 904-2293, Japan

[12]Department of Ophthalmology, St Thomas' Hospital, King's College London, London, SE1

7EH, United Kingdom.

[13]Department of Twin Research & Genetic Epidemiology, St Thomas' Hospital, King's College

London, London, SE1 7EH, United Kingdom.

[14]Department of Genetics and Genomics Science, Icahn School of Medicine at Mount Sinai,

New York, NY 10029, USA.

[15]Zuckerman Mind Brain Behavior Institute, Columbia University, New York, NY 10027, USA

[*]Co-first authors

[‡]Co-senior authors

Running Head: Pyruvate as a Modulator of Resilience Against High Genetic Risk for Glaucoma

Corresponding authors:

Louis R. Pasquale, MD, FARVO

Department of Ophthalmology, Icahn School of Medicine at Mount Sinai, New York, NY, USA

Louis.Pasquale@mssm.edu

Simon WM John, PhD

Department of Ophthalmology, Columbia University Irving Medical Center, New York, NY

10032, USA

Keywords: metabolite risk score; metabolites; genetic risk score; genetics; glaucoma; glaucoma

resilience; glycolysis; pyruvate; tricarboxylic acid cycle

Abbreviations and Acronyms:

AUC, area under the curve; BMI, body mass index; CI, confidence interval; HDL, high-density

lipoprotein; ICD, International Classification of Diseases; IOP, intraocular pressure; mRNFL,

macular retinal nerve fiber layer; MET, metabolic equivalent of task; MRS, metabolic risk score;

MTAG, Multi-Trait Analysis of Genome-wide association study data; NEF, number of effective;

NMR, nuclear magnetic resonance; OAG, open-angle glaucoma; OR, odds ratio; POAG, primary

open-angle glaucoma; PRS, polygenic risk score; RGC, retinal ganglion cell; ROC, receiver

operating characteristic; SD, standard deviation; SNP, single nucleotide polymorphism; TCA,

Tricarboxylic acid; ATP, Adenosine triphosphate; VCDR, vertical cup-disc ratio; VLDL, very

low-density lipoprotein.

Abstract

A glaucoma polygenic risk score (PRS) can effectively identify disease risk, but some individuals with high PRS do not develop glaucoma. Factors contributing to this resilience remain unclear. Using 4,658 glaucoma cases and 113,040 controls in a cross-sectional study of the UK Biobank, we investigated whether plasma metabolites enhanced glaucoma prediction and if a metabolomic signature of resilience in high-genetic-risk individuals existed. Logistic regression models incorporating 168 NMR-based metabolites into PRS-based glaucoma assessments were developed, with multiple comparison corrections applied. While metabolites weakly predicted glaucoma (Area Under the Curve=0.579), they offered marginal prediction improvement in PRS-only-based models (P=0.004). We identified a metabolomic signature associated with resilience in the top glaucoma PRS decile, with elevated glycolysis-related metabolites—lactate (P=8.8E-12), pyruvate (P=1.9E-10), and citrate (P=0.02)—linked to reduced glaucoma prevalence. These metabolites combined significantly modified the PRS-glaucoma relationship ($P_{interaction}$=0.011). Higher total resilience metabolite levels within the highest PRS quartile corresponded to lower glaucoma prevalence (Odds Ratio$_{highest\ vs.\ lowest\ total}$ $_{resilience\ metabolite\ quartile}$=0.71, 95% Confidence Interval=0.64–0.80). As pyruvate is a foundational metabolite linking glycolysis to tricarboxylic acid cycle metabolism and ATP generation, we pursued experimental validation for this putative resilience biomarker in a human-relevant Mus musculus glaucoma model. Dietary pyruvate mitigated elevated intraocular pressure (P=0.002) and optic nerve damage (P<0.0003) in $Lmx1b^{V265D}$ mice. These findings highlight the protective role of pyruvate-related metabolism against glaucoma and suggest potential avenues for therapeutic intervention.

Introduction

Glaucoma is a polygenic, progressive neurodegenerative disease and a leading cause of irreversible blindness.[1] The disease is typically asymptomatic until advanced visual field loss occurs, and around 50 to 70% of people affected remain undiagnosed.[2,3] Early detection and intervention are essential to stop disease progression and prevent visual impairment in glaucoma-affected individuals, as there is no cure available; however, population-based glaucoma screening is not cost-effective from a public health perspective.[4–6]

Glaucoma is well-suited for developing and applying a polygenic risk score (PRS) to facilitate disease identification and risk stratification as it is a condition with several endophenotypes, such as elevated intraocular pressure (IOP) and thinning of the retinal nerve fiber layer (RNFL), and it is highly heritable.[7–9] Glaucoma risk is influenced by both genetic and metabolic factors, with emerging evidence suggesting that gene-environment interactions may play a greater role in conferring disease risk than independent exposures alone.[10–15]

Glaucoma was five times more likely in the UK Biobank (UKBB) among participants with the highest genetic risk decile versus the lowest decile (7.4% vs 1.3%).[9] While the relatively low glaucoma prevalence in the highest decile group could be explained by disease under-ascertainment or a PRS that incompletely reflects the glaucoma genetic architecture, it is possible that resilience biomarkers could explain this result, in addition to protective genetic or epigenetic factors.

Recent advancements in metabolomics have opened up avenues to explore metabolites as potential biomarkers for glaucoma.[16] Metabolites are intermediate and end products of cellular processes critical for driving cellular growth and tissue homeostasis.[17] These small molecules provide a holistic measure of physiological status, reflecting both genetic predispositions and

environmental influences. Previous studies have indicated a potential role for plasma metabolites

to stratify glaucoma risk; however, these studies were limited by small sample sizes,[18–22] with

restricted coverage of metabolites. In addition, the potential benefits of the integration of

metabolomics with genetics to identify individuals at the highest risk of glaucoma remain

unexplored.

This study aims to evaluate plasma metabolites as risk factors for glaucoma. First, we

explored whether incorporating plasma metabolite data improved the predictive accuracy of a

PRS for glaucoma risk based on available data in the UKBB, where genetic and metabolomic

data were available for 117,698 participants. Additionally, we evaluated interactions between

glaucoma PRS and metabolomics for further stratifying individuals with high genetic risk but

without glaucoma. We undertook an agnostic approach to identify a metabolite signature

associated with resilience to high glaucoma genetic risk. Finally, we experimentally validated a

top resilience metabolite by assessing its ability to rescue the ocular phenotype in a human-

relevant, genetic mouse model of glaucoma.

Results

UK Biobank Study Characteristics

A total of 117,698 participants (4,658 glaucoma cases and 113,040 non-cases) were

included in this study. UKBB participants were predominately of European ancestry (85.8%) but

were also of admixed American (0.22%), Asian (2.77%), and African (1.73%) ancestry. The

design of the human studies is depicted graphically in **Figures 1 and 2**. There were significant

differences seen in baseline demographic and clinical characteristics between the glaucoma cases

and the non-cases presented in **Table 1**. Notably, characteristics associated with glaucoma

included male gender, older age, genetic African ancestry (Privé et al[23] describes the method for

determining genetically-inferred ancestry in the UKBB), prior history of smoking, slightly lower

cholesterol levels, higher body mass index (BMI), higher hemoglobin A1c (HbA1c), higher IOP,

thinner macular retinal nerve fiber layer (mRNFL) thickness, higher caffeine intake, higher

alcohol intake, higher oral steroid use, diabetes, and coronary artery disease.

Plasma metabolites marginally improve glaucoma risk prediction

We first sought to determine whether the addition of metabolites could improve the

prediction of glaucoma from basic demographics, clinical variables, and genetic data. We created

four logistic regression models, each with increasing predictive variables considered, to evaluate

their added utility for predicting glaucoma (see Methods and **Supplemental Table 2** for model

construction details). We analyzed two sets of metabolite variables: a comprehensive set of 168

metabolites measured by nuclear magnetic resonance (NMR) spectroscopy included in the

UKBB dataset and a limited set of 27 metabolites approved by the European Union (EU) for *in*

vitro diagnostic use.

To evaluate the performance of each model, we plotted receiver operating characteristic

(ROC) curves and calculated the area under the curve (AUC) metrics. In model 1, which utilized

only metabolite data, the highest performance was achieved using the full panel of 168

metabolites, yielding an AUC value of 0.602 (95% CI=0.592-0.612) compared to 0.579 (95%

CI=0.569-0.589; P=0.0003) using the 27 metabolites (**Figure 3**). Model 2, which incorporated

demographic information, and model 3, which also included clinical variables, exhibited similar

performance before the addition of metabolite data. In model 2, the addition of the 168

metabolite panel demonstrated the best performance, with an AUC value of 0.670 (95%

CI=0.660-0.680) compared to 0.664 (95% CI=0.654-0.674) for the model without metabolites

(P=0.002) and 0.666 (95% CI=0.656-0.676) for the limited panel of 27 metabolites. This

performance trend persisted in model 3 with the inclusion of 168 metabolites, producing an AUC

of 0.680 (95% CI=0.660-0.700), representing an increase in AUC (P=0.02) from model 3 with

the exclusion of metabolites (AUC 0.670; 95% CI=0.650-0.690). We subsequently examined

whether integrating metabolite data into a PRS-based model could improve glaucoma prediction

algorithms. In model 4, the panel of 168 metabolites yielded the best performance with an AUC

value of 0.806 (95% CI=0.796-0.816), while the model without metabolites (AUC=0.801, 95%

CI=0.791-0.811; P=0.004), as well as the model with a limited panel of 27 metabolites

(AUC=0.802, 95% CI=0.792-0.812; P=0.006) both had lower AUC values. Thus, these findings

suggest that the addition of metabolite data marginally enhances the prediction of glaucoma

beyond the use of demographic and genetic data.

We then investigated whether the inclusion of the 168 metabolites could improve the

predictive value for glaucoma by stratifying patients into specific subgroups and calculating

AUC curves for each stratum. Accordingly, we found that metabolites showed a modestly

improved predictive value for glaucoma among people of White ethnicity (P<0.001), those aged

55 and older (P=0.002), and males (P=0.002) (**Table 2**).

Metabolites associated with resilience to a high glaucoma polygenic risk score

As metabolite data only marginally augmented clinical or PRS predictions of glaucoma,

we hypothesized that metabolites might provide utility for differentiating patients within risk

groups. Specifically, we studied participants who possessed a high glaucoma PRS but did not

have glaucoma. For this study, we labeled them resilient while recognizing there are many

reasons they may not have glaucoma. Participants were stratified into PRS deciles, and

metabolite signatures were identified to differentiate patients with and without glaucoma in the

top decile (N=11,770) and the bottom half (N=58,358) of the glaucoma PRS distribution (**Figure**

4). Within the top decile of glaucoma PRS, compared to participants without glaucoma,

participants with glaucoma were more likely to be older, male, of White ethnicity, and were prior

smokers. Glaucoma participants with the highest PRS also had higher BMI, higher HbA1c,

higher spherical equivalent, consumed more caffeine and alcohol, and were more likely to have

diabetes and coronary artery disease (**Table 3**). Among participants in the bottom half of

glaucoma PRS, participants with glaucoma were also more likely to be older, male, of Black and

Asian ethnicity, prior smokers, had higher BMI and higher HbA1c, and were more likely to have

diabetes and coronary artery disease (**Table 4)**. As expected, participants with glaucoma in both

bins of glaucoma genetic risk had higher IOP and thinner mRNFL thickness.

Our multivariable-adjusted analysis revealed that higher probit-transformed levels of

lactate (adjusted P_{NEF}=8.8E-12), pyruvate (adjusted P_{NEF}=2.9E-10), and citrate (adjusted

P_{NEF}=0.018) were independently associated with no glaucoma in the top decile of glaucoma PRS

(**Table 5**). In addition, lower levels of triglycerides and higher levels of selected HDL analytes

had an adjusted $P_{NEF} < 0.2$ and were associated with no glaucoma among participants with high

genetic risk. Among the bottom half of the PRS distribution, higher levels of albumin were

associated with no glaucoma (adjusted P_{NEF}=0.047), while higher levels of small HDL, omega-3

fatty acids, docosahexaenoic acid, lactate, and citrate were associated with no glaucoma, albeit

with an adjusted $P_{NEF} < 0.2$.

Interaction of metabolic and genetic biomarkers

The three metabolites (lactate, pyruvate, and citrate) for which higher levels were

associated with reduced glaucoma prevalence in individuals with high genetic susceptibility had

no statistically significant relationship with glaucoma for participants in the lower half of the

PRS distribution at the adjusted $P_{NEF} < 0.05$ level. Thus, we hypothesized that there may be an

interaction between these metabolites and genetic risk. Such an interaction would suggest that

these glycolysis/tricarboxylic acid (TCA) cycle metabolites are relevant primarily in the setting

of high PRS, establishing them as resilience factors against elevated genetic risk of glaucoma.

Indeed, we observed a significant interaction between elevated levels of total lactate, pyruvate,

and citrate with glaucoma PRS quartile for predicting the risk of glaucoma ($P_{interaction}$=0.011;

Figure 5A).

To confirm and visualize this interaction, we calculated and plotted the glaucoma odds

ratio (OR) as a function of PRS quartile with each genetic predisposition bin stratified by the

sum of the resilience-metabolite probit score quartiles (**Figures 5A and 5B**). Notably, high

levels of resilience-associated metabolites were significantly associated with lower odds of

glaucoma, particularly within the higher PRS quartiles (Q3 and Q4). For example, for individuals

in PRS Q3 with total resilience-associated metabolite sum Q3, the glaucoma OR was 0.78 (95%

CI=0.65-0.93; P=0.0059), using metabolite sum Q1 as the reference. In the same PRS Q3 with a

higher quartile of total resilience-associated metabolite sum (Q4), the OR was 0.74 (95%

CI=0.62-0.88; P=0.0009). The lowest OR of glaucoma was observed in participants with both

the highest PRS (Q4) and highest resilience-associated metabolite sum (Q4), with an OR of 0.71

(95% CI=0.64-0.80; P<0.001). Although both age and PRS were significant predictors of

glaucoma prevalence (both are strongly associated with increased risk), there was no evidence of

a significant three-way interaction between the resilience-associated metabolite sum, PRS, and

age ($P_{3\text{-}way\text{-}interaction}$=0.65).

Next, we investigated whether a holistic metabolic risk score (MRS) incorporating

resilience metabolites and other measured metabolites (full panel of 168 metabolites; see

Methods and **Supplemental Table 3** for the beta coefficients used in MRS construction) could

be used in conjunction with a glaucoma PRS to improve risk prediction. To quantify the degree

of interaction of glaucoma PRS and MRS, we plotted glaucoma OR at various quartiles of

glaucoma PRS and MRS compared to the first MRS quartile within each PRS quartile. We found

a significant interaction between glaucoma MRS built from 168 metabolites and PRS

($P_{interaction}$=0.0012; see **Supplemental Figure 1A**).

In stratified analyses by quartile of PRS, compared to the first quartile of MRS, higher

MRS was associated with increased glaucoma prevalence across most PRS quartiles, with the

most pronounced effect observed in individuals in the highest PRS and MRS quartile with an OR

of 2.14 compared to those in the highest PRS and lowest MRS quartile (**Supplemental Figure**

1B). Among individuals in the highest MRS quartile (Q4), PRS Q1 had an OR of 1.66 (95%

CI=1.15-2.42; P=0.0086), PRS Q2 had an OR of 2.08 (95% CI=1.62-2.67; P<0.001), PRS Q3

had an OR of 2.20 (95% CI=1.83-2.65; P<0.001), and PRS Q4 had an OR of 2.14 (95%

CI=1.91-2.40; P<0.001). This suggests potential synergistic effects of genetic and metabolite risk

factors on glaucoma risk that transcend the impacts of the resilience metabolites.

Most importantly, using individuals in the lowest quartiles of PRS and MRS (PRS Q1,

MRS Q1) as the reference group for the entire population (n=117,698), we determined glaucoma

OR as a function of PRS quartile, with further stratification by holistic MRS in each PRS bin

(**Figure 6**). The OR for PRS Q1 with MRS Q4 relative to PRS Q1 with MRS Q1 was modest

(OR=1.66; 95% CI= 1.15–2.42; P=0.0086). However, those with PRS Q4 (highest genetic risk)

and MRS Q1 (lowest metabolic risk) had an OR of 11.7 (95% CI=8.72-16.0; P<0.001) compared

to PRS Q1 MRS Q1. This illustrates the importance of genetics in impacting glaucoma risk.

Strikingly, the glaucoma OR further increased to 25.1 (95% CI=18.8-34.1; P<0.001) for those

with PRS Q4, MRS Q4 (combination of highest genetic and highest metabolic risk;

P*interaction*=0.019). Altogether, this demonstrates the strong prognostic utility of combining both

PRS and MRS measurements for assessing glaucoma risk, particularly in individuals with a high

genetic predisposition.

Pyruvate supplementation lessens intraocular pressure and glaucoma

To functionally test the association between higher levels of pyruvate and resilience to

glaucoma, we experimentally tested whether treatment with pyruvate induces resilience to IOP

elevation and glaucoma in a human-relevant mouse model. Common variants in *LMX1B* are

associated with IOP variation and the most common form of human glaucoma, primary open-

angle glaucoma (POAG).[24–29] Rare Mendelian variants in *LMX1B* can produce early-onset ocular

hypertension and open-angle glaucoma (OAG).[30] We have previously demonstrated that mice

with a dominant mutation in *Lmx1b* (*Lmx1b$^{V265D/+}$*) develop IOP elevation and glaucoma.[30,31]

Depending on the genetic background, this *Lmx1b^{V265D}* mutation induces either early-onset or

later glaucoma[31]. Mutant mice with a C57BL/6J strain background develop severe, early-onset

IOP elevation and glaucoma. As pyruvate and its metabolites were associated with no glaucoma

despite strong genetic predisposition in the UKBB cohort (highest decile of PRS), we tested the

ability of dietary pyruvate to induce resilience against the *Lmx1b^{V265D}*-induced glaucoma on this

C57BL/6J genetic background. Pyruvate supplementation through drinking water substantially

protected mice from IOP elevation and glaucoma. Pyruvate significantly protected against both

anterior chamber deepening (ACD), a consequence of IOP elevation in mouse eyes (**Figure 7A-**

B), and IOP elevation itself (**Figure 7C**). Importantly, pyruvate treatment protected against

glaucomatous optic nerve degeneration (**Figure 7D**). Together, our findings strongly support the

role of endogenous pyruvate in conferring resilience against glaucoma even countering strong

genetic predisposition. Further, they show that pyruvate can act as a potent resilience factor

against IOP elevation and glaucoma when delivered orally.

Discussion
270

In this large cohort study, we found that the inclusion of metabolite data from an NMR

platform only marginally improved various glaucoma prediction algorithms **(Figure 3)**.

Nonetheless, using an agnostic approach, we found that lactate, pyruvate, and citrate levels

collectively were associated with a 29% reduced risk of glaucoma among those in the top

quartile of glaucoma genetic predisposition **(Figure 5B)**. We showed that pyruvate

supplementation reduced glaucoma incidence in a human-relevant genetic mouse model (**Figure**

7). Furthermore, the interaction between glaucoma PRS and an MRS based on 168 metabolites

provided synergistic predictive information regarding glaucoma risk (**Figure 6**). The statistical

interaction we demonstrate between metabolite scores and PRS-derived glaucoma risk helps

identify individuals who, despite high genetic risk, are less likely to develop glaucoma due to

favorable metabolic profiles or, on the contrary, are more likely to develop glaucoma and should

be monitored at an earlier age. Overall, these findings indicate that an MRS may provide

clinically useful measures of metabolic status and improve patient risk stratification in glaucoma.

While earlier studies[32–34] showed modest discriminatory powers for a glaucoma PRS,

recent larger genome-wide association studies reported significantly improved risk

prediction.[7,35–37] However, no studies have examined the utility of incorporating metabolite data

into glaucoma prediction algorithms combined with a PRS. Future work is needed to understand

how non-genetic factors, known to alter glaucoma risk or IOP levels, like air pollution,

psychological stress, physical activity, and dietary factors,[14] could change the MRS and alter

high glaucoma genetic predisposition.

Previous studies have explored metabolomics to identify biomarkers for OAG. A

systematic review analyzing 13 studies identified a total of 144 metabolites, of which 12 were

associated with OAG.[22] Among studies using plasma samples, four metabolic pathways were

significantly enriched: sphingolipid metabolism, arginine and proline metabolism, and beta-

alanine metabolism. In another study across three US cohorts and the UKBB, higher levels of

plasma diglycerides and triglycerides were adversely associated with glaucoma.[38] This is

consistent with our observation that lower triglyceride levels were associated with glaucoma

resilience, albeit with less significance compared to the three glycolysis-related metabolites. The

biological mechanism is suggested to be the association of hypertriglyceridemia with increased

blood viscosity and elevated IOP.[39–42] Consistent with this, a meta-analysis reported that

individuals with glaucoma had a mean triglyceride level 14.2 mg/dl higher than those without the

disease.[41]

While studies have identified biomarkers associated with glaucoma, to the best of our

knowledge, there has been no investigation on biomarkers that are associated with being resilient

to a high glaucoma genetic predisposition. We identified elevated levels of lactate, pyruvate, and

citrate among individuals without glaucoma despite high genetic risk. The retina commonly

utilizes energy produced from glycolysis and oxidative phosphorylation. Therefore, glycolysis

end products (pyruvate and lactate) are important energy sources for retinal ganglion cells

(RGCs).[43] RGCs express monocarboxylate transporters, which facilitate the uptake of

extracellular pyruvate and lactate, improving energy homeostasis, neuronal metabolism, and

survival.[44,45] Lactate is critical for RGC survival during periods of glucose deprivation,[46,47] and

bioenergetic insufficiency contributes to RGC loss in glaucoma.[48–51] Finally, a study found that

baseline lactate levels were significantly decreased in patients with normal-tension glaucoma

compared to non-glaucomatous controls.[52]

Pyruvate plays a central role in energy metabolism by linking glycolysis to the TCA

cycle, ultimately driving ATP production by the electron transport chain. Pyruvate is converted

into acetyl-CoA, which fuels the TCA cycle, and into lactate, a key molecule that regulates

metabolism and serves as an important energy source.[53] The TCA cycle is tightly regulated to

balance biosynthesis (producing metabolic precursors for lipids, carbohydrates, amino acids,

nucleic acids, and co-factors) and energy production, adjusting dynamically to cellular needs.[54]

Beyond its metabolic functions, pyruvate also acts as an antioxidant, scavenging reactive oxygen

species (ROS), induceing the expression of the antioxidant response control gene *Nrf2*, and

promoteing beneficial autophagy.[55–57] Its neuroprotective effects on RGCs have been

demonstrated both in cell culture and in an induced model of glaucoma in rats.[58] Additionally,

pyruvate combined with nicotinamide has been shown to enhance optic nerve protection in

mice.[59] Importantly, the current study extends this evidence by demonstrating that pyruvate alone

provides strong protection against glaucoma in a mouse model carrying a mutation in the

ortholog of a human POAG gene. Further, it extends pyruvate's protective effects to the ocular

drainage tissues, mitigating IOP elevation. Notably, although our validation data demonstrates

the neuroprotective effects of exogenous pyruvate, it remains possible that endogenously

produced pyruvate within ocular tissues may also contribute to RGC protection. A randomized

Phase II clinical trial that tested pyruvate and nicotinamide in POAG patients found improved

visual function in glaucoma patients compared to the placebo group.[60] Together, both preclinical

and clinical evidence support the role of pyruvate as a protective factor against glaucoma.

Citrate is a key intermediate in the TCA cycle and is important for aerobic energy

production. Elevated citrate levels might reflect enhanced energy supply and antioxidative

capacity of ocular tissues, potentially protecting RGCs against oxidative stress.[50,61–64] Decreased

levels of citrate may indicate increased usage or mitochondrial dysfunction.[65] Studies have found

lower plasma citrate levels in glaucoma patients among adults and children.[66,67] A study

measuring plasma citrate with a cutoff of 110 μmol/L to detect glaucoma had a sensitivity of

66.7% and a specificity of 71.4%.[66] These findings corroborate the protective effects of citrate in

conferring resilience to glaucoma. Similar to pyruvate, plasma citrate may also play a role in

retinal homeostasis, highlighting the need to disentangle serum-derived from intraocularly

synthesized metabolites.

We also identified several lipoproteins associated with resilience, most notably

cholesteryl esters in small high-density lipoprotein (HDL) and medium HDL. Although these

lipoproteins did not reach an adjusted P-value<0.05 and thus remain suggestive biomarkers

rather than definitive, this is consistent with studies describing the neuroprotective effect of HDL

in the glaucoma pathophysiology.[68,69] Cholesteryl esters and triglycerides make up the core of

lipoproteins, which transport lipids in the blood. Lipoproteins are characterized by their size,

where HDL ranges from 8-12 nm in diameter, and low-density lipoprotein (LDL) ranges from

18-25 nm in diameter. The retinal pigment epithelium can take up lipoproteins,[70,71] and

dysregulation of lipid metabolism can result in the accumulation of lipids and oxidative stress in

the retina, resulting in RNFL thinning.[38,72–75] Thus, HDL has antioxidant properties to protect

RGCs by facilitating cholesterol efflux from the retina.[76,77] Furthermore, as HDL analytes can

cross blood-ocular barriers,[78] there is a plausible route for serum-derived HDL to influence RGC

homeostasis. Nonetheless, the relative contributions of circulating lipoproteins versus local

synthesis within ocular tissues remain unclear and warrant further investigation. Notably, genes

implicated by large cross-ancestry and European subset GWAS meta-analyses of POAG risk are

enriched in processes related to apolipoprotein binding and lipid transport.[26,78] Incorporating

ocular physiology and blood-retinal barrier considerations when interpreting lipoproteins as

potential resilience biomarkers will be critical for future studies aimed at understanding and

therapeutically targeting increased glaucoma risk.

This study had several strengths. First, the UKBB is a large cohort study with detailed

covariate information on demographics, clinical characteristics, and glaucoma endophenotypes,

allowing for better glaucoma risk profiling. Second, we used a PRS calculated from the large

(~600k individuals) multi-trait analysis of genome-wide association studies on glaucoma and

endophenotypes, including optic nerve head structural features and IOP data.[7] Finally, we

supported our biobank analyses with functional tests in a disease-relevant mouse model,

identifying a protective effect of pyruvate supplementation in animals susceptible to glaucoma.

The data showing that pyruvate protected against IOP elevation (a key causative risk factor for

glaucoma) and glaucomatous nerve damage in a mouse model, as well as pyruvate's biochemical

relatedness to lactate and citrate regarding energy metabolism, mitigate against concerns about

false discovery, uncontrolled confounding, reverse causation, and collinearity among

biomarkers.

A major limitation of our study is that it includes a predominately European population,

with an underrepresentation of African, Asian, and mixed American genetic ancestral groups;

nonetheless, we chose to analyze Europeans and non-Europeans together to make the results as

generalizable as possible. A second limitation is that glaucoma ascertainment in the UK Biobank

is based on self-reported diagnoses and hospital records rather than comprehensive

ophthalmologic examinations. Nonetheless, it is reassuring that the prevalence of glaucoma in

our sample (~4%) is similar to a directly performed disease burden estimate in a comparable,

albeit slightly older, United Kingdom sample (2.7%).[79] In addition, a subset of UKBB

participants have received glaucoma treatment, which could alter metabolite levels and could

lead to an underestimate of the protective effects of metabolites. Moreover, the UKBB study

includes a limited set of 168 metabolites, which is not inclusive of the comprehensive set of

metabolites, known and unknown. This study also only focuses on plasma metabolites and does

not account for ocular metabolite profiles. Due to the cross-sectional design of our study,

establishing temporality is challenging. As a result, it remains unclear whether the observed

metabolic changes contribute to the onset of glaucoma or arise as a consequence of the disease.

However, the mouse data support a causal role for pyruvate as a factor that mitigates high

genetic risk. This study's exploratory approach– investigating metabolite associations in an

agnostic, hypothesis-free manner– underscores the need for replication. Regarding the mouse

studies, we performed ad-lib pyruvate dosing and examined only a single human-related

mutation, whereas individuals with a genetic predisposition to glaucoma typically carry a high

cumulative burden of multiple common glaucoma-associated variants. Lastly, neither the mouse

studies, nor the human data accurately inform the exact dosing of TCA cycle biomarkers needed

to mitigate glaucoma risk. We find considerable overlap between TCA cycle biomarkers

between those with and without glaucoma in the top decile of glaucoma risk (**Supplementary**

Table 1 and **Supplementary Figure 2**).

Overall, we have shown that plasma metabolites marginally enhance the predictive power

of a glaucoma PRS and may have a role in identifying those who may be resilient to glaucoma

despite high genetic predisposition. Our study identified protective metabolites linked to

glycolysis and mitochondrial function, suggesting important pathophysiology in glaucoma. One

of these metabolites of significance is pyruvate. Oral pyruvate supplementation substantially

protected against IOP elevation and conferred resilience against glaucoma. Thus, these results

open new avenues for therapeutic strategies of pyruvate and its metabolites as resilience factors

against glaucoma in genetically predisposed individuals.

Methods

Study Design

Our initial human analysis was conducted in three steps. First, we assessed whether

plasma metabolite data alone can predict glaucoma risk with receiver operating characteristic

(ROC) curves in the UK Biobank. We then integrated the metabolite data into a polygenic risk

score (PRS)-based glaucoma risk assessment model to see if metabolites could enhance

glaucoma prediction. Second, we focused on identifying a metabolomic signature of resilience to

high glaucoma PRS by comparing plasma metabolites in glaucoma cases versus participants

without glaucoma (henceforth referred to as resilient participants) from the top 10% of the

glaucoma PRS. Third, we explored the interactions between metabolic scores and PRS in

modifying glaucoma risk.

Study population and data collection

The UKBB is a prospective cohort study of over half a million participants aged 37 to 73

years at recruitment across the United Kingdom from 2006 to 2010. Participants were recruited

through the National Health Service registers from 22 assessment centers, where they signed

electronic informed consent to participate. Participants then completed in-depth touchscreen

questionnaires and trained staff-led interviews, performed body measurements, and provided

biological samples.[80] Additional outcomes are available from data linkage to hospital episode

statistics, the death register, and primary data.[80–82] Biological samples collected from these

participants, including blood, urine, and saliva specimens, were used to generate genetic,

metabolomic, and proteomic data.[83] Details of the UKBB study design and population are

described online (https://www.ukbiobank.ac.uk). The UKBB study received approval from the

National Health Service North West Multicentre Research Ethics Committee (reference number

06/MRE08/65) and the National Information Governance Board for Health and Social Care. This

research, conducted under UK Biobank application number 36741, adhered to the tenets of the

Declaration of Helsinki.

The UKBB included 502,613 participants, of which 173,679 participants had self-

reported glaucoma data, ICD-coded, or previous glaucoma laser/surgical therapy (34.6%) at

baseline. A subset consisting of 117,698 participants with metabolomic data and genetic

profiling comprise the study population (**Figure 1**). The glaucoma cases were defined based on

those with self-reported glaucoma, in which participants selected "glaucoma" when they

completed a touchscreen questionnaire with the question, "Has a doctor told you that you have

any of the following problems with your eyes?". Glaucoma cases also included participants who

reported a history of glaucoma surgery or laser therapy on the questionnaire or if an International

Classification of Diseases (ICD) code for glaucoma (ICD 9[th] revision: 365.* [excluding 365.0];

ICD 10[th] revision: H40.* [excluding H40.0] and H42.*) was carried in linked Hospital Episode

Statistics before the baseline assessment. The described approach to identifying glaucoma has

been strongly supported by prior research publications.[27,28]

Metabolite Profiling

In the UKBB, a high-throughput nuclear magnetic resonance (NMR)-based biomarker

platform (Nightingale Health Ltd; Helsinki, Finland) was used to measure the metabolomic

profile in the randomly selected non-fasting EDTA plasma samples from a subset of

participants.[84] In contrast to liquid chromatography/mass spectroscopy, NMR spectroscopy

produces distinctive spectral shapes for molecules containing hydrogen atoms.[85] The areas under

the curve (AUC) are proportional to the concentration of each molecule based on chemical shifts

and J coupling split patterns derived from quantum mechanics.[86,87] The NMR platform contains

data on 249 metabolic biomarkers (168 absolute levels and 81 ratio measures), including a subset

of 36 biomarkers (27 absolute levels and 9 ratio measures) certified for broad diagnostic use by

the European Union (EU).[84] In this study, we focused on the 168-metabolite and the 27-

metabolite sets for analysis (for a complete list of quantified metabolites, refer to the

Supplementary File). Individual metabolite values were transformed to probit scores to

standardize their range and reduce the impact of skewed distributions.

Construction of PRS for glaucoma

The glaucoma PRS used in this study was from a multi-trait analysis of GWAS (MTAG)

on glaucoma.[7] This PRS was derived based on GWAS data from glaucoma (7,947 cases and

119,318 controls) and its endophenotypes: optic nerve head structure using vertical cup-disc ratio

(VCDR) (including additional data from 67,040 UKBB and 23,899 International Glaucoma

Genetics Consortium (IGGC) participants), and intraocular pressure (IOP, including additional

data on 103,914 UKBB and 29,578 IGGC participants). The glaucoma PRS was constructed

using single nucleotide polymorphisms (SNPs) with MTAG P-values ≤ 0.001, resulting in 2,673

uncorrelated SNPs after LD-clumping at $r^2 = 0.1$, and was found to have a predictive ability with

an area under the curve (AUC) of 0.68 (95% confidence interval, CI, [0.67-0.70]) and 0.80

accounting for age, sex, family history, and PRS tested in the Australian and New Zealand

Registry of Advanced Glaucoma cohort.

Model building/covariates

We first assessed whether the inclusion of probit-transformed metabolite data enhances

glaucoma prediction algorithms. We adjusted for factors involving major determinants of

variability in metabolites, POAG-established and suspected risk factors, and other comorbidities.

In model 1, we considered metabolites only. Model 2 incorporated additional demographic

covariates, including age (years), sex, genetic ancestry, season, time of day of specimen

collection (morning, afternoon, night), and fasting time (hours). Model 3 incorporated covariates

in model 2 and additional variables accounting for comorbidities potentially related to glaucoma

including smoking status (never, past, and current smoker), alcohol intake (g/week),[10] caffeine

intake (mg/day),[88] physical activity (metabolic equivalent of task [MET], hours/week), body

mass index (kg/m^2), average systolic blood pressure (mm Hg), history of diabetes, hemoglobin

A1C (HbA1c, mmol/mol), self-reported history of coronary artery disease, systemic beta-blocker

use, oral steroid use, and mean spherical equivalent refractive error (diopters) across both eyes.

Model 4 incorporated covariates in model 3 and a glaucoma PRS. We compared these models

with and without the metabolites to determine the predictive ability of the metabolites above and

beyond a glaucoma PRS and known risk factors. Missing values were imputed with medians for

numeric variables and modes for factor variables.

Statistical analysis

Metabolite-Based Predictive Modeling

Multivariable logistic regression models with regularization (implemented with *glmnet*)

were built to investigate the associations of metabolic biomarkers and the risk of glaucoma. We

used four sequential models, adjusting for covariates as described above. For each model, we

examined three groupings: no metabolites, the 27 metabolites corresponding to EU-stamped

validated markers, and 168 metabolites corresponding to all the measured plasma metabolites

captured in the UKBB. We used regularization to address collinearity, which reduced the set of

metabolites and other covariates considered in each model (reported in **Supplementary Table**

2). To examine model accuracy for predicting glaucoma prevalence, we utilized ROC curves and

associated AUC measurements as a metric for model performance. We performed five-fold

cross-validation and split data into 80% used for training and 20% used for evaluation. All

metabolites' values were probit transformed and used as continuous variables (per 1 standard

deviation (SD) increase). Additionally, we stratified glaucoma by age, ethnicity, and sex. The

DeLong test was utilized to examine the statistical significance of AUC differences between

models, with a threshold of $P<0.05$ as statistically significant.

Identifying resilience metabolites to high glaucoma polygenic risk scores

507   To identify individual metabolites associated with resilience to a high glaucoma PRS, we

first performed a logistic regression model to obtain metabolite residuals from probit-

transformed concentration data adjusting for the following variables: age, age-squared, time

since the last meal/drink ($\leq$4, 5–8, and 9+ hours), sex, ethnicity (Asian, Black, White, and

mixed/other), season, time of day of specimen collection, smoking status (never, past, and

current smoker), alcohol intake, caffeine intake, physical activity (metabolic equivalent of task

[MET] hours/week), body mass index (kg/m^2), average systolic blood pressure (mm Hg), history

of diabetes, HbA1c (mmol/mol), history of coronary artery disease, systemic beta-blocker use,

oral steroid use, and spherical equivalent refractive error (diopters). Metabolites associated with

glaucoma were nominated by performing a t-test comparing residuals in participants with and

without glaucoma to identify metabolites significantly associated with glaucoma in both the top

10% and the bottom 50% of glaucoma PRS score. This stratification balanced case counts to

enable comparison of groups with adequate sample size (780 glaucoma cases in the bottom 50%

of PRS score and 1,693 glaucoma cases in the top 10% of PRS score). To account for covariance

in metabolite abundance, we corrected for multiple comparisons using the number of effective

tests (NEF) method.[89] First, a correlation matrix was generated using pairwise complete

observations to address any missing values. Eigenvalue decomposition was performed on the

correlation matrix to capture the principal components. Principal components contributing more

than 1% of the total variance were calculated, yielding 9 significant components, capturing

91.6% of the variance. Thus, we applied a Bonferroni adjustment based on the number of

significant components (n=9) to calculate the adjusted p-values accounting for multiple

hypotheses. P-values are considered statistically significant if the NEF-adjusted P-value was

<0.05.[89] Given this was an exploratory analysis, NEF-adjust P-value <0.2 were considered

worthy of additional analysis.

Construction of Metabolite Risk Scores

We constructed metabolite risk scores (MRS), which were calculated as a weighted sum

of each metabolic biomarker from coefficients in the 168-metabolite base model. Specifically,

we utilized a logistic regression model (implemented with a *binomial glm*) using probit

transformed metabolite values to predict the prevalence of glaucoma using five-fold cross-

validation. The resulting beta values were extracted and are reported in **Supplementary Table 3**.

To determine if there is an interaction between glaucoma MRS and PRS, we utilized extracted

model coefficients and corresponding significance from a binomial generalized linear model. We

fit this model using the formula *Glaucoma ~ PRS * MRS + PRS + MRS + Covariates*. We

classified participants into 16 groups by quartiles of MRS and PRS. Within each PRS quartile,

we used MRS quartile 1 as the reference for comparison. Additionally, we used the group in both

PRS quartile 1 and MRS quartile 1 as the overall reference group for all comparisons among the

groups. A chi-squared test was used to assess statistical significance between odds ratios (ORs)

with 95% confidence intervals (CIs).

To investigate whether age modifies the associations of resilience metabolites with

glaucoma, we conducted a stratified analysis and assessed a three-way interaction term of age ($\geq$

or <58 years old, based on median age), the sum of resilience metabolite levels, and PRS using a

Wald test for individuals in the top 10% PRS. All statistical analyses and plots were produced

using R version 4.2.1 (R Foundation for Statistical Computing, Vienna, Austria). All statistical

tests were two-sided.

Animal husbandry and ethics statement

Experimental mice had a C57BL/6J (Jackson Laboratory Stock #000664) genetic

background. The *Lmx1b^{V265D}* (alias *Lmx1b^{Icst}*) mutation is previously reported to cause high IOP

and glaucoma in mice, with IOP becoming elevated in some eyes during the first months after

birth.[30,31,90] The mutation was backcrossed to the C57BL/6J strain for at least 30 generations. All

mice were treated per the Association for Research in Vision and Ophthalmology's statement on

the use of animals in ophthalmic research. The Institutional Animal Care and Use Committee of

Columbia University approved all experimental protocols performed. Mice were maintained on

PicoLab Rodent Diet 20 (5053, 4.5% fat) and provided with reverse osmosis-filtered water.

Mutant and control littermates were housed together in cages containing ¼-inch corn cob

bedding, covered with polyester filters. The animal facility was maintained at a constant

temperature of 22°C with a 14-hour light and 10-hour dark cycle.

 Genotyping mice

        Lmx1b^{V265D} and *Lmx1b$^+$* genotypes were determined by direct Sanger sequencing of a

 specific PCR product. Genomic DNA was PCR amplified with forward primer 5′-

 CTTTGAGCCATCGGAGCTG-3′ and reverse primer 5′-ATCTCCGACCGCTTCCTGAT-3′

 using the following program: (1) 94 °C for 3 min; (2) 94 °C for 30 s; (3) 57 °C for 30 s; (4) 72 °C

 for 1 min; (5) repeat steps 2–4 35 times; and (6) 72 °C for 5 min. PCR products were purified and

 sequenced by the Genewiz (Azenta Life Sciences).

 Slit-lamp examination

        Anterior segments were examined approximately every 2 weeks between 1 month and 3

 months of age and examined at monthly intervals between 3 months of age and harvest age (6

 months). Balanced groups of males and females were examined as previously published.[30,31]

 Photographs were taken with a 40× objective lens. Anterior chamber deepening, a sign of

 exposure to high IOP, was graded based on a semiquantitative scale of no deepening (not

 present), mild, moderate, or severe, as previously published.[31] Under examination by either a slit-

 lamp or dissection scope (during IOP procedure), anterior chamber deepening was only detected

 in *Lmx1b* mutant groups. Groups were compared statistically by Fisher's exact test. n > 40 eyes

 were examined in each group.

 Intraocular pressure measurement

        IOP was measured with the microneedle method as previously described in detail.[91,92]

 Before cannulation, mice were acclimatized to the procedure room and anesthetized via an

 intraperitoneal injection of a mixture of ketamine (99 mg/kg; Ketlar, Parke-Davis, Paramus, NJ,

 USA) and xylazine (9 mg/kg; Rompun, Phoenix Pharmaceutical, St Joseph, MO, USA)

 immediately prior to IOP assessment, a procedure that does not alter IOP in the experimental

window.[92] IOP was measured at both 5-6 weeks and 8-10 weeks of age in WT and $Lmx1b^{V265D/+}$

eyes. Balanced groups of males and females were examined. During each IOP measurement

period, the eyes of independent WT B6 mice were assessed in parallel with experimental mice as

a methodological control to ensure proper calibration and equipment function. $Lmx1b$ mutant

groups were compared by ANOVA followed by Tukey's honestly significant difference. n > 30

eyes were examined in each $Lmx1b^{V265D}$ mutant group and n > 20 eyes were examined in WT

groups.

Pyruvate administration

Ethyl pyruvate (Sigma-Aldrich, St. Louis, Missouri) was dissolved in the standard

institutional drinking water to a dose of 2000 mg/kg/day for adult mice based on the average

volume mice consume. The mothers consumed this dose and delivered it to their suckling pups at

an unknown dose throughout the first 3 to 4 weeks of life. At 4 weeks of age, the mice were

weaned into separate cages. From 4 weeks on, the young mice consumed a dose of

approximately 2000 mg/kg/day based on their water consumption and body weight over weekly

intervals. Untreated groups received the same drinking water without pyruvate. The water was

changed once per week. Treatment was started on postnatal day 2. Births were checked daily

between 9 am and 12 pm to determine the pups' age.

Optic nerve assessment

We analyzed $Lmx1b^{V265D/+}$ and WT controls optic nerve for glaucomatous damage at 6

months of age (5.3-5.9 month range, majority are 5.8-5.9 months, sex balanced). Intracranial

portions of optic nerves were dissected, processed, and analyzed as previously described.[48,93,94]

Briefly, optic nerve cross-sections were stained with para-phenylenediamine (PPD) and

examined for glaucomatous damage. PPD stains all myelin sheaths, but differentially darkly

stains the myelin sheaths and the axoplasm of damaged axons. This allows for the sensitive

detection and quantification of axon damage and loss. Optic nerves were prepared for analysis

with a 48-h fixation in 0.8% paraformaldehyde and 1.2% glutaraldehyde in 0.08 M phosphate

buffer (pH 7.4) at 4 °C followed by overnight post-fix in osmium tetroxide at 4 °C. Nerves were

washed twice for 10 min in 0.1 M phosphate buffer, once in 0.1 M sodium-acetate buffer, and

dehydrated in graded ethanol concentrations. Tissues were then embedded in Embed 812 resin

(Electron Microscopy Sciences, Fort Washington, PA, USA), and 1 μm-thick sections were

stained in 1% PPD for ~40 minutes. Stained sections were compared using a damage scale that

is validated against axon counting.[93,95] Multiple sections of each nerve were considered when

determining damage level. Nerves were determined to have one of 4 damage levels: 1) No

glaucoma (NOE) – less than 5% axons damaged. This level of damage is seen in age- and sex-

matched non-glaucomatous mice and is not due to glaucoma. We named this level no or early

stage as some have early molecular changes when assessed with transcriptomics, but they cannot

be distinguished from control by morphology;[50,96] 2) Moderate damage (MOD) – average of

30% axon loss; 3) Severe (SEV) – greater than 50% axonal loss and extensive axon damage; and

4) Very severe (V. SEV) – glial scar over the vast majority of nerve with few remaining axons.

Acknowledgements

The authors would like to thank the participants who contributed their data to the UK Biobank

study for this study. The authors would also like to thank the members of the Simon John

Laboratory for experimental and technical assistance and the Institute of Comparative Medicine

for animal and veterinary care.

Consortia

UK Biobank Eye and Vision Consortium

Pirro Hysi & Anthony Khawaja

CRediT authorship contribution statement

Keva Li: Writing – original draft, Writing – review & editing, Methodology, Investigation, Formal analysis, Conceptualization. Nicholas Tolman: original draft, Writing – review & editing, Methodology, Investigation, Formal analysis, Conceptualization. Ayellet V. Segrè: Writing – review & editing, Methodology, Investigation. Kelsey V. Stuart: Data Curation, Methodology, Resources, Writing – review & editing, Methodology, Investigation. Oana A. Zeleznik: Writing – review & editing, Investigation. Neeru A. Vallabh: Writing – review & editing, Investigation. Kuang Hu: Writing – review & editing, Investigation. Nazlee Zebardast: Writing – review & editing, Investigation. Akiko Hanyuda: Writing – review & editing, Investigation. Yoshihiko Raita: Writing – review & editing, Investigation. Christa Montgomery: Writing – review & editing, Investigation. Chi Zhang: Writing – review & editing, Investigation. Pirro G. Hysi: Writing – review & editing, Resources, Project administration, Methodology, Investigation, Funding acquisition. Ron Do: Writing – review & editing, Investigation. Anthony Khawaja: Writing – review & editing, Supervision, Resources, Project administration, Methodology, Investigation, Funding acquisition, Conceptualization. Janey L. Wiggs: Writing – review & editing, Investigation, Funding acquisition. Jae H. Kang: Writing – review & editing, Supervision, Resources, Methodology, Investigation, Funding acquisition. Simon WM. John: Writing – review & editing, Supervision, Resources, Project administration, Methodology, Investigation, Conceptualization, Funding acquisition. Louis R. Pasquale: Writing – review & editing, Supervision, Resources, Project administration, Methodology, Investigation, Funding

653   acquisition, Conceptualization. All authors were involved in the interpretation of the results, and

654   the critical review and approval of the manuscript.

Data Availability

656   Data from the UK Biobank cannot be shared due to our Material Transfer Agreement. To request

657   access to the UK Biobank data, please make requests directly to the UK Biobank via

658   https://www.ukbiobank.ac.uk/enable-your-research/apply-for-access.

References

1. Soh Z, Yu M, Betzler BK, et al. The Global Extent of Undetected Glaucoma in Adults: A
Systematic Review and Meta-analysis. *Ophthalmology*. 2021;128(10):1393-1404.
doi:10.1016/j.ophtha.2021.04.009

2. Khawaja AP, Viswanathan AC. Are we ready for genetic testing for primary open-angle
glaucoma? *Eye*. 2018;32(5):877-883. doi:10.1038/s41433-017-0011-1

3. Jan C, He M, Vingrys A, Zhu Z, Stafford RS. Diagnosing glaucoma in primary eye care and
the role of Artificial Intelligence applications for reducing the prevalence of undetected
glaucoma in Australia. *Eye*. 2024;38(11):2003-2013. doi:10.1038/s41433-024-03026-z

4. Moyer VA, U. S. Preventive Services Task Force. Screening for Glaucoma: U.S. Preventive
Services Task Force Recommendation Statement. *Ann Intern Med*. 2013;159(7):484-489.
doi:10.7326/0003-4819-159-6-201309170-00686

5. US Preventive Services Task Force. Screening for Primary Open-Angle Glaucoma: US
Preventive Services Task Force Recommendation Statement. *JAMA*. 2022;327(20):1992-
1997. doi:10.1001/jama.2022.7013

6. Zukerman R, Harris A, Oddone F, Siesky B, Verticchio Vercellin A, Ciulla TA. Glaucoma
Heritability: Molecular Mechanisms of Disease. *Genes*. 2021;12(8):1135.
doi:10.3390/genes12081135

7. Craig JE, Han X, Qassim A, et al. Multitrait analysis of glaucoma identifies new risk loci and
enables polygenic prediction of disease susceptibility and progression. *Nat Genet*.
2020;52(2):160-166. doi:10.1038/s41588-019-0556-y

8. de Vries VA, Hanyuda A, Vergroesen JE, et al. The Clinical Usefulness of a Glaucoma
Polygenic Risk Score in 4 Population-Based European Ancestry Cohorts. *Ophthalmology*.
Published online August 14, 2024. doi:10.1016/j.ophtha.2024.08.005

9. Sekimitsu S, Xiang D, Smith SL, et al. Deep Ocular Phenotyping Across Primary Open-Angle
Glaucoma Genetic Burden. *JAMA Ophthalmol*. 2023;141(9):891-899.
doi:10.1001/jamaophthalmol.2023.3645

10. Stuart KV, Luben RN, Warwick AN, et al. The Association of Alcohol Consumption with
Glaucoma and Related Traits: Findings from the UK Biobank. *Ophthalmol Glaucoma*.
2022;6(4):366. doi:10.1016/j.ogla.2022.11.008

11. Kang JH, Willett WC, Rosner BA, Buys E, Wiggs JL, Pasquale LR. Association of Dietary
Nitrate Intake With Primary Open-Angle Glaucoma: A Prospective Analysis From the
Nurses' Health Study and Health Professionals Follow-up Study. *JAMA Ophthalmol*.
2016;134(3):294-303. doi:10.1001/jamaophthalmol.2015.5601

12. Kang JH, Ivey KL, Boumenna T, Rosner B, Wiggs JL, Pasquale LR. Prospective study of
flavonoid intake and risk of primary open-angle glaucoma. *Acta Ophthalmol (Copenh)*.
2018;96(6):e692. doi:10.1111/aos.13705

13. Kang JH, Wu J, Cho E, et al. Contribution of the Nurses' Health Study to the Epidemiology
of Cataract, Age-Related Macular Degeneration, and Glaucoma. *Am J Public Health*.
2016;106(9):1684. doi:10.2105/AJPH.2016.303317

14. Stuart KV, Pasquale LR, Kang JH, Foster PJ, Khawaja AP. Towards modifying the genetic
predisposition for glaucoma: An overview of the contribution and interaction of genetic and
environmental factors. *Mol Aspects Med*. 2023;93:101203. doi:10.1016/j.mam.2023.101203

15. Doucette LP, Rasnitsyn A, Seifi M, Walter MA. The interactions of genes, age, and
environment in glaucoma pathogenesis. *Surv Ophthalmol*. 2015;60(4):310-326.
doi:10.1016/j.survophthal.2015.01.004

16. Nazifova-Tasinova N, Radeva M, Galunska B, Grupcheva C. Metabolomic analysis in
ophthalmology. *Biomed Pap*. 2020;164(3):236-246. doi:10.5507/bp.2020.028

17. Johnson CH, Ivanisevic J, Siuzdak G. Metabolomics: beyond biomarkers and towards
mechanisms. *Nat Rev Mol Cell Biol*. 2016;17(7):451-459. doi:10.1038/nrm.2016.25

18. Javadiyan S, Burdon KP, Whiting MJ, et al. Elevation of Serum Asymmetrical and
Symmetrical Dimethylarginine in Patients with Advanced Glaucoma. *Invest Ophthalmol Vis*
Sci. 2012;53(4):1923-1927. doi:10.1167/iovs.11-8420

19. Burgess LG, Uppal K, Walker DI, et al. Metabolome-Wide Association Study of Primary
Open Angle Glaucoma. *Invest Ophthalmol Vis Sci*. 2015;56(8):5020-5028.
doi:10.1167/iovs.15-16702

20. Kouassi Nzoughet J, Guehlouz K, Leruez S, et al. A Data Mining Metabolomics Exploration
of Glaucoma. *Metabolites*. 2020;10(2):49. doi:10.3390/metabo10020049

21. Leruez S, Marill A, Bresson T, et al. A Metabolomics Profiling of Glaucoma Points to
Mitochondrial Dysfunction, Senescence, and Polyamines Deficiency. *Invest Ophthalmol Vis*
Sci. 2018;59(11):4355-4361. doi:10.1167/iovs.18-24938

22. Wang Y, Hou XW, Liang G, Pan CW. Metabolomics in Glaucoma: A Systematic Review.
Invest Ophthalmol Vis Sci. 2021;62(6):9. doi:10.1167/iovs.62.6.9

23. Privé F, Aschard H, Carmi S, et al. Portability of 245 polygenic scores when derived from
the UK Biobank and applied to 9 ancestry groups from the same cohort. *Am J Hum Genet*.
2022;109(1):12-23. doi:10.1016/j.ajhg.2021.11.008

24. Choquet H, Paylakhi S, Kneeland SC, et al. A multiethnic genome-wide association study of
primary open-angle glaucoma identifies novel risk loci. *Nat Commun*. 2018;9:2278.
doi:10.1038/s41467-018-04555-4

25. Gao XR, Huang H, Nannini DR, Fan F, Kim H. Genome-wide association analyses identify
new loci influencing intraocular pressure. *Hum Mol Genet*. 2018;27(12):2205-2213.
doi:10.1093/hmg/ddy111

26. Gharahkhani P, Jorgenson E, Hysi P, et al. Genome-wide meta-analysis identifies 127 open-
angle glaucoma loci with consistent effect across ancestries. *Nat Commun*. 2021;12:1258.
doi:10.1038/s41467-020-20851-4

27. Khawaja AP, Cooke Bailey JN, Wareham NJ, et al. Genome-wide analyses identify 68 new
loci associated with intraocular pressure and improve risk prediction for primary open-angle
glaucoma. *Nat Genet*. 2018;50(6):778-782. doi:10.1038/s41588-018-0126-8

28. MacGregor S, Ong JS, An J, et al. Genome-wide association study of intraocular pressure
uncovers new pathways to glaucoma. *Nat Genet*. 2018;50(8):1067-1071. doi:10.1038/s41588-
018-0176-y

29. Shiga Y, Akiyama M, Nishiguchi KM, et al. Genome-wide association study identifies seven
novel susceptibility loci for primary open-angle glaucoma. *Hum Mol Genet*. 2018;27(8):1486-
1496. doi:10.1093/hmg/ddy053

30. Cross SH, Macalinao DG, McKie L, et al. A Dominant-Negative Mutation of Mouse Lmx1b
Causes Glaucoma and Is Semi-lethal via LBD1-Mediated Dimerisation. *PLoS Genet*.
2014;10(5):e1004359. doi:10.1371/journal.pgen.1004359

31. Tolman NG, Balasubramanian R, Macalinao DG, et al. Genetic background modifies
vulnerability to glaucoma-related phenotypes in Lmx1b mutant mice. *Dis Model Mech*.
2021;14(2):dmm046953. doi:10.1242/dmm.046953

32. Mabuchi F, Mabuchi N, Sakurada Y, et al. Additive effects of genetic variants associated
with intraocular pressure in primary open-angle glaucoma. *PLOS ONE*. 2017;12(8):e0183709.
doi:10.1371/journal.pone.0183709

33. Tham YC, Liao J, Vithana EN, et al. Aggregate Effects of Intraocular Pressure and Cup-to-
Disc Ratio Genetic Variants on Glaucoma in a Multiethnic Asian Population. *Ophthalmology*.
2015;122(6):1149-1157. doi:10.1016/j.ophtha.2015.01.024

34. Zanon-Moreno V, Ortega-Azorin C, Asensio-Marquez EM, et al. A Multi-Locus Genetic
Risk Score for Primary Open-Angle Glaucoma (POAG) Variants Is Associated with POAG
Risk in a Mediterranean Population: Inverse Correlations with Plasma Vitamin C and E
Concentrations. *Int J Mol Sci*. 2017;18(11):2302. doi:10.3390/ijms18112302

35. Han X, Souzeau E, Ong JS, et al. Myocilin Gene Gln368Ter Variant Penetrance and
Association With Glaucoma in Population-Based and Registry-Based Studies. *JAMA
Ophthalmol*. 2019;137(1):28-35. doi:10.1001/jamaophthalmol.2018.4477

36. Gao XR, Huang H, Kim H. Polygenic Risk Score Is Associated With Intraocular Pressure
and Improves Glaucoma Prediction in the UK Biobank Cohort. *Transl Vis Sci Technol*.
2019;8(2):10. doi:10.1167/tvst.8.2.10

37. Nannini DR, Kim H, Fan F, Gao X. Genetic risk score is associated with vertical cup-disc
ratio and improves prediction of primary open angle glaucoma in Latinos. *Ophthalmology*.
2018;125(6):815-821. doi:10.1016/j.ophtha.2017.12.014

38. Zeleznik OA, Kang JH, Lasky-Su J, et al. Plasma metabolite profile for primary open-angle
glaucoma in three US cohorts and the UK Biobank. *Nat Commun*. 2023;14(1):2860.
doi:10.1038/s41467-023-38466-w

39. Madjedi KM, Stuart KV, Chua SY, et al. The Association between Serum Lipids and
Intraocular Pressure in 2 Large United Kingdom Cohorts. *Ophthalmology*. 2022;129(9):986.
doi:10.1016/j.ophtha.2022.04.023

40. Wang S, Bao X. Hyperlipidemia, Blood Lipid Level, and the Risk of Glaucoma: A Meta-
Analysis. *Invest Ophthalmol Vis Sci*. 2019;60(4):1028-1043. doi:10.1167/iovs.18-25845

41. Pertl L, Mossböck G, Wedrich A, et al. Triglycerides and Open Angle Glaucoma - A Meta-
analysis with meta-regression. *Sci Rep*. 2017;7(1):7829. doi:10.1038/s41598-017-08295-1

42. Rasoulinejad SA, Kasiri A, Montazeri M, et al. The Association Between Primary Open
Angle Glaucoma and Clustered Components of Metabolic Syndrome. *Open Ophthalmol J*.
2015;9:149-155. doi:10.2174/1874364101509010149

43. Williams PA, Casson RJ. Glycolysis and glucose metabolism as a target for bioenergetic and
neuronal protection in glaucoma. *Neural Regen Res*. 2024;19(8):1637. doi:10.4103/1673-
5374.389638

44. Harun-Or-Rashid M, Pappenhagen N, Zubricky R, Coughlin L, Jassim AH, Inman DM.
MCT2 Overexpression Rescues Metabolic Vulnerability and Protects Retinal Ganglion Cells
in Two Models of Glaucoma. *Neurobiol Dis*. 2020;141:104944.
doi:10.1016/j.nbd.2020.104944

45. Pappenhagen N, Harun-Or-Rashid M, Jassim Jaboori A, Inman DM. Upregulation of
monocarboxylate transporter 2 protects retinal ganglion cells in glaucoma. *Invest Ophthalmol*
Vis Sci. 2019;60(9):622.

46. Vohra R, Aldana BI, Bulli G, et al. Lactate-Mediated Protection of Retinal Ganglion Cells. *J*
Mol Biol. 2019;431(9):1878-1888. doi:10.1016/j.jmb.2019.03.005

47. Rajala RVS, Rajala A. Unlocking the role of lactate: metabolic pathways, signaling, and gene
regulation in postmitotic retinal cells. *Front Ophthalmol*. 2023;3.
doi:10.3389/fopht.2023.1296624

48. Williams PA, Harder JM, John SWM. Glaucoma as a metabolic optic neuropathy: making
the case for nicotinamide treatment in glaucoma. *J Glaucoma*. 2017;26(12):1161-1168.
doi:10.1097/IJG.0000000000000767

49. Inman DM, Harun-Or-Rashid M. Metabolic Vulnerability in the Neurodegenerative Disease
Glaucoma. *Front Neurosci*. 2017;11. doi:10.3389/fnins.2017.00146

50. Williams PA, Harder JM, Foxworth NE, et al. Vitamin B3 modulates mitochondrial
vulnerability and prevents glaucoma in aged mice. *Science*. 2017;355(6326):756-760.
doi:10.1126/science.aal0092

51. Chen S da, Wang L, Zhang X lan. Neuroprotection in glaucoma: present and future. *Chin*
Med J (Engl). 2013;126(8):1567-1577.

52. Vohra R, Dalgaard LM, Vibæk J, et al. Potential metabolic markers in glaucoma and their
regulation in response to hypoxia. *Acta Ophthalmol (Copenh)*. 2019;97(6):567-576.
doi:10.1111/aos.14021

53. Brooks GA. Lactate as a fulcrum of metabolism. *Redox Biol*. 2020;35:101454.
doi:10.1016/j.redox.2020.101454

54. Arnold PK, Finley LWS. Regulation and function of the mammalian tricarboxylic acid cycle.
J Biol Chem. 2022;299(2):102838. doi:10.1016/j.jbc.2022.102838

55. Jiménez-Moreno N, Kollareddy M, Stathakos P, et al. ATG8-dependent LMX1B-autophagy
crosstalk shapes human midbrain dopaminergic neuronal resilience. *J Cell Biol*.
2023;222(5):e201910133. doi:10.1083/jcb.201910133

56. Kim SW, Lee HK, Shin JH, Lee JK. Up-down Regulation of HO-1 and iNOS Gene
Expressions by Ethyl Pyruvate via Recruiting p300 to Nrf2 and Depriving It from p65. *Free*
Radic Biol Med. 2013;65:468-476. doi:10.1016/j.freeradbiomed.2013.07.028

57. Shin JH, Kim SW, Jin Y, Kim ID, Lee JK. Ethyl pyruvate-mediated Nrf2 activation and
hemeoxygenase 1 induction in astrocytes confer protective effects via autocrine and paracrine
mechanisms. *Neurochem Int*. 2012;61(1):89-99. doi:10.1016/j.neuint.2012.04.005

58. Harder JM, Guymer C, Wood JPM, et al. Disturbed glucose and pyruvate metabolism in
glaucoma with neuroprotection by pyruvate or rapamycin. *Proc Natl Acad Sci*.
2020;117(52):33619-33627. doi:10.1073/pnas.2014213117

59. Guymer C, Chidlow G, Wood JPM, Casson RJ. Oral pyruvate supplementation protects
against neurodegeneration in a rat model of glaucoma. *Invest Ophthalmol Vis Sci*.
2018;59(9):2618.

60. De Moraes CG, John SWM, Williams PA, Blumberg DM, Cioffi GA, Liebmann JM.
Nicotinamide and Pyruvate for Neuroenhancement in Open-Angle Glaucoma. *JAMA*
Ophthalmol. 2022;140(1):11-18. doi:10.1001/jamaophthalmol.2021.4576

61. Wu X, Dai H, Liu L, et al. Citrate reduced oxidative damage in stem cells by regulating
cellular redox signaling pathways and represent a potential treatment for oxidative stress-
induced diseases. *Redox Biol*. 2019;21:101057. doi:10.1016/j.redox.2018.11.015

62. Morganti C, Bonora M, Marchi S, et al. Citrate Mediates Crosstalk between Mitochondria
and the Nucleus to Promote Human Mesenchymal Stem Cell In Vitro Osteogenesis. *Cells*.
2020;9(4):1034. doi:10.3390/cells9041034

63. Catalina-Rodriguez O, Kolukula VK, Tomita Y, et al. The mitochondrial citrate transporter,
CIC, is essential for mitochondrial homeostasis. *Oncotarget*. 2012;3(10):1220-1235.
doi:10.18632/oncotarget.714

64. Ohanele C, Peoples JN, Karlstaedt A, et al. The mitochondrial citrate carrier SLC25A1
regulates metabolic reprogramming and morphogenesis in the developing heart. *Commun*
Biol. 2024;7(1):1-16. doi:10.1038/s42003-024-07110-8

65. Chhimpa N, Singh N, Puri N, Kayath HP. The Novel Role of Mitochondrial Citrate Synthase
and Citrate in the Pathophysiology of Alzheimer's Disease. *J Alzheimers Dis*. 94(Suppl
1):S453-S472. doi:10.3233/JAD-220514

66. Fraenkl SA, Muser J, Groell R, et al. Plasma Citrate Levels as a Potential Biomarker for
Glaucoma. *J Ocul Pharmacol Ther*. 2011;27(6):577-580. doi:10.1089/jop.2011.0062

67. Michalczuk M, Tadeusz P, Urban B, Anna W, Łazarczyk AB. Plasma citrate concentration: a
possible biomarker for glaucoma in children. *BMJ Paediatr Open*. 2017;1(1):e000023.
doi:10.1136/bmjpo-2017-000023

68. Nusinovici S, Li H, Thakur S, et al. High-Density Lipoprotein 3 Cholesterol and Primary
Open-Angle Glaucoma: Metabolomics and Mendelian Randomization Analyses.
Ophthalmology. 2022;129(3):285-294. doi:10.1016/j.ophtha.2021.09.013

69. Masson EAY, Serrano J, Leger-Charnay E, Acar N. Cholesterol and oxysterols in retinal
neuron-glia interactions: relevance for glaucoma. *Front Ophthalmol*. 2024;3:1303649.
doi:10.3389/fopht.2023.1303649

70. Increased High-Density Lipoprotein Levels Associated with Age-Related Macular
Degeneration: Evidence from the EYE-RISK and European Eye Epidemiology Consortia -
ScienceDirect. Accessed August 3, 2024.
https://www.sciencedirect.com/science/article/pii/S0161642018310911

71. Park SJ, Park DH. REvisiting Lipids in REtinal Diseases: A Focused Review on Age-related
Macular Degeneration and Diabetic Retinopathy. *J Lipid Atheroscler*. 2020;9(3):406.
doi:10.12997/jla.2020.9.3.406

72. Pasquale LR, Khawaja AP, Wiggs JL, et al. Metabolite and Lipid Biomarkers Associated
With Intraocular Pressure and Inner Retinal Morphology: [1] H NMR Spectroscopy Results
From the UK Biobank. *Investig Opthalmology Vis Sci*. 2023;64(11):11.
doi:10.1167/iovs.64.11.11

73. Ana R da, Gliszczyńska A, Sanchez-Lopez E, et al. Precision Medicines for Retinal Lipid
Metabolism-Related Pathologies. *J Pers Med*. 2023;13(4):635. doi:10.3390/jpm13040635

74. Fu Z, Chen CT, Cagnone G, et al. Dyslipidemia in retinal metabolic disorders. *EMBO Mol
Med*. 2019;11(10):e10473. doi:10.15252/emmm.201910473

75. Yuksel S, Ufret-Vincenty R, Butovich IA. The role of Lipe in lipid metabolism in the retina.
Invest Ophthalmol Vis Sci. 2023;64(8):5454.

76. Kelly UL, Grigsby D, Cady MA, et al. High-density lipoproteins are a potential therapeutic
target for age-related macular degeneration. *J Biol Chem*. 2020;295(39):13601-13616.
doi:10.1074/jbc.RA119.012305

77. Morvaridzadeh M, Zoubdane N, Heshmati J, Alami M, Berrougui H, Khalil A. High-Density
Lipoprotein Metabolism and Function in Cardiovascular Diseases: What about Aging and
Diet Effects? *Nutrients*. 2024;16(5):653. doi:10.3390/nu16050653

78. Hamel AR, Yan W, Rouhana JM, et al. Integrating genetic regulation and single-cell
expression with GWAS prioritizes causal genes and cell types for glaucoma. *Nat Commun*.
2024;15:396. doi:10.1038/s41467-023-44380-y

79. Minassian DC, Reidy A, Coffey M, Minassian A. Utility of predictive equations for
estimating the prevalence and incidence of primary open angle glaucoma in the UK. *Br J
Ophthalmol*. 2000;84(10):1159-1161. doi:10.1136/bjo.84.10.1159

80. Sudlow C, Gallacher J, Allen N, et al. UK Biobank: An Open Access Resource for
Identifying the Causes of a Wide Range of Complex Diseases of Middle and Old Age. *PLoS
Med*. 2015;12(3):e1001779. doi:10.1371/journal.pmed.1001779

81. Bycroft C, Freeman C, Petkova D, et al. The UK Biobank resource with deep phenotyping
and genomic data. *Nature*. 2018;562(7726):203-209. doi:10.1038/s41586-018-0579-z

82. Zeleznik O, Kang JH, Lasky-Su J, et al. Plasma metabolomics of primary open-angle
glaucoma in three prospective US cohorts and the UK Biobank. Published online February 28,
2022. doi:10.1101/2022.02.24.22271483

83. Elliott P, Peakman TC, on behalf of UK Biobank. The UK Biobank sample handling and
storage protocol for the collection, processing and archiving of human blood and urine. *Int J*
Epidemiol. 2008;37(2):234-244. doi:10.1093/ije/dym276

84. Julkunen H, Cichońska A, Slagboom PE, Würtz P, Nightingale Health UK Biobank
Initiative. Metabolic biomarker profiling for identification of susceptibility to severe
pneumonia and COVID-19 in the general population. Janus ED, Barton M, Karunajeewa H,
Janus ED, eds. *eLife*. 2021;10:e63033. doi:10.7554/eLife.63033

85. Takis PG, Ghini V, Tenori L, Turano P, Luchinat C. Uniqueness of the NMR approach to
metabolomics. *TrAC Trends Anal Chem*. 2019;120:115300. doi:10.1016/j.trac.2018.10.036

86. Soininen P, Kangas AJ, Würtz P, Suna T, Ala-Korpela M. Quantitative serum nuclear
magnetic resonance metabolomics in cardiovascular epidemiology and genetics. *Circ*
Cardiovasc Genet. 2015;8(1):192-206. doi:10.1161/CIRCGENETICS.114.000216

87. Mihaleva VV, Korhonen SP, van Duynhoven J, Niemitz M, Vervoort J, Jacobs DM.
Automated quantum mechanical total line shape fitting model for quantitative NMR-based
profiling of human serum metabolites. *Anal Bioanal Chem*. 2014;406(13):3091-3102.
doi:10.1007/s00216-014-7752-5

88. Kim J, Aschard H, Kang JH, et al. Intraocular Pressure, Glaucoma, and Dietary Caffeine
Consumption: A Gene—Diet Interaction Study from the UK Biobank. *Ophthalmology*.
2020;128(6):866. doi:10.1016/j.ophtha.2020.12.009

89. Gao X, Starmer J, Martin ER. A multiple testing correction method for genetic association
studies using correlated single nucleotide polymorphisms. *Genet Epidemiol*. 2008;32(4):361-
369. doi:10.1002/gepi.20310

90. Thaung C, West K, Clark BJ, et al. Novel ENU-induced eye mutations in the mouse: models
for human eye disease. *Hum Mol Genet*. 2002;11(7):755-767. doi:10.1093/hmg/11.7.755

91. John SW, Hagaman JR, MacTaggart TE, Peng L, Smithes O. Intraocular pressure in inbred
mouse strains. *Invest Ophthalmol Vis Sci*. 1997;38(1):249-253.

92. Savinova OV, Sugiyama F, Martin JE, et al. Intraocular pressure in genetically distinct mice:
an update and strain survey. *BMC Genet*. 2001;2:12. doi:10.1186/1471-2156-2-12

93. Howell GR, Libby RT, Jakobs TC, et al. Axons of retinal ganglion cells are insulted in the
optic nerve early in DBA/2J glaucoma. *J Cell Biol*. 2007;179(7):1523-1537.
doi:10.1083/jcb.200706181

94. Nair KS, Cosma M, Raghupathy N, et al. YBR/EiJ mice: a new model of glaucoma caused
by genes on chromosomes 4 and 17. *Dis Model Mech*. 2016;9(8):863-871.
doi:10.1242/dmm.024307

95. Howell GR, Soto I, Zhu X, et al. Radiation treatment inhibits monocyte entry into the optic
nerve head and prevents neuronal damage in a mouse model of glaucoma. *J Clin Invest*.
2012;122(4):1246-1261. doi:10.1172/JCI61135

96. Howell GR, Macalinao DG, Sousa GL, et al. Molecular clustering identifies complement and
endothelin induction as early events in a mouse model of glaucoma. *J Clin Invest*.
2011;121(4):1429-1444. doi:10.1172/JCI44646

Figures

936
937
938   **Figure 1. Participant flow chart describing inclusion and exclusion criteria from the UK**
939   **Biobank**

940
Figure 2. Study design from the UK Biobank. (a) 117,698 individuals had metabolomics data
available from the UK Biobank, which was divided into a training and test set to formulate a
metabolic risk score (MRS) model. (b) The inclusion of metabolites (either 168 metabolites on
the NMR platform or a subset of 27 metabolites with European Union (EU) certification) in
relation to prevalent glaucoma risk prediction was studied. (c) A histogram showing the
polygenic risk score (PRS) distribution is shown. Overall, 4,658 cases and 113,040 individuals
without glaucoma are available for analysis. The metabolomic signature of resilience to the top
10% of glaucoma PRS was assessed among 1,693 cases (14.4%) and 10,077 individuals without
glaucoma (85.6%). (d) Interactions of prevalent glaucoma with MRS and PRS quartiles were
examined.

Figure 3. Inclusion of metabolite data into glaucoma prediction algorithms. Model 1 includes metabolites only; Model 2 incorporates additional covariates including age, sex, genetic ancestry, season, time of day of specimen collection, and fasting time; Model 3 incorporates covariates in Model 2 and smoking status (never, past, and current smoker), alcohol intake (g/week), caffeine intake (mg/day), physical activity (metabolic equivalent of task [MET], hours/week), body mass index (kg/m^2), average systolic blood pressure (mm Hg), history of diabetes, HbA1c (mmol/mol), history of coronary artery disease, systemic beta-blocker use, oral steroid use, and spherical equivalent refractive error (diopters); Model 4 incorporates covariates in model 3 and a glaucoma polygenic risk score (PRS). Each color represents a different panel of metabolites (grey = no metabolites; light blue = 27 metabolites; and dark blue = 168 metabolites). The white text represents the AUC ± 95% confidence interval.

Abbreviations: ROC, receiver operator curve; AUC, area under the curve; EU, European Union.

A.

	Bottom 50%	Top 10%
PRS	[-0.0090,-0.0073]	[-0.0068,-0.0055]
n	58,358	11,770
Glaucoma cases	780 (1.3%)	1,693 (14.4%)
No Glaucoma	57,578	10,077

Figure 4. The distribution of glaucoma cases and no glaucoma stratified by polygenic risk score (PRS) deciles. Participants were divided into the bottom 50% and top 10% based on their glaucoma PRS, where the prevalence of glaucoma cases from (A) the bottom 50% (n=58,358) was 1.3% and from the top 10% (n=11,770) was 14.4%. (B) Box plot illustrating the distribution of participants with glaucoma (top) and no glaucoma (bottom) as a function of PRS decile. The blue line denotes participants at the bottom 50% of glaucoma PRS, the red line highlights the participants at the top decile of glaucoma PRS, and the red box represents the participants resilient to glaucoma despite high PRS.

A.

Quartiles	Q1	Q2	Q3	Q4
PRS	-0.0090, -0.0075	-0.0075, -0.0073	-0.0073, -0.0070	-0.0070, -0.0055
Metabolite sum	-8.2, -1.5	-1.5, -0.2	-0.2, 1.3	1.3, 32.2

B.

	Glaucoma cases	Glaucoma Metabolite Reference	Odds Ratio (95% CI)	p-value
PRS Q1				
Metabolite sum Q2	66	55	1.18 (0.82, 1.69)	0.42
Metabolite sum Q3	61	55	1.1 (0.77, 1.60)	0.66
Metabolite sum Q4	57	55	1.0 (0.69, 1.46)	1.00
PRS Q2				
Metabolite sum Q2	138	161	0.87 (0.69, 1.10)	0.27
Metabolite sum Q3	128	161	0.79 (0.63, 1.00)	0.057
Metabolite sum Q4	127	161	0.80 (0.63, 1.01)	0.069
PRS Q3				
Metabolite sum Q2	266	297	0.90 (0.76, 1.06)	0.23
Metabolite sum Q3	228	297	0.78 (0.65, 0.93)	0.0059
Metabolite sum Q4	219	297	0.74 (0.62, 0.88)	0.0009
PRS Q4				
Metabolite sum Q2	750	793	0.93 (0.84, 1.03)	0.19
Metabolite sum Q3	733	793	0.90 (0.81, 1.00)	0.054
Metabolite sum Q4	579	793	0.71 (0.64, 0.80)	<0.0001

Figure 5. Interaction of three putative resilient metabolites (lactate, pyruvate, and citrate) and polygenic risk score (PRS) on glaucoma risk. (A) The bar chart shows the interaction of resilient probit-transformed metabolite sum with glaucoma genetic predisposition in each PRS quartile. In each glaucoma PRS quartile, the lowest metabolic sum quartile (Q1) is the metabolite reference group used to calculate the odds ratios. Each color represents resilient-metabolite sum quartiles (red = second quartile; blue = third quartile; and green = fourth quartile). Error bars show 95% confidence interval (CI). The table under the bar chart shows the ranges for the PRS and metabolite sum value quartiles. (B) The table shows odds ratios for glaucoma by PRS and putative resilient metabolite sum within various quartiles. The number of glaucoma cases within each resilient metabolite sum quartile and the number of glaucoma cases in the first quartile of resilient metabolite sum (Q1, labeled as glaucoma metabolite reference) are used to calculate the odd ratios. This analysis is adjusted for time since the last meal/drink (hours), age, age-squared, sex, ethnicity (Asian, Black, White, and other), season, time of day of specimen collection (morning, afternoon, night), smoking status (never, past, and current smoker), alcohol intake, caffeine intake, physical activity (metabolic equivalent of task [MET] hours/week), body mass index (kg/m^2), average systolic blood pressure (mm Hg), history of diabetes (yes or no), HbA1c (mmol/mol), history of coronary artery disease, systemic beta-blocker use, oral steroid use, and spherical equivalent refractive error (diopters).

Figure 6. Interaction of the holistic metabolite risk score (n=168 metabolites) and polygenic risk score (PRS) on glaucoma risk. (A) The bar chart plots the odds ratio of glaucoma as a function of holistic probit-transformed MRS quartile with further stratification by glaucoma PRS in each MRS bin. The lowest quartile of glaucoma PRS and MRS is the reference group (see dotted red line) for the entire population. Each color represents the MRS quartiles (red = first quartile; blue = second quartile; green = third quartile; and purple = fourth quartile). Error bars show 95% confidence interval (CI). The table under the bar chart shows the ranges for the PRS and MRS quartiles. (B) Table showing odds ratios for glaucoma by polygenic risk score (PRS) and MRS within various quartiles. The number of glaucoma cases within each MRS and the number of glaucoma cases in PRS Q1 and MRS Q1 are used to calculate the odds ratios. This analysis is adjusted for time since the last meal/drink (hours), age, age-squared, sex, ethnicity (Asian, Black, White, and other), season, time of day of specimen collection (morning, afternoon, night), smoking status (never, past, and current smoker), alcohol intake, caffeine intake, physical activity (metabolic equivalent of task [MET] hours/week), body mass index (kg/m^2), average systolic blood pressure (mm Hg), history of diabetes (yes or no), HbA1c (mmol/mol), history of coronary artery disease, systemic beta-blocker use, oral steroid use, and spherical equivalent refractive error (diopters).

Figure 7. Pyruvate treatment protects from IOP elevation and glaucoma. (A) Representative photos of eyes from mice of the indicated genotypes and treatments (Unt = untreated, Pyr = pyruvate treated). *Lmx1b* is expressed in the iris and cornea, so *Lmx1b*^{V265D} mutant eyes have primary abnormalities of the iris and cornea. This includes corneal haze, which is present before IOP elevation in many eyes and likely reflects a direct transcriptional role of LMX1B in collagen gene expression. *Lmx1b*^{V265D} mutant eyes also develop anterior chamber deepening (ACD), a sensitive indicator of IOP elevation in mice. The WT and pyruvate-treated mutant eyes have shallow anterior chambers, while the untreated mutant eye has a deepened chamber (arrowheads). (B) Distributions of ACD are based on a previously defined scoring system.[31] Groups are compared by Fisher's exact test. n > 30 eyes were examined in each group. (C) Boxplots of IOP (interquartile range and median line) in WT and mutant eyes. Pyruvate treatment significantly lessens IOP elevation in mutants compared to untreated mutant controls. Groups were compared by ANOVA followed by Tukey's honestly significant difference. n > 30 eyes were examined in each *Lmx1b*^{V265D} mutant group, and n > 20 eyes were examined in WT groups. (D) Distributions of damage based on analysis of PPD-stained optic nerve cross sections from 6-month-old mice (Methods). Pyruvate treatment lessened the incidence of glaucoma (Fisher's exact test). No glaucoma was found in WT mice. Geno = genotype. n > 40 nerves examined per group. NOE = no glaucoma, MOD = moderate. SEV = severe, and V. SEV = very severe (see Methods).

 Tables

1031 **Table 1. Demographic and clinical characteristics of the UK Biobank study population**
1032 **assessed in 2006-2010**

1033

Characteristic	No Glaucoma	Glaucoma	P-value
Sample Size (%)	113,040 (96.0)	4,658 (4.0)	
Sex - Male (%)	52,497 (46.4)	2,493 (53.5)	<0.001
Age at recruitment (mean (SD))	56.7 (8.0)	60.9 (6.6)	<0.001
Ethnicity (%)			0.0010
White	105,912 (94)	4,331 (93)	
Asian	2,977 (2.6)	127 (2.7)	
Black	2,228 (2.0)	130 (2.8)	
Other	1,923 (1.7)	70 (1.5)	
Genetic Ancestry (%)			<0.001
African	1,911 (1.7)	123 (2.7)	
AMR	261 (0.2)	2 (0.0)	
Asian	3,125 (2.8)	134 (2.9)	
European	96,991 (86.6)	4,001 (86.7)	
Smoking Status (%)			<0.001
Never	63,240 (56)	2,339 (52)	
Prefer not to answer	430 (0.4)	16 (0.4)	
Previous	38,926 (34)	1,732 (39)	
Current	10,605 (9.4)	410 (9.1)	
Total Cholesterol, mmol/l (median [IQR])	4.6 [3.98, 5.22]	4.5 [3.88, 5.18]	<0.001
Physical Activity, MET-minutes per week (mean (SD))	2,465 (2430)	2,458 (2,438)	0.84
Body Mass Index, kg/m^2 (mean (SD))	27.4 (4.8)	27.7 (4.7)	<0.001
HbA1c, mmol/mol (mean (SD))	36.0 (5.8)	37.4 (6.8)	<0.001
Spherical Equivalent, diopter (mean (SD))	-0.1 (2.1)	-0.1 (1.9)	0.67
Intraocular pressure, mmHg (mean (SD))	15.9 (2.6)	17.8 (4.5)	<0.001
mRNFL thickness, μm (mean (SD))	28.7 (1.8)	28.5 (1.3)	<0.001
Beta blocker use (%)	8,398 (7.4)	447 (9.6)	<0.001
Caffeine intake, mg/day (mean (SD))	165 (67)	168 (59)	0.0020
Alcohol intake, g/week (median [IQR])	84 [40.3, 146.9]	84 [48.0, 154.6]	<0.001
Diabetes (%)	6,512 (5.8)	463 (9.9)	<0.001
Oral steroid use (%)	3,043 (2.7)	184 (4.0)	<0.001
Coronary Artery Disease (%)	5,153 (4.6)	377 (8.1)	<0.001

1034

1035 Abbreviations: SD, standard deviation; IQR, interquartile range; AMR, mixed American; MET
1036 metabolic equivalent of task mRNFL, macular retinal nerve fiber layer

Table 2. Stratification of glaucoma by ethnicity, age, and gender for predictive assessment with and without using metabolite data.

Stratification	Sample size	Glaucoma cases	No Metabolites (AUC)	Metabolites (AUC)	P-value
Ethnicity					
White	110,243	4,331	0.675	0.686	<0.001
Asian	3,104	127	0.780	0.768	0.062
Black	2,358	130	0.704	0.706	0.52
Age					
<55 years	43,648	788	0.576	0.566	0.52
≥55 years	74,050	3,870	0.569	0.596	0.002
Gender					
Female	62,708	2,165	0.689	0.696	0.25
Male	54,990	2,493	0.659	0.673	0.002

The area under the curve (AUC) of the receiver operating characteristic (ROC) curve was calculated for each demographic stratification to evaluate the predictive performance of models both with and without metabolite data. Differences in model AUC were tested using the DeLong test, and p-values were reported. For models excluding metabolite data, the predictors include as appropriate, age, sex, genetic ancestry, season, time of day of specimen collection, fasting time (hours), smoking status (never, past, and current smoker), alcohol intake (g/week), caffeine intake (mg/day), physical activity (metabolic equivalent of task [MET], hours/week), body mass index (kg/m^2), average systolic blood pressure (mm Hg), history of diabetes, HbA1c (mmol/mol), history of coronary artery disease, systemic beta-blocker use, oral steroid use, and spherical equivalent refractive error (diopters). Models including metabolite data incorporated the same predictors with the addition of the 168 metabolite measurements.

1052 **Table 3. Demographic and clinical characteristics in 2006-2010 of participants among the**
1053 **top 10% of glaucoma polygenic risk score (PRS).**
1054

Characteristic	No Glaucoma	Glaucoma	P-value
Sample size (%)	10,077 (85.6)	1,693 (14.4)	
Sex - Male (%)	4,667 (46.3)	893 (52.7)	<0.001
Age at recruitment (mean (SD))	56.3 (8.0)	61.0 (6.3)	<0.001
Ethnicity (%)			<0.001
White	9,256 (91.9)	1,636 (96.6)	
Asian	279 (2.8)	24 (1.4)	
Black	345 (3.4)	20 (1.2)	
Other	197 (2.0)	13 (0.8)	
Smoking Status (%)			0.0030
Never	5,806 (57.6)	899 (53.1)	
Prefer not to answer	48 (0.5)	5 (0.3)	
Previous	3,402 (33.8)	640 (37.8)	
Current	821 (8.1)	149 (8.8)	
Physical Activity, MET-minutes per week (mean (SD))	2,439 (2,352)	2,514 (2,480)	0.23
Body mass index, kg/m^2 (mean (SD))	27.4 (4.7)	27.7 (4.8)	0.0080
HbA1c, mmol/mol (mean (SD))	36.00 (5.9)	37.0 (6.1)	<0.001
Spherical Equivalent, diopter (mean (SD))	-0.3 (2.2)	-0.2 (1.9)	0.0090
Intraocular pressure, mmHg (mean (SD))	17.2 (3.0)	18.3 (5.0)	<0.001
mRNFL thickness, μm (mean (SD))	28.7 (1.8)	28.5 (1.2)	<0.001
Beta blocker use (%)	719 (7.1)	179 (10.6)	<0.001
Caffeine intake, mg/day (mean (SD))	164 (65)	170 (60)	0.0010
Alcohol intake, g/week (median [IQR])	84 [38.6, 145.3]	84 [51.5, 155.3]	0.002
Diabetes mellitus (%)	607 (6.0)	151 (8.9)	<0.001
Oral steroid use (%)	292 (2.9)	59 (3.5)	0.22
Coronary Artery Disease (%)	443 (4.4)	148 (8.7)	<0.001
Total cholesterol, mmol/l (median [IQR])	4.61 [4.00, 5.25]	4.54 [3.90, 5.21]	0.006
Lactate, mmol/l (median [IQR])	3.96 [3.27, 4.78]	3.79 [3.16, 4.58]	<0.001
Pyruvate, mmol/l (median [IQR])	0.080 [0.06, 0.10]	0.077 [0.06, 0.09]	<0.001
Citrate, mmol/l (median [IQR])	0.065 [0.06, 0.07]	0.065 [0.06, 0.07]	0.87
Cholesteryl Esters in Small HDL, mmol/l (median [IQR])	0.33 [0.30, 0.36]	0.33 [0.30, 0.36]	0.005
Triglycerides in Very Large VLDL, mmol/l (median [IQR])	0.091 [0.05, 0.15]	0.10 [0.05, 0.17]	<0.001
Alanine, mmol/l (median [IQR])	0.29 [0.24, 0.35]	0.29 [0.24, 0.35]	0.58
Triglycerides in Chylomicrons and extremely Large VLDL, mmol/l (median [IQR])	0.088 [0.03, 0.18]	0.10 [0.04, 0.21]	<0.001
Acetoacetate, mmol/l (median [IQR])	0.010 [0.01, 0.02]	0.011 [0.01, 0.02]	<0.001
Cholesteryl Esters in Medium HDL, mmol/l (median [IQR])	0.41 [0.34, 0.48]	0.40 [0.34, 0.47]	0.003
Triglycerides in Large VLDL, mmol/l (median [IQR])	0.15 [0.09, 0.22]	0.16 [0.10, 0.23]	<0.001
Cholesterol in Medium HDL, mmol/l (median [IQR])	0.49 [0.42, 0.58]	0.48 [0.41, 0.57]	0.005

1055   Abbreviations: SD, standard deviation; IQR, interquartile range; MET, metabolic equivalent of
1056   task HbA1c, hemoglobin A1C; mRNFL, macular retinal nerve fiber layer; HDL, high-density
1057   lipoprotein; VLDL, very low-density lipoprotein
1058

1059 **Table 4. Demographic and clinical characteristics in 2006-2010 of participants among the**
1060 **bottom 50% of glaucoma polygenic risk score (PRS).**
1061

Characteristic	No Glaucoma	Glaucoma	P-value
Sample size (%)	57,578 (98.7)	780 (1.3)	
Sex - Male (%)	26,887 (46.7)	409 (52.4)	0.002
Age at recruitment (mean (SD))	57 (8.0)	60 (7.0)	<0.001
Ethnicity (%)			<0.001
White	55,034 (96)	690 (89)	
Asian	1,129 (2.0)	41 (5.3)	
Black	574 (1.0)	27 (3.5)	
Other	841 (1.5)	22 (2.8)	
Smoking Status (%)			0.67
Never	200 (0.3)	4 (0.5)	
Prefer not to answer	31,593 (54.9)	418 (53.6)	
Previous	20,276 (35.2)	287 (36.8)	
Current	5,509 (9.6)	71 (9.1)	
MET, minutes per week (mean (SD))	2,483 (2,450)	2,612 (2,562)	0.14
Body mass index kg/m^2 (mean (SD))	27.4 (4.8)	27.8 (4.7)	0.021
HbA1c, mmol/mol (mean (SD))	35.9 (5.7)	38.1 (7.8)	<0.001
Spherical Equivalent, diopter (mean (SD))	-0.03 (2.1)	0.01 (1.9)	0.63
Intraocular pressure, mmHg (mean (SD))	15.4 (2.4)	17.1 (3.9)	<0.001
mRNFL thickness, μm (mean (SD))	28.7 (1.7)	28.6 (1.3)	0.02
Beta blocker use (%)	4,322 (7.5)	71 (9.1)	0.11
Caffeine intake, mg/day (mean (SD))	1,66.3 (67.4)	166.6 (61.5)	0.90
Alcohol intake, g/week (median [IQR])	84 [41.7, 149.3]	84 [42.5, 145.9]	0.91
Diabetes (%)	3209 (5.6)	104 (13.3)	<0.001
Oral steroid use (%)	1,538 (2.7)	39 (5.0)	<0.001
Coronary Artery Disease (%)	2,656 (4.6)	64 (8.2)	<0.001
Total cholesterol, mmol/l (median [IQR])	4.59 [3.98, 5.22]	4.49 [3.76, 5.13]	0.001
Lactate, mmol/l (median [IQR])	3.95 [3.24, 4.75]	3.84 [3.21, 4.69]	0.16
Concentration of Small HDL Particles, mmol/l (median [IQR])	0.0097 [0.0089, 0.011]	0.0095 [0.0088, 0.010]	0.002
Cholesteryl esters in small HDL, mmol/l (median [IQR])	0.33 [0.30, 0.36]	0.32 [0.30, 0.36]	0.001
Albumin, mmol/l (median [IQR])	39.4 [37.3, 41.45]	38.9 [36.7, 40.8]	<0.001
Total lipids in small HDL, mmol/l (median [IQR])	1.16 [1.06, 1.26]	1.15 [1.05, 1.25]	0.018
Citrate, mmol/l (median [IQR])	0.065 [0.057, 0.074]	0.065 [0.057, 0.073]	0.52
Pyruvate, mmol/l (median [IQR])	0.080 [0.06, 0.10]	0.079 [0.061, 0.098]	0.14
Alanine, mmol/l (median [IQR])	0.29 [0.24, 0.35]	0.29 [0.24, 0.35]	0.95
Phospholipids in Small HDL, mmol/l (median [IQR])	0.66 [0.60, 0.72]	0.66 [0.60, 0.71]	0.033

1062 Abbreviations: SD, standard deviation; MET, metabolic equivalents; HbA1c, hemoglobin A1C;
1063 mRNFL, macula region retinal nerve fiber layer; HDL, high density lipoprotein; VLDL, very
1064 low density lipoprotein
1065

1066 **Table 5. Metabolites associated with glaucoma among participants in the top decile and the**
1067 **bottom half of glaucoma polygenic risk score.**
1068

Top 10% of glaucoma polygenic risk score			
Metabolites (Probit score)	**Glaucoma**	**No Glaucoma**	**Adjusted P-value (NEF)**
Lactate	-0.146	0.0239	8.8E-12
Pyruvate	-0.137	0.0117	2.9E-10
Citrate	-0.0693	0.0079	0.018
Triglycerides in Very Large VLDL	0.0606	-0.0043	0.10
Triglycerides in Chylomicrons and Extremely Large VLDL	0.0572	-0.0076	0.11
Acetoacetate	0.0744	0.0011	0.11
Cholesteryl Esters in Small HDL	-0.0239	0.0376	0.12
Cholesteryl Esters in Medium HDL	-0.0261	0.0276	0.13
Alanine	-0.0612	-0.0006	0.14
Triglycerides in Large VLDL	0.0629	0.0005	0.15

1069
1070

Bottom 50% of glaucoma polygenic risk score			
Metabolites (Probit score)	**Glaucoma**	**No Glaucoma**	**Adjusted P-value (NEF)**
Albumin	-0.110	-0.0099	0.047
Concentration of Small HDL Particles	-0.0943	-0.0062	0.071
Cholesteryl Esters in Small HDL	-0.0925	-0.0055	0.085
Cholesterol in Small HDL	-0.0919	-0.0063	0.088
Omega-3 Fatty Acids	-0.0896	-0.0052	0.092
Docosahexaenoic Acid	-0.0876	-0.0061	0.10
Total Lipids in Small HDL	-0.0844	-0.0051	0.12
Lactate	-0.0823	0.0031	0.12
Citrate	-0.0882	-0.0023	0.16
Alanine	-0.0766	0.0011	0.18

Potential confounders adjusted by regression include time since the last meal/drink (hours), age,
age-squared, sex, ethnicity (Asian, Black, White, and other), season, time of day of specimen
collection (morning, afternoon, night), smoking status (never, past, and current smoker), alcohol
intake, caffeine intake, physical activity (metabolic equivalent of task [MET] hours/week), body
mass index (kg/m^2), average systolic blood pressure (mm Hg), history of diabetes (yes or no),
HbA1c (mmol/mol), history of coronary artery disease, systemic beta-blocker use, oral steroid
use, and spherical equivalent refractive error (diopters).

Financial Support:

This work is supported by NEI R01 015473 (LRP), NEI R01 032559 (LRP, JLW, AVS),

EY036460 (LRP, JHK), NEI P30 EY014104 (JLW, AVS), NEI R01EY031424 (AVS), NIGMS

R35-GM124836 (RD), NEI EY032507, EY032062, and EY018606 (SWMJ), an unrestricted

Challenge Grant from Research to Prevent Blindness (NYC), and The Glaucoma Foundation

(NYC). Further funding was provided by startup funds at Columbia University including the

Precision Medicine Initiative, and by the New York Fund for Innovation in Research and

Scientific Talent (NYFIRST; EMPIRE CU19-2660; SWMJ). This work is also supported by K23

(1K23EY032634) and the Research to Prevent Blindness Career Development Award (N.Z), a

Vision Core grant P30EY019007 (Columbia University) an unrestricted departmental award

from Research to Prevent Blindness (Columbia University). Supported by Fight for Sight (UK)

(1956A), and The Desmond Foundation (K.V.S.); UK Research and Innovation Future Leaders

Fellowship (MR/T040912/1), Alcon Research Institute Young Investigator Award and a Lister

Institute of Preventative Medicine Fellowship (A.P.K.); financial support from the UK

Department of Health through an award made by the National Institute for Health Research

(NIHR) to Moorfields Eye Hospital National Health Service (NHS) Foundation Trust and

University College London (UCL) Institute of Ophthalmology for a Biomedical Research Centre

(BRC) for Ophthalmology (A.P.K.). The sponsor or funding organizations had no role in the

design or conduct of this research. The content is solely the responsibility of the authors and does

not necessarily represent the official views of the National Institutes of Health.

Conflict of Interest:

The authors declare the following financial interests/personal relationships that may be

considered potential competing interests: LRP reports consulting honoraria from Twenty

Twenty. JLW reports consulting fees from Editas and CRISPR Therapeutics. NZ reports

consulting fees from Character Biosciences. APK has acted as a paid consultant or lecturer to

Abbvie, Aerie, Allergan, Google Health, Heidelberg Engineering, Novartis, Reichert, Santen,

Thea and Topcon. NV has acted as a paid consultant or lecturer to Santen, Thea, NovaEye,

Glaukos, and Elios. RD reports being a scientific co-founder, consultant, and equity holder for

Pensieve Health (pending) and being a consultant for Variant Bio and Character Bio. SWMJ is

founder of Myco Advising LLC, a founder of Qura Inc and advises Starlight Bio Inc with no

overlap to the currently presented research. SWMJ and NT are inventors on patents/patent

applications held by The Jackson Laboratory and Columbia University on the use of pyruvate to

treat glaucoma. No other disclosures were reported.