Integrative Gene Co-expression Network Analysis Reveals Protein-Coding and lncRNA Genes Associated with Alzheimer's Disease Pathology

Fatemeh Zebardast[1,2], Michael Peter Sascha Riethmüller[1], Katja Nowick[1] (corresponding author, katja.nowick@fu-berlin.de)

1 Human Biology and Primate Evolution Group; Institute for Zoology; Department of Biology, Chemistry and Pharmacy; Freie Universität Berlin, Berlin, Germany.
2 International Max Planck Research School for Biology and Computation IMPRS-BAC, Berlin, Germany

Abstract

Introduction

Alzheimer's disease (AD) is a complex neurodegenerative disorder involving widespread molecular disruptions, many of which remain poorly understood. While AD pathology progresses through distinct brain regions, it is unclear whether these regions are similarly affected. Long non-coding RNAs (lncRNAs) have emerged as key regulators in cellular processes, but their roles in AD remain largely unexplored. A comprehensive analysis of the complex molecular interactions underlying AD, particularly the roles of lncRNAs in their interactions with protein-coding genes and the comparison of networks across brain regions, could offer valuable insights into the disease's progression and underlying mechanisms.

Method

To address this, we applied consensus weighted gene co-expression network analysis using a meta-analytic approach to 396 postmortem brain RNA-seq samples to explore AD pathophysiology mechanisms.

Results

Our study revealed significant network rewiring in AD, with pronounced alterations in the temporal cortex compared to the frontal cortex. While the temporal cortex showed adaptive changes in gene interactions, the frontal cortex exhibited a loss of healthy correlations, potentially reflecting different levels of disease progression. We identified key players in the temporal cortex AD network, including 46 protein-coding genes and 27 lncRNAs. Using the established functions of protein-coding genes as anchors, we provided functional annotations for over 100 lncRNAs across two brain regions, identifying potential lncRNAs involved in AD pathology and highlighting their roles in both healthy and diseased states.

Conclusion

We provided novel insights into the molecular interactions underlying AD and introduced new candidate protein-coding and lncRNA genes for further experimental validation and therapeutic exploration.

Keywords: Alzheimer's disease, consensus gene co-expression network, weighted Topological Overlap (wTO), long non-coding RNAs (lncRNA), Random Walk with Restart (RWR).

Introduction

Alzheimer's disease (AD) is a complex, multifactorial neurodegenerative disorder characterized by progressive cognitive decline and underlying pathological mechanisms (1). Current knowledge indicates that the hallmark pathology of AD begins in the medial temporal lobe (2) and, as the disease progresses, extends to other areas of the temporal cortex before spreading to additional brain regions, including the frontal lobes, in the mid to late stages (2–4). Its pathology disrupts numerous cellular mechanisms, such as mitochondrial functions (5–7), endoplasmic reticulum (8,9), Golgi apparatus integrity (10,11), cellular transport processes (5,12–14), and synaptic signaling (15–18). These widespread dysfunctions closely align with

alterations in gene expression and molecular interactions, highlighting the complex biological

networks that underlie AD.

Non-coding RNAs have attracted attention due to their potential involvement in both normal

and disease-related cellular processes (19–22). Among these, long non-coding RNAs

(lncRNAs)—transcripts longer than 200 nucleotides without apparent protein-coding

capacity—have emerged as critical regulators of cellular functions (23–26). LncRNAs

introduce an additional layer of complexity to gene regulatory processes through their

interactions with protein-coding genes. Studies have implicated lncRNAs in several AD-

associated dysfunctions, including amyloid beta (Aβ) production (27–30), oxidative stress (31),

and synaptic impairments (32). Despite the annotation of numerous lncRNAs with known

sequences and genomic locations, only a fraction have been extensively studied to uncover

their functional roles in both healthy and disease conditions (24,26,33), leaving a significant

knowledge gap yet to be addressed. A promising approach to address this gap is the

construction of gene co-expression networks, which can reveal functional relationships

between genes (34). Current methodologies face challenges such as mitigating noise and

improving the reliability of gene-gene correlation detection (35,36). Overcoming these

limitations is essential for generating accurate and interpretable biological insights. To address

these challenges, we integrated gene co-expression networks derived from the same brain

region across independent studies to construct consensus networks. This approach

consolidates shared signals from multiple datasets, reducing the impact of dataset-specific

noise and variability. By focusing on interactions consistently observed across networks, this

method enhances the detection of biologically relevant signals, increasing confidence in their

accuracy and interpretability compared to signals supported by individual networks alone (37–

40).

In this study, we utilized RNA-seq data from two independent studies, comprising five datasets

with 396 individuals total in AD and control groups, with gene expression profiles from the

temporal cortex (TCX) and frontal cortex (FCX). These profiles were used to construct signed

weighted topological overlap (wTO) networks, which were subsequently integrated into consensus networks to enhance reliability and biological relevance (Figure 1). Our analysis focused on lncRNAs and protein-coding genes, examining their correlations within gene co-expression networks under both AD and non-AD (control) conditions. Comparative analysis revealed significant rewiring of gene-gene interactions in the temporal cortex in AD, contrasted with a marked loss of 'healthy' gene correlations in the frontal cortex. Building on these observations, we identified 46 protein-coding genes and 27 lncRNAs as critical components of the AD network. A key objective of this study was to predict the functions of lncRNAs and their associations with AD pathology. By leveraging the established functions of protein-coding genes as network anchors, we assigned over 100 lncRNAs to biological processes in both AD and control conditions.

Figure 1: Overview and illustration of the current study (41)**.** From left to right, the top panel illustrates the network construction and integration process. The bottom panel illustrates the network analysis, including topology assessment, identification of hub genes, functional enrichment analysis, assigning enriched GOs to lncRNAs within each cluster, and identifying key genes associated with AD.

Material and Methods

RNA-seq datasets

We downloaded the FASTQ files and metadata from the Synapse platform (42). A total of five RNA-seq datasets from two independent studies were used. The first study, known as the Mayo RNA-seq data (43) includes RNA-seq data for the temporal cortex human brain region. The FASTQ files for this study were downloaded from the Synapse repository with the Synapse ID syn3163039 (44). The second dataset, known as the Mount Sinai Brain Bank (MSBB) study (45), includes gene expression profiles for four brain regions: Brodmann area (BM) 36, BM44, BM22, and BM10. This dataset was downloaded from the Synapse repository with the Synapse ID syn3159438 (46). All samples were collected under precise clinical-pathological diagnoses and distinctive histologies, including Braak NFT stage (47), plaque density, CERAD (Consortium to Establish a Registry for Alzheimer's Disease) (48), and CDR (clinical dementia rating) (49). Each AD and the corresponding control dataset from both brain areas consisted of an equal number of individuals across sexes with identical gene sets. Furthermore, these datasets have been carefully balanced for age, disease stage, and other potential confounders. All data underwent standardized preprocessing steps, including filtering based on library size and gene expression levels, to ensure uniformity across the datasets.

Preprocessing of RNA-seq data

To ensure uniform data processing, we employed a unified pipeline that included standard alignment, quality control, and gene expression analyses on each dataset (Mayo and MSBB RNA-seq). The preprocessing steps included FASTQ quality control, adapter trimming, base quality control, and filtering out short reads. We excluded all FASTQ files with fewer than 12 million reads. Following this, reads were mapped to the human reference genome (hg38) using STAR (v 2.7.10b) (50). Samples with less than 60% unique mapped reads were also

excluded from further analysis. We used the `featureCounts()`function from the Subread

software (V 1.6.3) (51) for gene quantification, with the following parameters: `-s 0 -t gene`

`-g gene_id -M -O --fraction`. `-s 0` (un-stranded data), `-t gene` (gene feature

type), `-g gene_id` (grouped by gene ID), `-M` (allow multi-mapping reads), `-O` (count

overlapping reads), and `--fraction` (fractional assignment of multi-mapping reads). To

account for both sense and antisense lncRNAs, reads were mapped specifically to the gene

feature type (`-t gene`) and counted fractionally by `countMultiMappingReads` arguments

(`--fraction`).   Noteworthy,   the   human   gene   annotation   file   version   103

(*Homo_sapiens.GRCh38.103.gtf*) from Ensembl was used for gene annotation. Notably, both

datasets contained un-stranded sequence data, which required the use of the `-s 0` option.

To remove systematic effects in the RNA-seq data, we normalized all gene count data using

the `DESeq2` R package (52), which normalizes for both sequencing depth and RNA

composition. Moreover, genes with low expression levels can introduce noise into the

correlation calculation process. Hence, we excluded genes with low expression levels to

enhance the reliability of the correlation analysis. The criteria for excluding low-expressed

genes were set such that only genes with a count of at least 1 count per million (cpm) in 70%

of the samples were retained. The use of cpm avoids favoring genes expressed in larger

libraries.

Signed weighted co-expression network

Gene co-expression networks represent gene-gene interactions, with nodes representing

genes and edges indicating the strength of their co-expression (53). The weight of an edge

reflects the strength of the co-expression relationship between two genes, while the sign

distinguishes between positive and negative correlations, showing whether genes have similar

or opposing expression patterns. The signed weighted gene co-expression networks were

constructed using the weighted Topological Overlap (wTO) method separately for each RNA-

seq dataset. Using the wTO method, link weights for connected nodes are calculated through

an averaging process that accounts for all their common connections (54). The wTO

calculation was performed using the `wTO` R package. In summary, for each dataset, Pearson's

correlation coefficient (ρ) was calculated for all pairs of genes to generate the adjacency matrix

A = [$a_{i,j}$], where each element $a_{i,j}$ represents the correlation between gene i and gene j. The

weighted Topological Overlap ($\omega_{i,j}$) was then calculated for each pair of nodes i and j, reflecting

the shared network connectivity between the two genes. Gene co-expression correlations with

$|\omega i,j| < 0.5$ were discarded. This threshold was chosen to balance the inclusion of meaningful

interactions while reducing the influence of weak or spurious associations. This threshold,

higher than the third quartile value (mean of 0.4) and close to the maximum observed weight

(mean of 0.61), ensures that only the strongest links are preserved for further analysis (Figure

2e, supplementary Table S1).

Consensus network construction by integrating wTO networks

To obtain a single consensus network derived from multiple independent wTO networks, the

`intersection()` function from the `igraph` package was used. A custom function was

developed to subset the overlapping links across the given networks and calculate an average

weight for each link. Links with differing signs were excluded. The consensus weight between

nodes i and j ($wTO_{CN} = [\Omega_{i,j}]$) for common links across n replicate networks (`k=1,…,n`) was

calculated using the formula from the `wTO.Consensus()` function of the `wTO` R package,

as follows:

$$\Omega_{i,j} = \sum_{k=1}^{n} \alpha_{ij}^k \, \omega_{ij}^k$$

173   Where

175
$$\alpha_{ij}^k = |\omega_{ij}^k| / \sum_{k=1}^{n} |\omega_{ij}^k|$$

This approach ensures that stronger link weights ω_{ij}^k contribute more significantly to the

corresponding consensus weights through a greater weighting factor α_{ij}^k. This helps preserve

the impact of strong links, preventing their dilution by weaker ones. In contrast, a simple

average would treat all connections equally, potentially blurring the distinction between strong

and weak links and resulting in the loss of valuable information.

Differential co-expression networks analysis

To identify differences between AD and control consensus networks, we performed a network

comparison analysis using the `CoDiNA` R package (55). CoDiNA (Co-Expression Differential

Network Analysis) is a network comparison framework that systematically identifies shared

and condition-specific gene-gene interactions between multiple networks. Instead of simply

highlighting differentially connected genes, CoDiNA classifies links into three categories: (1)

conserved links that are common between two conditions with the same sign (α-link); (2)

condition-specific links, appearing only in one condition (β-link); and (3) reversed links,

representing common correlations with opposite signs in the two conditions (γ-links). This

qualitative framework enables a clear interpretation of how gene interactions differ between

conditions. For example, conserved links indicate stability in gene regulation, while condition-

specific and reversed links highlight shifts in regulatory relationships that may underpin

disease pathology. The package allows comparison of multiple networks, requiring input in a

table format where the first two columns specify node pairs, and the third column contains

their correlation values. We disabled weight normalization (stretch = FALSE) and applied the

default cutoff (0.33) to define the minimum weight threshold for a link to be considered present.

CoDiNA considers genes (nodes) present in both networks to avoid false associations, as the

absence of a gene in one network might be due to technical reasons. However, given the

nature of the gene expression data used in this study to construct the wTO consensus

networks, the gene lists were identical for both conditions. Thus, the absence of a gene in one

network did not necessarily indicate a technical issue but rather that it might not meet the

criteria for meaningful correlations, such as the weight cutoff and network integration. To

address this, we added the absent genes to each network with a zero-weight assigned to their

links, ensuring that genes present in one condition but absent in the other yet still exhibiting a

correlation were not excluded from the comparison. This approach enabled a comprehensive

analysis of network differences between AD and control conditions.

Hub core genes

To determine hub genes within each network, we used the K-core decomposition method (56),

which identifies the most interconnected nodes by progressively removing nodes with degrees

less than k until no more such nodes exist. K-cores are determined as subgraphs where every

vertex has a degree of at least k. Therefore, vertices with the highest coreness scores

represent the most tightly integrated and connected parts of the network. The k-core subgraph

is defined through the following process: first, all nodes and their links are removed if their

degree is less than k (starting with $k = 1$). The algorithm works iteratively, increasing k each

time, until no more nodes meet the criteria (Figure 8). This effectively peels off layers (shells)

of nodes until the graph has none left. For each network, all nodes were classified into shells

based on the coreness scores, and nodes with the highest score were assigned as core hub

genes of that network. We used the `coreness()` function from the `igraph` R package to

calculate the coreness scores.

Figure 8: Illustration of K-core decomposition method (57)**.** All nodes of the network are classified

into shells based on their coreness score. The inner shell (K-shell 3) with red-colored nodes represents

those with the highest connectivity within the network, considered as core hub nodes in this study.

Identifying sub-network clusters

Clusters are groups of genes with similar expression profiles and were explored based on the

topological structure of the gene co-expression network. The Louvain method (58) was used

for clustering. While this function can consider edge weights, it only accepts positive numeric

vectors for weights. Since the networks contain both positive and negative weights, the

weights argument was set to NULL, ensuring that all links are considered without their signs.

Additionally, the resolution argument was set to 1, as it corresponds to the original definition

of modularity defined by the algorithm. Essentially, a lower resolution value results in fewer

and larger clusters. Cluster smaller than 20 links were excluded from further analysis.

GO enrichment analysis

Gene Ontology (GO) enrichment analysis for biological process terms was performed using

the `compareCluster()` function from the `clusterProfiler` R package (59). The analysis

was conducted separately for each gene cluster, using the same background genes.

Background genes comprised the protein-coding genes that were included in the wTO

construction. P-values were adjusted using the false discovery rate (FDR) method, with a

significance threshold of 0.05.

Redundancy reduction of the enriched GO terms

Although GO enrichment is a powerful tool for understanding the biological processes and

pathways associated with a set of genes, this analysis is frequently challenged by the

redundancy inherent in GO terms. This redundancy occurs because of the hierarchical nature

of the GO hierarchy, where parent terms frequently encompass child terms. For example, a

gene may be annotated with both a parent term and a more specific term at different levels of

the hierarchy within the GO. Various methods have been developed to address this issue,

each with its own advantages and disadvantages. However, none of these methods work

flexlessly, and some incorrectly group GO terms with inappropriate representative parent

terms. Therefore, a manual approach was chosen to mitigate redundancy. Child terms were

manually grouped into broader cellular biological process terms to which they belong.

Additionally, GO terms that were irrelevant, included fewer than five genes, or represented

different terms for the same biological process were excluded.

Random Walk with Restart

The random walk on a graph is an iterative process where a walker transitions from its current

node to a randomly selected neighbor, starting from specified source node(s), known as

seeds. The Random Walk with Restart (RWR) algorithm extends this concept by allowing the

walker to restart at the seed node(s) with a probability r at each time step (60). In this study,

RWR was used to assign lncRNAs within each cluster to the enriched functional categories

(GO terms) identified for that cluster. Each co-expression network constructed in this study

included both protein-coding genes and lncRNAs as nodes. Knowing the involvement of

protein-coding genes in the enriched biological processes, we aimed to assign lncRNAs to

these processes by leveraging their connectivity with neighboring protein-coding genes within

the network. For each cluster, we performed iterative RWR calculations, using genes from

each functional category as seed genes and ranking lncRNAs based on their associations.

The process for each iteration included the following steps: (1) performing RWR calculations

within each network cluster, (2) using genes from each functional category within the cluster

as seed genes and computing RWR scores for lncRNAs in that cluster, (3) excluding link

weights when computing RWR scores due to the algorithm's inability to handle negative values

in signed networks, (4) computing the adjacency matrix for the network, (5) normalizing the

adjacency matrix by columns, and (6) setting the restart probability (r) to 0.7.

Assigning lncRNAs to the functional category based on RWR scores

LncRNAs were ranked based on their computed scores for each functional category, and

those scoring above the 10th percentile were assigned to that functional category. As cross-

validation, we calculated RWR scores for genes within each functional category using a leave-

one-out method. Within each cluster, for each protein-coding gene in a given GO group, we

treated all other genes except one as seeds and computed the RWR score for the excluded

gene. This process was repeated across all genes annotated to each functional category. The

assigned lncRNA was accepted only if its RWR score did not fall below the lowest RWR scores

observed in the leave-one-out calculation for that category (Figure S1).

Statistical evaluation tests

Szymkiewicz-Simpson coefficient and permutation test was specifically chosen to rigorously

assess whether the observed overlaps in lncRNA annotations were significant or merely due

to random chance. The similarity between assigned lncRNAs to a common functional category

across the two networks was initially measured using the Szymkiewicz-Simpson coefficient.

For each functional category, the overlap coefficient was calculated by determining the

intersection of lncRNAs assigned to the respective functional category in each network,

normalized by the minimum number of lncRNAs assigned to that functional category in either

network. This approach normalized the overlap relative to the size of the smaller set,

accommodating the significant disparity in the total number of lncRNAs between the FCX and

TCX control networks (1100 and 504, respectively). The statistical significance of the observed

overlap was assessed using a permutation test. The permutation test involved generating

10,000 random permutations of lncRNA sub-samples from the complete list of lncRNAs for

each network. During each permutation, lncRNAs were randomly sampled from the entire pool

of lncRNAs in each network to create a new, synthetic group of lncRNAs matching the original

count of lncRNAs assigned to that functional category. The significance of the observed

overlap was assessed by calculating the proportion of permuted overlap coefficients that were

equal to or greater than the observed coefficient, which was then divided by the number of

permutations to yield a p-value.

To assess the difference in lncRNA representation between TCX AD and control core

networks, we constructed a contingency table comparing the number of core lncRNAs in each

network and evaluated statistical significance using Fisher's exact test.

Differential gene expression analysis

The `DESeq2` R package was used to identify differentially expressed genes (DEGs) in AD compared to control. Genes with a false discovery rate (FDR) < 0.05 and absolute log fold change > 0.3 were considered significant DEGs. Due to the integration of several networks to make each consensus network, identifying DEGs presented a challenge. To address this, only DEGs that were consistently identified across all gene expression datasets used to construct the given network were selected. For example, the AD TCX consensus network integrates three networks: AD Mayo, AD MSBB BM36, and AD MSBB BM22. DEGs were identified separately for each dataset—AD Mayo, AD MSBB BM36, and AD MSBB BM22. Ultimately, the DEGs in the AD TCX consensus network were determined by selecting those DEGs that overlapped across all these three datasets.

Network visualization

The network visualizations presented in the manuscript were created using Cytoscape v.310.2 (61) and Gephi v.0.10.1 (62).

Single cell RNA-seq analysis

The single-cell RNA sequencing data was downloaded from the Synapse repository with the Synapse ID syn22079621. This dataset includes samples from the prefrontal cortex, comprising 7 control and 11 AD cases (63). Data preprocessing and analysis were performed using Seurat v 4.3.0.1 (64). To ensure high-quality single-cell data, we applied the following filtering criteria: cells with fewer than 200 detected genes or more than 7500 detected genes were removed, and cells with more than 5% of reads mapping to mitochondrial genes were excluded to eliminate low-quality or stressed cells. Following quality control, the dataset was normalized using the `NormalizeData` function in Seurat. Uniform Manifold Approximation and Projection (UMAP) was used for dimensionality reduction and visualization of cellular clusters. Clusters were identified and cell type were manually annotated based on established marker genes including *AQP4*, *SLC1A2*, and *GPC5* for astrocytes; *MBP*, *MOBP*, and *PLP1* for oligodendrocytes; *CD74*, *LPAR6*, *DOCK8*, and CSF1R for microglia; FLT1, *CLDN5*, and

LAMA2 for endothelial cells; *VCAN*, *DSCAM*, *PCDH15*, and *MEGF11* for oligodendrocyte

progenitor cells; *NRGN* for excitatory neurons; and *GAD1* for inhibitory neurons.

Bulk RNA-seq deconvolution

To estimate cell-type proportions in bulk RNA-seq data, deconvolution was performed using

`CIBERSORT` R package (65). The single-cell dataset described earlier was used to construct

a signature gene matrix, utilizing only control samples for this purpose. The expression data

was normalized using `SCTransform` (SCT), and the signature matrix was generated from

SCT-normalized data by averaging expression across cell types. The CIBERSORT algorithm

was run with 100 permutations, and only samples with a p-value < 0.05 were retained for

further analysis. The estimated cell-type proportions between AD and control samples were

compared using the Wilcoxon test, with false discovery rate-adjusted p-values < 0.05

considered significant.

Results

Construction and integration of gene co-expression networks

We used a total of five RNA-seq datasets from two independent studies: the Mayo RNA-seq

study (Mayo) (43) and the Mount Sinai Brain Bank study (MSBB) (45). The gene expression

data were derived from two brain regions: TCX and FCX. These datasets were generated

using samples with detailed clinical-pathological diagnoses, including Braak NFT stage (47),

plaque density, CERAD (48), and CDR (49).

To minimize the impact of technical and unwanted biological factors, we implemented several

steps: 1- Filtering individuals based on AD assessment criteria (Figure 2a). 2- Sex balance**:**

each dataset includes an equal number of males and females. 3- Age balance**:** the age range

was balanced within each group, as well as across all groups, as much as possible (Figure

2b-c). Furthermore, we uniformly processed all data from FASTQ files to minimize the impact

of technical factors.

Figure 2: Summary of data demographics. a) Parameters and thresholds used to classify individuals

into AD and control groups include Braak stage (based on neurofibrillary tangles distribution), CERAD

(scoring neuritic plaques) from 0 (none) to 3 (frequent), and CDR (Clinical Dementia Rating) 0 (no

dementia) to 3 (severe dementia). **b)** Illustration of brain areas used in this study that are temporal

(yellow) and frontal (blue) cortices. **c)** The datasets, sample size, and Brodmann (BM) brain area utilized

in this study. **d)** Data demography after data preparation: sample sizes were balanced between AD and

control groups, as well as between sexes. Efforts were made to maintain a balanced age distribution.

e) Distribution of wTO values for each network before filtering links for a cutoff. Red dashed lines

indicate weight cutoffs. **f,g)** The graphs illustrate the number of links and nodes before and after filtering

links and integration into consensus networks (x-axis: Before, After and Consensus). Each line and

node represent one network, distinguished by colors for AD and control networks (see Tables S1 and

S2 for more details).

Normalized gene count data were then used to construct signed weighted gene co-expression

networks using the wTO method (54) separately for each AD and control dataset. We used

gene count data for both protein-coding and lncRNA genes as input, resulting in networks that

capture correlations between protein-coding and lncRNA genes. To focus on the strongest

links, we considered the summary statistics of absolute weights of links in the networks (Figure

2e, supplementary Table S1) and excluded all weak links with an absolute weight below 0.5.

This substantially reduced the number of links from a median of 219 million to 7.2 million links

(Figure 2f). While the number of nodes also decreased, the reduction was less drastic,

dropping from 20,900 nodes before filtering to a median of 13,200 nodes afterward (Figure

2g). This implied that most genes remained in the network, but only the most robust

interactions were represented.

Subsequently, the filtered networks were integrated separately for AD and control groups

based on brain regions to further enhance the reliability of the networks, excluding links that

might be dataset/study specific. Specifically, the Mayo TCX network was integrated with two

MSBB networks, BM36 and BM22, all derived from different parts of the temporal cortex, into

a consensus TCX network. Similarly, the BM10 and BM44 networks, which originated from

two distinct areas of the frontal cortex, were integrated to form the consensus FCX network

(Figure 2 b-c, supplementary Table S2). The resulting consensus networks included only the

commonalities across the integrated networks for each brain region. This method ensured that

only gene relations observed in multiple studies were included in our further analyses. From

here on, we refer to these consensus networks as the "AD" and "control" networks of the TCX

and FCX, respectively.

Comparative analysis of the TCX networks

We began our analysis with the TCX networks. Using K-core decomposition, we identified

nodes in the innermost shell as core genes (hubs) in both AD and control networks

(supplementary Table S3 and S4). The TCX AD network had 306 core genes, compared to

235 in the TCX control network, with 18% (81 genes) shared between them (Figure 3a).

Notably, lncRNAs accounted for 15% of the core genes in the AD network, nearly double the

8% observed in the control network (Figure 3b). Statistical evaluation using Fisher's exact test

confirmed the significant enrichment of core lncRNAs in the AD network (p-value = 0.016,

odds ratio = 2.01). Among the TCX AD core genes, 22 were significantly differentially

expressed (DE, FDR < 0.05 and absolute log fold change> 0.3) across all three TCX datasets,

all of which were downregulated in AD (Figure 3 c and d). These included 17 protein-coding

and 5 lncRNA core genes. A complete list of DE genes is provided in supplementary Table

S5.

Figure 3: Demography and differential expression analysis of TCX core genes. a) Venn diagram

depicting the overlap of core genes between AD (orange) and control (green) networks in TCX. **b)**

Proportion of protein-coding and lncRNA genes within cores of TCX AD network (orange) and control

network (green). **c** and **d)** Heatmaps showing the log2 fold change (FC) for differentially expressed (DE)

core genes in AD, including protein-coding (c) and lncRNA (d) genes that are in the TCX AD network.

Each column represents the log2 FC for a single dataset used to construct the TCX networks, which

are Mayo, MSBB BM22 and BM36.

To examine the structure and underlying biological functions of the networks, we applied the

Louvain method for community detection to each consensus network separately. In the TCX

AD network, we observed three gene clusters, whereas the control network formed five

clusters (Figure 4a, Table 1). GO enrichment analysis was conducted on the protein-coding

genes within each cluster, using the same set of background genes (see Methods). Enriched

GO terms were manually curated and grouped into broader functional categories to address

GO term redundancy (see Methods for details). Significantly enriched biological processes fell

into eight broad functional categories, including 1- mitochondria energy production, 2-

intracellular trafficking, 3- synapse signaling, 4- neurogenesis, 5- apoptosis, 6- cytoskeleton

organization 7- gene expression, 8- cellular homeostasis, which were enriched to differing

degrees among the clusters (Figure 4b, supplementary Table S6). Additionally, four GO terms

were grouped as "others" due to their specificity or broad relevance (e.g., GTPase activity).

Among all clusters, only genes of control cluster 5 were not significantly enriched for any terms.

Cellular homeostasis was the only functional category that was enriched only in control

clusters.

	Cluster 1		Cluster 2		Cluster 3		Cluster 4		Cluster 5	
	Nodes	Links	Nodes	Links	Nodes	Links	Nodes	Links	Nodes	Links
control	961	18,583	1,662	31,509	836	4,702	113	169	894	30,439
AD	920	48,198	1,689	44,374	476	17,262				

429   **Table 1: The topological details of AD and control clusters in the temporal cortex**

430
431
Figure 4: Topological and differential co-expression analysis of TCX networks. a) Representation

of clusters for TCX AD and control networks, shown on the right and left, respectively. There are two

node sizes; larger nodes represent core hub genes, while smaller nodes represent the remaining genes.

Each color represents a distinct cluster. Gray backgrounds indicate links, which are faded to reduce

complexity for visualization. The networks were visualized by Gephi (62). **b)** The plot displays selected

enriched GO terms and their classification into broader functional categories. The significance of

enriched GOs was assessed using the false discovery rate, and terms with an adjusted p-value below

0.05 were selected. Dot colors indicate the adjusted p-value, and dot size is proportional to the number

of genes of interest in the input gene list for the given term. **c)** Distribution of differential co-expression link categories within each cluster. The percentages represent relative abundance, meaning the original proportion of link types within each cluster is normalized relative to the cluster size. **d)** Distribution of differential co-expression links categories connecting genes associated with each functional category. **e)** Distribution of protein-coding core genes annotated to the enriched GO terms.

In order to gain a deeper understanding of the differences between AD and control networks, a differential co-expression network analysis was performed (55). We classified links into three categories: 1- conserved links, which were common between the two conditions (i.e., AD and control); 2- condition-specific links, which appeared only in one condition (i.e., AD-specific); and 3- reversed links, which represented correlations with opposite signs in the two conditions. The results showed that 96%, 77%, and 83% of links in AD clusters 1, 2, and 3, respectively, were AD-specific (Figure 4c). This pattern was also reflected in links between genes of each functional category in AD clusters (Figure 4d). These findings indicated that AD-related changes are not confined to a single cluster or GO term; rather, they broadly impacted all three gene clusters and the identified underlying biological process. Similar observations were seen in the control network analysis, indicating that the majority of links per cluster and functional category were control-specific and, in other words, were absent in AD. There was only one exception: 51% of control cluster 1 included conserved links, which involved genes annotated to mitochondria energy production and apoptosis.

The observed rewiring of the AD network suggested a shift in the organization of molecular interactions. To investigate the key players of this rewiring, we examined all core genes, as these genes were likely central to the disrupted processes and extensive rewiring observed in the AD network. Notably, of the protein-coding core genes, a higher proportion was annotated to the functional categories in the AD network (18%, 46 genes) compared to the control network (2%, 5 genes) (Figure 4e). Among the core genes linked to the enriched GOs, the largest proportion (34%) were associated with mitochondrial energy production, followed by 22% with synaptic signaling and 15% with intracellular trafficking (Figure 4e). While the

majority of the protein-coding core genes in both networks were not annotated to enriched GO

terms, the increased representation of AD core genes in these categories underscored their

potential roles in key processes disrupted in AD pathology, particularly in energy production,

neural communication, and intracellular transport.

Among the 46 core genes of the AD network annotated to the enriched GO terms (Table 2,

Figure 5), nearly all were exclusively core to the AD network, except for four genes. Notably,

these four genes—*CHL1* and *USP33* (neurogenesis), *GRIA2* (synapse signaling), and *YARS2*

(mitochondrial gene expression)—were annotated to biological processes only within the AD

network and not in the control network. This observation suggests that these genes might gain

a disease-related role in the context of AD. It might be observed that a single gene was

annotated to different functional categories in the AD and control networks (Table 2). This was

explained by the fact that the gene clustered with different sets of genes in the two networks,

leading to its association with distinct GO terms. This observation reflected the differing

patterns of gene interactions between AD and control networks, likely linked to the disease

condition.

Table 2: Identified key protein-coding genes potentially associated with AD pathology in TCX.

Gene name	KEGG pathway	DEG	AD		Control		
			Cluster	Functional category	Cluster	Functional category	hub
ABL1	-	-	3	Cytoskeleton organization	1	-	-
ACSL3	-	-	1	Mitochondrial energy production	1	Mitochondrial energy production	-
AK6	-	-	1	Mitochondrial energy production	1	Mitochondrial energy production	-
AP3M2	-	Down	1	Synapse signaling	1	-	-
ATL1	-	-	2	Neurogenesis	-	-	-
ATP6AP2	-	-	1	Mitochondrial energy production, synapse signaling	1	Intracellular trafficking, synapse signaling	-
ATP6V1A	-	-	1	Intracellular trafficking, mitochondrial energy production, synapse signaling	1	Intracellular trafficking, mitochondrial energy production, synapse signaling	-

Gene							
ATP6V1B2	-	Down	1	Intracellular trafficking, mitochondrial energy production, synapse signaling	1	Intracellular trafficking, Mitochondrial energy production, synapse signaling	-
ATP6V1D	-	-	1	Intracellular trafficking, mitochondrial energy production, synapse signaling	1	Intracellular trafficking, synapse signaling	-
BLOC1S6	-	-	1	Synapse signaling	1	-	-
CCNB1	-	-	1	Mitochondrial energy production	1	Mitochondrial energy production	-
CHL1	-	-	2	Neurogenesis	5	-	core
CISD1	-	-	1	Mitochondrial energy production	1	Mitochondrial energy production	-
COQ10B	-	-	1	Mitochondrial energy production	1	Mitochondrial energy production	-
EPM2AIP1	-	-	1	Mitochondrial energy production	1	Mitochondrial energy production	-
ETFRF1	-	-	1	Mitochondrial energy production	1	Mitochondrial energy production	-
EXOC8	-	-	1	Intracellular trafficking	1	-	-
GHITM	-		1	Apoptosis, mitochondrial energy production, others (mitochondrial membrane organization)	1	Apoptosis, mitochondrial energy production	-
GLRX2	-		1	Mitochondrial energy production	1	Mitochondrial energy production	-
GRIA2	-		2	Synapse signaling	2	-	core
GUCY1B1	-		1	Mitochondrial energy production	1	Mitochondrial energy production	-
MDH1	-		1	Mitochondrial energy production	1	Mitochondrial energy production	-
MKKS	-		3	Cytoskeleton organization	1	-	-
MRPL42	-		1	Gene expression	1	-	-
MRPS30	-	-	1	Gene expression	1	-	-
MRPS35	-	-	1	Gene expression	1	-	-
NAPB	-	-	1	Synapse signaling	1	-	-
NDUFA5	AD pathway	Down	1	Mitochondrial energy production	1	Mitochondrial energy production	-
NDUFA9	AD pathway	-	1	Mitochondrial energy production	1	Mitochondrial energy production	-
NDUFB5	AD pathway	Down	1	Mitochondrial energy production	1	Mitochondrial energy production	-
NECAP1	-	Down	1	Synapse signaling	1	-	-
PFN2	-	Down	1	Synapse signaling	1	-	-

PREPL	-	-	1	Synapse signaling	1	-	-
PRPS2	-	-	1	Mitochondrial energy production	1	Mitochondrial energy production	-
RFK	-	Down	1	Mitochondrial energy production	1	Mitochondrial energy production	-
SCOC	-	-	1	Intracellular trafficking	1	-	-
SKP1	-	-	1	Intracellular trafficking	1	-	-
TAF9	-	-	1	Apoptosis	1	-	-
TMEM14A	-	-	1	Apoptosis, others (mitochondrial membrane organization)	1	Apoptosis	-
TMEM33	-	-	1	Intracellular trafficking, synapse signaling	1	-	-
TOMM70	-	-	1	others (mitochondrial membrane organization)	1	-	-
TRMT10C	-	-	1	Gene expression	1	-	-
UBE2T	-	-	1	Intracellular trafficking	1	-	-
USP33	-	-	2	Neurogenesis	5	-	core
VPS35	-	-	1	Apoptosis, intracellular trafficking, synapse signaling	1	Apoptosis	-
YARS2	-	-	1	Gene expression	1	-	core

The table lists 46 protein-coding key genes potentially associated with AD pathology in the TCX AD network. Various features of these genes are listed, including their involvement in the KEGG AD pathway, differential expressions in AD (DEG), cluster membership, as well as their annotation to the functional categories. All are AD hubs, while their hub status in control is indicated under the hub column.

lncRNA functional assignment and core lncRNAs in TCX networks

The gene co-expression networks in this study included both protein-coding and lncRNA genes, providing us with the opportunity to gather functional information for lncRNAs based on their network neighbors. For each functional category, Random Walk with Restart (RWR) was run using annotated protein-coding genes as seeds. LncRNAs were ranked based on their RWR scores, and the top 10th percentile were assigned to the corresponding functional category. To ensure the biological relevance of these assignments, a leave-one-out approach was applied to validate the rankings. A lncRNA was assigned to a category only if its RWR

score exceeded the lowest score from the leave-one-out simulation (Methods, supplementary

Figure S1 and Table S8). Using this approach, 54 lncRNAs in the TCX AD network and 103

in the TCX control network were assigned to functional categories within their respective

clusters (supplementary Table S8).

Out of the 46 AD core lncRNA genes, we focused on 27 lncRNAs that were assigned to at

least one functional category, as these lncRNAs likely played key roles in the dysregulation of

their assigned functional categories within the AD network (Table 3, Figure 5). All but six of

these lncRNAs were AD-specific core genes. Notably, *FBXW7-AS1*, *Lnc-PDE4D-2*, *Lnc-*

KLHL11-1,   *Lnc-NIF3L1-5*,   *Lnc-KALRN-1*,   *Lnc-CCDC68-1*,   *HSALNG0099005*,

HSALNG0105911, *ZRANB2-DT*, *ENSG00000287022*, and *SLC8A1-AS1* were functionally

assigned only within the AD network, suggesting a gain of function for these lncRNAs in the

context of AD. The remaining lncRNAs were functionally assigned in both networks, either to

similar or different functional categories (Table 3). Many of these lncRNAs were assigned to

all functional categories within their respective clusters, as expected given their status as core

genes with high connectivity to numerous nodes, which often results in high RWR scores.

However, a closer examination of their functional categories and comparisons with the control

network revealed considerable differences. This suggests that lncRNAs were incorporated

into different neighborhoods or changed the sets of genes they regulated in AD compared to

control brains. For example, *POC1B-AS1* was assigned to cytoskeleton organization-related

processes in the AD network, whereas it was associated with mitochondrial energy production

in the control network. Among these 27 core lncRNAs, five were downregulated in AD, namely

ITFG1-AS1, *Lnc-MYCN-6*, *Lnc-EFR3A-6*, *HSALNG0089044*, and *Lnc-CCDC68-1* (Figure 3d,

Table 3).

Table 3: Key lncRNA genes potentially associated with dysregulated GO terms in the TCX AD
network.

| Gene name | DEG | AD network | | Control network | | |
		Cluster	Functional category	Cluster	Functional category	Hub
FBXW7-AS1	-	1	Synapse signaling	1	-	-

Lnc-PDE4D-2	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	-	-
Lnc-KLHL11-1	-	1	Apoptosis, gene expression, mitochondrial energy production, others (mitochondrial membrane organization)	1	-	-
Lnc-NIF3L1-5	-	3	Cytoskeleton organization	1	-	-
Lnc-KALRN-1	-	3	Cytoskeleton organization	2	-	-
Lnc-CCDC68-1	Down	3	Cytoskeleton organization	1	-	-
HSALNG0099005	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	Absent	-	-
HSALNG0105911	-	2	Neurogenesis, synapse signaling	5	-	core
ZRANB2-DT	-	2	Neurogenesis, synapse signaling	5	-	core
ENSG00000287022	-	3	Cytoskeleton organization	5	-	core
SLC8A1-AS1	-	3	Cytoskeleton organization	5	-	core
HSALNG0096948	-	1	Intracellular trafficking	1	Intracellular trafficking, synapse signaling	-
ITFG1-AS1	Down	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Apoptosis, intracellular trafficking, mitochondrial energy production, others*, synapse signaling	-
Lnc-MYCN-6	Down	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Intracellular trafficking, synapse signaling	-
Lnc-EFR3A-6	Down	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Apoptosis, mitochondrial energy production, others*	-
HSALNG0089044	Down	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane	1	Intracellular trafficking, mitochondrial energy production, others*, synapse signaling	-

			organization), Synapse signaling			
Lnc-LRP11-1 (ENSG00000285 889)	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Apoptosis, intracellular trafficking, mitochondrial energy production, others[*], synapse signaling	-
Lnc-KBTBD3-1	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Apoptosis, intracellular trafficking, mitochondrial energy production, others[*], synapse signaling	-
SNX10-AS1	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Apoptosis, mitochondrial energy production	-
Lnc-LRP11-1 (ENSG00000231 760)	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Intracellular trafficking, others[*], synapse signaling	-
Lnc-KCNH5-3 ant	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Intracellular trafficking	-
Lnc-CLUH-2	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Others[*]	-
Lnc-SUMF1-5	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), Synapse signaling	1	Others[*]	-
HSALNG0144080	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Intracellular trafficking, mitochondrial energy production, others[*], Synapse signaling	-

POC1B-AS1	-	3	Cytoskeleton organization	1	Mitochondrial energy production	-
MYCBP2-AS1	-	3	Cytoskeleton organization	1	Apoptosis, intracellular trafficking, mitochondrial energy production, synapse signaling	core
HSALNG0018143	-	2	Neurogenesis, synapse signaling	1	Apoptosis, mitochondrial energy production, others*	core

The table shows the 27 selected AD core lncRNAs and their functional categories in the TCX AD and

control networks, as well as cluster membership and differential expression in AD. Down in the DEG

column indicates downregulation in AD. others*: positive regulation of cyclin-dependent protein

serine/threonine kinase activity.

Figure 5: AD cluster 1 network visualization. Node colors indicate functional categories; multicolored

nodes represent genes annotated to multiple categories. Larger nodes are identified as key genes in

the AD network, while smaller nodes represent other genes in AD cluster 1. Circles represent protein-

coding genes, and squares represent lncRNA genes. Link colors: blue links are conserved between the

TCX AD and control network, and black links are AD-specific links. Names of key genes are enlarged;

names of other genes are visible upon full zoom. Further details can be explored through the interactive

Shiny app we have provided.

Comparing TCX and FCX

After analyzing networks of the temporal cortex, we performed the same analysis for the frontal cortex and compared our results between the two brain areas. In the frontal cortex, we found that unlike in the temporal cortex, the AD network shows much higher similarity to the control network (Figure 6 a-d), where the majority of links are conserved in the AD network (86%, figure 6a). Conversely, the control network is predominantly composed of control-specific links (83%), with minimal overlap (14%) with links found in the AD network (Figure 6b). This observation implied that the AD network in the frontal cortex has lost a significant number of gene-gene correlations. This hypothesis was supported by the marked difference in the size of the networks: the AD network contained 1.17 million links, while the control network contained 5.93 million links. This pattern was also evident in the significant overlap between AD core hubs and control core genes (Figure 6e). While this overlap suggested that certain gene relationships were preserved in AD, despite the extensive loss of specific gene-gene interactions, a substantial number of core genes presented in the control network were absent in the AD network. Interestingly, the proportion of gene types, i.e., protein-coding or lncRNA genes, among the core genes {Citation}was very similar between AD and control networks, with 15% or 18% of cores being lncRNAs, respectively (Figure 6f).

We used the same method as for the TCX to cluster the FCX consensus networks (supplementary Table S9). As done for the TCX, significantly enriched GO terms were then manually classified into five broader functional categories (Figure 6g) including: 1-mitochondrial energy production, 2-intracellular trafficking, 3- synapse signaling, 4- apoptosis, 5- gene expression. Additionally, an "other" category was included, containing a single term, mitochondrial membrane organization. A complete list of enriched GO terms can be found in supplementary table S10. Genes in clusters 1 and 3 of both AD and control networks were significantly enriched for biological process GO terms, but not in the other clusters.

Differential gene co-expression analysis revealed a dominance of conserved links connecting genes associated with enriched GO terms within AD clusters (Figure 6h). Each functional

category showed 94% to 100% conserved links. In contrast, in the control network, the majority

of links were control-specific within the functional categories. One possible interpretation of

this tendency was that the pathogenic alterations associated with AD disrupted numerous

healthy gene expression correlations in the frontal cortex area. This might also suggest that

AD-specific correlations represent cellular compensation against AD-related disruptions.

In bulk RNA-Seq data, observed differences in gene expression can often be attributed to

variations in cell-type composition. Given that such differences might also contribute to

alterations in gene co-expression networks, we sought to assess the cellular composition of

the bulk RNA-seq data used in this study. To this end, we performed a deconvolution analysis

on both the TCX and FCX regions using CIBERSORT, utilizing human brain single-cell data

as a reference (Figure 6i). Across both brain regions and independent of disease status,

excitatory neurons were the predominant cell type, comprising 41–56% of the total population.

However, the relative abundance of other cell types varied between TCX and FCX, with

microglia consistently being the least abundant in both regions. The overrepresentation of

excitatory neurons and the underrepresentation of microglia align with previous findings in the

human brain, indicating that bulk RNA-seq data primarily capture transcriptional changes in

excitatory neurons, while microglia, due to their lower RNA content, are typically

underrepresented (66).

To assess whether AD pathology affects cell-type composition, we compared cell-type

proportions between AD and control samples within each brain region. Interestingly, we

observed significant alterations only in TCX, where excitatory neurons were markedly

decreased (p.adj = 9.0×10^{-5}), while astrocytes (p.adj = 4.66×10^{-5}), microglia (p.adj = $5.0 \times$

10^{-4}), oligodendrocytes (p.adj = 0.03), and endothelial cells (p.adj = 2.07×10^{-7}) were

significantly increased (Figure 6j). This shift in cellular composition may contribute to the gene

co-expression rewiring observed in the TCX AD network.

Figure 6: Results of comparative analysis between FCX and TCX. Pie chart showing proportions of

differential co-expressed links **a)** in the FCX AD network, **b)** FCX control network, **c)** TCX AD network

and **d)** TCX control network. **e** Venn diagram showing the overlap of core genes between AD and

control networks in FCX. **f)** Proportion of protein-coding and lncRNA genes within FCX core genes. **g)**

Significantly enriched GO terms for biological processes, grouped into five broader functional

categories: 1) mitochondrial energy production, 2) intracellular trafficking, 3) synapse signaling, 4) apoptosis, 5) gene expression, with an additional "other" category with a single term-mitochondrial membrane organization. **h)** Distribution of differential link categories within each functional category, with percentages representing relative abundance normalized to cluster size. **i)** Stacked bar plots showing the estimated cell-type proportions in FCX and TCX for AD and control samples. Each color represents one cell type. **j)** Significant cell-type proportion changes between AD and controls were observed only in TCX datasets. P-values were measured by the Wilcoxon test (*p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001).

In addition, the FCX control network was employed to evaluate the reliability and consistency of the lncRNA annotation method implemented in this study. We investigated the similarity of lncRNAs annotated to the same functional category between the TCX and FCX control networks (supplementary Table S11). Five functional categories common to both networks were analyzed, including mitochondrial energy production, intracellular trafficking, synapse signaling, apoptosis, and gene expression (supplementary Table S12). Significant overlaps of lncRNAs assigned to functional categories were observed between the FCX and TCX control networks, with p-values approaching zero (Table 4). The analysis accounted for the disparity in lncRNA totals between FCX and TCX control networks (1100 and 504, respectively) and a permutation test for statistical significance. Notably, the gene expression category was the only one without any overlapping lncRNAs. Additionally, permutation tests were conducted for lncRNAs not assigned to functional categories. These tests yielded a non-significant p-value of 0.71, highlighting that the functional assignments of annotated lncRNAs are significantly more consistent than would be expected by chance.

Table 4: Similarity of lncRNA annotations across FCX and TCX control networks.

Functional category	Overlap coefficient	p-value
Apoptosis	0.47	p < 1e-03
Synapse signaling	0.41	p < 1e-03
Mitochondrial energy production	0.3	p < 1e-03
Intracellular trafficking	0.2	p < 1e-03
Gene expression	0	1
Not assigned	0.12	0.71

619   The table displays the overlap coefficients between lncRNAs assigned to common functional categories

620   between FCX and TCX control networks, calculated using the Szymkiewicz-Simpson coefficient.

621   Permutation tests with 1000 random permutations were used to assess statistical significance. P-values

622   shown as '< 1e-03' indicate no permutation matched or exceeded the observed overlap, implying p-

623   values smaller than the test resolution (p < 1/1000).

Discussion

In this study, we presented robust consensus gene co-expression networks including protein-

coding and lncRNA genes of two brain regions (temporal cortex and frontal cortex) and

investigated how the networks differ in AD compared to control brains. Our deep comparative

analysis between the consensus networks of AD and control groups highlighted significant

rewiring of gene-gene interactions within the temporal cortex area (Figure 4c, d), in contrast

to a more pronounced loss of 'healthy' gene correlations in the frontal cortex area (Figure 6 a-

d). Prompted by these observations, we identified key players in the TCX AD network,

including 46 protein-coding genes and 27 lncRNAs (Table 2 and 3). Utilizing the established

functions of protein-coding genes as an anchor, we bridged lncRNAs to the enriched biological

processes, demonstrating the utility of network-based approaches. We not only nominated

potential key lncRNAs involved in AD pathology but also predicted their roles in biological

processes under both AD and non-AD conditions.

Networks provide a powerful framework for studying the structure of complex biological

systems, offering valuable insights into gene-gene interactions. However, network analysis

poses challenges, such as managing noise signals and enhancing the reliability of

correlations. To address these issues, we employed several strategies to mitigate biases and

increase robustness (Figure 1). First, we ensured that all datasets used had a minimum of 30

replicates per condition (Figure 2c), as larger sample sizes improve the accuracy of network

inference (67,68). Second, we applied the wTO method (69) to compute gene-gene

correlations. This approach accounts for shared network neighborhoods, producing more

robust scores compared to simple correlation calculations, and incorporates both positive and

negative correlations (70). The gene co-expression networks were subsequently filtered

based on the absolute value of gene pair correlations. Following this, we integrated the

networks within each brain region (TCX and FCX) (Figure 2b, c). This integration allowed us

to treat them as biological replicates, identifying common interactions across the integrated

networks and reducing the impacts of sampling and batch variability, a widely adopted

approach in gene co-expression studies (37–40,71,72). Ultimately, this consensus network

offers a high-resolution representation of the conserved cellular processes functioning under

AD and control conditions, providing a robust foundation for further investigation.

Co-expressed genes are often coregulated or involved in similar pathways, forming densely

interconnected clusters within the network. In this study, we identified gene clusters and their

functions for each consensus network. Through a comprehensive comparative analysis of

gene correlations and network topology between AD and control consensus networks, we

observed remarkable network alterations. Specifically, in the TCX networks, substantial

rewiring was evident within the AD clusters, which was also reflected in the links between

protein-coding genes associated with enriched terms (Figure 4c, d). The emergence of new

core genes (Figure 3a), along with their higher engagement in the AD network compared to

the control network (18% versus 2%, Figure 4e), indicated a reorganization of the molecular

landscape in AD, suggesting that specific genes adopted new roles within the altered

regulatory environment. As a result, we identified 46 core protein-coding genes as candidates

potentially playing a central role in the rewiring observed in enriched terms in the TCX AD

network (Table 2).

Several of these genes have been supported by prior experimental or network-based studies

as being linked to AD. Notably, 15 of these genes have been reported as hub genes in AD,

including *ATL1* (73), *AP3M2* (74)*, ATP6AP2* (75), *ATP6V1A* (76), *ATP6V1D* (77,78), *ACSL3*

(74,79), *CCNB1* (80)*, GHITM* (78), *GRIA2* (81), *MDH1* (82), *MRPS30* (83), *MRPS35* (82),

NAPB (84,85), *NDUFB5* (82) and *PFN2* (86). Several genes showed strong evidence

supporting their role in AD pathology. For example, *REPL* knockdown led to increased Aβ

levels and tau phosphorylation (87–89), *NAPB* was linked to neuroinflammatory neuronal

damage (90), *MDH1* correlated with total and phosphorylated tau levels (90,91) and *VPS35*

deficiencies or mutations caused altered endosomal trafficking and pathological tau

accumulation (13,92–94). Of particular interest, *ABL1* (c-Abl) contributed to tau pathology

through regulation of cytoskeletal signaling cascades (95,96). Our findings identified *ABL1* as

an AD-specific hub annotated to the stress fiber assembly term (Table 2). Notably, about half

of the 46 AD key genes were annotated to the mitochondrial energy production functional

category. Among them, *NDUFA5* and *NDUFB5*, encoding subunits of Complex I of the

mitochondrial respiratory chain, were consistently downregulated across TCX datasets

(Figure 3c), aligning with their known significant downregulation in AD (97). Recent findings

revealed that the endolysosomal V-ATPase acts as a receptor for Aβ and Tau in AD (14).

Among the 13 genes involved in the V-ATPase complex, four—*ATP6V1B2*, *ATP6V1A*,

ATP6AP2, and *ATP6V1D*—were in our key gene list. In addition to previously known AD-

associated genes, several key genes identified in this study have been pursued as promising

therapeutic targets for AD, including *ABL1* (98), *ACSL3* (79), *ATP6V1A* (76,99), *CISD* (100)

and *COQ10B* (101–104).

Having identified many previously known AD-associated protein-coding genes with our

network analysis, we leveraged the connectivity of lncRNAs with their neighboring protein-

coding genes within the network to infer potential functions for these lncRNAs. Through this

approach, 54 lncRNAs in the TCX AD network and 103 in the FCX control network were

assigned to functional categories identified within each cluster (supplementary Table S8 and

S11). The assigned biological functions were cross-validated by assessing the similarity of

lncRNAs assigned to common functional categories between the TCX and FCX control

networks, revealing significant overlap (Table 4). From the TCX AD network, 27 core lncRNAs

with functional assignments were selected for further investigation (Table 3). We observed

that all but six of these 27 lncRNAs were AD-specific core genes. Notably, *FBXW7-AS1*, *Lnc-*

PDE4D-2, Lnc-KLHL11-1, Lnc-NIF3L1-5, Lnc-KALRN-1, Lnc-CCDC68-1, HSALNG0099005,

HSALNG0105911, ZRANB2-DT, ENSG00000287022, and *SLC8A1-AS1* were functionally

assigned exclusively within the AD network, suggesting a gain of function for these lncRNAs

in the context of AD. Among them, *ZRANB2-DT*, assigned to neurogenesis and synapse

signaling, stood out. This lncRNA acts as an antisense RNA for *ZRANB2* (Zinc Finger

RANBP2-Type Containing 2), a gene previously reported to be associated with neuronal

responses to chronic oxidative stress—a process closely linked to neurodegenerative

diseases (105,106). We observed that the correlation between *ZRANB2-DT* and *ZRANB2* was

AD-specific, suggesting a gain regulatory role for *ZRANB2-DT* in the context of AD.

Additionally, differential gene expression analysis revealed that five of the candidate

lncRNAs—*ITFG1-AS1*, *Lnc-MYCN-6*, *Lnc-EFR3A-6*, *HSALNG0089044*, and *Lnc-CCDC68-*

1—were consistently dysregulated in AD across TCX datasets (Figure 3f, Table 3). Despite

comprehensive data collection from databases such as MalaCards (107), LncRNADisease

v3.0 (108), and existing literature (supplementary Table S13), none of the previously reported

AD-associated lncRNAs overlapped with our candidate lncRNAs. Nevertheless, the presence

of supportive evidence linking the candidate protein-coding genes to AD reinforced our

confidence that some of the identified key lncRNA genes were also likely to play a role in AD,

introducing promising new lncRNA candidates for further investigation and experimental

validation.

Aligning with the current knowledge that AD-related pathology begins in the medial temporal

lobe (2) and progresses to the frontal lobes in later stages (4), we had hypothesized that AD-

related network disruptions would differ across brain regions. In the TCX AD network, we

observed significant rewiring, while in the FCX AD network, there was a substantial loss of

healthy gene correlations. This suggests that AD-related damage initially disrupts cellular

interactions in the temporal lobe, as evidenced by the loss of control-specific links in the TCX

control network. As the disease progresses, cells in the temporal lobe establish new gene

expression patterns to compensate for the damage (AD-specific links). Conversely, the FCX

AD network shows a predominant loss of gene correlations, suggesting that the frontal lobe

may experience more extensive disruption before compensatory rewiring occurs. In line with

this, studies have shown that genes like *ACSL3* (79) and those involved in mitochondrial

respiratory complexes (97,109) display opposing patterns of dysregulation during the early

and late stages of AD, potentially reflecting mechanisms of gene expression compensation as

the disease progresses. This notion is further supported by findings from a single-cell study

across 19 cortical regions, which identified the temporal lobe as exhibiting the earliest and

most pronounced gene expression abnormalities (110).

While our study provided valuable insights into AD pathology through gene co-expression

network analysis, it is important to acknowledge certain limitations inherent to the

methodology. For instance, the thresholds applied for filtering links and annotating lncRNAs

to biological processes may have introduced some biases. The threshold for filtering links may

have removed interactions from one network if they fell below the chosen cutoff, leading to

their classification as lost links in that network. However, we reasoned that if a link was weaker

in one condition and stronger in another, it could still be considered an affected or altered

correlation. Similarly, the strict threshold for annotating lncRNAs may have excluded some

lncRNAs, as genes belonging to the same GO terms produced a wide range of RWR scores.

While this strict threshold helped maintain confidence in the assignments, we recognized that

non-annotated lncRNAs might still have been closely related to their neighboring protein-

coding genes. Additionally, other genes within the network might have been crucial to AD

pathology, but limitations in our methods may have prevented their identification. To address

this latter limitation, we made the TCX AD network publicly available as an interactive Shiny

app, enabling further exploration (see Data Availability). The app allows users to investigate

the entire TCX AD network, including all clusters, and search for specific genes of interest. It

also provides detailed information, such as gene neighbors, functional categories, differential

expression status, core membership, and more, offering a resource for further discoveries and

validations.

Conclusion

In summary, our network-based approach allowed us to provide insights into the alterations of gene co-expression networks in AD brains, revealing clusters of functionally related genes driven by key protein-coding and lncRNA genes. While early stages of AD, particularly in the FCX, appear to be characterized primarily by a loss of gene interactions, disease progression seems to trigger network rewiring, potentially aimed at preserving fundamental brain functions. Additionally, we predicted functional roles for numerous lncRNAs within these networks, identifying 27 lncRNAs with potential involvement in AD pathology. These findings highlight promising candidates for future studies, offering deeper insights into their roles in both normal brain function and AD development.

Data availability

The datasets used in this study are available in the Synapse online repositories under the following accession IDs: Mayo RNA-seq dataset (Synapse ID: syn3163039) (44) and MSBB RNA-seq dataset (Synapse ID: syn3159438) (46).

R script code availability

To enhance the exploration of our findings, we provided an interactive Shiny app that visualizes the gene co-expression networks. This tool enables researchers to investigate specific genes and their correlations and associated functions, offering deeper insights into the TCX AD and control networks. They can be downloaded from https://github.com/mpstar92/Shiny_App_ADnet.

The R script code used in this study is available on the GitHub repository at https://github.com/Tima-Ze/AD_network.

Acknowledgments

Data used in this project were obtained from the Synapse platform (https://www.synapse.org/), including the Mayo RNA-seq study (44), the Mount Sinai Brain Bank study (MSBB) dataset (46) and Human AD Single-nuclei Multi-Omics (63). We extend our gratitude to the data providers on the Synapse platform, as well as the researchers and authors involved in these studies. This study was supported by the International Max Planck Research School for Biology and Computation (IMPRS-BAC), Berlin, Germany. We would like to thank Vladimir M. Jovanovic, Melanie Sarfert, Yao-Chung Chen, and Rebecca Sinead Saager for their kind assistance and contributions, as well as Ralf Herwig for his scientific methodology advice.

Author contributions

Nowick supervised the research. Zebardast analyzed and visualized the data and wrote the manuscript. Riethmüller developed the Shiny app. All authors read and approved the final manuscript.

Conflict of Interest

The authors declare that there were no commercial or financial relationships involved in this research that could be interpreted as a potential conflict of interest.

Funding

This work is supported by funding from the Deutsche Forschungsgemeinschaft awarded to KN (NO 920/8-1).

References

1. Livingston G, Sommerlad A, Orgeta V, Costafreda SG, Huntley J, Ames D, et al. Dementia prevention, intervention, and care. The Lancet. 2017 Dec 16;390(10113):2673–734.

2.   Planche V, Manjon JV, Mansencal B, Lanuza E, Tourdias T, Catheline G, et al. Structural
progression of Alzheimer's disease over decades: the MRI staging scheme. Brain
Commun. 2022 Jun 1;4(3):fcac109.

3.   Ahmad F, Javed M, Athar M, Shahzadi S. Determination of affected brain regions at
various stages of Alzheimer's disease. Neurosci Res. 2023 Jul 1;192:77–82.

4.   Casanova R, Whitlow CT, Wagner B, Williamson J, Shumaker SA, Maldjian JA, et al.
High dimensional classification of structural MRI Alzheimer's disease data based on
large scale regularization. Front Neuroinformatics. 2011;5:22.

5.   Reddy PH. Abnormal tau, mitochondrial dysfunction, impaired axonal transport of
mitochondria, and synaptic deprivation in Alzheimer's disease. Brain Res. 2011 Sep
30;1415:136–48.

6.   Devi L, Prabhu BM, Galati DF, Avadhani NG, Anandatheerthavarada HK. Accumulation
of amyloid precursor protein in the mitochondrial import channels of human Alzheimer's
disease brain is associated with mitochondrial dysfunction. J Neurosci Off J Soc
Neurosci. 2006 Aug 30;26(35):9057–68.

7.   Moreira PI, Carvalho C, Zhu X, Smith MA, Perry G. Mitochondrial dysfunction is a trigger
of Alzheimer's disease pathophysiology. Biochim Biophys Acta BBA - Mol Basis Dis.
2010 Jan;1802(1):2–10.

8.   Plácido AI, Pereira CMF, Duarte AI, Candeias E, Correia SC, Santos RX, et al. The role
of endoplasmic reticulum in amyloid precursor protein processing and trafficking:
Implications for Alzheimer's disease. Biochim Biophys Acta BBA - Mol Basis Dis. 2014
Sep 1;1842(9):1444–53.

9.   Nishitsuji K, Tomiyama T, Ishibashi K, Ito K, Teraoka R, Lambert MP, et al. The E693Δ
Mutation in Amyloid Precursor Protein Increases Intracellular Accumulation of Amyloid β
Oligomers and Causes Endoplasmic Reticulum Stress-Induced Apoptosis in Cultured
Cells. Am J Pathol. 2009 Mar 1;174(3):957–69.

10.   Joshi G, Chi Y, Huang Z, Wang Y. Aβ-induced Golgi fragmentation in Alzheimer's
disease enhances Aβ production. Proc Natl Acad Sci. 2014 Apr;111(13):E1230–9.

11.   Joshi G, Bekier ME, Wang Y. Golgi fragmentation in Alzheimer's disease. Front Neurosci.
2015;9:340.

12.   Kim SH, Cho YS, Kim Y, Park J, Yoo SM, Gwak J, et al. Endolysosomal impairment by
binding of amyloid beta or MAPT/Tau to V-ATPase and rescue via the HYAL-CD44 axis
in Alzheimer disease. Autophagy. 2023 Aug;19(8):2318–37.

13.   Simoes S, Neufeld JL, Triana-Baltzer G, Moughadam S, Chen EI, Kothiya M, et al. Tau
and other proteins found in Alzheimer's disease spinal fluid are linked to retromer-
mediated endosomal traffic in mice and humans. Sci Transl Med. 2020 Nov
25;12(571):eaba6334.

14.   Gabrielli M, Prada I, Joshi P, Falcicchia C, D'Arrigo G, Rutigliano G, et al. Microglial large
extracellular vesicles propagate early synaptic dysfunction in Alzheimer's disease. Brain
J Neurol. 2022 Aug 27;145(8):2849–68.

15.   Karisetty BC, Bhatnagar A, Armour EM, Beaver M, Zhang H, Elefant F. Amyloid-β
Peptide Impact on Synaptic Function and Neuroepigenetic Gene Control Reveal New
Therapeutic Strategies for Alzheimer's Disease. Front Mol Neurosci. 2020 Nov 13;13.

16.   Du H, Guo L, Yan S, Sosunov AA, McKhann GM, Yan SS. Early deficits in synaptic
mitochondria in an Alzheimer's disease mouse model. Proc Natl Acad Sci U S A. 2010
Oct 26;107(43):18670–5.

17.   Scheff SW, Price DA, Ansari MA, Roberts KN, Schmitt FA, Ikonomovic MD, et al.
Synaptic Change in the Posterior Cingulate Gyrus in the Progression of Alzheimer's
Disease. J Alzheimers Dis. 2015 Jan 1;43(3):1073–90.

18.   Berchtold NC, Coleman PD, Cribbs DH, Rogers J, Gillen DL, Cotman CW. Synaptic
genes are extensively downregulated across multiple brain regions in normal human
aging and Alzheimer's disease. Neurobiol Aging. 2013 Jun;34(6):1653–61.

19.   Mattick JS, Makunin IV. Non-coding RNA. Hum Mol Genet. 2006 Apr
15;15(suppl_1):R17–29.

20.   Wang E, Lemos Duarte M, Rothman LE, Cai D, Zhang B. Non-coding RNAs in
Alzheimer's disease: perspectives from omics studies. Hum Mol Genet. 2022 Oct
15;31(R1):R54–61.

21.   Grillone K, Riillo C, Scionti F, Rocca R, Tradigo G, Guzzi PH, et al. Non-coding RNAs in
cancer: platforms and strategies for investigating the genomic "dark matter". J Exp Clin
Cancer Res. 2020 Jun 20;39(1):117.

22.   Zhang Y, Zhao Y, Ao X, Yu W, Zhang L, Wang Y, et al. The Role of Non-coding RNAs
in Alzheimer's Disease: From Regulated Mechanism to Therapeutic Targets and
Diagnostic Biomarkers. Front Aging Neurosci. 2021 Jul 2;13:654978.

23.   Statello L, Guo CJ, Chen LL, Huarte M. Gene regulation by long non-coding RNAs and
its biological functions. Nat Rev Mol Cell Biol. 2021 Feb;22(2):96–118.

24.   Fernandes JCR, Acuña SM, Aoki JI, Floeter-Winter LM, Muxel SM. Long Non-Coding
RNAs in the Regulation of Gene Expression: Physiology and Disease. Non-Coding RNA.
2019 Mar;5(1):17.

25.   Oo JA, Brandes RP, Leisegang MS. Long non-coding RNAs: novel regulators of cellular
physiology and function. Pflüg Arch - Eur J Physiol. 2022 Feb 1;474(2):191–204.

26.   DiStefano JK. The Emerging Role of Long Noncoding RNAs in Human Disease. In:
DiStefano JK, editor. Disease Gene Identification: Methods and Protocols. New York,
NY: Springer; 2018. p. 91–110.

27.   Wang X, Zhang M, Liu H. LncRNA17A regulates autophagy and apoptosis of SH-SY5Y
cell line as an in vitro model for Alzheimer's disease. Biosci Biotechnol Biochem. 2019
Apr 3;83(4):609–21.

28.   Zhang W, Zhao H, Wu Q, Xu W, Xia M. Knockdown of BACE1-AS by siRNA improves
memory and learning behaviors in Alzheimer's disease animal model. Exp Ther Med.
2018 Sep 1;16(3):2080–6.

29. Faghihi MA, Modarresi F, Khalil AM, Wood DE, Sahagan BG, Morgan TE, et al.
Expression of a noncoding RNA is elevated in Alzheimer's disease and drives rapid feed-
forward regulation of β-secretase expression. Nat Med. 2008 Jul;14(7):723–30.

30. Li F, Wang Y, Yang H, Xu Y, Zhou X, Zhang X, et al. The effect of BACE1-AS on β-
amyloid generation by regulating BACE1 mRNA expression. BMC Mol Biol. 2019 Oct
1;20(1):23.

31. Wang Q, Ge X, Zhang J, Chen L. Effect of lncRNA WT1-AS regulating WT1 on oxidative
stress injury and apoptosis of neurons in Alzheimer's disease via inhibition of the miR-
375/SIX4 axis. Aging. 2020 Nov 21;12(23):23974.

32. Mus E, Hof PR, Tiedge H. Dendritic BC200 RNA in aging and in Alzheimer's disease.
Proc Natl Acad Sci U S A. 2007 Jun 19;104(25):10679–84.

33. Quek XC, Thomson DW, Maag JL, Bartonicek N, Signal B, Clark MB, et al. lncRNAdb
v2.0: expanding the reference database for functional long noncoding RNAs. Nucleic
Acids Res. 2014 Oct 20;43(Database issue):D168.

34. Pavlopoulos GA, Secrier M, Moschopoulos CN, Soldatos TG, Kossida S, Aerts J, et al.
Using graph theory to analyze biological networks. BioData Min. 2011 Apr 28;4(1):10.

35. Liesecke F, De Craene JO, Besseau S, Courdavault V, Clastre M, Vergès V, et al.
Improved gene co-expression network quality through expression dataset down-
sampling and network aggregation. Sci Rep. 2019 Oct 8;9(1):14431.

36. Lee HK, Hsu AK, Sajdak J, Qin J, Pavlidis P. Coexpression Analysis of Human Genes
Across Many Microarray Data Sets. Genome Res. 2004 Jun 1;14(6):1085–94.

37. Berto S, Perdomo-Sabogal A, Gerighausen D, Qin J, Nowick K. A Consensus Network
of Gene Regulatory Factors in the Human Frontal Lobe. Front Genet. 2016 Mar 8;7:31.

38. Ghandikota S, Sharma M, Ediga HH, Madala SK, Jegga AG. Consensus Gene Co-
Expression Network Analysis Identifies Novel Genes Associated with Severity of Fibrotic
Lung Disease. Int J Mol Sci. 2022 May 13;23(10):5447.

39. Chai K, Zhang X, Tang H, Gu H, Ye W, Wang G, et al. The Application of Consensus
Weighted Gene Co-expression Network Analysis to Comparative Transcriptome Meta-
Datasets of Multiple Sclerosis in Gray and White Matter. Front Neurol. 2022 Feb 24;13.

40. Arshad Z, McDonald JF. A computational approach to generate highly conserved gene
co-expression networks with RNA-seq data. STAR Protoc. 2022 Jun 17;3(2):101432.

41. Zebardast F. Overview and illustration of the current study.Created in BioRender.
[Internet]. 2025 [cited 2024 Dec 25]. Available from: https://BioRender.com/k67x627

42. info@sagebase.org SB. Synapse | Sage Bionetworks [Internet]. [cited 2024 May 28].
Available from: https://www.synapse.org

43. Allen M, Carrasquillo MM, Funk C, Heavner BD, Zou F, Younkin CS, et al. Human whole
genome genotype and transcriptome data for Alzheimer's and other neurodegenerative
diseases. Sci Data. 2016 Oct 11;3(1):160089.

44. (Author Name Not Available). temporal cortex [Internet]. Synapse; 2016 [cited 2025 Jan
13].   Available   from: https://repo-
prod.prod.sagebase.org/repo/v1/doi/locate?id=syn3163039&type=ENTITY

45. Wang M, Beckmann ND, Roussos P, Wang E, Zhou X, Wang Q, et al. The Mount Sinai
cohort of large-scale genomic, transcriptomic and proteomic data in Alzheimer's disease.
Sci Data. 2018 Sep 11;5(1):180185.

46. (Author Name Not Available). MSBB [Internet]. Synapse; 2017 [cited 2025 Jan 13].
Available   from: https://repo-
prod.prod.sagebase.org/repo/v1/doi/locate?id=syn3159438&type=ENTITY

47. Braak H, Braak E. Neuropathological stageing of Alzheimer-related changes. Acta
Neuropathol. 1991;82(4):239–259.

48. Morris JC, Heyman A, Mohs RC, Hughes JP, van Belle G, Fillenbaum G, et al. The
Consortium to Establish a Registry for Alzheimer's Disease (CERAD). Part I. Clinical and
neuropsychological   assessment   of   Alzheimer's   disease.   Neurology.   1989
Sep;39(9):1159–65.

49. Morris JC. Clinical Dementia Rating: A Reliable and Valid Diagnostic and Staging
Measure for Dementia of the Alzheimer Type. Int Psychogeriatr. 1997 Dec;9(S1):173–6.

50. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinforma Oxf Engl. 2013 Jan 1;29(1):15–21.

51. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinforma Oxf Engl. 2014 Apr
1;30(7):923–30.

52. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550.

53. Stuart JM, Segal E, Koller D, Kim SK. A gene-coexpression network for global discovery
of conserved genetic modules. Science. 2003 Oct 10;302(5643):249–55.

54. Gysi DM, Voigt A, Fragoso T de M, Almaas E, Nowick K. wTO: an R package for
computing weighted topological overlap and a consensus network with integrated
visualization tool. BMC Bioinformatics. 2018 Oct 24;19(1):392.

55. Gysi DM, Fragoso T de M, Zebardast F, Bertoli W, Busskamp V, Almaas E, et al. Whole
transcriptomic network analysis using Co-expression Differential Network Analysis
(CoDiNA). PLOS ONE. 2020 Oct 15;15(10):e0240523.

56. Seidman SB. Network structure and minimum degree. Soc Netw. 1983 Sep 1;5(3):269–
87.

57. Zebardast F. Illustration of K-core decomposition method. Created in BioRender.
[Internet]. 2025 [cited 2024 Dec 25]. Available from: https://BioRender.com/w13e275

58. Traag VA, Waltman L, van Eck NJ. From Louvain to Leiden: guaranteeing well-connected
communities. Sci Rep. 2019 Mar 26;9(1):5233.

59. Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal enrichment
tool for interpreting omics data. Innov Camb Mass. 2021 Aug 28;2(3):100141.

60. Tong H, Faloutsos C, Pan JY. Random walk with restart: fast solutions and applications.
Knowl Inf Syst. 2008 Mar 1;14(3):327–46.

61. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a
software environment for integrated models of biomolecular interaction networks.
Genome Res. 2003 Nov;13(11):2498–504.

62. Bastian M, Heymann S, Jacomy M. Gephi: An Open Source Software for Exploring and
Manipulating Networks. Proc Int AAAI Conf Web Soc Media. 2009 Mar 19;3(1):361–2.

63. Morabito S, Miyoshi E, Michael N, Shahin S, Martini AC, Head E, et al. Single-nucleus
chromatin accessibility and transcriptomic characterization of Alzheimer's disease. Nat
Genet. 2021 Aug;53(8):1143–55.

64. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, et al.
Comprehensive Integration of Single-Cell Data. Cell. 2019 Jun 13;177(7):1888-
1902.e21.

65. Newman AM, Liu CL, Green MR, Gentles AJ, Feng W, Xu Y, et al. Robust enumeration
of cell subsets from tissue expression profiles. Nat Methods. 2015 May;12(5):453–7.

66. Mathys H, Davila-Velderrain J, Peng Z, Gao F, Mohammadi S, Young JZ, et al. Single-
cell transcriptomic analysis of Alzheimer's disease. Nature. 2019 Jun;570(7761):332–7.

67. Escorcia-Rodríguez JM, Gaytan-Nuñez E, Hernandez-Benitez EM, Zorro-Aranda A,
Tello-Palencia MA, Freyre-González JA. Improving gene regulatory network inference
and assessment: The importance of using network structure. Front Genet. 2023 Feb
27;14.

68. Ballouz S, Verleyen W, Gillis J. Guidance for RNA-seq co-expression network
construction and analysis: safety in numbers. Bioinforma Oxf Engl. 2015 Jul
1;31(13):2123–30.

69. Zhang B, Horvath S. A general framework for weighted gene co-expression network
analysis. Stat Appl Genet Mol Biol. 2005;4:Article17.

70. Voigt A, Almaas E. Assessment of weighted topological overlap (wTO) to improve fidelity
of gene co-expression networks. BMC Bioinformatics. 2019 Jan 28;20(1):58.

71. Morabito S, Reese F, Rahimzadeh N, Miyoshi E, Swarup V. hdWGCNA identifies co-
expression networks in high-dimensional transcriptomics data. Cell Rep Methods. 2023
Jun 12;3(6):100498.

72. He F, Maslov S. Pan- and core- network analysis of co-expression genes in a model
plant. Sci Rep. 2016 Dec 16;6(1):38956.

73. Kundra R, Ciryam P, Morimoto RI, Dobson CM, Vendruscolo M. Protein homeostasis of
a metastable subproteome associated with Alzheimer's disease. Proc Natl Acad Sci.
2017 Jul 11;114(28):E5703–11.

74. Zhang T, Liu N, Wei W, Zhang Z, Li H. Integrated Analysis of Weighted Gene
Coexpression Network Analysis Identifying Six Genes as Novel Biomarkers for
Alzheimer's Disease. Oxid Med Cell Longev. 2021 Jul 26;2021:9918498.

75. Lanke V, Moolamalla STR, Roy D, Vinod PK. Integrative Analysis of Hippocampus Gene
Expression Profiles Identifies Network Alterations in Aging and Alzheimer's Disease.
Front Aging Neurosci. 2018 May 23;10.

76. Wang M, Li A, Sekiya M, Beckmann ND, Quan X, Schrode N, et al. Transformative
Network Modeling of Multi-omics Data Reveals Detailed Circuits, Key Regulators, and
Potential Therapeutics for Alzheimer's Disease. Neuron. 2021 Jan 20;109(2):257-
272.e14.

77. Gu X, Lai D, Liu S, Chen K, Zhang P, Chen B, et al. Hub Genes, Diagnostic Model, and
Predicted Drugs Related to Iron Metabolism in Alzheimer's Disease. Front Aging
Neurosci. 2022 Jul 7;14:949083.

78. Duan K, Ma Y, Tan J, Miao Y, Zhang Q. Identification of genetic molecular markers and
immune infiltration characteristics of Alzheimer's disease through weighted gene co-
expression network analysis. Front Neurol. 2022 Aug 22;13:947781.

79. Majerníková N, den Dunnen WFA, Dolga AM. The Potential of Ferroptosis-Targeting
Therapies for Alzheimer's Disease: From Mechanism to Transcriptomic Analysis. Front
Aging Neurosci. 2021 Dec 20;13:745046.

80. Zhang Q, Yang P, Pang X, Guo W, Sun Y, Wei Y, et al. Preliminary exploration of the
co-regulation of Alzheimer's disease pathogenic genes by microRNAs and transcription
factors. Front Aging Neurosci. 2022 Dec 6;14.

81. Baglietto-Vargas D, Forner S, Cai L, Martini AC, Trujillo-Estrada L, Swarup V, et al.
Generation of a humanized Aβ expressing mouse demonstrating aspects of Alzheimer's
disease-like pathology. Nat Commun. 2021 Apr 23;12(1):2421.

82. Wang Z, Yan X, Zhao C. Dynamical differential networks and modules inferring disrupted
genes associated with the progression of Alzheimer's disease. Exp Ther Med. 2017 Oct
1;14(4):2969–75.

83. Finney CA, Delerue F, Gold WA, Brown DA, Shvetcov A. Artificial intelligence-driven
meta-analysis of brain gene expression identifies novel gene candidates and a role for
mitochondria in Alzheimer's disease. Comput Struct Biotechnol J. 2023 Jan 1;21:388–
400.

84. Neff RA, Wang M, Vatansever S, Guo L, Ming C, Wang Q, et al. Molecular subtyping of
Alzheimer's disease using RNA sequencing data reveals novel mechanisms and targets.
Sci Adv. 2021 Jan;

85. Zhang F, Zhong S, Yang S, Wei Y, Wang J, Huang J, et al. Identification of Potential
Therapeutic Targets of Alzheimer's Disease By Weighted Gene Co-Expression Network
Analysis. Chin Med Sci J. 2020 Dec 1;35(4):330–41.

86. Wu Z, Liu P, Huang B, Deng S, Song Z, Huang X, et al. A novel Alzheimer's disease
prognostic signature: identification and analysis of glutamine metabolism genes in
immunogenicity and immunotherapy efficacy. Sci Rep. 2023 Apr 27;13(1):6895.

87. Régal L, Mårtensson E, Maystadt I, Voermans N, Lederer D, Burlina A, et al. PREPL
deficiency: delineation of the phenotype and development of a functional blood assay.
Genet Med. 2018 Jan 1;20(1):109–18.

88. Männistö PT, García-Horsman JA. Mechanism of Action of Prolyl Oligopeptidase (PREP) in Degenerative Brain Diseases: Has Peptidase Activity Only a Modulatory Role on the Interactions of PREP with Proteins? Front Aging Neurosci. 2017 Feb 14;9.

89. Duarte ML, Wang M, Gomes I, Liu C, Sharma A, Fakira AK, et al. Multiomics Analyses Identify Proline Endopeptidase–Like Protein As a Key Regulator of Protein Trafficking, a Pathway Underlying Alzheimer's Disease Pathogenesis. Mol Pharmacol. 2023 Jul;104(1):1.

90. Heywood WE, Galimberti D, Bliss E, Sirka E, Paterson RW, Magdalinou NK, et al. Identification of novel CSF biomarkers for neurodegeneration and their validation by a high-throughput multiplexed targeted proteomic assay. Mol Neurodegener. 2015 Dec 1;10(1):64.

91. Shi Q, Gibson GE. Up-regulation of the mitochondrial malate dehydrogenase by oxidative stress is mediated by miR-743a. J Neurochem. 2011 Jun 24;118(3):440.

92. Rovelet-Lecrux A, Charbonnier C, Wallon D, Nicolas G, Seaman MNJ, Pottier C, et al. De novo deleterious genetic variations target a biological network centered on Aβ peptide in early-onset Alzheimer disease. Mol Psychiatry. 2015 Sep;20(9):1046–56.

93. Small SA, Kent K, Pierce A, Leung C, Kang MS, Okada H, et al. Model-guided microarray implicates the retromer complex in Alzheimer's disease. Ann Neurol. 2005 Dec;58(6):909–19.

94. Muhammad A, Flores I, Zhang H, Yu R, Staniszewski A, Planel E, et al. Retromer deficiency observed in Alzheimer's disease causes hippocampal dysfunction, neurodegeneration, and Abeta accumulation. Proc Natl Acad Sci U S A. 2008 May 20;105(20):7327–32.

95. Schlatterer SD, Tremblay MA, Acker CM, Davies P. Neuronal c-Abl Overexpression Leads to Neuronal Loss and Neuroinflammation in the Mouse Forebrain. J Alzheimer's Dis. 2011 Jan 1;25(1):119–33.

96. Lebouvier T, Scales TME, Williamson R, Noble W, Duyckaerts C, Hanger DP, et al. The Microtubule-Associated Protein Tau is Also Phosphorylated on Tyrosine. J Alzheimer's Dis. 2009 Jul 29;18(1):1–9.

97. Mastroeni D, Khdour OM, Delvaux E, Nolz J, Olsen G, Berchtold N, et al. Nuclear but not mitochondrial-encoded oxidative phosphorylation genes are altered in aging, mild cognitive impairment, and Alzheimer's disease. Alzheimers Dement. 2017 May 1;13(5):510–9.

98. La Barbera L, Vedele F, Nobili A, Krashia P, Spoleti E, Latagliata EC, et al. Nilotinib restores memory function by preventing dopaminergic neuron degeneration in a mouse model of Alzheimer's Disease. Prog Neurobiol. 2021 Jul;202:102031.

99. Zhou Z, Bai J, Zhong S, Zhang R, Kang K, Zhang X, et al. Downregulation of ATP6V1A Involved in Alzheimer's Disease via Synaptic Vesicle Cycle, Phagosome, and Oxidative Phosphorylation. Oxid Med Cell Longev. 2021 Apr 19;2021:5555634.

100. Geldenhuys WJ, Leeper TC, Carroll RT. mitoNEET as a novel drug target for mitochondrial dysfunction. Drug Discov Today. 2014 Oct 1;19(10):1601–6.

101. Dumont M, Kipiani K, Yu F, Wille E, Katz M, Calingasan NY, et al. Coenzyme Q10
decreases amyloid pathology and improves behavior in a transgenic mouse model of
Alzheimer's disease. J Alzheimers Dis JAD. 2011;27(1):211–23.

102. Choi H, Park HH, Koh SH, Choi NY, Yu HJ, Park J, et al. Coenzyme Q10 protects against
amyloid beta-induced neuronal cell death by inhibiting oxidative stress and activating the
P13K pathway. NeuroToxicology. 2012 Jan 1;33(1):85–90.

103. Sheykhhasan M, Amini R, Soleimani Asl S, Saidijam M, Hashemi SM, Najafi R.
Neuroprotective effects of coenzyme Q10-loaded exosomes obtained from adipose-
derived stem cells in a rat model of Alzheimer's disease. Biomed Pharmacother. 2022
Aug 1;152:113224.

104. Llanos-González E, Sancho-Bielsa FJ, Frontiñán-Rubio J, Rabanal-Ruíz Y, García-
Carpintero S, Chicano E, et al. Spatial and Temporal Protein Modules Signatures
Associated with Alzheimer Disease in 3xTg-AD Mice Are Restored by Early Ubiquinol
Supplementation. Antioxidants. 2023 Mar;12(3):747.

105. Li X, Xue Y, Liu X, Zheng J, Shen S, Yang C, et al. ZRANB2/SNHG20/FOXK1 Axis
regulates Vasculogenic mimicry formation in glioma. J Exp Clin Cancer Res CR. 2019
Feb 11;38:68.

106. Cohen OS, Weickert TW, Hess JL, Paish LM, McCoy SY, Rothmond DA, et al. A splicing-
regulatory polymorphism in DRD2 disrupts ZRANB2 binding, impairs cognitive
functioning and increases risk for schizophrenia in six Han Chinese samples. Mol
Psychiatry. 2016 Jul;21(7):975–82.

107. Malacards: The Human Disease Database - PMC [Internet]. [cited 2024 Nov 15].
Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC5764584/

108. Lin X, Lu Y, Zhang C, Cui Q, Tang YD, Ji X, et al. LncRNADisease v3.0: an updated
database of long non-coding RNA-associated diseases. Nucleic Acids Res. 2024 Jan
5;52(D1):D1365–9.

109. Adav SS, Park JE, Sze SK. Quantitative profiling brain proteomes revealed mitochondrial
dysfunction in Alzheimer's disease. Mol Brain. 2019 Jan 28;12(1):8.

110. Wang M, Roussos P, McKenzie A, Zhou X, Kajiwara Y, Brennand KJ, et al. Integrative
network analysis of nineteen brain regions identifies molecular signatures and networks
underlying selective regional vulnerability to Alzheimer's disease. Genome Med. 2016
Nov 1;8(1):104.

1121