Integrative Gene Co-expression Network Analysis Reveals Protein-Coding and lncRNA Genes Associated with Alzheimer's Disease Pathology

Fatemeh Zebardast[1,2], Michael Peter Sascha Riethmüller[1], Katja Nowick[1] (corresponding author, katja.nowick@fu-berlin.de)

1 Human Biology and Primate Evolution Group; Institute for Zoology; Department of Biology, Chemistry and Pharmacy; Freie Universität Berlin, Berlin, Germany.
2 International Max Planck Research School for Biology and Computation IMPRS-BAC, Berlin, Germany

Abstract

Alzheimer's disease (AD) is a complex neurodegenerative disorder marked by widespread molecular changes, many of which remain poorly understood. While AD pathology progresses through specific brain regions, it is unclear whether these regions are affected similarly. Long non-coding RNAs (lncRNAs), emerging as key cellular regulators, remain largely uncharacterized in AD. Understanding how lncRNAs interact with protein-coding genes across brain regions could shed light on AD mechanisms and progression. To investigate this, we performed consensus weighted gene co-expression network analysis on 396 postmortem brain RNA-seq samples using a meta-analytic approach. Our analysis revealed substantial network rewiring in AD, particularly in the temporal cortex compared to the frontal cortex. The temporal cortex exhibited adaptive changes in gene interactions, while the frontal cortex showed a breakdown of healthy correlations—possibly reflecting regional differences in disease progression. We identified 46 protein-coding genes and 27 lncRNAs as key components in the AD network of the temporal cortex. Using known functions of protein-coding genes as reference points, we inferred potential functions for over 100 lncRNAs across both regions. These findings highlight novel lncRNA candidates

32 potentially involved in AD and provide insights into their roles in both healthy and diseased

33 brain states.

35 **Keywords:** Alzheimer's disease, consensus gene co-expression network, weighted

36 Topological Overlap (wTO), long non-coding RNAs (lncRNA), Random Walk with Restart

37 (RWR).

Introduction

Alzheimer's disease (AD) is a complex, multifactorial neurodegenerative disorder

characterized by progressive cognitive decline and underlying pathological mechanisms (1).

Current knowledge indicates that the hallmark pathology of AD begins in the medial temporal

lobe (2) and, as the disease progresses, extends to other areas of the temporal cortex

before spreading to additional brain regions, including the frontal lobes, in the mid to late

stages (2–4). Its pathology disrupts numerous cellular mechanisms, such as mitochondrial

functions (5–7), endoplasmic reticulum (8,9), Golgi apparatus integrity (10,11), cellular

transport processes (5,12–14), and synaptic signaling (15–18). These widespread

dysfunctions closely align with alterations in gene expression and molecular interactions,

highlighting the complex biological networks that underlie AD.

Non-coding RNAs have attracted attention due to their potential involvement in both normal

and disease-related cellular processes (19–22). Among these, long non-coding RNAs

(lncRNAs)—transcripts longer than 200 nucleotides without apparent protein-coding

capacity—have emerged as critical regulators of cellular functions (23–26). LncRNAs

introduce an additional layer of complexity to gene regulatory processes through their

interactions with protein-coding genes. Studies have implicated lncRNAs in several AD-

associated dysfunctions, including amyloid beta (Aβ) production (27–30), oxidative stress

(31), and synaptic impairments (32). Despite the annotation of numerous lncRNAs with

known sequences and genomic locations, only a fraction have been extensively studied to

uncover their functional roles in both healthy and disease conditions (24,26,33), leaving a

significant knowledge gap yet to be addressed. A promising approach to address this gap is

the construction of gene co-expression networks, which can reveal functional relationships

between genes (34). Current methodologies face challenges such as mitigating noise and

improving the reliability of gene-gene correlation detection (35,36). Overcoming these

limitations is essential for generating accurate and interpretable biological insights. To

address these challenges, we integrated gene co-expression networks derived from the

same brain region across independent studies to construct consensus networks. This

approach consolidates shared signals from multiple datasets, reducing the impact of

dataset-specific noise and variability. By focusing on interactions consistently observed

across networks, this method enhances the detection of biologically relevant signals,

increasing confidence in their accuracy and interpretability compared to signals supported by

individual networks alone (37–40).

In this study, we utilized RNA-seq data from two independent studies, comprising five

datasets with 396 individuals total in AD and control groups, with gene expression profiles

from the temporal cortex (TCX) and frontal cortex (FCX). These profiles were used to

construct signed weighted topological overlap (wTO) networks, which were subsequently

integrated into consensus networks to enhance reliability and biological relevance (Figure 1).

Our analysis focused on lncRNAs and protein-coding genes, examining their correlations

within gene co-expression networks under both AD and non-AD (control) conditions.

Comparative analysis revealed significant rewiring of gene-gene interactions in the temporal

cortex in AD, contrasted with a marked loss of 'healthy' gene correlations in the frontal

cortex. Building on these observations, we identified 46 protein-coding genes and 27

lncRNAs as critical components of the AD network. A key objective of this study was to

predict the functions of lncRNAs and their associations with AD pathology. By leveraging the

established functions of protein-coding genes as network anchors, we assigned over 100

lncRNAs to biological processes in both AD and control conditions.

Figure 1: Overview and illustration of the current study (41)**.** From left to right, the top panel

illustrates the network construction and integration process. The bottom panel illustrates the network

analysis, including topology assessment, identification of hub genes, functional enrichment analysis,

assigning enriched GOs to lncRNAs within each cluster, and identifying key genes associated with

AD.

Results

Construction and integration of gene co-expression networks

We used a total of five RNA-seq datasets from two independent studies: the Mayo RNA-seq

study (Mayo) (42) and the Mount Sinai Brain Bank study (MSBB) (43). The gene expression

data were derived from two brain regions: TCX and FCX. These datasets were generated

using samples with detailed clinical-pathological diagnoses, including Braak NFT stage (44),

plaque density, CERAD (45), and CDR (46).

To minimize the impact of technical and unwanted biological factors, we implemented

several steps: 1- Filtering individuals based on AD assessment criteria (Figure 2a). 2- Sex

balance**:** each dataset includes an equal number of males and females. 3- Age balance**:** the

age range was balanced within each group, as well as across all groups, as much as

possible (Figure 2b-c). Furthermore, we uniformly processed all data from FASTQ files to

minimize the impact of technical factors.

Figure 2: Summary of data demographics. a) Parameters and thresholds used to classify

individuals into AD and control groups include Braak stage (based on neurofibrillary tangles

distribution), CERAD (scoring neuritic plaques) from 0 (none) to 3 (frequent), and CDR (Clinical

Dementia Rating) 0 (no dementia) to 3 (severe dementia). **b)** Illustration of brain areas used in this

study that are temporal (yellow) and frontal (blue) cortices. **c)** The datasets, sample size, and

Brodmann (BM) brain area utilized in this study. **d)** Data demography after data preparation: sample

sizes were balanced between AD and control groups, as well as between sexes. Efforts were made to

maintain a balanced age distribution. **e)** Distribution of wTO values for each network before filtering

links for a cutoff. Red dashed lines indicate weight cutoffs. **f,g)** The graphs illustrate the number of

links and nodes before and after filtering links and integration into consensus networks (x-axis:

Before, After and Consensus). Each line and node represent one network, distinguished by colors for

AD and control networks (see Tables S1 and S2 for more details).

Normalized gene count data were then used to construct signed weighted gene co-

expression networks using the wTO method (47) separately for each AD and control dataset.

We used gene count data for both protein-coding and lncRNA genes as input, resulting in

networks that capture correlations between protein-coding and lncRNA genes. To focus on

the strongest links, we considered the summary statistics of absolute weights of links in the

networks (Figure 2e, supplementary Table S1) and excluded all weak links with an absolute

weight below 0.5. This substantially reduced the number of links from a median of 219

million to 7.2 million links (Figure 2f). While the number of nodes also decreased, the

reduction was less drastic, dropping from 20,900 nodes before filtering to a median of

13,200 nodes afterward (Figure 2g). This implied that most genes remained in the network,

but only the most robust interactions were represented.

Subsequently, the filtered networks were integrated separately for AD and control groups

based on brain regions to further enhance the reliability of the networks, excluding links that

might be dataset/study specific. Specifically, the Mayo TCX network was integrated with two

MSBB networks, BM36 and BM22, all derived from different parts of the temporal cortex, into

a consensus TCX network. Similarly, the BM10 and BM44 networks, which originated from

two distinct areas of the frontal cortex, were integrated to form the consensus FCX network

(Figure 2 b-c, supplementary Table S2). The resulting consensus networks included only the

commonalities across the integrated networks for each brain region. This method ensured

that only gene relations observed in multiple studies were included in our further analyses.

From here on, we refer to these consensus networks as the "AD" and "control" networks of

the TCX and FCX, respectively.

Comparative analysis of the TCX networks

We began our analysis with the TCX networks. Using K-core decomposition, we identified

nodes in the innermost shell as core genes (hubs) in both AD and control networks

(supplementary Table S3 and S4). The TCX AD network had 306 core genes, compared to

235 in the TCX control network, with 18% (81 genes) shared between them (Figure 3a).

Notably, lncRNAs accounted for 15% of the core genes in the AD network, nearly double the

8% observed in the control network (Figure 3b). Statistical evaluation using Fisher's exact

test confirmed the significant enrichment of core lncRNAs in the AD network (p-value =

0.016, odds ratio = 2.01). Among the TCX AD core genes, 22 were significantly differentially

expressed (DE, FDR < 0.05 and absolute log fold change> 0.3) across all three TCX

datasets, all of which were downregulated in AD (Figure 3 c and d). These included 17

protein-coding and 5 lncRNA core genes. A complete list of DE genes is provided in

supplementary Table S5.

Figure 3: Demography and differential expression analysis of TCX core genes. a) Venn diagram

depicting the overlap of core genes between AD (orange) and control (green) networks in TCX. **b)**

Proportion of protein-coding and lncRNA genes within cores of TCX AD network (orange) and control

network (green). **c** and **d)** Heatmaps showing the log2 fold change (FC) for differentially expressed

(DE) core genes in AD, including protein-coding (c) and lncRNA (d) genes that are in the TCX AD

network. Each column represents the log2 FC for a single dataset used to construct the TCX

networks, which are Mayo, MSBB BM22 and BM36.

To examine the structure and underlying biological functions of the networks, we applied the

Louvain method for community detection to each consensus network separately. In the TCX

AD network, we observed three gene clusters, whereas the control network formed five

clusters (Figure 4a, Table 1). GO enrichment analysis was conducted on the protein-coding

genes within each cluster, using the same set of background genes (see Methods). Enriched

GO terms were manually curated and grouped into broader functional categories to address

GO term redundancy (see Methods for details). Significantly enriched biological processes

fell into eight broad functional categories, including 1- mitochondria energy production, 2-

intracellular trafficking, 3- synapse signaling, 4- neurogenesis, 5- apoptosis, 6- cytoskeleton

organization 7- gene expression, 8- cellular homeostasis, which were enriched to differing

degrees among the clusters (Figure 4b, supplementary Table S6). Additionally, four GO

terms were grouped as "others" due to their specificity or broad relevance (e.g., GTPase

activity). Among all clusters, only genes of control cluster 5 were not significantly enriched

for any terms. Cellular homeostasis was the only functional category that was enriched only

in control clusters.

Table 1: The topological details of AD and control clusters in the temporal cortex

	Cluster 1		Cluster 2		Cluster 3		Cluster 4		Cluster 5	
	Nodes	Links	Nodes	Links	Nodes	Links	Nodes	Links	Nodes	Links
control	961	18,583	1,662	31,509	836	4,702	113	169	894	30,439
AD	920	48,198	1,689	44,374	476	17,262				

Figure 4: Topological and differential co-expression analysis of TCX networks. a)

Representation of clusters for TCX AD and control networks, shown on the right and left, respectively.

There are two node sizes; larger nodes represent core hub genes, while smaller nodes represent the

remaining genes. Each color represents a distinct cluster. Gray backgrounds indicate links, which are

faded to reduce complexity for visualization. The networks were visualized by Gephi (48). **b)** The plot

displays selected enriched GO terms and their classification into broader functional categories. The

significance of enriched GOs was assessed using the false discovery rate, and terms with an adjusted

p-value below 0.05 were selected. Dot colors indicate the adjusted p-value, and dot size is

proportional to the number of genes of interest in the input gene list for the given term. **c)** Distribution

of differential co-expression link categories within each cluster. The percentages represent relative

abundance, meaning the original proportion of link types within each cluster is normalized relative to

the cluster size. **d)** Distribution of differential co-expression links categories connecting genes

associated with each functional category. **e)** Distribution of protein-coding core genes annotated to the

enriched GO terms.

In order to gain a deeper understanding of the differences between AD and control networks,

a differential co-expression network analysis was performed (49). We classified links into

three categories: 1- conserved links, which were common between the two conditions (i.e.,

AD and control); 2- condition-specific links, which appeared only in one condition (i.e., AD-

specific); and 3- reversed links, which represented correlations with opposite signs in the two

conditions. The results showed that 96%, 77%, and 83% of links in AD clusters 1, 2, and 3,

respectively, were AD-specific (Figure 4c). This pattern was also reflected in links between

genes of each functional category in AD clusters (Figure 4d). These findings indicated that

AD-related changes are not confined to a single cluster or GO term; rather, they broadly

impacted all three gene clusters and the identified underlying biological process. Similar

observations were seen in the control network analysis, indicating that the majority of links

per cluster and functional category were control-specific and, in other words, were absent in

AD. There was only one exception: 51% of control cluster 1 included conserved links, which

involved genes annotated to mitochondria energy production and apoptosis.

The observed rewiring of the AD network suggested a shift in the organization of molecular

interactions. To investigate the key players of this rewiring, we examined all core genes, as

these genes were likely central to the disrupted processes and extensive rewiring observed

in the AD network. Notably, of the protein-coding core genes, a higher proportion was

annotated to the functional categories in the AD network (18%, 46 genes) compared to the

control network (2%, 5 genes) (Figure 4e). Among the core genes linked to the enriched

GOs, the largest proportion (34%) were associated with mitochondrial energy production,

followed by 22% with synaptic signaling and 15% with intracellular trafficking (Figure 4e).

While the majority of the protein-coding core genes in both networks were not annotated to

enriched GO terms, the increased representation of AD core genes in these categories

underscored their potential roles in key processes disrupted in AD pathology, particularly in

energy production, neural communication, and intracellular transport.

Among the 46 core genes of the AD network annotated to the enriched GO terms (Table 2,

Figure 5), nearly all were exclusively core to the AD network, except for four genes. Notably,

these four genes—*CHL1* and *USP33* (neurogenesis), *GRIA2* (synapse signaling), and

YARS2 (mitochondrial gene expression)—were annotated to biological processes only within

the AD network and not in the control network. This observation suggests that these genes

might gain a disease-related role in the context of AD. It might be observed that a single

gene was annotated to different functional categories in the AD and control networks (Table

2). This was explained by the fact that the gene clustered with different sets of genes in the

two networks, leading to its association with distinct GO terms. This observation reflected the

differing patterns of gene interactions between AD and control networks, likely linked to the

disease condition.

234
235 **Table 2: Identified key protein-coding genes potentially associated with AD pathology in TCX.**
236

Gene name	KEGG pathway	DEG	AD		Control		hub
			Cluster	Functional category	Cluster	Functional category	
ABL1	-	-	3	Cytoskeleton organization	1	-	-
ACSL3	-	-	1	Mitochondrial energy production	1	Mitochondrial energy production	-
AK6	-	-	1	Mitochondrial energy production	1	Mitochondrial energy production	-
AP3M2	-	Down	1	Synapse signaling	1	-	-
ATL1	-	-	2	Neurogenesis	-	-	-
ATP6AP2	-	-	1	Mitochondrial energy production, synapse signaling	1	Intracellular trafficking, synapse signaling	-
ATP6V1A	-	-	1	Intracellular trafficking, mitochondrial energy production, synapse signaling	1	Intracellular trafficking, mitochondrial energy production, synapse signaling	-

ATP6V1B2	-	Down	1	Intracellular trafficking, mitochondrial energy production, synapse signaling	1	Intracellular trafficking, Mitochondrial energy production, synapse signaling	-
ATP6V1D	-	-	1	Intracellular trafficking, mitochondrial energy production, synapse signaling	1	Intracellular trafficking, synapse signaling	-
BLOC1S6	-	-	1	Synapse signaling	1	-	-
CCNB1	-	-	1	Mitochondrial energy production	1	Mitochondrial energy production	-
CHL1	-	-	2	Neurogenesis	5	-	core
CISD1	-	-	1	Mitochondrial energy production	1	Mitochondrial energy production	-
COQ10B	-	-	1	Mitochondrial energy production	1	Mitochondrial energy production	-
EPM2AIP1	-	-	1	Mitochondrial energy production	1	Mitochondrial energy production	-
ETFRF1	-	-	1	Mitochondrial energy production	1	Mitochondrial energy production	-
EXOC8	-	-	1	Intracellular trafficking	1	-	-
GHITM	-		1	Apoptosis, mitochondrial energy production, others (mitochondrial membrane organization)	1	Apoptosis, mitochondrial energy production	-
GLRX2	-		1	Mitochondrial energy production	1	Mitochondrial energy production	-
GRIA2	-		2	Synapse signaling	2	-	core
GUCY1B1	-		1	Mitochondrial energy production	1	Mitochondrial energy production	-
MDH1	-		1	Mitochondrial energy production	1	Mitochondrial energy production	-
MKKS	-		3	Cytoskeleton organization	1	-	-
MRPL42	-		1	Gene expression	1	-	-
MRPS30	-	-	1	Gene expression	1	-	-
MRPS35	-	-	1	Gene expression	1	-	-
NAPB	-	-	1	Synapse signaling	1	-	-
NDUFA5	AD pathway	Down	1	Mitochondrial energy production	1	Mitochondrial energy production	-
NDUFA9	AD pathway	-	1	Mitochondrial energy production	1	Mitochondrial energy production	-
NDUFB5	AD pathway	Down	1	Mitochondrial energy production	1	Mitochondrial energy production	-
NECAP1	-	Down	1	Synapse signaling	1	-	-
PFN2	-	Down	1	Synapse signaling	1	-	-

Gene	KEGG	DEG	cluster	functional category	cluster	functional category	hub
PREPL	-	-	1	Synapse signaling	1	-	-
PRPS2	-	-	1	Mitochondrial energy production	1	Mitochondrial energy production	-
RFK	-	Down	1	Mitochondrial energy production	1	Mitochondrial energy production	-
SCOC	-	-	1	Intracellular trafficking	1	-	-
SKP1	-	-	1	Intracellular trafficking	1	-	-
TAF9	-	-	1	Apoptosis	1	-	-
TMEM14A	-	-	1	Apoptosis, others (mitochondrial membrane organization)	1	Apoptosis	-
TMEM33	-	-	1	Intracellular trafficking, synapse signaling	1	-	-
TOMM70	-	-	1	others (mitochondrial membrane organization)	1	-	-
TRMT10C	-	-	1	Gene expression	1	-	-
UBE2T	-	-	1	Intracellular trafficking	1	-	-
USP33	-	-	2	Neurogenesis	5	-	core
VPS35	-	-	1	Apoptosis, intracellular trafficking, synapse signaling	1	Apoptosis	-
YARS2	-	-	1	Gene expression	1	-	core

237 The table lists 46 protein-coding key genes potentially associated with AD pathology in the TCX AD

238 network. Various features of these genes are listed, including their involvement in the KEGG AD

239 pathway, differential expressions in AD (DEG), cluster membership, as well as their annotation to the

240 functional categories. All are AD hubs, while their hub status in control is indicated under the hub

241 column.

lncRNA functional assignment and core lncRNAs in TCX networks

The gene co-expression networks in this study included both protein-coding and lncRNA

genes, providing us with the opportunity to gather functional information for lncRNAs based

on their network neighbors. For each functional category, Random Walk with Restart (RWR)

was run using annotated protein-coding genes as seeds. LncRNAs were ranked based on

their RWR scores, and the top 10th percentile were assigned to the corresponding functional

category. To ensure the biological relevance of these assignments, a leave-one-out

approach was applied to validate the rankings. A lncRNA was assigned to a category only if

its RWR score exceeded the lowest score from the leave-one-out simulation (Methods,

supplementary Figure S1 and Table S8). Using this approach, 54 lncRNAs in the TCX AD

network and 103 in the TCX control network were assigned to functional categories within

their respective clusters (supplementary Table S8).

Out of the 46 AD core lncRNA genes, we focused on 27 lncRNAs that were assigned to at

least one functional category, as these lncRNAs likely played key roles in the dysregulation

of their assigned functional categories within the AD network (Table 3, Figure 5). All but six

of these lncRNAs were AD-specific core genes. Notably, *FBXW7-AS1*, *Lnc-PDE4D-2*, *Lnc-*

KLHL11-1,   *Lnc-NIF3L1-5*,   *Lnc-KALRN-1*,   *Lnc-CCDC68-1*,   *HSALNG0099005*,

HSALNG0105911, *ZRANB2-DT*, *ENSG00000287022*, and *SLC8A1-AS1* were functionally

assigned only within the AD network, suggesting a gain of function for these lncRNAs in the

context of AD. The remaining lncRNAs were functionally assigned in both networks, either to

similar or different functional categories (Table 3). Many of these lncRNAs were assigned to

all functional categories within their respective clusters, as expected given their status as

core genes with high connectivity to numerous nodes, which often results in high RWR

scores. However, a closer examination of their functional categories and comparisons with

the control network revealed considerable differences. This suggests that lncRNAs were

incorporated into different neighborhoods or changed the sets of genes they regulated in AD

compared to control brains. For example, *POC1B-AS1* was assigned to cytoskeleton

organization-related processes in the AD network, whereas it was associated with

mitochondrial energy production in the control network. Among these 27 core lncRNAs, five

were   downregulated   in   AD,   namely   *ITFG1-AS1*,   *Lnc-MYCN-6*,   *Lnc-EFR3A-6*,

HSALNG0089044, and *Lnc-CCDC68-1* (Figure 3d, Table 3).

Table 3: Key lncRNA genes potentially associated with dysregulated GO terms in the TCX AD
network.

| Gene name | DEG | AD network | | Control network | | |
		Cluster	Functional category	Cluster	Functional category	Hub
FBXW7-AS1	-	1	Synapse signaling	1	-	-
Lnc-PDE4D-2	-	1	Apoptosis, gene	1	-	-

			expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling		
Lnc-KLHL11-1	-	1	Apoptosis, gene expression, mitochondrial energy production, others (mitochondrial membrane organization)	1	-	-
Lnc-NIF3L1-5	-	3	Cytoskeleton organization	1	-	-
Lnc-KALRN-1	-	3	Cytoskeleton organization	2	-	-
Lnc-CCDC68-1	Down	3	Cytoskeleton organization	1	-	-
HSALNG0099005	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	Absent	-	-
HSALNG0105911	-	2	Neurogenesis, synapse signaling	5	-	core
ZRANB2-DT	-	2	Neurogenesis, synapse signaling	5	-	core
ENSG00000287022	-	3	Cytoskeleton organization	5	-	core
SLC8A1-AS1	-	3	Cytoskeleton organization	5	-	core
HSALNG0096948	-	1	Intracellular trafficking	1	Intracellular trafficking, synapse signaling	-
ITFG1-AS1	Down	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Apoptosis, intracellular trafficking, mitochondrial energy production, others , synapse signaling	-
Lnc-MYCN-6	Down	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Intracellular trafficking, synapse signaling	-
Lnc-EFR3A-6	Down	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Apoptosis, mitochondrial energy production, others	-
HSALNG0089044	Down	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), Synapse	1	Intracellular trafficking, mitochondrial energy production, others , synapse signaling	-

			signaling			
Lnc-LRP11-1 (ENSG00000285889)	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Apoptosis, intracellular trafficking, mitochondrial energy production, others[*], synapse signaling	-
Lnc-KBTBD3-1	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Apoptosis, intracellular trafficking, mitochondrial energy production, others[*], synapse signaling	-
SNX10-AS1	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Apoptosis, mitochondrial energy production	-
Lnc-LRP11-1 (ENSG00000231760)	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Intracellular trafficking, others[*], synapse signaling	-
Lnc-KCNH5-3 ant	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Intracellular trafficking	-
Lnc-CLUH-2	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Others[*]	-
Lnc-SUMF1-5	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), Synapse signaling	1	Others[*]	-
HSALNG0144080	-	1	Apoptosis, gene expression, intracellular trafficking, mitochondrial energy production, others (mitochondrial membrane organization), synapse signaling	1	Intracellular trafficking, mitochondrial energy production, others[*], Synapse signaling	-
POC1B-AS1	-	3	Cytoskeleton organization	1	Mitochondrial energy production	-

MYCBP2-AS1	-	3	Cytoskeleton organization	1	Apoptosis, intracellular trafficking, mitochondrial energy production, synapse signaling	core
HSALNG0018143	-	2	Neurogenesis, synapse signaling	1	Apoptosis, mitochondrial energy production, others	core

275
276 The table shows the 27 selected AD core lncRNAs and their functional categories in the TCX AD and

277 control networks, as well as cluster membership and differential expression in AD. Down in the DEG

278 column indicates downregulation in AD. others[*]: positive regulation of cyclin-dependent protein

279 serine/threonine kinase activity.

280
Figure 5: AD cluster 1 network visualization. Node colors indicate functional categories;

multicolored nodes represent genes annotated to multiple categories. Larger nodes are identified as

key genes in the AD network, while smaller nodes represent other genes in AD cluster 1. Circles

represent protein-coding genes, and squares represent lncRNA genes. Link colors: blue links are

conserved between the TCX AD and control network, and black links are AD-specific links. Names of

key genes are enlarged; names of other genes are visible upon full zoom. Further details can be

explored through the interactive Shiny app we have provided.

Comparing TCX and FCX

After analyzing networks of the temporal cortex, we performed the same analysis for the

frontal cortex and compared our results between the two brain areas. In the frontal cortex,

we found that unlike in the temporal cortex, the AD network shows much higher similarity to

the control network (Figure 6 a-d), where the majority of links are conserved in the AD

network (86%, figure 6a). Conversely, the control network is predominantly composed of

control-specific links (83%), with minimal overlap (14%) with links found in the AD network

(Figure 6b). This observation implied that the AD network in the frontal cortex has lost a

significant number of gene-gene correlations. This hypothesis was supported by the marked

difference in the size of the networks: the AD network contained 1.17 million links, while the

control network contained 5.93 million links. This pattern was also evident in the significant

overlap between AD core hubs and control core genes (Figure 6e). While this overlap

suggested that certain gene relationships were preserved in AD, despite the extensive loss

of specific gene-gene interactions, a substantial number of core genes presented in the

control network were absent in the AD network. Interestingly, the proportion of gene types,

i.e., protein-coding or lncRNA genes, among the core genes {Citation}was very similar

between AD and control networks, with 15% or 18% of cores being lncRNAs, respectively

(Figure 6f).

We used the same method as for the TCX to cluster the FCX consensus networks

(supplementary Table S9). As done for the TCX, significantly enriched GO terms were then

manually classified into five broader functional categories (Figure 6g) including: 1-

mitochondrial energy production, 2-intracellular trafficking, 3- synapse signaling, 4-

apoptosis, 5- gene expression. Additionally, an "other" category was included, containing a

single term, mitochondrial membrane organization. A complete list of enriched GO terms can

be found in supplementary table S10. Genes in clusters 1 and 3 of both AD and control

networks were significantly enriched for biological process GO terms, but not in the other

clusters.

Differential gene co-expression analysis revealed a dominance of conserved links

connecting genes associated with enriched GO terms within AD clusters (Figure 6h). Each

functional category showed 94% to 100% conserved links. In contrast, in the control

network, the majority of links were control-specific within the functional categories. One

possible interpretation of this tendency was that the pathogenic alterations associated with

AD disrupted numerous healthy gene expression correlations in the frontal cortex area. This

might also suggest that AD-specific correlations represent cellular compensation against AD-

related disruptions.

In bulk RNA-Seq data, observed differences in gene expression can often be attributed to

variations in cell-type composition. Given that such differences might also contribute to

alterations in gene co-expression networks, we sought to assess the cellular composition of

the bulk RNA-seq data used in this study. To this end, we performed a deconvolution

analysis on both the TCX and FCX regions using CIBERSORT, utilizing human brain single-

cell data as a reference (Figure 6i). Across both brain regions and independent of disease

status, excitatory neurons were the predominant cell type, comprising 41–56% of the total

population. However, the relative abundance of other cell types varied between TCX and

FCX, with microglia consistently being the least abundant in both regions. The

overrepresentation of excitatory neurons and the underrepresentation of microglia align with

previous findings in the human brain, indicating that bulk RNA-seq data primarily capture

transcriptional changes in excitatory neurons, while microglia, due to their lower RNA

content, are typically underrepresented (50).

To assess whether AD pathology affects cell-type composition, we compared cell-type

proportions between AD and control samples within each brain region. Interestingly, we

observed significant alterations only in TCX, where excitatory neurons were markedly

decreased (p.adj = $9.0 \times 10^{-\square}$), while astrocytes (p.adj = $4.66 \times 10^{-\square}$), microglia (p.adj = 5.0

$\times 10^{-\square}$), oligodendrocytes (p.adj = 0.03), and endothelial cells (p.adj = $2.07 \times 10^{-\square}$) were

significantly increased (Figure 6j). This shift in cellular composition may contribute to the

gene co-expression rewiring observed in the TCX AD network.

Figure 6: Results of comparative analysis between FCX and TCX. Pie chart showing proportions

of differential co-expressed links **a)** in the FCX AD network, **b)** FCX control network, **c)** TCX AD

network and **d)** TCX control network. **e** Venn diagram showing the overlap of core genes between AD

and control networks in FCX. **f)** Proportion of protein-coding and lncRNA genes within FCX core

genes. **g)** Significantly enriched GO terms for biological processes, grouped into five broader

functional categories: 1) mitochondrial energy production, 2) intracellular trafficking, 3) synapse

signaling, 4) apoptosis, 5) gene expression, with an additional "other" category with a single term-

mitochondrial membrane organization. **h)** Distribution of differential link categories within each

functional category, with percentages representing relative abundance normalized to cluster size. **i)**

Stacked bar plots showing the estimated cell-type proportions in FCX and TCX for AD and control

samples. Each color represents one cell type. **j)** Significant cell-type proportion changes between AD

and controls were observed only in TCX datasets. P-values were measured by the Wilcoxon test (*p <

0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001).

In addition, the FCX control network was employed to evaluate the reliability and consistency

of the lncRNA annotation method implemented in this study. We investigated the similarity of

lncRNAs annotated to the same functional category between the TCX and FCX control

networks (supplementary Table S11). Five functional categories common to both networks

were analyzed, including mitochondrial energy production, intracellular trafficking, synapse

signaling, apoptosis, and gene expression (supplementary Table S12). Significant overlaps

of lncRNAs assigned to functional categories were observed between the FCX and TCX

control networks, with p-values approaching zero (Table 4). The analysis accounted for the

disparity in lncRNA totals between FCX and TCX control networks (1100 and 504,

respectively) and a permutation test for statistical significance. Notably, the gene expression

category was the only one without any overlapping lncRNAs. Additionally, permutation tests

were conducted for lncRNAs not assigned to functional categories. These tests yielded a

non-significant p-value of 0.71, highlighting that the functional assignments of annotated

lncRNAs are significantly more consistent than would be expected by chance.

Table 4: Similarity of lncRNA annotations across FCX and TCX control networks.

Functional category	Overlap coefficient	p-value
Apoptosis	0.47	p < 1e-03
Synapse signaling	0.41	p < 1e-03
Mitochondrial energy production	0.3	p < 1e-03
Intracellular trafficking	0.2	p < 1e-03
Gene expression	0	1

| Not assigned | 0.12 | 0.71 |

373　The table displays the overlap coefficients between lncRNAs assigned to common functional

374　categories between FCX and TCX control networks, calculated using the Szymkiewicz-Simpson

375　coefficient. Permutation tests with 1000 random permutations were used to assess statistical

376　significance. P-values shown as '< 1e-03' indicate no permutation matched or exceeded the observed

377　overlap, implying p-values smaller than the test resolution (p < 1/1000).

Discussion

In this study, we presented robust consensus gene co-expression networks including

protein-coding and lncRNA genes of two brain regions (temporal cortex and frontal cortex)

and investigated how the networks differ in AD compared to control brains. Our deep

comparative analysis between the consensus networks of AD and control groups highlighted

significant rewiring of gene-gene interactions within the temporal cortex area (Figure 4c, d),

in contrast to a more pronounced loss of 'healthy' gene correlations in the frontal cortex area

(Figure 6 a-d). Prompted by these observations, we identified key players in the TCX AD

network, including 46 protein-coding genes and 27 lncRNAs (Table 2 and 3). Utilizing the

established functions of protein-coding genes as an anchor, we bridged lncRNAs to the

enriched biological processes, demonstrating the utility of network-based approaches. We

not only nominated potential key lncRNAs involved in AD pathology but also predicted their

roles in biological processes under both AD and non-AD conditions.

Networks provide a powerful framework for studying the structure of complex biological

systems, offering valuable insights into gene-gene interactions. However, network analysis

poses challenges, such as managing noise signals and enhancing the reliability of

correlations. To address these issues, we employed several strategies to mitigate biases

and increase robustness (Figure 1). First, we ensured that all datasets used had a minimum

of 30 replicates per condition (Figure 2c), as larger sample sizes improve the accuracy of

network inference (51,52). Second, we applied the wTO method (53) to compute gene-gene

correlations. This approach accounts for shared network neighborhoods, producing more

robust scores compared to simple correlation calculations, and incorporates both positive

and negative correlations (54). The gene co-expression networks were subsequently filtered

based on the absolute value of gene pair correlations. Following this, we integrated the

networks within each brain region (TCX and FCX) (Figure 2b, c). This integration allowed us

to treat them as biological replicates, identifying common interactions across the integrated

networks and reducing the impacts of sampling and batch variability, a widely adopted

approach in gene co-expression studies (37–40,55,56). Ultimately, this consensus network

offers a high-resolution representation of the conserved cellular processes functioning under

AD and control conditions, providing a robust foundation for further investigation.

Co-expressed genes are often coregulated or involved in similar pathways, forming densely

interconnected clusters within the network. In this study, we identified gene clusters and their

functions for each consensus network. Through a comprehensive comparative analysis of

gene correlations and network topology between AD and control consensus networks, we

observed remarkable network alterations. Specifically, in the TCX networks, substantial

rewiring was evident within the AD clusters, which was also reflected in the links between

protein-coding genes associated with enriched terms (Figure 4c, d). The emergence of new

core genes (Figure 3a), along with their higher engagement in the AD network compared to

the control network (18% versus 2%, Figure 4e), indicated a reorganization of the molecular

landscape in AD, suggesting that specific genes adopted new roles within the altered

regulatory environment. As a result, we identified 46 core protein-coding genes as

candidates potentially playing a central role in the rewiring observed in enriched terms in the

TCX AD network (Table 2).

Several of these genes have been supported by prior experimental or network-based studies

as being linked to AD. Notably, 15 of these genes have been reported as hub genes in AD,

including *ATL1* (57), *AP3M2* (58)*, ATP6AP2* (59), *ATP6V1A* (60), *ATP6V1D* (61,62), *ACSL3*

(58,63), *CCNB1* (64), *GHITM* (62), *GRIA2* (65), *MDH1* (66), *MRPS30* (67), *MRPS35* (66),

NAPB (68,69), *NDUFB5* (66) and *PFN2* (70). Several genes showed strong evidence

supporting their role in AD pathology. For example, *REPL* knockdown led to increased Aβ

levels and tau phosphorylation (71–73), *NAPB* was linked to neuroinflammatory neuronal

damage (74), *MDH1* correlated with total and phosphorylated tau levels (74,75) and *VPS35*

deficiencies or mutations caused altered endosomal trafficking and pathological tau

accumulation (13,76–78). Of particular interest, *ABL1* (c-Abl) contributed to tau pathology

through regulation of cytoskeletal signaling cascades (79,80). Our findings identified *ABL1*

as an AD-specific hub annotated to the stress fiber assembly term (Table 2). Notably, about

half of the 46 AD key genes were annotated to the mitochondrial energy production

functional category. Among them, *NDUFA5* and *NDUFB5*, encoding subunits of Complex I of

the mitochondrial respiratory chain, were consistently downregulated across TCX datasets

(Figure 3c), aligning with their known significant downregulation in AD (81). Recent findings

revealed that the endolysosomal V-ATPase acts as a receptor for Aβ and Tau in AD (14).

Among the 13 genes involved in the V-ATPase complex, four—*ATP6V1B2*, *ATP6V1A*,

ATP6AP2, and *ATP6V1D*—were in our key gene list. In addition to previously known AD-

associated genes, several key genes identified in this study have been pursued as promising

therapeutic targets for AD, including *ABL1* (82), *ACSL3* (63), *ATP6V1A* (60,83), *CISD* (84)

and *COQ10B* (85–88).

Having identified many previously known AD-associated protein-coding genes with our

network analysis, we leveraged the connectivity of lncRNAs with their neighboring protein-

coding genes within the network to infer potential functions for these lncRNAs. Through this

approach, 54 lncRNAs in the TCX AD network and 103 in the FCX control network were

assigned to functional categories identified within each cluster (supplementary Table S8 and

S11). The assigned biological functions were cross-validated by assessing the similarity of

lncRNAs assigned to common functional categories between the TCX and FCX control

networks, revealing significant overlap (Table 4). From the TCX AD network, 27 core

lncRNAs with functional assignments were selected for further investigation (Table 3). We

observed that all but six of these 27 lncRNAs were AD-specific core genes. Notably,

FBXW7-AS1, *Lnc-PDE4D-2*, *Lnc-KLHL11-1*, *Lnc-NIF3L1-5*, *Lnc-KALRN-1*, *Lnc-CCDC68-1*,

HSALNG0099005, *HSALNG0105911*, *ZRANB2-DT*, *ENSG00000287022*, and *SLC8A1-AS1*

were functionally assigned exclusively within the AD network, suggesting a gain of function

for these lncRNAs in the context of AD. Among them, *ZRANB2-DT*, assigned to

neurogenesis and synapse signaling, stood out. This lncRNA acts as an antisense RNA for

ZRANB2 (Zinc Finger RANBP2-Type Containing 2), a gene previously reported to be

associated with neuronal responses to chronic oxidative stress—a process closely linked to

neurodegenerative diseases (89,90). We observed that the correlation between *ZRANB2-DT*

and *ZRANB2* was AD-specific, suggesting a gain regulatory role for *ZRANB2-DT* in the

context of AD. Additionally, differential gene expression analysis revealed that five of the

candidate lncRNAs—*ITFG1-AS1*, *Lnc-MYCN-6*, *Lnc-EFR3A-6*, *HSALNG0089044*, and *Lnc-*

CCDC68-1—were consistently dysregulated in AD across TCX datasets (Figure 3f, Table 3).

Despite comprehensive data collection from databases such as MalaCards (91),

LncRNADisease v3.0 (92), and existing literature (supplementary Table S13), none of the

previously reported AD-associated lncRNAs overlapped with our candidate lncRNAs.

Nevertheless, the presence of supportive evidence linking the candidate protein-coding

genes to AD reinforced our confidence that some of the identified key lncRNA genes were

also likely to play a role in AD, introducing promising new lncRNA candidates for further

investigation and experimental validation.

Aligning with the current knowledge that AD-related pathology begins in the medial temporal

lobe (2) and progresses to the frontal lobes in later stages (4), we had hypothesized that AD-

related network disruptions would differ across brain regions. In the TCX AD network, we

observed significant rewiring, while in the FCX AD network, there was a substantial loss of

healthy gene correlations. This suggests that AD-related damage initially disrupts cellular

interactions in the temporal lobe, as evidenced by the loss of control-specific links in the TCX

control network. As the disease progresses, cells in the temporal lobe establish new gene

expression patterns to compensate for the damage (AD-specific links). Conversely, the FCX

AD network shows a predominant loss of gene correlations, suggesting that the frontal lobe

may experience more extensive disruption before compensatory rewiring occurs. In line with

this, studies have shown that genes like *ACSL3* (63) and those involved in mitochondrial

respiratory complexes (81,93) display opposing patterns of dysregulation during the early

and late stages of AD, potentially reflecting mechanisms of gene expression compensation

as the disease progresses. This notion is further supported by findings from a single-cell

study across 19 cortical regions, which identified the temporal lobe as exhibiting the earliest

and most pronounced gene expression abnormalities (94).

While our study provided valuable insights into AD pathology through gene co-expression

network analysis, it is important to acknowledge certain limitations inherent to the

methodology. For instance, the thresholds applied for filtering links and annotating lncRNAs

to biological processes may have introduced some biases. The threshold for filtering links

may have removed interactions from one network if they fell below the chosen cutoff, leading

to their classification as lost links in that network. However, we reasoned that if a link was

weaker in one condition and stronger in another, it could still be considered an affected or

altered correlation. Similarly, the strict threshold for annotating lncRNAs may have excluded

some lncRNAs, as genes belonging to the same GO terms produced a wide range of RWR

scores. While this strict threshold helped maintain confidence in the assignments, we

recognized that non-annotated lncRNAs might still have been closely related to their

neighboring protein-coding genes. Additionally, other genes within the network might have

been crucial to AD pathology, but limitations in our methods may have prevented their

identification. To address this latter limitation, we made the TCX AD network publicly

available as an interactive Shiny app, enabling further exploration (see Data Availability).

The app allows users to investigate the entire TCX AD network, including all clusters, and

search for specific genes of interest. It also provides detailed information, such as gene

neighbors, functional categories, differential expression status, core membership, and more,

offering a resource for further discoveries and validations.

Conclusion

In summary, our network-based approach allowed us to provide insights into the alterations

of gene co-expression networks in AD brains, revealing clusters of functionally related genes

driven by key protein-coding and lncRNA genes. While early stages of AD, particularly in the

FCX, appear to be characterized primarily by a loss of gene interactions, disease

progression seems to trigger network rewiring, potentially aimed at preserving fundamental

brain functions. Additionally, we predicted functional roles for numerous lncRNAs within

these networks, identifying 27 lncRNAs with potential involvement in AD pathology. These

findings highlight promising candidates for future studies, offering deeper insights into their

roles in both normal brain function and AD development.

Material and Methods

RNA-seq datasets

We downloaded the FASTQ files and metadata from the Synapse platform (95). A total of

five RNA-seq datasets from two independent studies were used. The first study, known as

the Mayo RNA-seq data (42) includes RNA-seq data for the temporal cortex human brain

region. The FASTQ files for this study were downloaded from the Synapse repository with

the Synapse ID syn3163039 (96). The second dataset, known as the Mount Sinai Brain

Bank (MSBB) study (43), includes gene expression profiles for four brain regions: Brodmann

area (BM) 36, BM44, BM22, and BM10. This dataset was downloaded from the Synapse

repository with the Synapse ID syn3159438 (97). All samples were collected under precise

clinical-pathological diagnoses and distinctive histologies, including Braak NFT stage (44),

plaque density, CERAD (Consortium to Establish a Registry for Alzheimer's Disease) (45),

and CDR (clinical dementia rating) (46). Each AD and the corresponding control dataset

from both brain areas consisted of an equal number of individuals across sexes with

identical gene sets. Furthermore, these datasets have been carefully balanced for age,

disease stage, and other potential confounders. All data underwent standardized

preprocessing steps, including filtering based on library size and gene expression levels, to

ensure uniformity across the datasets.

Preprocessing of RNA-seq data

To ensure uniform data processing, we employed a unified pipeline that included standard

alignment, quality control, and gene expression analyses on each dataset (Mayo and MSBB

RNA-seq). The preprocessing steps included FASTQ quality control, adapter trimming, base

quality control, and filtering out short reads. We excluded all FASTQ files with fewer than 12

million reads. Following this, reads were mapped to the human reference genome (hg38)

using STAR (v 2.7.10b) (98). Samples with less than 60% unique mapped reads were also

excluded from further analysis. We used the `featureCounts()` function from the Subread

software (V 1.6.3) (99) for gene quantification, with the following parameters: `-s 0 -t`

`gene -g gene_id -M -O --fraction`. `-s 0` (un-stranded data), `-t gene` (gene

feature type), `-g gene_id` (grouped by gene ID), `-M` (allow multi-mapping reads), `-O`

(count overlapping reads), and `--fraction` (fractional assignment of multi-mapping

reads). To account for both sense and antisense lncRNAs, reads were mapped specifically

to the gene feature type (`-t gene`) and counted fractionally by

`countMultiMappingReads` arguments (`--fraction`). Noteworthy, the human gene

annotation file version 103 (*Homo_sapiens.GRCh38.103.gtf*) from Ensembl was used for

gene annotation. Notably, both datasets contained un-stranded sequence data, which

required the use of the `-s 0` option.

To remove systematic effects in the RNA-seq data, we normalized all gene count data using

the `DESeq2` R package (100), which normalizes for both sequencing depth and RNA

composition. Moreover, genes with low expression levels can introduce noise into the

correlation calculation process. Hence, we excluded genes with low expression levels to

enhance the reliability of the correlation analysis. The criteria for excluding low-expressed

genes were set such that only genes with a count of at least 1 count per million (cpm) in 70%

of the samples were retained. The use of cpm avoids favoring genes expressed in larger

libraries.

Signed weighted co-expression network

Gene co-expression networks represent gene-gene interactions, with nodes representing

genes and edges indicating the strength of their co-expression (101). The weight of an edge

reflects the strength of the co-expression relationship between two genes, while the sign

distinguishes between positive and negative correlations, showing whether genes have

similar or opposing expression patterns. The signed weighted gene co-expression networks

were constructed using the weighted Topological Overlap (wTO) method separately for each

RNA-seq dataset. Using the wTO method, link weights for connected nodes are calculated

through an averaging process that accounts for all their common connections (47). The wTO

calculation was performed using the `wTO` R package. In summary, for each dataset,

Pearson's correlation coefficient (ρ) was calculated for all pairs of genes to generate the

adjacency matrix $A = [a_{i,j}]$, where each element $a_{i,j}$ represents the correlation between gene i

and gene j. The weighted Topological Overlap ($\omega_{i,j}$) was then calculated for each pair of

nodes i and j, reflecting the shared network connectivity between the two genes. Gene co-

expression correlations with $|\omega i, j| < 0.5$ were discarded. This threshold was chosen to

balance the inclusion of meaningful interactions while reducing the influence of weak or

spurious associations. This threshold, higher than the third quartile value (mean of 0.4) and

close to the maximum observed weight (mean of 0.61), ensures that only the strongest links

are preserved for further analysis (Figure 2e, supplementary Table S1).

Consensus network construction by integrating wTO networks

To obtain a single consensus network derived from multiple independent wTO networks, the

`intersection()`function from the `igraph` package was used. A custom function was

developed to subset the overlapping links across the given networks and calculate an

average weight for each link. Links with differing signs were excluded. The consensus

weight between nodes i and j ($wTO_{CN} = [\Omega_{i,j}]$) for common links across n replicate networks

(`k=1,…,n`) was calculated using the formula from the `wTO.Consensus()` function of the

`wTO` R package, as follows:

$$\Omega_{i,j} = \sum_{k=1}^{n} \alpha_{ij}^{k} \, \omega_{ij}^{k}$$

592 Where

$$\alpha_{ij}^{k} = \left| \omega_{ij}^{k} \right| / \sum_{k=1}^{n} \left| \omega_{ij}^{k} \right|$$

This approach ensures that stronger link weights ω_{ij}^{k} contribute more significantly to the

corresponding consensus weights through a greater weighting factor α_{ij}^{k}. This helps

preserve the impact of strong links, preventing their dilution by weaker ones. In contrast, a

simple average would treat all connections equally, potentially blurring the distinction

between strong and weak links and resulting in the loss of valuable information.

Differential co-expression networks analysis

To identify differences between AD and control consensus networks, we performed a

network comparison analysis using the `CoDiNA` R package (49). CoDiNA (Co-Expression

Differential Network Analysis) is a network comparison framework that systematically

identifies shared and condition-specific gene-gene interactions between multiple networks.

Instead of simply highlighting differentially connected genes, CoDiNA classifies links into

three categories: (1) conserved links that are common between two conditions with the same

sign ($α$-link); (2) condition-specific links, appearing only in one condition ($β$-link); and (3)

reversed links, representing common correlations with opposite signs in the two conditions

($γ$-links). This qualitative framework enables a clear interpretation of how gene interactions

differ between conditions. For example, conserved links indicate stability in gene regulation,

while condition-specific and reversed links highlight shifts in regulatory relationships that may

underpin disease pathology. The package allows comparison of multiple networks, requiring

input in a table format where the first two columns specify node pairs, and the third column

contains their correlation values. We disabled weight normalization (stretch = FALSE) and

applied the default cutoff (0.33) to define the minimum weight threshold for a link to be

considered present. CoDiNA considers genes (nodes) present in both networks to avoid

false associations, as the absence of a gene in one network might be due to technical

reasons. However, given the nature of the gene expression data used in this study to

construct the wTO consensus networks, the gene lists were identical for both conditions.

Thus, the absence of a gene in one network did not necessarily indicate a technical issue

but rather that it might not meet the criteria for meaningful correlations, such as the weight

cutoff and network integration. To address this, we added the absent genes to each network

with a zero-weight assigned to their links, ensuring that genes present in one condition but

absent in the other yet still exhibiting a correlation were not excluded from the comparison.

This approach enabled a comprehensive analysis of network differences between AD and

control conditions.

Hub core genes

To determine hub genes within each network, we used the K-core decomposition method

(102), which identifies the most interconnected nodes by progressively removing nodes with

degrees less than k until no more such nodes exist. K-cores are determined as subgraphs

where every vertex has a degree of at least k. Therefore, vertices with the highest coreness

scores represent the most tightly integrated and connected parts of the network. The k-core

subgraph is defined through the following process: first, all nodes and their links are

removed if their degree is less than k (starting with   ). The algorithm works iteratively,

increasing k each time, until no more nodes meet the criteria (Figure 7). This effectively

peels off layers (shells) of nodes until the graph has none left. For each network, all nodes

were classified into shells based on the coreness scores, and nodes with the highest score

were assigned as core hub genes of that network. We used the `coreness()` function from

the `igraph` R package to calculate the coreness scores.

Figure 7: Illustration of K-core decomposition method (103)**.** All nodes of the network are

classified into shells based on their coreness score. The inner shell (K-shell 3) with red-colored nodes

represents those with the highest connectivity within the network, considered as core hub nodes in

this study.

Identifying sub-network clusters

Clusters are groups of genes with similar expression profiles and were explored based on

the topological structure of the gene co-expression network. The Louvain method (104) was

used for clustering. While this function can consider edge weights, it only accepts positive

numeric vectors for weights. Since the networks contain both positive and negative weights,

the weights argument was set to NULL, ensuring that all links are considered without their

signs. Additionally, the resolution argument was set to 1, as it corresponds to the original

definition of modularity defined by the algorithm. Essentially, a lower resolution value results

in fewer and larger clusters. Cluster smaller than 20 links were excluded from further

analysis.

GO enrichment analysis

Gene Ontology (GO) enrichment analysis for biological process terms was performed using the `compareCluster()`function from the `clusterProfiler` R package (105). The analysis was conducted separately for each gene cluster, using the same background genes. Background genes comprised the protein-coding genes that were included in the wTO construction. P-values were adjusted using the false discovery rate (FDR) method, with a significance threshold of 0.05.

Redundancy reduction of the enriched GO terms

Although GO enrichment is a powerful tool for understanding the biological processes and pathways associated with a set of genes, this analysis is frequently challenged by the redundancy inherent in GO terms. This redundancy occurs because of the hierarchical nature of the GO hierarchy, where parent terms frequently encompass child terms. For example, a gene may be annotated with both a parent term and a more specific term at different levels of the hierarchy within the GO. Various methods have been developed to address this issue, each with its own advantages and disadvantages. However, none of these methods work flawlessly, and some incorrectly group GO terms with inappropriate representative parent terms. Therefore, a manual approach was chosen to mitigate redundancy. Child terms were manually grouped into broader cellular biological process terms to which they belong. Additionally, GO terms that were irrelevant, included fewer than five genes, or represented different terms for the same biological process were excluded.

Random Walk with Restart

The random walk on a graph is an iterative process where a walker transitions from its current node to a randomly selected neighbor, starting from specified source node(s), known as seeds. The Random Walk with Restart (RWR) algorithm extends this concept by allowing the walker to restart at the seed node(s) with a probability r at each time step (106). In this study, RWR was used to assign lncRNAs within each cluster to the enriched functional categories (GO terms) identified for that cluster. Each co-expression network constructed in this study included both protein-coding genes and lncRNAs as nodes. Knowing the

involvement of protein-coding genes in the enriched biological processes, we aimed to

assign lncRNAs to these processes by leveraging their connectivity with neighboring protein-

coding genes within the network. For each cluster, we performed iterative RWR calculations,

using genes from each functional category as seed genes and ranking lncRNAs based on

their associations. The process for each iteration included the following steps: (1) performing

RWR calculations within each network cluster, (2) using genes from each functional category

within the cluster as seed genes and computing RWR scores for lncRNAs in that cluster, (3)

excluding link weights when computing RWR scores due to the algorithm's inability to handle

negative values in signed networks, (4) computing the adjacency matrix for the network, (5)

normalizing the adjacency matrix by columns, and (6) setting the restart probability (r) to

0.7.

Assigning lncRNAs to the functional category based on RWR scores

LncRNAs were ranked based on their computed scores for each functional category, and

those scoring above the 10th percentile were assigned to that functional category. As cross-

validation, we calculated RWR scores for genes within each functional category using a

leave-one-out method. Within each cluster, for each protein-coding gene in a given GO

group, we treated all other genes except one as seeds and computed the RWR score for the

excluded gene. This process was repeated across all genes annotated to each functional

category. The assigned lncRNA was accepted only if its RWR score did not fall below the

lowest RWR scores observed in the leave-one-out calculation for that category (Figure S1).

Statistical evaluation tests

Szymkiewicz-Simpson coefficient and permutation test was specifically chosen to rigorously

assess whether the observed overlaps in lncRNA annotations were significant or merely due

to random chance. The similarity between assigned lncRNAs to a common functional

category across the two networks was initially measured using the Szymkiewicz-Simpson

coefficient. For each functional category, the overlap coefficient was calculated by

determining the intersection of lncRNAs assigned to the respective functional category in

each network, normalized by the minimum number of lncRNAs assigned to that functional

category in either network. This approach normalized the overlap relative to the size of the

smaller set, accommodating the significant disparity in the total number of lncRNAs between

the FCX and TCX control networks (1100 and 504, respectively). The statistical significance

of the observed overlap was assessed using a permutation test. The permutation test

involved generating 10,000 random permutations of lncRNA sub-samples from the complete

list of lncRNAs for each network. During each permutation, lncRNAs were randomly sampled

from the entire pool of lncRNAs in each network to create a new, synthetic group of lncRNAs

matching the original count of lncRNAs assigned to that functional category. The significance

of the observed overlap was assessed by calculating the proportion of permuted overlap

coefficients that were equal to or greater than the observed coefficient, which was then

divided by the number of permutations to yield a p-value.

To assess the difference in lncRNA representation between TCX AD and control core

networks, we constructed a contingency table comparing the number of core lncRNAs in

each network and evaluated statistical significance using Fisher's exact test.

Differential gene expression analysis

The `DESeq2` R package was used to identify differentially expressed genes (DEGs) in AD

compared to control. Genes with a false discovery rate (FDR) $<$ 0.05 and absolute log fold

change > 0.3 were considered significant DEGs. Due to the integration of several networks

to make each consensus network, identifying DEGs presented a challenge. To address this,

only DEGs that were consistently identified across all gene expression datasets used to

construct the given network were selected. For example, the AD TCX consensus network

integrates three networks: AD Mayo, AD MSBB BM36, and AD MSBB BM22. DEGs were

identified separately for each dataset—AD Mayo, AD MSBB BM36, and AD MSBB BM22.

Ultimately, the DEGs in the AD TCX consensus network were determined by selecting those

DEGs that overlapped across all these three datasets.

Network visualization

The network visualizations presented in the manuscript were created using Cytoscape v.310.2 (107) and Gephi v.0.10.1 (48).

Single cell RNA-seq analysis

The single-cell RNA sequencing data was downloaded from the Synapse repository with the Synapse ID syn22079621. This dataset includes samples from the prefrontal cortex, comprising 7 control and 11 AD cases (108). Data preprocessing and analysis were performed using Seurat v 4.3.0.1 (109). To ensure high-quality single-cell data, we applied the following filtering criteria: cells with fewer than 200 detected genes or more than 7500 detected genes were removed, and cells with more than 5% of reads mapping to mitochondrial genes were excluded to eliminate low-quality or stressed cells. Following quality control, the dataset was normalized using the `NormalizeData` function in Seurat. Uniform Manifold Approximation and Projection (UMAP) was used for dimensionality reduction and visualization of cellular clusters. Clusters were identified and cell type were manually annotated based on established marker genes including *AQP4*, *SLC1A2*, and *GPC5* for astrocytes; *MBP*, *MOBP*, and *PLP1* for oligodendrocytes; *CD74*, *LPAR6*, *DOCK8*, and CSF1R for microglia; FLT1, *CLDN5*, and *LAMA2* for endothelial cells; *VCAN*, *DSCAM*, *PCDH15*, and *MEGF11* for oligodendrocyte progenitor cells; *NRGN* for excitatory neurons; and *GAD1* for inhibitory neurons.

Bulk RNA-seq deconvolution

To estimate cell-type proportions in bulk RNA-seq data, deconvolution was performed using `CIBERSORT` R package (110). The single-cell dataset described earlier was used to construct a signature gene matrix, utilizing only control samples for this purpose. The expression data was normalized using `SCTransform` (SCT), and the signature matrix was generated from SCT-normalized data by averaging expression across cell types. The CIBERSORT algorithm was run with 100 permutations, and only samples with a p-value < 0.05 were retained for further analysis. The estimated cell-type proportions between AD and

control samples were compared using the Wilcoxon test, with false discovery rate-adjusted

p-values < 0.05 considered significant.

Data availability

The datasets used in this study are available in the Synapse online repositories under the

following accession IDs: Mayo RNA-seq dataset (Synapse ID: syn3163039) (96) and MSBB

RNA-seq dataset (Synapse ID: syn3159438) (97).

R script code availability

To enhance the exploration of our findings, we provided an interactive Shiny app that

visualizes the gene co-expression networks. This tool enables researchers to investigate

specific genes and their correlations and associated functions, offering deeper insights into

the TCX AD and control networks. They can be downloaded from

https://github.com/mpstar92/Shiny_App_ADnet.

The R script code used in this study is available on the GitHub repository at

https://github.com/Tima-Ze/AD_network.

Acknowledgments

Data used in this project were obtained from the Synapse platform

(https://www.synapse.org/), including the Mayo RNA-seq study (96), the Mount Sinai Brain

Bank study (MSBB) dataset (97) and Human AD Single-nuclei Multi-Omics (108). We extend

our gratitude to the data providers on the Synapse platform, as well as the researchers and

authors involved in these studies. This study was supported by the International Max Planck

Research School for Biology and Computation (IMPRS-BAC), Berlin, Germany. We would

like to thank Vladimir M. Jovanovic, Melanie Sarfert, Yao-Chung Chen, and Rebecca Sinead

Saager for their kind assistance and contributions, as well as Ralf Herwig for his scientific

methodology advice.

Author contributions

Nowick supervised the research. Zebardast analyzed and visualized the data and wrote the manuscript. Riethmüller developed the Shiny app. All authors read and approved the final manuscript.

Conflict of Interest

The authors declare that there were no commercial or financial relationships involved in this research that could be interpreted as a potential conflict of interest.

Funding

This work is supported by funding from the Deutsche Forschungsgemeinschaft awarded to KN (NO 920/8-1).

References

1. Livingston G, Sommerlad A, Orgeta V, Costafreda SG, Huntley J, Ames D, et al. Dementia prevention, intervention, and care. The Lancet. 2017 Dec 16;390(10113):2673–734.

2. Planche V, Manjon JV, Mansencal B, Lanuza E, Tourdias T, Catheline G, et al. Structural progression of Alzheimer's disease over decades: the MRI staging scheme. Brain Commun. 2022 Jun 1;4(3):fcac109.

3. Ahmad F, Javed M, Athar M, Shahzadi S. Determination of affected brain regions at various stages of Alzheimer's disease. Neurosci Res. 2023 Jul 1;192:77–82.

4. Casanova R, Whitlow CT, Wagner B, Williamson J, Shumaker SA, Maldjian JA, et al. High dimensional classification of structural MRI Alzheimer's disease data based on large scale regularization. Front Neuroinformatics. 2011;5:22.

5. Reddy PH. Abnormal tau, mitochondrial dysfunction, impaired axonal transport of mitochondria, and synaptic deprivation in Alzheimer's disease. Brain Res. 2011 Sep 30;1415:136–48.

6. Devi L, Prabhu BM, Galati DF, Avadhani NG, Anandatheerthavarada HK. Accumulation of amyloid precursor protein in the mitochondrial import channels of human Alzheimer's disease brain is associated with mitochondrial dysfunction. J Neurosci Off J Soc Neurosci. 2006 Aug 30;26(35):9057–68.

7.　Moreira PI, Carvalho C, Zhu X, Smith MA, Perry G. Mitochondrial dysfunction is a
trigger of Alzheimer's disease pathophysiology. Biochim Biophys Acta BBA - Mol Basis
Dis. 2010 Jan;1802(1):2–10.

8.　Plácido AI, Pereira CMF, Duarte AI, Candeias E, Correia SC, Santos RX, et al. The role
of endoplasmic reticulum in amyloid precursor protein processing and trafficking:
Implications for Alzheimer's disease. Biochim Biophys Acta BBA - Mol Basis Dis. 2014
Sep 1;1842(9):1444–53.

9.　Nishitsuji K, Tomiyama T, Ishibashi K, Ito K, Teraoka R, Lambert MP, et al. The E693Δ
Mutation in Amyloid Precursor Protein Increases Intracellular Accumulation of Amyloid
β Oligomers and Causes Endoplasmic Reticulum Stress-Induced Apoptosis in Cultured
Cells. Am J Pathol. 2009 Mar 1;174(3):957–69.

10.　Joshi G, Chi Y, Huang Z, Wang Y. Aβ-induced Golgi fragmentation in Alzheimer's
disease enhances Aβ production. Proc Natl Acad Sci. 2014 Apr;111(13):E1230–9.

11.　Joshi G, Bekier ME, Wang Y. Golgi fragmentation in Alzheimer's disease. Front
Neurosci. 2015;9:340.

12.　Kim SH, Cho YS, Kim Y, Park J, Yoo SM, Gwak J, et al. Endolysosomal impairment by
binding of amyloid beta or MAPT/Tau to V-ATPase and rescue via the HYAL-CD44 axis
in Alzheimer disease. Autophagy. 2023 Aug;19(8):2318–37.

13.　Simoes S, Neufeld JL, Triana-Baltzer G, Moughadam S, Chen EI, Kothiya M, et al. Tau
and other proteins found in Alzheimer's disease spinal fluid are linked to retromer-
mediated endosomal traffic in mice and humans. Sci Transl Med. 2020 Nov
25;12(571):eaba6334.

14.　Gabrielli M, Prada I, Joshi P, Falcicchia C, D'Arrigo G, Rutigliano G, et al. Microglial
large extracellular vesicles propagate early synaptic dysfunction in Alzheimer's disease.
Brain J Neurol. 2022 Aug 27;145(8):2849–68.

15.　Karisetty BC, Bhatnagar A, Armour EM, Beaver M, Zhang H, Elefant F. Amyloid-β
Peptide Impact on Synaptic Function and Neuroepigenetic Gene Control Reveal New
Therapeutic Strategies for Alzheimer's Disease. Front Mol Neurosci. 2020 Nov 13;13.

16.　Du H, Guo L, Yan S, Sosunov AA, McKhann GM, Yan SS. Early deficits in synaptic
mitochondria in an Alzheimer's disease mouse model. Proc Natl Acad Sci U S A. 2010
Oct 26;107(43):18670–5.

17.　Scheff SW, Price DA, Ansari MA, Roberts KN, Schmitt FA, Ikonomovic MD, et al.
Synaptic Change in the Posterior Cingulate Gyrus in the Progression of Alzheimer's
Disease. J Alzheimers Dis. 2015 Jan 1;43(3):1073–90.

18.　Berchtold NC, Coleman PD, Cribbs DH, Rogers J, Gillen DL, Cotman CW. Synaptic
genes are extensively downregulated across multiple brain regions in normal human
aging and Alzheimer's disease. Neurobiol Aging. 2013 Jun;34(6):1653–61.

19.　Mattick JS, Makunin IV. Non-coding RNA. Hum Mol Genet. 2006 Apr
15;15(suppl_1):R17–29.

20.　Wang E, Lemos Duarte M, Rothman LE, Cai D, Zhang B. Non-coding RNAs in
Alzheimer's disease: perspectives from omics studies. Hum Mol Genet. 2022 Oct
15;31(R1):R54–61.

21. Grillone K, Riillo C, Scionti F, Rocca R, Tradigo G, Guzzi PH, et al. Non-coding RNAs
865   in cancer: platforms and strategies for investigating the genomic "dark matter". J Exp
866   Clin Cancer Res. 2020 Jun 20;39(1):117.

867   22. Zhang Y, Zhao Y, Ao X, Yu W, Zhang L, Wang Y, et al. The Role of Non-coding RNAs
868   in Alzheimer's Disease: From Regulated Mechanism to Therapeutic Targets and
869   Diagnostic Biomarkers. Front Aging Neurosci. 2021 Jul 2;13:654978.

870   23. Statello L, Guo CJ, Chen LL, Huarte M. Gene regulation by long non-coding RNAs and
871   its biological functions. Nat Rev Mol Cell Biol. 2021 Feb;22(2):96–118.

872   24. Fernandes JCR, Acuña SM, Aoki JI, Floeter-Winter LM, Muxel SM. Long Non-Coding
873   RNAs in the Regulation of Gene Expression: Physiology and Disease. Non-Coding
874   RNA. 2019 Mar;5(1):17.

875   25. Oo JA, Brandes RP, Leisegang MS. Long non-coding RNAs: novel regulators of
876   cellular physiology and function. Pflüg Arch - Eur J Physiol. 2022 Feb 1;474(2):191–
877   204.

878   26. DiStefano JK. The Emerging Role of Long Noncoding RNAs in Human Disease. In:
879   DiStefano JK, editor. Disease Gene Identification: Methods and Protocols. New York,
880   NY: Springer; 2018. p. 91–110.

881   27. Wang X, Zhang M, Liu H. LncRNA17A regulates autophagy and apoptosis of SH-SY5Y
882   cell line as an in vitro model for Alzheimer's disease. Biosci Biotechnol Biochem. 2019
883   Apr 3;83(4):609–21.

884   28. Zhang W, Zhao H, Wu Q, Xu W, Xia M. Knockdown of BACE1◻AS by siRNA improves
885   memory and learning behaviors in Alzheimer's disease animal model. Exp Ther Med.
886   2018 Sep 1;16(3):2080–6.

887   29. Faghihi MA, Modarresi F, Khalil AM, Wood DE, Sahagan BG, Morgan TE, et al.
888   Expression of a noncoding RNA is elevated in Alzheimer's disease and drives rapid
889   feed-forward regulation of β-secretase expression. Nat Med. 2008 Jul;14(7):723–30.

890   30. Li F, Wang Y, Yang H, Xu Y, Zhou X, Zhang X, et al. The effect of BACE1-AS on β-
891   amyloid generation by regulating BACE1 mRNA expression. BMC Mol Biol. 2019 Oct
892   1;20(1):23.

893   31. Wang Q, Ge X, Zhang J, Chen L. Effect of lncRNA WT1-AS regulating WT1 on
894   oxidative stress injury and apoptosis of neurons in Alzheimer's disease via inhibition of
895   the miR-375/SIX4 axis. Aging. 2020 Nov 21;12(23):23974.

896   32. Mus E, Hof PR, Tiedge H. Dendritic BC200 RNA in aging and in Alzheimer's disease.
897   Proc Natl Acad Sci U S A. 2007 Jun 19;104(25):10679–84.

898   33. Quek XC, Thomson DW, Maag JL, Bartonicek N, Signal B, Clark MB, et al. lncRNAdb
899   v2.0: expanding the reference database for functional long noncoding RNAs. Nucleic
900   Acids Res. 2014 Oct 20;43(Database issue):D168.

901   34. Pavlopoulos GA, Secrier M, Moschopoulos CN, Soldatos TG, Kossida S, Aerts J, et al.
902   Using graph theory to analyze biological networks. BioData Min. 2011 Apr 28;4(1):10.

903   35.   Liesecke F, De Craene JO, Besseau S, Courdavault V, Clastre M, Vergès V, et al.
904   Improved gene co-expression network quality through expression dataset down-
905   sampling and network aggregation. Sci Rep. 2019 Oct 8;9(1):14431.

906   36.   Lee HK, Hsu AK, Sajdak J, Qin J, Pavlidis P. Coexpression Analysis of Human Genes
907   Across Many Microarray Data Sets. Genome Res. 2004 Jun 1;14(6):1085–94.

908   37.   Berto S, Perdomo-Sabogal A, Gerighausen D, Qin J, Nowick K. A Consensus Network
909   of Gene Regulatory Factors in the Human Frontal Lobe. Front Genet. 2016 Mar 8;7:31.

910   38.   Ghandikota S, Sharma M, Ediga HH, Madala SK, Jegga AG. Consensus Gene Co-
911   Expression Network Analysis Identifies Novel Genes Associated with Severity of
912   Fibrotic Lung Disease. Int J Mol Sci. 2022 May 13;23(10):5447.

913   39.   Chai K, Zhang X, Tang H, Gu H, Ye W, Wang G, et al. The Application of Consensus
914   Weighted Gene Co-expression Network Analysis to Comparative Transcriptome Meta-
915   Datasets of Multiple Sclerosis in Gray and White Matter. Front Neurol. 2022 Feb 24;13.

916   40.   Arshad Z, McDonald JF. A computational approach to generate highly conserved gene
917   co-expression networks with RNA-seq data. STAR Protoc. 2022 Jun 17;3(2):101432.

918   41.   Zebardast F. Overview and illustration of the current study.Created in BioRender.
919   [Internet]. 2025 [cited 2024 Dec 25]. Available from: https://BioRender.com/k67x627

920   42.   Allen M, Carrasquillo MM, Funk C, Heavner BD, Zou F, Younkin CS, et al. Human
921   whole genome genotype and transcriptome data for Alzheimer's and other
922   neurodegenerative diseases. Sci Data. 2016 Oct 11;3(1):160089.

923   43.   Wang M, Beckmann ND, Roussos P, Wang E, Zhou X, Wang Q, et al. The Mount Sinai
924   cohort of large-scale genomic, transcriptomic and proteomic data in Alzheimer's
925   disease. Sci Data. 2018 Sep 11;5(1):180185.

926   44.   Braak H, Braak E. Neuropathological stageing of Alzheimer-related changes. Acta
927   Neuropathol. 1991;82(4):239–259.

928   45.   Morris JC, Heyman A, Mohs RC, Hughes JP, van Belle G, Fillenbaum G, et al. The
929   Consortium to Establish a Registry for Alzheimer's Disease (CERAD). Part I. Clinical
930   and neuropsychological assessment of Alzheimer's disease. Neurology. 1989
931   Sep;39(9):1159–65.

932   46.   Morris JC. Clinical Dementia Rating: A Reliable and Valid Diagnostic and Staging
933   Measure for Dementia of the Alzheimer Type. Int Psychogeriatr. 1997 Dec;9(S1):173–
934   6.

935   47.   Gysi DM, Voigt A, Fragoso T de M, Almaas E, Nowick K. wTO: an R package for
936   computing weighted topological overlap and a consensus network with integrated
937   visualization tool. BMC Bioinformatics. 2018 Oct 24;19(1):392.

938   48.   Bastian M, Heymann S, Jacomy M. Gephi: An Open Source Software for Exploring and
939   Manipulating Networks. Proc Int AAAI Conf Web Soc Media. 2009 Mar 19;3(1):361–2.

940   49.   Gysi DM, Fragoso T de M, Zebardast F, Bertoli W, Busskamp V, Almaas E, et al.
941   Whole transcriptomic network analysis using Co-expression Differential Network
942   Analysis (CoDiNA). PLOS ONE. 2020 Oct 15;15(10):e0240523.

50. Mathys H, Davila-Velderrain J, Peng Z, Gao F, Mohammadi S, Young JZ, et al. Single-
cell transcriptomic analysis of Alzheimer's disease. Nature. 2019 Jun;570(7761):332–7.

51. Escorcia-Rodríguez JM, Gaytan-Nuñez E, Hernandez-Benitez EM, Zorro-Aranda A,
Tello-Palencia MA, Freyre-González JA. Improving gene regulatory network inference
and assessment: The importance of using network structure. Front Genet. 2023 Feb
27;14.

52. Ballouz S, Verleyen W, Gillis J. Guidance for RNA-seq co-expression network
construction and analysis: safety in numbers. Bioinforma Oxf Engl. 2015 Jul
1;31(13):2123–30.

53. Zhang B, Horvath S. A general framework for weighted gene co-expression network
analysis. Stat Appl Genet Mol Biol. 2005;4:Article17.

54. Voigt A, Almaas E. Assessment of weighted topological overlap (wTO) to improve
fidelity of gene co-expression networks. BMC Bioinformatics. 2019 Jan 28;20(1):58.

55. Morabito S, Reese F, Rahimzadeh N, Miyoshi E, Swarup V. hdWGCNA identifies co-
expression networks in high-dimensional transcriptomics data. Cell Rep Methods. 2023
Jun 12;3(6):100498.

56. He F, Maslov S. Pan- and core- network analysis of co-expression genes in a model
plant. Sci Rep. 2016 Dec 16;6(1):38956.

57. Kundra R, Ciryam P, Morimoto RI, Dobson CM, Vendruscolo M. Protein homeostasis of
a metastable subproteome associated with Alzheimer's disease. Proc Natl Acad Sci.
2017 Jul 11;114(28):E5703–11.

58. Zhang T, Liu N, Wei W, Zhang Z, Li H. Integrated Analysis of Weighted Gene
Coexpression Network Analysis Identifying Six Genes as Novel Biomarkers for
Alzheimer's Disease. Oxid Med Cell Longev. 2021 Jul 26;2021:9918498.

59. Lanke V, Moolamalla STR, Roy D, Vinod PK. Integrative Analysis of Hippocampus
Gene Expression Profiles Identifies Network Alterations in Aging and Alzheimer's
Disease. Front Aging Neurosci. 2018 May 23;10.

60. Wang M, Li A, Sekiya M, Beckmann ND, Quan X, Schrode N, et al. Transformative
Network Modeling of Multi-omics Data Reveals Detailed Circuits, Key Regulators, and
Potential Therapeutics for Alzheimer's Disease. Neuron. 2021 Jan 20;109(2):257-
272.e14.

61. Gu X, Lai D, Liu S, Chen K, Zhang P, Chen B, et al. Hub Genes, Diagnostic Model, and
Predicted Drugs Related to Iron Metabolism in Alzheimer's Disease. Front Aging
Neurosci. 2022 Jul 7;14:949083.

62. Duan K, Ma Y, Tan J, Miao Y, Zhang Q. Identification of genetic molecular markers and
immune infiltration characteristics of Alzheimer's disease through weighted gene co-
expression network analysis. Front Neurol. 2022 Aug 22;13:947781.

63. Majerníková N, den Dunnen WFA, Dolga AM. The Potential of Ferroptosis-Targeting
Therapies for Alzheimer's Disease: From Mechanism to Transcriptomic Analysis. Front
Aging Neurosci. 2021 Dec 20;13:745046.

64.　Zhang Q, Yang P, Pang X, Guo W, Sun Y, Wei Y, et al. Preliminary exploration of the
co-regulation of Alzheimer's disease pathogenic genes by microRNAs and transcription
factors. Front Aging Neurosci. 2022 Dec 6;14.

65.　Baglietto-Vargas D, Forner S, Cai L, Martini AC, Trujillo-Estrada L, Swarup V, et al.
Generation of a humanized Aβ expressing mouse demonstrating aspects of
Alzheimer's disease-like pathology. Nat Commun. 2021 Apr 23;12(1):2421.

66.　Wang Z, Yan X, Zhao C. Dynamical differential networks and modules inferring
disrupted genes associated with the progression of Alzheimer's disease. Exp Ther
Med. 2017 Oct 1;14(4):2969–75.

67.　Finney CA, Delerue F, Gold WA, Brown DA, Shvetcov A. Artificial intelligence-driven
meta-analysis of brain gene expression identifies novel gene candidates and a role for
mitochondria in Alzheimer's disease. Comput Struct Biotechnol J. 2023 Jan 1;21:388–
400.

68.　Neff RA, Wang M, Vatansever S, Guo L, Ming C, Wang Q, et al. Molecular subtyping of
Alzheimer's disease using RNA sequencing data reveals novel mechanisms and
targets. Sci Adv. 2021 Jan;

69.　Zhang F, Zhong S, Yang S, Wei Y, Wang J, Huang J, et al. Identification of Potential
Therapeutic Targets of Alzheimer's Disease By Weighted Gene Co-Expression
Network Analysis. Chin Med Sci J. 2020 Dec 1;35(4):330–41.

70.　Wu Z, Liu P, Huang B, Deng S, Song Z, Huang X, et al. A novel Alzheimer's disease
prognostic signature: identification and analysis of glutamine metabolism genes in
immunogenicity and immunotherapy efficacy. Sci Rep. 2023 Apr 27;13(1):6895.

71.　Régal L, Mårtensson E, Maystadt I, Voermans N, Lederer D, Burlina A, et al. PREPL
deficiency: delineation of the phenotype and development of a functional blood assay.
Genet Med. 2018 Jan 1;20(1):109–18.

72.　Männistö PT, García-Horsman JA. Mechanism of Action of Prolyl Oligopeptidase
(PREP) in Degenerative Brain Diseases: Has Peptidase Activity Only a Modulatory
Role on the Interactions of PREP with Proteins? Front Aging Neurosci. 2017 Feb 14;9.

73.　Duarte ML, Wang M, Gomes I, Liu C, Sharma A, Fakira AK, et al. Multiomics Analyses
Identify Proline Endopeptidase–Like Protein As a Key Regulator of Protein Trafficking,
a Pathway Underlying Alzheimer's Disease Pathogenesis. Mol Pharmacol. 2023
Jul;104(1):1.

74.　Heywood WE, Galimberti D, Bliss E, Sirka E, Paterson RW, Magdalinou NK, et al.
Identification of novel CSF biomarkers for neurodegeneration and their validation by a
high-throughput multiplexed targeted proteomic assay. Mol Neurodegener. 2015 Dec
1;10(1):64.

75.　Shi Q, Gibson GE. Up-regulation of the mitochondrial malate dehydrogenase by
oxidative stress is mediated by miR-743a. J Neurochem. 2011 Jun 24;118(3):440.

76.　Rovelet-Lecrux A, Charbonnier C, Wallon D, Nicolas G, Seaman MNJ, Pottier C, et al.
De novo deleterious genetic variations target a biological network centered on Aβ
peptide in early-onset Alzheimer disease. Mol Psychiatry. 2015 Sep;20(9):1046–56.

77. Small SA, Kent K, Pierce A, Leung C, Kang MS, Okada H, et al. Model-guided
microarray implicates the retromer complex in Alzheimer's disease. Ann Neurol. 2005
Dec;58(6):909–19.

78. Muhammad A, Flores I, Zhang H, Yu R, Staniszewski A, Planel E, et al. Retromer
deficiency observed in Alzheimer's disease causes hippocampal dysfunction,
neurodegeneration, and Abeta accumulation. Proc Natl Acad Sci U S A. 2008 May
20;105(20):7327–32.

79. Schlatterer SD, Tremblay MA, Acker CM, Davies P. Neuronal c-Abl Overexpression
Leads to Neuronal Loss and Neuroinflammation in the Mouse Forebrain. J Alzheimer's
Dis. 2011 Jan 1;25(1):119–33.

80. Lebouvier T, Scales TME, Williamson R, Noble W, Duyckaerts C, Hanger DP, et al.
The Microtubule-Associated Protein Tau is Also Phosphorylated on Tyrosine. J
Alzheimer's Dis. 2009 Jul 29;18(1):1–9.

81. Mastroeni D, Khdour OM, Delvaux E, Nolz J, Olsen G, Berchtold N, et al. Nuclear but
not mitochondrial-encoded oxidative phosphorylation genes are altered in aging, mild
cognitive impairment, and Alzheimer's disease. Alzheimers Dement. 2017 May
1;13(5):510–9.

82. La Barbera L, Vedele F, Nobili A, Krashia P, Spoleti E, Latagliata EC, et al. Nilotinib
restores memory function by preventing dopaminergic neuron degeneration in a mouse
model of Alzheimer's Disease. Prog Neurobiol. 2021 Jul;202:102031.

83. Zhou Z, Bai J, Zhong S, Zhang R, Kang K, Zhang X, et al. Downregulation of ATP6V1A
Involved in Alzheimer's Disease via Synaptic Vesicle Cycle, Phagosome, and Oxidative
Phosphorylation. Oxid Med Cell Longev. 2021 Apr 19;2021:5555634.

84. Geldenhuys WJ, Leeper TC, Carroll RT. mitoNEET as a novel drug target for
mitochondrial dysfunction. Drug Discov Today. 2014 Oct 1;19(10):1601–6.

85. Dumont M, Kipiani K, Yu F, Wille E, Katz M, Calingasan NY, et al. Coenzyme Q10
decreases amyloid pathology and improves behavior in a transgenic mouse model of
Alzheimer's disease. J Alzheimers Dis JAD. 2011;27(1):211–23.

86. Choi H, Park HH, Koh SH, Choi NY, Yu HJ, Park J, et al. Coenzyme Q10 protects
against amyloid beta-induced neuronal cell death by inhibiting oxidative stress and
activating the P13K pathway. NeuroToxicology. 2012 Jan 1;33(1):85–90.

87. Sheykhhasan M, Amini R, Soleimani Asl S, Saidijam M, Hashemi SM, Najafi R.
Neuroprotective effects of coenzyme Q10-loaded exosomes obtained from adipose-
derived stem cells in a rat model of Alzheimer's disease. Biomed Pharmacother. 2022
Aug 1;152:113224.

88. Llanos-González E, Sancho-Bielsa FJ, Frontiñán-Rubio J, Rabanal-Ruíz Y, García-
Carpintero S, Chicano E, et al. Spatial and Temporal Protein Modules Signatures
Associated with Alzheimer Disease in 3xTg-AD Mice Are Restored by Early Ubiquinol
Supplementation. Antioxidants. 2023 Mar;12(3):747.

89. Li X, Xue Y, Liu X, Zheng J, Shen S, Yang C, et al. ZRANB2/SNHG20/FOXK1 Axis
regulates Vasculogenic mimicry formation in glioma. J Exp Clin Cancer Res CR. 2019
Feb 11;38:68.

90. Cohen OS, Weickert TW, Hess JL, Paish LM, McCoy SY, Rothmond DA, et al. A
1067   splicing-regulatory polymorphism in DRD2 disrupts ZRANB2 binding, impairs cognitive
1068   functioning and increases risk for schizophrenia in six Han Chinese samples. Mol
1069   Psychiatry. 2016 Jul;21(7):975–82.

1070   91. Malacards: The Human Disease Database - PMC [Internet]. [cited 2024 Nov 15].
1071   Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC5764584/

1072   92. Lin X, Lu Y, Zhang C, Cui Q, Tang YD, Ji X, et al. LncRNADisease v3.0: an updated
1073   database of long non-coding RNA-associated diseases. Nucleic Acids Res. 2024 Jan
1074   5;52(D1):D1365–9.

1075   93. Adav SS, Park JE, Sze SK. Quantitative profiling brain proteomes revealed
1076   mitochondrial dysfunction in Alzheimer's disease. Mol Brain. 2019 Jan 28;12(1):8.

1077   94. Wang M, Roussos P, McKenzie A, Zhou X, Kajiwara Y, Brennand KJ, et al. Integrative
1078   network analysis of nineteen brain regions identifies molecular signatures and networks
1079   underlying selective regional vulnerability to Alzheimer's disease. Genome Med. 2016
1080   Nov 1;8(1):104.

1081   95. info@sagebase.org SB. Synapse | Sage Bionetworks [Internet]. [cited 2024 May 28].
1082   Available from: https://www.synapse.org

1083   96. (Author Name Not Available). temporal cortex [Internet]. Synapse; 2016 [cited 2025 Jan
1084   13]. Available from: https://repo-
1085   prod.prod.sagebase.org/repo/v1/doi/locate?id=syn3163039&type=ENTITY

1086   97. (Author Name Not Available). MSBB [Internet]. Synapse; 2017 [cited 2025 Jan 13].
1087   Available from: https://repo-
1088   prod.prod.sagebase.org/repo/v1/doi/locate?id=syn3159438&type=ENTITY

1089   98. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
1090   universal RNA-seq aligner. Bioinforma Oxf Engl. 2013 Jan 1;29(1):15–21.

1091   99. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
1092   assigning sequence reads to genomic features. Bioinforma Oxf Engl. 2014 Apr
1093   1;30(7):923–30.

1094   100. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
1095   RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550.

1096   101. Stuart JM, Segal E, Koller D, Kim SK. A gene-coexpression network for global
1097   discovery of conserved genetic modules. Science. 2003 Oct 10;302(5643):249–55.

1098   102. Seidman SB. Network structure and minimum degree. Soc Netw. 1983 Sep 1;5(3):269–
1099   87.

1100   103. Zebardast F. Illustration of K-core decomposition method. Created in BioRender.
1101   [Internet]. 2025 [cited 2024 Dec 25]. Available from: https://BioRender.com/w13e275

1102   104. Traag VA, Waltman L, van Eck NJ. From Louvain to Leiden: guaranteeing well-
1103   connected communities. Sci Rep. 2019 Mar 26;9(1):5233.

105. Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal
enrichment tool for interpreting omics data. Innov Camb Mass. 2021 Aug
28;2(3):100141.

106. Tong H, Faloutsos C, Pan JY. Random walk with restart: fast solutions and
applications. Knowl Inf Syst. 2008 Mar 1;14(3):327–46.

107. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a
software environment for integrated models of biomolecular interaction networks.
Genome Res. 2003 Nov;13(11):2498–504.

108. Morabito S, Miyoshi E, Michael N, Shahin S, Martini AC, Head E, et al. Single-nucleus
chromatin accessibility and transcriptomic characterization of Alzheimer's disease. Nat
Genet. 2021 Aug;53(8):1143–55.

109. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, et al.
Comprehensive Integration of Single-Cell Data. Cell. 2019 Jun 13;177(7):1888-
1902.e21.

110. Newman AM, Liu CL, Green MR, Gentles AJ, Feng W, Xu Y, et al. Robust enumeration
of cell subsets from tissue expression profiles. Nat Methods. 2015 May;12(5):453–7.

1120