Title: R-loop imbalance compromises virulence in *Salmonella enterica*

Authors: Diego A. Pedraza-Delgado[1], Julia Jiménez-Espadafor[1,2], Miriam Ortiz-Padilla[1,3], Joaquín Bernal-Bayard[1], Francisco Ramos-Morales[1#] and Roberto Balbontín[1#].

Affiliations:

[1]Departamento de Genética, Facultad de Biología, Universidad de Sevilla. Avda. Reina Mercedes, 6. 41012 Sevilla. Spain.

[2]Current address: Department of Biology, Science4Life, Utretch University. Padualaan 8, Utretch. 3584 CH. The Netherlands.

[3]Current address: Centro Andaluz de Biología del Desarrollo, Universidad Pablo de Olavide/Consejo Superior de Investigaciones Científicas/Junta de Andalucía, Ctra Utrera Km. 1, Seville, 41013. Spain.

[#]Correspondence to: rbalbontin@us.es and framos@us.es.

Abstract:

Bacterial virulence and antibiotic resistance are interconnected global threats that can synergize to multiply their adverse effects on health and economy. Two proposed strategies to address this dual challenge are non-biocidal inhibition of virulence traits (anti-virulence strategy) and manipulation of eco-evolutionary dynamics in bacterial populations to hinder dissemination of antibiotic resistance (anti-resistance strategy). Both strategies require identifying factors involved in bacterial gene expression, fitness and evolution. R-loops strongly influence these traits, as well as fitness of bacteria carrying antibiotic resistance mutations. This makes the enzyme responsible for R-loop degradation, the RNase HI, a promising target for anti-resistance approaches. Interestingly, the involvement of R-loops in gene expression could also make RNase HI a potential target for anti-virulence strategies. In this study, we explored this possibility by investigating the effects of RNase HI deficiency on the pathogenicity of *Salmonella enterica*. We found that the absence of RNase HI alters the expression of genes associated with virulence both at the population and single-cell levels, and both *ex vivo* and during infection of mammalian cells. Furthermore, we observed that RNase HI depletion causes defects in phenotypes associated with virulence, such as motility and biofilm formation. Lack of RNase HI also reduces the ability of *Salmonella* to survive within macrophages. However, lack of RNase HI does not significantly affect the invasion of mammalian epithelial cells or their immune response. Overall, our results demonstrate a pleiotropic influence of R-loops on bacterial virulence, suggesting that RNase HI could be a target for anti-virulence strategies. These findings provide a theoretical framework for the development of dual anti-resistance and anti-virulence interventions based on RNase HI targeting.

45   **INTRODUCTION**

The world is currently entering a so-called post-antibiotic era[1]. With a reduced palette of effective

antibiotics, bacterial pathogens may cause an enormous health and economic burden worldwide[2]. For

instance, mortality associated to antibiotic-resistant infections is estimated to exceed that of cancer by

50   2050[3], and was almost 5 millions in 2019[4] . Moreover, there is growing evidence base linking antibiotic

resistance and climate change[5], increasing the scopes of these global challenges. The capacity of

bacteria to cause disease is a critical aspect of the antibiotic resistance crisis. Indeed, antibiotic

resistance and bacterial virulence can be conceived as two sides of the same coin, as bacterial

pathogens represent a smaller threat when can be treated with effective antibiotics, and antibiotic

55   resistant bacteria pose a lesser challenge if are not able to develop pathogenesis. In addition, antibiotic

resistance and bacterial virulence can influence one another, multiplying their combined detrimental

effects on global health and economy[6]. Thus, there is an imperative need to develop novel strategies to

tackle bacterial pathogens and antibiotic resistance, as well as to address their reciprocal influence[1].

60   *Salmonella enterica* is a Gram-negative bacterial species that encompasses over 2500 serovars[7],

able to infect a broad range of hosts, causing from gastroenteritis to systemic infection, depending on

the serovar and the host[8]. Many *S. enterica* pathogenic functions are encoded by virulence genes, some

clustered in horizontally acquired genomic regions known as *Salmonella* pathogenicity islands (SPIs)[9–

12], which can encode secretion systems able to inject into the host cells bacterial proteins known as

65   effectors[13]. These effectors promote invasion of host cells by bacteria and interfere with host functions

in order to facilitate pathogenesis[14]. Two of the most relevant and well-studied SPIs are SPI-1 and SPI-

2, which encode two distinct Type III Secretion Systems (T3SS) involved in invasion of the intestinal

epithelium and proliferation within mammalian cells[15], respectively. *Salmonellae* are responsible for a

large mortality and morbidity in human populations, as well as for a vast economic loss due to

70   infections of farming animals[16,17]. Thus, novel therapeutical interventions against infections caused by

S. enterica, particularly by antibiotic-resistant strains, are urgently needed.

Two promising approaches recently proposed to tackle the double threat of bacterial virulence

and antibiotic resistance are anti-virulence therapy (non-biocidal suppression of pathogenic traits)[18] and

75   anti-resistance strategy (selection against resistant bacteria by exploitation of eco-evolutionary

dynamics in bacterial populations)[19]. A key aspect for the development of either strategy is identifying

specific targets involved in bacterial pathogenesis and/or evolution. Interestingly, bacterial cells contain

molecular structures, called R-loops, which influence both regulation of gene expression and bacterial

evolution[20–23]. Thus, mechanisms involved in R-loop biology could potentially serve as targets for both

80   anti-virulence and anti-resistance therapies.

R-loops are RNA-DNA hybrids formed during transcription, upon the invasion of the DNA

template strand by the nascent RNA[24], which participate in key physiological processes, such as

transcription initiation and termination, regulation of gene expression, DNA repair, etc[20–23]. However, if

85   elevated, R-loops can cause transcription stalling, blocks to replication fork progression, inhibition of

DNA repair, replication-transcription conflicts and double-strand DNA breaks[25]. Thus, R-loops need to

be maintained within an optimal range to remain harmless for the cell[26–28]. RNase HI, whose function is

conserved along the phylogenetic tree, is one of the key factors involved in R-loop homeostasis, as it

specifically degrades RNA in R-loops[29,30].

We recently showed that depleting RNase HI function (either genetically or chemically)

strongly reduces the fitness of *Escherichia coli* carrying mutations conferring resistance to rifampicin

and/or streptomycin[31]. Indeed, lack of RNase HI leads to extinction of resistant clones when competing

against susceptible bacteria even at high initial frequency of resistance, abolishing compensatory

95 evolution, both *ex vivo* and in the mammalian gut[31]. Interestingly, depletion of RNase HI can enhance an antimycobacterial treatment[32], confirming RNase HI as a promising target for anti-resistance approaches.

The involvement of R-loops in transcription[33,34], and therefore in gene expression, makes
100 possible that R-loop imbalance caused by lack of RNase HI affects expression of virulence genes, potentially altering the capacity of bacterial pathogens to timely develop the program of gene expression that leads to efficient pathogenesis. Indeed, other ribonucleases have been shown to play central roles in the control of bacterial virulence[35,36], including regulation of SPIs in *Salmonella enterica*[37,38].

Thus, RNase HI function, which is susceptible to chemical inhibition[31], could potentially constitute a convergent target for both anti-resistance and anti-virulence approaches that remained unexplored to date. Here we investigated the involvement of R-loops in *S. enterica* virulence, and found that expression of genes associated with pathogenesis are significantly altered in the absence of
110 RNase HI, both at the population and single-cell levels. We also demonstrate that phenotypes related to virulence, such as biofilm formation, motility and growth in low pH, are reduced in a strain lacking RNase HI. Finally, we showed that, albeit lack of RNase HI does not significantly change *Salmonella*'s ability to invade mammalian epithelial cells not their immune response to the bacterium, it negatively affects its capacity to survive within mammalian macrophages. Taken together, our results indicate that
115 RNase HI represents a promising target for anti-virulence strategies and may constitute a novel dual target for both anti-virulence and anti-resistance approaches.

RESULTS

120 *Expression of genes associated with virulence is altered in* Salmonella *lacking RNase HI*

To study the effects of R-loop imbalance on pathogenesis, we first constructed a *Salmonella* strain carrying a scarless in frame deletion in the gene encoding RNase HI: *rnhA*. The deletion was designed in a way that does not affect a promoter of the downstream gene *dnaQ*, which is located within the

125 coding sequence of *rnhA*[39], and rendered a shorter version of the RNase HI lacking amino acids 44-153, which encompass three out of the four highly conserved carboxylates that form the so-called DEDD motif comprising the active-site core required for its catalytic activity[30]. We then measured the fitness of the *ΔrnhA* mutant strain and wild-type bacteria by running both growth curves and competitions in Lysogeny Broth (LB)[40], and observed a small but significant fitness defect in the *ΔrnhA* mutant with

130 respect to wild-type bacteria (Figure S1).

In order to test if R-loop imbalance affects expression of genes important for virulence, we took advantage of a collection of luciferase reporter fusions available in the laboratory[41], and selected a set of 15 genes involved in processes relevant to virulence, such as SPIs, iron uptake, quorum sensing or

135 Type VI Secretion Systems (Table 1). We measured the expression levels of these genes in wild-type and *ΔrnhA* strains under three different conditions: i) standard laboratory conditions (LB with shaking), ii) conditions of induction of SPI-1 (static cultures in LB with 0.3M of NaCl)[42], and iii) in a medium that mimics the intracellular milieu of mammalian cells (Low Phosphate and Magnesium medium, LPM) and induces SPI-2[43]. As Figure 1 shows, expression of most genes is significantly altered in the

140 *ΔrnhA* mutant in at least one of the conditions tested (Figure 1A), causing in most cases a negative effect (Figure 1B-D). However, upregulation can be observed in certain genes and conditions; for instance, genes encoding the SPI-2 T3SS effector SseK3, the structural component of the system SsaV, and the enterobactin non-ribosomal peptide synthetase EntF in conditions of SPI-2 induction, and the

synthase of the quorum molecule autoinducer-2 LuxS in LB and conditions of SPI-1 induction (Figure

145   1A).

<u>*The absence of RNase HI affects expression of SPI-1 at the single-cell level*</u>

150   Expression of the SPI-1 is not homogeneous across the entire population of *S. enterica*. Instead, it

shows bistability; that is, genes in the SPI-1 are expressed only in a subset of the total population,

whereas another fraction of the population does not express them[44]. Since lack of RNase HI affects

expression of a SPI-1 gene (*iagB*), we next tested if RNAse HI depletion influences expression of SPI-

1 at the single-cell level. We constructed wild-type and *ΔrnhA* strains carrying: i) a copy of the gene

155   encoding the superfolder green fluorescent protein (*sfGFP* hereafter) under the control of the promoter

and ribosome binding site (RBS) of the SPI-1 gene *sicA,* inserted in the pseudogene locus *malYX*, and

ii) a copy of the gene encoding the *mCherry* red fluorescent protein under the control of a constitutive

promoter, inserted in the pseudogene STM14_2013. These strains allowed us to distinguish viable

bacteria in a regular transcriptional status (those expressing *mCherry*) from compromised or dead cells

160   (those not expressing it), and to simultaneously measure by flow cytometry percentages of viable

bacteria expressing or not the SPI-1 (bacteria either showing both red and green fluorescences or only

red fluorescence, respectively) in each genetic background (Figure 2A). Interestingly, we observed that

lack of RNase HI significantly affects the percentage of bacteria that expresses SPI-1 both in rich

(Figure 2B, top panel) and minimal medium (Figure 2B, bottom panel), demonstrating that R-loop

165   imbalance indeed influences bistability in the expression of SPI-1.

The results described above confirm that lack of RNase HI alters expression of genes encoding

proteins directly involved in virulence, such as T3SS effectors, but also influences genes coding for

proteins involved in mechanisms relevant for bacterial physiology that can influence pathogenesis,

170 such as synthesis of enterobactin, signal transduction, and synthesis and processing of the quorum sensing molecule autoinducer-2[45–47]. Particularly, some of these regulatory circuits can influence phenotypes previously connected with bacterial pathogenesis. For instance, quorum sensing has been shown to affect both biofilm formation, bacterial motility and virulence[48,49]. Conversely, SPI-1 has been reported to modulate *Salmonella* motility[50]. This intricate cross-regulation opens the possibility that

175 RNase HI depletion pleiotropically affects multiple phenotypes associated with virulence.

RNase HI depletion negatively influences biofilm formation and bacterial motility

Biofilms play a relevant role in pathogenicity of *S. enterica*[51]. We thus first used the well-established

180 colony morphology assay[52], to measure the ability to form biofilms of the *ΔrnhA* mutant, using wild-type bacteria and a mutant lacking the master regulator of biofilm formation CsgD (*ΔcsgD*) as controls. Figure 3A shows that the strain lacking RNase HI is able to form wrinkly structures similar to those of the wild type, and different from the smooth colonies formed by the *ΔcsgD* control. However, colonies of *ΔrnhA* are unable to expand radially as wild-type colonies do, presenting a somewhat intermediate

185 phenotype, not as smooth as the *ΔcsgD* control but not as structured and expanded as wild-type bacteria. In order to quantify these differences, we compared the same genotypes using the crystal violet assay[53,54]. This allowed us to ratify that the mutant lacking RNase HI is not able to form biofilms as efficiently as wild-type bacteria, but still forms better biofilms than the *ΔcsgD* control (Figure 3B). Using controls as references, the ability to form biofilms in the *ΔrnhA* mutant was estimated to be

190 reduced approximately 65% with respect to the wild type (Figure 3C).

As bacterial motility is also linked to *Salmonella*'s virulence[55], and connected with some of the genes whose expression is altered by RNase HI depletion, we next tested the effects of lack of RNase HI on it, using the classic bacterial motility assay[56]. We compared motility of bacteria lacking RNase HI with those of the wild type and a non-flagellated *ΔfliGHI* mutant[57], as controls, and found that the

195   mutant lacking RNase HI is significantly less motile than the wild type, albeit not completely non-
       motile, as it is the *ΔfliGHI* mutant (Figure 4A). According to controls, the *ΔrnhA* mutant shows a
       reduction of around 40% in motility compared to wild-type bacteria (Figure 4B).

       Expression of SPI-2 is altered in the ΔrnhA mutant during infection of mammalian epithelial cells

       In order to test if the influence of R-loops on expression of virulence genes also occurs *in vivo*, we then
       constructed wild-type and *ΔrnhA* strains carrying the P_{sicA}-*sfGFP* reporter fusion of the SPI-1 described
       above and also an additional reporter fusion, P_{ssaB}-*mCherry*, in which the *mCherry* is under the control
       of the promoter and RBS of the SPI-2 gene *ssaB,* inserted in the pseudogene locus *STM14_2013.* These
205   double reporter strains allow to simultaneously measure expression of both SPI-1 and SPI-2 by flow
       cytometry using wild-type and *ΔrnhA* strains carrying constitutively expressed *sfGFP* or *mCherry* as
       controls. We infected HeLa human epithelial cells with these wild-type and *ΔrnhA* strains carrying the
       double reporter for SPI-1 and SPI-2 or constitutively expressing *sfGFP* and *mCherry,* and analyzed the
       number of bacterial cells expressing each island along the infection. Strains constitutively expressing
210   each fluorescence allowed us: i) to monitor progression of the infection in each genetic background and
       ii) to set up a threshold to distinguish fluorescent from non-fluorescent bacterial cells, and therefore
       which ones can be considered to show SPI-1 and/or SPI-2 activation. Interestingly, initially both
       genotypes seem equally able to invade host cells, but differences in number of bacteria can be observed
       as the infection progresses (Figure 5A), suggesting that mid-term survival of the strain lacking RNase
215   HI may be compromised. Regarding expression of SPIs, we observed similar expression patterns in the
       wild type and the mutant lacking RNase HI at 30 minutes after infection. However, at 24 hours the
       ΔrnhA shows a significantly greater percentage of cells expressing SPI-2 than wild-type bacteria
       (Figure 5B).

220 *Invasion of mammalian epithelial cells by* Salmonella *is not affected by loss of RNase HI*

The fact that lack of RNase HI alters expression of SPI-1, SPI-2, and that of other genes associated with virulence, together with the observed effects on phenotypes relevant for pathogenesis, such as motiliy and biofilm formation, may cause the *ΔrnhA* mutant to be defective in virulence. To test this,
225 we first measured the ability to invade human HeLa human epithelial cells of wild-type bacteria, the *ΔrnhA* mutant, and a double mutant where SPI-1 and SPI-2 are deleted (*ΔSPI-1 ΔSPI-2* hereafter)[58]. Consistent with the results shown in Figure 5, we did not observe any significant difference in invasion by wild-type bacteria or by mutants lacking RNase HI (Figure 6), indicating that R-loop imbalance does not affect the ability of *Salmonella* to invade epithelial cells.

Lack of RNase HI negatively affects Salmonella's ability to proliferate within mammalian macrophages

In order to confirm or refute the seeming decline of mutants without RNase HI over the course of infection (Figure 5), we decided to specifically test potential effects of R-loop imbalance on
235 intracellular proliferation. We thus next performed competitions during infection of RAW264.7 murine macrophages[59] of either the wild type, the *ΔrnhA* mutant or the *ΔSPI-1 ΔSPI-2* mutant against a reference wild-type strain labeled with a constitutively expressed *lacZ*. As Figure 7 shows, the capacity of the mutant lacking RNase HI is severely impaired with respect to the wild type, showing proliferation similar to that of the *ΔSPI-1 ΔSPI-2* control.

Secretion of interleukin 8 by mammalian epithelial cells in response to Salmonella *is not significantly affected by RNase HI depletion*

Infection of host cells by bacterial pathogens can trigger immune responses[60]. In turn, bacterial pathogens have evolved a complex weaponry of virulence factors to evade these immune defenses, at least temporarily[14,61]. Secretion of the pro-inflammatory cytokine interleukin-8 (IL-8) by human epithelial cells has been shown to rapidly increase in response to *S. enterica*[62–65], and to be downmodulated by the bacterium[66]. In order to test the effects of R-loop imbalance on both cell host immune response and immunomodulation by *Salmonella*, we measured by ELISA levels of IL-8 in the supernatant of HeLa human epithelial cells infected by the *ΔrnhA* mutant, and compared these amounts with those from supernantants of cells infected by either wild-type bacteria or the *ΔSPI-1 ΔSPI-2* mutant, or from non-infected cells, as controls. We did not detect any significant difference in IL-8 secretion between wild-type and *ΔrnhA* bacteria, whereas the *ΔSPI-1 ΔSPI-2* mutant shows IL-8 levels slightly above those in non-infected cells (Figure 8).

DISCUSSION

The advent of the post-antibiotic era poses the greatest challenge to modern medicine in human history. Numerous medical interventions require efficient antibiotics (e.g. surgery, dental care, treatment of traumatic injuries, etc.), and multiple medical conditions enhance susceptibility to bacterial pathogens (e. g. cancer, AIDS, etc.), making functional antibiotics a necessity to protect the population affected by these conditions[67,68]. Thus, novel strategies to treat bacterial infections and/or prevent the development and dissemination of antibiotic resistances are urgently needed. Anti-resistance and anti-virulence strategies have been proposed as innovative approaches to tackle the increasing threat of antibiotic resistant pathogens[18,19]. We recently reported that the enzyme responsible for degradation of R-loops, the RNase HI, is involved in maintenance of fitness in bacteria carrying antibiotic resistance mutations[31], being thus a promising candidate as target for anti-resistance approaches. Moreover, the role of R-loops in transcription made plausible that depletion of RNase HI function could affect

expression of genes relevant for virulence and therefore cause reduced pathogenicity. We tested this
hypothesis by analyzing the effects of lacking RNase HI on phenotypes linked to *Salmonella*'s
virulence. We found that, despite not causing a great fitness defect (Figure S1), lack of RNase HI
significantly affects expression of multiple genes related to virulence, including genes involved in
signal transduction, metal uptake, quorum sensing and genes in SPI-1 and SPI-2 (Figure 1). These
effects on expression of virulence genes also occurs at single-cell level (Figure 2), and during infection
of mammalian cells (Figure 5). Moreover, several phenotypes relevant for pathogenicity, such as
biofilm formation (Figure 3), and motility (Figure 4), are also negatively affected by RNase HI
depletion. This suggests that altered expression of genes associated with virulence caused by the excess
of R-loops in the absence of RNase HI renders *Salmonella* unable to properly develop the
transcriptional program required for efficiently develop the complete pathogenic course. For instance,
reduced motility decreases bacterial ability to swim towards mammalian cells[69], and defective biofilms
affects attachment to mammalian tissues[70]. In addition, excess of R-loops has consequences for
bacterial physiology that are particularly disadvantageous during infection. For instance, preliminary
results indicate that the mutant lacking RNase HI shows reduced growth at low pH (Figure S2), which
is a condition characteristic of the intracellular millieu. Moreover, this strain shows a reduction in
proliferation within mammalian macrophages comparable to that of a double SPI-1/SPI-2 deletion
mutant (Figure 7). This defect is likely due to misregulation of virulence genes, leading to poor
coordination between secretion machinery assembly and effector translocation[71], as well as to pervasive
RNA polymerase backtracking during intracellular growth[72], which promotes formation of R-loops[33].
Interestingly, invasion of human epithelial cells is not affected in *Salmonella* undergoing R-loop
imbalance (Figure 6), nor it is the immune response of epithelial cells to this genotype (Figure 8). Thus,
the inability to fully develop pathogenesis while inducing a normal immune response in the host opens
the possibility of using RNase HI mutants as live vaccines[73].

In the onset of a post-antibiotics era, where one of the biggest achievements of modern medicine is under siege, novel approaches to tackle bacterial infections and to reduce antibiotic resistance are desperately needed. In summary, our results demonstrated that R-loop imbalance pleiotropically affects multiple traits important for bacterial pathogenesis, making it a suitable target for anti-virulence strategies. Importantly, as previous results revealed the potential of RNase HI as a target for anti-resistance interventions[31,32], RNase HI has thus the potential to become a dual target for both anti-virulence and anti-resistance strategies, representing a promising intervention against one of the most pressing challenges to global health.

MATERIALS AND METHODS

Bacterial strains, media, and growth conditions

All strains in this study (Table S1) are *Salmonella enterica* subspecies *enterica* serovar Typhimurium strain ATCC 14028s[74]. Mutant strains were constructed by Lambda-Red recombineering[75] and/or scarless DNA recombineering[76], followed by transduction to a clean genetic background using P22 HT 105/1 *int201*[77]. Constitutively fluorescent strains harbor a copy of either superfolder green fluorescent protein (*sfGFP*) or mCherry fluorescent protein (*mCherry*), under the control of the $P_{LTetO-1}$ constitutive promoter and a strong RBS[78], inserted in the pseudogene locus STM14_2013. These constructions were generated by Lambda-Red recombineering using preexisting constructions[79,31] as templates. Fluorescent reporter fusions were generated by one-step recombineering[80] and P22 HT 105/1 *int201* transduction. Luciferase fusions were derivatives resulting from P22 HT 105/1 *int201* transduction of fusions obtained in a previous study[41]. Cultures were grown in either Lysogeny Broth[40] (LB, Lennox formulation), carbon-free minimal medium (NCE)[81] broth supplemented with 0.4% glucose, LB 0.3M NaCl[42], or low phosphate, low magnesium-containing medium (LPM)[43], and incubated at 37ºC in either

glass inocule tubes with shaking at 200 r.p.m or without shaking, in a INFORs HT Multitron Standard

320   shaker, or in round-bottom 96-well plates with shaking (700 rpm) in a Grant-bio PHMP-4 benchtop

incubator. Except for motility and colony morphology assays, solid medium was LB containing 1.5%

agar. Media were supplemented when necessary with antibiotics at the following concentrations:

ampicillin (100 µg/mL), kanamycin (50 µg/mL), chloramphenicol (10 µg/mL), and tetracycline (20 µg/

mL).

Growth curves and luciferase expression assays

Strains were streaked individually onto LB agar plates supplemented with the appropriate antibiotic,

and incubated overnight at 37 °C. The next day, independent colonies from each strain were inoculated

330   separately in LB supplemented with the appropriate antibiotic (100 µl per well) in a 96-well plate and

incubated overnight at 37 °C with shaking (700 rpm). The next day, 1:20 dilutions of the overnight

cultures (10µl of culture in 190µl medium) were made, and 10µl of that diluted solution was used to

inoculate round-bottom 96-well plates containing 90µl of the corresponding medium. Then plates were

incubated at 37 °C with double orbital shaking set as "fast" in a Biotek SYNERGY HTX® multi-mode

335   benchtop microplate reader (Agilent, Santa Clara, CA, United States of America), and

OD_{600}/luminiscence were measured every 30 min during 24 h.

Competitive fitness assays

340   The relative fitness (selection coefficient per generation) analyzed genotype was measured by

competitive growth against an isogenic strain carrying a Mu*d*J element (which encodes the *lacZ* gene)

inserted in the gene *trg,* which was shown not to affect bacterial virulence[59], as reference. Competitor

strains were first streaked out of their respective frozen vials to LB supplemented with the appropriate

antibiotic, then individual colonies were inoculated separately in medium without antibiotics and

345 incubated overnight (approximately 16 h); the next morning, the number of cells in each culture was

estimated by OD_{600} and 1:1 mixtures of Lac+ and Lac- bacteria were inoculated in glass tubes and

incubated at 37ºC with shaking (200 r.p.m) for 24 h. Initial and final frequencies of the strains were

estimated by plating dilutions onto plates supplemented with X-gal, and counting Colony Forming

Units (CFUs). The number of generations was estimated from that of the reference sensitive strain, and

350 the selection coefficient was determined as described below for each independent competition. The

formula used to calculate the selection coefficient was $s = \left[\ln\left(NSf/NRf\right) - \ln\left(NSi/NRi\right)\right]/\ln\left(NRf/NRi\right)$,

being NRi and NRf the initial and final number of reference bacteria, and NSi and NSf the initial and

final number of the sample bacteria that is being analyzed.

355 *Flow cytometry*

Flow cytometry was performed using either a MACSQuant® Analyzer VYB flow cytometer (Miltenyi

Biotec, Bergisch Gladbach, Germany) or a CytoFlex S® flow cytometer (Beckman Coulter, Brea, CA,

United States of America), equipped with 488 nm and 561 nm lasers used for scatter parameters and

360 sfGFP and mCherry excitation. Regarding optical configuration for detection, sfGFP was detected

using bandpass filters of 525/50 nm (MACSQuant®) or 525/40 nm (CytoFlex S®), and mCherry was

detected using bandpass filters of 615/20 nm (MACSQuant®) or 610/20 nm (CytoFlex S®). To detect

bacteria, the analyzers were equipped with either a photomultiplier tube (PMT) in the case of the

MACSQuant® and an avalanche photodiode (APD) in the case of the CytoFlex S®. Samples were

365 acquired using either MACSQuantify™ 2.13.3 software (MACSQuant®) or CytExpert version 2.3

software (CytoFlex S®), and analyzed using FlowJo version 10.9.0 software (Tree Star, Inc., Ashland,

OR, United States of America). Prior to flow cytometry measurement, bacterial cultures were diluted in filtered phosphate-buffered saline (PBS).

370 *Colony morphology assays*

Strains were streaked individually onto LB agar plates supplemented with the appropriate antibiotic, and glass tubes containing LB were inoculated with six independent colonies of each strain and incubated overnight at 37 °C. The next day, 5 µL of each culture were used to inoculate a LB agar

375 without salt (Yeast Extract 5g/L, Tryptone 10g/L)[52] plate. Images of bacterial colonies after 7 days of incubation at 30°C were acquired using a Gel Doc XR+® Molecular Imager (Bio-Rad, Hercules, CA, United States of America) and Image Lab version 5.1 software.

Crystal violet assays

Strains were streaked individually onto LB agar plates supplemented with the appropriate antibiotic, and glass tubes containing LB without salt (Yeast Extract 5g/L, Tryptone 10g/L)[52] were inoculated with three independent colonies of each strain and incubated overnight at 37 °C. The next day, overnight cultures were diluted 1:100 in LB without salt, six technical replicates of 100 µL from each subculture

385 were placed in wells of a round-bottom 96-well plate to inoculate a LB without salt (Yeast Extract 5g/L, Tryptone 10g/L), and the plate was incubated at 25ºC for 48 h, without shaking. Then OD_{600} of the cultures were measured, the medium containing planctonic bacteria was removed, wells were washed three times with sterile PBS, 200 µL of methanol was added, and the plate was incubated 15 minutes at room temperature in static to fix the bacterial biofilm. After the methanol was removed, the

390 plate was left to dry for 10 minutes at room temperature in static, 125 µL of 0.1 crystal violet was added, and the plate was incubated 15 minutes at room temperature in static to stain bacterial biofilms.

Then wells were washed again three times with PBS, and the plate was left to dry for at least 2 h. Once dry, 250 µL of 33% acetic acid was added to each well, and the plate was incubated 15 minutes at room temperature in static, to dissolve the crystal violet attached to the biofilms, and then OD_{580} was measured.

Motility assays

Strains were streaked individually onto LB agar plates supplemented with the appropriate antibiotic, and glass tubes containing LB were inoculated with six independent colonies of each strain and incubated overnight at 37 °C. The next day, a sterile toothpick was soaked in each culture and used to inoculate a motility agar plate (5 g/L NaCl, 10 g/L Tryptone, 3 g/L agar)[56]. Bacterial growth halos were compared after 14 h of incubation at 37°C.

Infection of HeLa human epithelial cells

HeLa (human epithelial; ECACC no. 93021013) cells were seeded in 24-well plates at 1.5×10^5 cells per well and incubated in DMEM supplemented with 10% fetal calf serum for 24 h at 37 °C with 5% CO2. Bacterial strains were streaked individually onto LB agar plates supplemented with the appropriate antibiotic, and glass tubes containing LB were inoculated with independent colonies of each strain and incubated overnight at 37 °C. Then bacteria were added to the cell culture at a multiplicity of infection of 100. To remove noninternalized bacteria, cell cultures were washed twice with sterile PBS 30 minutes after infection, overlaid with DMEM containing 100 µg/mL gentamicin, and incubated for another hour and a half. At this point, supernatant samples for invasion assays and ELISA were taken. For measurement of gene expression *in vivo*, cultures were washed twice with sterile PBS, covered with DMEM with gentamicin (16 µg/mL), and incubated for additional 22 h. For

cell invasion assays, then infected cells were lysed with 1% Triton X-100–PBS (pH 7.4) (10 min, room temperature), and the number of viable intracellular bacteria was determined by plating of appropriate dilutions in LB agar, and colony counting.

Competitive proliferation assays within RAW murine macrophages

RAW264.7 (murine macrophages; ECACC no. 91062702) cells were seeded in 24-well plates at 1.5 × 10^5 cells per well and incubated in DMEM supplemented with 10% fetal calf serum for 24 h at 37 °C

425   with 5% CO2. Bacterial strains were streaked individually onto LB agar plates supplemented with the appropriate antibiotic, and glass tubes containing LB were inoculated with each strain, incubated overnight at 37 °C with shaking, and mixtures of different genotypes at 1:1 ratio were added to eucaryotic cells cultured to up to 60 to 70% confluency in 24-well plastic plates (approximately 3 x 10^5 eucaryotic cells/well) at a multiplicity of infection of 50. To remove noninternalized bacteria, cell

430   cultures were washed twice with sterile PBS 30 minutes after infection, overlaid with DMEM containing 100 µg/mL gentamicin, and incubated for another hour and a half. The culture was then washed twice with sterile PBS, covered with DMEM with gentamicin (16 µg/mL), and incubated for 22 h. Then infected cells were lysed with 1% Triton X-100–PBS (pH 7.4) (10 min, room temperature), and the number of viable intracellular bacteria was determined by plating of appropriate dilutions in LB

435   agar, and colony counting. Calculation of competitive index was done as described elsewhere[59].

Detection of IL-8 secretion by ELISA

Supernatants of HeLa human epithelial cells either non-infected or infected by the different bacterial

440   genotypes, were assayed for IL-8 secretion using the Human IL-8 Uncoated ELISA kit (Invitrogen), following manufacturer's instructions. Briefly, 96-well polysorbent plates previously incubated

overnight at room temperature with human IL-8 capture antibody were washed, and then blocked with Reagent Diluent overnight at 4°C, and washed again. Then, 100 µL of appropriately diluted samples of supernatant (and serial dilutions of an appropriate human IL-8 standard, for IL-8 quantification) were

445 transferred to the plates and incubated for 2 h at room temperature. Then, plates were washed again, followed by 2 h of incubation at room temperature with IL-8 detection antibody, a new wash, and 20 min of incubation, at room temperature and in the dark, with streptavidin-HRP B. Plates were then washed one final time, and incubated for 20 min at room temperature in the dark with substrate solution, upon which STOP solution was added, and signal development and detection were carried out

450 using a Biotek SYNERGY HTX® multi-mode benchtop microplate reader (Agilent, Santa Clara, CA, United States of America), at a wavelength of 450 nm, with a subtraction wavelength of 570 nm. Serially diluted IL-8 standards were used for the extrapolation of protein concentration from absorbance data.

455 *Viability spot assays at neutral and low pH*

For viability assays at normal and low pH, strains were first streaked out of their respective frozen vials, then individual colonies were inoculated separately in medium without antibiotics and incubated overnight (approximately 16 h); the next morning, 10^{-1}, 10^{-2}, 10^{-3}, 10^{-4}, 10^{-5}, 10^{-6} and 10^{-7} dilutions were

460 prepared by serial dilutions in sterile PBS. Then 5 µl aliquots of these bacterial solutions (approximately 5×10^6 to 0.5 cells) were spotted onto plain LB agar plates at pHs of either 6.8 or 4.9, and these plates were incubated overnight at 37 °C.

Statistical analyses

All analyses were conducted using Libreoffice Calc version 6.0.7.3 (www.libreoffice.org) software, R version 3.6.3 (www.r-project.org) software via RStudio version 1.2.1578 interface (www.rstudio.com), and GraphPad Prism version 8.0.2 (www.graphpad.com). In competitive fitness and luminiscence measurement assays, for each set of competitions, one-sample two-tailed Student's t-tests were used to

470 determine whether the data set corresponding to each genotype is significantly different than 0. For comparing different conditions or genotypes, two-tailed unpaired Student's t-tests were used.

ACKNOWLEDGEMENTS

475 We are grateful to Modesto Carballo and Laura Navarro of the "Servicio de Biología" of the "Centro de Investigación, Tecnología e Innovación de la Universidad de Sevilla (CITIUS)", for their assistance with experiments performed at CITIUS facilities.

AUTHOR CONTRIBUTIONS

Conceptualization: R.B., F.R.-M., J.B.-B.; validation: R.B., J.B.-B., F.R.-M.; formal analysis: R.B., J.B.-B., F.R.-M; investigation: R.B., D.A.P.-D., J.J.-E., M.O.-P; resources: F.R.-M., J.B.-B., R.B.; writing—original draft preparation: R.B.; writing—review and editing: R.B., D.A.P.-D, J.J.-E., M.O.-P, J.B.-B., F.R.-M.; supervision: F.R.-M., J.B.-B., R.B.; project administration: F.R.-M., J.B.-B., R.B.;

485 funding acquisition, F.R.-M., J.B.-B., R.B. All authors have read and agreed to the published version of the manuscript.

FUNDING

490 This research was funded by the grants PID2019-106132RB-I00 and PID2022-136863NB-I00, granted by MCIN/AEI/10.13039/501100011033 and by "ERDF a way of making Europe" (to F.R-M. and J.B-B.), María Zambrano fellowship MZAMBRANO-2021-20045 (to R.B.), by grant CNS2022-135955, from Plan de Recuperación, Transformación y Resiliencia «European Union-NextGenerationEU» (to R.B.), by Emergia fellowship EMC21_00154, from Consejería de Transformación Económica,

495 Industria, Conocimiento y Universidades from Junta de Andalucía (to R.B.), and by grant VII PP PRECOM C-2025/00000353, from the VII Plan propio de Investigación y Transferencia of Universidad de Sevilla (to R.B.). The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

500 **REFERENCES**

1. Okeke, I. N. *et al.* The scope of the antimicrobial resistance challenge. *Lancet* **403**, 2426–2438 (2024).

2. Ho, C. S. *et al.* Antimicrobial resistance: a concise update. *Lancet Microbe* **6**, 100947 (2025).

3. Tackling drug-resistant infections globally : final report and recommendations / the Review on Antimicrobial Resistance chaired by Jim O'Neill. *Wellcome Collection* https://wellcomecollection.org/works/thvwsuba/items.

4. Murray, C. J. *et al.* Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *The Lancet* **0**, (2022).

5. van Bavel, B. *et al.* Intersections between climate change and antimicrobial resistance: a systematic scoping review. *Lancet Planet Health* **8**, e1118–e1128 (2024).

6. Cepas, V. & Soto, S. M. Relationship between Virulence and Resistance among Gram-Negative Bacteria. *Antibiotics (Basel)* **9**, E719 (2020).

7. Tindall, B. J., Grimont, P. a. D., Garrity, G. M. & Euzéby, J. P. Nomenclature and taxonomy of the genus *Salmonella*. *Int J Syst Evol Microbiol* **55**, 521–524 (2005).

8. Garai, P., Gnanadhas, D. P. & Chakravortty, D. *Salmonella enterica* serovars Typhimurium and Typhi as model organisms: revealing paradigm of host-pathogen interactions. *Virulence* **3**, 377–388 (2012).

9. Galán, J. E. & Curtiss, R. Cloning and molecular characterization of genes whose products allow *Salmonella* Typhimurium to penetrate tissue culture cells. *Proc Natl Acad Sci U S A* **86**, 6383–6387 (1989).

10. Ochman, H., Soncini, F. C., Solomon, F. & Groisman, E. A. Identification of a pathogenicity island required for *Salmonella* survival in host cells. *Proc Natl Acad Sci U S A* **93**, 7800–7804 (1996).

11. Shea, J. E., Hensel, M., Gleeson, C. & Holden, D. W. Identification of a virulence locus encoding a second type III secretion system in *Salmonella* Typhimurium. *Proc Natl Acad Sci U S A* **93**, 2593–2597 (1996).

12. Gerlach, R. G. & Hensel, M. Salmonella pathogenicity islands in host specificity, host pathogen-interactions and antibiotics resistance of *Salmonella enterica*. *Berl Munch Tierarztl Wochenschr* **120**, 317–327 (2007).

13. Azimi, T., Zamirnasta, M., Sani, M. A., Soltan Dallal, M. M. & Nasser, A. Molecular Mechanisms of *Salmonella* Effector Proteins: A Comprehensive Review. *Infect Drug Resist* **13**, 11–26 (2020).

14. Bernal-Bayard, J. & Ramos-Morales, F. Molecular Mechanisms Used by *Salmonella* to Evade the Immune System. *Curr Issues Mol Biol* **25**, 133–168 (2018).

15. Gerlach, R. G. & Hensel, M. Protein secretion systems and adhesins: the molecular armory of Gram-negative pathogens. *Int J Med Microbiol* **297**, 401–415 (2007).

16. Kim, S., Kang, H., Excler, J.-L., Kim, J. H. & Lee, J.-S. The Economic Burden of Non-Typhoidal Salmonella and Invasive Non-Typhoidal *Salmonella* Infection: A Systematic Literature Review. *Vaccines (Basel)* **12**, 758 (2024).

17. Perez-Sepulveda, B. M. & Hinton, J. C. D. Microbe Profile: *Salmonella* Typhimurium: the master of the art of adaptation. *Microbiology (Reading)* **171**, (2025).

18. Ogawara, H. Possible drugs for the treatment of bacterial infections in the future: anti-virulence drugs. *J Antibiot (Tokyo)* **74**, 24–41 (2021).

19. Baquero, F., Coque, T. M. & de la Cruz, F. Ecology and evolution as targets: the need for novel eco-evo drugs and strategies to fight antibiotic resistance. *Antimicrob Agents Chemother* **55**, 3649–3660 (2011).

20. Cheng, B. *et al.* RNase H overproduction allows the expression of stress-induced genes in the absence of topoisomerase I. *FEMS Microbiol Lett* **221**, 237–242 (2003).

21. Naka, K., Koga, M., Yonesaki, T. & Otsuka, Y. RNase HI stimulates the activity of RnlA toxin in *Escherichia coli*. *Mol Microbiol* **91**, 596–605 (2014).

22. Maduike, N. Z., Tehranchi, A. K., Wang, J. D. & Kreuzer, K. N. Replication of the *Escherichia coli* chromosome in RNase HI-deficient cells: multiple initiation regions and fork dynamics. *Mol Microbiol* **91**, 39–56 (2014).

23. Drolet, M. & Brochu, J. R-loop-dependent replication and genomic instability in bacteria. *DNA Repair (Amst)* **84**, 102693 (2019).

24. Thomas, M., White, R. L. & Davis, R. W. Hybridization of RNA to double-stranded DNA: formation of R-loops. *Proc Natl Acad Sci U S A* **73**, 2294–2298 (1976).

25. Sollier, J. & Cimprich, K. A. Breaking bad: R-loops and genome integrity. *Trends Cell Biol* **25**, 514–522 (2015).

26. Crossley, M. P., Bocek, M. & Cimprich, K. A. R-Loops as Cellular Regulators and Genomic Threats. *Mol Cell* **73**, 398–411 (2019).

27. García-Muse, T. & Aguilera, A. R Loops: From Physiological to Pathological Roles. *Cell* **179**, 604–618 (2019).

28. Hegazy, Y. A., Fernando, C. M. & Tran, E. J. The balancing act of R-loop biology: The good, the bad, and the ugly. *J Biol Chem* **295**, 905–913 (2020).

29. Tadokoro, T. & Kanaya, S. Ribonuclease H: molecular diversities, substrate binding domains, and catalytic mechanism of the prokaryotic enzymes. *FEBS J* **276**, 1482–1493 (2009).

30. Hyjek, M., Figiel, M. & Nowotny, M. RNases H: Structure and mechanism. *DNA Repair (Amst)* **84**, 102672 (2019).

31. Balbontín, R., Frazão, N. & Gordo, I. DNA Breaks-Mediated Fitness Cost Reveals RNase HI as a New Target for Selectively Eliminating Antibiotic-Resistant Bacteria. *Mol Biol Evol* **38**, 3220–3234 (2021).

32. Al-Zubaidi, A. *et al.* RNase HI Depletion Strongly Potentiates Cell Killing by Rifampicin in Mycobacteria. *Antimicrob Agents Chemother* **66**, e0209121 (2022).

33. Gowrishankar, J., Leela, J. K. & Anupama, K. R-loops in bacterial transcription: their causes and consequences. *Transcription* **4**, 153–157 (2013).

34. Raghunathan, N. *et al.* Genome-wide relationship between R-loop formation and antisense transcription in *Escherichia coli*. *Nucleic Acids Res* **46**, 3400–3411 (2018).

35. Lawal, A., Jejelowo, O., Chopra, A. K. & Rosenzweig, J. A. Ribonucleases and bacterial virulence. *Microb Biotechnol* **4**, 558–571 (2011).

36. Matos, R. G. *et al.* The Role of Ribonucleases and sRNAs in the Virulence of Foodborne Pathogens. *Front Microbiol* **8**, 910 (2017).

37. López-Garrido, J. & Casadesús, J. Regulation of *Salmonella enterica* pathogenicity island 1 by DNA adenine methylation. *Genetics* **184**, 637–649 (2010).

38. López-Garrido, J., Puerta-Fernández, E. & Casadesús, J. A eukaryotic-like 3' untranslated region in *Salmonella enterica hilD* mRNA. *Nucleic Acids Res* **42**, 5894–5906 (2014).

39. Kröger, C. *et al.* An infection-relevant transcriptomic compendium for *Salmonella enterica* Serovar Typhimurium. *Cell Host Microbe* **14**, 683–695 (2013).

40. Bertani, G. Studies on lysogenesis. I. The mode of phage liberation by lysogenic *Escherichia coli*. *J Bacteriol* **62**, 293–300 (1951).

41. de la Rosa-Altura, J. J. *et al.* Generation of Random *luxCDABE* Transcriptional Fusions in the Genome of *Salmonella enterica*. *Methods Mol Biol* **2182**, 141–151 (2021).

42. Galán, J. E. & Curtiss, R. Expression of *Salmonella* Typhimurium genes required for invasion is regulated by changes in DNA supercoiling. *Infect Immun* **58**, 1879–1885 (1990).

43. Deiwick, J., Nikolaus, T., Erdogan, S. & Hensel, M. Environmental regulation of *Salmonella* pathogenicity island 2 gene expression. *Mol Microbiol* **31**, 1759–1773 (1999).

44. Hautefort, I., Proença, M. J. & Hinton, J. C. D. Single-copy green fluorescent protein gene fusions allow accurate measurement of *Salmonella* gene expression *in vitro* and during infection of mammalian cells. *Appl Environ Microbiol* **69**, 7480–7491 (2003).

45. Ratledge, C. Iron metabolism and infection. *Food Nutr Bull* **28**, S515-523 (2007).

46. de Pina, L. C. *et al.* The role of two-component regulatory systems in environmental sensing and virulence in *Salmonella*. *Crit Rev Microbiol* **47**, 397–434 (2021).

47. Zhang, X. *et al.* Regulatory Mechanisms between Quorum Sensing and Virulence in *Salmonella*. *Microorganisms* **10**, 2211 (2022).

48. Sholpan, A., Lamas, A., Cepeda, A. & Franco, C. M. *Salmonella* spp. quorum sensing: an overview from environmental persistence to host cell invasion. *AIMS Microbiol* **7**, 238–256 (2021).

49. Rana, K. *et al.* Association of quorum sensing and biofilm formation with *Salmonella* virulence: story beyond gathering and cross-talk. *Arch Microbiol* **203**, 5887–5897 (2021).

50. Saleh, D. O. *et al.* SPI-1 virulence gene expression modulates motility of *Salmonella* Typhimurium in a proton motive force- and adhesins-dependent manner. *PLoS Pathog* **19**, e1011451 (2023).

51. Espinoza-Erazo, V. P., Vela-Chauvin, M. G., Collantes-Vela, J. C., Zapata-Mena, S. & Machado, A. Biofilms of *Salmonella*: Implications for Food Safety and Public Health. *Foodborne Pathog Dis* https://doi.org/10.1177/15353141251389597 (2025) doi:10.1177/15353141251389597.

52. Römling, U., Sierralta, W. D., Eriksson, K. & Normark, S. Multicellular and aggregative behaviour of *Salmonella* Typhimurium strains is controlled by mutations in the *agfD* promoter. *Mol Microbiol* **28**, 249–264 (1998).

53. Fletcher, M. The effects of culture concentration and age, time, and temperature on bacterial attachment to polystyrene. *Can. J. Microbiol.* **23**, 1–6 (1977).

54. O'Toole, G. A. & Kolter, R. Initiation of biofilm formation in Pseudomonas fluorescens WCS365 proceeds via multiple, convergent signalling pathways: a genetic analysis. *Mol Microbiol* **28**, 449–461 (1998).

55. Stecher, B. *et al.* Motility allows S. Typhimurium to benefit from the mucosal defence. *Cell Microbiol* **10**, 1166–1180 (2008).

56. Gillen, K. L. & Hughes, K. T. Negative regulatory loci coupling flagellin synthesis to flagellar assembly in *Salmonella* Typhimurium. *J Bacteriol* **173**, 2301–2310 (1991).

57. Lockman, H. A. & Curtiss, R. *Salmonella* Typhimurium mutants lacking flagella or motility remain virulent in BALB/c mice. *Infect Immun* **58**, 137–143 (1990).

58. Baisón-Olmo, F., Cardenal-Muñoz, E. & Ramos-Morales, F. PipB2 is a substrate of the *Salmonella* pathogenicity island 1-encoded type III secretion system. *Biochem Biophys Res Commun* **423**, 240–246 (2012).

59. Segura, I., Casadesús, J. & Ramos-Morales, F. Use of mixed infections to study cell invasion and intracellular proliferation of *Salmonella enterica* in eukaryotic cell cultures. *J Microbiol Methods* **56**, 83–91 (2004).

60. Amadou Amani, S. & Lang, M. L. Bacteria That Cause Enteric Diseases Stimulate Distinct Humoral Immune Responses. *Front Immunol* **11**, 565648 (2020).

61. Aguilera-Herce, J. *et al. Salmonella* Type III Secretion Effector SrfJ: A Glucosylceramidase Affecting the Lipidome and the Transcriptome of Mammalian Host Cells. *Int J Mol Sci* **24**, 8403 (2023).

62. McCormick, B. A., Colgan, S. P., Delp-Archer, C., Miller, S. I. & Madara, J. L. *Salmonella* Typhimurium attachment to human intestinal epithelial monolayers: transcellular signalling to subepithelial neutrophils. *J Cell Biol* **123**, 895–907 (1993).

63. Hobbie, S., Chen, L. M., Davis, R. J. & Galán, J. E. Involvement of mitogen-activated protein kinase pathways in the nuclear responses and cytokine production induced by *Salmonella* Typhimurium in cultured intestinal epithelial cells. *J Immunol* **159**, 5550–5559 (1997).

64. Gewirtz, A. T. *et al. Salmonella* Typhimurium induces epithelial IL-8 expression via Ca(2+)-mediated activation of the NF-kappaB pathway. *J Clin Invest* **105**, 79–92 (2000).

65. He, H., Tan, H., Ge, C. & Xu, Y. Secretion analysis during the invasion process of *Salmonella* Typhimurium into HT-29 intestinal epithelial cells on a microfluidic chip. *Analyst* **150**, 3871–3879 (2025).

66. Haraga, A. & Miller, S. I. A *Salmonella enterica* serovar Typhimurium translocated leucine-rich repeat effector protein inhibits NF-kappa B-dependent gene expression. *Infect Immun* **71**, 4052–4058 (2003).

67. Olaru, I. D. *et al.* The association between antimicrobial resistance and HIV infection: a systematic review and meta-analysis. *Clin Microbiol Infect* **27**, 846–853 (2021).

68. Danielsen, A. S. *et al.* Clinical outcomes of antimicrobial resistance in cancer patients: a systematic review of multivariable models. *BMC Infect Dis* **23**, 247 (2023).

69. Jones, G. W., Rabert, D. K., Svinarich, D. M. & Whitfield, H. J. Association of adhesive, invasive, and virulent phenotypes of *Salmonella* Typhimurium with autonomous 60-megadalton plasmids. *Infect Immun* **38**, 476–486 (1982).

70. Römling, U. *et al.* Occurrence and regulation of the multicellular morphotype in *Salmonella* serovars important in human disease. *Int J Med Microbiol* **293**, 273–285 (2003).

71. Hébrard, M., Kröger, C., Sivasankaran, S. K., Händler, K. & Hinton, J. C. D. The challenge of relating gene expression to the virulence of *Salmonella enterica* serovar Typhimurium. *Curr Opin Biotechnol* **22**, 200–210 (2011).

72. Browning, K. R. & Merrikh, H. Pathogenic bacteria experience pervasive RNA polymerase backtracking during infection. *mBio* **15**, e0273723 (2024).

73. Bansal, G., Ghanem, M., Sears, K. T., Galen, J. E. & Tennant, S. M. Genetic engineering of *Salmonella* spp. for novel vaccine strategies and therapeutics. *EcoSal Plus* **12**, eesp00042023 (2024).

74. Jarvik, T., Smillie, C., Groisman, E. A. & Ochman, H. Short-term signatures of evolutionary change in the *Salmonella enterica* serovar Typhimurium 14028 genome. *J Bacteriol* **192**, 560–567 (2010).

75. Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc Natl Acad Sci U S A* **97**, 6640–6645 (2000).

76. Figueroa-Bossi, N., Balbontín, R. & Bossi, L. Scarless DNA Recombineering. *Cold Spring Harb Protoc* **2023**, 638–650 (2023).

77. Schmieger, H. Phage P22-mutants with increased or decreased transduction abilities. *Mol Gen Genet* **119**, 75–88 (1972).

78. Lutz, R. & Bujard, H. Independent and tight regulation of transcriptional units in *Escherichia coli* via the LacR/O, the TetR/O and AraC/I1-I2 regulatory elements. *Nucleic Acids Res* **25**, 1203–1210 (1997).

79. Balbontín, R., Vlamakis, H. & Kolter, R. Mutualistic interaction between *Salmonella enterica* and *Aspergillus niger* and its effects on *Zea mays* colonization. *Microb Biotechnol* **7**, 589–600 (2014).

80. Balbontín, R., Figueroa-Bossi, N. & Bossi, L. Construction of Single-Copy Fluorescent Protein Fusions by One-Step Recombineering. *Cold Spring Harb Protoc* **2023**, 651–662 (2023).

81. Maloy, S. R. & Roth, J. R. Regulation of proline utilization in *Salmonella* Typhimurium: characterization of *put::*Mu*d*(Ap, lac) operon fusions. *J Bacteriol* **154**, 561–568 (1983).

TABLES AND FIGURES

Table 1. List of genes in the set of luciferase reporter fusions

Gene	Location	Function
iagB	SPI-1	SPI-1 type III secretion system invasion protein
pipB2	Orphan	SPI-1 and SPI-2 effector
sseK3	ST64B	SPI-2 effector
sseC	SPI-2	SPI-2 effector
ssaV	SPI-2	Structural component of the SPI-2 type III secretion system
spvB	pSLT	Effector
kdpD	Orphan	Two-component system, sensor histidine kinase
rfaB	Orphan	Galactosyltransferase
proV	Orphan	ATP-binding protein of glycine betaine/L-proline ABC transporter
nixA	Orphan	Putative nickel/cobalt transporter
entF	Orphan	Enterobactin synthetase
hupA	Orphan	HU nucleoid-associated protein
lsrF	Orphan	Quorum Sensing, autoinducer 2 aldolase
luxS	Orphan	Quorum sensing, autoinducer 2 synthase
rhs	SPI-6	Type VI secretion system

Figure 1. Expression of luciferase reporter fusions to genes associated with *S. enterica* virulence.
A. Ratio *ΔrnhA*/WT of the area under the curve of luminiscence normalized to OD_{600} of different reporter fusions in LB (grey bars), LB 0.3M NaCl (red bars) or LPM (blue bars). Bars represent average and error bars standard deviation (n = 3 independent biological replicates). N.S. non-significant; *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001 (one-sample two-tailed Student's t test). **B-D.** Logarithmic representation of the expression of luciferase reporter fusions in the WT (X axis) and the *ΔrnhA* (Y axis) backgrounds, either in LB (**B**), LB 0.3M NaCl (**C**) or LPM (**D**). The black diagonal line represents the linear regression if expression levels were identical.

Figure 2. Lack of RNase HI affects fluorescence intensity of bacteria carrying the P_{sicA}-sfGFP fusion. A. Distribution of sfGFP fluorescence intensity in populations of wild-type (blue line) and $\Delta rnhA$ (red line) bacteria carrying the P_{sicA}-sfGFP fusion, and in a control not carrying the fusion (black line), in LB. The vertical green dashed line represents the threshold of fluorescent versus non-fluorescent bacteria, determined using non-fluorescent bacteria as control. **B.** Percentage of cells with the P_{sicA}-sfGFP fusion induced in wild-type and $\Delta rnhA$ bacteria, either in LB (top panel) or in NCE (bottom panel) media. Bars represent average and error bars standard deviation (n = 6 independent biological replicates). N.S. non-significant; *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001 (two-tailed Student's t test).

Figure 3. Lack of RNase HI causes reduction in biofilm formation. A. Colony morphology of wild-type bacteria (left panel), the *ΔrnhA* mutant (center panel) and the *ΔcsgD* mutant, unable to form biofilms (right panel), after 7 days of incubation at 30 ºC in LB agar without salt. **B.** Ratio between the OD_{580} (measuring concentration of crystal violet) and OD_{600} (measuring cell density) of wild-type bacteria, the *ΔrnhA* mutant and the *ΔcsgD* mutant, unable to form biofilms, after 48 h of incubation at 25 ºC in LB without salt. **C.** Percentage of biofilm formation of the *ΔrnhA* mutant with respect to wild-type bacteria (considered to form 100%) and the *ΔcsgD* mutant (considered to form 0%). Bars represent average and error bars standard deviation (n = 9 independent biological replicates). N.S. non-significant; *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001 (two-tailed Student's t test).

565 **Figure 4. RNase HI depletion negatively affects bacterial motility. A.** Motility (expressed in milimeters per hour) of wild-type bacteria, the *ΔrnhA* mutant, and the non-motile *ΔfliGHI* mutant. Blue lines represent mean values. **B.** Percentage of motility of the *ΔrnhA* mutant with respect to wild-type bacteria (considered to move 100%) and the *ΔfliGHI* mutant (considered to move 0%). Bars represent average and error bars standard deviation (n ≥ 20 independent biological replicates). N.S. non-

570 significant; *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001 (two-tailed Student's t test).

Figure 5. Lack of RNase HI affects expression of a P_{ssaB}-mCherry fusion during infection of epithelial cells. A. Fluorescence microscopy image of HeLa cells infected with wild-type (left panels) or *ΔrnhA* (right panels) bacteria constitutively expressing sfGFP at 2, 7 and 24 h after infection. **B.** Distribution of sfGFP and mCherry fluorescence intensity in wild-type (top panel) and *ΔrnhA* (bottom panel) bacteria carrying the P_{sicA}-*sfGFP* and P_{ssaB}-*mCherry* fusions, within HeLa epithelial cells at 30 minutes and 24 h after infection. Thresholds for sfGFP and mCherry fluorescences were determined using strains expressing either fluorescence constitutively and non-fluorescent control strains. Percentages represent the average of two independent experiments.

Figure 6. R-loop imbalance does not affect invasion of human epithelial cells. Rate of invasion

during individual infection of HeLa human epithelial cells of either wild-type bacteria (grey circles), a

mutant lacking the two major *Salmonella* Pathogenicity Islands SPI-1 and SPI-2 (black squares), as

controls, and the *ΔrnhA* mutant (red diamonds). Horizontal bars represent the mean (n ≥ 20). N.S. non-

significant; * = P < 0.05; ** = P < 0.01; *** = P < 0.001;**** = P < 0.0001 (one-sample two-tailed

605   Student's t-test).

Figure 7. Mutants lacking RNase HI are defective in proliferation within macrophages.
Competitive index in proliferation within murine macrophages of either wild-type bacteria (grey circles), a mutant lacking the two major *Salmonella* Pathogenicity Islands SPI-1 and SPI-2 (black squares), as negative control, and the *ΔrnhA* mutant (red diamonds). Horizontal bars represent the mean (n ≥ 3 independent biological replicates). The red dashed line indicates the value of mean competitive proliferation index when both genotypes show similar ability to proliferate. N.S. non-significant; * = P < 0.05; ** = P < 0.01; *** = P < 0.001;**** = P < 0.0001 (one-sample two-tailed Student's t-test).

Figure 8. Mutants lacking RNAse HI trigger a normal immune response. Ratio of IL-8 secretion by HeLa epithelial cells at 2 hours after infection by wild-type bacteria (grey circles), a mutant lacking the two major *Salmonella* Pathogenicity Islands SPI-1 and SPI-2 (black squares), as controls, the *ΔrnhA* mutant (red diamonds), and non-infected cells (green triangles). Data are presented as ratios between the corresponding value and the mean values of cells infected by wild-type bacteria. Blue lines represent mean values of six independent biological replicates in each group, with two technical replicates each. Red dashed lines indicate mean levels in non-infected cells and in cells infected by wild-type bacteria. N.S. non-significant; * = P < 0.05; ** = P < 0.01; *** = P < 0.001;**** = P < 0.0001 (one-sample two-tailed Student's t-test).

SUPPLEMENTARY FIGURES

Figure S1. Fitness of the *ΔrnhA* mutant compared to wild-type bacteria. A. Growth curves of wild-type (blue dots and line) and *ΔrnhA* mutant (red dots and line) bacteria in LB. Dots represent the average and error bars represent standard deviation (n ≥ 7 independent biological replicates, with two technical replicates each). **Inset:** Area Under the Curve (AUC) of wild-type (blue bar) and *ΔrnhA* mutant (red bar) in the growth curves shown. Bars represent average and error bars standard deviation (n ≥ 8 independent biological replicates, with two technical replicates each). N.S. non-significant; *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001 (one-sample two-tailed Student's t test). **B.** Competitive fitness of wild-type and *ΔrnhA* mutant bacteria against a labeled wild-type reference strain in LB. Bars represent average and error bars standard deviation (n ≥ 3 independent biological replicates). N.S. non-significant; *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001 (one-sample two-tailed Student's t test).

Figure S2. Lack of RNase HI increases sensitivity to low pH. Serially diluted wild-type (top lanes) and *ΔrnhA* (bottom lanes) in LB at pH 6.8 (top panel) or 4.9 (bottom panel). The experiment included

705   four biological replicates and four technical replicates; representative datasets are shown.