Density-dependent expression of epitranscriptomic, stress, and appetite regulating genes in Atlantic salmon

Authors

Morgane Frapin[1], Laura Quispe[1], Joana Troka[1], Jenni M. Prokkola[1,2], Ossi Laurikainen[3,4], Pekka Hyvärinen[5], Craig R. Primmer[1,6], Tutku Aykanat*[1], Ehsan Pashay Ahi*[1]

Affiliations

[1] *Organismal and Evolutionary Biology Research Program, Faculty of Biological and Environmental Sciences, PO Box 56, FI-00014 University of Helsinki, Finland*
[2] *Natural Resources Institute Finland (Luke), Oulu, Finland*
[3] *Natural Resources Institute Finland (Luke), Viikki, Helsinki, Finland*
[4] *Department of Environmental and Biological Sciences, Faculty of Science, Forestry and Technology, University of Eastern Finland*
[5] *Natural Resources Institute Finland (Luke), Paltamo, Finland*
[6] *Institute of Biotechnology, Helsinki Institute of Life Sciences (HiLIFE) University of Helsinki, Finland*

* co-senior authors.

Corresponding author: Morgane Frapin, morgane.frapin@helsinki.fi

Abstract

Intraspecific competition due to e.g., density, has substantial influence on fitness dynamics and life histories, but the underlying physiological mechanisms are often complex, and the molecular basis unclear. Further, designing laboratory experiments to measure physiological responses that reflect natural conditions is challenging. Here, we reared Atlantic salmon juveniles in semi-wild conditions in two naturally occurring densities to investigate the molecular mechanism of density-related changes in the hypothalamus, a key brain region regulating stress and energy homeostasis. We measured density-dependent changes in the expression of 12 genes involved in appetite and stress regulation, and 16 genes involved in post-transcriptional regulation of gene expression via m^6A RNA methylation. This gene set includes paralog pairs, to account for potential functional divergence associated with salmonid genome duplication. We also quantified genotype-environment interactions between density and two major life-history loci, *vgll3* and *six6*. We found significant density-related differences in the expression of genes coding for corticotropin-releasing factors, appetite stimulators and inhibitors, and m^6A RNA methylation actors. Moreover, a paralogue of an appetite inhibitor showed a density-dependent pattern that was the opposite of what was expected. *Six6* locus was also associated with changes in the expression of epitranscriptomic markers, including two writers and one eraser. Our results highlight that individuals' response to density in natural conditions is shaped by a complex interplay between stress, appetite and epitranscriptomic pathways in the hypothalamus. In addition, the functional divergence of paralogs indicates a potential role of genome duplication shaping such a response. We emphasize the value of integrating different physiological responses at the molecular level to better understand ecological processes affected by environmental change.

Keywords

life-history, teleost, semi-natural environment, m^6A RNA methylation, hypothalamus

1. Introduction

Population density is an important factor in ecology, affecting resource availability and social interactions among conspecifics (Nislow et al., 2010; Reznick, 2016). An increase in population density intensifies the intraspecific competition for resources, leading to alterations in energy homeostasis and stress levels, and subsequently impaired growth and reproductive success (Auer et al., 2012; Grossman & Simon, 2020; Reznick, 2016). Fishes are good models for studying density-dependent physiological responses, as density has a significant effect on their growth and survival (Grossman & Simon, 2020; Matte et al., 2020). In the wild, riverine habitats are important ecosystems that are particularly affected by density-dependent processes, due to the limited area and resources, providing a valuable case for studying soft selection and the effects of density changes (Jonsson et al., 1998). For example, the strong territorial behavior of Atlantic salmon to occupy favorable habitats can be beneficial in poor habitats, but may not be energetically efficient in resource rich habitats (Blanchet et al., 2006; Jonsson et al., 1998; Nislow et al., 2010). In laboratory or hatchery settings, fish are reared at much higher densities, and competition and food acquisition dynamics do not reflect the natural conditions. This is likely limiting individuals from displaying natural behaviors. As such, two habitats (*wild versus* laboratory conditions) are likely to result in different physiological responses to density changes (Blanchet et al., 2006; Höjesjö et al., 2004).

An individual's response to density may invoke multiple physiological and/or stress pathways, which can result in complex responses at the population level (Birk et al., 2020; Hodgson et al., 2017), with potential impacts on life histories (Archer et al., 2020). At the organismal level, such a response is regulated by hypothalamus, which exhibits distinct molecular pathways for various environmentally induced responses (Schwartz et al., 2000; Smith & Vale, 2006). The hypothalamus is a key region of the brain that connects the central nervous system to the endocrine system. It regulates feeding behavior by balancing appetite-stimulating genes (orexigenic genes), such as Agouti-related protein (AgRP) and neuropeptide Y (NPY), and appetite-suppressing genes (anorexigenic genes), such as cocaine- and amphetamine-regulated transcript (CART) and pro-opiomelanocortin (POMC). Together, these mechanisms help the body regulate energy intake and expenditure in response to nutritional status (Parker & Bloom, 2012). The hypothalamus is also functioning as the primary interface for stress detection by orchestrating the glucocorticoid receptor signaling pathway and initiating the secretion of corticotropin-releasing hormone/factor (CRH or CRF). This activates the hypothalamic-pituitary-adrenal axis, leading to the release of cortisol, a key stress hormone that prepares the body to respond to stressors (Smith and Vale 2006). Additionally, the hypothalamus plays a pivotal role in sexual maturation by releasing gonadotropin-releasing hormone, influencing the onset of puberty and reproductive capacity (Sisk & Foster, 2004). These regulatory roles of the hypothalamus are conserved

across vertebrates, underscoring its fundamental importance in maintaining homeostasis and enabling
responses to environmental changes and internal conditions (Denver, 2009). Thus, understanding
hypothalamic regulation may be critical in partitioning the relative contribution of different
physiological processes, such as stress and appetite pathways, which may help to pinpoint ecological
functions that underlie density-dependent dynamics in the wild.

Emerging research highlights the role of m^6A RNA methylation in modulating hypothalamic signals
related to feeding, stress, and sexual maturation. Thus, m^6A RNA methylation may play a role in
density-dependent regulation by controlling gene expression in response to environmental changes.
m^6A RNA methylation is a widespread, reversible, and evolutionarily conserved mRNA modification
(Ahi & Singh, 2024; Shi et al., 2022). The addition of m^6A RNA methylation is facilitated by the m^6A
methyltransferase complex, which involves writer proteins (such as Mettl3, Mettl14, and Wtap), and
removal is carried out by eraser demethylase proteins (such as Fto and Alkbh5) (Zaccara et al., 2019).
Reader proteins (e.g., Ythdf1/2/3 and Ythdc1/2) recognize the m^6A modifications, directing the
methylated mRNA towards various outcomes, including degradation, stabilization, transportation, and
either promotion or inhibition of translation (Liao et al., 2018; Zhao et al., 2017). Investigating whether
m^6A RNA modification affects hypothalamic signaling that controls stress and feeding in response to
environmental changes, such as population density, could provide insight into growth flexibility and
the proximal mechanism of phenotypic plasticity in changing environmental conditions (Jonsson et al.,
1998).

The Atlantic salmon (S*almo salar*) is a culturally and economically important anadromous fish species
that spends its juvenile phase in the freshwater. The freshwater phase is highly influenced by density
effects (Thorstad et al., 2010). While molecular and/or physiological response to density have been
widely investigated in Atlantic salmon reared in artificial conditions (Liu et al., 2015; Sundh et al., 2019;
Sveen et al., 2016; Y. Wang et al., 2019), these hardly reflect natural settings, and conclusions from
these studies can't necessarily be extrapolated to wild conditions. Studying the molecular/physiological
pathways of density regulation is feasible in Atlantic salmon in semi-natural environment, and
therefore more natural-like physiological responses can be obtained. Furthermore, Atlantic salmon
may also be a good model to study molecular mechanisms of life history variation and potential
interplay with hypothalamic regulation of density. Genetic variants located by *vgll3* gene in
chromosome 25 and *six6* gene in chromosome 9 have been associated with a sea age at maturity in
salmon (Barson et al., 2015; Sinclair-Waters et al., 2020). This simple genetic architecture is also
associated with various behavioral and physiological pathways (e.g., Aykanat et al., 2020; Bangura et
al., 2022; Prokkola et al., 2022), and a recent study reported that the hypothalamic expression of genes
involved in m^6A RNA methylation is influenced by *six6* locus, sex and maturation status in Atlantic

salmon (Ahi et al., 2024). As such these genes may partly be associated with hypothalamic endocrine pathways that respond to environmental, dietary, and physiological cues (Gerisch & Antebi, 2011; Stearns, 2011).

In this study, our primary aim is to investigate how changes in density affect fundamental molecular physiological processes in an environment that closely mimics natural conditions. We explore the hypothalamic expression of genes related to appetite regulation, stress hormone signaling pathways and m^6A RNA methylation in 2-year old Atlantic salmon parr reared in semi-natural stream conditions, and naturally fed on mainly benthic prey items, at two different densities similar to those found naturally in the wild (Finstad et al., 2009). In addition, the presence of the genetic variation at the two loci associated with sea age at maturity, *vgll3* and *six6*, allowed us to assess their effect, as well as their interaction with density, on the gene expression of the targeted hypothalamic pathways. Given the whole-genome duplication which occurred approximately 80 million years ago in salmonids (Dysin et al., 2022; Lien et al., 2016), we also investigate potential functional evolution of the duplicated genes by characterizing the expression pattern of paralogs (Force et al., 1999; Kassahn et al., 2009).

2. Materials and Methods

2.1. Fish rearing

All fish experiments were conducted in accordance with the Finnish Project Authorisation Board approval (ESAVI/4511/2020). In this experiment, Atlantic salmon were obtained from broodstocks reared by the Natural Resources Institute Finland (LUKE, Laukaa, Finland). The parents of the broodstocks, i.e., the grandparents of the experimental fish originated from the Kymijoki River in Southern Finland. From the broodstock, three unrelated males and females heterozygous for the alleles associated with earlier (E) and later (L) maturation at both the *vgll3* and *six6* loci were crossed in October 2019 to obtain three unrelated families each containing offspring with all combination of genotypes for these loci. Fertilized eggs were incubated in the dark at 7°C at the University of Helsinki until the alevin stage as described earlier (Debes et al., 2021), and then transferred late January 2020 to the Kainuu Fisheries Research Station of LUKE in Paltamo (Finland). The fish were reared in a seasonal temperature and daylight cycle and fed based on their size (Veronesi vita 0.2-0.8mm, Raisioagro Oy) in a flow-through system supplied with water from a nearby lake. In January 2021, fish were PIT-tagged (12 mm) and small fin clips were taken for genotyping individuals for sex and *vgll3* and *six6* loci (Sinclair-Waters et al., 2022). In May 2021, a subset of fish was released in six parallel outdoor channels (length: 25 meters, width: 1.5 meters, approx. 39 m^2) with concrete walls and bottoms covered by 15 cm layer of gravel-to-pebble-sized particles (8–50 mm in diameter). These

semi-natural stream channels were fed by water from the same lake as the indoor tanks to maintain a

flow rate of 0.15-0.25 ms^{-1}. The fish were naturally fed on mainly benthic prey items from the bottom

substrate (insects, arachnids, and crustaceans), but also some surface prey (such as adult midges,

mosquitoes), similar to their natural habitat. All fish selected in this experiment were smolts based on

silvery coloration. Each of the three families was divided between two stream channels with two

density treatments, which initially had 48 (1.2 fish/m^2) and 112 (2.8 fish/m^2) fish per low-density and

high-density treatment streams, respectively. However, an infection outbreak caused by *Saprolegnia*

spp. right after the transfer to the outdoor tanks inflicted high mortalities to the experimental fish (as

high as 46 % mortality in less than a month). To bolster the density effect, high-density tanks were

supplemented with additional fish, which were not included in subsequent analysis. These additional

fish originated from the same families and were reared in the same indoor tanks as the experimental

fish prior to the transfer to the natural stream channels. In August 2021, due to practical reasons, all

fish were transferred from the channels to six circular stream channels (Outer circumference: 26.15

meters, width: 1.5 meters, approx. 39.2 m^2) with concrete walls and a bottom covered with a 15 cm

layer of gravel (8-50 mm in diameter) and a gravity driven flow of 0.11 ms^{-1}. The densities by the end

of the experiments ranged from 0.35-0.51 fish/m^2 and 1.62-2.05 fish/m^2 for low-density and high-

density stream channels, respectively (see Supplementary table 1 for details).

2.2. Sampling and RNA extraction

On 25-27 April 2022, fish were caught from the streams by netting after water level in each stream was

lowered and placed in aerated 40-L buckets until sampling the same day. Fish were euthanized with

an overdose of benzocaine (200 mg/L), weighed, and measured. The hypothalamus of fish from both

densities and all genotypes were sampled, snap-frozen in liquid nitrogen and stored at -70°C prior to

processing.

To avoid introducing variation associated with maturation status, only immature individuals were

selected for this study. Maturity was assessed by checking the presence of milt in males (females are

immature at that age). The hypothalamus of 81 individuals (40 males and 41 females) was selected.

The individuals were either homozygous (EE or LL) or heterozygous (EL) for *six6* and *vgll3* loci (N = 3 –

10 samples per genotype, sex and density) (see, Supplementary table 2 for details). Total RNA was

extracted using Nucleospin RNA kit (Machery-Nagel). Homogenization and cell lysis steps were

performed simultaneously by transferring the frozen hypothalamus in a 2 mL screw cap tube

containing 1.4 mm ceramic beads (OMNI), 3.5 µL of DTT 1M and 350 µL of RA1 buffer and placing the

tubes in a Bead Ruptor Elite (OMNI) at 4 ms^{-1} for 3x20 seconds with a 10 second break in between.

The subsequent steps followed the manufacturer's instructions and RNA was eluted in 40 µL of RNase-

free water. RNA quantity and quality were assessed using Nanodrop 2000 (Thermo Fisher Scientific)

and agarose gel electrophoresis.

2.3. cDNA synthesis and qPCR

Synthesis of cDNA was performed using iScript cDNA Synthesis Kit (BioRad) with 1 µg of RNA. The

genes *ef1a* and *hprt1* were used as housekeeping genes (Ahi et al., 2024). The expression of six genes

related to stress (*nr3c1*, *nr3c2*, *crf1a1*, *crf1a2*, *crf1b1*, *crf1b2*), six genes related to appetite regulation

(*agrp1*, *cart2a*, *cart2b*, *pomca1*, *pomca2*, *npya1*) and 16 genes related to epitranscriptomic

mechanisms (*mettl3*, *mettl14*, *wtap*, *alkbh5-1*, *alkbh5-2*, *fto-1*, *fto-2*, *ythdc1-1*, *ythdc1-2*, *ythdc2*,

ythdf1-1, *ythdf1-2*, *ythdf1-3*, *ythdf2-1*, *ythdf2-2*, *ythdf3*) were measured using quantitative real-time

PCR (qPCR) (genes and primer sequences are described in Supplementary table 3). The qPCR reactions

were performed in a final volume of 10 µL containing 1 µL of cDNA diluted 1:10 in nuclease-free water,

250 nM of forward and reverse primers and 5 µL of PowerUp™ SYBR™ Green Master Mix

(ThermoFisher Scientific). The qPCRs were performed on 384-well plates using the BioRad CFX384 Real

Time PCR System, with each plate measuring the expression of a single target gene across all samples

that were present in triplicate. The Cqs (quantification cycle) were obtained with a manually fixed

threshold located in the exponential part of the amplification. The qPCR program and primer efficiency

calculation were performed as described in Ahi, Richter, et al. (2019). For each sample, the Cq values

of the triplicates obtained for each gene were averaged. The ΔCq values were calculated by subtracting

the geometric mean of the Cq values of the housekeeping genes, *ef1a* and *hprt1*, from the Cq values

of the target genes. Then, for each gene, the ΔCq values were subtracted by the ΔCq value of the

sample with the lowest expression to obtain the ΔΔCq. The relative quantifications (RQ) were

calculated as $2^{-\Delta\Delta Cq}$ and statistical analyses used the $\log_2$ RQ (Livak & Schmittgen, 2001).

2.4. m^{6}A RNA methylation quantification

Quantification of m^{6}A RNA methylation was performed on the 81 total RNA extracted from the

hypothalamus using the EpiQuikTM m6A RNA Methylation Quantification Kit (Epigentek). Since one

preparation kit can accommodate a maximum of 96 tests, two kits were used to include all samples in

duplicate, along with negative and positive controls. Samples were randomly assigned between the

two kits to account for potential batch effects in the statistical model. RNA samples were diluted to

33.33 ng/µL in RNase-free water, and the concentration was verified with Nanodrop 2000 (Thermo

Scientifc). m^{6}A RNA methylation quantification was performed according to the manufacturer's

protocol with 6 µL of the diluted total RNA (approximately 200 ng). Negative and positive controls from

the kit were included in each plate. The absorbance was then read at 450 nm with the EnSpire

Multimode Plate Reader (PerkinElmer). m^{6}A percentages were calculated according to the

manufacturer's protocol based on the standard curve and with the amount of total RNA input adjusted based on the actual concentrations of the dilutions used.

2.5. Statistical analysis

All the statistical analysis were done in R (version 4.2.2). A linear mixed effect model was fitted using lmer function in the *lme4* package (version 1.1-31) (Bates et al., 2015), with REML=FALSE option, using a general formula as below:

$$\text{response} \sim \text{density} * (\text{sex} + \textit{six6} + \textit{vgll3}) + \varepsilon_{\text{Family}} + \varepsilon_{\text{Batch}} + \varepsilon_{\text{error}}$$

In which, response is either *body mass, gene expression or m^6A %*. Density was modelled as a fixed effect interacting with sex, *six6* and *vgll3* genotypes. Genotypes were coded categorically to account for non-additivity. $\varepsilon_{\text{Family}}$, $\varepsilon_{\text{Batch}}$ are error terms associated with family and batch effect, and $\varepsilon_{\text{error}}$ is the residual normal variation. $\varepsilon_{\text{Batch}}$ was only included for the *m^6A % response* to account for the kit effect. The significance of the fixed effects terms was estimated using type 1 F-test using the anova function in the stats package (version 4.2.2) in R. The normal distribution of the model residuals was checked using the simulateResiduals function in the *DHARMa* package (version 0.4.6, (Hartig et al., 2024). P-values for the gene expression analysis were adjusted for multiple comparisons per model term (i.e., density, sex, *six6* genotype, *vgll3* genotype, and the interaction terms: density × sex, density × *six6*, density × *vgll3*) using the Benjamini-Hochberg method to control the false discovery rate (Benjamini & Hochberg, 1995). An adjusted p-value below 0.05 was considered statistically significant. For terms with significant effects, post-hoc pairwise comparisons were performed using estimated marginal means (EMMs) via the *emmeans* package (version 1.8.2, (Lenth et al., 2024). These pairwise comparisons allowed us to examine specific contrasts between the levels of significant variables. The p-values presented in the graphs are the adjusted p-values, obtained from these post-hoc pairwise comparisons and corrected using the Benjamini-Hochberg method. The p-values in the text correspond to the adjusted p-values obtained after adjusting the ANOVA results for the false discovery rate.

To investigate potential interactions between the target genes, body mass, and m^6A percentage, we performed a correlation analysis. Pearson correlations were calculated using cor.test function (*stats* package version 4.2.2) on the residuals of the fitted body mass, gene expression and m^6A percentage models. This analysis was carried out separately for the high- and low-density groups.

3. Results

3.1. Fish body mass was influenced by density treatment

After one year in different densities, fish body mass was significantly lower in high-density (p<0.001; $F_{1;78.37}$=80.425) (Figure 1). However, neither sex nor the *six6* and *vgll3* genotypes had a significant effect on fish body mass.

Figure 1: *Fish body mass (in grams) measured after one year under density treatment. Each dot represents the body mass of an individual fish in the high-density (HD) group (n = 44, orange dots) and low-density (LD) group (n = 37, blue dots). The black points display the estimated marginal means for each treatment group. Error bars indicate 95% confidence intervals. ***: p-value<0.001.*

3.2. Density and genetic factors affect expression levels of Corticotropin-Releasing factor genes

Density, and *six6* and *vgll3* genotypes influenced the expression of corticotropin-releasing factor (*crf*) genes but not the expression of glucocorticoid receptor (*nr3c*) genes (Supplementary table 4). The level of expression of *crf1a1* was significantly higher in the high-density treatment (p=0.025; $F_{1;79.26}$=8.598) while *crf1b1* expression was significantly lower (p=0.003; $F_{1;81}$=13.801). Moreover, *crf1a1* expression was also significantly influenced by *six6* genotype (p=0.018, $F_{2,78}$=6.78) with a higher level of expression in *six*6*EE (95% CI=1.64-2.66) compared to *six6**EL genotype individuals (95% CI=0.84-1.91). An interaction effect between density and *vgll3* genotype was found for one of the corticotropin-releasing factor paralogs, *crf1b2* (p=0.039, $F_{2,79}$=7.15), with a higher level of expression of the gene in *vgll3**LL in low-density (95% CI=1.02–1.43) compared to *vgll3**EE (95% CI=0.44-0.91) and *vgll3**EL (95% CI=0.49-1.02), and compared to *vgll3**LL in high density (95% CI=0.51-0.91, Figure 2).

Figure 2: *Relative mRNA expression (log₂RQ) level of stress-related genes affected by density **(a,b)**, six6 **(c)** and the interaction between vgll3 and density **(d)**. Error bars indicate 95% confidence intervals. *: p-value<0.05; **: p-value<0.01; ***: p-value<0.001. **(e)** Diagram summarizing the implication of the studied genes in stress regulation. Stress is sensed by the hypothalamus. This triggers a hormonal cascade initiated by the release of corticotropin-releasing hormone/factor (CRH/CRF). Secretion of CRH stimulates the release of adrenocorticotropic hormone (ACTH) into the bloodstream, which then acts on the interrenal cells located in the fish head kidney, stimulating the release of cortisol (Gorissen & Flik, 2016). Cortisol binds to glucocorticoid or mineralocorticoid receptors present in various tissues, e.g. NR3C, leading to physiological and metabolic adaptations in response to the stressor and affecting fish growth* (Barton & Iwama, 1991; Takahashi & Sakamoto, 2013)

3.3. Density influences the expression of both anorexigenic and orexigenic peptide-coding genes

The expression levels of genes encoding both appetite-stimulating (Agrp) and appetite-inhibiting (Cart2a/b) peptides were affected by density treatment (Figure 3) (Supplementary table 4). The level of expression of *agrp1* was significantly lower at low-density (p<0.001; $F_{1;79}$=51.64). The paralogs *cart2a* (p=0.05; $F_{1;81}$=6.51) and *cart2b* (p=0.004; $F_{1;81}$=12.67) showed opposite patterns, with *cart2a* showing higher expression and *cart2b* lower expression at low-density. The level of expression of the

cart2a was also influenced by *six6* genotype (p=0.006; $F_{2;81}$=9.38), where heterozygote (95% CI=0.81-

1.43) exhibited significantly lower expression compared to homozygotes (95% CI=1.62-2.15 and 1.31-

1.77 for *six6**EE and *six6**LL, respectively).

Figure 3: *Relative mRNA expression (log$_2$RQ) level of appetite-related genes affected by density **(a,b,c)***

*and six6 **(d)**. Error bars indicate 95% confidence intervals. *: p-value<0.05; **: p-value<0.01; ***: p-*

*value<0.001. **(e)** Diagram summarizing the implication of the studied genes in the appetite regulation.*

Orexigenic (AGRP and NPY) and anorexigenic (POMC and CART) neurons in the arcuate nucleus of the

hypothalamus detect the presence of hormones (insulin, leptin) in the bloodstream and induce an

increase (orexigenic response) or a decrease (anorexigenic response) in food intake respectively

(Schneeberger et al., 2014).

3.4. Density and *six6* genotype affect the expression of genes encoding m⁶A regulators without

altering overall m⁶A levels

Both density and *six6* genotype influenced the expression of genes encoding m⁶A RNA methylation

writers (Supplementary table 4). The level of expression of *mettl3* (p=0.034; $F_{1;81}$=7.55) and *ythdf1.2*

(p=0.003; $F_{1;80}$=14.82) was high in low-density group. The *six6* genotype influenced the expression

levels of two writers, *mettl3* (p=0.006; $F_{2;81}$=8.52) and *wtap* (p=0.031; $F_{2;81}$=5.81). For *mettl3*, expression was lower in *six6*EL* (95% CI=0.35-0.62) compared to *six6*LL* (95% CI=0.69-0.89). *wtap* expression was reduced in *six6*EL* (95% CI=0.66-1.04) compared to both *six6*EE* and *six6*LL* (95% CI=1.05-1.38 and 1.01-1.30 for *six6*EE* and *six6*LL*, respectively). The level of expression of *fto1*, an m^6A eraser, was also significantly affected by the *six6* genotype (p=0.041; $F_{2;78}$=5.25) with a lower level of expression in *six6*EL* (95% CI=0.58-1.25) compared to either of the homozygous genotypes (95% CI=0.84-1.52 and 0.81-1.48 for *six6*EE* and *six6*LL*, respectively, Figure 4). The analysis of the percentage of m^6A in total hypothalamic RNA showed no significant effects for density (p=0.834; $F_{1;80}$=0.04), sex (p=0.832; $F_{1;78}$=0.05), *six6* (p=0.054; $F_{2;79}$=3.02) and *vgll3* (p=0.237; $F_{2;81}$=1.46) genotypes.

Figure 4: *Relative mRNA expression (log₂RQ) level of m^6A RNA methylation-related genes affected by density **(a,b)** and six6 **(c,d,e)**. Error bars indicate 95% confidence intervals. *: p-value<0.05; **: p-value<0.01; ***: p-value<0.001. **(f)** Diagramm summarizing the implication of the studied genes in m^6A RNA methylation regulation. m6A RNA methylation is added by writers (METTL3, METTL14, WTAP), forming a complex and removed by erasers (FTO and ALKBH5). The fate of RNA is determined by readers (such as the YTH family of proteins) that direct RNA to be translated, transported or degraded (Zaccara et al., 2019).*

3.5. Correlation between gene expression levels and m⁶A RNA methylation quantity

The expression levels of *nr3c1* and *nr3c2* were significantly correlated with several genes related to m^{6}A RNA methylation in both high-density (Supplementary figure 1) and low-density (Supplementary figure 2). Furthermore, *nr3c1* was negatively correlated with m^{6}A RNA methylation percentages only in the low-density group (R=-0.48 (p=0.002)) and strongly positively associated with the level of expression of *fto1* in both densities (R=0.7 (p<0.001) in high-density and R=0.8 (p<0.001) in low-density, Figure 5). In the low-density treatment, *cart2b* was negatively correlated with several m^{6}A readers (*ythdc1.2*: R=-0.72 (p<0.001); *ythdc2.1*: R=-0.6 (p<0.001); *ythdf2.2*: R=-0.4 (p=0.01); *ythdf3.1*: R=-0.58 (p<0.001)) but was strongly positively correlated with *alkbh5.1* (R=0.74 (p<0.001)). No strong correlation was found between *alkbh5* paralogs, *alkbh5.1* and *alkbh5.2*, however in low-density condition these two genes showed an opposite correlation with m^{6}A RNA percentage (*alkbh5.1*: R=0.1 (p=0.50) in high-density and R=0.38 (p=0.02) in low-density; *alkbh5.2*: R=0.11 (p=0.48) in high-density and R=-0.38 (p=0.02) in low-density)

A positive correlation between two appetite related genes, *agrp1* and *cart2b,* was observed exclusively in the high-density group (R=0.64 (*p*<0.001), Figure 5). These two genes were also negatively correlated with body mass in the high-density treatment (*agrp1*-body mass: R=-0.56 (p<0.001); *cart2b*-body mass: R=-0.62 (p<0.001)).

Figure 5: *Correlation plots of residual expression levels at different densities **(a)** between fto1 (m^{6}A-related gene) and nr3c1 (stress-related gene), and **(b)** between agrp1 and cart2b (appetite-related genes). R: Pearson correlation coefficient.*

4. Discussion

Individuals can respond to changes in population density by adjusting appetite and stress levels (Nislow et al., 2010; Reznick, 2016), both of which would be regulated via the hypothalamus. The degree of their response to density may also influence growth and, consequently, life-history traits such as timing of maturation. However, little is known about how changes in density affect related molecular pathways in the wild. In this study, we assessed, in a semi-natural environment, the effects of population density on the expression levels of genes related to stress and appetite, and m^6A RNA methylation in the hypothalamus of Atlantic salmon, in relation to the genotypes of two loci associated with age at maturity (*six6* and *vgll3*). We found that density and *six6* genotype influenced the expression of several genes related to stress, appetite and m^6A RNA methylation. A *vgll3* genotype effect was also found on a stress-related gene (*crf1b2*) in low-density condition. Our results also suggested widespread expression differences in paralogue gene pairs, which suggests functional divergence in these molecular pathways since the recent genome duplication event (Lien et al., 2016).

4.1. Density effects on stress-related genes suggest functional divergence of the *crf* paralogs

Since population density may influence stress levels in fish by affecting food availability and social interactions, we tested the effect of density on the expression of genes encoding corticotropin-releasing factor, CRF, and glucocorticoid receptors, NR3C, which are important ubiquitous stress modulators. We found paralog-specific expression differences; while *crf1a1* was upregulated in the high-density, *crf1a2* expression was not altered, and *crf1b1* was upregulated in the low-density group. Furthermore, in contrast to the two *crf1b* paralogs, the expression of the two *crf1a* paralogs was not significantly correlated. Previous research in aquaculture stocks has shown that the expression of *crf* increases significantly under stress conditions in the hypothalamus of rainbow trout (Conde-Sieira et al., 2010) and in the brain of the Senegalese sole, *Solea senegalensis,* (Wunderink et al., 2011). Interestingly, some studies have shown the opposite trend, with a decrease in *crf* expression under stress conditions, for example in the whole brain of zebrafish exposed to acute restraint stress (Ghisleni et al., 2012) and in a cichlid fish (*Astatotilapia burtoni*) after a month of social stress (Chen & Fernald, 2008). This could be due to a different response depending on the stress conditions, but also to different functions of the *crf* paralogs given that the paralogue studied in teleost fish is not always clearly stated (Grone & Maruska, 2015). Indeed, a recent study showed that all the *crf* paralogs (*crf1a1/2*, *crf1b1/2*) found in Atlantic salmon do not exhibit the same changes in expression patterns under stress conditions, suggesting paralog specificity (Lai et al., 2021). Lai et al., (2021) showed that hypothalamic *crf1a1* expression increased in response to hypoxia and/or hunting stress, in a pattern similar to serum cortisol levels, while *crf1a2* expression remained unchanged. The expression of the *crf1b1* and *crf1b2* paralogs was only altered during the combination of these two stress stimuli (Lai et

369 al., 2021). This potential functional divergence of the paralogs may indicate that certain paralogs are
370 more closely associated with the anorexigenic function previously described in teleosts (Bernier, 2006),
371 while others have a function more related to stress response.

4.2.*cart2b* may have an orexigenic effect in high-density conditions

The effect of population density was also investigated on genes involved in appetite regulation,
including those encoding orexigenic peptides that stimulate appetite (*agrp* and *npy*), as well as
anorexigenic peptides that inhibit appetite (*pomc* and *cart*). We found that *agrp1* showed the strongest
response to density, with a highly significant, approximately 2-fold increase in expression at high-
density compared to low-density. This might be due to its orexigenic role in fish and/or its
responsiveness to stressors. As in many vertebrates, a*grp* has orexigenic effects in several fish species
and its expression is induced in response to lack of food availability (Ahi, Brunel, et al., 2019; Cerdá-
Reverter & Peter, 2003; Wei et al., 2013). Recent studies of Atlantic salmon have also shown that the
hypothalamic expression levels of *agrp1* was increased after exposure to fasting (Kalananthan et al.,
2020, 2023), suggesting a similar orexigenic role in salmonids. On the other hand, the expression of
agrp1 also significantly increased in response to stress in zebrafish (Cortés et al., 2018). Therefore, the
strong induction of *agrp1* in high density conditions might be an orexigenic response due to lower food
availability as well as its response to the stress related to inter-individual competition in higher
densities.

We found paralog-specific responses of *cart* to density; *cart2b* had a significantly higher expression in
high-density, whereas *cart2a* showed an increased expression in low-density. In zebrafish and many
other teleost fishes, *cart* plays anorexigenic role (Volkoff, 2016), however, a study in zebrafish
suggested that not all *cart* genes follow a similar expression pattern in response to changes in food
intake (Akash et al., 2014). The increased expression of *cart2b* observed in high-density does not
support the general anorexigenic role of *cart* genes, as less food availability in high-density is expected
to reduce expression of anorexigenic genes. However, the tendency of *cart2a* towards increased
expression in low-density follows the expected anorexigenic role, thus consistent with a hypothesis
that suggests a potential functional divergence for *cart2a* and *cart2b* paralogs in the hypothalamus of
Atlantic salmon. Interestingly, previous studies of Atlantic salmon also revealed a similar unexpected
expression pattern for *cart2b* paralog, where its hypothalamic expression has been found significantly
higher in fasted groups in response to both short- and long-term fasting (Kalananthan et al., 2021,
2023). In these studies, the authors suggested a potential orexigenic role for *cart2b* in the
hypothalamus of Atlantic salmon, which is uncommon across teleost fish and remained to be
functionally elucidated. Moreover, in our study, *cart2b* expression was strongly correlated with the

orexigenic gene, *agrp1*, but only in high-density. Taken together, these results suggest that the neo-functionalized *cart2b* has obtained a role that appears to be opposite to the function previously described for CART, and which may be a mechanism that helps salmon adapt to changing environmental conditions. Furthermore, both *agrp1* and *cart2b* expression levels were negatively correlated with body mass, but only under high-density conditions. The negative correlation with *agrp1* is consistent with expectations, as reduced food availability in such conditions would likely lead to lower body mass and an increase in *agrp1* expression, a gene known to stimulate appetite. In contrast, the negative correlation with *cart2b* is unexpected for a gene that encodes an anorexigenic peptide. This finding aligns with previous observation that *cart2b* may have undergone functional divergence following genome duplication in Atlantic salmon (Kalananthan et al., 2021, 2023) resulting in a modified or divergent role from its original function.

4.3. Density affects the expression of m^6A RNA methylation writer and reader genes

Given that m^6A RNA methylation has been shown to be a mechanism sensitive to environmental changes (Frapin et al., 2020; Yang et al., 2022), such as variation in food availability and stress, we investigated the impact of population density on the expression of genes coding for writers, erasers, and readers of m^6A RNA methylation.

We found a reduction in the hypothalamic expression of a writer, *mettl3*, and a reader, *ythdf1* (*ythdf1-2* paralog), in high-density. This suggests the potential role of these factors in mediating density response to stress and/or food availability in hypothalamus of Atlantic salmon. In recent years, the function of m^6A RNA methylation factors in mediating stress signals has gained significant interest, with discoveries pointing to their versatile, tissue-specific, and context-dependent roles (Wilkinson et al., 2021). *mettl3* is a well-characterized and highly conserved m^6A writer, the expression of which was reduced in high-density compared to low-density. Decreased *mettl3* expression under acute stress was earlier reported in the hypothalamus of mice (Engel et al., 2018). Strikingly, it seems that the stress mediated role of *mettl3* in the central nervous system is also highly conserved as a similar role was proposed in the *Drosophila* brain (Perlegos et al., 2022). The second density-affected m^6A RNA methylation marker, *ythdf1-2*, encodes an m^6A reader protein that facilitates translation initiation (X. Wang et al., 2015). Interestingly, findings in mammals have already revealed that *ythdf1* plays a role in various stress signals such as oxidative stress, hypoxia, heat and inflammatory stresses (He et al., 2022; Lu et al., 2019; Wilkinson et al., 2021). A recent study in rainbow trout also found several *ythdf1* paralogs to be responsive to heat and inflammatory stresses in various tissues (Yu et al., 2023). Although, to our knowledge, no study has indicated its role in stress signals of social competition, a novel discovery has shown that Ythdf1 is the main mediator of microbiota-dependent brain-gut

crosstalk in mice (Huang et al., 2023). This raises a topic for future research, to understand whether

ythdf1-2 plays a role in translating gut signals triggered by food availability to the hypothalamus in

salmon kept in different densities. The answer to this question is particularly interesting because it has

been already demonstrated that microbiota-dependent brain-gut crosstalk is essential for regulation

of both appetite and body temperature (Gabanyi et al., 2022).

4.4. Interconnection of stress, appetite regulation and m^{6}A RNA methylation mechanism

In this study, several lines of evidence suggest that the processes/mechanisms targeted are

interrelated. We observed significant correlations between *nr3c* paralogs and several m^{6}A RNA

methylation actors in both low- and high-density conditions. Notably, the strongest correlation was

found between the gene coding for the main glucocorticoid receptor, *nr3c1,* and the m6A eraser *fto1,*

and we also detected a negative correlation between *nr3c1* expression and the percentage of m^{6}A RNA

modifications only in the low-density condition. Interestingly, an association between *nr3c1* and *fto1,*

has also been reported in recent studies, suggesting a dynamic and reciprocal regulation between FTO

and NR3C1, mediated by m^{6}A modifications (Roy et al., 2022; T. Wu et al., 2023). Moreover, our data

suggest that this regulatory mechanism responds to the environment. Under high-density condition,

while a correlation between *fto1* and *nr3c1* was still observed, there was no significant correlation

between *nr3c1* and m^{6}A methylation levels. This suggests that external factors, such as density-related

stress, may alter the m^{6}A-mediated regulation of these genes. This highlights the potential for a more

complex, condition-specific interaction between *nr3c1* and *fto* that could influence m^{6}A modifications

and their downstream effects on gene expression.

Although we did not find any clear density-related difference in m^{6}A RNA methylation levels, we cannot

exclude the possibility that it is affected by density. Indeed, based on our results, this can be argued

by (1) the effect of density on the expression of the gene encoding the core of the writing complex,

mettl3, and (2) the effect of density on genes previously shown to be direct targets of m^{6}A RNA

methylation, such as CART mRNA in mice (Song et al., 2020; P.-F. Wu et al., 2021) and *crh* in chicken

(Yang et al., 2022). Furthermore, we observed that *cart2b* was correlated with several m^{6}A RNA readers

under low-density conditions, including paralogs of genes involved in alternative splicing (*ythdc1.2*), in

fate transition (*ythdc2.1*), in mRNA clearance (*ythdf2.2*) and translation (*ythdf3*.1) (Li et al., 2017; Liao

et al., 2018). This suggests that *cart2b* may be regulated by a variety of m6A-modifying proteins, which

play important roles in post-transcriptional processes. However, further investigation is required to

understand what the mechanism behind the observed difference in *cart2b* expression between the

two densities.

4.5. Gene expression associations with maturation locus genotypes

In Atlantic salmon, two loci near/across the genes *vgll3* and *six6* have been associated with sea age at maturity (Barson et al., 2015; Sinclair-Waters et al., 2020). Investigating whether genotypes at these loci influence gene expression in response to environmental factors such as population density, which can affect body mass, could provide insight into how genetic and environmental factors together regulate maturation.

In this study*, six6* genotype was associated with the expression of one gene related to stress, *crf1a1*, one gene related to appetite, *cart2a*, and three genes related to m^{6}A RNA methylation, *mettl3, wtap* and *fto1*. For all these genes, the *six6**EL individuals had lower expression compared with *six6**EE and/or *six6**LL. Previous study of gene expression influenced by *six6* genotypes has only compared homozygous individuals (Ahi et al., 2024). In this study, we included heterozygotes and found notable differences in gene expression between heterozygous and homozygous individuals which may reflect a potential heterozygote advantage that warrants further investigation (Hedrick, 2012). As *six6* is coding for a transcription factor, it has the potential to directly regulate the expression of other genes. In addition, it remains an interesting area for future research to investigate whether six6 might influence gene expression through m^{6}A methylation mechanisms. Although no differences in overall m^{6}A RNA methylation levels were found between the *six6* genotypes, a more targeted analysis focusing on m^{6}A methylation at the gene-specific level - particularly for genes affected by the *six6* genotype - would provide more informative insights.

We also found lower hypothalamic expression of *crf1b2* in the high-density compared to low-density condition, specifically in individuals with the *vgll3**LL genotype. It has been shown in Atlantic salmon that *crf1b2* paralogue is significantly induced in the telencephalon and hypothalamus in response to chasing stress (Lai et al., 2021). Moreover, in rainbow trout, CRF administration reduces aggressive behaviors but increases anxiety-like behaviors in a dose-dependent manner, suggesting a dual role of CRF1 in regulating both aggression and anxiety in the brain (Backström et al., 2011). Similarly, in rodents, lower doses of CRF can induce aggression, while higher doses tend to trigger anxiety-like responses (Hostetler & Ryabinin, 2013). Notably, a recent study showed that Atlantic salmon with the *vgll3**LL genotype exhibit higher aggression than those with the *vgll3**EE genotype (Bangura et al., 2022). This suggests that individuals with different *vgll3* genotypes may show distinct behavioral responses under stress, such as density pressure. It is possible that *vgll3* influences aggression in salmon by differentially regulating *crf1b2* in the hypothalamus, where the balance between aggression and anxiety-like behaviors may shift more drastically in *vgll3**LL individuals under high-density conditions due to altered *crf1b2* expression. Although no direct regulatory connection between *vgll3* (or the Hippo pathway) and *crf* genes has been identified in any species, the Hippo-vgll3 signal may

indirectly modulate *CRF* gene expression through crosstalk with stress-related pathways, such as the
glucocorticoid pathway, which acts upstream of *CRF* genes (Stepan et al., 2018).

5. Conclusions

Due to the importance of Atlantic salmon in aquaculture, the effects of density on gene expression
have mainly been studied under farmed conditions (Conde-Sieira et al., 2010; Wunderink et al., 2011).
Therefore, studies such as our that use near-natural densities can shed light on gene expression
responses in more natural conditions. In this study, we found that the expression of genes related to
stress, appetite and m^6A RNA methylation were affected by density under semi-natural conditions. Our
results suggest an interconnection between these three pathways. In addition, both *six6* and *vgll3*
genotypes, which are associated with sea age at maturity in Atlantic salmon, appear to influence the
expression of some of the genes studied. An intriguing finding regarding the *vgll3* genotype is that it
may influence anxiety-related genes in a density-independent manner.

Our results have broader implications for other species, especially those with comparable ecological
or evolutionary pressures, by highlighting how environmental factors and genetic variation interact to
influence gene expression. Furthermore, the observed interconnection between the stress and
appetite regulation, and m^6A RNA methylation, which are well-conserved mechanisms across
vertebrates (Ahi & Singh, 2024; Denver, 2009; Rønnestad et al., 2017), provided a framework for
understanding these relationships in other species. Finally, our study highlights the importance of a
deeper investigation of paralogs in salmonids, and other species with recent genome duplication, to
better understand their specific functions, especially in adapting to environmental changes.

Author Contributions

Experimental design (TA, JMP, OL, PH), conducting experiment (TA, JMP, OL, CRP), genotyping (TA,
JMP), molecular experiment design (MF, EPA), molecular experiment (LQ, TA, MF), statistical analysis
(MF, TA), funding (TA, CRP), writing - original draft (MF, EPA), writing – review and editing (MF, TA, EPA,
JMP, CRP, PH, JT, OL).

Acknowledgments

Funding was provided to CRP by Academy of Finland (grant numbers 307593, 302873, 327255 and
342851), the University of Helsinki, and the European Research Council under the European Articles
Union's Horizon 2020 and Horizon Europe research and innovation programs (grant nos. 742312 and
101054307). Views and opinions expressed are however those of the author(s) only and do not

necessarily reflect those of the European Union or the European Research Council Executive Agency. Neither the European Union nor the granting authority can be held responsible for them. Funding was provided to TA by the Research Council of Finland (grant nos. 328860, 353388, and 325964). We would like to thank the Bodossaki Foundation for their financial support, which enabled Joana Troka to realize her MSc studies at the University of Helsinki. We thank Matyas Paszko, Atakan Yildiz, Paul Bangura, Ari Leinonen, Lison Gueguen, Ronan O'Sullivan, Nikolai Piavchenko, and Tommi Junnonaho for their help with fish rearing and experiment.

Data Accessibility and Benefit-Sharing

All the data generated in this study are included in this article as supplementary file.

References

Ahi, E. P., Brunel, M., Tsakoumis, E., & Schmitz, M. (2019). Transcriptional study of appetite regulating genes in the brain of zebrafish (Danio rerio) with impaired leptin signalling. *Scientific Reports*, *9*(1), 20166. https://doi.org/10.1038/s41598-019-56779-z

Ahi, E. P., Frapin, M., Hukkanen, M., & Primmer, C. R. (2024). Hypothalamic expression analysis of m6A RNA methylation associated genes suggests a potential role of epitranscriptomics in sexual maturation of Atlantic salmon. *Aquaculture*, *579*, 740151. https://doi.org/10.1016/j.aquaculture.2023.740151

Ahi, E. P., Richter, F., Lecaudey, L. A., & Sefc, K. M. (2019). Gene expression profiling suggests differences in molecular mechanisms of fin elongation between cichlid species. *Scientific Reports*, *9*(1), 9052. https://doi.org/10.1038/s41598-019-45599-w

Ahi, E. P., & Singh, P. (2024). Emerging Orchestrator of Ecological Adaptation: m6A Regulation of Post-Transcriptional Mechanisms. *Molecular Ecology*, e17545. https://doi.org/10.1111/mec.17545

Akash, G., Kaniganti, T., Tiwari, N. K., Subhedar, N. K., & Ghose, A. (2014). Differential distribution and energy status–dependent regulation of the four CART neuropeptide genes in the zebrafish brain. *Journal of Comparative Neurology*, *522*(10), 2266–2285. https://doi.org/10.1002/cne.23532

Archer, L. C., Hutton, S. A., Harman, L., McCormick, S. D., O'Grady, M. N., Kerry, J. P., Poole, W. R., Gargan, P., McGinnity, P., & Reed, T. E. (2020). Food and temperature stressors have opposing effects in determining flexible migration decisions in brown trout (Salmo trutta). *Global Change Biology*, *26*(5), 2878–2896. https://doi.org/10.1111/gcb.14990

Auer, S. K., Lopez-Sepulcre, A., Heatherly II, T., Kohler, T. J., Bassar, R. D., Thomas, S. A., & Reznick, D. N. (2012). Life histories have a history: Effects of past and present conditions on adult somatic growth rates in wild Trinidadian guppies. *Journal of Animal Ecology*, *81*(4), 818–826. https://doi.org/10.1111/j.1365-2656.2012.01964.x

Aykanat, T., Rasmussen, M., Ozerov, M., Niemelä, E., Paulin, L., Vähä, J.-P., Hindar, K., Wennevik, V., Pedersen, T., Svenning, M.-A., & Primmer, C. R. (2020). Life-history genomic regions explain

differences in Atlantic salmon marine diet specialization. *The Journal of Animal Ecology*, *89*(11), 2677–2691. https://doi.org/10.1111/1365-2656.13324

Backström, T., Pettersson, A., Johansson, V., & Winberg, S. (2011). CRF and urotensin I effects on aggression and anxiety-like behavior in rainbow trout. *Journal of Experimental Biology*, *214*(6), 907–914. https://doi.org/10.1242/jeb.045070

Bangura, P. B., Tiira, K., Niemelä, P. T., Erkinaro, J., Liljeström, P., Toikkanen, A., & Primmer, C. R. (2022). Linking *vgll3* genotype and aggressive behaviour in juvenile Atlantic salmon (*Salmo salar*). *Journal of Fish Biology*, *100*(5), 1264–1271. https://doi.org/10.1111/jfb.15040

Barson, N. J., Aykanat, T., Hindar, K., Baranski, M., Bolstad, G. H., Fiske, P., Jacq, C., Jensen, A. J., Johnston, S. E., Karlsson, S., Kent, M., Moen, T., Niemelä, E., Nome, T., Næsje, T. F., Orell, P., Romakkaniemi, A., Sægrov, H., Urdal, K., … Primmer, C. R. (2015). Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. *Nature*, *528*(7582), Article 7582. https://doi.org/10.1038/nature16062

Barton, B. A., & Iwama, G. K. (1991). Physiological changes in fish from stress in aquaculture with emphasis on the response and effects of corticosteroids. *Annual Review of Fish Diseases*, *1*, 3–26. https://doi.org/10.1016/0959-8030(91)90019-G

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *67*, 1–48. https://doi.org/10.18637/jss.v067.i01

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, *57*(1), 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Bernier, N. J. (2006). The corticotropin-releasing factor system as a mediator of the appetite-suppressing effects of stress in fish. *General and Comparative Endocrinology*, *146*(1), 45–55. https://doi.org/10.1016/j.ygcen.2005.11.016

Birk, S., Chapman, D., Carvalho, L., Spears, B. M., Andersen, H. E., Argillier, C., Auer, S., Baattrup-Pedersen, A., Banin, L., Beklioğlu, M., Bondar-Kunze, E., Borja, A., Branco, P., Bucak, T., Buijse, A. D., Cardoso, A. C., Couture, R.-M., Cremona, F., de Zwart, D., … Hering, D. (2020). Impacts of multiple stressors on freshwater biota across spatial scales and ecosystems. *Nature Ecology & Evolution*, *4*(8), 1060–1068. https://doi.org/10.1038/s41559-020-1216-4

Blanchet, S., Dodson, J. J., & Brosse, S. (2006). Influence of habitat structure and fish density on Atlantic salmon *Salmo salar* L. territorial behaviour. *Journal of Fish Biology*, *68*(3), 951–957. https://doi.org/10.1111/j.0022-1112.2006.00970.x

Cerdá-Reverter, J. M., & Peter, R. E. (2003). Endogenous melanocortin antagonist in fish: Structure, brain mapping, and regulation by fasting of the goldfish agouti-related protein gene. *Endocrinology*, *144*(10), 4552–4561. https://doi.org/10.1210/en.2003-0453

Chen, C.-C., & Fernald, R. D. (2008). Sequences, expression patterns and regulation of the corticotropin-releasing factor system in a teleost. *General and Comparative Endocrinology*, *157*(2), 148–155. https://doi.org/10.1016/j.ygcen.2008.04.003

Conde-Sieira, M., Agulleiro, M. J., Aguilar, A. J., Míguez, J. M., Cerdá-Reverter, J. M., & Soengas, J. L. (2010). Effect of different glycaemic conditions on gene expression of neuropeptides involved in control of food intake in rainbow trout; interaction with stress. *The Journal of Experimental Biology*, *213*(Pt 22), 3858–3865. https://doi.org/10.1242/jeb.048439

Cortés, R., Teles, M., Oliveira, M., Fierro-Castro, C., Tort, L., & Cerdá-Reverter, J. M. (2018). Effects of acute handling stress on short-term central expression of orexigenic/anorexigenic genes in

zebrafish. *Fish Physiology and Biochemistry*, *44*(1), 257–272.
https://doi.org/10.1007/s10695-017-0431-7

Debes, P. V., Piavchenko, N., Ruokolainen, A., Ovaskainen, O., Moustakas-Verho, J. E., Parre, N.,
Aykanat, T., Erkinaro, J., & Primmer, C. R. (2021). Polygenic and major-locus contributions to
sexual maturation timing in Atlantic salmon. *Molecular Ecology*, *30*(18), 4505–4519.
https://doi.org/10.1111/mec.16062

Denver, R. J. (2009). Structural and Functional Evolution of Vertebrate Neuroendocrine Stress
Systems. *Annals of the New York Academy of Sciences*, *1163*(1), 1–16.
https://doi.org/10.1111/j.1749-6632.2009.04433.x

Dysin, A. P., Shcherbakov, Y. S., Nikolaeva, O. A., Terletskii, V. P., Tyshchenko, V. I., & Dementieva, N. V.
(2022). Salmonidae Genome: Features, Evolutionary and Phylogenetic Characteristics. *Genes*,
13(12), 2221. https://doi.org/10.3390/genes13122221

Engel, M., Eggert, C., Kaplick, P. M., Eder, M., Röh, S., Tietze, L., Namendorf, C., Arloth, J., Weber, P.,
Rex-Haffner, M., Geula, S., Jakovcevski, M., Hanna, J. H., Leshkowitz, D., Uhr, M., Wotjak, C. T.,
Schmidt, M. V., Deussing, J. M., Binder, E. B., & Chen, A. (2018). The Role of m6A/m-RNA
Methylation in Stress Response Regulation. *Neuron*, *99*(2), 389-403.e9.
https://doi.org/10.1016/j.neuron.2018.07.009

Finstad, A. G., Einum, S., Ugedal, O., & Forseth, T. (2009). Spatial distribution of limited resources and
local density regulation in juvenile Atlantic salmon. *Journal of Animal Ecology*, *78*(1), 226–
235. https://doi.org/10.1111/j.1365-2656.2008.01476.x

Force, A., Lynch, M., Pickett, F. B., Amores, A., Yan, Y., & Postlethwait, J. (1999). Preservation of
Duplicate Genes by Complementary, Degenerative Mutations. *Genetics*, *151*(4), 1531–1545.
https://doi.org/10.1093/genetics/151.4.1531

Frapin, M., Guignard, S., Meistermann, D., Grit, I., Moullé, V. S., Paillé, V., Parnet, P., & Amarger, V.
(2020). Maternal Protein Restriction in Rats Alters the Expression of Genes Involved in
Mitochondrial Metabolism and Epitranscriptomics in Fetal Hypothalamus. *Nutrients*, *12*(5),
Article 5. https://doi.org/10.3390/nu12051464

Gabanyi, I., Lepousez, G., Wheeler, R., Vieites-Prado, A., Nissant, A., Chevalier, G., Wagner, S.,
Moigneu, C., Dulauroy, S., Hicham, S., Polomack, B., Verny, F., Rosenstiel, P., Renier, N.,
Boneca, I. G., Eberl, G., & Lledo, P.-M. (2022). Bacterial sensing via neuronal Nod2 regulates
appetite and body temperature. *Science (New York, N.Y.)*, *376*(6590), eabj3986.
https://doi.org/10.1126/science.abj3986

Gerisch, B., & Antebi, A. (2011). Molecular basis of life history regulation in C. elegans and other
organisms. In T. Flatt & A. Heyland (Eds.), *Mechanisms of Life History Evolution: The Genetics
and Physiology of Life History Traits and Trade-Offs* (p. 0). Oxford University Press.
https://doi.org/10.1093/acprof:oso/9780199568765.003.0022

Ghisleni, G., Capiotti, K. M., Da Silva, R. S., Oses, J. P., Piato, Â. L., Soares, V., Bogo, M. R., & Bonan, C.
D. (2012). The role of CRH in behavioral responses to acute restraint stress in zebrafish.
Progress in Neuro-Psychopharmacology & Biological Psychiatry, *36*(1), 176–182.
https://doi.org/10.1016/j.pnpbp.2011.08.016

Gorissen, M., & Flik, G. (2016). 3 - The Endocrinology of the Stress Response in Fish: An Adaptation-
Physiological View. In C. B. Schreck, L. Tort, A. P. Farrell, & C. J. Brauner (Eds.), *Fish Physiology*
(Vol. 35, pp. 75–111). Academic Press. https://doi.org/10.1016/B978-0-12-802728-8.00003-5

Grone, B. P., & Maruska, K. P. (2015). Divergent evolution of two corticotropin-releasing hormone (CRH) genes in teleost fishes. *Frontiers in Neuroscience*, *9*. https://doi.org/10.3389/fnins.2015.00365

Grossman, G. D., & Simon, T. N. (2020). Density-dependent effects on salmonid populations: A review. *Ecology of Freshwater Fish*, *29*(3), 400–418. https://doi.org/10.1111/eff.12523

Hartig, F., Lohse, L., & leite, M. de S. (2024). *DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models* (Version 0.4.7) [Computer software]. https://cran.r-project.org/web/packages/DHARMa/index.html

He, M., Li, D., Fang, C., & Xu, Q. (2022). YTHDF1 regulates endoplasmic reticulum stress, NF-κB, MAPK and PI3K-AKT signaling pathways in inflammatory osteoclastogenesis. *Archives of Biochemistry and Biophysics*, *732*, 109464. https://doi.org/10.1016/j.abb.2022.109464

Hedrick, P. W. (2012). What is the evidence for heterozygote advantage selection? *Trends in Ecology & Evolution*, *27*(12), 698–704. https://doi.org/10.1016/j.tree.2012.08.012

Hodgson, E. E., Essington, T. E., & Halpern, B. S. (2017). Density dependence governs when population responses to multiple stressors are magnified or mitigated. *Ecology*, *98*(10), 2673–2683. https://doi.org/10.1002/ecy.1961

Höjesjö, J., Johnsson, J., & Bohlin, T. (2004). Habitat Complexity Reduces the Growth of Aggressive and Dominant Brown Trout (Salmo trutta) Relative to Subordinates. *Behavioral Ecology and Sociobiology*, *56*(3), 286–289.

Hostetler, C. M., & Ryabinin, A. E. (2013). The CRF system and social behavior: A review. *Frontiers in Neuroscience*, *7*. https://doi.org/10.3389/fnins.2013.00092

Huang, P., Liu, M., Zhang, J., Zhong, X., & Zhong, C. (2023). YTHDF1 Attenuates TBI-Induced Brain-Gut Axis Dysfunction in Mice. *International Journal of Molecular Sciences*, *24*(4), Article 4. https://doi.org/10.3390/ijms24044240

Jonsson, N., Jonsson, B., & Hansen, L. P. (1998). The relative role of density-dependent and density-independent survival in the life cycle of Atlantic salmon Salmo salar. *Journal of Animal Ecology*, *67*(5), 751–762. https://doi.org/10.1046/j.1365-2656.1998.00237.x

Kalananthan, T., Folkedal, O., Gomes, A. S., Lai, F., Handeland, S. O., Tolås, I., Gelebart, V., & Rønnestad, I. (2023). Impact of long-term fasting on the stomach-hypothalamus appetite regulating genes in Atlantic salmon postsmolts. *Aquaculture*, *563*, 738917. https://doi.org/10.1016/j.aquaculture.2022.738917

Kalananthan, T., Gomes, A. S., Lai, F., Tolås, I., Jordal, A.-E. O., Norland, S., Murashita, K., & Rønnestad, I. (2021). Brain Distribution of 10 cart Transcripts and Their Response to 4 Days of Fasting in Atlantic Salmon (Salmo salar L.). *Frontiers in Marine Science*, *8*. https://doi.org/10.3389/fmars.2021.763766

Kalananthan, T., Murashita, K., Rønnestad, I., Ishigaki, M., Takahashi, K., Silva, M. S., Wakabayashi, Y., Lai, F., Shimizu, M., Nilsen, T. O., Pino Martinez, E., & Gomes, A. S. (2020). Hypothalamic agrp and pomc mRNA Responses to Gastrointestinal Fullness and Fasting in Atlantic Salmon (Salmo salar, L.). *Frontiers in Physiology*, *11*, 61. https://doi.org/10.3389/fphys.2020.00061

Kassahn, K. S., Dang, V. T., Wilkins, S. J., Perkins, A. C., & Ragan, M. A. (2009). Evolution of gene function and regulatory control after whole-genome duplication: Comparative analyses in vertebrates. *Genome Research*, *19*(8), 1404–1418. https://doi.org/10.1101/gr.086827.108

Lai, F., Royan, M. R., Gomes, A. S., Espe, M., Aksnes, A., Norberg, B., Gelebart, V., & Rønnestad, I. (2021). The stress response in Atlantic salmon (*Salmo salar* L.): Identification and functional

characterization of the corticotropin-releasing factor (*crf*) paralogs. *General and Comparative*
Endocrinology, *313*, 113894. https://doi.org/10.1016/j.ygcen.2021.113894

Lenth, R. V., Banfai, B., Bolker, B., Buerkner, P., Giné-Vázquez, I., Herve, M., Jung, M., Love, J., Miguez,
F., Piaskowski, J., Riebl, H., & Singmann, H. (2024). *emmeans: Estimated Marginal Means, aka*
Least-Squares Means (Version 1.10.5) [Computer software]. https://cran.r-
project.org/web/packages/emmeans/index.html

Li, A., Chen, Y.-S., Ping, X.-L., Yang, X., Xiao, W., Yang, Y., Sun, H.-Y., Zhu, Q., Baidya, P., Wang, X.,
Bhattarai, D. P., Zhao, Y.-L., Sun, B.-F., & Yang, Y.-G. (2017). Cytoplasmic m6A reader YTHDF3
promotes mRNA translation. *Cell Research*, *27*(3), 444–447.
https://doi.org/10.1038/cr.2017.10

Liao, S., Sun, H., & Xu, C. (2018). YTH Domain: A Family of N6-methyladenosine (m6A) Readers.
Genomics, Proteomics & Bioinformatics, *16*(2), 99–107.
https://doi.org/10.1016/j.gpb.2018.04.002

Lien, S., Koop, B. F., Sandve, S. R., Miller, J. R., Kent, M. P., Nome, T., Hvidsten, T. R., Leong, J. S.,
Minkley, D. R., Zimin, A., Grammes, F., Grove, H., Gjuvsland, A., Walenz, B., Hermansen, R. A.,
von Schalburg, K., Rondeau, E. B., Di Genova, A., Samy, J. K. A., … Davidson, W. S. (2016). The
Atlantic salmon genome provides insights into rediploidization. *Nature*, *533*(7602), 200–205.
https://doi.org/10.1038/nature17164

Liu, B., Liu, Y., & Wang, X. (2015). The effect of stocking density on growth and seven physiological
parameters with assessment of their potential as stress response indicators for the Atlantic
salmon (Salmo salar). *Marine and Freshwater Behaviour and Physiology*, *48*(3), 177–192.
https://doi.org/10.1080/10236244.2015.1034956

Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods (San Diego, Calif.)*, *25*(4),
402–408. https://doi.org/10.1006/meth.2001.1262

Lu, Z., Ma, Y., Li, Q., Liu, E., Jin, M., Zhang, L., & Wei, C. (2019). The role of N6-methyladenosine RNA
methylation in the heat stress response of sheep (*Ovis aries*). *Cell Stress and Chaperones*,
24(2), 333–342. https://doi.org/10.1007/s12192-018-00965-x

Matte, J.-M., Fraser, D. J., & Grant, J. W. A. (2020). Density-dependent growth and survival in
salmonids: Quantifying biological mechanisms and methodological biases. *Fish and Fisheries*,
21(3), 588–600. https://doi.org/10.1111/faf.12448

Nislow, K. H., Armstrong, J. D., & Grant, J. W. A. (2010). The Role of Competition in the Ecology of
Juvenile Atlantic Salmon. In *Atlantic Salmon Ecology* (pp. 171–197). John Wiley & Sons, Ltd.
https://doi.org/10.1002/9781444327755.ch7

Parker, J. A., & Bloom, S. R. (2012). Hypothalamic neuropeptides and the regulation of appetite.
Neuropharmacology, *63*(1), 18–30. https://doi.org/10.1016/j.neuropharm.2012.02.004

Perlegos, A. E., Shields, E. J., Shen, H., Liu, K. F., & Bonini, N. M. (2022). Mettl3-dependent m6A
modification attenuates the brain stress response in Drosophila. *Nature Communications*,
13(1), 5387. https://doi.org/10.1038/s41467-022-33085-3

Prokkola, J. M., Åsheim, E. R., Morozov, S., Bangura, P., Erkinaro, J., Ruokolainen, A., Primmer, C. R., &
Aykanat, T. (2022). Genetic coupling of life-history and aerobic performance in Atlantic
salmon. *Proceedings. Biological Sciences*, *289*(1967), 20212500.
https://doi.org/10.1098/rspb.2021.2500

Reznick, D. (2016). Hard and Soft Selection Revisited: How Evolution by Natural Selection Works in
the Real World. *Journal of Heredity*, *107*(1), 3–14. https://doi.org/10.1093/jhered/esv076

Rønnestad, I., Gomes, A. S., Murashita, K., Angotzi, R., Jönsson, E., & Volkoff, H. (2017). Appetite-
Controlling Endocrine Systems in Teleosts. *Frontiers in Endocrinology*, *8*, 73.
https://doi.org/10.3389/fendo.2017.00073

Roy, B., Ochi, S., & Dwivedi, Y. (2022). M6A RNA Methylation-Based Epitranscriptomic Modifications
in Plasticity-Related Genes via miR-124-C/EBPα-FTO-Transcriptional Axis in the Hippocampus
of Learned Helplessness Rats. *The International Journal of Neuropsychopharmacology*,
25(12), 1037–1049. https://doi.org/10.1093/ijnp/pyac068

Schneeberger, M., Gomis, R., & Claret, M. (2014). Hypothalamic and brainstem neuronal circuits
controlling homeostatic energy balance. *Journal of Endocrinology*, *220*(2), T25–T46.
https://doi.org/10.1530/JOE-13-0398

Schwartz, M. W., Woods, S. C., Porte, D., Seeley, R. J., & Baskin, D. G. (2000). Central nervous system
control of food intake. *Nature*, *404*(6778), 661–671. https://doi.org/10.1038/35007534

Shi, H., Xu, Y., Tian, N., Yang, M., & Liang, F.-S. (2022). Inducible and reversible RNA N6-
methyladenosine editing. *Nature Communications*, *13*(1), 1958.
https://doi.org/10.1038/s41467-022-29665-y

Sinclair-Waters, M., Ødegård, J., Korsvoll, S. A., Moen, T., Lien, S., Primmer, C. R., & Barson, N. J.
(2020). Beyond large-effect loci: Large-scale GWAS reveals a mixed large-effect and polygenic
architecture for age at maturity of Atlantic salmon. *Genetics Selection Evolution*, *52*(1), 9.
https://doi.org/10.1186/s12711-020-0529-8

Sinclair-Waters, M., Piavchenko, N., Ruokolainen, A., Aykanat, T., Erkinaro, J., & Primmer, C. R. (2022).
Refining the genomic location of single nucleotide polymorphism variation affecting Atlantic
salmon maturation timing at a key large-effect locus. *Molecular Ecology*, *31*(2), 562–570.
https://doi.org/10.1111/mec.16256

Sisk, C. L., & Foster, D. L. (2004). The neural basis of puberty and adolescence. *Nature Neuroscience*,
7(10), 1040–1047. https://doi.org/10.1038/nn1326

Smith, S. M., & Vale, W. W. (2006). The role of the hypothalamic-pituitary-adrenal axis in
neuroendocrine responses to stress. *Dialogues in Clinical Neuroscience*, *8*(4), 383–395.
https://doi.org/10.31887/DCNS.2006.8.4/ssmith

Song, N., Du, J., Gao, Y., & Yang, S. (2020). Epitranscriptome of the ventral tegmental area in a deep
brain-stimulated chronic unpredictable mild stress mouse model. *Translational Neuroscience*,
11(1), 402–418. https://doi.org/10.1515/tnsci-2020-0146

Stearns, S. C. (2011). Does impressive progress on understanding mechanisms advance life history
theory? In T. Flatt & A. Heyland (Eds.), *Mechanisms of Life History Evolution: The Genetics*
and Physiology of Life History Traits and Trade-Offs (p. 0). Oxford University Press.
https://doi.org/10.1093/acprof:oso/9780199568765.003.0027

Stepan, J., Anderzhanova, E., & Gassen, N. C. (2018). Hippo Signaling: Emerging Pathway in Stress-
Related Psychiatric Disorders? *Frontiers in Psychiatry*, *9*.
https://doi.org/10.3389/fpsyt.2018.00715

Sundh, H., Finne-Fridell, F., Ellis, T., Taranger, G. L., Niklasson, L., Pettersen, E. F., Wergeland, H. I., &
Sundell, K. (2019). Reduced water quality associated with higher stocking density disturbs
the intestinal barrier functions of Atlantic salmon (*Salmo salar* L.). *Aquaculture*, *512*, 734356.
https://doi.org/10.1016/j.aquaculture.2019.734356

Sveen, L. R., Timmerhaus, G., Torgersen, J. S., Ytteborg, E., Jørgensen, S. M., Handeland, S.,
Stefansson, S. O., Nilsen, T. O., Calabrese, S., Ebbesson, L., Terjesen, B. F., & Takle, H. (2016).
Impact of fish density and specific water flow on skin properties in Atlantic salmon (*Salmo*

salar L.) post-smolts. *Aquaculture*, *464*, 629–637.
https://doi.org/10.1016/j.aquaculture.2016.08.012

Takahashi, H., & Sakamoto, T. (2013). The role of 'mineralocorticoids' in teleost fish: Relative importance of glucocorticoid signaling in the osmoregulation and 'central' actions of mineralocorticoid receptor. *General and Comparative Endocrinology*, *181*, 223–228. https://doi.org/10.1016/j.ygcen.2012.11.016

Thorstad, E. B., Whoriskey, F., Rikardsen, A. H., & Aarestrup, K. (2010). Aquatic Nomads: The Life and Migrations of the Atlantic Salmon. In *Atlantic Salmon Ecology* (pp. 1–32). John Wiley & Sons, Ltd. https://doi.org/10.1002/9781444327755.ch1

Volkoff, H. (2016). The Neuroendocrine Regulation of Food Intake in Fish: A Review of Current Knowledge. *Frontiers in Neuroscience*, *10*. https://doi.org/10.3389/fnins.2016.00540

Wang, X., Zhao, B. S., Roundtree, I. A., Lu, Z., Han, D., Ma, H., Weng, X., Chen, K., Shi, H., & He, C. (2015). N(6)-methyladenosine Modulates Messenger RNA Translation Efficiency. *Cell*, *161*(6), 1388–1399. https://doi.org/10.1016/j.cell.2015.05.014

Wang, Y., Chi, L., Liu, Q., Xiao, Y., Ma, D., Xiao, Z., Xu, S., & Li, J. (2019). Effects of stocking density on the growth and immunity of Atlantic salmon salmo salar reared in recirculating aquaculture system (RAS). *Journal of Oceanology and Limnology*, *37*(1), 350–360. https://doi.org/10.1007/s00343-019-7350-7

Wei, R., Yuan, D., Wang, T., Zhou, C., Lin, F., Chen, H., Wu, H., Yang, S., Wang, Y., Liu, J., Gao, Y., & Li, Z. (2013). Characterization, tissue distribution and regulation of agouti-related protein (AgRP) in a cyprinid fish (Schizothorax prenanti). *Gene*, *527*(1), 193–200. https://doi.org/10.1016/j.gene.2013.06.003

Wilkinson, E., Cui, Y.-H., & He, Y.-Y. (2021). Context-Dependent Roles of RNA Modifications in Stress Responses and Diseases. *International Journal of Molecular Sciences*, *22*(4), Article 4. https://doi.org/10.3390/ijms22041949

Wu, P.-F., Han, Q.-Q., Chen, F.-F., Shen, T.-T., Li, Y.-H., Cao, Y., Chen, J.-G., & Wang, F. (2021). Erasing m6A-dependent transcription signature of stress-sensitive genes triggers antidepressant actions. *Neurobiology of Stress*, *15*, 100390. https://doi.org/10.1016/j.ynstr.2021.100390

Wu, T., Shao, Y., Li, X., Wu, T., Yu, L., Liang, J., Zhang, Y., Wang, J., Sun, T., Zhu, Y., Chang, X., Wang, S., Chen, F., & Han, X. (2023). NR3C1/Glucocorticoid receptor activation promotes pancreatic β-cell autophagy overload in response to glucolipotoxicity. *Autophagy*, *19*(9), 2538–2557. https://doi.org/10.1080/15548627.2023.2200625

Wunderink, Y. S., Engels, S., Halm, S., Yúfera, M., Martínez-Rodríguez, G., Flik, G., Klaren, P. H. M., & Mancera, J. M. (2011). Chronic and acute stress responses in Senegalese sole (Solea senegalensis): The involvement of cortisol, CRH and CRH-BP. *General and Comparative Endocrinology*, *171*(2), 203–210. https://doi.org/10.1016/j.ygcen.2011.01.010

Yang, Y., Han, W., Zhang, A., Zhao, M., Cong, W., Jia, Y., Wang, D., & Zhao, R. (2022). Chronic corticosterone disrupts the circadian rhythm of CRH expression and m6A RNA methylation in the chicken hypothalamus. *Journal of Animal Science and Biotechnology*, *13*(1), 29. https://doi.org/10.1186/s40104-022-00677-4

Yu, H., Gao, Q., Wang, W., Liu, D., He, J., & Tian, Y. (2023). Comprehensive Analysis of YTH Domain-Containing Genes, Encoding m6A Reader and Their Response to Temperature Stresses and Yersinia ruckeri Infection in Rainbow Trout (Oncorhynchus mykiss). *International Journal of Molecular Sciences*, *24*(11), Article 11. https://doi.org/10.3390/ijms24119348

Zaccara, S., Ries, R. J., & Jaffrey, S. R. (2019). Reading, writing and erasing mRNA methylation. *Nature*

Reviews. Molecular Cell Biology, *20*(10), 608–624. https://doi.org/10.1038/s41580-019-

0168-5

Zhao, B. S., Roundtree, I. A., & He, C. (2017). Post-transcriptional gene regulation by mRNA

modifications. *Nature Reviews Molecular Cell Biology*, *18*(1), 31–42.

https://doi.org/10.1038/nrm.2016.132