Acute temporal, regional, and cell-type specific NKCC1 disruption following severe TBI in the developing gyrencephalic brain

Authors:

Alexandra Hochstetler[1,2], Ya'el Courtney[2,3], Peace Oloko[1], Benjamin Baskin[1], Andrew Ding-Su [1], Tawny Stinson[1], Declan McGuone[4], Robin Haynes[2], Maria K. Lehtinen[2,3], and Beth Costine-Bartell[1]*

*Corresponding Author, bcostinebartell@mgh.harvard.edu

Affiliations:

[1]Department of Neurosurgery, Massachusetts General Hospital, Harvard Medical School, Boston, MA 02115

[2]Department of Pathology, Boston Children's Hospital and Harvard Medical School, Boston, MA 02115

[3]Graduate Program in Neuroscience, Harvard Medical School, Boston, MA 02115

[4]Department of Pathology, Yale University School of Medicine, New Haven, CT, 06510

Conflict of Interest statement:

The authors declare no competing financial interests.

Acknowledgements:

This work is supported by NIH NICHD K01HD083759 and R01 HD099397 (BCB), by the Pappendick Family Translational Research Program (MKL), the Hydrocephalus Association, and R01 HD090064 (RH). AH was supported by T32NS007473 and F32NS134588 training fellowships from the National Institute for Neurological Disorders and Stroke. YC was supported by the Howard Hughes Medical Institute through the James J. Gilliam Fellowships for Advanced Study program. PO was supported by a Mass General Summer Research Trainee Program (SRTP). We thank the Boston Children's Hospital IDDRC Cellular Imaging Core supported by S10OD016453. We thank the past and present team of undergraduate interns in the Costine-Bartell laboratory who assisted with the surgeries and 30 hours of intensive care to create the brain injury model and tissue collection as well as the Knight Surgical Research Lab for assistance with anesthesia to create the brain injuries. We are especially grateful for the fetal piglet tissue obtained from Dr. Ryann Fame (Stanford University Department of Neurosurgery), Dr. Mercedes Paredes (University of California, San Francisco (UCSF) Department of Neurology), and Dr. Misty Williams-Fritze (CBSET) and postnatal piglet tissue obtained from Dr. Corinna Beale (Center for Comparative Medicine, MGH). We thank Dr. Cameron Sadegh (UC Davis Department of Neurosurgery) and members of the Lehtinen Laboratory for assistance with prenatal swine brain dissections. We are deeply grateful to participating families for their courage and generosity in consenting to the use of their children in research. We also thank the clinicians, medical examiners, and pathologists who helped in the accrual of cases, including Drs. Gary Dale, Jeffrey Goldstein, Steven Campman, Catherine Stoos, and Colin Smith. We thank Elisabeth Haas and Richard Goldstein for the processing and analysis of infant cases. We thank Jane's Trust Initiative, the NIH NeuroBioBank, and the Edinburgh Brain Bank.

Author contributions:

AH: Conceptualization, investigation, methodology, formal analysis, writing-original draft, visualization; YC: Investigation, methodology, formal analysis, writing-original draft; PO: Investigation, methodology, formal analysis, visualization; BB: Investigation, methodology, formal analysis, visualization; AD: Investigation, methodology; DM: Investigation, formal analysis, writing-revising; TS: Investigation, methodology; RH: Investigation, methodology, formal analysis, writing-revising, funding acquisition; MKL: Writing-revising, funding acquisition; BCB: Conceptualization, investigation, methodology, visualization, writing-original draft, project administration, funding acquisition

Abstract: Traumatic brain injury (TBI) in children is a leading cause of morbidity and mortality, with no effective treatment and limited clinical management. We developed a multifactorial traumatic brain injury model in piglets which mirrors the evolving pathophysiology of severe pediatric TBI, showing age-dependent hypoxic-ischemic cerebral cortical injury and matrix metalloproteinase-driven vasogenic edema, with infant piglets experiencing less tissue damage than toddler piglets. Extracellular matrix breakdown can precipitate neuronal dysfunction, disrupting chloride homeostasis and the reversal potential for GABA. We hypothesized that ongoing tissue damage might be related to markers of "immature GABA", evaluated by changes to the expression and phosphorylation of sodium-potassium-2-chloride cotransporter 1 (NKCC1), potassium-chloride cotransporter 2 (KCC2), and a regulatory kinase, (STE20/SPS1-related proline-alanine-rich protein kinase) SPAK. We mapped these markers in developing swine and infant human brain, identifying a postnatal pNKCC1 decrease in human infant hippocampus, and a perinatal cortical and hippocampal GABA switch in pigs, with no change in the thalamus. In infant piglets with severe TBI, upregulation of neuronal pNKCC1 correlated with hypoxic-ischemic injury and seizure duration. We also observed dysregulation of NKCC1, KCC2, and SPAK in cortex and hippocampus in infant and toddler piglets with severe TBI, with thalamus unchanged. We noted ectopic, non-apical localization of pNKCC1 signal in choroid plexus epithelium across ages in piglets and humans with severe TBI, indicating acute dysregulation of the CSF chloride milieu. These findings position swine as a useful model for pediatric TBI research and suggest that SPAK or NKCC1 inhibition in infants may be therapeutic.

Significance statement: Severe TBI in early childhood, the majority of which is due to abuse, remains an understudied area of neurotrauma. Our piglet model effectively replicates the pathophysiology of severe pediatric TBI, capturing age-dependent injury patterns and mechanisms of spreading hypoxic-ischemic injury throughout the cortical ribbon. We identified upregulation of neuronal-pNKCC1 in infant piglets, but not toddler piglets with severe TBI, and found this correlates with injury severity, seizure duration, and subarachnoid hemorrhage. Our findings indicate that treatments targeted to inhibit neuronal NKCC1 might alleviate evolving brain injury in infants with severe TBI. This model provides a valuable platform for studying mechanisms of TBI and testing new interventions, potentially advancing therapeutic strategies for pediatric brain injury where stopping traumatic seizures is difficult.

INTRODUCTION:

In childhood, severe traumatic brain injury (TBI) commonly arises from abusive head trauma (AHT), causing high mortality and persistent neurological comorbidities (Keenan et al., 2003). Children of different ages show distinct structural, biomechanical, and physiological vulnerabilities, influencing both injury severity and long-term outcomes. Animal models which enable scaling of injuries to brain volume can help us better understand and treat TBI in pediatric populations.

Swine models are especially valuable as models of age-dependent traumatic brain injury and neurological comorbidities (Clancy et al., 2007; Lind, 2007; Swindle, 1998; Dobbing and Sands 1979). For example, a standardized focal cortical impact produces a larger contusion in piglets approximating human toddlers than in those approximating human infants. Conversely, younger piglets are more prone to diffuse injuries from non-impact rotational forces, similar to clinical observations (Missios et al., 2009; Ibrahim et al., 2010).

We developed a multifactorial brain injury model in piglets which recapitulates a "hemispheric hypodensity" (HH) pattern akin to that observed in pediatric TBI patients, where infant piglets show bilateral HH and toddler piglets show unilateral patterns (Costine-Bartell et al., 2019; Duhaime and Durham 2007; Foster et al., 2014). We have found that subarachnoid hemorrhage with at least 60 minutes of seizure are key drivers of age-dependent hypoxic-ischemic type injury and vasogenic edema (Costine-Bartell et al., 2021; Hochstetler, Pret al., 2024). Maintaining proper ion homeostasis in the developing brain is crucial for neurodevelopment and may be disrupted in TBI. A critical factor in these age-dependent differences may be the "GABA switch"—the developmental transition of GABAergic signaling from primarily excitatory to inhibitory as the brain matures, governed by an increase in the expression and membrane phosphorylation/activity of KCC2 over NKCC1, chloride cotransporters which control neuronal $[Cl^-]_i$ (Ben-Ari 2014; LoTurco et al., 1995). TBI might disrupt this balance, thereby exacerbating injury and its consequences (Bortone and Polleux, 2009).

Anti-seizure medications (GABA agonists) are the standard of care for children with severe TBI to halt traumatic seizures (Smith et al., 2016). In cases of high extracellular chloride such as in early development or injury, GABA agonists may potentiate neuronal excitability and ongoing tissue damage.

The timing of the developmental "GABA switch" from primarily excitatory to inhibitory varies widely across

species, but always depends on the relative abundance and activity of two cation chloride cotransporters,

NKCC1 (sodium-potassium-2-chloride cotransporter) and KCC2 (potassium-chloride cotransporter) which work

in antagonism to control neuronal $[Cl]_i$. In precocial mammals like sheep and guinea pigs, the shift occurs largely

before birth, whereas in altricial mammals such as rabbits and rodents, or hatched species such as birds, turtles,

and frogs, the switch takes place predominantly after birth (Ben-Ari et al., 2007; Ben-Ari 2014; Sedmak et al.,

2016; Tang et al., 2020; Peerboom and Wierenga, 2021). Humans and non-human primates appear to follow a

more perinatal pattern (Khazipov et al., 2001; Sedmak et al., 2016). Swine are considered sometimes precocial,

and sometimes semi-altricial, thus, pinpointing when the GABA switch occurs is critical for understanding their

suitability as a model for pediatric brain injury and for testing age-appropriate anticonvulsant therapies (Young

et al., 2023; Bienboire-Frosini et al., 2023). The timing of the GABA switch is also important when considering

other diagnoses where seizure activity may play a role in pathogenesis (ex. Sudden Unexpected Death in

Childhood) (Goldstein and Poduri 2024; Leitner et al., 2022; Harowitz et al., 2021; Haynes et al., 2021; Patodia

et al., 2021).

In this study, we characterized the developmental expression and phosphorylation of NKCC1 and KCC2

in the neonatal human hippocampus and the developing piglet brain and examined how severe TBI disrupts this

balance. We hypothesized that developmental differences in NKCC1 expression may contribute to distinct TBI

injury patterns in infant and toddler piglets. To address this hypothesis, we employ spatial transcriptomics,

immunohistochemistry, and Western blot analyses across developmental stages, brain regions, and TBI

conditions. We also examined human TBI autopsy samples from various ages, assessing pNKCC1 expression

in different brain compartments and cell types.

MATERIALS AND METHODS:

Sample procurement

Animal procedures

All protocols and procedures performed were in accordance with the guidelines of the American Veterinary

Association, the National Institutes of Health, and the United States Department of Agriculture. They were

approved by the Institutional Animal Care and Use Committee at Massachusetts General Hospital.

Naïve swine tissue collection

Prenatal collections

Preterm piglets were collected from euthanized Yorkshire sows at approximately 30% gestation (ED36), 75%

gestation (ED90), and 90% gestation (ED105). Pig fetuses were surgically removed from the amniotic sac

following maternal euthanasia, rinsed under cold running water, brains were dissected on ice, and tissues were

immediately flash frozen. In all cases, the time from euthanasia to sample collection was less than 4 hours.

Postnatal collections

7 day old (PND7, "infant") and 30 day old (PND30, "toddler") male Yorkshire piglets were selected for naïve

tissue collection. Surgery and anesthesia protocols were employed as previously described (BCB et al., 2019a)

adjusted to the age of the piglets. In brief, piglets were sedated with ketamine/xylazine/atropine and anesthetized

by 1-1.5% isoflurane with medical oxygen. Piglets were intubated and mechanically ventilated with room air. An

ear intravenous (IV), or external jugular vein catheter was placed to access venous blood and deliver saline

infusion. In all cases, mean arterial blood pressure, oxygen saturation, heart rate and end tidal CO_2, respiratory

rate, and core body temperature were managed for the duration of the procedures. A craniotomy was performed,

and the dura was removed. Euthasol was delivered intravenously, and then fresh tissue dissection was

performed immediately on wet ice. Samples (n=4/age) were collected from August 2023 to October 2023 and

flash-frozen on dry ice.

Pediatric severe TBI swine tissue collection

Samples were requisitioned as freezer aliquots from prior studies but are distinct aliquots from the same piglets

published in Hochstetler, Price *et al.* 2024. In brief, a series of PND7 and PND30 male Yorkshire piglets from

September 2016 to October 2019 were exposed to surgery and anesthesia protocols as detailed above in

"postnatal collections". *Immediately prior to injury*, piglets were switched to a seizure-permissive anesthesia

protocol and subjected to a combination of injuries scaled to brain volume as previously described (Costine-

Bartell et al., 2019; Costine-Bartell et al., 2021) including cortical impact, mass effect, subdural

hematoma/subarachnoid hemorrhage supplemented with kainic acid, one minute of apnea, and 10 minutes of

hypoventilation. In the sham piglets, only the burr holes and duration of anesthesia were modeled. Piglets were

managed in an intensive care monitoring unit for 24 hours after injury. For fresh frozen tissue, euthasol was

delivered intravenously, and then fresh tissue dissection followed immediately on wet ice, and samples were

flash-frozen on dry ice. For perfused-fixed tissue, piglets were transcardially perfused with room temperature

normal saline at a rate of ~1L/min until complete surface vessel clearing followed by room temperature 10%

neutral buffered formalin at the same rate until total body stiffness. Brains were obtained and post-fixed in 10%

neutral buffered formalin for 3-5 days until appropriate tissue stiffness was achieved.

Human tissue collection

Lysates from frozen hippocampus at the level of the lateral geniculate nucleus were collected from either the

NIH NeuroBioBank or the San Diego County Medical Examiner's office, the latter available for research under

the auspice of the California Code, Section 27491.41. All infants died suddenly and unexpectedly with the cause

of death determined at autopsy and adjudicated within the laboratory for research purposes (Table 1). Unstained

tissue sections from FFPE blocks were procured from autopsy brain samples (TBI or non-TBI control) performed

at the Neuropathology Department at the University of Edinburgh. Permission to use the brain tissue was given

according to the guidelines required by the respective ethics committees of these institutions: the Boston

Children's Hospital and Massachusetts General Hospital Institutional Review Boards. Demographic information

of all human samples is detailed in Table 1.

Standard laboratory techniques

Western Blotting

Fresh frozen samples were manually homogenized in N-PER lysis buffer (ThermoScientific #87792)

supplemented with 1X Halt (ThermoScientific #78440) protease and phosphatase inhibitor cocktail and lysed at

4°C for 60 minutes with end-over-end agitation. Lysates were spun at 16,000 x g for 10 minutes to pellet debris.

Supernatants were aliquoted and Pierce BCA assay was used protein concentration (ThermoScientific #23250).

20µg of the sample was reduced with 5X Laemmli Sample Buffer (BioLegend #426311) supplemented with B-

mercaptoethanol (5%) and diluted with double distilled water to an equivalent volume. Samples were heated at

70°C for 5 minutes, loaded into 4-12% Bis-Tris gels (NuPage #NP0321) and separated at 140V for 65 minutes.

Gels were transferred to 0.2 µm nitrocellulose membranes at 1.0A for 30 minutes using the Bio-Rad Trans-

Blotter® system. Membranes were stained with Ponceau-S total protein stain blocked with 5% milk dissolved in

1X Tris Buffered Saline-0.1% Tween-20 (TBST) with gentle agitation. Membranes were incubated with primary

antibody diluted in 5% milk-TBST, washed with TBST, and then incubated with secondary antibody diluted in

5% milk-TBST. Membranes were washed with TBST and TBS, exposed with chemiluminescent reagents

(Cytiva-Amersham #RPN2235), and visualized on a Bio-Rad Chemi-Doc Imager. For antibody validation, full

complete membranes were first probed with each antibody to evaluate non-specific bands, specific bands and

dimers, and to optimize dilutions. In the case of pNKCC1 and NKCC1 antibodies, our laboratory has performed

in-house knock-down validation. All validation methods are presented in Table 2. For publication gels, technical

duplicate membranes were cut at 80kDa to allow probing of 1) pSPAK, 2) SPAK, and 3) beta-actin (below 80kDa)

simultaneously with 1) pNKCC1, 2) NKCC1, and 1) pKCC2, 2) KCC2 (above 80kDa). In all cases, band

intensities were normalized to total protein loaded (Ponceau stain) for the graphs, since beta-actin and beta-

tubulin were found to be poor loading controls across age and region. Membrane stripping was performed with

ReStore™ Western Blot Stripping Solution and protocol (ThermoScientific #21059).

Quantitative, Real-Time PCR

Samples were processed for RNA extraction using the Total RNA Miniprep Kit (New England BioLabs #T2010),

evaluated for concentration >50ng/µL and a quality ratio of A260/280 of >2.0. RNA was prepared for cDNA

synthesis with +reverse transcriptase and –reverse transcriptase reactions using the LunaScript® RT Supermix

Kit (New England Biolabs #E3010) using quality control metrics of concentration >100ng/µL and A260/280 ratio

>1.8. Samples were diluted to equivalent cDNA loading and probed with Taqman™ reagents for quantitative

gene expression using fast run chemistry on an Applied Biosystems machine. In all cases, -RT and no template

controls were run for each sample. Probe information is detailed in Table 3.

Spatial Transcriptomics

Spatial transcriptomics was conducted on anatomically matched formalin-fixed, paraffin-embedded (FFPE) brain

sections from a sham 30-day-old piglet and a 30-day-old piglet with severe traumatic brain injuries using

NanoString GeoMx® Digital Spatial Profiling technology, facilitated by the Comparative Pathology and Genomics

Shared Resource core facility at The Cummings School of Veterinary Medicine at Tufts University. Regions of

interest (ROIs) were selected to capture key anatomical areas relevant to age-dependent TBI pathology and

neuroinflammation. Following ROI selection, samples underwent GeoMx® platform analysis, where regions were

scanned, and digital counts were obtained for gene targets of interest. Signal thresholding at a 3% cutoff was

applied to exclude low-expression signals, ensuring that only significant transcript data were included in

downstream analysis. Normalization was performed using internal controls (External RNA Control Consortium)

provided by the GeoMx® platform to account for technical variation across slides and samples. Upper quartile

normalization was performed after filtering the dataset to remove targets below the limit of quantitation.

Histology

Following immersion fixation, brains were washed with 1X phosphate-buffered saline (PBS), paraffin embedded

and sectioned at 5-μm thickness and mounted onto silanized glass slides. Immunofluorescence was performed

on these sections using antibodies detailed in Table 2. Briefly, sections were deparaffinized in xylenes,

rehydrated, and then subjected to antigen retrieval in boiling 10mM citric acid-Tween pH 6.0 for 20 minutes.

Staining protocols are presented in more detail below. In all cases, negative control sections were incubated in

corresponding species-specific IgG at an equivalent concentration to the primary antibody dilution during the

primary antibody incubation step. For figures where quantitation is presented, all slides evaluated were stained

together in a single batch to minimize signal variability. Slides were imaged on a Zeiss Axio Observer D1 inverted

microscope (fluorescent, 20x air objective), an Olympus BX51 light microscope (chromogenic, 10x or 20x air

objective), or a Zeiss LSM980 (20x air or 63x oil objective) at the Boston Children's Hospital Cellular Imaging

Core. Images were processed and assembled using ZEN Blue, ImageJ software, and Adobe Illustrator.

Chromogenic Immunohistochemistry

For chromogenic stains, a peroxidase quenching step was performed immediately prior to antigen retrieval.

Sections were then permeabilized for 5 minutes with 1% PBS-Tween-20 and blocked for 1 hour using normal

horse serum diluted in Super Sensitive Wash Buffer (BioGenex, HK583-5KEGP). Sections were incubated in

primary antibodies diluted in blocking buffer for 2 hours at room temperature. Slides were washed with 1% PBS-

Tween-20 and then incubated in the secondary antibody diluted in blocking buffer for 30 minutes. Sections were

washed and subjected to antibody visualization with Vectastain® ABC and DAB reagents (VectorLabs), counter

stained with Mayer's hematoxylin plus bluing reagent, and then dehydrated, cleared, and mounted with resin-

based medium (Cytoseal™ 60 Mounting Medium, Epredia).

Fluorescent Immunohistochemistry

Sections were permeabilized for at least 10 minutes with 0.3% Triton X-100 in PBSand subsequently blocked

for at least 40 minutes at room temperature with blocking solution (3% donkey serum and 2% BSA). Following,

sections were washed with PBS and incubated in primary antibody diluted in blocking solution for 2 hours at

room temperature. Next, sections were washed again and incubated in secondary antibody diluted in blocking

solution for 1 hour at room temperature. Sections were washed and exposed to Hoechst 33342 nuclear stain for

five minutes and mounted with an aqueous mounting medium (Fluoromount-G™).

<u>Study design</u>

The evaluation of the GABA switch in the naïve swine brain was planned prospectively and the maximum

biological replicates were collected as allowed by protocol restrictions. Sample size, demographics, and

characteristics are detailed in Figure 1A'. Postnatal samples were analyzed first to determine if the study was

sufficiently powered. Following this power analysis, all naïve swine samples were analyzed together to generate

heat maps and scatterplots in Figure 1. For the spatial transcriptomics experiment, Figure 3, the region of interest

was the experimental unit.

The evaluation of frozen human hippocampus was performed in the BCH Department of Pathology. A series of

infants who died from non-TBI causes was used. Samples with an extended postmortem interval (PMI) (>30

hours) were excluded because prolonged PMI positively correlated with NKCC1 expression, likely a reflection of

tissue autolysis in our pilot analysis. The effect of TBI injuries in swine on the cation chloride cotransporters was

characterized using aliquots from a prior cohort where the main findings of age, injury, seizure medications, and

mechanisms of vasogenic edema had been published. The effect of TBI injuries in humans on cation chloride

cotransporters was examined using autopsy samples and semi-quantitatively evaluated by a board-certified

forensic neuropathologist (DM).

Statistical analysis

Graphing and statistical analyses were completed in GraphPad Prism. Parametric statistical tests included

student's t-test, two- or three-way ANOVA considering age, treatment, co-variates, and/or region as applicable

and described in the figure legends. Single variate analysis utilized Pearson correlation and curve fit (R^2) and/or

p values are displayed on the graph where applicable. Multivariate analysis was performed when applicable and

multiple linear regressions (Pearson's) correlation matrices are displayed on the figure. Exact p values are

included on the graphs. For non-parametric data, either Spearman correlation, Mann-Whitney, Wilcoxson Signed

Rank Test or Kruskal-Wallis Test by Ranks considering age, treatment, and/or region as applicable were

performed as described in the figure legends.

RESULTS:

Establishing the swine "GABA switch" with high spatial and temporal resolution.

We examined the expression and abundance of key chloride cotransporters (*Slc12a2*/NKCC1 and

Slc12a5/KCC2) and a regulatory kinase (*Stk39/*SPAK) in the American Yorkshire pig (*Sus scrofa domesticus*)

across several developmental stages, aiming to map how these changes might align with a prenatal, perinatal,

or postnatal GABA switch. In many studies, the relative mRNA expression of these transporters, and a ratio of

Slc12a5 / Scl12a2 have been utilized as a marker for the "GABA switch" (ex. Sedmak et al., 2016).

mRNA expression of Slc12a2, Slc12a5, and Stk39 across development and region in swine

Our intent was to determine whether transcriptional patterns recapitulate those described in other species

and to gain initial insight into the developmental timing of the GABA switch in swine. Transcriptional analysis

suggests the swine GABA switch occurs perinatally, with region-specific trends in *Slc12a2* (NKCC1) and *Slc12a5*

(KCC2) expression differing from altricial species, while *Stk39* (SPAK) levels showed no significant

developmental changes.

We began by quantifying mRNA expression of *Slc12a2*, *Slc12a5*, and *Stk39* in somatosensory cortex ,

hippocampus, and thalamus at embryonic day 36 (ED36), ED90, ED105, postnatal day 7 ("infant"), and postnatal

day 30 ("toddler") (**Figure 1A**; demographic data in **1A'**). As shown in **Figures 1B and 1B'**, *Slc12a2* expression

remained relatively constant in hippocampus and thalamus, while cortex levels exhibited a moderate upward

trend over time. Although these findings were somewhat variable, this complexity may reflect underlying cellular

heterogeneity between samples, as NKCC1 is expressed in both neurons and non-neuronal cells (Kurki et al.,

2023*)*. For *Slc12a5* (**Figures 1C and 1C'**), expression in the hippocampus and thalamus remained relatively

stable, but the cortex showed a marked developmental decrease (Spearman non-parametric R^2=0.89,

p=0.0235). This downward cortical trend contrasts with the postnatal increase reported in altricial species,

suggesting that, in swine, the GABA switch might occur earlier, closer to birth—more in line with precocial and

semi-altricial mammals (Ben-Ari 2014). Lastly, *Stk39* (**Figure 1D and 1D'**) did not show significant age- or region-

related trends, indicating that *Stk39* transcript levels may not be the most sensitive indicator of developmental

shifts in GABAergic signaling at this age.

Taken together, our transcriptional analysis provided initial clues that the swine GABA switch may have

a perinatal timing, similar to humans and non-human primates, and differs from the patterns observed in highly

altricial species such as rodents (Ben-Ari 2014; Khazipov et al., 2001; Sedmak et al., 2016). However, the

complexity and variability in these data demonstrate the limitations of relying solely on transcriptional profiles to

resolve developmental changes in chloride cotransporter function. To gain a clearer understanding we next

turned to protein-level analyses.

Protein abundances and phosphorylation state for NKCC1, KCC2, and SPAK across development and region in

swine

In protein-level analyses, we observed age-related increases in KCC2/NKCC1 ratios in swine cortex and

hippocampus, stable ratios in the thalamus, and opposite trends in SPAK abundance and phosphorylation.

Representative western blots, summarized in **Figure 1E**, reflect consistent findings from multiple technical

replicates and mean abundances in each biological replicate group. To provide a clearer overview, we visualized

mean abundances and phosphorylation ratios as summary heat maps in **Figures 1F–K**, and we considered

established interpretations like the ratio of KCC2/NKCC1 (**Figure 1L**) to better contextualize these changes. We

found that in naive swine brain, the dimer band of KCC2 and pKCC2 (~250kDa) was more predominant across

all ages, so we quantified that band (Agez et al., 2017).

In the cortex, the KCC2/NKCC1 ratio increased with postnatal age (Spearman R^2=0.7253, p=0.0671,

Figure 1L), aligning with a developmental shift toward inhibitory GABA signaling. However, when examining the

ratio of phosphorylated (i.e., "activated") KCC2 to phosphorylated NKCC1, we found no clear age effect. This

finding suggests that while overall protein levels of these transporters adjust with development, changes in their

phosphorylation states might follow a more complex pattern, potentially influenced by other post-translational

modifications of these transporters or other modulators of intraneuronal chloride and is beyond the scope of this

study.

Regionally, hippocampal KCC2/NKCC1 ratios also increased with age (R^2=0.8784, non-linear fit, **Figure**

1N), mirroring known patterns in rodent models and underscoring KCC2's essential role in maturing hippocampal

neurons. In contrast, thalamic ratios remained stable across the developmental time points examined, perhaps

indicating that the switch had occurred earlier than in the hippocampus and cortex (**Figure 1O**). Furthermore,

although we observed an overall increase in SPAK abundance with age, its phosphorylation state declined—a

dynamic that may reflect evolving regulatory mechanisms as the brain matures.

Postnatal pNKCC1 downregulation in human hippocampus.

To compare these developmental patterns with those observed in humans, we examined hippocampal

samples from infants who died unexpectedly from known causes of death. The study design and tissue validation

steps are outlined in **Figure 2A.** Previous transcriptional data from qPCR and microarray studies suggest that

the GABA switch in humans occurs around birth in the limbic system and neocortex (Sedmak et al., 2016; Tao

et al., 2012; Hyde et al., 2011). Furthermore, evidence from slice electrophysiology, Western blot analysis, and

immunolocalization has indicated hyperpolarizing GABA activity in the adult temporal lobe cortex and

hippocampus (Huberfeld et al., 2015; Wang et al., 2010; Munoz et al., 2007), which is altered in epilepsies.

However, our study offers a more detailed resolution and analysis of the activity state (via phosphorylation) of

these cotransporters during the immediate postnatal period in infants with normal brain development.

Like our approach in swine, we began by measuring mRNA levels of *Slc12a2, Slc12a5*, and *Stk39* in

relation to adjusted postconceptional age (PCA) [gestational age plus postnatal age] and found no significant

age regulation. As shown in **Figure 2B**, transcriptional levels of these genes remained relatively stable across

postnatal ages, and no positive correlation emerged between the ratio of *Slc12a5/Slc12a2* and PCA (**Figure**

2C). These transcriptional results indicate that major transcriptional shifts in these transporters are minimal

during the postnatal period in humans. Similar to swine, transcriptional variability was low in the hippocampus.

We then assessed protein abundances and phosphorylation states in these hippocampal samples. Unlike

in swine, both the monomer and dimer band of KCC2 were prominent, so we quantified both bands additively.

However, the pKCC2 band was only visualized at the monomer weight in this cohort. We found that while total

NKCC1 and KCC2 levels increased with age, the ratio of KCC2/NKCC1 decreased with age (**Figure 2D–G**). In

contrast, the phosphorylation state of NKCC1 (p-NKCC1) declined significantly over time, while p-KCC2 levels

remained low and unchanged. When we considered the ratio of activated KCC2 to activated NKCC1 (**Figure**

2H), we found a clear upward trend with increasing age. These results suggest that post-translational

modifications—in particular, phosphorylation—may be a key driver of the GABA switch during human postnatal

development. The increase in relative KCC2 activation aligns with the gradual shift toward inhibitory GABA

signaling that characterizes brain maturation (Watanabe et al., 2019; Kahle et al., 2015).

TBI injuries in swine affect cation chloride cotransporter transcriptional levels in an age x treatment

manner.

We next examined the influence of TBI in piglets at different developmental stages and found that severe

TBI induced diverse transcriptional, total protein, and phospho-protein changes to NKCC1, KCC2, and SPAK

across regions and ages. Across these analyses, specific age- and injury-related patterns emerged. In the cortex,

transcripts for *Slc12a2*, *Slc12a5*, and *Stk39* showed main effects of age or treatment (TBI injuries versus sham

injuries), as well as age x treatment interactions in some cases (**Supplemental Figure 1**).

Spatial transcriptomics reveals tissue subtype-specific alterations in chloride transporter expression

and pathway dysregulation following severe TBI in swine.

To assess whether NKCC1 could serve as a novel therapeutic target for traumatic brain injury, we

assessed *Slc12a2* and *Slc12a5* expression spatially using the GeoMx platform in injured cortical gray and white

matter in a toddler piglet with severe brain injury compared to a sham, **Figure 3A**. **Figure 3B** displays 2-fold

changes in transcript abundance for *Slc12a2* and *Slc12a5* relative to sham gray matter from the GeoMx data.

We observed a significant increase in *Slc12a2* transcript levels in both sham and injured white matter compared

to gray matter, with injured white matter showing the highest expression. This upregulation may reflect the

presence of NKCC1 in glial cells of the white matter or in injured axonal projections, as NKCC1 typically localizes

to neuronal somas and axon initial segments. In contrast, *Slc12a5* transcript levels decreased in both sham and

injured white matter relative to gray matter, with a slight, non-significant reduction in injured white matter

compared to sham. This pattern suggests a region-specific downregulation of KCC2 in injured axons, potentially

contributing to chloride dysregulation in TBI-affected areas. We find a significant interaction between gene

expression, brain region, and injury, indicating differential responses to TBI in cortical gray and white matter for

both *Slc12a2* and *Slc12a5* (Three-Way ANOVA, **Figure 3B**). These findings suggest that TBI induces distinct

region-specific changes in chloride transporter expression, which may impact chloride balance and

excitatory/inhibitory signaling post-injury in toddler piglets.

Pathway analyses (**Figures 3C–D**) revealed that TBI-associated changes in gene expression in gyral

cortical gray matter encompassed both upregulated pathways related to cellular signaling, neural development,

and energy metabolism and downregulated pathways involved in cell adhesion, apoptosis, differentiation, and

ion transport. Although neuroinflammatory signatures were mixed and did not meet gene ontology enrichment

criteria in these damaged regions, these findings are consistent with known complexities underlying

neuroinflammation in TBI pathophysiology. Future exploration of additional regions, ages, and longer recovery

intervals will be critical to fully elucidate the molecular drivers of injury progression and identify novel therapeutic

targets.

TBI injuries in swine changes cation chloride cotransporter abundance and phosphorylation across

cortex and hippocampus.

Western blotting of cortical lysates (**Figures 4A–C**, quantified in **4D–E**), where the severe spreading

hypoxic-ischemic injury is typically most profound revealed that younger, infant piglets with TBI exhibited

elevated NKCC1, KCC2, and SPAK protein levels compared to older, "toddler" piglets. NKCC1 phosphorylation

(pNKCC1) also increased with TBI injury in both ages, indicating that TBI may influence the functional state of

this transporter in younger brains. Additionally, where the dimer of KCC2 and pKCC2 was the prominent band

in naive samples, in the TBI samples, the monomer band was equally, or sometimes more, abundant and was

distinct in pattern by region (**Figure 4A**). In these quantifications, the band intensities were added together (Agez

et al., 2017). This difference in blotting patterns for KCC2 compared to naive animals could be explained by the

24-hour anesthesia and intensive care regimens experienced by both sham and TBI injury animals, which was

absent in the naive cohort. Even though the piglets were anesthetized using a sedation regimen that largely

avoided the GABA receptor, anesthetic effects might still have been present. The significant upregulation of

SPAK in infant piglets with TBI, but not toddler piglets, is interesting for two reasons. 1) Our previous work

identified that while both ages demonstrate cerebral edema due to TBI, toddler piglets show more vasogenic

edema than infant piglets (Hochstetler, Price, et al., 2024). An upregulation of both SPAK and NKCC1 in infant

piglets with TBI injuries might be driving evolving cytotoxic edema that does not result in vasogenic edema

(Glykys et al., 2017; Zhang et al., 2020). 2) SPAK abundance can be promoted by NFkB signaling (Bhuiyan et

al., 2022), which could be activated by either the subarachnoid hemorrhage or cortical impact-induced

mechanical stress in this model.

In the hippocampus—an area typically spared from hypoxic-ischemic injury in this model—we observed

increased NKCC1, KCC2, and SPAK in infant TBI piglets compared to toddler TBI piglets (**Figure 4F**). This

suggests that even regions not displaying overt injury can exhibit early changes in chloride transporter

expression following TBI. Indeed, structural hippocampus damage might require additional time to manifest

beyond the 24-hour paradigm of our study. By contrast, no significant age- or treatment-related changes were

detected in the thalamus (**Figure 4G**), underscoring thalamic resiliency to this injury type.

These observations provide evidence that TBI influences chloride transporter expression and activation

differently across brain regions and developmental stages. Younger piglets appear more prone to TBI-related

shifts in these key ionic regulators, raising the possibility that early therapeutic interventions targeting chloride

cotransporters could be beneficial.

Severe TBI in swine produces region- and cell-type specific alterations in pNKCC1 that are correlated

with damage, subarachnoid hemorrhage, and seizure duration.

TBI significantly increased pNKCC1 expression in non-parvalbumin-expressing cortical neurons

(NeuN$^+$/pNKCC1$^+$/PV$^-$) of infant piglets positively correlated with seizure duration and SAH area, while cortical

PV$^+$ populations and hippocampal neuronal populations showed no significant changes, indicating region- and

cell-specific effects. We utilized immunohistochemistry to assess phosphorylated NKCC1 (pNKCC1) expression

in NeuN co-labeled mature neurons in the somatosensory cortex layer 4, revealing more age-specific effects of

injury (**Figure 5A**). Apoptotic neurons (**Supplemental Figure 2A**, assessed by nuclear chromatin condensation

via Hoechst/NeuN$^+$) in gyral layer 4 increased in toddlers but not infants (**Figure 5B).** Also in gyral layer 4, in the

remaining healthy NeuN$^+$ neurons, pNKCC1 counts increased in infants but not toddlers (**Figure 5B**). There was

no effect of age nor injury on the number of pNKCC1 in all NeuN$^+$ neurons in the dorsal or ventral hippocampus

dentate gyrus or CA2 **(Figure 5C)**. In all hippocampal regions, less than 5% of total neurons were apoptotic

(**Supplemental Figure 2B**).

In gyral cortical neurons, TBI significantly increased pNKCC1 expression (main effect of treatment

p=0.0017, **Figure 5B**). When comparing pNKCC1 levels to univariate injury metrics, **visualized in Figure 5D/E**,

seizure duration, % damage, and SAH show positive correlation with pNKCC1 in infant piglets (Pearson

R^2=0.838, p=0.0093, Spearman R^2=0.833, p=0.0738; Pearson R^2=0.6514, p=0.0801; respectively, **Figure 5E**)

Multivariate analysis further confirmed the relationship between neuronal-pNKCC1 expression and SAH% and

seizure duration as positive predictors across both ages (**Figure 5F**).

Further cell-specific analyses indicated that PV$^+$ interneurons did not show enhanced pNKCC1 after TBI,

suggesting that these inhibitory interneurons remain relatively unaffected. Non-neuronal cells in the cortex

displayed nominal, heterogeneous changes in pNKCC1. In the hippocampus, no significant pNKCC1 changes

were detected in either neuronal or non-neuronal populations, indicating a regionally selective effect of TBI on

pNKCC1 expression. Collectively, these data suggest that TBI-induced elevations in pNKCC1 predominantly

occur in surviving cortical PV⁻ neurons of younger piglets, possibly contributing to a pro-excitatory state

associated with seizures.

NKCC1 expression is also high in the choroid plexus of many mammals, including piglets. We observed

pNKCC1 downregulation and signal fragmentation in the choroid plexus due to severe TBI (**Figure 6A**). We

denote signal fragmentation as an ectopic, non-apical pNKCC1 signal in the epithelial cells, indicating the

presence, but potential lack of function, of this transporter. This is in opposition to a prior study where rats with

cortical contusion injuries showed an upregulation of NKCC1 in the choroid plexus within 24 hours following

injury (Lu et al., 2006). Two factors may explain this result. The first is the severity of the injury, where the rats

with contusion injuries were a much less severe model than our piglet model, which requires intensive care to

sustain. The second factor is age, where our piglet model is accomplished in young animals, the rat model

utilized adult animals.

Region- and cell-type-specific alterations in pNKCC1 expression are heterogeneous in an autopsy series

of pediatric TBI.

To bridge our findings from the piglet model to humans, we examined pNKCC1 expression in autopsy

samples from non-TBI and TBI cases covering a range of ages (detailed neuropathological findings in

Supplemental File 1). In control, human tissue, pNKCC1 was consistently present at low to moderate levels in

neurons and glia across multiple regions, with relatively robust expression in ependyma and choroid plexus. In

TBI cases, pNKCC1 expression varied according to injury severity and location. Regions with significant hypoxic-

ischemic damage, including the parietal lobe, corpus callosum, and hippocampus, showed increased pNKCC1

in glia and axons relative to controls. The temporal lobe, especially ependyma of the striatum terminalis, exhibited

pronounced pNKCC1 upregulation in cases with severe white matter injuries. Meanwhile, the choroid plexus

demonstrated reduced pNKCC1 expression, mirroring signal fragmentation patterns seen in piglets (**Figure 6B**).

In human cases, profound damage to the choroid plexus (temporal lobe, lateral ventricle) accompanied this

phenomenon. The atrium of the lateral ventricle was chosen in the piglets as a match for the human cases due

to the proximity to the hippocampus.

These human data align closely with the piglet findings, suggesting that TBI prompts region- and cell-

specific changes in chloride transporter phosphorylation. The parallels in pNKCC1 distribution and injury-related

shifts point toward a conserved mechanism by which TBI alters ionic balance and potentially contributes to

secondary injury processes in the developing brain in neurons that survive the brain injury.

DISCUSSION:

A central feature of normal brain development is the "GABA switch," in which inhibitory transduction

matures as the balance shifts from NKCC1-driven chloride influx toward KCC2-mediated chloride efflux. This

switch helps mature neural circuits, limit hyperexcitability, and prevent seizures. Our swine data show that this

switch occurs around the perinatal period in a pattern more closely resembling semi-altricial species (including

primates) than altricial rodents. In humans, we similarly observed a gradual postnatal decrease in NKCC1

phosphorylation aligned with a relative increase in potential inhibitory tone over time. Together, these findings

suggest that swine provide a useful experimental bridge for modeling human neurodevelopment and the timing

of the GABA switch.

We hypothesized that regions of tissue damage (i.e. cerebral cortex) would exhibit changes to the

NKCC1/KCC2 balance and regions without tissue damage (hippocampus and thalamus) would not. We did find

upregulation of NKCC1 in the cortex as predicted, but we also found changes in hippocampus as well, a region

which does not have acute tissue damage in our model at the 24 post-injury timepoint, but perhaps would exhibit

structural damage chronically (Costine-Bartell et al., 2019). We saw no change to transporter abundance in

thalamus, an area we expect to be spared in our model both acutely and chronically as the thalamus is spared

in children with Hemispheric Hypodensity (Durham and Duhaime 2007).

We further hypothesized that toddler piglets with more pervasive, hemispheric spreading injury with and

more pronounced vasogenic edema would exhibit upregulated NKCC1 compared to infant piglets with patchy,

less tissue damage. However, we found the opposite to be true where greater upregulation of NKCC1 and SPAK

in response to injury was not in the age with greater tissue damage. Younger animals displayed a more

pronounced TBI-induced upregulation of NKCC1 and SPAK, a kinase that promotes NKCC1 phosphorylation

and activity. This shift toward elevated NKCC1 function can increase neuronal excitability and seizure risk—

clinical challenges frequently observed in pediatric TBI. This is perhaps because apoptotic (dead) neurons,

perhaps due to cytotoxic edema, which are abundant in the injured toddler piglets, are non-functional and will

not be affected by changes to the Cl^- milieu, whereas surviving neurons might exhibit ongoing dysfunction. The

correlation between pNKCC1 counts and seizure duration and hemorrhage in piglets further reinforces the notion

that disrupted chloride regulation might contribute to excitatory/inhibitory imbalances after injury. Notably, these

TBI-induced changes were not uniform across cell types or regions: parvalbumin-positive cortical interneurons

were spared, and hippocampal neurons remained relatively unaffected by these specific injuries. Since

hippocampal neurons didn't show significantly increased neuronal-pNKCC1, but lysates showed upregulated

NKCC1, perhaps non-neuronal NKCC1 upregulation contributes to early responses to TBI before neuron

dysfunction. Moreover, changes in chloride transporter expression extended beyond neurons and non-neuronal

cortical cells, such as the downregulation in the choroid plexus. Spatial transcriptomics indicated that white

matter alterations in NKCC1 and KCC2 expression likely involve glial cells and support matrix, influencing

extracellular ion balance and potentially contributing to edema and barrier dysfunction. Indeed, we previously

demonstrated that toddlers, which have the most damage, also have ongoing upregulation in abundance and

activity of matrix metalloproteinase, whereas matrix metalloproteinases are either not increased or even

downregulated in infants, who have less damage (Hochstetler et al., 2024). This works supports the hypothesis

that chloride transporters alone do not control intraneuronal chloride concentrations, especially after pediatric

TBI (Normoyle et al., 2024). Human autopsy data corroborated these animal findings. In TBI cases, we saw

similar region- and cell-type-specific shifts in pNKCC1, particularly in damaged white matter tracts and in the

choroid plexus. We believe this dysregulation of choroidal pNKCC1 signal could likely be a consequence of

severe injury, including hemorrhages into the CSF space – specifically subarachnoid hemorrhage (SAH) and

intraventricular hemorrhage (IVH), and perhaps CSF production by choroid plexus transporters is shunted. An

alternative explanation could be that the choroid plexus is mechanically damaged in our injury model and in

severe TBI in humans. Together, our finding of choroid plexus pNKCC1 dysregulation demonstrates how severe

TBI disturbs ionic homeostasis at brain-fluid interfaces, potentially affecting cerebrospinal fluid dynamics, cell

volume, bulk tissue shifts, and intracranial pressure.

Taken together, after TBI, toddler piglets display vasogenic edema likely from MMP upregulation, while

infant piglets might instead experience acute cytotoxic edema with rapid, clean cell death without leading to

vasogenic edema, resulting in ongoing neuronal dysfunction in surviving neurons, which might make them

unique candidates for therapeutic intervention (Balena and Staley, 2024). This postulate is supported by Level

III clinical evidence and recent age-dependent mechanisms of cerebral edema in small diameter blood vessels

following severe TBI in an autopsy case-series investigation (Fullerton et al., 2021). Given the robust age- and

treatment-effects of TBI injuries on SPAK abundance, we propose that treatment of our infant piglets with a

SPAK inhibitor immediately following TBI might offer protective and antiepileptic effects (Gong et al., 2021; Wu

et al. 2020). Indeed, as pathophysiology is age-specific, successful therapeutics might also need to be age-

specific after severe TBI according to developmental stage: targeting NKCC1 or SPAK in infants and targeting

MMP's in toddlers.

Limitations:

Our study has several limitations that should be acknowledged. Firstly, we were not powered to examine

the distinct effect of sex as a covariate on NKCC1 and KCC2 levels in the developing swine brain due to protocol

limitations, which may influence the observed outcomes (Missios et al., 2009). Additionally, we did not assess

the levels of brain-derived neurotrophic factor (BDNF) or insulin-like growth factor (IGF), both of which are known

to promote the GABA switch. Other intracellular regulators such as protein kinase C (PKC), Src kinases, and

with no-lysine (WNK) kinases, the latter of which are chloride-sensitive and may play a role in the switch, were

also not considered in this study (Watanabe and Fukuda 2015). Furthermore, we did not investigate the potential

contributions of maternal estrogen or taurine as positive sensory regulators of the GABA switch. Finally, and

perhaps most importantly, we did not perform electrophysiology or chloride ion imaging either *in vivo* or in

organotypic slice, thus, our observations are limited to when the switch is *likely* occurring, however, we cannot

say with certainty when GABA switches from excitatory to inhibitory in the developing swine brain.

Our study, while leveraging a translationally relevant swine model and human autopsy samples, cannot

fully capture the clinical complexity and heterogeneity of pediatric TBI. The standardized injury approach in pigs,

though informative, may not reflect the multifaceted trauma mechanisms, variable interventions, and long

recovery trajectories seen in patients. These factors represent important avenues for future research to better

understand the mechanisms underlying the excitatory/inhibitory balance in the developing swine brain and in the

context of TBI. Donnan effects with the loss of the extracellular matrix also likely affect the amount of

extraneuronal and intraneuronal chloride concentrations post-TBI as demonstrated in rodent hippocampal

explants (Glykys et al., 2014; Normoyle et al., 2024). Indeed, in our own severe TBI model, we have described

an age-dependent upregulation of matrix metalloproteinases where there was an ongoing upregulation in

toddlers with worse injury, compared to a resolution in infants with lower injury burden scaled to brain volume.

The human autopsy samples, while invaluable, represent end-stage conditions and lack information on the

temporal progression of TBI pathology. Further, it is unclear to what extent non-traumatic pediatric autopsy

samples are representative of a physiologically normal pediatric population.

We observed notable variability in transcript-level measurements, likely due to biological heterogeneity

and technical factors such as RNA quality, sampling consistency, and cellular composition of lysates. Though

immunoblotting analyses were particularly informative, phosphorylation state was less informative in swine. It is

important to note that our study did not evaluate additional post-translational modifications, such as glycosylation,

though these are certainly indicated in the activity of these transporters. Finally, while we identified potential

mechanistic links between chloride transporter dysregulation, seizures, and injury severity, further studies are

needed to establish causal relationships and to explore how interventions that target these pathways might

improve long-term outcomes in pediatric TBI.

Conclusions and future directions:

By integrating findings from a swine model—whose brain development and structure closely parallel

those of humans—with postmortem human tissue analyses, our work supports the translational value of targeting

chloride regulation to improve TBI outcomes, especially in pediatric populations (Glykys et al., 2017; Soul et al.,

2021; Staley 2022). Pharmacologically modulating NKCC1 or KCC2—potentially with agents like bumetanide or

a SPAK inhibitor—could help restore chloride homeostasis and reduce hyperexcitability and swelling/tissue shifts

after pediatric TBI. Such treatments would need to consider the patient's developmental stage and the specific

brain regions affected. Early intervention, when the brain is most vulnerable to ionic imbalances, may be

especially beneficial. Our findings emphasize that TBI's impact on the developing brain cannot be fully

understood by focusing solely on neuronal excitability or a single brain area. Instead, we must consider a

dynamic interplay among developmentally regulated ion transporters, glial function, neuroinflammatory

processes, and structural integrity. Future work should address 1) how these factors interact over longer recovery

periods, potentially via *in vivo* monitoring of neuronal chloride (Costine-Bartell et al., 2023), 2) explore targeted

therapies that normalize chloride balance, and 3) consider how interventions might influence long-term

neurodevelopmental outcomes and post-traumatic sequelae such as epilepsy.

References:

Agez M, Schultz P, Medina I, Baker DJ, Burnham MP, Cardarelli RA, Conway LC, Garnier K, Geschwindner S, Gunnarsson A, McCall EJ, Frechard A, Audebert S, Deeb TZ, Moss SJ, Brandon NJ, Wang Q, Dekker N, Jawhari A. (2017) Molecular architecture of potassium chloride co-transporter KCC2. Sci Rep. Nov 28;7(1):16452.

Balena T, Staley K. (2024). Neuronal Death: Now You See It, Now You Don't. Neuroscientist. Sep 24:10738584241282632.

Ben-Ari Y, Gaiarsa JL, Tyzio R, Khazipov R. (2007). GABA: a pioneer transmitter that excites immature neurons and generates primitive oscillations. Physiol Rev. Oct;87(4):1215-84.

Ben-Ari Y. (2014). The GABA excitatory/inhibitory developmental sequence: a personal journey. Neuroscience. Oct 24;279:187-219.

Bhuiyan MIH, Young CB, Jahan I, Hasan MN, Fischer S, Meor Azlan NF, Liu M, Chattopadhyay A, Huang H, Kahle KT, Zhang J, Poloyac SM, Molyneaux BJ, Straub AC, Deng X, Gomez D, Sun D. (2022) NF-κB Signaling-Mediated Activation of WNK-SPAK-NKCC1 Cascade in Worsened Stroke Outcomes of Ang II-Hypertensive Mice. Stroke. May;53(5):1720-1734.

Bienboire-Frosini C, Marcet-Rius M, Orihuela A, Domínguez-Oliva A, Mora-Medina P, Olmos-Hernández A, Casas-Alvarado A, Mota-Rojas D. (2023) Mother-Young Bonding: Neurobiological Aspects and Maternal Biochemical Signaling in Altricial Domesticated Mammals. Animals (Basel). Feb 2;13(3):532

Bortone D, Polleux F. (2009). KCC2 expression promotes the termination of cortical interneuron migration in a voltage-sensitive calcium-dependent manner. Neuron. Apr 16;62(1):53-71.

Clancy B, Finlay BL, Darlington RB, Anand KJ. (2007). Extrapolating brain development from experimental species to humans. Neurotoxicology. Sep;28(5):931-7.

Costine-Bartell B, Price G, Shen J, McGuone D, Staley K, Duhaime AC. (2021). A perfect storm: The distribution of tissue damage depends on seizure duration, hemorrhage, and developmental stage in a gyrencephalic, multi-factorial, severe traumatic brain injury model. Neurobiol Dis. Jul;154:105334.

Costine-Bartell BA, Martinez-Ramirez L, Normoyle K, Stinson T, Staley KJ, Lillis KP. (2023). 2-photon imaging of fluorescent proteins in living swine. Sci Rep. Aug 29;13(1):14158.

Costine-Bartell BA, McGuone D, Price G, Crawford E, Keeley KL, Munoz-Pareja J, Dodge CP, Staley K, Duhaime AC. (2019). Development of a Model of Hemispheric Hypodensity ("Big Black Brain"). J Neurotrauma. Mar 1;36(5):815-833.

Dobbin J and Sands J. (1979) Early Human Development. Vol. 311 Pages 79-83.

Duhaime AC, Durham S. (2007) Traumatic brain injury in infants: the phenomenon of subdural hemorrhage with hemispheric hypodensity ("Big Black Brain") Prog Brain Res. 161:293-302.

Foster KA, Recker M, Lee P, Bell MJ, Tyler-Kabara E. (2014) Factors Associated with Hemispheric Hypodensity after Subdural Hematoma following Abusive Head Trauma in Children. J Neurotrauma. 2014 Oct 1;31(19):1625-31.

Fullerton JL, Hay J, Bryant-Craig C, Atkinson J, Smith DH, Stewart W. (2024) Pediatric Traumatic Brain Injury and Microvascular Blood-Brain Barrier Pathology. JAMA Netw Open. Nov 4;7(11):e2446767.

Glykys J, Dzhala V, Egawa K, Balena T, Saponjian Y, Kuchibhotla KV, Bacskai BJ, Kahle KT, Zeuthen T, Staley KJ. (2014) Local impermeant anions establish the neuronal chloride concentration. Science. Feb 7;343(6171):670-5.

Glykys J, Dzhala V, Egawa K, Kahle KT, Delpire E, Staley K. (2017) Chloride Dysregulation, Seizures, and Cerebral Edema: A Relationship with Therapeutic Potential. Trends Neurosci. May;40(5):276-294.

Goldstein RD, Poduri A. (2024) Seizures and Sudden Death Beyond SUDEP. Neurology. Feb 13;102(3):e208119.

Gong Y, Wu M, Gao F, Shi M, Gu H, Gao R, Dang BQ, Chen G. (2021) Inhibition of the p-SPAK/p-NKCC1 signaling pathway
protects the blood-brain barrier and reduces neuronal apoptosis in a rat model of surgical brain injury. Mol Med Rep.
Oct;24(4):717.

Harowitz J, Crandall L, McGuone D, Devinsky O. (2021) Seizure-related deaths in children: The expanding spectrum.
Epilepsia. Mar;62(3):570-582.

Haynes RL, Kinney HC, Haas EA, Duncan JR, Riehs M, Trachtenberg F, Armstrong DD, Alexandrescu S, Cryan JB, Hefti
MM, Krous HF, Goldstein RD, Sleeper LA. (2021) Medullary Serotonergic Binding Deficits and Hippocampal Abnormalities
in Sudden Infant Death Syndrome: One or Two Entities? Front Pediatr. Dec 21;9:762017.

Hochstetler A, Price G, Baohan A, Li M, Rodriguez Lara F, Lok J, Costine-Bartell B. (2024). Developmental Regulation of
Matrix Metalloproteinases in Response to Multifactorial, Severe Traumatic Brain Injuries during Immaturity. Dev Neurosci.
46(5):319-332.

Huberfeld G, Blauwblomme T, Miles R. (2015) Hippocampus and epilepsy: Findings from human tissues. Rev Neurol
(Paris). Mar;171(3):236-51.

Hyde TM, Lipska BK, Ali T, Mathew SV, Law AJ, Metitiri OE, Straub RE, Ye T, Colantuoni C, Herman MM, Bigelow LB,
Weinberger DR, Kleinman JE. (2011) Expression of GABA signaling molecules KCC2, NKCC1, and GAD1 in cortical
development and schizophrenia. J Neurosci. Jul 27;31(30):11088-95.

Ibrahim NG, Natesh R, Szczesny SE, Ryall K, Eucker SA, Coats B, Margulies SS. (2010). In situ deformations in the
immature brain during rapid rotations. J Biomech Eng. Apr;132(4):044501.

Kahle KT, Deeb TZ, Puskarjov M, Silayeva L, Liang B, Kaila K, Moss SJ. (2013) Modulation of neuronal activity by
phosphorylation of the K-Cl cotransporter KCC2. Trends Neurosci. Dec;36(12):726-737.

Keenan HT, Runyan DK, Marshall SW, Nocera MA, Merten DF, Sinal SH. (2003). A population-based study of inflicted
traumatic brain injury in young children. JAMA. Aug 6;290(5):621-6.

Khazipov R, Esclapez M, Caillard O, Bernard C, Khalilov I, Tyzio R, Hirsch J, Dzhala V, Berger B, Ben-Ari Y. (2001) Early
development of neuronal activity in the primate hippocampus in utero. J Neurosci. Dec 15;21(24):9770-81.

Kurki SN, Uvarov P, Pospelov AS, Trontti K, Hübner AK, Srinivasan R, Watanabe M, Hovatta I, Hübner CA, Kaila K, Virtanen
MA. (2023). Expression patterns of NKCC1 in neurons and non-neuronal cells during cortico-hippocampal development.
Cereb Cortex. May 9;33(10):5906-5923.

Leitner DF, William C, Faustin A, Askenazi M, Kanshin E, Snuderl M, McGuone D, Wisniewski T, Ueberheide B, Gould L,
Devinsky O. (2022) Proteomic differences in hippocampus and cortex of sudden unexplained death in childhood. Acta
Neuropathol. May;143(5):585-599.

Lind NM, Moustgaard A, Jelsing J, Vajta G, Cumming P, Hansen AK. (2007). The use of pigs in neuroscience: modeling
brain disorders. Neurosci Biobehav Rev. 31(5):728-51.

LoTurco JJ, Owens DF, Heath MJ, Davis MB, Kriegstein AR. (1995). GABA and glutamate depolarize cortical progenitor
cells and inhibit DNA synthesis. Neuron. Dec;15(6):1287-98.

Lu KT, Wu CY, Cheng NC, Wo YY, Yang JT, Yen HH, Yang YL. (2006). Inhibition of the Na+ -K+ -2Cl- -cotransporter in
choroid plexus attenuates traumatic brain injury-induced brain edema and neuronal damage. Eur J Pharmacol. Oct
24;548(1-3):99-105. doi: 10.1016/j.ejphar.2006.07.048.

Missios S, Harris BT, Dodge CP, Simoni MK, Costine BA, Lee YL, Quebada PB, Hillier SC, Adams LB, Duhaime AC. (2009).
Scaled cortical impact in immature swine: effect of age and gender on lesion volume. J Neurotrauma. Nov;26(11):1943-51.

Muñoz A, Méndez P, DeFelipe J, Alvarez-Leefmans FJ. (2007) Cation-chloride cotransporters and GABA-ergic innervation
in the human epileptic hippocampus. Epilepsia. Apr;48(4):663-73.

Normoyle KP, Lillis KP, Egawa K, McNally MA, Paulchakrabarti M, Coudhury BP, Lau L, Shiu FH, Staley KJ. (2024)
Displacement of extracellular chloride by immobile anionic constituents of the brain's extracellular matrix. J Physiol. Dec 2.

Patodia S, Somani A, Thom M. (2021) Review: Neuropathology findings in autonomic brain regions in SUDEP and future
research directions. Auton Neurosci. Nov;235:102862.

Peerboom C, Wierenga CJ. (2021). The postnatal GABA shift: A developmental perspective. Neurosci Biobehav Rev.
May;124:179-192.

Sauro KM, Wiebe S, Dunkley C, Janszky J, Kumlien E, et al. 2016. The current state of epilepsy guidelines: A systematic
review. Epilepsia 57: 13-23

Sedmak G, Jovanov-Milošević N, Puskarjov M, Ulamec M, Krušlin B, Kaila K, Judaš M. (2016) Developmental Expression
Patterns of KCC2 and Functionally Associated Molecules in the Human Brain. Cereb Cortex. Dec;26(12):4574-4589.

Smith DM, McGinnis EL, Walleigh DJ, Abend NS. 2016. Management of Status Epilepticus in Children. J Clin Med 5.

Soul JS, Bergin AM, Stopp C, Hayes B, Singh A, Fortuno CR, O'Reilly D, Krishnamoorthy K, Jensen FE, Rofeberg V, Dong
M, Vinks AA, Wypij D, Staley KJ; Boston Bumetanide Trial Group. (2021) A Pilot Randomized, Controlled, Double-Blind
Trial of Bumetanide to Treat Neonatal Seizures. Ann Neurol. Feb;89(2):327-340.

Staley KJ. (2022) Clarifications regarding bumetanide for neonatal seizures. Epilepsia. Jul;63(7):1863-1867. doi:
10.1111/epi.17278. Epub 2022 May 26.

Swindle MM, Smith AC, Hepburn BJ. (1988). Swine as models in experimental surgery. J Invest Surg. 1(1):65-79.

Tang BL. (2020) The Expanding Therapeutic Potential of Neuronal KCC2. Cells. Jan 17;9(1):240.

Tao R, Li C, Newburn EN, Ye T, Lipska BK, Herman MM, Weinberger DR, Kleinman JE, Hyde TM. (2012) Transcript-
specific associations of SLC12A5 (KCC2) in human prefrontal cortex with development, schizophrenia, and affective
disorders. J Neurosci. Apr 11;32(15):5216-22.

Wang WZ, Hoerder-Suabedissen A, Oeschger FM, Bayatti N, Ip BK, Lindsay S, Supramaniam V, Srinivasan L, Rutherford
M, Møllgård K, Clowry GJ, Molnár Z. (2010) Subplate in the developing cortex of mouse and human. J Anat. Oct;217(4):368-
80.

Watanabe M, Fukuda A. (2015) Development and regulation of chloride homeostasis in the central nervous system. Front
Cell Neurosci. Sep 24;9:371.

Watanabe M, Zhang J, Mansuri MS, Duan J, Karimy JK, Delpire E, Alper SL, Lifton RP, Fukuda A, Kahle KT. (2019).
Developmentally regulated KCC2 phosphorylation is essential for dynamic GABA-mediated inhibition and survival. Sci
Signal. Oct 15;12(603):eaaw9315.

Wu D, Lai N, Deng R, Liang T, Pan P, Yuan G, Li X, Li H, Shen H, Wang Z, Chen G. (2020) Activated WNK3 induced by
intracerebral hemorrhage deteriorates brain injury maybe via WNK3/SPAK/NKCC1 pathway. Exp Neurol. Oct;332:113386.

Young JW, Mayerl CJ, Mannava A, Lewis C, Fan T, Nair M, Mamone C, Schapker NM, Mossor AM, German RZ. (2023)
Evaluating The Precocial-altricial Axis of Motor Skill at Birth in A Preterm Pig Model. Integr Comp Biol. Sep 15;63(3):625-
640.

Zhang J, Bhuiyan MIH, Zhang T, Karimy JK, Wu Z, Fiesler VM, Zhang J, Huang H, Hasan MN, Skrzypiec AE, Mucha M,
Duran D, Huang W, Pawlak R, Foley LM, Hitchens TK, Minnigh MB, Poloyac SM, Alper SL, Molyneaux BJ, Trevelyan AJ,
Kahle KT, Sun D, Deng X. (2020) Modulation of brain cation-Cl- cotransport via the SPAK kinase inhibitor ZT-1a. Nat
Commun. Jan 7;11(1):78.

Figure 1:

Figure 1: The expression of KCC2 in cortex and hippocampus matures postnatally in the developing
swine brain.
A, study design, sample acquisition (**A'**), and approximate age of swine across the developmental axis. **B**,
expression of *Slc12a2* across development and region in swine presented as heat map for mean fold change
relative to ED90. All data points presented in **B'**. **C**, expression of *Slc12a5* across development and region in
swine presented as heat map for mean fold change relative to ED90. All data points presented in **C'**. **D**,
expression of *Stk39* across development and region in swine presented as heat map for mean fold change
relative to ED90. All data points presented in **D'**. **E**, representative western blots for each protein with total protein
stain presented as loading control. **F**, NKCC1 protein presented as heat map for mean a.u. normalized to total
protein across developmental ages and regions. **G**, phosphorylated NKCC1 protein presented as heat map for
mean pNKCC1 a.u. normalized to total NKCC1 across developmental ages and regions. **H**, KCC2 protein
presented as heat map for mean a.u. normalized to total protein across developmental ages and regions. **I**,
phosphorylated KCC2 protein presented as heat map for mean pKCC2 a.u. normalized to KCC2 a.u. across
developmental ages and regions. **H**, SPAK protein presented as heat map for mean a.u. normalized to total
protein across developmental ages and regions. **I**, phosphorylated SPAK protein presented as heat map for
mean pSPAK a.u. normalized to SPAK a.u. across developmental ages and regions. **L**, Ratio of KCC2/NKCC1
in cortex over development. **M**, Ratio of phosphoKCC2/phosphoNKCC1 in cortex over development. **N**, Ratio of
KCC2/NKCC1 in hippocampus over development. **O**, Ratio of KCC2/NKCC1 in thalamus over development. For
all correlative relationships, significant findings (Pearson or nonlinear fit) are noted on the graphs with a pink box
and Pearson fits and 95% Confidence interval bands are shown for all.

Figure 2:

Figure 2: Phosphorylated KCC2 increases with postnatal age in human hippocampus.
A, study design, acquisition, and demographics of fresh frozen infantile human hippocampus samples. *B*,
expression of *Slc12a2*, *Slc12a5*, and *Stk39* over postnatal development. *C*, Ratio of *Slc12a5/Slc12a2* transcript
over postnatal development. *D*, representative western blots from selected samples showing protein abundance
over developmental age with total protein stain as loading control. *E*, Protein abundances for NKCC1, KCC2,
and SPAK graphed over age for all samples. *F*, Abundance of KCC2 / NKCC1 over postnatal age. *G*, Ratio of
phosphorylated to total protein over developmental age for NKCC1 and KCC2. *H*, Ratio of activated KCC2 /
activated NKCC1 plotted over developmental age. For all linear relationships, Pearson correlations are noted on
the graphs and 95% Confidence interval bands are shown.

Figure 3:

Figure 3: Spatial transcriptomic profiling of toddler piglets reveals region- and injury- specific changes
to *Slc12a2* and *Slc12a5* and identifies mechanisms of ongoing tissue damage.
A, Study design and ROI selection for GeoMx Spatial Transcriptomics. ***B***, Selected relative expression of
Slc12a2 (encoding NKCC1) and *Slc12a5* (encoding KCC2) across regions from sham animals and animals with
severely injured tissue. Three-Way ANOVA performed to analyze transcriptional levels across gene, region, and
injury. ***C***, Gene ontology enrichment and KEGG (Kyoto Encyclopedia of Genes and Genomes) hierarchical
clustering for top 10% of upregulated genes in superficial gray matter. ***D***, Gene ontology enrichment and KEGG
clustering for top 10% of downregulated genes in superficial gray matter.

Figure 4:

735
736

Figure 4: The cortex and hippocampus were most sensitive to severe TBI injuires with upregulation of
NKCC1, KCC2, and SPAK abundance and phosphorylation unchanged, and thalamus was unaffected.
Western blots for all biological replicates for infant and toddler piglets receiving sham or severe TBI injuries in
cortex (**A**), hippocampus, (**B**), or thalamus (**C**). NKCC1, KCC2, and SPAK increased in cortex of infant piglets,
but not toddler piglets with TBI injuries, and the increase of KCC2/NKCC1 is driven by age, not treatment (**D**).
Levels of phosphorylated NKCC1 were affected by age and treatment, but not pKCC2, pKCC2/pNKCC1, or
pSPAK in cortex (**E**). NKCC1, KCC2, and SPAK increased in hippocampus of infant piglets, but not toddler
piglets with TBI injuries, and with no age or treatment effects on KCC2/NKCC1 (**F**). The levels of NKCC1, KCC2,
KCC2/NKCC1, and SPAK were unchanged in thalamus across ages and injury (**G**). Two-way ANOVA performed
with posthoc multiple t-tests, exact p-values reported on graph.

Figure 5:

749
750

Figure 5: Cortical non-apoptotic, PV⁻ neurons upregulate pNKCC1 acutely following TBI injuries and
are associated with burden of tissue damage across ages and burden of subarachnoid hemorrhage
and seizure duration in infant piglets with TBI.
Immunohistochemical staining for pNKCC1 (red), parvalbumin (green), NeuN (magenta), and nuclei (white), (**A**).
(**B**), percentage of apoptotic neurons, % of neuronal pNKCC1 expression, and % PV+ neuronal pNKCC1
expression due to age and injury in cortex in gyral or sulcal layer IV populations. Two-way ANOVA performed
with posthoc multiple t-tests, exact p-values reported on graph. (**C**), percentage of neuronal pNKCC1 expression
in dorsal or ventral CA2 (cornu ammnionis region 2) and DG (dentate gyrus) due to age and injury. Two-way
ANOVA performed with posthoc multiple t-tests, exact p-values reported on graph. (**D**), Injury characteristics as
unique variables, single variate analysis (**E**) performed across age and correlations (Pearson parametric or
Spearman non-parametric) shown on graphs. Multivariate analysis performed and correlation matrices
presented (**F**).

763
764

Figure 6:

Figure 6: pNKCC1 signal becomes irregular in choroid plexus following TBI in piglets and humans.
Representative fluorescent immunohistochemical staining in piglets for choroid plexus pNKCC1 (red) and nuclei
(blue) in sham, normal animals compared to infant and toddler pigs with TBI injuries, ***A***. White arrow shows shift
from apical to cytosolic localization, green arrows show regions of "fragmented" or "irregular" signal. ***B***,
Representative chromogenic immunohistochemical staining of pNKCC1 in choroid plexus in humans with or
without TBI. Non-TBI infants and toddlers show pNKCC1 restricted to epithelial cells of ChP, localization is
cytosolic and apical, with infants having more pNKCC1 signal than toddlers. In infant TBI, pNKCC1 signal
distribution is altered (red arrows). In toddler TBI, pNKCC1 signal is infrequently observed. In TBI, damage to
the ChP is apparent, particularly in toddlers.

777
Table 1: Human Sample Demographics

	Sample IDs	PNA (GA)	Sex	Diagnosis	PMI (hr)
Frozen/FFPE Hippocampus (Figure 2)	1	0.4 (40)	Female	Fatty acid oxidation disorder	15
	2	0 (41)	Female	Neonatal bradycardia with arrest and meconium aspiration	12.8
	3	17 (26.5)	Female	Acute asphyxiation	9
	4	12 (32)	Male	Acute asphyxiation	9
	5	5 (40)	Male	Acute asphyxiation	5
	6	11 (36)	Female	Acute asphyxiation	27
	7	13.5 (37.5)	Female	Acute asphyxiation	29
	8	16 (40)	Female	Acute asphyxiation	9
	9	26 (36)	Male	Acute asphyxiation	9
	10	33.4 (38.5)	Male	Intussuseption of inverted Merkel's diverticulum	13
	11	35 (39)	Female	Acute asphyxiation	29
FFPE Non-TBI Control (Figure 6, Supplemental File 1)	Non-TBI-1	0 (23)	Unknown	Normal brain development; labor/pregnancy complications	<24
	Non-TBI-2	0 (27)			
	Non-TBI-3	0 (39)			
	Non-TBI-4	0 (40)			
	Non-TBI-5	4 month	Male	Sudden unexpected death in infancy	24
	Non-TBI-6	5 month	Male	Cystic fibrosis	3.33
	Non-TBI-7	4 year	Female	Acute peritonitis	24
FFPE TBI (Figure 6, Supplemental File 1)	TBI - 1	3 month	Female	Blunt force head injury	23
	TBI - 2	3 year	Female	Traumatic axonal injury, subdural hematoma	24
	TBI - 3	3 year	Male	Road traffic accident, contusions, cerebral edema	24
	TBI - 4	9 year	Female	Road traffic accident, contusions, cerebral edema, subdural hematoma, global ischemia	24
	TBI - 5	9 year	Male	Road traffic accident, contusions, cerebral edema, subdural hematoma, global ischemia	24
	TBI - 6	10 year	Female	Road traffic accident, contusions, cerebral edema	72

778
779

Table 2: Antibody information.

Antibody	Species	Target	Company	Catalog #	Validation Method	WB	IHC
NKCC1	Rabbit	Synthetic Arg77 of mouse NKCC1	Cell Signaling Technology	14581	KO lysate	1:1000	
pNKCC1	Rabbit	Human pNKCC1 Th212/217	Millipore	ABS1004	KO lysate	1:500	1:250
SPAK	Mouse	GST-tagged c-terminus of human STK39	Millipore	MABS178	(+) ctrl human brain tissue	1:1000	
pSPAK	Rabbit	Human pSPAK Ser373	Millipore	07-2273	Hypotonic HEK293 cells	1:500	
KCC2	Rabbit	Synthetic peptide	Abcam	ab259969	(+) and (-) ctrl tissue lysates	1:2000	
pKCC2	Rabbit	Synthetic Ser940	Invitrogen	PA5-95678	KO lysate	1:1000	
Parvalbumin	Sheep	E. coli-derived recombinant human Parvalbumin alpha Ser2-Ser110	Invitrogen	PA5-47693	(+) ctrl brain tissue		1:100
NeuN	Conj 647	Synthetic peptide	Abcam	ab190565	(+) ctrl brain tissue		1:500

Table 3: Taqman probe information

Target	Species	Sequence
Nkcc1	*Human*	*Hs01074325_m1*
Kcc2	*Human*	*Hs00221168_m1*
Stk39	*Mouse*	*Mm01270753_m1*
Reference Gene (18s)	*N/A*	*Hs99999901_s1*

784 **Supplemental Figure 1:** Transcriptional changes to *Slc12a2, Slc12a5*, and *Stk39* due to age and injury in
785 piglets.
786 Quantitative PCR analysis for transcripts of *Stk39*, *Slc12a5*, and *Slc12a2* due to age, injury, and region in piglets,
787 presented as fold change from sham PND7 piglets. Two-way ANOVA results detailed in table on right and post-
788 hoc multiple t-tests exact p values included on graphs where applicable.

	Rostral Gyrus	Hippocampus	Thalamus
Age	0.0086	0.0104	0.2886
Treatment	0.0688	0.0176	0.0073
Age x Treatment	0.0352	0.0097	0.1703

	Rostral Gyrus	Hippocampus	Thalamus
Age	0.0039	0.00001	0.0038
Treatment	0.0636	0.2677	0.9538
Age x Treatment	0.0535	0.3256	0.0293

	Rostral Gyrus	Hippocampus	Thalamus
Age	0.0314	0.6891	0.5661
Treatment	0.2754	0.0085	0.0002
Age x Treatment	0.124	0.5932	0.5661

791   **Supplemental Figure 2:**

792   Examples of healthy neurons (peach arrowhead) and dead neurons (white arrowhead) in cortex as assessed by
793   nuclear stain (cyan) and NeuN[+] (magenta), **A**. Counts for apoptotic neurons in dorsal or ventral CA2 and DG
794   regions in hippocampus due to age and injury, **B**.

795