Research Article

Title: Temporal and cell-type specific SPAK-NKCC1 disruption following severe TBI in the developing gyrencephalic brain

Abbreviated title: Age-dependent changes to NKCC1 after severe TBI

Authors:

Alexandra Hochstetler[1,2], Ya'el Courtney[2,3], Peace Oloko[1], Benjamin Baskin[1], Andrew Ding-Su[1], Tawny Stinson[1], Declan McGuone[4], Robin Haynes[2], Maria K. Lehtinen[2,3], and Beth Costine-Bartell[1]*

*Corresponding Author, bcostinebartell@mgh.harvard.edu

Affiliations:

[1]Department of Neurosurgery, Massachusetts General Hospital, Harvard Medical School, Boston, MA 02115
[2]Department of Pathology, Boston Children's Hospital and Harvard Medical School, Boston, MA 02115
[3]Graduate Program in Neuroscience, Harvard Medical School, Boston, MA 02115
[4]Department of Pathology, Yale University School of Medicine, New Haven, CT, 06510

Conflict of Interest statement:

The authors declare no competing financial interests.

Acknowledgements:

This work is supported by NIH NICHD K01HD083759 and R01 HD099397 (BCB), by the Pappendick Family Translational Research Program (MKL), the Hydrocephalus Association, and R01 HD090064 (RH). AH was supported by T32NS007473 and F32NS134588 training fellowships from the National Institute for Neurological Disorders and Stroke. YC was supported by the Howard Hughes Medical Institute through the James J. Gilliam Fellowships for Advanced Study program. PO was supported by a Mass General Summer Research Trainee Program (SRTP). We thank the Boston Children's Hospital IDDRC Cellular Imaging Core supported by S10OD016453. Spatial transcriptomics was facilitated by the Comparative Pathology and Genomics Shared Resource core facility at The Cummings School of Veterinary Medicine at Tufts University. We thank the past and present team of undergraduate interns in the Costine-Bartell laboratory who assisted with the surgeries and 30 hours of intensive care to create the brain injury model and tissue collection as well as the Knight Surgical Research Lab for assistance with anesthesia to create the brain injuries. We are especially grateful for the fetal piglet tissue obtained from Dr. Ryann Fame (Stanford University Department of Neurosurgery), Dr. Mercedes Paredes (University of California, San Francisco (UCSF) Department of Neurology), and Dr. Misty Williams-Fritze (CBSET) and postnatal piglet tissue obtained from Dr. Corinna Beale (Center for Comparative Medicine, MGH). We thank Dr. Cameron Sadegh (UC Davis Department of Neurosurgery) and members of the Lehtinen Laboratory for assistance with prenatal swine brain dissections. We are deeply grateful to participating families for their courage and generosity in consenting to the use of their children in research. We also thank the clinicians, medical examiners, and pathologists who helped in the accrual of cases, including Drs. Gary Dale, Jeffrey Goldstein, Steven Campman, Catherine Stoos, and Colin Smith. We thank Elisabeth Haas and Richard Goldstein for the processing and analysis of infant cases. We thank Jane's Trust Initiative, the NIH NeuroBioBank, and the Edinburgh Brain Bank. We thank Kevin Staley for careful reading and suggestions on the research article.

Author contributions:

AH: Conceptualization, investigation, methodology, formal analysis, writing-original draft, visualization; YC: Investigation, methodology, formal analysis, writing-original draft; PO: Investigation, methodology, formal analysis, visualization; BB: Investigation, methodology, formal analysis, visualization; AD: Investigation, methodology; DM: Investigation, formal analysis, writing-revising; TS: Investigation, methodology; RH: Investigation, methodology, formal analysis, writing-revising, funding acquisition; MKL: Writing-revising, funding acquisition; BCB: Conceptualization, investigation, methodology, visualization, writing-original draft, project administration, funding acquisition

Abstract: Traumatic brain injury (TBI) is a leading cause of morbidity and mortality in infants and toddlers, with limited treatment options and persistent neurological sequelae. We developed a multi-pathoanatomic lesion multi-insult (MuLMI) severe TBI model in piglets that replicates age-dependent damage patterns to the cortical ribbon observed in human patients with less injury in postnatal day (PND) 7 "infant" piglets and more extensive tissue damage in PND30 "toddler" piglets. Given that neuronal chloride homeostasis influences excitability, seizure susceptibility, and edema, we examined the developmental and injury-induced regulation of key cation-chloride cotransporters and modulators: NKCC1 (sodium-potassium-2-chloride cotransporter), KCC2 (potassium-chloride cotransporter), and the regulatory kinase SPAK, which are biomarkers of neuronal chloride concentrations. This study is the first to define the spatiotemporal expression and phosphorylation profiles of these proteins in the developing piglet brain. We found a perinatal shift in the ratio of KCC2:NKCC1 across the brain, driven primarily by protein abundance, rather than transcriptional levels. We hypothesized that toddler piglets would exhibit an increase in cortical NKCC1 and SPAK causing hyperexcitability and perhaps explaining their more severe, unilateral cortical damage. Severe TBI induced a transcriptional increase in *Slc12a2* and *Stk39,* and a decrease in *Slc12a5* in toddler piglets, but not infant piglets. We further found that infant piglets, not toddler piglets, upregulated SPAK and Tyrosine Receptor Kinase B (TRKB) protein in cortex after TBI, with minimal changes in NKCC1 and KCC2. However, phosphorylated NKCC1 (pNKCC1) was significantly upregulated in surviving cortical neurons after TBI in infant piglets and was unchanged in toddlers, despite more severe injury. These findings suggest that cortical neuronal NKCC1 activation may play a role in post-traumatic excitability or resilience in the immature brain and identify NKCC1 and/or SPAK as a potential therapeutic target. In human tissue, the KCC2:NKCC1 ratio also increased postnatally, and TBI caused region and cell-type specific dysregulation of pNKCC1. Our results establish piglets as a valuable model for investigating age-specific mechanisms of pediatric TBI and for testing targeted interventions, particularly for infant populations where seizure control remains a major clinical challenge.

INTRODUCTION:

In infants and toddlers, most severe traumatic brain injury (TBI) results from abusive head trauma and is associated with high mortality and long-term neurological comorbidities [1]. Despite its prevalence, pediatric TBI remains understudied, and therapeutic options are limited. Young children of different ages exhibit distinct structural, biomechanical, and physiological vulnerabilities that shape injury severity and outcomes. Animal models must scale injury appropriately to brain volume, which increases significantly across postnatal development, especially in species like pigs and humans that undergo a perinatal brain growth spurt. Piglets are a high-fidelity model for pediatric TBI due to their gyrencephalic brain anatomy, developmental trajectory, and capacity to undergo intensive care interventions [1-5].

We employ a multi-pathoanatomic lesion, multi-insult TBI model (MuLMI) in piglets incorporating cortical impact, mass effect, subdural/subarachnoid hemorrhage, traumatic seizures, and transient apnea and hypoventilation under non-GABA-acting sedation to study the pathophysiology of severe TBI in young children. One example of MuLMI is the "hemispheric hypodensity" (HH) injury pattern observed in pediatric TBI patients: patchy, bilateral HH in infants and more unilateral HH in toddlers [6-8].

A critical contributor to these age-dependent injury patterns may be the "GABA switch"—the developmental transition in GABAergic signaling from excitatory to inhibitory, in part, driven by shifting expression and phosphorylation of two cation-chloride cotransporters: NKCC1 and KCC2. NKCC1 maintains high intracellular chloride ($[Cl^-]i$), promoting depolarizing GABA responses, whereas KCC2 lowers $[Cl^-]i$, enabling inhibitory GABA action [9, 10]. TBI may disrupt chloride homeostasis and exacerbate excitotoxicity [11-13]. Since standard seizure management for pediatric TBI often relies on GABA agonists [14, 15], an immature or disrupted chloride gradient may render these treatments ineffective or even harmful.

The timing of the GABA switch varies widely across species. In precocial species (e.g., sheep, guinea pigs), it occurs before birth; in altricial species (e.g., rodents), after birth; and in humans and swine, likely during the perinatal period [16-21]. Pinpointing this transition is essential for evaluating the translational validity of piglet models and for identifying vulnerable windows in pediatric brain injury. This is particularly relevant in diagnoses

where seizure plays a pathogenic role, such as sudden unexpected death in childhood and early-life

encephalopathies [22-26].

In this study, we defined the spatiotemporal expression of NKCC1 and KCC2 in the human and piglet

brain and investigated their regulation following severe TBI. We found that TBI induces upregulation of

phosphorylated NKCC1 (pNKCC1) in neurons of infant but not toddler piglets, with similar patterns observed in

human tissue. Notably, neuronal pNKCC1 expression correlated positively with seizure duration, subarachnoid

hemorrhage, and the extent of tissue damage: key features of our severe TBI model. These findings suggest

that aberrant NKCC1 activity contributes to excitability and injury in early development, and that NKCC1 inhibition

may represent a promising therapeutic target for mitigating damage after infant TBI. This work provides a

valuable platform for understanding age-dependent mechanisms of pediatric brain injury and evaluating novel

interventions in a translationally relevant large-animal model.

MATERIALS AND METHODS:

Sample procurement

Animal procedures

All protocols and procedures performed were in accordance with the guidelines of the American Veterinary Association, the National Institutes of Health, and the United States Department of Agriculture. They were approved by the Institutional Animal Care and Use Committee at Massachusetts General Hospital.

Naïve swine tissue collection

Prenatal collections

Preterm piglets were collected from euthanized Yorkshire sows at approximately 30% gestation (ED36), 75% gestation (ED90), and 90% gestation (ED105). Pig fetuses were surgically removed from the amniotic sac following maternal euthanasia, rinsed under cold running water, brains were dissected on ice, and tissues were immediately flash frozen. In all cases, the time from euthanasia to sample collection was less than 4 hours.

Postnatal collections

7-day-old (PND7, "infant") and 30-day-old (PND30, "toddler") male Yorkshire piglets were selected for naïve tissue collection. Surgery and anesthesia protocols were employed as previously described (Costine-Bartell et al., 2019), adjusted to the age of the piglets. In brief, piglets were sedated with ketamine/xylazine/atropine and anesthetized by 1-1.5% isoflurane with medical oxygen. Piglets were intubated and mechanically ventilated with room air. An ear intravenous or external jugular vein catheter was placed to access venous blood and deliver saline infusion. In all cases, mean arterial blood pressure, oxygen saturation, heart rate and end-tidal CO_2, respiratory rate, and core body temperature were managed for the duration of the procedures. A craniotomy was performed, and the dura was removed. Euthasol was delivered intravenously, and then fresh tissue dissection was performed immediately on wet ice. Samples (n=4/age) were collected from August 2023 to October 2023 and flash-frozen on dry ice.

Multi-focal severe TBI model injuries

In brief, a series of PND7 and PND30 male Yorkshire piglets from September 2016 to October 2019 were exposed to surgery and anesthesia protocols as previously described [6, 27]. A total of 23 piglets were used for

fixed tissue and a total of 14 for fresh tissue. Immediately prior to the injury, piglets were switched to a seizure-

permissive anesthesia protocol and subjected to a combination of injuries scaled to brain volume as previously

described, including cortical impact, mass effect, subdural hematoma/subarachnoid hemorrhage supplemented

with kainic acid, one-minute of apnea, and 10 minutes of hypoventilation. In the sham piglets, only the burr holes

and duration of anesthesia were modeled. Piglets were managed in an intensive care unit for 24 hours after

injury, where video EEG was recorded, beginning with scalp EEGs and then eventually with two 4-electrode

epidural EEG strips (1 per hemisphere), resulting in a 4-channel bipolar montage as previously described [27].

Seizure length, subdural hematoma area, subarachnoid hemorrhage area, and tissue damage were

analyzed as previously described [27]. Briefly, sections from FFPE blocks in 5 mm increments rostral to caudal

in each hemisphere were analyzed for damage that included areas of "red neurons," areas adjacent to red

neurons with vacuolization of neuropil and blood vessels, and areas with acute hemorrhage. "Red neurons" were

classified by the presence of all three of the following features: (1) cell shrinkage with perineuronal vacuolization,

(2) cytoplasmic hypereosinophilia, and (3) nuclear pyknosis. This pattern of damage is termed "hypoxic-ischemic

type injury," though a myriad of pathways can lead to this common end-stage pattern of apoptosis/necrosis.

Vacuolization alone was not included in the calculation of brain damage areas. Areas of damage were found to

mostly overlap with albumin extravasation, expression of amyloid precursor protein, and matrix

metalloproteinase-9, demonstrating vasogenic edema that extends far beyond the area of cortical impact [27,

28] where tissue edema might eventually lead to capillary and blood flow collapse. The percent of the hemisphere

damaged presented here is a subset of previously published cohorts for the purpose of displaying the model and

damage in the subjects used here for analyses of chloride salts membrane transporters.

Swine fresh tissue collection

Samples were requisitioned as freezer aliquots from prior studies (both sham and injured piglets) but are distinct

aliquots from the same piglets published in [28]. For fresh frozen tissue, Euthasol was delivered intravenously,

and then fresh tissue dissection followed immediately on wet ice, and samples were flash-frozen on dry ice. For

perfused-fixed tissue, piglets were transcardially perfused with room temperature normal saline at a rate of

~1L/min until complete surface vessel clearing followed by room temperature 10% neutral buffered formalin at

the same rate until total body stiffness. Brains were obtained and post-fixed in 10% neutral buffered formalin for

3-5 days until appropriate tissue stiffness was achieved.

Human tissue collection

Lysates from frozen hippocampus at the level of the lateral geniculate nucleus were collected from either the

NIH NeuroBioBank or the San Diego County Medical Examiner's office, the latter available for research under

the auspice of the California Code, Section 27491.41. All infants died suddenly and unexpectedly with the cause

of death determined at autopsy and adjudicated within the laboratory for research purposes (**Table 1**). Unstained

tissue sections from FFPE blocks were procured from autopsy brain samples (TBI or non-TBI control) performed

at the Neuropathology Department at the University of Edinburgh. Permission to use the brain tissue was given

according to the guidelines required by the respective ethics committees of these institutions: the Boston

Children's Hospital and Massachusetts General Hospital Institutional Review Boards. Demographic information

of all human samples is detailed in **Table 1**.

Standard laboratory techniques

Western Blotting

Fresh frozen samples were manually homogenized in N-PER lysis buffer (ThermoScientific #87792)

supplemented with 1X Halt (ThermoScientific #78440) protease and phosphatase inhibitor cocktail and lysed at

4°C for 60 minutes with end-over-end agitation. Lysates were spun at 16,000 x g for 10 minutes to pellet debris.

Supernatants were aliquoted and Pierce BCA assay was used protein concentration (ThermoScientific #23250).

20μg of the sample was reduced with 5X Laemmli Sample Buffer (BioLegend #426311) supplemented with B-

mercaptoethanol (5%) and diluted with double distilled water to an equivalent volume. Samples were heated at

70°C for 5 minutes, loaded into 4-15% TGX Stain-Free gels (Biorad # #4568086) and separated at 180V for 45

minutes. Gels were transferred to 0.2 μm nitrocellulose membranes at 1.0A for 30 minutes using the Bio-Rad

Trans-Blotter® system. Membranes were stained with Ponceau-S total protein stain blocked with EveryBlot

blocking buffer with gentle agitation. Membranes were incubated with primary antibody diluted in EveryBlot

blocking buffer washed with 1X Tris Buffered Saline-0.1% Tween-20 (TBST) and then incubated with secondary

antibody diluted in EveryBlot blocking buffer. Membranes were washed with TBST and TBS, exposed with

chemiluminescent reagents (Cytiva-Amersham #RPN2235), and visualized on a Bio-Rad Chemi-Doc Imager.

For antibody validation, full complete membranes were first probed with each antibody to evaluate non-specific

bands, specific bands and dimers, and to optimize dilutions. In the case of pNKCC1 and NKCC1 antibodies, our

laboratory has performed in-house knock-down validation. All validation methods are presented in **Table 2**. For

publication gels, technical duplicate membranes were cut at 80kDa to allow probing of 1) pSPAK, 2) SPAK, and

3) beta-actin (below 80kDa) simultaneously with 1) pNKCC1, 2) NKCC1, and 1) pKCC2, 2) KCC2 (above 80kDa).

In Figure 1 and 6, band intensities were normalized to total protein loaded (Ponceau stain) for the graphs, since

beta-actin and beta-tubulin were found to be poor loading controls across age and region. In Figures 3 and 5,

band intensities were normalized to both beta actin and total protein, and beta actin normalization is shown in

the figure. Membrane stripping was performed with ReStore™ Western Blot Stripping Solution and protocol

(ThermoScientific #21059). All western blot data presented and quantified in this manuscript was the result of a

minimum of three technical replications on different days.

Quantitative, Real-Time PCR

Samples were processed for RNA extraction using the Total RNA Miniprep Kit (New England BioLabs #T2010),

evaluated for concentration >50ng/μL and a quality ratio of A260/280 of >2.0. RNA was prepared for cDNA

synthesis with +reverse transcriptase and –reverse transcriptase reactions using the LunaScript® RT Supermix

Kit (New England Biolabs #E3010) using quality control metrics of concentration >100ng/μL and A260:280 ratio

>1.8. Samples were diluted to equivalent cDNA loading and probed in triplicate with Taqman™ reagents for

quantitative gene expression using fast run chemistry on an Applied Biosystems machine. In all cases, duplicate

-RT and singlet no template controls were run for each sample. Probe information is detailed in **Table 3**.

Spatial Transcriptomics

Spatial transcriptomics was conducted on anatomically matched formalin-fixed, paraffin-embedded (FFPE) brain

sections from a sham 30-day-old piglet and a 30-day-old piglet with severe traumatic brain injuries using

NanoString GeoMx® Digital Spatial Profiling technology. Regions of interest (ROIs) were selected to capture

key anatomical areas relevant to age-dependent TBI pathology and neuroinflammation. Following ROI selection,

samples underwent GeoMx® platform analysis, where regions were scanned, and digital counts were obtained

for gene targets of interest. Signal thresholding at a 3% cutoff was applied to exclude low-expression signals,

ensuring that only significant transcript data were included in downstream analysis. Normalization was performed

using internal controls (External RNA Control Consortium) provided by the GeoMx® platform to account for

technical variation across slides and samples. Upper quartile normalization was performed after filtering the

dataset to remove targets below the limit of quantitation.

Histology

Following immersion fixation, brains were washed with 1X phosphate-buffered saline (PBS), paraffin embedded

and sectioned at 5-µm thickness and mounted onto silanized glass slides. Immunofluorescence was performed

on these sections using antibodies detailed in **Table 2**. Briefly, sections were deparaffinized in xylenes,

rehydrated, and then subjected to antigen retrieval in boiling 10mM citric acid-Tween pH 6.0 for 20 minutes.

Staining protocols are presented in more detail below. In all cases, negative control sections were incubated in

corresponding species-specific IgG at an equivalent concentration to the primary antibody dilution during the

primary antibody incubation step. For figures where quantitation is presented, all slides evaluated were stained

together in a single batch to minimize signal variability. Slides were imaged on a Zeiss Axio Observer D1 inverted

microscope (fluorescent, 20x air objective) or a Zeiss LSM980 (20x air or 63x oil objective) at the Boston

Children's Hospital Cellular Imaging Core. Images were processed and assembled using ZEN Blue, ImageJ

software, and Adobe Illustrator. For quantitation presented in **Figures 4, 7, and S2**, the examiner performed cell

counts on images that had been blinded and randomized.

Chromogenic Immunohistochemistry

For chromogenic stains, a peroxidase quenching step was performed immediately prior to antigen retrieval.

Sections were then permeabilized for 5 minutes with 1% PBS-Tween-20 and blocked for 1 hour using normal

horse serum diluted in Super Sensitive Wash Buffer (BioGenex, HK583-5KEGP). Sections were incubated in

primary antibodies diluted in blocking buffer for 2 hours at room temperature. Slides were washed with 1% PBS-

Tween-20 and then incubated in the secondary antibody diluted in blocking buffer for 30 minutes. Sections were

washed and subjected to antibody visualization with Vectastain® ABC and DAB reagents (VectorLabs), counter

stained with Mayer's hematoxylin plus bluing reagent, and then dehydrated, cleared, and mounted with resin-

based medium (Cytoseal™ 60 Mounting Medium, Epredia). Slides were sent coded to a blinded pediatric

neuropathologist (DM) for analysis.

Fluorescent Immunohistochemistry

Sections were permeabilized for at least 10 minutes with 0.3% Triton X-100 in PBS and subsequently blocked

for at least 40 minutes at room temperature with blocking solution (3% donkey serum and 2% BSA). Following,

sections were washed with PBS and incubated in primary antibody diluted in blocking solution for 2 hours at

room temperature. Next, sections were washed again and incubated in secondary antibody diluted in blocking

solution for 1 hour at room temperature. Sections were washed and exposed to Hoechst 33342 nuclear stain for

five minutes and mounted with an aqueous mounting medium (Fluoromount-G™).

Study design and rigor

The evaluation of the GABA switch in the naïve swine brain was planned prospectively, and the maximum

biological replicates were collected as allowed by protocol restrictions. Postnatal samples were analyzed first to

determine if the study was sufficiently powered. Following this power analysis, all naïve swine samples were

analyzed together to generate bar plots as presented. The effect of TBI injuries in swine on the cation chloride

cotransporters was characterized using aliquots and sections from a prior cohort where the main findings of age,

injury, seizure medications, and mechanisms of vasogenic edema had been published. The original cohorts were

randomized to treatment and exclusion criteria as previously published. Real-time PCR was performed in

triplicate with the appropriate controls described above. Western blots were validated by running full length blots

with the appropriate controls described above for antibody specificity and were run in technical duplicate

(minimum of three times on two duplicate gels per run). For the spatial transcriptomics experiment, Figure 3, the

region of interest was the experimental unit. Cell counts from Figures 4, 7, and S2 were performed by counters

blinded to treatment as described above, with auditing on every 10th section. Counters were within 10% of each

other. The evaluation of frozen human brain sections was performed in the BCH Department of Pathology. A

series of infants who died from non-TBI causes was used. Samples with an extended postmortem interval (PMI)

(>30 hours) were excluded because prolonged PMI positively correlated with NKCC1 expression, likely a

reflection of tissue autolysis in our pilot analysis. The effect of TBI injuries in humans on cation chloride

cotransporters was examined using autopsy samples (**Table 1**) and semi-quantitatively evaluated by a board-

certified forensic neuropathologist (DM) blinded to cause of death and age.

Statistical analysis

Graphing and statistical analyses were completed in GraphPad Prism. Parametric statistical tests two- or three-

way ANOVA considering age, treatment, co-variates, and/or region as applicable and described in the figure

legends. Single variate analysis utilized Pearson correlation and linear fit (R^2) and/or p values are displayed on

the graph where applicable. Multivariate analysis was performed when applicable and multiple linear regressions

correlation matrices are displayed on the figure. Exact p values are included on the graphs. For non-parametric

data, Spearman correlation, Wilcoxson Signed Rank Test or Kruskal-Wallis Test by Ranks considering age,

treatment, and/or region as applicable were performed as described in the figure legends.

RESULTS:

Establishing the swine "GABA switch" with high spatial and temporal resolution.

We examined the expression and abundance of key chloride cotransporters (*Slc12a2*: NKCC1 and

Slc12a5: KCC2) and a regulatory kinase (*Stk39:* SPAK) in the American Yorkshire pig (*Sus scrofa domesticus*)

across several developmental stages and brain regions, aiming to map how these changes might align with a

prenatal, perinatal, or postnatal GABA switch (**Figure 1A, B**). We used whole tissue lysates of specific brain

regions (**Figure 1C**) and applied quantitative PCR of tissue lysates, and western blot of total protein and

phospho-protein (**Figure 1D, E**) to assess this question.

mRNA expression of Slc12a2, Slc12a5, and Stk39 across development and region in swine

In many studies, the relative mRNA expression of these transporters and a ratio of *Slc12a5: Scl12a2*

have been utilized as a marker for the "GABA switch" (ex. [17]). Our intent was to determine whether

transcriptional patterns recapitulate those described in other species and to gain initial insight into the

developmental timing of the GABA switch in swine.

We began by quantifying mRNA expression of *Slc12a2*, *Slc12a5*, and *Stk39* in the somatosensory cortex,

hippocampus, and thalamus at embryonic day 36 (ED36), ED90, ED105 (sows gestate on average for 115-118

days) postnatal day 7 ("infant"), and postnatal day 30 ("toddler"). As shown in **Figure 1C**, *Slc12a2* expression

remained relatively constant in the hippocampus and thalamus, while cortex levels exhibited a moderate upward

trend over time. The level of *Slc12a2* transcript across biological replicates, regions, and ages was variable; this

complexity may reflect underlying cellular heterogeneity as NKCC1 is expressed in both neurons and non-

neuronal cells [29]. *Slc12a5* expression across all regions remained relatively stable. *Stk39* declined in

hippocampus and thalamus between late gestation and infancy (2-way ANOVA main effect of age p=0.0007,

main effect of region p=0.0008; post-hoc multiple t-test **p<0.01). Taken together, transcriptional analysis was

not sufficient to resolve developmental changes in chloride cotransporters, though the drop in *Stk39* around birth

might indicate a perinatal GABA switch. To gain a clearer understanding, we next turned to protein-level

analyses.

Protein abundances and phosphorylation state for NKCC1, KCC2, and SPAK across development and region in

swine

Representative western blots from randomly selected biological replicates in each group are shown in

Figure 1D. Band densitometry was normalized to the total protein stain, as β-actin, β-III-Tubulin, and GAPDH

were found to be poor loading controls across ages. Specifically, β-actin was more abundant in younger samples,

and β-III-Tubulin and GAPDH were more abundant in older samples. Phosphorylated proteins were probed first,

followed by total proteins, and in the case of NKCC1 and KCC2 exhibiting similar molecular weights, technical

replicate full gels were run, and each sample was normalized to total protein in that gel. For each region, all ages

and biological replicates were performed within the same experiment with a select replicate serving as an internal

loading control for each membrane to account for any additional variance. In **Figure 1E**, total protein abundance

of NKCC1, KCC2, and SPAK (left) and the phosphorylated ratios of those proteins are visualized (right).

Strikingly, levels of NKCC1 remain relatively constant across ages and regions, while SPAK and KCC2 are

significantly upregulated postnatally (KCC2: 2-way ANOVA main effect of age p=0.0002, main effect of region

p=0.0043; post-hoc multiple t-test **p<0.01; SPAK: 2-way ANOVA main effect of age p<0.0001, main effect of

region p-0.008; post-hoc multiple t-test *p<0.05,****p<0.0001). Phosphorylation ratios for SPAK and KCC2 all

decline postnatally (pKCC2/KCC2: 2-way ANOVA main effect of age p=0.0015, main effect of region n.s.; post-

hoc multiple t-test **p<0.01; pSPAK/SPAK: 2-way ANOVA main effect of age p<0.0001, main effect of region

p=0.0104; post-hoc multiple t-test **p<0.01). Phosphorylation ratio for NKCC1 declines perinatally

(pNKCC1/NKCC1: 2-way ANOVA main effect of age p=0.0012, main effect of region p=0.018; post-hoc multiple

t-test p=0.0649). Taken together, the mRNA and protein data suggest that the developmental "GABA switch" in

swine follows a largely perinatal trajectory and both PND7 and 30 piglets have undergone the GABA switch. This

is supported by the general trend for increased total KCC2 and decreased pNKCC1 and pKCC2, pointing toward

a maturation of inhibitory signaling. These findings are consistent with swine occupying a semi-altricial

developmental niche and establishes a high-resolution baseline for these proteins in swine across brain regions.

Given that our transcriptional data alone did not reliably reflect the abundance state of these transporters, this

underscores the importance of assessing phosphorylation status and protein abundance when studying markers

of GABAergic maturation.

With this framework in place, we investigated how these developmental differences might shape the

brain's response to injury. Rather than using a single-insult model like controlled cortical impact (CCI), which

primarily captures focal mechanical injury, we employed a MuLMI TBI model designed to more closely replicate

the complex pathophysiology observed in infants and toddlers with severe traumatic brain injury [27, 30]. This

model has multiple pathoanatomic lesions (mass effect, subdural and subarachnoid hemorrhage) as well as

insults (traumatic seizures and brief apnea and hypoventilation) under non-GABAergic sedation. The MuLMI

model mimics components frequently observed in abusive head trauma [30]. By simulating this multifaceted

injury environment, we aimed to better understand how developmental regulation of chloride transporters might

contribute to age-specific patterns of tissue damage and excitability following trauma.

Creation of multi-factorial, severe TBI injuries and resulting age-specific pathology patterns.

Implementing a MuLMI TBI model in a large animal is resource- and labor-intensive, but essential for

studying complex pediatric brain injuries with high translational fidelity to human pathophysiology. **Figure 2A**

illustrates the model, including the location of injuries, the placement of EEG electrodes for seizure monitoring,

and the timeline of insults, injuries, and intensive care. Following GABA agonist weaning, the pathophysiology

was allowed to evolve over a 24-hour period under intensive care conditions where the tissue damage develops

beyond the area of the cortical impact. The right panel displays the ICU equipment used to sedate, intubate, and

ventilate piglets. Examples of gross pathology (**Figure 2B)** and stylized maps of the damaged regions (yellow,

Figure 2C) are provided, demonstrating that damage extends beyond the site of the cortical impact (arrow). The

total damaged area was greater in piglets subjected to severe TBI injuries compared to sham injuries (Two-way

ANOVA, main effect of treatment p = 0.02, **Figure 2D**). Furthermore, damage distribution exhibited age-

dependent patterns (**Figure 2E**) where the percentage of hemisphere damaged due to TBI injuries was greater

in the ipsilateral hemisphere compared to the contralateral hemisphere in toddlers compared to infants (Two-

way ANOVA, main effect of treatment p=0.05, post-hoc t-test p = 0.0765). These findings highlight a clear age-

specific vulnerability to this severe model of TBI and motivated our desire to understand why toddler piglets might

have more tissue damage, possibly via upregulation of neuronal NKCC1.

TBI-associated shifts of transcription of cation chloride cotransporters were characteristic of reversion

of the GABA switch in toddler piglets

We thus examined the effects of severe TBI on chloride transporter gene expression in piglets at different

developmental stages, focusing on age- and region-specific transcriptional responses in the cortex. The cortex

is where the differential patterns of injury are most obvious (**Figure 3A**). Quantitative PCR revealed *Slc12a2*

(encoding NKCC1) expression was largely unchanged in cortex but exhibited a modest effect of age (**Figure 3B**;

main effect of age p=0.0314). *Slc12a5* (encoding KCC2) was significantly downregulated in the cortex of injured

toddler piglets (**Figure 3C**, main effect of age p=0.0039, main effect of treatment p=0.0636, age x treatment p =

0.0535). Stk39 (encoding SPAK), a kinase that regulates both NKCC1 and KCC2, was significantly upregulated

of toddler piglets with TBI in cortex (**Figure 3D**, main effect of age p=0.0066, main effect of treatment p=0.0686,

age x treatment p=0.0352).

Given that cortical transcriptional effects were most consistent across toddler piglets, we applied GeoMx

spatial transcriptomics to cortical regions from toddler piglets with sham or severe TBI injuries. Regions of interest

(ROIs) were chosen from superficial gray matter (teal) and gyral white matter (orange), as shown in the

schematic (**Figure 3E**). We found that TBI-induced reductions in *Slc12a5* and increases in *Slc12a2* were more

pronounced in white matter than gray matter, specifically in areas of structural damage (**Figure 3F**). These

results suggest that white matter is particularly vulnerable to transcriptional dysregulation of chloride homeostasis following TBI in toddlers.

Taken together, these findings support the hypothesis that severe TBI disrupts chloride transporter mRNA expression in a developmentally and regionally specific manner. The apparent acute (24 hours after injury) shift toward excitatory GABA signaling in surviving toddler neurons may contribute to ongoing tissue vulnerability and highlights a potential window for targeting NKCC1 activity beyond the acute post-injury period.

TBI increases SPAK abundance in the cortex of infant piglets greater than toddler piglets, driven by acute response after injury and independent of hemisphere.

Western blotting of cortical lysates (**Figures 3G, H**, quantified in **3I-J**), rostral and lateral to the area of cortical impact but still damaged, revealed that infant piglets with TBI exhibited relatively unchanged levels of NKCC1 and KCC2 due to TBI injuries, but SPAK protein levels greatly increased with TBI in infant piglets only (**Figure 3I**; Two-way ANOVA effect of treatment p=0.117; post-hoc multiple t-test infant sham v TBI p=0.0834; infant TBI v sham TBI p=0.0338). This effect became more apparent when considering the timeframe injuries were allowed to develop in piglets with TBI injuries (1 hours of injury versus 24 hours), wherein SPAK demonstrated a robust effect of age and time (Two-way ANOVA, time x age effect p=0.0007). This increase in SPAK was mirrored in abundance of TRKB, a kinase receptor known to influence NKCC1 and KCC2 expression [31] (Two-way ANOVA effect of time p=0.0067, time x age p=0.201). There was no effect of hemisphere across any proteins examined (**Figure 3K**). The significant upregulation of SPAK and TRKB in infant piglets with TBI, but not toddler piglets, is interesting for several reasons. 1) Our previous work identified that while both ages demonstrate vasogenic edema due to TBI, toddler piglets demonstrate a greater upregulation of matrix metalloproteinases and more vasogenic edema than infant piglets [28]. Upregulation of both SPAK and TRKB in infant piglets with TBI injuries might be driving evolving cytotoxic edema that does not result in vasogenic edema [13, 32]. 2) SPAK abundance can be promoted by NFkB signaling [33], which could be activated by either the subarachnoid hemorrhage or cortical impact-induced mechanical stress in this model. 3) Upregulation of SPAK and TRKB acutely after injury, but less so as the injuries progress, could profoundly impact cell-type

phosphorylation states of the transporters which are missed at the level of resolution offered by whole tissue

lysates, thus, requiring investigation of phosphorylation state of NKCC1 via immunohistochemistry.

Severe TBI in swine produces region- and cell-type specific alterations in pNKCC1 that are correlated

with damage, subarachnoid hemorrhage, and seizure duration.

At 24 hrs post-injury, TBI significantly increased pNKCC1 expression in non-parvalbumin-expressing

cortical neurons (NeuN$^+$/pNKCC1$^+$/PV$^-$) of infant piglets positively correlated with seizure duration and SAH area,

while cortical PV$^+$ populations and hippocampal neuronal populations showed no changes, indicating region-

and cell-specific effects. We utilized immunohistochemistry to assess phosphorylated NKCC1 (pNKCC1)

expression in NeuN co-labeled mature neurons in the cortex lateral or caudal to the cortical impact (**Figure 4A**),

and each ROIs for each animal are presented in **Figure S1** where the pattern of damage assessed by H&E is

highlighted in yellow (same cohort of piglets as summary data presented in Figure 2). Apoptotic neurons

assessed by nuclear chromatin condensation via Hoechst/NeuN$^+$ across cortical ROIs generally increased in

toddlers but not infants (**Figure 4C-E**), consistent with our previous reports [6, 27]. In cortical ROI 2, the site

caudo-medial to contusion (not the area of impact, but where the neuronal ischemia spreads beyond the

contusion site in this MuLMI mode), TBI increased pNKCC1 expression in NeuN$^+$ neurons, without a preference

for PV$^+$/NeuN$^+$ interneurons (main effect of treatment p=0.0017, **Figure 4C**). When comparing pNKCC1 levels

to % hemispheric damage, **visualized in Figure 4D, far right**, % damage shows a significant correlation with

pNKCC1 in both ages (infant- Spearman r = 0.833; p=0.0154; toddler- Spearman r = 0.943; p=0.0167); however,

only infant piglets demonstrated a positive (non-zero) correlation assessed by linear regression (p=0.0736).

Multivariate analysis further confirmed the relationship between neuronal-pNKCC1 expression caudo-medial to

the cortical impact site and spreading damage, SAH%, and seizure duration as positive predictors across both

ages (**Figure 4F;** SAH x pNKCC1 Spearman r = 0.56; p=0.031; seizure duration x pNKCC1 Spearman r = 0.61,

p=0.018). Further cell-specific analyses indicated that PV$^+$ interneurons did not show enhanced pNKCC1 after

TBI, suggesting that these inhibitory interneurons remain relatively unaffected. Expression of pNKCC1 in non-

neuronal cells in the cortex was not different according to age or injury. In this early time post-injury (24 hrs),

gliosis and re-building of the extracellular matrix are not expected. We have previously demonstrated an age-

dependent increase in matrix metalloproteinase activity as early as 1 hr post-injury with extensive matrix

metalloproteinase activity at 24 hrs, indicating active loss of the extracellular matrix at this time point [28].

Transcriptional, but not translational or post-translational changes to cation chloride cotransporters are

observed in regions with minimal structural damage.

In the hippocampus and thalamus, areas typically spared from hypoxic-ischemic damage in this model

(**Figure 5A**), we again tested the effect of TBI injuries on age-dependent cation-chloride cotransporter

dysregulation. In hippocampus, *Slc12a2* increased following TBI across both ages (**Figure 5B**, main effect of

treatment p=0.0085). Conversely, *Slc12a5* was downregulated with age, but there was no TBI effect (**Figure 5C**,

main effect of age p=0.0001). Stk39 exhibited a significant upregulation due to TBI injuries in the toddler piglets

(**Figure 5D**, main effect of age p=0.0104, main effect of treatment p=0.0176, age x treatment p=0.0097). In

thalamus, *Slc12a2* was significantly downregulated in the thalamus in both age groups (**Figure 5E**, main effect

of treatment p=0.0002). *Slc12a5* exhibited a significant age and TBI effect, with infant piglets with TBI

upregulating *Slc12a5* and toddler piglets with TBI downregulating *Slc12a5* (**Figure 5F**, main effect of age

p=0.0038, main effect of treatment p=0.9538, age x treatment p=0.0293). Stk39 demonstrated a modest effect

of TBI, but not by age (**Figure 5G**).

We observed minimal changes to NKCC1, KCC2, and SPAK in infant TBI piglets compared to toddler

TBI piglets (**Figure 5H-J**). Therefore, regions not displaying overt injury can exhibit early changes in chloride

transporter mRNA expression following TBI, but this may not be reflected in protein abundance or

phosphorylation within this acute window. Further, structural damage might require additional time to manifest

beyond the 24-hour paradigm of our study.

To be thorough in our investigation, we also employed immunohistochemistry across the dorso-ventral

axis of the hippocampus (**Figure S2A-B**). Intriguingly, we observed pNKCC1 expression across postnatal ages

in granule cells, neurons and glia of the polymorph layer, and neurons throughout the cornu ammonius, which

seems to be different than expression typically reported in rodents. In all hippocampal regions, less than 5% of

total neurons were apoptotic (**Figure S2C-F**). There was only an effect of injury on neuronal-pNKCC1 expression

in the dorsal dentate gyrus (**Figure S2C**), but not in the dorsal CA2 or ventral DG or CA2 (**Figure S2D-F**).

However, unlike in cortex, the effect on neuronal-pNKCC1 reflected a decrease in expression, rather than an

increase. This likely reflects a downregulation of excitatory transduction proximal to the damage spreading

distant from the cortical impact, perhaps as a compensatory mechanism.

Collectively, these data suggest that TBI-induced elevations in pNKCC1 predominantly occur in surviving

cortical PV[-] neurons of younger piglets, possibly contributing to a pro-excitatory state associated with seizures.

These observations provide evidence that TBI influences cation chloride transporter expression and activation

differently across brain regions and developmental stages. Younger piglets appear more prone to TBI-related

shifts in these key proteins, raising the possibility that early therapeutic interventions targeting chloride

cotransporters could be beneficial. An alternative possibility is that these changes were not detected in toddler

piglets due to advanced tissue damage, but we attempted to resolve this with cell-specific expression in apoptotic

vs. healthy cells.

Postnatal pNKCC1 downregulation in human hippocampus.

To compare these developmental patterns with those observed in humans, we examined hippocampal

samples from infants with known causes of unexpected death other than from TBI to determine developmental

changes around the time of birth. The study design and tissue validation steps are outlined in **Figure 6A.**

Previous transcriptional data from qPCR and microarray studies suggest that the GABA switch in humans occurs

around birth in both the limbic system and neocortex [17, 34, 35]. Furthermore, evidence from slice

electrophysiology, western blot analysis, and immunolabeling indicates hyperpolarizing GABA activity in the adult

temporal lobe cortex and hippocampus [36, 37], which is altered in epilepsies [38].

We did not find a significant effect of age on the abundance of mRNA for *Slc12a2, Slc12a5*, and *Stk39*

in relation to adjusted postconceptional age (PCA) [gestational age plus postnatal age]. As shown in **Figure 6B**,

transcriptional levels of these genes remained relatively stable across postnatal ages, and no positive correlation

emerged between the ratio of *Slc12a5: Slc12a2* and PCA (**Figure 6C**). These transcriptional results indicate that

major transcriptional shifts in these transporters are minimal during the postnatal period in humans similar to

development in piglets.

We then assessed protein abundances and phosphorylation states in these hippocampal samples

(**Figure 6D**). Our study offers a more detailed resolution and analysis of the activity state (via phosphorylation)

of these cotransporters during the immediate postnatal period in infants with normal brain development. Total

protein abundances for NKCC1, KCC2, and SPAK all increased with PCA (**Figure 6E**; linear regression NKCC1

R^2 = 0.7019, p=0.0025 non-zero; KCC2 linear regression R^2 = 0.5247, p=0.0178 non-zero; SPAK linear

regression R^2 = 0.5953, p=0.0089 non-zero). An assessment of the ratio of active KCC2:NKCC1 (KCC2 /

(pNKCC1/NKCC1) increased postnatally (linear regression R^2=0.5610, p=0.0127 non-zero, **Figure 6F**), like

swine. Postnatal KCC2 dominance aligns with the gradual shift toward inhibitory GABA signaling that

characterizes brain maturation [9, 16, 17, 39, 40].

Upregulation of pNKCC1 expression is associated with tissue damage in autopsy cases of pediatric TBI.

To bridge our findings from the piglet model to humans, we examined pNKCC1 expression in autopsy

samples from non-TBI and TBI cases covering a range of ages (**Figure 7**). In non-TBI "control" human tissue,

pNKCC1 was consistently present at low to moderate levels in neurons and glia across multiple regions (**Figure**

7A). In TBI cases, pNKCC1 expression varied according to injury severity and location. Regions with significant

hypoxic-ischemic damage (the same type of tissue damage we model here), including the parietal lobe, corpus

callosum, and hippocampus, showed increased pNKCC1 in glia and axons relative to controls (**Figure 7B, 7C**).

The temporal lobe, especially the ependyma of the striatum terminalis, exhibited pronounced subependymal

pNKCC1 upregulation in cases with severe white matter injuries.

DISCUSSION:

A central feature of normal brain development is the "GABA switch," in which signaling becomes

predominantly hyperpolarizing because of a change in the direction of chloride flux from outward (depolarizing)

to inward (hyperpolarizing). This switch is correlated with shifts in NKCC1 and KCC2 expression, although a

myriad of other factors affects the concentration of chloride such as impermeant anion moieties [41]. Our swine

data show that this switch occurs around the perinatal period in a pattern more closely resembling semi-altricial

species (including primates) than altricial rodents. In humans, we similarly observed a gradual postnatal

decrease in NKCC1 phosphorylation aligned with a relative increase in potential inhibitory tone over time.

Together, these findings suggest that swine provide a useful experimental bridge for modeling human

neurodevelopment and the timing of the GABA switch.

The standard of care for seizures in infants and toddlers after TBI is attempting to stop seizures with

GABA agonists, even though it is not known if seizures drive pathophysiology or are a symptom of TBI. Data on

seizure incidence are confounded with TBI severity: worse brain injury results in more seizures. Furthermore,

stopping seizures is difficult in infants with both levetiracetam and phenobarbital only working half the time [42].

We hypothesized that regions of tissue damage (i.e. cerebral cortex) would exhibit changes to the NKCC1 and

KCC2 and regions without tissue damage (hippocampus and thalamus) would not, which clinically, would be

expected to make stopping seizures even more difficult. We did find upregulation of neuronal pNKCC1 in the

cortex following TBI injuries as predicted.

We further hypothesized that toddler piglets with more pervasive, hemispheric spreading injury with more

pronounced vasogenic edema would exhibit upregulated NKCC1 compared to infant piglets with patchy, less

tissue damage. However, we found the opposite to be true where greater upregulation of SPAK in response to

injury was not in the age with greater tissue damage. Younger animals displayed a more pronounced TBI-

induced upregulation of SPAK, a kinase that promotes NKCC1 phosphorylation and activity. This shift toward

elevated NKCC1 function can increase neuronal excitability and seizure risk—clinical challenges frequently

observed in pediatric TBI. This is perhaps because apoptotic (dead) neurons, possibly due to cytotoxic edema,

which are abundant in the injured toddler piglets, are non-functional and will not be affected by changes to the

Cl^- milieu, whereas surviving neurons might exhibit ongoing dysfunction. The correlation between pNKCC1

counts and seizure duration and hemorrhage in piglets further reinforces the notion that disrupted chloride

regulation might contribute to post-traumatic neuronal excitation. Notably, these TBI-induced changes were not

uniform across cell types or regions: parvalbumin-positive cortical interneurons were spared, and hippocampal

neurons remained relatively unaffected by these specific injuries. Spatial transcriptomics indicated that white

matter alterations in NKCC1 and KCC2 expression likely involve glial cells and support matrix, influencing

extracellular ion concentration [41] and potentially contributing to edema and barrier dysfunction. Indeed, we

previously demonstrated that toddlers, which have the most damage, also have ongoing upregulation in abundance and activity of matrix metalloproteinase, whereas matrix metalloproteinases are either not increased or even downregulated in infants, who have less damage [28]. This work supports the hypothesis that chloride transporters alone do not control intraneuronal chloride concentrations, especially after pediatric TBI [41, 43]. Human autopsy data corroborated these animal findings. In TBI cases, we saw similar heterogeneous region- and cell-type-specific shifts in pNKCC1, particularly in damaged white matter tracts.

An alternative hypothesis is that reversion back to NKCC1, a biomarker for increased intraneuronal chloride, is protective in PND7 piglets, who demonstrate less tissue damage, and perhaps at the expense of future hyper-excitability and development of post-traumatic epilepsy. Indeed, the only way to determine cause vs. effect in the developmental differences in response to injury is to provide therapies that would reduce MMP upregulation such as matrix metalloproteinase inhibitors and/or NKCC1 an antagonist such as bumetanide. Another level of complexity is that NKCC1 expression in non-neuronal cells have been shown to mediate cell death and vasogenic edema in models of ischemia (reviewed by [44]). Here, the differential neuronal-specific NKCC1 increase after severe TBI was not detectable via Western blots of mixed cell lysates.

Taken together, after TBI, toddler piglets display vasogenic edema likely from MMP upregulation, while infant piglets might instead experience acute cytotoxic edema with rapid, clean cell death without leading to vasogenic edema, resulting in ongoing neuronal dysfunction in surviving neurons, which might make them unique candidates for therapeutic intervention [45]. This postulate is supported by Level III clinical evidence and recent age-dependent mechanisms of cerebral edema in small-diameter blood vessels following severe TBI in an autopsy case-series investigation [46]. Given the robust age- and treatment-effects of TBI injuries on SPAK abundance, we propose that treatment of our infant piglets with a SPAK inhibitor immediately following TBI might offer protective and antiepileptic effects [47, 48]. Indeed, as pathophysiology is age-specific, successful therapeutics might also need to be age-specific after severe TBI according to developmental stage: targeting NKCC1 or SPAK in infants and targeting MMP's in toddlers.

Limitations:

Our study has several limitations that should be acknowledged. Firstly, we were not powered to examine

the distinct effect of sex as a covariate on NKCC1 and KCC2 levels in the developing swine brain due to protocol

limitations, which may influence the observed outcomes [49]. Additionally, we did not assess the levels of brain-

derived neurotrophic factor (BDNF) or insulin-like growth factor (IGF), both of which are known to promote the

GABA switch. Other intracellular regulators such as protein kinase C (PKC), Src kinases, and with no-lysine

(WNK) kinases, the latter of which are chloride-sensitive and may play a role in the switch, were also not

considered in this study [48, 50]. Furthermore, we did not investigate the potential contributions of maternal

estrogen or taurine as positive sensory regulators of the GABA switch. Finally, and perhaps most importantly,

we did not perform electrophysiology or chloride ion imaging either *in vivo* or in organotypic slice, thus, our

observations are limited to when the switch is *likely* occurring, however, we cannot say with certainty when GABA

switches from excitatory to inhibitory in the developing swine brain.

Our study, while leveraging a translationally relevant swine model and human autopsy samples, cannot

fully capture the clinical complexity and heterogeneity of pediatric TBI. The standardized injury approach in pigs,

though informative, may not reflect the multifaceted trauma mechanisms, variable interventions, and long

recovery trajectories seen in patients. These factors represent important avenues for future research to better

understand the mechanisms underlying the changes in excitation and inhibition in the developing swine brain

and in the context of TBI. Donnan effects with the loss of the extracellular matrix also likely affect the amount of

extraneuronal and intraneuronal chloride concentrations post-TBI as demonstrated in rodent hippocampal

explants [41, 43]. Indeed, in our own severe TBI model, we have described an age-dependent upregulation of

matrix metalloproteinases where there was an ongoing upregulation in toddlers with worse injury, compared to

a resolution in infants with lower injury burden scaled to brain volume [28]. The human autopsy samples, while

invaluable, represent end-stage conditions and lack information on the temporal progression of TBI pathology.

Further, it is unclear to what extent non-traumatic pediatric autopsy samples are representative of a

physiologically normal pediatric population.

We observed notable variability in transcript-level measurements, likely due to biological heterogeneity

and technical factors such as RNA quality, sampling consistency, and cellular composition of lysates. Though

immunoblotting analyses were particularly informative, phosphorylation state was less informative in swine. It is

important to note that our study did not evaluate additional post-translational modifications, such as glycosylation,

though these are certainly indicated in the activity of these transporters. Finally, while we identified potential

mechanistic links between chloride transporter dysregulation, seizures, and injury severity, further studies are

needed to establish causal relationships and to explore how interventions that target these pathways might

improve long-term outcomes in pediatric TBI.

Conclusions and future directions:

By integrating findings from a swine model, whose brain development and structure closely parallel those

of humans, with postmortem human tissue analyses, our work supports the translational value of targeting

chloride regulation to improve TBI outcomes, especially in pediatric populations [13, 51, 52]. Pharmacologically

modulating NKCC1 or KCC2, potentially with agents like a SPAK inhibitor, could help restore chloride

homeostasis and reduce hyperexcitability and swelling/tissue shifts after pediatric TBI. Such treatments would

need to consider the patient's developmental stage and the specific brain regions affected. Early intervention,

when the brain is most vulnerable to ionic changes that might exacerbate tissue damage, may be especially

beneficial. Our findings emphasize that TBI's impact on the developing brain cannot be fully understood by

focusing solely on neuronal excitability or a single brain area. Instead, we must consider a dynamic interplay

among developmentally regulated ion transporters, glial function, neuroinflammatory processes, and structural

integrity. Future work should address 1) how these factors interact over longer recovery periods, potentially via

in vivo monitoring of neuronal chloride [53], 2) explore targeted therapies that normalize chloride concentrations,

and 3) consider how interventions might influence long-term neurodevelopmental outcomes and post-traumatic

sequelae such as epilepsy.

600
601

References:
1. Keenan HT, Runyan DK, Marshall SW, Nocera MA, Merten DF, Sinal SH. (2003). A population-based study of inflicted traumatic brain injury in young children. JAMA. Aug 6;290(5):621-6.
2. Clancy B, Finlay BL, Darlington RB, Anand KJ. (2007). Extrapolating brain development from experimental species to humans. Neurotoxicology. Sep;28(5):931-7.
3. Lind NM, Moustgaard A, Jelsing J, Vajta G, Cumming P, Hansen AK. (2007). The use of pigs in neuroscience: modeling brain disorders. Neurosci Biobehav Rev. 31(5):728-51.
4. Swindle MM, Smith AC, Hepburn BJ. (1988). Swine as models in experimental surgery. J Invest Surg. 1(1):65-79.
5. Dobbing J and Sands J. (1979) Comparative aspects of the brain growth spurt. Early Human Development. Vol. 311 Pages 79-83.
6. Costine-Bartell BA, McGuone D, Price G, Crawford E, Keeley KL, Munoz-Pareja J, Dodge CP, Staley K, Duhaime AC. (2019). Development of a Model of Hemispheric Hypodensity ("Big Black Brain"). J Neurotrauma. Mar 1;36(5):815-833.
7. Duhaime AC, Durham S. (2007) Traumatic brain injury in infants: the phenomenon of subdural hemorrhage with hemispheric hypodensity ("Big Black Brain") Prog Brain Res. 161:293-302.
8. Foster KA, Recker M, Lee P, Bell MJ, Tyler-Kabara E. (2014) Factors Associated with Hemispheric Hypodensity after Subdural Hematoma following Abusive Head Trauma in Children. J Neurotrauma. 2014 Oct 1;31(19):1625-31.
9. Ben-Ari Y. (2014). The GABA excitatory/inhibitory developmental sequence: a personal journey. Neuroscience. Oct 24; 279:187-219.
10. LoTurco JJ, Owens DF, Heath MJ, Davis MB, Kriegstein AR. (1995). GABA and glutamate depolarize cortical progenitor cells and inhibit DNA synthesis. Neuron. Dec;15(6):1287-98.
11. Lu KT, Cheng NC, Wu CY, Yang YL. NKCC1-mediated traumatic brain injury-induced brain edema and neuron death via Raf/MEK/MAPK cascade. Crit Care Med. 2008 Mar;36(3):917-22. doi: 10.1097/CCM.0B013E31816590C4. PMID: 18431281.
12. Dzhala V, Valeeva G, Glykys J, Khazipov R, Staley K. Traumatic alterations in GABA signaling disrupt hippocampal network activity in the developing brain. J Neurosci. 2012 Mar 21;32(12):4017-31. doi: 10.1523/JNEUROSCI.5139-11.2012. PMID: 22442068; PMCID: PMC3333790.
13. Glykys J, Dzhala V, Egawa K, Kahle KT, Delpire E, Staley K. (2017) Chloride Dysregulation, Seizures, and Cerebral Edema: A Relationship with Therapeutic Potential. Trends Neurosci. May;40(5):276-294.
14. Carney N, Totten AM, O'Reilly C, Ullman J, Hawryluck GW, Bell MJ, Bratton SL, Chestnut R, Harriss OA, Kissoon NK, Rubiano AM, Shutter L, Tasker RC, Vavilala MS, Wiberger J, Wright DW, Ghajar J. Guidelines for the Management of Severe Traumatic Brain Injury, 4th Edition. (2016) Brain Trauma Foundation.
15. Kochanek PM, Tasker RC, Carney N, Totten AM, Adelson PD, Selden NR, Davis-O'Reilly CD, Hart EL, Bell MJ, Bratton SL, Grant GA, Kissoo M, Reuter-Rice KE, Vavilala MS, Wainwright MS. 2019 Guidelines for the Management of Pediatric Severe Traumatic Brain Injury, Third Edition: Update of the Brain Trauma Foundation Guidelines, Executive Summary. Neurosrugery 84 (6): 1169-1178.
16. Khazipov R, Esclapez M, Caillard O, Bernard C, Khalilov I, Tyzio R, Hirsch J, Dzhala V, Berger B, Ben-Ari Y. (2001) Early development of neuronal activity in the primate hippocampus in utero. J Neurosci. Dec 15;21(24):9770-81.
17. Sedmak G, Jovanov-Milošević N, Puskarjov M, Ulamec M, Krušlin B, Kaila K, Judaš M. (2016) Developmental Expression Patterns of KCC2 and Functionally Associated Molecules in the Human Brain. Cereb Cortex. Dec;26(12):4574-4589.
18. Tang BL. (2020) The Expanding Therapeutic Potential of Neuronal KCC2. Cells. Jan 17;9(1):240.
19. Peerboom C, Wierenga CJ. (2021). The postnatal GABA shift: A developmental perspective. Neurosci Biobehav Rev. May;124:179-192.
20. Young JW, Mayerl CJ, Mannava A, Lewis C, Fan T, Nair M, Mamone C, Schapker NM, Mossor AM, German RZ. (2023) Evaluating The Precocial-altricial Axis of Motor Skill at Birth in A Preterm Pig Model. Integr Comp Biol. Sep 15;63(3):625-640.
21. Bienboire-Frosini C, Marcet-Rius M, Orihuela A, Domínguez-Oliva A, Mora-Medina P, Olmos-Hernández A, Casas-Alvarado A, Mota-Rojas D. (2023) Mother-Young Bonding: Neurobiological Aspects and Maternal Biochemical Signaling in Altricial Domesticated Mammals. Animals (Basel). Feb 2;13(3):532.
22. Goldstein RD, Poduri A. (2024) Seizures and Sudden Death Beyond SUDEP. Neurology. Feb 13;102(3):e208119.
23. Leitner DF, William C, Faustin A, Askenazi M, Kanshin E, Snuderl M, McGuone D, Wisniewski T, Ueberheide B, Gould L, Devinsky O. (2022) Proteomic differences in hippocampus and cortex of sudden unexplained death in childhood. Acta Neuropathol. May;143(5):585-599.
24. Harowitz J, Crandall L, McGuone D, Devinsky O. (2021) Seizure-related deaths in children: The expanding spectrum. Epilepsia. Mar;62(3):570-582.

25. Haynes RL, Kinney HC, Haas EA, Duncan JR, Riehs M, Trachtenberg F, Armstrong DD, Alexandrescu S, Cryan JB, Hefti MM, Krous HF, Goldstein RD, Sleeper LA. (2021) Medullary Serotonergic Binding Deficits and Hippocampal Abnormalities in Sudden Infant Death Syndrome: One or Two Entities? Front Pediatr. Dec 21;9:762017.

26. Patodia S, Somani A, Thom M. (2021) Review: Neuropathology findings in autonomic brain regions in SUDEP and future research directions. Auton Neurosci. Nov;235:102862.

27. Costine-Bartell B, Price G, Shen J, McGuone D, Staley K, Duhaime AC. (2021). A perfect storm: The distribution of tissue damage depends on seizure duration, hemorrhage, and developmental stage in a gyrencephalic, multi-factorial, severe traumatic brain injury model. Neurobiol Dis. Jul;154:105334.

28. Hochstetler A, Price G, Baohan A, Li M, Rodriguez Lara F, Lok J, Costine-Bartell B. (2024). Developmental Regulation of Matrix Metalloproteinases in Response to Multifactorial, Severe Traumatic Brain Injuries during Immaturity. Dev Neurosci. 46(5):319-332.

29. Kurki SN, Uvarov P, Pospelov AS, Trontti K, Hübner AK, Srinivasan R, Watanabe M, Hovatta I, Hübner CA, Kaila K, Virtanen MA. (2023). Expression patterns of NKCC1 in neurons and non-neuronal cells during cortico-hippocampal development. Cereb Cortex. May 9;33(10):5906-5923.

30. Ferguson NM, Rebsamen S, Field AS, Guerrero JM, Rosario BL, Broman AT, Rathouz PJ, Bell MJ, Alexander AL, Ferrazzano PA on behalf of the ADAPT MRI Investigators. (2022) Magnetic Resonance Imaging Findings in Infants with Severe Traumatic Brain Injury and Associations with Abusive Head Trauma. Children 9(7): 1092.

31. Szymanski J, Minichiello L. NKCC1 Deficiency in Forming Hippocampal Circuits Triggers Neurodevelopmental Disorder: Role of BDNF-TrkB Signalling. Brain Sci. 2022 Apr 15;12(4):502. doi: 10.3390/brainsci12040502. PMID: 35448033; PMCID: PMC9030861.

32. Zhang J, Bhuiyan MIH, Zhang T, Karimy JK, Wu Z, Fiesler VM, Zhang J, Huang H, Hasan MN, Skrzypiec AE, Mucha M, Duran D, Huang W, Pawlak R, Foley LM, Hitchens TK, Minnigh MB, Poloyac SM, Alper SL, Molyneaux BJ, Trevelyan AJ, Kahle KT, Sun D, Deng X. (2020) Modulation of brain cation-Cl- cotransport via the SPAK kinase inhibitor ZT-1a. Nat Commun. Jan 7;11(1):78.

33. Bhuiyan MIH, Young CB, Jahan I, Hasan MN, Fischer S, Meor Azlan NF, Liu M, Chattopadhyay A, Huang H, Kahle KT, Zhang J, Poloyac SM, Molyneaux BJ, Straub AC, Deng X, Gomez D, Sun D. (2022) NF-κB Signaling-Mediated Activation of WNK-SPAK-NKCC1 Cascade in Worsened Stroke Outcomes of Ang II-Hypertensive Mice. Stroke. May;53(5):1720-1734.

34. Tao R, Li C, Newburn EN, Ye T, Lipska BK, Herman MM, Weinberger DR, Kleinman JE, Hyde TM. (2012) Transcript-specific associations of SLC12A5 (KCC2) in human prefrontal cortex with development, schizophrenia, and affective disorders. J Neurosci. Apr 11;32(15):5216-22.

35. Hyde TM, Lipska BK, Ali T, Mathew SV, Law AJ, Metitiri OE, Straub RE, Ye T, Colantuoni C, Herman MM, Bigelow LB, Weinberger DR, Kleinman JE. (2011) Expression of GABA signaling molecules KCC2, NKCC1, and GAD1 in cortical development and schizophrenia. J Neurosci. Jul 27;31(30):11088-95.

36. Wang WZ, Hoerder-Suabedissen A, Oeschger FM, Bayatti N, Ip BK, Lindsay S, Supramaniam V, Srinivasan L, Rutherford M, Møllgård K, Clowry GJ, Molnár Z. (2010) Subplate in the developing cortex of mouse and human. J Anat. Oct;217(4):368-80.

37. Muñoz A, Méndez P, DeFelipe J, Alvarez-Leefmans FJ. (2007) Cation-chloride cotransporters and GABA-ergic innervation in the human epileptic hippocampus. Epilepsia. Apr;48(4):663-73.

38. Huberfeld G, Blauwblomme T, Miles R. (2015) Hippocampus and epilepsy: Findings from human tissues. Rev Neurol (Paris). Mar;171(3):236-51.

39. Watanabe M, Zhang J, Mansuri MS, Duan J, Karimy JK, Delpire E, Alper SL, Lifton RP, Fukuda A, Kahle KT. (2019). Developmentally regulated KCC2 phosphorylation is essential for dynamic GABA-mediated inhibition and survival. Sci Signal. Oct 15;12(603): eaaw9315.

40. Kahle KT, Deeb TZ, Puskarjov M, Silayeva L, Liang B, Kaila K, Moss SJ. (2013) Modulation of neuronal activity by phosphorylation of the K-Cl cotransporter KCC2. Trends Neurosci. Dec;36(12):726-737.

41. Glykys J, Dzhala V, Egawa K, Balena T, Saponjian Y, Kuchibhotla KV, Bacskai BJ, Kahle KT, Zeuthen T, Staley KJ. (2014) Local impermeant anions establish the neuronal chloride concentration. Science. Feb 7;343(6171):670-5.

42. Qiao, MY, Cui HT, Zhao LZ, Miao JK, and Chen QX. 2021. Efficacy and Safety of Levetiracetam vs. Phenobarbital for Neonatal Seizures: A Systematic Review and Meta-Analysis. Front Neurol 12: 747745.

43. Normoyle KP, Lillis KP, Egawa K, McNally MA, Paulchakrabarti M, Coudhury BP, Lau L, Shiu FH, Staley KJ. (2024) Displacement of extracellular chloride by immobile anionic constituents of the brain's extracellular matrix. J Physiol. Dec 2.

44. Vernantin MA, Uvarov P, Hubner CA, Kaila K. (2020) NKCC1, an Elusive Molecular Target in Brain Development: Making Sense of the Existing Data. Cell 9(12):2607.

45. Balena T, Staley K. (2024). Neuronal Death: Now You See It, Now You Don't. Neuroscientist. Sep 24:10738584241282632.

46. Fullerton JL, Hay J, Bryant-Craig C, Atkinson J, Smith DH, Stewart W. (2024) Pediatric Traumatic Brain Injury and
Microvascular Blood-Brain Barrier Pathology. JAMA Netw Open. Nov 4;7(11):e2446767.
47. Gong Y, Wu M, Gao F, Shi M, Gu H, Gao R, Dang BQ, Chen G. (2021) Inhibition of the p-SPAK/p-NKCC1 signaling
pathway protects the blood-brain barrier and reduces neuronal apoptosis in a rat model of surgical brain injury. Mol Med
Rep. Oct;24(4):717.
48. Wu D, Lai N, Deng R, Liang T, Pan P, Yuan G, Li X, Li H, Shen H, Wang Z, Chen G. (2020) Activated WNK3 induced
by intracerebral hemorrhage deteriorates brain injury may be via WNK3/SPAK/NKCC1 pathway. Exp Neurol.
Oct;332:113386.
49. Missios S, Harris BT, Dodge CP, Simoni MK, Costine BA, Lee YL, Quebada PB, Hillier SC, Adams LB, Duhaime AC.
(2009). Scaled cortical impact in immature swine: effect of age and gender on lesion volume. J Neurotrauma.
Nov;26(11):1943-51.
50. Watanabe M, Fukuda A. (2015) Development and regulation of chloride homeostasis in the central nervous system.
Front Cell Neurosci. Sep 24;9:371.
51. Soul JS, Bergin AM, Stopp C, Hayes B, Singh A, Fortuno CR, O'Reilly D, Krishnamoorthy K, Jensen FE, Rofeberg V,
Dong M, Vinks AA, Wypij D, Staley KJ; Boston Bumetanide Trial Group. (2021) A Pilot Randomized, Controlled, Double-
Blind Trial of Bumetanide to Treat Neonatal Seizures. Ann Neurol. Feb;89(2):327-340.
52. Staley KJ. (2022) Clarifications regarding bumetanide for neonatal seizures. Epilepsia. Jul;63(7):1863-1867. doi:
10.1111/epi.17278. Epub 2022 May 26.
53. Costine-Bartell BA, Martinez-Ramirez L, Normoyle K, Stinson T, Staley KJ, Lillis KP. (2023). 2-photon imaging of
fluorescent proteins in living swine. Sci Rep. Aug 29;13(1):14158.

Figure 1:

736
Figure 1: Proxy markers of transition to inhibitory GABA increase perinatally in swine, particularly across cortex.
A, schematic of samples represented across gestation and postnatal ages. *B*, regions collected for analysis. *C,* quantitative PCR analysis
of samples across ages (ED36, ED90, ED105, Infant, Toddler) and regions (cortex, hippocampus, thalamus) for *Slc12a2* (encoding
NKCC1), *Slc12a5* (encoding KCC2), and *Stk39* (encoding SPAK). 2-way ANOVA for *Stk39* expression across ages and regions: main
effect of age p=0.0007, main effect of region p=0.0008; post-hoc multiple t-test p<0.01. D, representative western blots for protein and
phospho protein abundance, total protein stain as loading control. E, quantitation of western lots for samples across ages and regions for
total protein (left) and ratio of phosphorylated to total protein (right). KCC2: 2-way ANOVA main effect of age p=0.0002, main effect of

region p=0.0043; post-hoc multiple t-test **p<0.01. SPAK: 2-way ANOVA main effect of age p<0.0001, main effect of region p-0.008;
post-hoc multiple t-test *p<0.05,****p<0.0001. pKCC2/KCC2: 2-way ANOVA main effect of age p=0.0015, main effect of region n.s.; post-
hoc multiple t-test **p<0.01. pSPAK/SPAK: 2-way ANOVA main effect of age p<0.0001, main effect of region p=0.0104; post-hoc multiple
t-test **p<0.01. pNKCC1/NKCC1: 2-way ANOVA main effect of age p=0.0012, main effect of region p=0.018; post-hoc multiple t-test
p=0.0649.
749

Figure 2:

Figure 2: Multi-factorial, severe TBI piglet model injuries result in age-specific pathology patterns.
***A*, Left:** The array of injuries displaying the location of various injuries and insults per hemisphere and the electrode EEG strips. **Center:** The timeline of the injuries and insults performed after weaned from GABA agonists, along with intensive care, lasting approximately 24 hours to allow the injury to evolve. **Right:** Photograph of our ICU equipment where piglets are sedated, intubated, and ventilated. ***B*,** 5 mm brain slabs (right is right), and the subarachnoid hemorrhage can be seen along with dusky tissue and a swollen right hemisphere. ***C*,** Maps displaying the damaged areas (yellow) rostral to caudal and site of cortical impact (arrow). Damage spreads beyond the site of the cortical impact as we now hypothesize that seizures and subarachnoid hemorrhage synergistically drive tissue damage. ***D*,** In the cohort studied here, the amount of damage was greater in piglets receiving severe TBI injuries in both infants and toddlers (p=0.02). ***E*,** The percentage of the hemisphere damaged (microscopically determined in one section per 5 mm slab and mapped) exhibited age-dependent patterns where the ipsilateral hemisphere had greater damage than the contralateral hemisphere in toddlers (p=0.05).

764
765 **Figure 3:**

767 **Figure 3: Severe TBI induces a reversion of markers of immature GABA in "infants" and upregulation of**
768 **maturation in "toddlers".**
769 Representative patterns of hypoxic-ischemic injury at the level of the somatosensory cortex between ages (***A***, yellow = injured tissue).
770 Quantitative PCR analysis for transcripts *Slc12a2 (**B**)*, *Slc12a5 (**C**)*, and *Stk39 (**D**)* due to age and TBI injuries in piglets, presented as
771 fold change from sham infant piglets. Two-way ANOVA results are below graph, and post-hoc multiple t-tests are included on graphs
772 where applicable (*$p < 0.05$, **$p < 0.01$). ***E***, Study design and ROI selection for GeoMx Spatial Transcriptomics. ***F***, selected relative
773 expression of *Slc12a2* and *Slc12a5* across regions from sham animals and animals with severely injured tissue. Three-Way ANOVA
774 performed to analyze transcriptional levels across gene, region, and injury. Western blots from cortical lysates in infant (G) and toddler
775 (H) piglets probed for pNKCC1, NKCC1, TrkB, SPAK, pKCC2, and KCC2, normalized to beta actin, quantified by band densiometry and
776 graphed with regard to sham v. TBI (I), time for TBI injuries (J), and hemisphere for TBI injuries (K). Two-way ANOVA results are below
777 graph, and post-hoc multiple t-tests are included on graphs where applicable with exact p values.

778 **Figure 4:**

Figure 4: Cortical non-apoptotic, PV⁻ neurons upregulate pNKCC1 caudo-medial to contusion site in TBI-induced spreading hypoxic-ischemic injury and are associated with burden of tissue damage across ages and burden of subarachnoid hemorrhage and seizure duration in infant piglets with TBI.

Schematic of ROIs for analysis in proximity to contusion site, *A*. Immunohistochemical staining for pNKCC1 (red), parvalbumin (green), NeuN (magenta), and nuclei (white), *B*. Cell counts expressed as percentages per ROI, with ROI 1 in *C*, ROI 2 in *D*, and ROI 3 in *E*. In all cases, the **left** dot plot presents % of neurons that were apoptotic, the **middle** dot plot presents the % of healthy neurons expressing pNKCC1, and the **right** dot plot presents % of healthy neurons co-expressing parvalbumin and pNKCC1. The scatter plot with spearman correlation on **far right** examines whether pNKCC1 neuronal expression correlates with damage. Two-way ANOVA with post-hoc multiple t tests employed for all comparisons and significant findings for main effect of age or treatment reported in graphs. Since a significant correlation with damage was found in region 2, *F* shows the multivariate analysis considering seizure duration, subarachnoid hemorrhage %, and damage % with Spearman correlation matrix presented and significant interactions marked with asterisk scheme, ** $p<0.01$, * $p<0.05$.

Figure 5:

794
Figure 5: Protein expression of NKCC1, KCC2, and SPAK in the hippocampus and thalamus were minimally
sensitive to severe TBI injuries in piglets despite persistent transcriptional changes.
Representative patterns of minimal hypoxic-ischemic injury at the levels of the hippocampus and thalamus between ages (*A*, yellow =
injured tissue). Quantitative PCR analysis for transcripts *Slc12a2 (B, E)*, *Slc12a5 (C, F)*, and *Stk39 (D, G)* due to age and TBI injuries in

piglets, presented as fold change from sham infant piglets in hippocampus (**B-D**) and thalamus (**E-G**). Two-way ANOVA results are below
graph, and post-hoc multiple t-tests are included on graphs where applicable (*p<0.05, **p<0.01). Western blots for all biological replicates
for infant and toddler piglets receiving sham or severe TBI injuries in hippocampus and thalamus (**H**) and quantitated below (**I, J**).
802

803 **Figure 6:**

804
805
Figure 6: Postnatal GABAergic maturation occurs in human hippocampus.
Schematic of study design, sample collection protocol, and sample demographics (**A**). Quantitative PCR analysis for transcripts *Slc12a2,*
Slc12a5, and *Stk39 (**B**)* across postconceptional age (PCA) presented as fold change from youngest sample. Ratio of Slc12a5 to Slc12a2
as a function of PCA (**C**). Representative western blots of hippocampal lysates (**D**), quantified in (**E**) as a function of PCA. SPAK, NKCC1,
and KCC2 all demonstrate positive linear correlations with PCA. Ratio of KCC2 : (pNKCC1/NKCC1) (**F**) also demonstrates a significant
positive correlation with PCA. In all graphs, linear regression analysis was used. R^2 values are shown on the graphs along with a p-value
for whether the slope is non-zero.
813
814
815

816 **Figure 7:**
817

818
819
820 **Figure 7: TBI induces heterogeneous cellular and regional changes to pNKCC1 expression and localization.**
821 Representative pNKCC1 staining patterns throughout human brain (***A***), semiquantitated as an effect of cell type (***B***) and cell type x region
822 (***C***). Two-Way ANOVA in (***B***) with effects listed below graph.
823

824
825

Table 1: Human Sample Demographics

	Sample IDs	PNA (GA)	Sex	Diagnosis	PMI (hr)
Frozen/FFPE Hippocampus (Figure 2)	1	0.4 (40)	Female	Fatty acid oxidation disorder	15
	2	0 (41)	Female	Neonatal bradycardia with arrest and meconium aspiration	12.8
	3	17 (26.5)	Female	Acute asphyxiation	9
	4	12 (32)	Male	Acute asphyxiation	9
	5	5 (40)	Male	Acute asphyxiation	5
	6	11 (36)	Female	Acute asphyxiation	27
	7	13.5 (37.5)	Female	Acute asphyxiation	29
	8	16 (40)	Female	Acute asphyxiation	9
	9	26 (36)	Male	Acute asphyxiation	9
	10	33.4 (38.5)	Male	Intussuseption of inverted Merkel's diverticulum	13
	11	35 (39)	Female	Acute asphyxiation	29
FFPE Non-TBI Control (Figure 6, Supplemental File 1)	Non-TBI-1	0 (23)	Unknown	Normal brain development; labor/pregnancy complications	<24
	Non-TBI-2	0 (27)			
	Non-TBI-3	0 (39)			
	Non-TBI-4	0 (40)			
	Non-TBI-5	4 month	Male	Sudden unexpected death in infancy	24
	Non-TBI-6	5 month	Male	Cystic fibrosis	3.33
	Non-TBI-7	4 year	Female	Acute peritonitis	24
FFPE TBI (Figure 6, Supplemental File 1)	TBI - 1	3 month	Female	Blunt force head injury	23
	TBI - 2	3 year	Female	Traumatic axonal injury, subdural hematoma	24
	TBI - 3	3 year	Male	Road traffic accident, contusions, cerebral edema	24
	TBI - 4	9 year	Female	Road traffic accident, contusions, cerebral edema, subdural hematoma, global ischemia	24
	TBI - 5	9 year	Male	Road traffic accident, contusions, cerebral edema, subdural hematoma, global ischemia	24
	TBI - 6	10 year	Female	Road traffic accident, contusions, cerebral edema	72

826
827

Table 2: Antibody information.

Antibody	Species	Target	Company	Catalog #	Validation Method	WB	IHC
NKCC1	Rabbit	Synthetic Arg77 of mouse NKCC1	Cell Signaling Technology	14581	KO lysate	1:1000	
pNKCC1	Rabbit	Human pNKCC1 Th212/217	Millipore	ABS1004	KO lysate	1:500	1:250
SPAK	Mouse	GST-tagged c-terminus of human STK39	Millipore	MABS178	(+) ctrl human brain tissue	1:1000	
pSPAK	Rabbit	Human pSPAK Ser373	Millipore	07-2273	Hypotonic HEK293 cells	1:500	
KCC2	Rabbit	Synthetic peptide	Abcam	ab259969	(+) and (-) ctrl tissue lysates	1:2000	
pKCC2	Rabbit	Synthetic Ser940	Invitrogen	PA5-95678	KO lysate	1:1000	
Parvalbumin	Sheep	E. coli-derived recombinant human Parvalbumin alpha Ser2-Ser110	Invitrogen	PA5-47693	(+) ctrl brain tissue		1:100
NeuN	Conj 647	Synthetic peptide	Abcam	ab190565	(+) ctrl brain tissue		1:500

Table 3: Taqman probe information

Target	Species	Sequence
Nkcc1	*Human*	*Hs01074325_m1*
Kcc2	*Human*	*Hs00221168_m1*
Stk39	*Mouse*	*Mm01270753_m1*
Reference Gene (18s)	*N/A*	*Hs99999901_s1*

A

C

833 **Supplemental Figure 1:**

834 Damage maps for each animal and region used for cell counts in **Figures 4 and S2**. Yellow map shows damage
835 visible on H&E. Red box indicates approximate ROI used for counts. *A*. ROIs for Cortex 1 *B* ROIs for Cortex 2
836 and 3. *C,* ROIs for hippocampus counts.

838 **Figure S2:**

A   ROI selection for hippocampus counts

1: dorsal DG, 2: dorsal CA2
3: ventral DG, 4: ventral CA2

B   ROI 1 (Dorsal DG)   ROI 2 (Dorsal CA2)

PV / pNKCC1 / NeuN / Nuclei

839
Figure S2: The expression of neuronal pNKCC1 expression in the hippocampus of swine was not affected by age
or injury.
Schematic of ROIs for analysis, **A**. Immunohistochemical staining for pNKCC1 (red), parvalbumin (green), NeuN (magenta), and nuclei
(white), **B**. Cell counts expressed as percentages per ROI, with ROI 1 in **C**, ROI 2 in **D**, ROI 3 in **E,** and ROI 4 in **F**. In all cases, the **left**
dot plot presents % of neurons that were apoptotic, and the **right** dot plot presents the % of healthy neurons expressing pNKCC1. Two-
way ANOVA with post-hoc multiple t tests employed for all comparisons and significant findings for main effect of age or treatment
reported in graphs.
847