In silico evidence suggests that the SARS-CoV-2 Spike protein may target coiled-coil regions of numerous cytoskeletal and cytoskeleton-associated proteins

Pierre Brézellec*[1,2]

[1] Institut Systématique Evolution Biodiversité (ISYEB UMR 7205), Sorbonne Université, MNHN, CNRS, EPHE, UA – Paris, France
[2] Université de Versailles Saint Quentin, 45 avenue des Etats Unis – Versailles, France

*Corresponding author
Correspondence: pierre.brezellec@uvsq.fr

ABSTRACT
Understanding the interactions between host and viral envelope proteins is essential to get insights into the dynamics of viral infection. To investigate more closely the mechanisms governing SARS-CoV-2 entry and intracellular trafficking, I reanalyzed the most extensive SARS-CoV-2–human protein–protein interactome dataset currently available. My investigation centered on the Spike S protein, a key player in initiating viral infection by binding to the host cell membrane receptor Angiotensin-Converting Enzyme 2 (ACE2). I first present evidence demonstrating the statistical overrepresentation of actin-binding proteins among the Spike S partners/interactors. Next, I show that a majority of these partners contains a structural domain sharing high similarity with the C-terminal region of the Myosin II heavy chain, Myosin II being known for its roles in muscle contraction and various cellular motility processes. I subsequently demonstrate that this domain is particularly prevalent in actin-binding proteins, intermediate filaments proteins and kinesins, which all are related to the cytoskeleton known to be involved in diverse cellular functions, including endocytosis and intracellular transport — processes crucial for viral infections. Finally, I highlight that the structural domain mentioned above is a bonafide coiled-coil region. I therefore conclude that Spike S might target proteins possessing such regions. Collectively, my findings suggest that the interactions between SARS-CoV-2 Spike S and human proteins, potentially mediated by coiled-coil regions, may have been underestimated. As this work relies on *in silico* evidence, direct biological extrapolations require caution.

Keywords: SARS-CoV-2–human protein–protein interactome, Protein Structure Prediction, Phyre2, BackPhyre, Bioinformatics

Introduction

Exploring protein-protein interactions (PPI) between host proteins and viral proteins has emerged as a relevant approach in viral infection research (Calderwood *et al.*, 2007). In the context of COVID-19, such approaches have been used to identify pharmacological agents with antiviral activity or potential host therapeutic targets (Gordon *et al.*, 2020, Zhou *et al.*, 2022). In this study, I focused on the most comprehensive and reliable SARS-CoV-2–human PPI data available to date (Zhou *et al.*, 2022), based on both two-hybrid (Y2H) and affinity purification-mass spectrometry (AP-MS), to enhance our understanding of virus infection dynamics (encompassing entry and propagation both intra- and extracellularly).

Large-scale protein interaction studies using techniques like Y2H (Rajagopala *et al.*, 2015) and AP-MS (Causier *et al.*, 2004) have well-documented limitations. For example, they might miss transient interactions or detect interactions between proteins that are not normally found in the same cellular compartment or not expressed at the same time in the cell, *etc.* (EBI Training Online Courses, (Millan, 2022)). Despite these weaknesses, however, Y2H and AP-MS often generate high-quality data which can be used as a reliable and robust basis for further investigation. In our context, at first glance, this assertion might seem surprising given a well-known major limitation of the aforementioned interactome: the critical interaction between the Spike S binding domain (RBD) and the host Angiotensin-Converting Enzyme 2 (ACE2), which allows for the initiation of viral infection, has not been identified. However, while this latter interaction is clearly of great importance, this interactome can potentially help to detect or reveal other types of interesting and meaningful interactions, such as those involving other regions of Spike S, or those taking place at other stages of the infection process.

Relying on the Spike S sub-network of the aforementioned interactome, I first demonstrate that actin-binding proteins are statistically overrepresented within the set of Spike S interactors/partners. Using Phyre2 (Kelley *et al.*, 2015), a popular bioinformatics method for protein structure prediction, I then show that a majority of Spike S partners share structural similarities with the C-terminal part of the Myosin II heavy chain, which is a well-documented coiled-coil region (Rahmani *et al.*, 2021). Next, using BackPhyre, a tool that identifies proteins harboring a given structural domain, I identify a set of human proteins containing with high confidence the C-terminal part of the Myosin II heavy chain and demonstrate that this set is enriched in cytoskeletal and associated cytoskeletal proteins, with actin-binding proteins, intermediate filament proteins, and kinesins all being statistically overrepresented compared to their distribution in the human proteome (refer to the Figure for an overall view of my results).

My work thus provides *in silico* evidence suggesting that SARS-CoV-2 Spike S might target proteins harboring coiled-coil regions, which, as I show here (see also (Simm *et al.*, 2021)), are particularly abundant in cytoskeletal and cytoskeletal-associated proteins. This suggestion is in line with numerous studies that have highlighted the crucial role of the host cell cytoskeleton for coronavirus infection, including SARS-CoV-2 (Wen *et al.*, 2020; Kloc *et al.*, 2022). For example, the actin-binding protein Myosin 9 (Chen *et al.*, 2021) and the intermediate filament vimentin (Amraei *et al.*, 2022) have been shown to facilitate SARS-CoV-2 host cell entry.

          # Materials and methods

Interactome data

The interactions between the Spike S protein and human proteins used in this paper are extracted from a protein-protein interactome published in Nature Biotechnology (Zhou *et al*., 2022). They involved 22 human proteins, hereafter denoted as interactors or partners, as shown in Table 1 below.

Table 1 - Each line represents a Spike S partner. The 'UniProtID' column identifies the protein using its unique identifier in the UniProt database. The 'Protein Name' column indicates the name of the protein. The 'Technique of identification' column refers to the approach used to identify the considered interaction, where 'Y2H' denotes/stands for 'Yeast Two-Hybrid' and 'AP-MS' stands for 'Affinity Purification-Mass Spectrometry'.

UniProt ID	Protein Name	Technique of identification
ACTN4_HUMAN	Alpha-actinin-4	Y2H
MYH9_HUMAN	Myosin-9	AP-MS
MYH10_HUMAN	Myosin-10	AP-MS
MYH14_HUMAN	Myosin-14	AP-MS
TPM1_HUMAN	Tropomyosin alpha-1 chain	AP-MS
TPM2_HUMAN	Tropomyosin beta chain	AP-MS
TPM3_HUMAN	Tropomyosin alpha-3 chain	AP-MS
TPM4_HUMAN	Tropomyosin alpha-4	AP-MS
COR1C_HUMAN	Coronin-1C	AP-MS
TMOD3_HUMAN	Tropomodulin-3	AP-MS
BICL2_HUMAN	BICD family-like cargo adapter 2	Y2H
CYTSA_HUMAN	Cytospin-A	AP-MS
EPIPL_HUMAN	Epiplakin	AP-MS
ITBP2_HUMAN	Integrin beta-1-binding protein 2	Y2H
LZTS2_HUMAN	Leucine zipper putative tumor suppressor 2	Y2H
LMO7_HUMAN	LIM domain only protein 7	AP-MS
ML12B_HUMAN	Myosin regulatory light chain 12B	AP-MS
MYL9_HUMAN	Myosin regulatory light polypeptide 9	AP-MS
MYL6B_HUMAN	Myosin light chain 6B 208	AP-MS
MYL6_HUMAN	Myosin light polypeptide 6	AP-MS
PARP1_HUMAN	Poly(ADP-ribose) polymerase 1	AP-MS
STON2_HUMAN	Stonin 2905	AP-MS;Y2H

Note that partners associated with 'S active' and 'S active RBD1' were excluded from the analysis because the paper by (Zhou *et al*., 2022) provided insufficient information regarding the precise nature of these two terms in both the main text and the Materials and Methods section. Including these interactions would have resulted in an expanded set of 25 proteins (adding ZNF274, ZDHHC5, and ELF4).

Protein Data Bank entry 6XE9

Structure 6XE9 (https://www.rcsb.org/structure/6XE9) represents the heavy chain of Myosin II complex (Yang *et al.*, 2020) of *Meleagris gallopavo* (wild turkey). Its UniProt accession number is G1N5L2 and its sequence length is 1979 amino acids. This protein, also referred to as "class II myosin heavy chain," is a part of the "myosin II complex" which enables muscle cells to contract and facilitates movement and shape changes in non-muscle cells. This complex contains two class II myosin heavy chains, two myosin essential light chains, and two myosin regulatory light chains. These chains assemble to form an alpha-helical, coiled-coil tail.

G1N5L2 is not fully modeled; the protein segments 1-23, 205-210, 635-655, 955-1396, and 1676-1979 are missing. As a consequence:

116
117   - Protein segment 24-204 aligns with structure segment 1-181.
118   - Protein segment 211-634 aligns with structure segment 182-605.
119   - Protein segment 656-954 aligns with structure segment 606-904.
120   - Protein segment 1397-1675 aligns with structure segment 905-1183.
121
122   G1N5L2 is made of two InterPro domains: a Myosin Head Motor Domain (IPR001609,
123 positions 78-790) and a Myosin Tail Domain (IPR002928, positions 854-1934). The alignments
124 between the Myosin Head domain and the modeled part of this domain are the following:
125
126   - Protein segment 78-204 aligns with structure segment 55-181,
127   - Protein segment 211-634 aligns with structure segment 182-605,
128   - Protein segment 656-790 aligns with structure segment 606-740.
129
130   The alignments between the Myosin Tail domain and the modeled part of this domain are the
131 following:
132
133   - Protein segment 854-954 aligns with structure segments 804-904.
134   - Protein segment 1397-1675 aligns with structure segments 905-1183.
135
136   Thus, the modeled Myosin Head covers structure segment 55-740, and the modeled Myosin
137 Tail spans structure segment 804-1183.
138

Phyre2: Protein structure prediction

Phyre2 (Protein Homology/AnalogY Recognition Engine), a free web-based service for
protein structure prediction (Kelley *et al.*, 2015), has been used for protein modeling (*i.e.*,
inferring the three-dimensional structure of a protein from its amino acid sequence).

BackPhyre: Identification of proteins containing a given structural domain

The identification of potential Spike S interaction partners in the human proteome was carried
out using the BackPhyre program (see BackPhyre documentation). BackPhyre takes a known 3D
structure and searches a given proteome for proteins which harbor that specific structure (or a
segment of it). I queried the human proteome (NCBI Reference Sequence database) using PDB
structure 6XE9 (the heavy chain of myosin II complex). BackPhyre returned 999 protein entries
with probabilities ranging from 95.8% to 100%. According to BackPhyre's documentation, if a
match probability exceeds 90%, one can generally be very confident that the protein adopts the
overall fold shown and that the core of the protein is modelled at high accuracy (2-4Å rmsd from
native, true structure).
153
154   From the original 999 entries, I excluded 195 sequences prefixed with "XP_," indicating they
155 were not curated RefSeq records. This left us with a set of 804 curated protein sequences
156 (starting with "NP_"). I then grouped together the different isoforms of each protein-coding gene,
157 resulting in 622 distinct protein clusters (see Supplemental file 2).
158
Given that I fed BackPhyre with the entire PDB structure 6XE9, and to the extent that the
feature identified as common to 12 out of the 22 Spike S' partners (see Results) concerns the C-
terminal part of 6XE9, I performed a filtering step to preserve clusters that encompass at least
one isoform possessing this characteristic. To achieve this, I identified the boundaries of the
largest motif of 6XE9 common to all of my 12 previously referred proteins; I found 853-1067. This

step narrowed down the set to 488 protein clusters. Finally, to represent each protein cluster
uniquely, I selected a single isoform from those with multiple isoforms.

166
Note that BackPhyre's human protein-coding gene set is based on the NCBI36/hg18 genome
assembly (released March 3, 2006), an older version than the current GRCh38.p14 assembly
(released February 3, 2022). Although incomplete, this release is sufficiently accurate, as only 5
of my set of 488 identified human proteins have no UniProt ID or correspond to an obsolete entry
(see Supplemental file 3), though the annotations of the majority of the remaining 483 proteins
were obviously subsequently updated. Consequently, my analysis remains sound, though
BackPhyre's protein set may not be fully representative of human coiled-coil proteins.

Statistical analysis

To evaluate whether a given set of proteins of size N was statistically enriched for a specific
feature F compared to its expected distribution within the human proteome, I performed a
Fisher's exact test (significance level = 0.05). Specifically, considering that a number n of
proteins from the given set share a common feature F, I tested the null hypothesis that this set is
a random sample of size N drawn from the human proteome. Defining N_H as the size of the
human proteome and n_H as the number of human proteins possessing feature F, the structure
of the contingency table and the corresponding formulas for the counts are as follows:

182
	Observed Count (in set N)	Expected Count (in set N)
Proteins with feature F	n	$(n_H/N_H)*N$
Proteins wihtout feature F	N-n	$((N_H-n_H)/N_H)*N$

186
187   Fisher's exact test was conducted on the website BiostatTGV (https://biostatgv.sentiweb.fr/?
188 module=tests/fisher).

189
190   According to UniProt (UniProt Consortium, 2024), the total number of proteins in the human
191 proteome is 82518 (query: (proteome:UP000005640), where UP000005640 is the UniProt ID of
192 the human proteome).

193
I used UniProt Gene Ontology queries to define the feature F and to compute the number of
human proteins with this feature. The human proteins with the feature 'intermediate filament
protein' were identified using the query: (proteome:UP000005640) AND (go:0005882), where
0005882 is the GO term for 'intermediate filament'. Similarly, human proteins with the feature
'kinesin' were identified using the query: (proteome:UP000005640) AND (go:0005871), where
0005871 is the GO term for 'kinesin complex'.

200
201   For the feature 'actin-binding protein', I relied on a collection of 418 actin-associated proteins
202 (AAPs) whose direct interactions with actin have been experimentally verified (Gao & Nakamura,
203 2022).

204   # Results

Spike S partners are enriched in cytoskeleton-associated proteins, particularly actin-
binding proteins, which are statistically overrepresented compared to their distribution in
the human proteome

Among the 22 Spike S partners (see Table 2 for a summary), 10 are annotated as "Actin
Binding Proteins" according to a study which gathered a collection of 418 actin-associated
proteins (AAPs) whose direct interactions with actin have been experimentally verified (Gao &

Nakamura, 2022). These 10 proteins play various roles in actin dynamics (see UniProt
annotations (UniProt Consortium, 2025), (Gao & Nakamura, 2022) and (Cooper, 2000)):

- "Alpha-actinin-4" (UniProt ID ACTN4_HUMAN) is a component of the actin cytoskeleton.

- "Myosin-9" (UniProt ID MYH9_HUMAN), "Myosin-10" (MYH10_HUMAN), and "Myosin-14"
(MYH14_HUMAN) are cellular myosin.

- "Tropomyosin alpha-1 chain" (UniProt ID TPM1_HUMAN), "Tropomyosin beta chain"
(TPM2_HUMAN), "Tropomyosin alpha-3 chain" (TPM3_HUMAN), and "Tropomyosin alpha-4
chain" (TPM4_HUMAN) are involved in stabilizing actin filaments (e.g., protecting F-actin from
severing).

- "Coronin-1C" (UniProt ID COR1C_HUMAN) and "Tropomodulin-3" (TMOD3_HUMAN) are
involved in stabilizing, bundling, and capping actin, respectively.

In the 'Statistical Analysis' section, I demonstrate a statistically significant enrichment of actin-
binding proteins among the set of Spike S protein interactors compared to their expected
distribution within the human proteome.

The remaining 12 Spike S partners interact directly or indirectly with the cytoskeleton, an
intricate network of filaments within cells comprising actin filaments (AFs), microtubules (MTs),
and intermediate filaments (IFs) (Cooper, 2000):

- "BICD family-like cargo adapter 2" (UniProt ID BICL2_HUMAN) belongs to the BICD family-
like. "BICD family-like cargo adapter 1" (BICL1_HUMAN), a member of this family, is predicted to
connect the dynein motor complex to various cargos and enhance dynein's ability to move along
microtubules (MTs) for extended distances without falling off (see UniProt annotations).

- "Cytospin-A" (UniProt ID CYTSA_HUMAN) is documented as a "cross-linking" protein that
functionally interacts with both the actin cytoskeleton and MTs (Saadi *et al.*, 2011).

- "Epiplakin" (UniProt ID EPIPL_HUMAN) is a cytoskeletal linker protein that connects to IFs
and controls their reorganization in response to stress ((Jang *et al.*, 2005), (Shimada *et al.*,
2013), (Kokado *et al.*, 2016)).

- "Integrin beta-1-binding protein 2" (UniProt ID ITBP2_HUMAN), also referred as Melusin, is
a muscle-specific interactor for the beta(1) integrin's cytoplasmic domain, which interacts with the
actin-binding protein alpha-actinin (Otey *et al.*, 1990)

- "Leucine zipper putative tumor suppressor 2" (UniProt ID LZTS2_HUMAN) is a beta-
catenin-interacting protein that modulates beta-catenin signaling and localization (Thyssen *et al.*,
2006). Beta-catenin, a crucial regulator of cell-cell adhesion, forms a complex with alpha-catenin
that bridges cadherins (adherens junction components) and the actin cytoskeleton (Kobielak &
Elaine Fuchs, 2004).

- "LIM domain only protein 7" (UniProt ID LMO7_HUMAN) belongs to a group of proteins that
bind to stressed actin filaments (Winkelman *et al.*, 2020).

- "Myosin regulatory light chain 12B" (UniProt ID ML12B_HUMAN) and "Myosin regulatory
light polypeptide 9" (MYL9_HUMAN) are part of the Myosin II complex, as are "myosin light chain

6B" (MYL6B_HUMAN) and "myosin light polypeptide 6" (MYL6_HUMAN). These light chains do
not bind to actin but interact with the myosin heavy chain, which allow Myosin II complex to bind
to actin.
265
- "Poly(ADP-ribose) polymerase 1" (UniProt ID PARP1_HUMAN), which is mainly and well
documented to play a key role in DNA repair, also interacts with Transgelin, an actin-binding
protein, in human colon cancer cells (Lew *et al.*, 2020). Note that several studies underscore the
functional importance of lamins, actin, myosin, spectrin, and the linker of nucleoskeleton and
cytoskeleton (LINC) complex in DNA repair (Lambert, 2019). Finally note that while PARP1 is
involved in multiple antiviral mechanisms, several viruses can inhibit these antiviral functions and
recruit and utilize PARP1 for their own benefit (Sobotka & Tempera, 2024).
273
274 - "Stonin 2" is an adaptor protein that interacts with components of the endocytic machinery
275 (Martina *et al.*, 2020), which are involved in regulating coat assembly, vesicle budding, and signal
276 transduction pathways along with the actin cytoskeleton (Chakrabarti *et al.*, 2021).
277
278 A Fisher's exact test was performed to assess the enrichment of 'actin-binding' proteins within
279 the set of 22 Spike S partners (10 of which are actin-binding proteins). With a calculated Fisher's
280 exact p-value of 0.000521 (significance level = 0.05), the null hypothesis is rejected, indicating
281 that actin-binding proteins are significantly overrepresented in this set relative to the entire
282 human proteome.
283
In conclusion, the set of 22 Spike S partners is enriched in "cytoskeleton-associated
proteins", particularly in actin-binding proteins which are statistically over-represented.

Table 2 - Each line represents a Spike S partner. The 'UniProtID' column identifies
the protein using its unique identifier in the UniProt database. The 'Protein Name'
column indicates the name of the protein. The 'Length (AA)' column indicates the
total length of the protein sequence (in amino acids). The 'Link with Cytoskeleton'
column indicates whether a documented link exists between the considered Spike
S partner and the cytoskeleton. The 'Similarities with the C-terminal part of Myosin
II heavy chain (PDB entry 6XE9)' column indicates whether the considered Spike
S partner shares similarities (+) or does not share similarities (-) with the C-
terminal tail of the Myosin II heavy chain (this latter column refers to subsection 'A
majority of Spike S partners share similarities with the C-terminal myosin II heavy
chain domain').

UniProt ID	Protein Name	Length (AA)	Link with Cytoskeleton	Similarities with the C-terminal part of the Myosin II heavy chain (PDB entry 6XE9)
ACTN4_HUMAN	Alpha-actinin-4	911	actin-binding	+
MYH9_HUMAN	Myosin-9	1960	actin-binding	+
MYH10_HUMAN	Myosin-10	1976	actin-binding	+
MYH14_HUMAN	Myosin-14	1995	actin-binding	+
TPM1_HUMAN	Tropomyosin alpha-1 chain	284	actin-binding	+
TPM2_HUMAN	Tropomyosin beta chain	284	actin-binding	+
TPM3_HUMAN	Tropomyosin alpha-3 chain	285	actin-binding	+
TPM4_HUMAN	Tropomyosin alpha-4	248	actin-binding	+
COR1C_HUMAN	Coronin-1C	474	actin-binding	-
TMOD3_HUMAN	Tropomodulin-3	352	actin-binding	-
BICL2_HUMAN	BICD family-like cargo adapter 2	508	connects the dynein motor complex to various cargos and enhances dynein's ability to move along microtubules	+
CYTSA_HUMAN	Cytospin-A	1117	interacts with both the actin cytoskeleton and microtubules	+
EPIPL_HUMAN	Epiplakin	5088	cytoskeletal linker protein that connects	-

			to Intermediate Filaments	
ITBP2_HUMAN	Integrin beta-1-binding protein 2	347	Interacts with alpha-actinin	-
LZTS2_HUMAN	Leucine zipper putative tumor suppressor 2	669	forms a complex with alpha-catenin that bridges cadherins (adherens junction components) and the actin cytoskeleton	+
LMO7_HUMAN	LIM domain only protein 7	1683	binds to stressed Actin Filaments	+
ML12B_HUMAN	Myosin regulatory light chain 12B	172	myosin regulatory subunit	-
MYL9_HUMAN	Myosin regulatory light polypeptide 9	172	myosin regulatory subunit	-
MYL6B_HUMAN	Myosin light chain 6B 208	208	regulatory light chain of Myosin	-
MYL6_HUMAN	Myosin light polypeptide 6	151	regulatory light chain of Myosin	-
PARP1_HUMAN	Poly(ADP-ribose) polymerase 1	1014	not really linked although it has been shown to interact with Transgelin, an actin-binding protein, in human colon cancer cells	-
STON2_HUMAN	Stonin 2905	905	interacts with components of the endocytic machinery - actin assembly being an essential element of endocytosis	-

A majority of Spike S partners share similarities with the C-terminal Myosin II heavy chain domain

The observation that a protein interacts with a set of proteins often leads to the speculation that these proteins share a common domain mediating the identified interactions. This is why I investigated whether a majority of the 22 Spike S protein partners would share a common domain. Using InterPro domain annotations, I found no such domain. Therefore, I decided to use Phyre2, a protein 3D structure prediction method (Kelley *et al.*, 2015), to address this question.

I ran Phyre2 on each of the 22 Spike S partners (see Supplemental file 1). Interestingly, a careful manual examination of the results revealed that 12 out of the 22 Spike S partners shared a high-confidence similarities with the C-terminal part of the PDB entry 6XE9, *i.e.*, the myosin II heavy chain of the myosin II complex of *Meleagris gallopavo* (wild turkey) (see Materials and Methods section 'Protein Data Bank entry 6XE9' for a detailed presentation of 6XE9). Its UniProt accession number is G1N5L2 and its sequence length is 1979 amino acids (see (Yang *et al.*, 2020)). It is made of two InterPro domains: a Myosin Head Motor Domain (IPR001609, positions 78-790) and a Myosin Tail Domain (IPR002928, positions 854-1934). As highlighted in subsection 'Protein Data Bank entry 6XE9', 6XE9 is not fully modeled. Specifically, the modeled Myosin Head covers structure segments 55-740, and the modeled Myosin Tail spans structure segments 804-1183.

The 12 Spike S partners that share similarities with the C terminal part of the PDB entry 6XE9 are listed in the Table 3 below. For each of these latter, this table summarizes some protein characteristics (UniProt ID, name, length) as well as its alignment with PDB entry 6XE9 and Phyre2's confidence score.

The results reveal a clear fit between protein structure 6XE9 and myosin heavy chains MYH9, 10, and 14, which is consistent as these proteins belong to the same family. The tropomyosins (TPM1, 2, 3, and 4) exhibit a significant fit with the tail of 6XE9, leaving only a minor portion of their sequences not covered. A substantial portion of the actin sequence ACTN4 also aligns with 6XE9 tail. Finally, the similarity shared between 6XE9 and BICL2 (resp., CYSTA, LMO7, and LZTS2) is localized in the tail region of 6XE9. In conclusion, these 12 proteins share similarities with the Myosin tail domain of 6XE9.

Table 3 - Each line represents a Spike S partner exhibiting similarities with the C-terminal part of PDB entry 6XE9 (Myosin II heavy chain). The 'ID' column provides the protein's UniProt identifier. The 'Length' column indicates the protein's length (in amino acids). The 'Query/6XE9' column shows the aligned amino acid positions between the query protein and the reference structure (6XE9), in the format start-end (query)/start-end (6XE9). The 'Confidence' column displays the Phyre2 confidence score for the similarity, indicating the probability of accurate alignment.

ID	Length	Query / 6XE9	Confidence
ACTN4	911	299-758 / 736-1183	98.44
BICL2	508	30-463 / 793-1182	98.65
CYTSA	1117	341-800 / 730-1183	98.91
LMO7	1683	714-1096 / 803-1183	97.05
LZTS2	669	331-663 / 853-1182	99.58
MYH9	1960	23-1223 / 3-1183	100.00
MYH10	1976	26-1230 / 2-1183	100.00
MYH14	1995	47-1247 / 3-1183	100.00
TPM1	284	1-282 / 810-1091	99.74
TPM2	284	1-282 / 810-1091	99.72
TPM3	285	2-284 / 810-1092	99.74
TPM4	248	4-243 / 828-1067	99.26

The set of human proteins identified by BackPhyre is enriched in cytoskeletal-associated proteins, with actin-binding proteins, intermediate filament proteins, and kinesins all being statistically overrepresented compared to their distribution in the human proteome

To identify potential human interactors/partners of Spike S, I employed BackPhyre which, given a 3D structure, determines if this structure (or part of this structure) is present in proteins of a genome/proteome of interest. I queried the human proteome (NCBI Reference Sequence database) using PDB structure 6XE9 (the heavy chain of myosin II complex), see Materials and Methods, 'BackPhyre: Identification of proteins containing a given structural domain' section.

BackPhyre returned a set of 999 high-confidence hits (probability greater than 0.97). Following a cleaning step (excluding RefSeq entries annotated as 'XP' and grouping together isoforms), I obtained a set of 488 human proteins, which I then clustered into 7 major groups.

1./ 'Link To Actin' cluster:

I relied on a list of manually curated Actin-Associated Proteins (AAPs) (Gao & Nakamura, 2022), which contain Actin-Binding Proteins (ABPs), to identify the actin-binding proteins of my set. This resulted in a cluster that includes proteins known to interact with actin, such as actin, myosin, and tropomyosin. Additionally, I also included proteins functionally linked to actin but not explicitly annotated as actin-binding proteins. The 'Link To Actin' cluster contains 67 proteins, among which 56 are ABPs; I demonstrate (see 'Statistical analysis') that my set of 488 proteins is statistically enriched with ABPs compared to their distribution in the human proteome.

2./ 'Link to Intermediate Filaments' cluster:

I relied on UniProt annotations to identify Intermediate Filament (IF) proteins within the human proteome. (query: "(proteome:UP000005640) AND (go:0005882) where '0005882' refers to 'intermediate filament' in Gene Ontology). 'Link to Intermediate Filaments' cluster groups together IFs, such as keratins, vimentin, neurofilament proteins, etc., along with proteins linked to intermediate filaments. 'Link to Intermediate Filaments' contains 71 proteins among which 68 are

369 IFs. I show in 'Statistical analysis' that my set of 488 proteins is statistically enriched with IFs
370 compared to their distribution in the human proteome.
371
372   3./ 'Link To Kinesin and Dynein' cluster:
373
My analysis of my set of 488 proteins revealed an absence of tubulins, the building blocks of
microtubules. Despite this, I identified several kinesin and dynein motor proteins, known for their
interaction with microtubules to facilitate intracellular transport of various cargo like vesicles,
organelles, and chromosomes. I additionally put four 'Tektins' in this cluster. According to UniProt
annotations, these latter are microtubule inner proteins (MIPs) that are components of the
dynein-decorated doublet microtubules (DMTs) in cilia and flagellar axonemes. The 'Link To
Kinesin and Dynein' cluster contains 46 proteins, among which 24 are kinesins (query:
"(proteome:UP000005640) AND (go:0005871)" where '0005871' refers to 'kinesin complex' in
Gene Ontology). I prove in section 'Statistical analysis' that my set of 488 proteins is statistically
enriched with kinesins compared to their distribution in the human proteome.
384
385   4./ 'Link To Microtubule' cluster:
386
The identification of microtubule motor proteins (kinesins and dyneins, as shown in the cluster
above) prompted us to search for microtubule-related proteins in my set of 488 proteins.
Microtubules, anchored at the centrosome (the MTOC), extend towards the cell periphery and
play a vital role in various cellular processes, e.g., during mitosis, they reorganize into the mitotic
spindle, ensuring proper chromosome segregation. Additionally, cilia and flagella are
microtubule-based structures that contribute to cellular functions. Not surprisingly, my analysis
identified many proteins associated with these structures (centrosome, mitotic spindle, cilia, and
flagella). These proteins were subsequently grouped together into 5 different subclusters,
namely, 'Link to MT', Link To Centrosome', 'Link To Cilia', 'Link To "Sperm Flagellar"', and 'Link
To Spindle'. The 'Link To Microtubule' cluster contains 104 proteins.
397
398   5./ 'Link To Trafficking' cluster:
399
400   Functional analysis of the remaining proteins allowed us to identify an enrichment in proteins
401 associated with transport and trafficking pathways. The 'Link To Trafficking' cluster contains 51
402 proteins, among which are several members of the BICD family. Members of this family are
403 known to link the dynein motor complex to various cargos.
404
405   6. 'Various' cluster:
406
407   I next identified several clusters with a moderate number of proteins, associated for instance
408 with 'defense' or 'LINC (Linker of Nucleoskeleton and Cytoskeleton)', this latter grouping 3
409 proteins involved in the connection between the nuclear lamina and the cytoskeleton. The
410 'Various' cluster contains 106 proteins.
411
412   7. 'Lacking UniProt Summary' cluster:
413
414   This cluster contains 43 proteins lacking a UniProt 'Function' field, which was one of my main
415 source of information for performing my clustering.
416
To evaluate whether the set of 488 human proteins identified by BackPhyre was statistically
enriched for specific functional categories, I performed Fisher's exact tests. The analysis
demonstrated that three key cytoskeletal protein classes, namely actin-binding proteins (grouped

in the 'Link To Actin' cluster, p-value = 1.307E-14), intermediate filament proteins (grouped in the
'Link to Intermediate Filaments' cluster, p-value = 3.935E-19), and kinesins (grouped in the 'Link
To Kinesin and Dynein' cluster, p-value = 8.921E-8), are significantly overrepresented
(significance level 0.05) (see Materials and Methods, Statistical analysis section).
424

Discussion

426   To gain deeper insights into SARS-CoV-2 infection dynamics, I re-examined the most
427 comprehensive SARS-CoV-2–human protein interactome (Zhou *et al*., 2022), which leverages
428 both Y2H (Rajagopala *et al*., 2015) and AP-MS (Causier *et al*., 2004) methods. I focused on the
429 Spike S protein, the viral entry protein that binds to the host cell receptor ACE2.
430
While Y2H and AP-MS have limitations (see Introduction), they generally produce high-quality
data. AP-MS identifies protein complexes while Y2H reveals binary interactions, providing
complementary insights. Nevertheless, initially, I decided to treat Y2H and AP-MS equally as two
probes of interaction. This decision was primarily driven by the observation that Human Myosin-
9, identified as a Spike S partner through AP-MS in the interactome I utilized (Zhou *et al.*, 2022),
had previously been demonstrated to be in close proximity to the SARS-CoV-2 Spike S protein
(using an APEX2 proximity-labeling technique) and subsequently showed to directly interact with
it (Chen *et al.*, 2021).
439
Zhou and colleagues' interactome identified 22 Spike S interactors/partners (see Materials
and Methods section 'Interactome data' and Table 1). Analysis of this set of 22 partners revealed
a predominance of cytoskeleton-associated proteins, with a statistically significant
overrepresentation of actin-binding proteins (10 out of 22) compared to their distribution in the
human proteome (see Results, 'Spike S partners are enriched in cytoskeleton-associated
proteins, particularly actin-binding proteins, which are statistically overrepresented compared to
their distribution in the human proteome' section, and Table 2).
447
To further investigate the interactome results, I hypothesized that the human Spike S partners
might possess a common domain mediating these interactions. Unable to identify such domains
using InterPro annotations (Paysan-Lafosse *et al.*, 2022), I used Phyre2 (Kelley *et al.*, 2015) to
predict the structures of the 22 Spike S partners. I then observed that 12 of these 22 structures
exhibited high-confidence similarities with the tail region/domain of the Myosin II heavy chain
(PDB entry 6XE9, see Results ,'A majority of Spike S partners share similarities with the C-
terminal Myosin II heavy chain domain' section, and Table 3). Among these 12 proteins, 8 are
actin-binding proteins, including human Myosin-9, which was previously shown to interact with
the Spike S protein via its C-terminal domain (as mentioned in the previously).
457
My analysis reveals several key findings: 1) Actin-binding proteins are significantly
overrepresented among the 22 Spike S partners (10 among 22), 2) Eight of these actin-binding
proteins exhibited structural similarities to the C-terminal region of Myosin II heavy chain, 3)
Myosin-9, a cellular myosin and one of these eight proteins, was previously shown to interact
with the Spike S protein through its C-terminal domain (Chen *et al.*, 2021). Importantly, this C-
terminal domain of Myosin-9 is homologous to the C-terminal region of Myosin II heavy chain
which is a well-documented coiled-coil region (Rahmani *et al.*, 2021). These findings led us to
hypothesize that domains homologous to the C-terminal region of Myosin II heavy chain may
play a crucial role in mediating interactions with the Spike S protein.
467

While proteins lacking this type of domain might still interact with the Spike S protein through
other, yet unidentified motifs or domains, my hypothesis provides the most parsimonious
explanation for the observed data. Specifically, this hypothesis clarifies why Affinity Purification-
Mass Spectrometry (AP-MS), which detects protein complexes rather than individual protein-
protein interactions, might identify Spike S partners that don't exhibit similarities to the C-terminal
region of Myosin II heavy chain. This is exemplified by Myosin II light chains. Myosin II light and
heavy chains are integral components of the Myosin II complex. Given that the Myosin II complex
binds to actin through its heavy chains, the identification of light chains by AP-MS likely reflects
their association with the heavy chains within the complex, rather than indicating a direct
interaction between the Spike S protein and the light chains themselves.
478
To identify potential human Spike S protein partners, I decided to search for human proteins
sharing similarities with the C-terminal part of Myosin II heavy chain. To achieve this, I used
BackPhyre ('Phyre2 in reverse'). Given a 3D protein structure, this tool searches for homologous
sequences across a broad range of genomes/proteomes. I ran BackPhyre with the PDB entry
6XE9 (corresponding to the Myosin II heavy chain) to explore the human proteome (see
Materials and methods, section BackPhyre: Identification of proteins containing a given structural
domain).
486
BackPhyre returned a set of 999 high-confidence hits (probability >= 0.97). Following a
cleaning step (excluding RefSeq entries annotated as 'XP' and grouping together isoforms), I
obtained a set of 488 human proteins, which I then clustered into 7 major groups. I next
demonstrated that three major classes of cytoskeletal proteins are statistically overrepresented in
this set compared to their distribution in the human proteome: Actin-binding proteins (grouped in
the 'Link To Actin' cluster), Intermediate filament proteins (grouped in the 'Link to Intermediate
Filaments' cluster), and Kinesins (grouped in the 'Link To Kinesin and Dynein' cluster) (see
Results, 'The set of human proteins identified by BackPhyre is enriched in cytoskeletal-
associated proteins, with actin-binding proteins, intermediate filament proteins, and kinesins all
being statistically overrepresented compared to their distribution in the human proteome'
section).
498
These findings are consistent with existing research emphasizing the importance of the
cytoskeleton in viral infections (Kloc *et al*., 2022; Wen *et al*., 2022). Hence, the 'Link To Actin'
cluster contains the non-muscle myosin heavy chain IIA (MYH9), which has been shown to
interact with Spike S and to facilitate SARS-CoV-2 infection in human pulmonary cells (Chen *et*
al., 2021). The 'Link to Intermediate Filaments' cluster contains vimentin, which has also been
documented to interact with Spike S and to facilitate entry into human endothelial cells (Amraei *et*
al., 2022). The 'Link To Kinesin and Dynein' cluster contains kinesin-1, a protein documented to
be hijacked by several viruses for transport along microtubules (to reach their destinations in the
highly crowded cellular environment). For instance, the intercellular transport of Porcine
Epidemic Diarrhea Virus (PEDV), which is a coronavirus, is driven along microtubules by both
dynein and kinesin-1 (Hou *et al*., 2021). Furthermore, cytoplasmic dynein and kinesin-1 have
both been found to bind Adenovirus (AdV) through direct interactions with the capsid proteins
(Scherer *et al*., 2020). Although, to the best of my knowledge, no study has yet reported that
SARS-CoV-2 hijacks kinesin-1 via an interaction with Spike S, exploring this possibility
represents an interesting avenue for future research.
514
515   In conclusion, my work provides *in silico* evidence suggesting that SARS-CoV-2 Spike S
516   might target proteins harboring regions homologous to the C-terminal region of Myosin II heavy
517   chain, which is a coiled-coil domain. These regions, as shown here, are particularly prevalent in
518   cytoskeletal and cytoskeleton-associated proteins. These regions, as shown here, are

particularly prevalent in cytoskeletal and cytoskeleton-associated proteins (for example, myosins, intermediate filament proteins, and kinesins; see also Simm *et al*., 2021) and are documented to be hijacked by several viruses. However, since this work is based primarily on a large-scale protein-protein interaction dataset, which may contain false positives or interactions irrelevant in physiological contexts (e.g., non-co-localization), direct biological extrapolations require caution. Further biochemical and cell biological investigations are necessary to study the interaction between Spike S and proteins containing coiled-coil domains, and the involvement of such domains in these interactions.

Acknowledgements

I warmly thank Joël Pothier (ABI, ISYEB) for his careful review of a draft of this manuscript. I am also grateful to the two referees of my paper, whose comments and suggestions significantly improved the quality of the final submission.

Conflict of interest disclosure

The author declares that he complies with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Supplementary information availability

Supplemental Files 1, 2, and 3 are available online (https://doi.org/10.5281/zenodo.14718017).

References

Amraei R, Xia C, Olejnik J, White M R , Napoleon M A , Lotfollahzadeh S, Hauser B M , Schmidt A G, Chitalia V, Mühlberger E, Costello C E , Rahimi N. Extracellular vimentin is an attachment factor that facilitates SARS-CoV-2 entry into human endothelial cells. Proc Natl Acad Sci U S A. 2022 Feb 8;119(6):e2113874119. https://doi.org/10.1073/pnas.2113874119.

Calderwood M A , Venkatesan K, Xing L, Chase M R , Vazquez A, Holthaus A M, Ewence A E, Li N, Hirozane-Kishikawa T, Hill D E, Vidal M, Kieff E, Johannsen E. Epstein-Barr virus and virus human protein interaction maps. Proc Natl Acad Sci U S A. 2007 May 1;104(18):7606-11. https://doi.org/10.1073/pnas.0702332104.

Causier B. Studying the interactome with the yeast two-hybrid system and mass spectrometry. Mass Spectrom Rev. 2004 Sep-Oct;23(5):350-67. https://doi.org/10.1002/mas.10080.

Chakrabarti R, Lee M, Higgs H N. Multiple roles for actin in secretory and endocytic pathways. Curr Biol. 2021 May 24;31(10):R603-R618. https://doi.org/10.1016/j.cub.2021.03.038.

Chen J, Fan J, Chen Z, Zhang M, Peng H, Liu J, Ding L, Liu M, Zhao C, Zhao P, Zhang S, Zhang X, Xu J. Nonmuscle myosin heavy chain IIA facilitates SARS-CoV-2 infection in human pulmonary cells. Proc Natl Acad Sci U S A. 2021 Dec 14;118(50):e2111011118. https://doi.org/10.1073/pnas.2111011118.

Cooper G M. The Cell: A Molecular Approach. 2nd edition. Sunderland (MA): Sinauer Associates; 2000.

Gao J, Nakamura F. Actin-Associated Proteins and Small Molecules Targeting the Actin Cytoskeleton. Int J Mol Sci. 2022 Feb 14;23(4):2118. doi: https://doi.org/10.3390/ijms23042118.

2016 Apr 26:6:25120. https://doi.org/10.1038/srep25120.

Gordon D E, Jang G M, Bouhaddou M, *et al*. A SARS-CoV-2 protein interaction map reveals
targets for drug repurposing. Nature. 2020 Jul;583(7816):459-468.
https://doi.org/10.1038/s41586-020-2286-9.

Jang S-I, Kalinin A, Takahashi K, Marekov L N, Steinert P M. Characterization of human
epiplakin: RNAi-mediated epiplakin depletion leads to the disruption of keratin and vimentin IF
networks. J Cell Sci. 2005 Feb 15;118(Pt 4):781-93. https://doi.org/10.1242/jcs.01647.

Kelley L A, Mezulis S, Yates C M, Wass M N, Sternberg Michael J E. The Phyre2 web portal for
protein modeling, prediction and analysis. Nat Protoc. 2015 Jun;10(6):845-58.
https://doi.org/10.1038/nprot.2015.053.

Kloc M, Uosef A, Wosik J, Kubiak J Z, Ghobrial R M. Virus interactions with the actin
cytoskeleton-what we know and do not know about SARS-CoV-2. Arch Virol. 2022
Mar;167(3):737-749. https://doi.org/10.1007/s00705-022-05366-1.

Kobielak A, Fuchs E. α-catenin: at the junction of intercellular adhesion and actin dynamics. Nat
Rev Mol Cell Biol. 2004 Aug;5(8):614–625. https://doi.org/10.1038/nrm1433.

Kokado M, Okada Y, Miyamoto T, Yamanaka O, Saika S. Effects of epiplakin-knockdown in
cultured corneal epithelial cells. BMC Res Notes. 2016 May 20:9:278.
https://doi.org/10.1186/s13104-016-2082-7.

Lambert M W. The functional importance of lamins, actin, myosin, spectrin and the LINC complex
in DNA repair. Exp Biol Med (Maywood). 2019 Nov;244(15):1382-1406.
https://doi.org/10.1177/1535370219876651.

Lew Z-X, Zhou H-M, Fang Y-Y, Ye Z, Zhong W, Yang X-Y, Yu Z, Chen D-Y, Luo S-M, Chen L-F,
Lin Y. Transgelin interacts with PARP1 in human colon cancer cells. Cancer Cell Int. 2020
Aug 3:20:366. https://doi.org/10.1186/s12935-020-01461-y.

Martina J A , Bonangelino C J, Aguilar R C, Bonifacino J S. Stonin 2: an adaptor-like protein that
interacts with components of the endocytic machinery. J Cell Biol. 2001 May 28;153(5):1111-
20. https://doi.org/10.1083/jcb.153.5.1111.

Otey C A, Pavalko F M, Burridge K. An interaction between alpha-actinin and the beta 1 integrin
subunit in vitro. J Cell Biol. 1990 Aug 111(2):721-9. https://doi.org/10.1083/jcb.111.2.721.

Millan P P. Protein interactions and their importance, an introduction. 2022 (last reviewed).
https://www.ebi.ac.uk/training/online/courses/protein-interactions-and-their-importance/.

Paysan-Lafosse T, Blum M, Chuguransky S *et al.* InterPro in 2022. Nucleic Acids Res. 2023 Jan
6;51(D1):D418-D427. https://doi.org/10.1093/nar/gkac993.

Rahmani H, Ma W, Hu Z, Daneshparvar N, Taylor D W, McCammon J A, Irving T C, Edwards R
J, Taylor K A. The myosin II coiled-coil domain atomic structure in its native environment.
Proc Natl Acad Sci U S A. 2021 Apr 6;118(14):e2024151118.
https://doi.org/10.1073/pnas.2024151118.

Rajagopala S V. Mapping the protein–protein interactome networks using yeast two-hybrid
screens. Adv Exp Med Biol. 2015:883:187-214. https://doi.org/10.1007/978-3-319-23603-
2_11.

Saadi I, Alkuraya F S, Gisselbrecht S S, Goessling W *et al.* Deficiency of the cytoskeletal protein
SPECC1L leads to oblique facial clefting. Am J Hum Genet. 2011 Jul 15;89(1):44-55.
https://doi.org/10.1016/j.ajhg.2011.05.023.

Shimada H, Nambu-Niibori A, Wilson-Morifuji M *et al.* Epiplakin modifies the motility of the HeLa
cells and accumulates at the outer surfaces of 3-D cell clusters. J Dermatol. 2013
Apr;40(4):249-58. https://doi.org/10.1111/1346-8138.12076.

Sobotka A A, Tempera I. PARP1 as an Epigenetic Modulator: Implications for the Regulation of
Host-Viral Dynamics. Pathogens. 2024 Jan 30;13(2):131.
https://doi.org/10.3390/pathogens13020131.

Simm D, Hatje K, Waack S, Kollmar Martin. Critical assessment of coiled-coil predictions based
on protein structure data. Sci Rep. 2021 Jun 14;11(1):12439. https://doi.org/10.1038/s41598-
021-91886-w.
Thyssen G, Li T-H, Lehmann L, Zhuo M, Sharma M, Sun Z. LZTS2 is a novel beta-catenin-
interacting protein and regulates the nuclear export of beta-catenin. Mol Cell Biol. 2006
Dec;26(23):8857-67. https://doi.org/10.1128/mcb.01031-06.
UniProt Consortium. UniProt: the Universal Protein Knowledgebase in 2025. Nucleic Acids Res.
2024 Nov 18:gkae1010. https://doi.org/10.1093/nar/gkae1010.
Wen Z, Zhang Y, Lin Z, Shi K, Jiu Y. Cytoskeleton-a crucial key in host cell for coronavirus
infection. J Mol Cell Biol. 2020 Jul 1;12(12):968-979. https://doi.org/10.1093/jmcb/mjaa042.
620
621 Winkelman J D, Anderson C A, Suarez C, Kovar D R, Gardel M L. Evolutionarily diverse LIM
622   domain-containing proteins bind stressed actin filaments through a conserved mechanism.
623   Proc   Natl   Acad   Sci   U   S   A.   2020   Oct   13;117(41):25532-25542.
624   https://doi.org/10.1073/pnas.2004656117.
625
Yang S, Tiwari P , Lee K-H, Sato O, Ikebe M, Padrón R, Craig R. Cryo-EM structure of the
inhibited   (10S)   form   of   myosin   II.   *Nature*   588,   521–525   (2020).
https://doi.org/10.1038/s41586-020-3007-0.
Zhou Y, Liu Y, Gupta S *et al.* A comprehensive SARS-CoV-2–human protein–protein interactome
reveals COVID-19 pathobiology and potential host therapeutic targets. Nat Biotechnol. 2022
Oct 10;41(1):128–139. https://doi.org/10.1038/s41587-022-01474-0.
632

633
634

Figure - A. Focusing on a human 'Spike S interactors/partners' set of 22 proteins (Zhou *et al.*, 2022), I show that actin-binding proteins are statistically overrepresented (10 among 22) compared to their distribution in the human proteome. Next, using Phyre2 (Kelley *et al.*, 2015), a bioinformatics tool for protein structure prediction, I identify that 12 out of 22 Spike S partners contain a structural domain sharing similarities with the C-terminal part of the Myosin II heavy chain. B./ Using BackPhyre (Kelley *et al.*, 2015), a tool that identifies proteins harboring a given structural domain, I identified a set of 488 human proteins that, with high probability, share similarites with the C-terminal part of the Myosin II heavy chain. This set of proteins is particularly abundant in human cytoskeletal and associated proteins (statistically overrepresented in actin-binding proteins, Intermediate Filament proteins, and kinesins), as well as in proteins involved in microtubule dynamics. C. Given that the C-terminal domain of the Myosin II heavy chain is documented as a coiled-coil region, I infer that Spike S

649 might target proteins containing such regions. * means statistically
650 overrepresented.