1

Neuropeptidergic circuit modulation of
developmental sleep in *Drosophila*

Chikayo Hemmi[1], Kenichi Ishii[1, #], Mana Motoyoshi[1], Masato Tsuji[1]

and Kazuo Emoto[1, 2, #, $]

[1]Department of Biological Sciences, Graduate School of Science, The University of
Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033 Japan

[2]International Research Center for Neurointelligence (WPI-IRCN), The University of
Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033 Japan

[#]Correspondence: emoto@bs.s.u-tokyo.ac.jp (lead contact), ishii-k@g.ecc.u-tokyo.ac.jp

[$]Lead contact

Abstract

Sleep-wakefulness regulation dynamically evolves along development in flies, fish, and humans. While the mechanisms regulating sleep in adults are relatively well understood, little is known about their counterparts in early developmental stages. Here, we report a neuropeptidergic circuitry that modulates sleep in developing *Drosophila* larvae. Through an unbiased screen, we identified the neuropeptide Hugin and its receptor PK2-R1 as critical regulators of larval sleep. Our data suggest that HugPC neurons secrete Hugin peptides to activate insulin-producing cells (IPCs), which express a Hugin receptor PK2-R1. IPCs, in turn, release *Drosophila* insulin-like peptides (Dilps) to regulate sleep. We further show that the Hugin/PK2-R1 axis is dispensable for adult sleep control. Our findings thus reveal the neuromodulatory circuitry regulating larval sleep, highlighting differential impacts of the same modulatory axis on developmental sleep and adult sleep.

Introduction

Sleep is a fundamental state with significant impacts on many aspects of cognition and metabolism (Blumberg et al., 2005; Davis et al., 2004; Sorribes et al., 2013). While sleep remains essential throughout life, its regulation unfolds dynamically over the course of development. For instance, the pace of sleep-wakefulness cycle is relatively fast in infants, but then gradually slows down and stabilizes in adults (Blumberg et al., 2005; Davis et al., 2004; Sorribes et al., 2013). Furthermore, sleep-wake cycle in infants is independent of circadian rhythm, but it gradually comes under the circadian regulation (Davis et al., 2004; Frank et al., 2017; Poe et al., 2023). Such developmental

evolution of sleep-wakefulness cycles appears to be conserved across flies and

mammals, including humans (Blumberg et al., 2005; Davis et al., 2004; Poe et al., 2023;

Szuperak et al., 2018). In contrast to the deep insight into the molecular and neural

mechanisms underlying adult sleep regulation (Barlow & Rihel, 2017; Scammell et al.,

2017; Shafer & Keene, 2021), little is known about regulatory mechanisms underlying

infant sleep, partially due to a lack of convenient models to study developmental sleep.

Drosophila larvae have recently emerged as a suitable model for studying sleep

regulation mechanisms. Recent studies have reported that the second instar larvae show

short periods (< 6 seconds) of an "inactive state" during locomotion. Importantly, this

"inactive state" is consistent with the general definition of sleep: reduced

responsiveness to noxious stimuli, homeostatic responses to sleep deprivation, and rapid

reversibility upon stimulation (Szuperak et al., 2018), suggesting that the "inactive

state" might correspond to the sleep state. Interestingly, loss-of-function mutations in

the clock genes *clock* and *cyc* fail to impact larval sleep, while significantly altering the

sleep patterns in adult flies (Dubowy & Sehgal, 2017; Hendricks et al., 2003; Szuperak

et al., 2018). Likewise, mutations in the dopamine transporter *DAT* do not influence

larval sleep, while reducing sleep amounts in adults (Kume et al., 2005; Szuperak et al.,

2018). These observations imply that sleep regulation mechanisms may be, at least in

part, distinct between larvae and adults, yet the neural mechanism of larval sleep

remains vastly understudied compared to that of adult sleep.

In this study, we performed an unbiased genetic screen in second-instar larvae, and

identified the neuropeptide Hugin and its receptor PK2-R1 as a pair critical for larval

sleep control. At the circuit level, Hugin-producing HugPC neurons directly stimulate

the insulin-producing cells (IPCs) via PK2-R1 to regulate sleep. Surprisingly, we found

that regulators of larval sleep are either irrelevant or exert the opposite effects on adult

sleep. Overall, our findings uncover the neuropeptidergic circuitry that regulates larval

sleep, and highlight the mechanistic differences between larval and adult sleep.

Results

PK2-R1 is required for sleep control in *Drosophila* larvae

To understand the molecular and neural mechanisms of larval sleep, we modified a

previously reported system (Churgin et al., 2019; Szuperak et al., 2018) to automatically

quantify the sleep amount of second instar larvae (Figure 1—figure supplement 1) (see

Materials & Methods for details). Our system automatically annotates larvae in each

frame as either "inactive" or "active" states on the basis of pixel changes, with ~90%

accuracy compared to manual annotation (Figure 1—figure supplement 1C). We found

that bouts of ≥ 12 inactive frames are consistent with the general criteria for sleep:

reduced responsiveness to noxious stimuli, homeostatic responses to sleep loss, and

rapid reversibility upon stimulation (Raizen et al., 2008; Szuperak et al., 2018) (Figure

1—figure supplement 2. Therefore, the present study defines sleep as an inactive state

of ≥ 12 consecutive frames (Figure 1—figure supplement 1A).

Using this system, we next carried out an unbiased screen for genes involved in sleep

regulation. To this end, we tested 47 CRISPR knock-out lines of genes encoding

enzymes or receptors for neuropeptides and monoamines (Deng et al., 2019). As

illustrated in Figure 1—figure supplement 3, 13 out of 47 homozygous null mutants

exhibited significant changes in sleep amounts compared to the control group. To

further test the functions of these candidate genes, we next took advantage of the

CRISPR knock-in drivers in which GAL4 sequence is inserted into each gene locus.

With these drivers, we silenced neurons expressing each candidate gene by expressing

the inward-rectifying potassium channel Kir2.1 (Baines et al., 2001). Silencing neurons

expressing *Oamb* and *PK2-R1* resulted in a significant sleep decrease, phenocopying the

knock-out of these genes (Figure 1A, 1B, and Figure 1—figure supplement 4D). *Oamb*

encodes the octopamine receptor and has been reported to be involved in larval sleep

(Szuperak et al., 2018). The role of *PK2-R1* in larval sleep, on the other hand, has been

unknown to date, prompting us to focus on this gene.

Insulin-producing cells express PK2-R1 and regulate sleep in developing larvae

Since the *PK2-R1* driver broadly labeled neurons (~1000 cells) throughout the CNS

(Figure 1C), we next attempted to identify a subpopulation of these neurons relevant for

sleep regulation. Notably, IPCs in the pars intercerebralis (PI), the neurosecretory center

of *Drosophila*, have been reported to express *PK2-R1* (Schlegel et al., 2016).

Consistently, the *PK2-R1$^{2A-LexA}$* knock-in driver labeled all IPCs (Figure 2A and 2B). To

assess whether IPCs are involved in larval sleep, we silenced IPCs with Kir2.1

expression using the IPC-specific drivers *Dilp3-* and *Dilp5-GAL4* (Figure 2C). We

found that silencing of IPCs significantly increased sleep amounts compared to the

control, phenocopying *PK2-R1* neuron silencing and *PK2-R1* knock-out (Figure 1A, 1B,

and 2C). Consistently, both *Dilp3* and *Dilp5* null mutants exhibited larger sleep amounts

compared to the control (Figure 2D). These phenotypes of *Dilp* mutants IPC silencing

are unlikely to be attributed to locomotion defects, as the travel distances during wake

periods were unaffected (Figure 2—figure supplement 1B and 1C). Collectively, these

findings suggest that *PK2-R1* in IPCs regulates larval sleep.

Hugin-expressing neurons control larval sleep

Given that PK2-R1 is a receptor for the neuropeptide Hugin (Rosenkilde et al., 2003),

we next examined whether Hugin is involved in larval sleep. We found that *Hug* mutant

larvae exhibited significantly larger sleep amounts compared to the control (Figure 3A),

consistent with the *PK2-R1* knock-out (Figure 1A). When we silenced *Hug*-expressing

neurons by driving the expression of Kir2.1 with the *Hug* CRISPR-knock-in driver

$Hug^{2A\text{-}GAL4}$ (Deng et al., 2019), it resulted in a significant increase in sleep amounts

without detectable changes in locomotion speed (Figure 3B and Figure 3—figure

supplement 1B). Conversely, activation of *Hug*-expressing neurons with the red-shifted

channel rhodopsin ReaChR (Inagaki et al., 2014) or the heat-sensitive ion channel

TrpA1 (Hamada et al., 2008) both caused a significant reduction in sleep amount

(Figure 3C and 3D). These data together indicate that *Hug*-expressing neurons are

responsible for downregulating larval sleep.

Hug-expressing neurons are comprised of 20 cells in the subesophageal zone (SEZ) of

the brain, and are classified into four subpopulations based on their projection patterns

(Bader et al., 2007). Of these, one subpopulation called HugPC neurons (Bader et al.,

2007) has been reported to form synaptic connections with IPCs (Hückesfeld et al.,

2016). We thus suspected that HugPC neurons may be responsible for sleep regulation.

In line with this notion, axons of HugPC neurons labeled by a newly generated *HugPC-*

LexA driver (Figure 3—figure supplement 2) and IPCs labeled by *Dilp3-GAL4* are

located in close proximity to each other (Figure 3E). Indeed, silencing of HugPC

neurons by Kir2.1 significantly increased larval sleep (Figure 3F), phenocopying those

observed in *Hug* and *PK2-R1* mutants. These results hint at the possibility that HugPC

neurons activate IPCs via Hugin/PK2-R1 signaling to regulate sleep.

Hugin triggers Ca^{2+} elevation and Dilp secretion in larval IPCs via PK2-R1

A previous report showed that IPCs exhibit increased firing rates and intracellular

calcium (Ca^{2+}) levels upon stimulation (Kréneisz et al., 2010). This motivated us to test

whether Hugin activates IPCs. To this end, we utilized the recently developed calcium

indicator CRTC::GFP, which can report Ca^{2+} dynamics in the timescale of several

minutes (Bonheur et al., 2023). As glucose reportedly triggers Ca^{2+} dynamics in IPCs in

a range of minutes (Oh et al., 2019), we wondered if CRTC::GFP can detect these

dynamics as a positive control. To this end, we applied D-glucose to isolated larval

brains and found a significant cytosol-to-nuclear translocation of CRTC::GFP in IPCs

within ~5 min, confirming that this reporter can indeed detect Ca^{2+} responses in IPCs

upon stimulation (Figure 5A and 5B). We then examined whether *Hug* neurons act

upstream of *PK2-R1*-expressing IPCs, by measuring Ca^{2+} responses of IPCs while

thermogenetically activating *Hug* neurons with TrpA1 (Figure 4A). In both isolated

brains (*ex vivo*) and living larvae (*in vivo*), we found that activation of *Hug* neurons

significantly increased Ca^{2+} levels in IPCs (Figure 4B and 4C), suggesting that *Hug*

neurons act upstream of IPCs (Figure 4D). Next, we tested whether Hug peptides can

induce Ca^{2+} responses in IPCs using *ex vivo* preparations. The *Hug* gene encodes a

precursor that potentially produces at least two distinct peptides, Hug-γ and PK-2, both

of which can bind PK2-R1 to induce Ca^{2+} responses (Rosenkilde et al., 2003). We thus

incubated the CRTC::GFP-expressing larval brain with chemically synthesized Hug-γ or

PK-2 (Figure 5A). We found that bath application of either Hug-γ or PK-2 induced

significant Ca^{2+} responses in IPCs (Figure 5B–D). In agreement with the idea that Hug

peptides activate IPCs via *PK2-R1*, knock-out of *PK2-R1* blocked the Hug peptide-

evoked Ca^{2+} elevation in IPCs (Figure 5G). Similar results were obtained when we bath-

applied the synthetic Neuromedin U (NMU), the mammalian homlog of Hugin (Figure

5A, 5C, and 5G), suggesting that the role of Hugin/PK2-R1 signaling may be conserved

across species. Furthermore, Dilp3 accumulated in IPCs was significantly reduced after

applying Hug peptides, suggesting that Hugin causes IPCs to release Dilp3 (Figure 5E).

Collectively, these data suggest that *Hug* neurons activate IPCs via Hugin/PK2-R1

signaling to regulate sleep (Figure 5H).

Hugin/PK2-R1 axis has distinct impacts on sleep control in larvae and adults

Given the differences between larval versus adult sleep in temporal pattern and

circadian influence, we next wondered if the Hugin/PK2-R1/Dilps axis functions in

adult sleep as well. As in larvae, *PK2-R1* knock-out adults showed increased sleep

amount compared to the control (Figure 6—figure supplement 1B). In contrast, *Hug*

knock-out or silencing of HugPC neurons failed to influence adult sleep (Figure 6A and

6B). This suggests that, unlike in larvae, Hug is dispensable for adult sleep.

Interestingly, the expression patterns of *PK2-R1* and *Hug*, as well as the morphology of

HugPC neurons and IPCs, were almost comparable between larvae and adults (Figure

6—figure supplement 2). This implies that the differential roles of Hug in larvae versus

adults are likely due to physiological differences in HugPC neurons and/or IPCs. We

thus examined how Hug peptides evoke Ca^{2+} responses in adult IPCs and found that

neither Hug-γ nor PK-2 could induce Ca^{2+} responses in adult IPCs (Figure 6D and 6E),

unlike in larvae. Consistently, Hug peptides failed to reduce Dilp3 levels in adult IPCs

(Figure 6F). These data indicate that Hug induces Ca^{2+} responses and Dilp secretion in

larval IPCs, but not in adult IPCs. Surprisingly, we further found that *Dilp3* or *Dilp5*

null mutations reduced sleep amounts in adults (Figure 6C), while in larvae, these

mutations in fact increased the sleep amounts. These data unexpectedly reveal how the

same set of neuronal circuits and molecular signaling therein can evolve over

development in a complex manner.

Discussion

In this study, we have identified the neuropeptide Hugin and its receptor PK2-R1 as a

ligand/receptor pair critical for sleep regulation in *Drosophila* larvae. We have further

shown that larval IPCs express PK2-R1, respond to and release Dilp3 upon Hugin

stimulation. Surprisingly, the Hugin/PK2-R1 axis is dispensable for sleep regulation in

adults, even though gene expressions as well as the circuit structure appeared to be

conserved between larval and adult brains. Furthermore, Dilps appear to modulate sleep

in opposite directions in larvae and adults. Our findings thus uncover the

neuropeptidergic modulation circuitry that specifically regulates developmental sleep,

and suggest divergent usage of the same molecules and circuitry for sleep modulation in

larvae and adults.

Hugin/PK2-R1/Dilps axis negatively regulates sleep in larvae

The present study shows that the neuropeptide Hugin and its receptor PK2-R1 are

required for sleep regulation in the early larval stage. This notion is supported by the

following lines of evidence. First, knock-out mutations in *Hug* or *PK2-R1* increased

sleep amounts compared to the control (Figure 1A and 3A). Consistently, silencing a

subpopulation of *Hug*-expressing neurons, called HugPC neurons, or *PK2-R1* neurons

increased sleep amounts (Figure 1B and 3F). Second, thermogenetic activation of *Hug*

neurons decreased sleep amounts (Figure 3D). Third, HugPC neurons project their

axons to IPCs in the brain that express PK2-R1 receptors (Figure 3E). Fourth, Hug

peptides induced Ca^{2+} elevation and promoted Dilp3 release from IPCs in a PK2-R1-

dependent manner (Figure 5). Last, *Dilp3* null mutations increased sleep amounts

(Figure 2D). Based on these data, we propose a model in which the Hugin/PK2-

R1/Dilps axis along the HugPC-IPC circuitry modulates larval sleep (Figure 7).

Examples of Hugin acting through PK2-R1 are prominent in the context of feeding

regulation in larvae and adults (Schoofs et al., 2014). For example, a recent study has

suggested that Hugin regulates the timing of pupariation and body wall contraction

during pupariation through PK2-R1 in PTTH neurons (Ohhara et al., 2024). In the

present study, we showed that HugPC in the larval brain modulates sleep (Figure 3F). A

previous electron microscopic analysis suggested that larval HugPC neurons likely form

synaptic connections to downstream neurons, including IPCs (Hückesfeld et al., 2021;

Schlegel et al., 2016). We here showed that larval sleep control requires PK2-R1, and

that thermogenetic activation of *Hug* neurons causes Ca^{2+} responses in PK2-R1-

expressing IPCs (Figure 4). Importantly, our *ex vivo* data indicated that Hugin peptide-

induced Ca^{2+} response in IPCs was largely abolished by *PK2-R1* mutation (Figure 5),

indicating the critical role of the Hugin/PK2-R1 axis for the HugPC-mediated IPC

activation. Together, we propose that the HugPC-IPCs circuity mobilizes Hugin/PK2-

R1 signaling to selectively modulate larval sleep. Interestingly, our *ex vivo* data

indicated that Neuromedin U (NMU), a mammalian ortholog of Hugin, can trigger Ca^{2+}

responses in larval IPCs in a PK2-R1-dependent fashion (Figure 5). Given that NMU

has been implicated in sleep regulation in fish and mammals (Ahnaou & Drinkenburg,

2011; Chiu et al., 2016), an analogous ligand/receptor pair might be involved in sleep

modulation in mammals as well.

Divergent roles of insulin in regulation of larval sleep and feeding

IPCs in larvae responded to exogenously applied Hug peptides, both by exhibiting

elevated Ca^{2+} levels and decreases in intracellular Dilp3 storage (Figure 5C–E). In

contrast to Dilp3, however, the amount of Dilp2 accumulated in IPCs did not decrease

following Hug peptide application (Figure 5—figure supplement 1B). On the other

hand, Dilp2 accumulation in IPCs decreased after glucose treatment (Figure 5—figure

supplement 1A) (Oh et al., 2019). These results unexpectedly suggest that larval IPCs

form parallel channels inside, one for Hug-Dilp3 and the other for glucose-Dilp2.

Mechanistically, Hug acts directly on IPCs via the PK2-R1 receptor. As for glucose, on

the other hand, glucose-sensing CN neurons in the brain take up glucose and

subsequently secrete another neuropeptide to activate IPCs (Oh et al., 2019). Such

parallelism inside IPCs might provide insights into why insect species broadly express

multiple insulin-like peptide genes, in some cases even as many as 40 genes in the

silkworm *Bombyx mori* (Okamoto, 2021). *Drosophila* genome harbors eight *Dilp* genes

(Nässel et al., 2015), of which *Dilp2*, *Dilp3*, and *Dilp5* are considered nutrient-related.

Nevertheless, mRNA expression or release patterns of these Dilps are not identical

under different nutrient conditions (Géminard et al., 2024; Nässel et al., 2015; Oh et al.,

2019; Ikeya et al., 2002). Furthermore, these Dilps exhibit different binding affinities

for the insulin receptor (InR), the secreted decoy of InR (SDR), and the ecdysone-

inducible gene L2 (Imp-L2) (Okamoto et al., 2013). These reports hint at the differential

roles of different insulin-like peptides; however, mechanistic insights remain largely

lacking. Our data on larval IPCs forming parallel channels for Hug-Dilp3 and glucose-

Dilp2 may provide a point of entry for tackling this mystery.

Differential impacts of the same neuropeptidergic modulations on larval and adult

sleep

Our data show that the Hugin/PK2-R1/Dilps axis likely downregulates sleep amounts in

larvae. Despite the significant increase in sleep amount in *Hug* mutant larvae, *Hug*

mutant adults showed sleep amounts comparable to wild-type control (Figure 6A),

consistent with a previous report that Hugin is dispensable for control of baseline sleep

levels (Schwarz et al., 2021). According to our histological analyses, the Hugin/PK2-R1

expression patterns, as well as the HugPC-IPCs circuitry, appear to be largely conserved

in adults (Figure 3E and Figure 6—figure supplement 2). Unlike larval brains, however,

Hug peptides failed to trigger Ca^{2+} responses in adults IPCs (Figure 6E). It is thus

possible that Hugin/PK2-R1 signaling along the HugPC-IPCs circuitry is suppressed in

adults. IPCs in adults receive multiple positive and negative modulatory inputs through

GPCRs including the metabotropic $GABA_B$ receptors (Enell et al., 2010), which

suppress IPC activity and Dilp release in adult IPCs (Enell et al., 2010). It is thus

plausible that such negative modulatory inputs to IPCs in adults might counteract the

Hugin/PK2-R1 axis to suppress Dilp release. In addition, our data suggest that Dilps

modulate sleep amount in the opposite directions in larvae and adults (Figure 7).

Comparing the expression levels and activities of GPCRs in larval and adult IPCs would

be essential to better understand how same modulatory signals over the course of

development come to exert differential impacts on sleep. Interestingly, Hugin in adults

appears irrelevant for the baseline sleep amount but is required for homeostatic

regulation of sleep (Schwarz et al., 2021). Thus, testing whether the Hugin/PK2-R1 axis

is involved in the homeostatic regulation of larval sleep, and how this system compares

to its adult counterpart, may further provide mechanistic insights into how homeostatic

sleep regulation maturates over development.

In summary, we identified the Hugin/PK2-R1/Dilps signaling along the HugPC/IPCs

circuit as the molecular and circuitry basis for larval sleep regulation. Our study begins

to fill the knowledge gap in larval sleep regulation, and sheds light on mechanistic

differences in sleep regulation between early developmental stages and adults.

Materials and Methods

Key resource table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Gene (*Drosophila melanogaster*)	*PK2-R1*	NA	FLYB:FBgn0038140	

Gene (*D. melanogaster*)	*Dilp3*	NA	FLYB:FBgn0044050	
Gene (*D. melanogaster*)	*Dilp5*	NA	FLYB:FBgn0044048	
Gene (*D. melanogaster*)	*Hug*	NA	FLYB:FBgn0028374	
Gene (*Homo sapiens*)	*NMU*	NA	HGNC:7859	
Genetic reagent (*D. melanogaster*)	w^{1118}	Other	NA	Provided by Dr. Yi Rao, Peking University.
Genetic reagent (*D. melanogaster*)	*iso31*	Bloomington *Drosophila* Stock Center	BDSC:5905; RRID:BDSC_5905	Isogenized for chr 1;2;3, with w[1118] line.
Genetic reagent (*D. melanogaster*)	$PK2\text{-}R1^{attP}$	Bloomington *Drosophila* Stock Center	BDSC:84563; RRID:BDSC_84563	FlyBase symbol: TI{TI}PK2-R1[attP]
Genetic reagent (*D. melanogaster*)	$PK2\text{-}R1^{2A\text{-}GAL4}$	Bloomington *Drosophila* Stock Center	BDSC:84686; RRID:BDSC_84686	FlyBase symbol: TI{2A-GAL4}PK2-R1[2A-GAL4]
Genetic reagent (*D. melanogaster*)	$PK2\text{-}R1^{2A\text{-}LexA}$	Bloomington *Drosophila* Stock Center	BDSC:84431; RRID:BDSC_84431	FlyBase symbol: TI{2A-lexA::GAD}PK2-R1[2A-lexA]
Genetic reagent (*D. melanogaster*)	$Dilp3^{1}$	Bloomington *Drosophila* Stock Center	BDSC:30882; RRID:BDSC_30882	FlyBase symbol: TI{TI}Ilp3[1]
Genetic reagent (*D. melanogaster*)	$Dilp5^{1}$	Bloomington *Drosophila* Stock Center	BDSC:30884; RRID:BDSC_30884	FlyBase symbol: TI{TI}Ilp5[1]
Genetic reagent (*D. melanogaster*)	*Dilp3-GAL4*	Bloomington *Drosophila* Stock Center	BDSC:52660; RRID:BDSC_52660	FlyBase symbol: P{Ilp3-GAL4.C}2

Genetic reagent (*D. melanogaster*)	*Dilp5-GAL4*	Bloomington *Drosophila* Stock Center	BDSC:66007; RRID:BDSC_66007	FlyBase symbol: P{Ilp5-GAL4.L}8
Genetic reagent (*D. melanogaster*)	*Hug^attP*	Bloomington *Drosophila* Stock Center	BDSC:84514; RRID:BDSC_84514	FlyBase symbol: TI{TI}Hug[attP]
Genetic reagent (*D. melanogaster*)	*Hug^2A-GAL4*	Bloomington *Drosophila* Stock Center	BDSC:84646; RRID:BDSC_84646	FlyBase symbol: TI{2A-GAL4}Hug[2A-GAL4]
Genetic reagent (*D. melanogaster*)	*Hug^2A-LexA*	Bloomington *Drosophila* Stock Center	BDSC:84397; RRID:BDSC_84397	FlyBase symbol: TI{2A-lexA::GAD}Hug[2A-lexA]
Genetic reagent (*D. melanogaster*)	*HugPC-GAL4*	Hückesfeld et al., 2016 (DOI:10.1038/ncomms12796)	NA	Provided by Dr. Michael J. Pankratz, University of Bonn.
Genetic reagent (*D. melanogaster*)	*HugPC-LexA*	This paper	NA	The 544-bp enhancer region of Hug was placed upstream of nlsLexA::p65, and the construst was inserted into attP2 site by ΦC31 integrase-mediated transgenesis. See Methods for details.
Genetic reagent (*D. melanogaster*)	*UAS-Kir2.1::EGFP*	Bloomington *Drosophila* Stock Center	BDSC:6596; RRID:BDSC_6596	FlyBase symbol: P{UAS-

				Hsap¥KCNJ2.EGFP}1
Genetic reagent (*D. melanogaster*)	*UAS-Stinger, LexAop-tdTomato.nls*	Bloomington *Drosophila* Stock Center	BDSC:66680; RRID:BDSC_66680	FlyBase symbol: P{UAS-Stinger}2; PBac{13XLexAop2-IVS-tdTomato.nls}VK00022
Genetic reagent (*D. melanogaster*)	*UAS-mCD8::mCherry, UAS-CRTC::GFP*	Bloomington *Drosophila* Stock Center	BDSC:99657; RRID:BDSC_99657	FlyBase symbol: P{UAS-mCD8.mCherry-T2A-lacZ.nls}JK22C; P{UAS-Crtc.GFP}attP40
Genetic reagent (*D. melanogaster*)	*UAS-ReaChR*	Bloomington *Drosophila* Stock Center	BDSC:53741; RRID:BDSC_53741	FlyBase symbol: P{UAS-ReaChR}attP40
Genetic reagent (*D. melanogaster*)	*UAS-TrpA1*	Bloomington *Drosophila* Stock Center	BDSC:26263; RRID:BDSC_26263	FlyBase symbol: P{UAS-TrpA1(B).K}attP16
Genetic reagent (*D. melanogaster*)	*LexAop-TrpA1*	Burke et al., 2012 (DOI:10.1038/nature11614)	NA	Provided by Dr. Scott Waddel, University of Oxford.
Genetic reagent (*D. melanogaster*)	*LexAop-rCD2::RFP-p10.UAS-mCD8::GFP-p10*	Bloomington *Drosophila* Stock Center	BDSC:67093; RRID:BDSC_67093	FlyBase symbol: P{lexAop-rCD2::RFP-p10.UAS-mCD8::GFP-p10}su(Hw)attP5

Genetic reagent (*Escherichia coli*)	One Shot™ *ccdB* Survival™ 2 T1^R competent cells	Thermo Fisher Scientific	Thermo Fisher Scientific:A10460	
Antibody	Anti-GFP (chicken polyclonal)	Aves Labs	Aves Labs:GFP-1020; RRID:AB_10000240	IHC (1:1000)
Antibody	Anti-mCherry (mouse monoclonal)	Takara Bio	Takara Bio:632543; RRID:AB_2307319	IHC (1:1000)
Antibody	Anti-Dilp3 (rabbit polyclonal)	Veenstra et al., 2009 (DOI:10.1007/s00441-009-0769-y)	NA	IHC (1:250) Provided by Dr. Jan A. Veenstra, Université de Bordeaux.
Antibody	Anti-Dilp2 (rabbit polyclonal)	Okamoto et al., 2012 (DOI:10.1073/pnas.1116050109)	NA	IHC (1:2000) Provided by Dr. Takashi Nishimura, RIKEN.
Antibody	Anti-brp (mouse monoclonal)	Developmental Studies Hybridoma Bank	DSHB:nc82; RRID: AB_2314866	IHC (1:100)
Antibody	Anti-histone H3 phospho S28 (rat monoclonal)	Abcam	Abcam:ab10543; RRID:AB_2295065	IHC (1:1000)
Antibody	Goat anti-chicken Alexa Fluor 488	Thermo Fisher Scientific	Thermo Fisher Scientific:A-11039; RRID:AB_2534096	IHC (1:200)

Antibody	Goat anti-mouse Alexa Fluor 555	Thermo Fisher Scientific	Thermo Fisher Scientific:A-21424; RRID:AB_141780	IHC (1:200)
Antibody	Goat anti-rabbit Alexa Fluor 633	Thermo Fisher Scientific	Thermo Fisher Scientific:A-21071; RRID:AB_2535732	IHC (1:200)
Antibody	Goat anti-rat Alexa Fluor 633	Thermo Fisher Scientific	Thermo Fisher Scientific:A-21094; RRID:AB_2535749	IHC (1:200)
Recombinant DNA reagent	pBPnlsLexA::p65Uw	Addgene	Addgene:26230; RRID:Addgene_26230	
Sequence-based reagent	*HugPC* enhancer	This paper	NA	A 544-bp fragment PCR-amplified from the fly genomic DNA. See Methods for details.
Peptide, recombinant protein	Hug-γ	This paper	NA	Chemically synthesized *Drosophila* Hug-γ peptide with amino acid sequences of Acetyl-QLQSNGEPAYRVRTPRL-CONH2

Peptide, recombinant protein	PK-2	This paper	NA	Chemically synthesized *Drosophila* PK-2 peptide with amino acid sequences of Acetyl-SVPFKPRL-CONH2
Peptide, recombinant protein	NMU	This paper	NA	Chemically synthesized human NMU peptide with amino acid sequences of Acetyl-FRVDEEFQSPFASQSRGYFLFRPRN-CONH2
Commercial assay or kit	PrimeSTAR Max DNA Polymerase	Takara Bio	Takara Bio:R045A	
Commercial assay or kit	In-Fusion HD Cloning Kit w/Cloning Enhancer	Takara Bio	Takara Bio:639635	
Chemical compound, drug	D(+)-Glucose	FUJIFILM Wako Pure Chemical Corporation	FUJIFILM Wako Pure Chemical Corporation:041-00595	
Software, algorithm	Adobe Illustrator 2023	Adobe	RRID:SCR_010279	
Software, algorithm	Adobe Photosop 2023	Adobe	RRID:SCR_014199	

Software, algorithm	ImageJ v1.53t	Schneider et al., 2012 (DOI:10.1038/nmeth.2089)	RRID:SCR_003070	
Software, algorithm	Inkscape v1.2	The Inkscape Team	RRID:SCR_014479	
Software, algorithm	Prism v9.5.1	GraphPad	RRID:SCR_002798	
Software, algorithm	Python Programming Language v3.10.0	Python Software Foundation	RRID:SCR_008394	
Software, algorithm	R Project for Statistical Computing v4.1.2	R Core Team 2021	RRID:SCR_001905	
Software, algorithm	SnapGene v5.3.3	GSL Biotech LLC	RRID:SCR_015052	
Software, algorithm	Arduino UNO	Arduino	RRID:SCR_017284	
Software, algorithm	TriKinetics DAMSystem3 Software	TriKinetics Inc. (available at https://trikinetics.com/)	RRID:SCR_021809	
Other	TriKinetics Drosophila Activity Monitoring System	TriKinetics Inc.	RRID:SCR_021798	
Other	Fan-less Peltier-type incubator	Mitsubishi Electric Engineering Co., Ltd.	Mitsubishi Electric Engineering Co., Ltd.:SLC-25	Used to keep flies for DAM experiments.

Fly strains

The following strains of *Drosophila melanogaster* were obtained from the Bloomington

Drosophila Stock Center: *iso31* (an isogenic w^{1118} strain; BDSC# 5905), *PK2-R1^{attP}*

(BDSC# 84563), *PK2-R1$^{2A-GAL4}$* (BDSC# 84686), *PK2-R1$^{2A-LexA}$* (BDSC# 84431),

Dilp3^1 (BDSC# 30882), *Dilp5^1* (BDSC# 30884), *Dilp3-GAL4* (BDSC# 52660), *Dilp5-*

GAL4 (BDSC# 66007), *Hug^{attP}* (BDSC# 84514), *Hug$^{2A-GAL4}$* (BDSC# 84646), *Hug$^{2A-LexA}$*

(BDSC# 84397), *UAS-Kir2.1::EGFP* (BDSC# 6596), *UAS-Stinger, LexAop-*

tdTomato.nls (BDSC# 66680), *UAS-mCD8::mCherry, UAS-CRTC::GFP* (BDSC#

99657), *UAS-ReaChR* (BDSC# 53741), *UAS-TrpA1* (BDSC# 26263), *LexAop-ReaChR*

(BDSC# 53746), *LexAop-rCD2::RFP-p10.UAS-mCD8::GFP-p10* (BDSC# 67093).

w^{1118} was from Y. Rao (Peking University); *HugPC-GAL4* (Hückesfeld et al., 2016) was

from M. Pankratz (University of Bonn); *LexAop-TrpA1* (Burke et al., 2012) was from S.

Waddel (University of Oxford). *HugPC-LexA* was generated in this study (see the

following section, "Generation of transgenic lines"). Detailed lists of fly genotypes can

be found in the key resource table and Source Data files.

General fly maintenance

Flies were maintained on conventional fly food under a 9AM:9PM light/ dark cycle at

25°C and 60 +/- 5 % humidity.

Egg collection

Agar substrates for egg collection were prepared by microwaving a mixture of 2.0 g

sucrose (Fujifilm Wako Pure Chemicals, cat# 196-00015), 3.0 g agar (BD, cat#

214010), 2.5 mL apple juice (Dole), and Milli-Q water up to 100 mL. Melted agar was

poured into Petri dishes with a diameter of 4 cm (approximately 3 mL per dish).

Solidified agar was scratched with tweezers to make the surface rough. Agar plates were

stored inside a sealed container at 4°C. Upon egg collection, yeast paste (dried yeast

(Nippon Beet Sugar Co., Ltd.) kneaded with Milli-Q water at 40% (w/w)) was placed in

the center of the agar plates to further promote oviposition.

Larval staging

First-instar larvae were collected two days after egg collection. Among them, second-

instar larvae that molted within 2 h were used for video recording. The distinction

between first- and second-instar larvae was based on morphological characteristics:

while anterior spiracles were not evident in the first-instar, fist-shaped spiracles were

observed in the second-instar.

Larval sleep analysis

Twenty-four-well silicone chambers for sleep analysis, described in Figure 1—figure

supplement 1B, were made by pouring dimethylpolysiloxane (PDMS) into molds based

on a previous study (Churgin et al., 2019; Szuperak et al., 2018). As a substrate, 120 µL

of a mixture of 3% agar and 2% sucrose was poured into each well. After the gel

solidified, 10 µL of 60 mg/mL yeast suspension was applied to the surface of the gel in

each well. One larva was transferred into each well, and a glass plate (Azone, cat# 1-

4540-01, 5 mm × 200 mm × 200 mm) was placed on top of the chamber to prevent the

larvae from escaping. The chamber was placed inside a dark box equipped with a ring-

type infrared LED (CCS, cat# LDR2-132IR940-LA) and a camera (The Imaging

Source, cat# DMK27BUP031), and recording was started immediately without

acclimation. Images were captured through the IC Capture software (The Imaging

Source, version 2.5) placed above the chamber under the following conditions: MJPEG

Compressor for codec, a frame rate of 0.87 fps, and an image size of 2592 × 1944. For

each well and frame, the image of larva was blurred and binarized to automatically

locate the larva. Consecutive frames were then compared so that changes in values were

summed to estimate the larval motion. This motion estimate was then used to classify

every consecutive pair of frames as representing "active" or "inactive." We found that

inactivity across 12 consecutive frames meets the sleep criteria, as shown in Figure 1—

figure supplement 2.

Sleep deprivation

Blue light irradiation was performed using a 455-nm LED (Thorlabs, cat# M455L4)

with a collimating adapter (Thorlabs, cat# SM1U25-A). The irradiation intensity was set

to the maximum scale using a T-cube LED driver (Thorlabs, cat# LEDD1B). Temporal

control of the LED illumination was achieved using an Arduino UNO microcontroller

board (Arduino, RRID:SCR_017284). For sleep deprivation experiments, the first 1 h

was recorded without LED illumination, followed by another 1 h consisting of repeated

cycles of "LED ON for 90 s and OFF for 30 s."

Immunohistochemistry

The following steps were carried out in succession: dissection in phosphate-buffered

saline (PBS); fixation in 4% paraformaldehyde/PBS at room temperature for either 90

min in CRTC::GFP experiments or 30 min in the others; washing with 0.3% Triton X-

100/PBS (hereafter referred to as PBST) for 10 min × 3; blocking in 5% normal goat

serum (NGS)/PBST at room temperature for 30 min; primary antibody treatment at 4°C

overnight; washing with PBST for 10 min × 3; secondary antibody treatment at 4°C

overnight; washing with PBST for 10 min × 3; and mounting in VECTASHIELD

medium (Vector Laboratories, cat# H-1000, RRID:AB_2336789). The above steps were

performed either in Terasaki 96-well plates (stem Corp., cat# P96R37N) for adults or in

48-well culture plates (Corning, cat# 3548) for larvae. The following primary antibodies

were diluted in 5% NGS/PBST: chicken polyclonal anti-GFP (1:1000, Aves Labs, cat#

GFP-1020, RRID:AB_10000240), mouse monoclonal anti-mCherry (1:1000, Takara

Bio, cat# 632543, RRID:AB_2307319), rabbit polyclonal anti-Dilp3 (1:250, (Veenstra

et al., 2008)), rabbit polyclonal anti-Dilp2 (1:2000, (Okamoto et al., 2012)), mouse

monoclonal anti-BRP (1:100, Developmental Studies Hybridoma Bank, clone nc82;

RRID:AB_2314866), and rat monoclonal anti-histone H3 phospho S28 (1:1000,

Abcam, cat# ab10543; RRID:AB_2295065). The following secondary antibodies were

diluted in 5% NGS/PBST: goat anti-chicken Alexa Fluor 488 (1:200, Thermo Fisher

Scientific, cat# A-11039, RRID:AB_2534096), goat anti-mouse Alexa Fluor 555 (1:200,

Thermo Fisher Scientific, cat# A-21424, RRID:AB_141780), goat anti-rabbit Alexa

Fluor 633 (1:200, Thermo Fisher Scientific, cat# A-21071, RRID:AB_2535732), and

goat anti-rat Alexa Fluor 633 (1:200, Thermo Fisher Scientific, cat# A-21094;

RRID:AB_2535749). Fluorescence images were obtained using a confocal microscope

(Leica, TCS SP8) with 20× (Leica, HC PL APO CS2 20×/0.75 IMM) and 63× (Leica,

HC PL APO CS2 63×/1.40 OIL) objectives. Images were captured at 512 × 512 pixel

resolution in 1 μm-steps along the z-axis. Z-stack images of the maximum projections

were generated using the ImageJ v1.53t software ((Schneider et al., 2012);

RRID:SCR_003070).

Generation of transgenic lines

HugPC-LexA lines were generated by ΦC31 integrase-mediated transgenesis, as

previously described (Groth et al., 2004). The 544-bp enhancer region of *Hug,* identical

region to that used for *HugPC-GAL4* (Hückesfeld et al., 2016), was PCR-amplified

from fly genomic DNA. The gel-purified amplicon was subjected to another round of

PCR to add hangout sequences. The backbone suicide-vector pBPnlsLexA::p65Uw

(RRID:Addgene_26230) was initially amplified by transforming the *ccdB*-resistant *E.*

coli strain, One Shot™ *ccdB* Survival™ 2 T1^R competent cells (Thermo Fisher

Scientific, cat# A10460). The PCR product and the vector were digested by Aat II

(NEB, cat# R0117S) and Fse I (NEB, cat# R0588S), followed by ligation using In-

Fusion HD Cloning Kit w/Cloning Enhancer (Takara Bio, cat# 639635). The integrity of

the resulting plasmid was confirmed by DNA sequencing. Plasmid insertion was

targeted to four attP sites (attP2, attP40, VK00005, and su(Hw)attP5) individually

through ΦC31 integrase-mediated transgenesis by BestGene Inc. (Chino Hills, CA).

Transformants were selected based on the eye color marker. Among the generated

strains harboring *HugPC-LexA* in each insertion site, labeling patterns in the larval brain

were checked by driving fluorescent marker expression under the control of *LexAop*

sequence (Figure 3—figure supplement 2). As the *HugPC-LexA* inserted in attP2

closely resembled the labeling pattern of the original *HugPC-GAL4*, we chose this

driver in further experiments.

The following primers were used for cloning:

Forward primer for the *HugPC* enhancer; 1st round PCR against genomic DNA:

5'-AAGGGTTTGGTTTAATTTATTTATGTCATA-3'

Reverse primer for the *HugPC* enhancer; 1st round PCR against the genomic DNA:

5'-GGACAACTGATGCCAGCAGC-3'

Forward primer for the *HugPC* enhancer with hangout sequences (underlined); 2nd

round PCR against the 1st round amplicon:

5'-gaaaagtgccacctgacgtAAGGGTTTGGTTTAATTTATTTATGTCATA-3'

Reverse primer for the *HugPC* enhancer with hangout sequences (underlined); 2nd round

PCR against the 1st round amplicon:

5'-cccgggcgagctcggccggGGACAACTGATGCCAGCAGC-3'

Measurement of neuronal activity and Dilp3 accumulation in IPCs

The GFP-tagged CRTC (CRTC::GFP) reporter was applied to quantify the activities of

larval IPCs, essentially according to the previous report in adult flies (Bonheur et al.,

2023). Parental *Dilp3-GAL4* and the reporter line harboring *UAS-CRTC::GFP* and

UAS-mCD8::mCherry (BDSC# 99657) were crossed, and eggs were collected within an

8–10 h time window. Offspring larvae were reared for 4 days after egg laying (AEL) at

25°C until the early third instar. Collected larvae were washed with water and starved

on water-soaked paper for 12–14 h. For *in vivo* thermogenetic activation, larva-

containing Petri dishes were warmed at 30°C for 60 min. For *ex vivo* thermogenetic

experiments, larvae were dissected in calcium- and sugar-free HL3.1 buffer (Nakamizo-

Dojo et al., 2023) and each brain was incubated in the same buffer at 30°C for 5 min.

For *ex vivo* glucose or peptide application assays, larvae were first dissected in the same

buffer as above at room temperature, and brain samples were then treated with either 20

mM D-glucose or 1 mM of chemically synthesized peptides dissolved in HL3.1 buffer

for 5 min. After incubation, samples were subjected to immunohistochemical analyses

to visualize GFP, mCherry, and anti-Dilp3 signals (see the section

"Immunohistochemistry").

Image processing was performed using ImageJ v1.53t ((Schneider et al., 2012);

RRID:SCR_003070). Nuclear ROIs and cell body outlines were determined based on

the mCD8::mCherry fluorescence insensitive to neuronal activity, and the mean

intensities of CRTC::GFP signals within the nucleus (F_{nuc}) and the cytoplasm (F_{cyto} =

$F_{whole\ cell}$ - F_{nuc}) were quantified. NLI was calculated using the following formula: (F_{nuc} -

F_{cyto})/(F_{nuc} + F_{cyto}). To assess Dilp3 accumulation in IPCs, the anti-Dilp3 signal intensity

was measured within the cytosol of each cell, excluding the nucleus. Data from 4–14

Dilp3-positive IPCs per brain, collected from 3–7 larvae, were pooled for each

genotype/treatment group.

Chemical synthesis of Hug peptides

Peptides with the following amino acid sequences were chemically synthesized using

the Fmoc solid phase method, and their purity was assessed by HPLC to be over 95%.

All procedures were carried out by Eurofins Genomics K.K. (Tokyo, Japan). The amino

acid sequences of the peptides used were as follows:

Drosophila Hug-γ:

Acetyl-QLQSNGEPAYRVRTPRL-CONH2 (17 a.a.)

Drosophila PK-2:

Acetyl-SVPFKPRL-CONH2 (8 a.a.)

Human NMU:

Acetyl-FRVDEEFQSPFASQSRGYFLFRPRN-CONH2 (25 a.a.)

Sleep quantification in adult flies

The following parts of the *Drosophila* activity monitor (DAM) system were purchased

from TriKinetics Inc. (Waltham, MA): the *Drosophila* Activity Monitor (cat# DAM2),

the power supply interface unit (cat# PSIU9), monitor tubes with an outside diameter of

5 mm and a length of 65 mm (cat# PGT5x65), and tube caps (cat# CAP5-Black). Newly

eclosed adult flies were collected under brief CO_2 anesthesia and kept as groups of 10–

15 in standard food vials. Males and females were kept separately to avoid mating. To

minimize disturbance by noise and vibration, flies were placed inside a fan-less Peltier-

type incubator (Mitsubishi Electric Engineering Co., Ltd., cat# SLC-25) set to 25°C,

externally connected with 5-mm diameter LED lights (MY-CRAFT, Ltd., cat# 5W0601)

and a timer device (Panasonic Corp., cat# WH3311WP) to generate a 9AM:9PM

light/dark cycle. After 3 days of eclosion, each fly was gently aspirated into a Pyrex

glass monitor tube, one end of which was stuffed with agar food (2% (w/w) agar and

5% (w/w) sucrose), and the other was plugged with a cotton string. The tubes were set

to the monitor device and placed inside the same fan-less incubator. The DAM system

was connected to an external PC running DAMSystem3 Data Collection Software

(RRID:SCR_021809; available at https://trikinetics.com/), which counted beam-

crossings by each fly for two successive days. Time windows longer than 5 min without

beam-crossing were judged as sleep bouts, following the conventional criteria for adult

fly sleep (Paul et al., 2000).

Statistical analysis

Statistical analyses by Kruskal–Wallis one-way ANOVA, Mann–Whitney U-test, or chi-

squared test were performed using Prism 9.5.1 (GraphPad, RRID:SCR_002798).

Bonferroni correction was applied for multiple comparisons. Asterisks (*) represent p-

values as indicated within each figure legend. All data necessary to reproduce the figure

panels and statistical analyses are available as Source Data files.

Acknowledgements

We thank Y. Rao, M. Pankratz, S. Waddel, and the Bloomington *Drosophila* Stock

Center for fly stocks; J. A. Veenstra for the anti-Dilp3 antibody; T. Nishimura for the

anti-Dilp2 antibody; Y. Nishizuka for valuable suggestions on writing codes for

automated larval detection and sleep quantification; H. Ito, N. Yamaji, M. Maetani, A.

Ogasawara, E. Kato, T. Rikiishi, S. Ando, M. Hayashi, and S. Miyazaki for technical

assistance and fly maintenance; and the members of Emoto Lab for critical comments

and discussion. This work is supported by MEXT Grants-in-Aid for Scientific Research

on Innovative Areas "Dynamic regulation of brain function by Scrap and Build system"

(KAKENHI 16H06456), JSPS (KAKENHI 16H02504), WPI-IRCN, AMED-CREST

(JP22gm310010), JSTCREST, Toray Foundation, Naito Foundation, Takeda Science

Foundation, and Uehara Memorial Foundation to K.E.; the Leading Initiative for

Excellent Young Researchers (LEADER) from MEXT, JSPS (KAKENHI 22K06309),

and AMED-PRIME (JP22gm6510011) to K.I.; Grant-in-Aid for JSPS Fellows

(24KJ0911) to M.M.; and Grant-in-Aid for JSPS Fellows (22J21096; 22KJ1042) to

C.H.

Author contributions

C.H., K.I., and M.M. performed experiments and data analyses; M.T. wrote the codes

for automated larval detection and sleep quantification; C.H., K.I., and K.E. are

responsible for study conception, experimental design, and data interpretation; C.H.,

K.I., and K.E. wrote the original manuscript, and M.T. edited the paper.

Competing interests

The authors declare no conflicts of interest.

Figure legends

Figure 1. PK2-R1 is required for larval sleep control

(A) Sleep amounts in *PK2-R1* or *Oamb* knock-out mutants. In this and the following

panels, 'N' indicates the number of biologically independent animals used for each

group. Box plots are generated so that the center line indicates the median, the box

limits indicate the upper and lower quartiles, and the whiskers indicate the minimum-to-

maximum range. *** $p < 0.001$ (Mann–Whitney U-test with Bonferroni correction). (B)

Sleep amounts in larvae in which *PK2-R1* neurons were silenced. *** $p < 0.001$ (Mann–

Whitney U-test with Bonferroni correction). (C) The expression pattern of *PK2-R1^{2A-}*

^{GAL4} > UAS-Kir2.1::EGFP in the larval CNS. Similar results were obtained across three

independent samples.

Figure 1—figure supplement 1. Detailed experimental setup for analyzing larval

sleep

(A) Schematic flow of automated larval sleep quantification. A second instar larva was

placed in each of 24 wells, and all 24 wells were videotaped from above. For each well

and frame, the image of larva was blurred and binarized to automatically locate the

larva. Consecutive frames were then compared so that changes in pixel values were

summed to estimate larval motion. This motion estimate was then used to classify every

consecutive pair of frames as representing "active" or "inactive." We found that

inactivity across 12 consecutive frames meets the sleep criteria, as shown in Figure 1—

figure supplement 2. (B) Schematic representation of the 24-well chamber used for

larval sleep assays. (C) Flow of automated processes of larval motion detection and

active/inactive state determination. For each well and frame, the image of larva was

blurred and binarized. Consecutive frames were then compared so that changes in pixel

values were summed to estimate larval motion. This motion estimate was then used to

classify every consecutive pair of frames as representing "active" or "inactive."

Figure 1—figure supplement 2. Larval "sleep state" in this study is consistent with

behavioral criteria for sleep

(A) Relationship between "inactive" state duration and spontaneous arousal probability.

Sustained quiescence for ≥ 12 frames (open circles) significantly reduced arousal

probability compared to 1-frame inactivity. * $p < 0.05$, NS: $p \geq 0.05$ (chi-squared test).

(B) The percentage of larvae that showed a response within the next 10 frames after

high-intensity blue light. The number of events analyzed was 36 and 60 with and

without blue light irradiation, respectively. * $p < 0.05$ (Fisher's exact test). (C) Sleep

amounts before, during, and after blue light irradiation. The number of larvae tested was

21 and 21 for the control and experimental groups, respectively. ** $p < 0.01$, NS: $p \geq$

0.05 (Mann–Whitney U-test with Bonferroni correction). (D) Percentage of larvae that

showed a response in the next 1 or 12 frames after blue light stimulation. Number of

events analyzed: 1960 or 111 for 1- or 12-frame continuous inactivity prior to blue light

illumination, respectively. **** $p < 0.0001$ (chi-squared test).

Figure 1—figure supplement 3. CRISPR-knock-out screen for genes that regulate

larval sleep.

Total sleep amounts in CRISPR-knock-out mutants were measured for 18 h from the

beginning of the second instar. The data for Control, *PK2-R1,* and *Oamb* mutants are

identical to those shown in Figure 1. Box plots are generated so that the center line

indicates the median, the box limits indicate the upper and lower quartiles, and the

whiskers indicate the minimum-to-maximum range. *** p < 0.005 (Kruskal–Wallis one-

way ANOVA and post hoc Mann–Whitney U-test with Bonferroni correction). Number

of larvae tested: 475 for the control and 10–52 for each mutant.

Figure 1—figure supplement 4. Effects of neuronal manipulation on larval sleep

Sleep amounts in larvae in which distinct neuronal populations were silenced. Kir2.1

expression was driven by each enhancer-GAL4 for "hit" genes identified in the

CRISPR-knock-out screen. 'N' indicates the number of biologically independent

animals used for each group. Box plots are generated so that the center line indicates the

median, the box limits indicate the upper and lower quartiles, and the whiskers indicate

the minimum-to-maximum range. *** p < 0.001, ** p < 0.01, * p < 0.05, NS: ≥ 0.05

(Mann–Whitney U-test with Bonferroni correction).

Figure 2. IPCs and Dilps negatively regulate larval sleep

(A) Triple labeling of *PK2-R1* neurons expressing nuclear-localized RFP (magenta),

Dilp3 neurons expressing nuclear-localized GFP (green), and anti-Dilp2 positive cells

(blue). Top panels show signals in the larval brain and the VNC. The bottom panels

show magnified images of the white-squared area in the top panels, where the cell

bodies of the IPCs are located. Note that all IPCs labeled by *Dilp3-GAL4* are also

labeled by *PK2-R1$^{2A-LexA}$*. Similar results were obtained across five independent

samples. (B) Triple labeling of *PK2-R1* neurons expressing nuclear-localized RFP

(magenta), *Dilp5* neurons expressing nuclear-localized GFP (green), and anti-Dilp2

positive cells (blue). Similar results were obtained across five independent samples. (C)

Effect of IPCs silencing on larval sleep. In this and the following panels, 'N' indicates

the number of biologically independent animals used for each group. Box plots are

generated so that the center line indicates the median, the box limits indicate the upper

and lower quartiles, and the whiskers indicate the minimum-to-maximum range. *** p

< 0.001, ** p < 0.01, * p < 0.05 (Mann–Whitney U-test with Bonferroni correction).

(D) Sleep amounts in *Dilp3* or *Dilp5* null mutant larvae. *** p < 0.001, ** p < 0.01

(Mann–Whitney U-test with Bonferroni correction).

Figure 2—figure supplement 1. Locomotion speeds are not consistently affected by

genetical manipulations of *PK2-R1*, IPCs, or *Dilps*

(A–B) Effects of silencing *PK2-R1* neurons (A) and IPCs (B) on larval locomotion

speed. (C) Larval locomotion speed in null mutants of *Dilp3* or *Dilp5*. The average

locomotion speed during the wake periods was measured from the data presented in

Figures 1B, 2C, and 2D. In this and the following panels, 'N' indicates the number of

biologically independent animals used for each group. Box plots are generated so that

the center line indicates the median, the box limits indicate the upper and lower

quartiles, and the whiskers indicate the minimum-to-maximum range. *** p < 0.0001, *

p < 0.05, NS: p ≥ 0.05 (Mann–Whitney U-test with Bonferroni correction).

Figure 3. *Hug* neurons negatively regulate larval sleep

(A) Sleep amounts in *Hug* CRISPR-knock-out mutant larvae. In this and the following

panels, 'N' indicates the number of biologically independent animals used for each

group. Box plots are generated so that the center line indicates the median, the box

limits indicate the upper and lower quartiles, and the whiskers indicate the minimum-to-

maximum range. **** p < 0.0001 (Mann–Whitney U-test). (B) Effect of silencing *Hug*

neurons on larval sleep amount. *** p < 0.001 (Mann–Whitney U-test with Bonferroni

correction). (C) Larval sleep changes induced by optogenetic activation of *Hug* neurons.

For each genotype, sleep duration in the 1-h light-ON period was normalized to that in

the 1-h light-OFF phase. * p < 0.05 (Mann–Whitney U-test). (D) Larval sleep during

thermogenetic activation of *Hug* neurons. *** p < 0.001 (Mann–Whitney U-test with

Bonferroni correction). (E) Images of HugPC neurons and IPCs labeled by rCD2::RFP

(magenta) and mCD8::GFP (green), respectively. The bottom panels show magnified

images of the white-squared region in the top panels, in which HugPC neurons project

their axons close to the cell bodies of the IPCs. While the top panels are z-stacks of 122

image slices (1 μm interval) covering the entire brain, the bottom panels are projections

of 60 slices around the cell bodies of IPCs. Similar results were obtained across five

independent samples. (F) Effect of silencing HugPC neurons on larval sleep. *** p <

0.001, ** p < 0.01 (Mann–Whitney U-test with Bonferroni correction).

Figure 3—figure supplement 1. Neither *Hug* knock-out mutation nor *Hug* neuron

silencing significantly affects larval locomotion speed

(A) Average locomotion speed of *Hug* knock-out mutants during waking periods. The

data presented in Figures 3A and 3B were reanalyzed. In this and the following panels,

'N' indicates the number of biologically independent animals used for each group. Box

plots are generated so that the center line indicates the median, the box limits indicate

the upper and lower quartiles, and the whiskers indicate the minimum-to-maximum

range. NS: p ≥ 0.05 (Mann–Whitney U-test). (B) Locomotion speed of larvae in which

Hug neurons were silenced. NS: p ≥ 0.05 (Mann–Whitney U-test with Bonferroni

correction).

Figure 3—figure supplement 2. Generation of *HugPC-LexA* transgenic lines

(A–D) Labeling patterns of four *HugPC-LexA* strains in larval CNS visualized by

rCD2::RFP. These transgenic lines were generated by injecting the identical plasmid

into the following landing sites, respectively: su(Hw)attP5, VK00005, attP2, and attP40.

Arrowheads indicate cell bodies of non-HugPC neurons. Filled and unfilled arrowheads

indicate cell bodies within the VNC and near the protocerebrum, respectively.

Figure 4. Activation of *Hug* neurons triggers Ca^{2+} responses in larval IPCs

(A) Schematic flow of assessing intracellular Ca^{2+} levels in IPCs while performing the

thermogenetic activation of *Hug* neurons. (B–C) The level of nuclear-localized

CRTC::GFP signal, calculated as NLI (see the Methods section for a detailed explanation

of NLI), in IPCs during *Hug* neurons stimulation *ex vivo* (B) or *in vivo* (C). In these panels,

'N' indicates the number of cells used for each group. Box plots are generated so that the

center line indicates the median, the box limits indicate the upper and lower quartiles, and

the whiskers indicate the minimum-to-maximum range. **** $p < 0.0001$ (Mann–Whitney

U-test). (D) The proposed neuronal connection that regulates larval sleep.

Figure 5. Hug peptides induce Ca^{2+} responses in larval IPCs via PK2-R1

(A) Schematic flow of assessing intracellular Ca^{2+} levels in IPCs and bath application of

peptides. (B–C) Ca^{2+} responses in IPCs to *ex vivo* bath application of either glucose (B)

or Hug peptides (C). In this and the following panels, 'N' indicates the number of cells

used for each group. Box plots are generated so that the center line indicates the median,

667 the box limits indicate the upper and lower quartiles, and the whiskers indicate the

668 minimum-to-maximum range. **** $p < 0.0001$, *** $p < 0.001$ (Mann–Whitney U-test

669 with Bonferroni correction). (D) Representative images of larval IPCs after peptide

670 application. Scale bars, 2 μm. (E) Anti-Dilp3 signal intensity measured within the

671 cytosolic areas of IPCs. *** $p < 0.001$, * $p < 0.05$ (Mann–Whitney U-test with

672 Bonferroni correction). (F–G) Ca^{2+} responses in IPCs of *PK2-R1* knock-out mutants,

673 measured following either glucose (F) or Hug peptide application (G). ** $p < 0.01$, NS:

674 $p \geq 0.05$ (Mann–Whitney U-test with Bonferroni correction). (H) A model where Hug

675 peptides, but not glucose, activate IPCs via the PK2-R1 receptor.

677 **Figure 5—figure supplement 1. D-glucose but not Hug peptide cause Dilp2**

678 **reduction in IPCs**

679 (A–B) Anti-Dilp2 signal intensity measured in IPCs after glucose (A) or Hug-γ

680 application (B). In this and the following panels, 'N' indicates the number of cells used

681 for each group. Box plots are generated so that center line indicates median, box limits

682 indicate upper and lower quartiles, and whiskers indicate the minimum-to-maximum

683 range. **** $p < 0.0001$, * $p < 0.05$ (Mann–Whitney U-test). (C) Anti-Dilp3 signal

684 intensity measured within the cytosolic areas of IPCs. NS: ≥ 0.05 (Mann–Whitney U-

685 test).

687 **Figure 6. Distinct impacts of Hugin/PK2-R1 axis on wake/sleep control in larvae**

688 **and adults**

689 (A) Total sleep amounts in *Hug* knock-out mutant adults. NS: $p \geq 0.05$ (Mann–Whitney

U-test). In these and the following panels, 'N' indicates the number of biologically

independent animals used for each group. Box plots are generated so that the center line

indicates the median, the box limits indicate the upper and lower quartiles, and the

whiskers indicate the minimum-to-maximum range. NS: $p \geq 0.05$ (Mann–Whitney U-

test). (B) Sleep amounts of adults in which HugPC neurons were silenced. *** $p <$

0.001, NS: $p \geq 0.05$ (Mann–Whitney U-test with Bonferroni correction). (C) Sleep

amounts in *Dilp3* or *Dilp5* null mutant adults. *** $p < 0.001$ (Mann–Whitney U-test

with Bonferroni correction). (D) Schematic flow of peptide application experiments

using adult brains. (E–F) Ca^{2+} responses (E) and anti-Dilp3 signal intensity (F) in adult

IPCs after Hug peptide application. ** $p < 0.01$, NS: $p \geq 0.05$ (Mann–Whitney U-test

with Bonferroni correction).

Figure 6—figure supplement 1. Sleep patterns of adult flies following

manipulations of Hugin and insulin pathways

(A) Experimental setup for monitoring adult sleep using the *Drosophila* Activity

Monitor (DAM). (B) Sleep duration in adult CRISPR-knock-out mutants. These

mutants showed significant changes in the sleep amounts in larvae (Figure 1—figure

supplement 3). In panels B, C, and E, sleep amounts were calculated as the ratio relative

to the median sleep amount of the control group. Box plots are generated so that the

center line indicates the median, the box limits indicate the upper and lower quartiles,

and the whiskers indicate the minimum-to-maximum range. *** $p < 0.001$, ** $p < 0.01$

(Kruskal–Wallis one-way ANOVA and post hoc Mann–Whitney U-test with Bonferroni

correction). (C, E) Sleep amounts relative to the median value of the control group. Data

presented in Figures 6B and 6C were reanalyzed. *** $p < 0.001$, * $p < 0.05$, NS: $p \geq$

0.05 (Mann–Whitney U-test with Bonferroni correction). (D, F) Temporal sleep patterns

over two consecutive days.

Figure 6—figure supplement 2. Morphology of *PK2-R1* neurons, HugPC neurons,

and IPCs in the adult brain

(A) Dual labeling of *PK2-R1* neurons and IPCs expressing nuclear-localized RFP

(stinger; magenta) and GFP (stinger; green), respectively. The top panels show signals

in the entire adult brain. The bottom panels show magnified images of the white-

squared area in each corresponding top panel, where the cell bodies of IPCs are located.

Similar results were obtained across five independent samples. (B) Visualization of

HugPC neurons and IPCs labeled by mCD8::GFP (magenta) and the anti-Dilp3

antibody (green), respectively. Similar results were obtained across four independent

samples.

Figure 7. A schematic model of the Hugin/PK2-R1/Dilps axis in larval and adult

sleep regulation

References

Ahnaou, A., & Drinkenburg, W. H. I. M. (2011). Neuromedin U2 receptor signaling
mediates alteration of sleep-wake architecture in rats. *Neuropeptides*, 45(2),
165–174. Retrieved from https://doi.org/10.1016/j.npep.2011.01.004

Bader, R., Colomb, J., Pankratz, B., Schröck, A., Stocker, R. F., & Pankratz, M. J.
(2007). Genetic dissection of neural circuit anatomy underlying feeding
behavior in Drosophila: Distinct classes of hugin-expressing neurons. *Journal*
of Comparative Neurology, 502(5), 848–856. Retrieved from
https://doi.org/10.1002/cne.21342

Baines, R. A., Uhler, J. P., Thompson, A., Sweeney, S. T., & Bate, M. (2001). *Altered*
Electrical Properties in Drosophila Neurons Developing without Synaptic
Transmission.

Barlow, I. L., & Rihel, J. (2017, June 1). Zebrafish sleep: from geneZZZ to
neuronZZZ. *Current Opinion in Neurobiology*. Elsevier Ltd. Retrieved from
https://doi.org/10.1016/j.conb.2017.02.009

Blumberg, M. S., H Seelke, A. M., Lowen, S. B., & Karlsson, K. A. (2005).
Dynamics of sleep–wake cyclicity in developing rats. *PNAS*, 102(41).
Retrieved from www.pnas.orgcgidoi10.1073pnas.0506340102

Bonheur, M., Swartz, K. J., Metcalf, M. G., Wen, X., Zhukovskaya, A., Mehta, A.,
… Hattori, D. (2023). A rapid and bidirectional reporter of neural activity
reveals neural correlates of social behaviors in Drosophila. *Nature*
Neuroscience, 26(7), 1295–1307. Retrieved from
https://doi.org/10.1038/s41593-023-01357-w

Burke, C. J., Huetteroth, W., Owald, D., Perisse, E., Krashes, M. J., Das, G., …
Waddell, S. (2012). Layered reward signalling through octopamine and
dopamine in Drosophila. *Nature*, 492(7429), 433–437. Retrieved from
https://doi.org/10.1038/nature11614

Chiu, C. N., Rihel, J., Lee, D. A., Singh, C., Mosser, E. A., Chen, S., … Prober, D.
A. (2016). A Zebrafish Genetic Screen Identifies Neuromedin U as a Regulator
of Sleep/Wake States. *Neuron*, 89(4), 842–856. Retrieved from
https://doi.org/10.1016/j.neuron.2016.01.007

Churgin, M. A., Szuperak, M., Davis, K. C., Raizen, D. M., Fang-Yen, C., & Kayser,
M. S. (2019). Quantitative imaging of sleep behavior in Caenorhabditis elegans
and larval Drosophila melanogaster. *Nature Protocols 2019 14:5*, 14(5), 1455–
1488. Retrieved 25 December 2021 from https://doi.org/10.1038/s41596-019-
0146-6

Davis, K. F., Parker, K. P., & Montgomery, G. L. (2004). *Sleep in infants and young children: Part one: normal sleep. Journal of Pediatric Health Care* (Vol. 18, pp. 65–71). Mosby. Retrieved 25 December 2021 from https://doi.org/10.1016/S0891-5245(03)00149-4

Deng, B., Li, Q., Liu, X., Cao, Y., Li, B., Qian, Y., … Rao, Y. (2019). Chemoconnectomics: Mapping Chemical Transmission in Drosophila. *Neuron*, 101(5), 876-893.e4. Retrieved from https://doi.org/10.1016/j.neuron.2019.01.045

Dubowy, C., & Sehgal, A. (2017). Circadian rhythms and sleep in Drosophila melanogaster. *Genetics*, 205(4), 1373–1397. Retrieved from https://doi.org/10.1534/genetics.115.185157

Enell, L. E., Kapan, N., Söderberg, J. A. E., Kahsai, L., & Nässe, D. R. (2010). Insulin signaling, lifespan and stress resistance are modulated by metabotropic GABA receptors on insulin producing cells in the brain of Drosophila. *PLoS ONE*, 5(12). Retrieved from https://doi.org/10.1371/journal.pone.0015780

Frank, M. G., Ruby, N. F., Heller, H. C., & Franken, P. (2017). Development of circadian sleep regulation in the rat: A longitudinal study under constant conditions. *Sleep*, 40(3). Retrieved from https://doi.org/10.1093/sleep/zsw077

Géminard, C., Rulifson, E. J., & Léopold, P. (2009). Remote Control of Insulin Secretion by Fat Cells in Drosophila.
Cell Metabolism, 10(3), 199–207. Retrieved from https://doi.org/10.1016/j.cmet.2009.08.002

Groth, A. C., Fish, M., Nusse, R., & Calos, M. P. (2004). Construction of Transgenic Drosophila by Using the Site-Specific Integrase From Phage φC31. *Genetics*, (166), 1775–1782. Retrieved from https://academic.oup.com/genetics/article/166/4/1775/6052665

Hamada, F. N., Rosenzweig, M., Kang, K., Pulver, S. R., Ghezzi, A., Jegla, T. J., & Garrity, P. A. (2008). An internal thermal sensor controlling temperature preference in Drosophila. *Nature*, 454(7201), 217–220. Retrieved from https://doi.org/10.1038/nature07001

Hendricks, J. C., Lu, S., Kume, K., Yin, J. C. P., Yang, Z., & Sehgal, A. (2003). Gender dimorphism in the role of cycle (BMAL1) in rest, rest regulation, and longevity in Drosophila melanogaster. *Journal of Biological Rhythms*, 18(1), 12–25. Retrieved from https://doi.org/10.1177/0748730402239673

Hückesfeld, S., Peters, M., & Pankratz, M. J. (2016). Central relay of bitter taste to the protocerebrum by peptidergic interneurons in the Drosophila brain. *Nature*

Communications, 7. Retrieved from https://doi.org/10.1038/ncomms12796

Hückesfeld, S., Schlegel, P., Miroschnikow, A., Schoofs, A., Zinke, I., Haubrich, A.
N., … Pankratz, M. J. (2021). Unveiling the sensory and interneuronal
pathways of the neuroendocrine connectome in Drosophila. *ELife*. Retrieved
from https://doi.org/10.7554/eLife

Inagaki, H. K., Jung, Y., Hoopfer, E. D., Wong, A. M., Mishra, N., Lin, J. Y., …
Anderson, D. J. (2014). Optogenetic control of Drosophila using a red-shifted
channelrhodopsin reveals experience-dependent influences on courtship.
Nature Methods, 11(3), 325–332. Retrieved from
https://doi.org/10.1038/nmeth.2765

Kauffman, M. R., & DiAngelo, J. R. (2024). Glut1 Functions in Insulin-Producing
Neurons to Regulate Lipid and Carbohydrate
Storage in Drosophila. *Biomolecules*, 14(8). Retrieved from
https://doi.org/10.3390/biom14081037

Kréneisz, O., Chen, X., Fridell, Y. W. C., & Mulkey, D. K. (2010). Glucose increases
activity and Ca2+ in insulin-producing cells of adult Drosophila. *NeuroReport*,
21(17), 1116–1120. Retrieved from
https://doi.org/10.1097/WNR.0b013e3283409200

Kume, K., Kume, S., Park, S. K., Hirsh, J., & Jackson, F. R. (2005). Dopamine Is a
Regulator of Arousal in the Fruit Fly. *Journal of Neuroscience*, 25(32), 7377–
7384. Retrieved 10 December 2024 from
https://doi.org/10.1523/JNEUROSCI.2048-05.2005

Nakamizo-Dojo, M., Ishii, K., Yoshino, J., Tsuji, M., & Emoto, K. (2023).
Descending GABAergic pathway links brain sugar-sensing to peripheral
nociceptive gating in Drosophila. *Nature Communications*, 14(1). Retrieved
from https://doi.org/10.1038/s41467-023-42202-9

Nässel, D. R., Liu, Y., & Luo, J. (2015). Insulin/IGF signaling and its regulation in
Drosophila. *General and Comparative Endocrinology*, 221, 255–266.
Retrieved from https://doi.org/10.1016/j.ygcen.2014.11.021

Oh, Y., Lai, J. S. Y., Mills, H. J., Erdjument-Bromage, H., Giammarinaro, B.,
Saadipour, K., … Suh, G. S. B. (2019). A glucose-sensing neuron pair regulates
insulin and glucagon in Drosophila. *Nature*, 574(7779), 559–564. Retrieved
from https://doi.org/10.1038/s41586-019-1675-4

Ohhara, Y., Blick, M., Park, D., Yoon, S. E., Kim, Y. J., Pankratz, M. J., … Yamanaka,
N. (2024). A Neuropeptide Signaling Network That Regulates Developmental
Timing and Systemic Growth in Drosophila. *The Journal of Comparative*

Neurology, 532(10), e25677. Retrieved from https://doi.org/10.1002/cne.25677

Okamoto, N. (2021). Insect insulin-like peptides. In *Handbook of Hormones:*
Comparative Endocrinology for Basic and Clinical
Research (pp. 673–675). Elsevier. Retrieved from
https://doi.org/10.1016/B978-0-12-820649-2.00177-7

Okamoto, N., Nakamori, R., Murai, T., Yamauchi, Y., Masuda, A., & Nishimura, T.
(2013). A secreted decoy of InR antagonizes insulin/IGF signaling to restrict
body growth in drosophila. *Genes and Development*, 27(1), 87–97. Retrieved
from https://doi.org/10.1101/gad.204479.112

Okamoto, N., Nishimori, Y., & Nishimura, T. (2012). Conserved role for the
Dachshund protein with Drosophila Pax6 homolog eyeless in insulin
expression. *Proceedings of the National Academy of Sciences of the United*
States of America, 109(7), 2406–2411. Retrieved from
https://doi.org/10.1073/pnas.1116050109

Paul J. Shaw, Chiara Cirelli, Ralph J. Greenspan, & Giulio Tononi. (2000).
Correlates of Sleep and Waking in Drosophila melanogaster. *SCIENCE*, (287),
1834–1837.

Poe, A. R., Zhu, L., Szuperak, M., Mcclanahan, P. D., Anafi, R. C., Scholl, B., …
Kayser, M. S. (2023). *Developmental emergence of sleep rhythms enables long-*
term memory in Drosophila. Retrieved from https://www.science.org

Raizen, D. M., Zimmerman, J. E., Maycock, M. H., Ta, U. D., You, Y. J., Sundaram,
M. V., & Pack, A. I. (2008). Lethargus is a Caenorhabditis elegans sleep-like
state. *Nature*, 451(7178), 569–572. Retrieved from
https://doi.org/10.1038/nature06535

Rosenkilde, C., Cazzamali, G., Williamson, M., Hauser, F., Søndergaard, L.,
DeLotto, R., & Grimmelikhuijzen, C. J. P. (2003). Molecular cloning,
functional expression, and gene silencing of two Drosophila receptors for the
Drosophila neuropeptide pyrokinin-2. *Biochemical and Biophysical Research*
Communications, 309(2), 485–494. Retrieved from
https://doi.org/10.1016/j.bbrc.2003.08.022

Scammell, T. E., Arrigoni, E., & Lipton, J. O. (2017, February 22). Neural Circuitry
of Wakefulness and Sleep. *Neuron*. Cell Press. Retrieved from
https://doi.org/10.1016/j.neuron.2017.01.014

Schlegel, P., Texada, M. J., Miroschnikow, A., Schoofs, A., Hückesfeld, S., Peters,
M., … Pankratz, M. J. (2016a). Synaptic transmission parallels
neuromodulation in a central food-intake circuit. *ELife*, 5(NOVEMBER2016).

Retrieved from https://doi.org/10.7554/eLife.16799.001

Schlegel, P., Texada, M. J., Miroschnikow, A., Schoofs, A., Hückesfeld, S., Peters, M., … Pankratz, M. J. (2016b). Synaptic transmission parallels neuromodulation in a central food-intake circuit. *ELife*, 5(NOVEMBER2016). Retrieved from https://doi.org/10.7554/eLife.16799.001

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nature Methods*, 9(7), 671–675. Retrieved from https://doi.org/10.1038/nmeth.2089

Schoofs, A., Hückesfeld, S., Schlegel, P., Miroschnikow, A., Peters, M., Zeymer, M., … Pankratz, M. J. (2014). Selection of Motor Programs for Suppressing Food Intake and Inducing Locomotion in the Drosophila Brain. *PLoS Biology*, 12(6). Retrieved from https://doi.org/10.1371/journal.pbio.1001893

Schwarz, J. E., King, A. N., Hsu, C. T., Barber, A. F., & Sehgal, A. (2021). Hugin + neurons provide a link between sleep homeostat and circadian clock neurons. *PNAS*. Retrieved from https://doi.org/10.1073/pnas.2111183118/-/DCSupplemental

Shafer, O. T., & Keene, A. C. (2021, January 11). The Regulation of Drosophila Sleep. *Current Biology*. Cell Press. Retrieved from https://doi.org/10.1016/j.cub.2020.10.082

Sorribes, A., Porsteinsson, H., Arnardóttir, H., Jóhannesdóttir, I. P., Sigurgeirsson, B., de Polavieja, G. G., & Karlsson, K. (2013). The ontogeny of sleep-wake cycles in zebrafish: A comparison to humans. *Frontiers in Neural Circuits*, 7(NOV). Retrieved from https://doi.org/10.3389/fncir.2013.00178

Szuperak, M., Churgin, M. A., Borja, A. J., Raizen, D. M., Fang-Yen, C., & Kayser, M. S. (2018a). A sleep state in Drosophila larvae required for neural stem cell proliferation. *ELife*, 7. Retrieved 25 December 2021 from https://doi.org/10.7554/ELIFE.33220

Szuperak, M., Churgin, M. A., Borja, A. J., Raizen, D. M., Fang-Yen, C., & Kayser, M. S. (2018b). A sleep state in Drosophila larvae required for neural stem cell proliferation. *ELife*. Retrieved from https://doi.org/10.7554/eLife.33220.001

Tomoatsu Ikeya, Milos Galic, Priyanka Belawat, Knud Nairz, & Ernst Hafen. (2002). *Nutrient-Dependent Expression of Insulin-like Peptides from Neuroendocrine Cells in the CNS Contributes to Growth Regulation in Drosophila median neurosecretory cells (m-NSCs) possessing axon*.

Veenstra, J. A., Agricola, H. J., & Sellami, A. (2008). Regulatory peptides in fruit fly

911   midgut. *Cell and Tissue Research*, 334(3), 499–516. Retrieved from
912   https://doi.org/10.1007/s00441-008-0708-3

915 **Figures**

Figure 1

Figure 1. PK2-R1 is required for larval sleep control

(A) Sleep amounts in *PK2-R1* or *Oamb* knock-out mutants. In this and the following

panels, 'N' indicates the number of biologically independent animals used for each

group. Box plots are generated so that the center line indicates the median, the box

limits indicate the upper and lower quartiles, and the whiskers indicate the minimum-to-

maximum range. *** $p < 0.001$ (Mann–Whitney U-test with Bonferroni correction). (B)

Sleep amounts in larvae in which *PK2-R1* neurons were silenced. *** $p < 0.001$ (Mann–

Whitney U-test with Bonferroni correction). (C) The expression pattern of *PK2-R1²A-*

GAL4 > UAS-Kir2.1::EGFP in the larval CNS. Similar results were obtained across three

independent samples.

Figure 1—figure supplement 1

Figure 1—figure supplement 1. Detailed experimental setup for analyzing larval

sleep

(A) Schematic flow of automated larval sleep quantification. A second instar larva was

placed in each of 24 wells, and all 24 wells were videotaped from above. For each well

and frame, the image of larva was blurred and binarized to automatically locate the

larva. Consecutive frames were then compared so that changes in pixel values were

summed to estimate larval motion. This motion estimate was then used to classify every

consecutive pair of frames as representing "active" or "inactive." We found that

inactivity across 12 consecutive frames meets the sleep criteria, as shown in Figure 1—

figure supplement 2. (B) Schematic representation of the 24-well chamber used for

larval sleep assays. (C) Flow of automated processes of larval motion detection and

active/inactive state determination. For each well and frame, the image of larva was

blurred and binarized. Consecutive frames were then compared so that changes in pixel

values were summed to estimate larval motion. This motion estimate was then used to

classify every consecutive pair of frames as representing "active" or "inactive."

Figure 1—figure supplement 2

Figure 1—figure supplement 2. Larval "sleep state" in this study is consistent with behavioral criteria for sleep

(A) Relationship between "inactive" state duration and spontaneous arousal probability. Sustained quiescence for ≥ 12 frames (open circles) significantly reduced arousal

probability compared to 1-frame inactivity. * $p < 0.05$, NS: $p \geq 0.05$ (chi-squared test). (B) The percentage of larvae that showed a response within the next 10 frames after high-intensity blue light. The number of events analyzed was 36 and 60 with and without blue light irradiation, respectively. * $p < 0.05$ (Fisher's exact test). (C) Sleep amounts before, during, and after blue light irradiation. The number of larvae tested was 21 and 21 for the control and experimental groups, respectively. ** $p < 0.01$, NS: $p \geq 0.05$ (Mann–Whitney U-test with Bonferroni correction). (D) Percentage of larvae that showed a response in the next 1 or 12 frames after blue light stimulation. Number of events analyzed: 1960 or 111 for 1- or 12-frame continuous inactivity prior to blue light illumination, respectively. **** $p < 0.0001$ (chi-squared test).

Figure 1—figure supplement 3. CRISPR-knock-out screen for genes that regulate larval sleep.

Total sleep amounts in CRISPR-knock-out mutants were measured for 18 h from the beginning of the second instar. The data for Control, *PK2-R1,* and *Oamb* mutants are identical to those shown in Figure 1. Box plots are generated so that the center line indicates the median, the box limits indicate the upper and lower quartiles, and the whiskers indicate the minimum-to-maximum range. *** $p < 0.005$ (Kruskal–Wallis one-way ANOVA and post hoc Mann–Whitney U-test with Bonferroni correction). Number of larvae tested: 475 for the control and 10–52 for each mutant.

Figure 1—figure supplement 4. Effects of neuronal manipulation on larval sleep
Sleep amounts in larvae in which distinct neuronal populations were silenced. Kir2.1
expression was driven by each enhancer-GAL4 for "hit" genes identified in the
CRISPR-knock-out screen. 'N' indicates the number of biologically independent
animals used for each group. Box plots are generated so that the center line indicates the
median, the box limits indicate the upper and lower quartiles, and the whiskers indicate
the minimum-to-maximum range. *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, NS: ≥ 0.05
(Mann–Whitney U-test with Bonferroni correction).

Figure 2

Figure 2. IPCs and Dilps negatively regulate larval sleep

(A) Triple labeling of *PK2-R1* neurons expressing nuclear-localized RFP (magenta),

Dilp3 neurons expressing nuclear-localized GFP (green), and anti-Dilp2 positive cells

(blue). Top panels show signals in the larval brain and the VNC. The bottom panels

show magnified images of the white-squared area in the top panels, where the cell

bodies of the IPCs are located. Note that all IPCs labeled by *Dilp3-GAL4* are also

labeled by *PK2-R1^{2A-LexA}*. Similar results were obtained across five independent

samples. (B) Triple labeling of *PK2-R1* neurons expressing nuclear-localized RFP

(magenta), *Dilp5* neurons expressing nuclear-localized GFP (green), and anti-Dilp2

positive cells (blue). Similar results were obtained across five independent samples. (C)

Effect of IPCs silencing on larval sleep. In this and the following panels, 'N' indicates

the number of biologically independent animals used for each group. Box plots are

generated so that the center line indicates the median, the box limits indicate the upper

and lower quartiles, and the whiskers indicate the minimum-to-maximum range. *** p

< 0.001, ** $p < 0.01$, * $p < 0.05$ (Mann–Whitney U-test with Bonferroni correction).

(D) Sleep amounts in *Dilp3* or *Dilp5* null mutant larvae. *** $p < 0.001$, ** $p < 0.01$

(Mann–Whitney U-test with Bonferroni correction).

1004

1005

Figure 2—figure supplement 1

Figure 2—figure supplement 1. **Locomotion speeds are not consistently affected by genetical manipulations of *PK2-R1*, IPCs, or *Dilps***

(A–B) Effects of silencing *PK2-R1* neurons (A) and IPCs (B) on larval locomotion speed. (C) Larval locomotion speed in null mutants of *Dilp3* or *Dilp5*. The average locomotion speed during the wake periods was measured from the data presented in Figures 1B, 2C, and 2D. In this and the following panels, 'N' indicates the number of biologically independent animals used for each group. Box plots are generated so that the center line indicates the median, the box limits indicate the upper and lower quartiles, and the whiskers indicate the minimum-to-maximum range. *** $p < 0.0001$, * $p < 0.05$, NS: $p \geq 0.05$ (Mann–Whitney U-test with Bonferroni correction).

Figure 3

1018

Figure 3. *Hug* neurons negatively regulate larval sleep

(A) Sleep amounts in *Hug* CRISPR-knock-out mutant larvae. In this and the following

panels, 'N' indicates the number of biologically independent animals used for each

group. Box plots are generated so that the center line indicates the median, the box

limits indicate the upper and lower quartiles, and the whiskers indicate the minimum-to-

maximum range. **** $p < 0.0001$ (Mann–Whitney U-test). (B) Effect of silencing *Hug*

neurons on larval sleep amount. *** $p < 0.001$ (Mann–Whitney U-test with Bonferroni

correction). (C) Larval sleep changes induced by optogenetic activation of *Hug* neurons.

For each genotype, sleep duration in the 1-h light-ON period was normalized to that in

the 1-h light-OFF phase. * $p < 0.05$ (Mann–Whitney U-test). (D) Larval sleep during

thermogenetic activation of *Hug* neurons. *** $p < 0.001$ (Mann–Whitney U-test with

Bonferroni correction). (E) Images of HugPC neurons and IPCs labeled by rCD2::RFP

(magenta) and mCD8::GFP (green), respectively. The bottom panels show magnified

images of the white-squared region in the top panels, in which HugPC neurons project

their axons close to the cell bodies of the IPCs. While the top panels are z-stacks of 122

image slices (1 µm interval) covering the entire brain, the bottom panels are projections

of 60 slices around the cell bodies of IPCs. Similar results were obtained across five

independent samples. (F) Effect of silencing HugPC neurons on larval sleep. *** $p <$

0.001, ** $p < 0.01$ (Mann–Whitney U-test with Bonferroni correction).

1038

Figure 3—figure supplement 1

1039

Figure 3—figure supplement 1. Neither *Hug* knock-out mutation nor *Hug* neuron silencing significantly affects larval locomotion speed

(A) Average locomotion speed of *Hug* knock-out mutants during waking periods. The data presented in Figures 3A and 3B were reanalyzed. In this and the following panels, 'N' indicates the number of biologically independent animals used for each group. Box plots are generated so that the center line indicates the median, the box limits indicate the upper and lower quartiles, and the whiskers indicate the minimum-to-maximum range. NS: p ≥ 0.05 (Mann–Whitney U-test). (B) Locomotion speed of larvae in which *Hug* neurons were silenced. NS: p ≥ 0.05 (Mann–Whitney U-test with Bonferroni correction).

1050

Figure 3—figure supplement 2

A

Figure 3—figure supplement 2. Generation of *HugPC-LexA* transgenic lines

(A–D) Labeling patterns of four *HugPC-LexA* strains in larval CNS visualized by rCD2::RFP. These transgenic lines were generated by injecting the identical plasmid into the following landing sites, respectively: su(Hw)attP5, VK00005, attP2, and attP40. Arrowheads indicate cell bodies of non-HugPC neurons. Filled and unfilled arrowheads indicate cell bodies within the VNC and near the protocerebrum, respectively.

Figure 4

A

Figure 4. Activation of *Hug* neurons triggers Ca²⁺ responses in larval IPCs

(A) Schematic flow of assessing intracellular Ca²⁺ levels in IPCs while performing the thermogenetic activation of *Hug* neurons. (B–C) The level of nuclear-localized CRTC::GFP signal, calculated as NLI (see the Methods section for a detailed explanation of NLI), in IPCs during *Hug* neurons stimulation *ex vivo* (B) or *in vivo* (C). In these panels, 'N' indicates the number of cells used for each group. Box plots are generated so that the center line indicates the median, the box limits indicate the upper and lower quartiles, and the whiskers indicate the minimum-to-maximum range. **** $p < 0.0001$ (Mann–Whitney U-test). (D) The proposed neuronal connection that regulates larval sleep.

Figure 5

Figure 5. Hug peptides induce Ca^{2+} responses in larval IPCs via PK2-R1

(A) Schematic flow of assessing intracellular Ca^{2+} levels in IPCs and bath application of peptides. (B–C) Ca^{2+} responses in IPCs to *ex vivo* bath application of either glucose (B) or Hug peptides (C). In this and the following panels, 'N' indicates the number of cells used for each group. Box plots are generated so that the center line indicates the median, the box limits indicate the upper and lower quartiles, and the whiskers indicate the minimum-to-maximum range. **** $p < 0.0001$, *** $p < 0.001$ (Mann–Whitney U-test with Bonferroni correction). (D) Representative images of larval IPCs after peptide application. Scale bars, 2 μm. (E) Anti-Dilp3 signal intensity measured within the

cytosolic areas of IPCs. *** $p < 0.001$, * $p < 0.05$ (Mann–Whitney U-test with Bonferroni correction). (F–G) Ca^{2+} responses in IPCs of *PK2-R1* knock-out mutants, measured following either glucose (F) or Hug peptide application (G). ** $p < 0.01$, NS: $p \geq 0.05$ (Mann–Whitney U-test with Bonferroni correction). (H) A model where Hug peptides, but not glucose, activate IPCs via the PK2-R1 receptor.

Figure 5—figure supplement 1

Figure 5—figure supplement 1. D-glucose but not Hug peptide cause Dilp2 reduction in IPCs

(A–B) Anti-Dilp2 signal intensity measured in IPCs after glucose (A) or Hug-γ application (B). In this and the following panels, 'N' indicates the number of cells used for each group. Box plots are generated so that center line indicates median, box limits indicate upper and lower quartiles, and whiskers indicate the minimum-to-maximum range. **** $p < 0.0001$, * $p < 0.05$ (Mann–Whitney U-test). (C) Anti-Dilp3 signal intensity measured within the cytosolic areas of IPCs. NS: ≥ 0.05 (Mann–Whitney U-test).

Figure 6

Figure 6. Distinct impacts of Hugin/PK2-R1 axis on wake/sleep control in larvae and adults

(A) Total sleep amounts in *Hug* knock-out mutant adults. NS: $p \geq 0.05$ (Mann–Whitney U-test). In these and the following panels, 'N' indicates the number of biologically independent animals used for each group. Box plots are generated so that the center line indicates the median, the box limits indicate the upper and lower quartiles, and the whiskers indicate the minimum-to-maximum range. NS: $p \geq 0.05$ (Mann–Whitney U-test). (B) Sleep amounts of adults in which HugPC neurons were silenced. *** $p < 0.001$, NS: $p \geq 0.05$ (Mann–Whitney U-test with Bonferroni correction). (C) Sleep amounts in *Dilp3* or *Dilp5* null mutant adults. *** $p < 0.001$ (Mann–Whitney U-test with Bonferroni correction). (D) Schematic flow of peptide application experiments using adult brains. (E–F) Ca^{2+} responses (E) and anti-Dilp3 signal intensity (F) in adult

1113   IPCs after Hug peptide application. ** $p < 0.01$, NS: $p \geq 0.05$ (Mann–Whitney U-test

1114   with Bonferroni correction).

1115

1116

1117

Figure 6—figure supplement 1. Sleep patterns of adult flies following manipulations of Hugin and insulin pathways

(A) Experimental setup for monitoring adult sleep using the *Drosophila* Activity Monitor (DAM). (B) Sleep duration in adult CRISPR-knock-out mutants. These mutants showed significant changes in the sleep amounts in larvae (Figure 1—figure supplement 3). In panels B, C, and E, sleep amounts were calculated as the ratio relative to the median sleep amount of the control group. Box plots are generated so that the center line indicates the median, the box limits indicate the upper and lower quartiles, and the whiskers indicate the minimum-to-maximum range. *** $p < 0.001$, ** $p < 0.01$ (Kruskal–Wallis one-way ANOVA and post hoc Mann–Whitney U-test with Bonferroni

correction). (C, E) Sleep amounts relative to the median value of the control group. Data presented in Figures 6B and 6C were reanalyzed. *** $p < 0.001$, * $p < 0.05$, NS: $p \geq 0.05$ (Mann–Whitney U-test with Bonferroni correction). (D, F) Temporal sleep patterns over two consecutive days.

Figure 6—figure supplement 2

1136

1137

Figure 6—figure supplement 2. Morphology of *PK2-R1* neurons, HugPC neurons, and IPCs in the adult brain

(A) Dual labeling of *PK2-R1* neurons and IPCs expressing nuclear-localized RFP (stinger; magenta) and GFP (stinger; green), respectively. The top panels show signals in the entire adult brain. The bottom panels show magnified images of the white-squared area in each corresponding top panel, where the cell bodies of IPCs are located. Similar results were obtained across five independent samples. (B) Visualization of HugPC neurons and IPCs labeled by mCD8::GFP (magenta) and the anti-Dilp3 antibody (green), respectively. Similar results were obtained across four independent samples.

Figure 7

Figure 7. A schematic model of the Hugin/PK2-R1/Dilps axis in larval and adult sleep regulation