1

Neuropeptidergic circuit modulation
of developmental sleep in *Drosophila*

Chikayo Hemmi[1], Kenichi Ishii[1, #], Mana Motoyoshi[1], Masato Tsuji[1]

and Kazuo Emoto[1, 2, #, $]

[1]Department of Biological Sciences, The University of Tokyo

[2]International Research Center for Neurointelligence (WPI-IRCN), The University of Tokyo

[#]Corresponding authors

[$]Lead contact: emoto@bs.s.u-tokyo.ac.jp

Abstract

Sleep-wakefulness regulation dynamically evolves along development in a wide range of organisms. While the mechanism regulating sleep in adults are relatively well understood, little is known about its counterpart in early developmental stages. Here we report a neuropeptidergic circuitry that modulates sleep in developing *Drosophila* larvae. Through an unbiased screen, we identified the neuropeptide Hugin and its receptor PK2-R1 as critical regulators of larval sleep. Our genetic and behavioral data suggest that HugPC neurons secrete Hugin peptides to activate Insulin-producing cells (IPCs), which expresses a Hugin receptor PK2-R1. IPCs, in turn, release insulin-like peptides (Dilps) to regulate sleep. We further show that the Hugin/PK2-R1 axis is dispensable for adult sleep. Our findings thus reveal the neuromodulatry circuit that regulates developmental sleep in larvae, and highlight differential impacts of the same modulatory axis on early-life sleep and adult sleep.

Introduction

Sleep is a fundamental state with significant impacts on many aspects of cognition and metabolism (Blumberg et al., 2005; Davis et al., 2004; Sorribes et al., 2013). While sleep remains essential throughout life, its regulation dynamically unfolds over the course of development. For instance, the pace of sleep-wakefulness cycle is relatively fast in infants, while gradually slows down and stabilize in adults (Blumberg et al., 2005; Davis et al., 2004; Sorribes et al., 2013). Furthermore, sleep-wake cycle in infants is typically independent of circadian rhythm, but it gradually comes under the circadian regulation (Davis et al., 2004; Frank et al., 2017; Poe et al., 2023). Such developmental evolution of sleep-wakefulness cycles appears to be conserved across a wide range of organisms including flies, fishes, mammals and humans (Blumberg et al., 2005; Davis et al., 2004; Poe et al., 2023; Sorribes et al., 2013; Szuperak et al., 2018a). In contrast to deep insights into the molecular and neural mechanisms underlying adult sleep regulation (Barlow & Rihel, 2017; Scammell et al., 2017; Shafer & Keene, 2021), little is known about regulatory mechanisms underlying developmental sleep in infants, partially due to a lack of convenient models to study developmental sleep.

Drosophila larvae have recently emerged as a suitable model to study sleep regulation
mechanisms. Recent studies have reported that the second instar larvae show short
periods (< 6 seconds) of "inactive state" during locomotion. Importantly, such "inactive
state" is consistent with the general definition of sleep: reduced responsiveness to
noxious stimuli, homeostatic response to sleep deprivation, and rapid reversibility upon
stimulation (Szuperak et al., 2018), suggesting that the "inactive state" in larvae likely
corresponds to the sleep state. Interestingly, loss-of-function mutations in the clock
genes *clock* and *cyc* fail to impact the larval sleep, while significantly altering the sleep
patterns in adult flies (Dubowy & Sehgal, 2017; Hendricks et al., 2003; Szuperak et al.,
2018). Likewise, mutations in the dopamine transporter *DAT* fail to influence larval
sleep, while reducing sleep amounts in adult (Kume et al., 2005; Szuperak et al., 2018).
These observations imply that sleep regulation mechanisms might be at least in part
distinct between larvae and adults, yet the neural mechanism of larval sleep remains
vastly understudied compared to that of adult sleep.

In this study, we performed an unbiased genetic screen in the second instar larvae, and
identified the neuropeptide Hugin and its receptor PK2-R1 as a pair critical for larval
sleep. At the circuit level, Hugin-producing HugPC neurons directly stimulate the
insulin-producing cells (IPCs) via PK2-R1 to regulate sleep. We further found that
regulators of larval sleep are either irrelevant or exert the opposite effects on adult sleep.
Overall, our findings uncover the neuropeptidergic circuitry that regulates
developmental sleep in larvae, and highlight mechanistic differences between larval and
adult sleep.

Results

PK2-R1 is required for wake/sleep control in Drosophila larvae

To understand molecular and neural mechanisms of larval sleep, we modified a
previously reported system (Churgin et al., 2019; Szuperak et al., 2018) to automatically
quantify the sleep amount of second instar larvae (Figure 1-figure supplement 1A-1C) (see
Materials & Methods for details). Our system automatically annotates larvae in each frame as

either "inactive" or "active" states on the basis of pixel changes, with ~90% accuracy compared

to manual annotation (Figure 1-figure supplement 1C; see Methods for details). We found that

bouts of ≥ 12 inactive frames are consistent with the general criteria for sleep; reduced

responsiveness to noxious stimuli, homeostatic response to sleep loss, and rapid reversibility

upon stimulation (Raizen et al., 2008; Szuperak et al., 2018) (Figure 1-figure supplement

2B–D). Therefore, the present study defines sleep as inactive state of ≥ 12 consecutive

frames (Figure 1-figure supplement 1A).

Using this system, we next carried out an unbiased screen for genes involved in sleep

regulation. To this end, we tested 47 CRISPR knock-out lines of genes encoding

enzymes or receptors for neuropeptides and monoamines (Deng et al.,

2019)(Supplemental Table 1). As illustrated in Figure 1-figure supplement 3, 13 out of

47 homozygous null mutants exhibited significant changes in sleep amounts compared

to the control group. To further test the functions of these candidate genes, we next took

advantage of the CRISPR knock-in drivers in which GAL4 sequence is inserted into

each gene locus. With these drivers, we silenced neurons expressing each candidate

gene by expressing the inward-rectifying potassium channel Kir2.1 (Baines et al.,

2001). Silencing neurons expressing *Oamb* and *PK2-R1* resulted in significant sleep

increase, phenocopying the knock-out of these genes (Figure 1A and 1B and Figure 1-

figure supplement 4D). *Oamb* encodes the octopamine receptor and has been reported to

be involved in larval sleep (Szuperak et al., 2018). The role of PK2-R1 in larval sleep,

on the other hand, has been unknown to date. Given its strong expression in insulin-

producing cells (Schlegel et al., 2016) and its function as a receptor for the neuropeptide

Hugin, which modulates feeding (Schoofs et al., 2014), we hypothesized that PK2-R1

might mediate neuropeptidergic signaling that links metabolic and sleep regulation

during development. We thus focused on this gene as a candidate connecting behavioral

and endocrine sleep control.

Insulin-producing cells express PK2-R1 and regulate sleep states in developing

larvae

Since the *PK2-R1* driver broadly labeled neurons (~1000 cells) throughout the CNS

(Figure 1C), we next attempted to identify a subpopulation of these neurons relevant for

sleep regulation. To narrow down the candidates within PK2-R1-positive populations,

we examined subsets labeling GABAergic, cholinergic, or glutamatergic neurons, as

well as those co-expressing other neuropeptides identified in our screen, but none of

these manipulations reproduced the sleep phenotype caused by PK2-R1 perturbation.

Notably, insulin-producing cells (IPCs) in the pars intercerebralis (PI), the

neurosecretory center of *Drosophila*, have been reported to express *PK2-R1* (Schlegel et

al., 2016). Consistently, the *PK2-R1²ᴬ⁻ᴸᵉˣᴬ* knock-in driver labeled all IPCs (Figure 2A

and B). To assess whether IPCs are involved in larval sleep, we silenced IPCs with

Kir2.1 expression, using the IPC-specific drivers *Dilp3-* and *Dilp5-GAL4* (Figure 2C).

We found that silencing of IPCs significantly increased the sleep amounts compared to

the control, phenocopying *PK2-R1* neuron silencing and *PK2-R1* knockout (Figure 1A).

Consistently, both *Dilp3* and *Dilp5* null mutants exhibited larger sleep amounts

compared to the control (Figure 2D). These phenotypes of *Dilp* mutants and IPC

silencing are unlikely to be accounted for by locomotion defects, as the travel distances

during wake periods were unaffected (Fig. 2—figure supplement 1). Through these

trials, insulin-producing cells (IPCs) emerged as one of the populations whose silencing

phenocopied PK2-R1 loss, supporting their key role in larval sleep regulation while not

excluding contributions from other PK2-R1-expressing neurons. Collectively, these

findings suggest that *PK2-R1* in IPCs regulates larval sleep.

Hugin-expressing neurons control larval sleep states

Given that PK2-R1 is a receptor of the neuropeptide Hugin (Rosenkilde et al., 2003), we

next examined whether Hugin is involved in the larval sleep. We found that *Hug* mutant

larvae exhibited significantly larger sleep amounts compared to the control (Figure 3A),

consistent with *PK2-R1* knockouts (Figure 1A). When we silenced Hug-expressing

neurons by driving the expression of Kir2.1 with the *Hug* CRISPR-knock-in driver

Hug²ᴬ⁻ᴳᴬᴸ⁴ (Deng et al., 2019), it resulted in a significant increase of sleep amounts

without detectable changes in locomotion speed (Figure 3B and Figure 3—figure

supplement 1B). Conversely, activation of *Hug*-expressing neurons with the red-shifted

channel rhodopsin ReaChR (Inagaki et al., 2014) or heat-sensitive ion channel TrpA1

(Hamada et al., 2008) both caused a significant reduction of sleep amount (Figure 3C

and D) and increased locomotor activity. However, locomotion changes were not

consistently observed upon either activation or suppression of Hug neurons (Figure 3—

figure supplement 1), suggesting that changes in sleep cannot be simply explained by

locomotor alterations. We further quantified larval feeding using a dye-based ingestion

assay and found that silencing HugPC neurons reduced food intake (Figure 3—figure

supplement 3), indicating that the sleep phenotype is unlikely to be explained by

feeding behavior being misclassified as sleep. These data together indicate that *Hug*-

expressing neurons are responsible for downregulating larval sleep.

Hug-expressing neurons are comprised of 20 cells in the subesophageal zone (SEZ) of

the brain, and are classified into four subpopulations based on their projection patterns

(Bader et al., 2007). Of these, one subpopulation called HugPC neurons (Bader et al.,

2007) have been reported to form synaptic connections with IPCs (Hückesfeld, Peters,

& Pankratz, 2016). We thus suspected that HugPC neurons may be responsible for sleep

regulation. In line with this notion, axons of HugPC neurons, labeled by a newly

generated *HugPC-LexA* driver (Figure 3—figure supplement 2), and IPCs, labeled by

Dilp3-GAL4 (Figure 3E), exist in space close to each other. Indeed, silencing of HugPC

neurons by Kir2.1 significantly increased larval sleep (Figure 3F), phenocopying those

observed in *Hugin* and *PK2-R1* mutants. These results hint at the possibility that HugPC

neurons activate IPCs via Hugin/PK2-R1 signaling to regulate sleep.

Hugin triggers Ca^{2+} elevation and Dilp secretion in larval IPCs via PK2-R1

A previous report showed that IPCs exhibit increased firing rates and intracellular

calcium (Ca^{2+}) levels upon stimulation (Kréneisz et al., 2010). This motivated us to next

test if Hugin activates IPCs. To further visualize the output pattern of HuginPC neurons,

we expressed the presynaptic marker Syt-eGFP in HuginPC neurons (Figure 4—figure

supplement 1).To test whether Hugin activates IPCs, we utilized the recently developed

calcium indicator CRTC::GFP, which can report Ca^{2+} dynamics in the timescale of

several minutes (Bonheur et al., 2023). As glucose reportedly triggers Ca^{2+} dynamics in

IPCs in a range of minutes (Oh et al., 2019), we wondered if CRTC::GFP can detect this

dynamics, as a positive control. To this end, we applied D-glucose to isolated larval

brains and found a significant cytosol-to-nuclear translocation of CRTC::GFP in IPCs

within ~5 min, confiming that this reporter can indeed detect Ca^{2+} responses in IPCs

upon stimulation (Figure 5B). We then examined whether *Hug* neurons act upstream of

PK2-R1-expressing IPCs by measuring Ca^{2+} responses of IPCs while thermogenetically

activating *Hug* neurons with TrpA1. We found that activation of *Hug* neurons tended to

increase Ca^{2+} levels in IPCs (Figure 4B and C), consistent with the HuginPC-to-IPCs

axis (Figure 4D). Next, we tested whether Hug peptides can induce Ca^{2+} responses in

IPCs using *ex vivo* preparations. The *Hug* gene encodes a precursor that potentially

produces at least two distinct peptides, Hug-γ and PK-2, both of which can bind PK2-

R1 to induce Ca^{2+} responses (Rosenkilde et al., 2003). We thus incubated the

CRTC::GFP-expressing larval brain with chemically synthesized Hug-γ or PK-2

(Kréneisz et al., 2010; Oh et al., 2019) (Figure 5A). We found that bath application of

either Hug-γ or PK-2 induced significant Ca^{2+} responses in IPCs (Figure 5B–D). In

agreement with the idea that Hug peptides activate IPCs via *PK2-R1*, knock-out of *PK2-

R1* blocked the Hug peptide-evoked Ca^{2+} elevations in IPCs (Figure 5F and G). Similar

results were obtained when we bath-applied the synthetic Neuromedin U (NMU), the

mammlaian homlog of Hugin (Figure 5A and 5G), suggesting the role of Hugin/PK2-

R1 signaling may be conserved across species. Furthermore, Dilp3 accumulated in IPCs

were significantly reduced after applying Hug peptides, suggesting that Hugin causes

IPCs to release Dilp3 (Figure 5E). Collectively, these data suggest that *Hug* neurons

activate IPCs via the Hug/PK2-R1 signaling to regulate sleep (Figure 5H).

Hugin/PK2-R1 axis has distinct impacts on wake/sleep control in larvae and adults

Given the differences between larval vs adult sleep in temporal pattern and circadian

influence, we next wondered if Hug/PK2-R1/Dilps axis functions in adult sleep as well.

As in larvae, *PK2-R1* knockout adults showed increased sleep amount compared to the

control (Figure 6—figure supplement 1B). In contrast, *Hug* knockout or silencing of

HugPC neurons failed to influence adult sleep (Figure 6A and 6B). This suggests that,

unlike larvae, Hug is dispensable for adult sleep. Interestingly, we observed that the

expression patterns of PK2-R1 and Hug, and the morphology of HugPC neurons and

IPCs, showed a broadly similar distribution in larvae and adults, although we did not

directly track individual neurons across development (Figure 6—figure supplement 2).

This implies that the differential roles of Hug in larvae vs adults are likely due to

physiological differences in HugPC neurons and/or IPCs. We thus examined how Hug

peptides evoke Ca^{2+} responses in adult IPCs and found that neither Hug-γ nor PK-2

could induce Ca^{2+} responses in adult IPCs (Figure 6D and 6E), unlike in larvae.

Although PK-2 treatment even led to a modest increase in Dilp3 immunoreactivity in

adult IPCs (Figure 6F), the physiological significance of this effect remains unclear; at

present, we consider it most likely that PK-2 acts on larval and adult IPCs in a different

manner. These data indicate that Hug induces Ca^{2+} responses and Dilps secretion in

larval IPCs but not in adult IPCs. Surprisingly, we further found that *Dilp3* or *Dilp5* null

mutations reduced sleep amounts in adults, consistent with previous reports (Cong et al.,

Sleep 2015) (Figure 6C), while in larvae, these mutations in fact increased the sleep

amounts. These data unexpectedly reveal how the same set of neuronal circuits and

molecular signaling therein can evolve over development in a complex manner.

Discussion

In this study, we have identified the neuropeptide Hugin and its receptor PK2-R1 as a

ligand/receptor pair critical for sleep regulation in *Drosophila* larvae. We have further

shown that larval IPCs express PK2-R1, respond and release Dilp3 upon Hugin

stimulation. Surprisingly, the Hugin/PK2-R1 axis is dispensable for sleep regulation in

adults, even though the gene expressions as well as the circuit structure appeared to be

conserved between larval and adult brains. Furthermore, Dilps appear to modulate sleep

in opposite directions in larvae and adults. Our findings thus uncover the

neuropeptidergic modulation circuitry that specifically regulates developmental sleep,

and suggest divergent usage of the same molecules and circuitry for sleep modulation in

larvae and adults.

Hugin/PK2-R1/Dilps axis negatively regulates sleep in larvae

The present study shows that the neuropeptide Hugin and its receptor PK2-R1 are

required for sleep regulation in the early larval stage. This notion is supported by

following lines of evidence. First, knock-out mutations in *Hugin* or *PK2-R1* increased

sleep amounts compared to the control (Figure 1A and 3A). Consistently, silencing a

subpopulation of *Hugin*-expressing neurons, called HugPC neurons, or *PK2-R1* neurons

increased sleep amounts (Figure 1B, 3B and 3F). Second, thermogenetic activation of

Hugin neurons decreased sleep amounts (Figure 3D). Third, HugPC neurons project their axons to IPCs in the brain that express PK2-R1 receptors (Figure 3E). Fourth, Hugin peptides induced Ca^{2+} elevation and promoted Dilp3 release from IPCs in a PK2-R1-dependent manner (Figure 5). Last, *Dilp3* null mutations increased sleep amounts (Figure 2D). Based on these data, we propose a model in which the Hugin/PK2-R1/Dilps axis along the HugPC-IPC circuitry modulates larval sleep (Figure 7).

Examples of Hugin acting through PK2-R1 are prominent in the context of feeding regulation in larvae and adults (Schoofs et al., 2014). For example, a recent study has suggested that Hugin regulates the timing of pupariation and body wall contraction during pupariation through PK2-R1 in PTTH neurons (Ohhara et al., 2024). In the present study, we showed that HugPC in the larval brain modulates sleep (Figure 3F). A previous electron microscopic analysis suggested that larval HugPC neurons likely form synaptic connections to downstream neurons including IPCs (Hückesfeld et al., 2021; Schlegel et al., 2016). We here showed that PK2-R1 is required for larval sleep control and that thermogenetic activation of Hug neurons is associated with Ca^{2+} responses in PK2-R1-expressing IPCs *in vivo* (Figure 4). Importantly, our *ex vivo* data indicated that Hugin peptide-induced Ca^{2+} response in IPCs was largely abolished by *PK2-R1* mutation (Figure 5), indicating the critical role of the Hugin/PK2-R1 axis for the HugPC-mediated IPC activation. Together, we propose that the HugPC-IPCs circuity mobilizes Hugin/PK2-R1 signaling to selectively modulate larval sleep. Interestingly, our *ex vivo* data indicated that Neuromedin U (NMU), a mammalian ortholog of Hugin, can trigger Ca^{2+} responses in larval IPCs in a PK2-R1-dependent fashion (Figure 5). Given that NMU has been implicated in sleep regulation in fishes and mammals (Ahnaou & Drinkenburg, 2011; Chiu et al., 2016), an analogous ligand/receptor pair might be involved in sleep modulation in mammals as well.

In this study, activation of Hugin–PK2-R1 signaling by bath application of Hugin peptides led to a significant reduction in Dilp3 immunoreactivity in IPCs, whereas Dilp2 immunoreactivity did not show a decrease under these conditions, suggesting that Hugin–PK2-R1 activation promotes Dilp3 secretion without strongly enhancing Dilp2 release in our assay. One plausible explanation is that individual Dilps are stored in

partially distinct vesicular pools within IPCs and are released through stimulus-specific

mechanisms. Consistent with this idea, previous work has shown that Dilp release from

IPCs can be selective, with Dilp2 and Dilp3 responding to distinct nutritional cues (Kim

and Neufeld, 2015). Confocal analyses in that study further demonstrated that Dilp2 and

Dilp3 segregate into different intracellular granules, supporting the notion that

individual Dilps can be targeted to distinct secretory pathways within the same neurons.

In light of these observations, the selective effect of Hugin–PK2-R1 activation on

Dilp3, but not Dilp2, is readily interpreted as another example of non-uniform Dilp

regulation. Rather than uniformly driving secretion of all IPC-derived Dilps, we propose

that Hugin–PK2-R1 signaling preferentially mobilizes the Dilp3 secretory pathway.

Such selective regulation is reminiscent of piecemeal degranulation in mammalian

eosinophils and mast cells, as well as stimulus-dependent mobilization of specific

vesicle pools in pancreatic β cells. In this model, different Dilps are stored in partially

distinct vesicular pools and are mobilized by different upstream inputs. While

ultrastructural approaches such as electron microscopy could provide further insight

into the anatomical segregation of Dilp-containing vesicles, these analyses are

technically beyond the scope of the present study, and we therefore highlight them as an

important direction for future work.

PK2-R1 knockout larvae exhibited both an increase in sleep and a reduction in

locomotion speed. Because our locomotor metric is defined as travel distance per unit

time during epochs classified as wake, the observed decrease in locomotion speed

cannot be fully explained by the increased fraction of time spent in sleep. Nevertheless,

we cannot exclude the possibility that part of the apparent sleep increase in the knockout

reflects broader motor impairments in PK2-R1–expressing circuits. PK2-R1 expression

is not confined to IPCs but is broadly distributed in the nervous system, including

populations in the ventral nerve cord (Figure 1C), making it likely that PK2-R1–positive

neurons outside IPCs contribute to locomotor control. Consistent with this idea, when

we restricted our manipulations to IPCs, we observed robust changes in sleep without

significant alterations in locomotor activity (Figure 2—figure supplement 1). These

findings suggest that IPCs are more directly involved in sleep regulation, whereas other

PK2-R1–expressing neurons may predominantly influence locomotion.

Dh44 neurons promote arousal in second-instar larvae (Poe et al., 2023), raising the

possibility that hugin signaling may influence sleep through Dh44 in addition to IPCs.

Consistent with this idea, the hugin receptor PK2-R1 is expressed not only in IPCs but

also in Dh44 and DMS neurons (Schlegel et al., 2016). Thus, an extension of our model

is that HuginPC-derived hugin may act on multiple PK2-R1–positive targets (including

IPCs and potentially Dh44/DMS neurons) to shape larval sleep. Future cell type-

specific perturbations of PK2-R1 in Dh44 and DMS neurons will be important to test

the contribution of these pathways.

Divergent roles of insulin signaling in regulation of larval sleep and feeding

IPCs in larvae respond to exogenously applied Hug peptides, both by exhibiting

elevated Ca^{2+} levels and decreases in intracellular Dilp3 storage (Figure 5C–E). In

contrast to Dilp3, however, the amount of Dilp2 accumulated in IPCs remains

unchanged following Hug peptide application (Figure 5—figure supplement 1B). On

the other hand, Dilp2 accumulation in IPCs decreases after glucose treatment (Figure

5—figure supplement 1A) (Oh et al., 2019). These results unexpectedly suggest that

larval IPCs form parallel channels inside, one for Hug-Dilp3 and the other for glucose-

Dilp2. Mechanistically, Hug acts directly on IPCs via the PK2-R1 receptor. As for

glucose, on the other hand, glucose-sensing CN neurons in the brain take up glucose

and subsequently secrete another neuropeptide to activate IPCs (Oh et al., 2019). Such

parallelism inside IPCs might provide insight into why insect species broadly express

multiple insulin-like peptide genes, in some cases even as many as 40 genes in the

silkworm Bombyx mori (Okamoto, 2021). Drosophila genome harbors eight Dilp genes

(Nässel et al., 2015), of which Dilp2, Dilp3, and Dilp5 are considered nutrient-related.

Nevertheless, mRNA expression or release patterns of these Dilps are not identical

under different nutrient conditions (Géminard et al., 2009; Kauffman & DiAngelo,

2024; Nässel et al., 2015; Oh et al., 2019; Tomoatsu et al., 2002). Furthermore, these

Dilps exhibit different binding affinities for the insulin receptor (InR), the secreted

decoy of InR (SDR), the ecdysone-inducible gene L2 (Imp-L2) (Okamoto et al., 2013).

These reports hint at the differential roles of different insulin-like peptides, but

mechanistic insights remain largely lacking. Our data of larval IPCs forming parallel

channels for Hug-Dilp3 and glucose-Dilp2 may provide a point of entry for tackling this
mystery.
339

340 Differential impacts of the same neuropeptidergic modulations on larval and adult
341 sleep

Our data show that the Hugin/PK2-R1/Dilps axis likely downregulates sleep amounts in
larvae. Despite the significant increase of sleep amount in *Hugin* mutant larvae, *Hugin*
mutant adults showed sleep amounts comparable to wild-type control (Figure 6A),
consistent with a previous report that Hugin is dispensable for control of baseline sleep
levels (Schwarz et al., 2021). According to our histological analyses, the Hugin/PK2-R1
expression patterns as well as the HugPC-IPCs circuitry appear to be largely conserved
in adults (Figure 3E and Figure 6—figure supplement 2). Unlike larval brain, however,
Hugin peptides failed to trigger Ca^{2+} responses in IPCs in adults (Figure 6E). It is thus
possible that Hugin/PK2-R1 signaling along the HugPC-IPCs circuitry is suppressed in
adults. IPCs in adults receive multiple positive and negative modulatory inputs through
GPCRs including the metabotropic GABA_B receptors (Enell et al., 2010), which
suppresses IPC activity and Dilp release in adult IPCs (Enell et al., 2010). It is thus
plausible that such negative modulatory inputs to IPCs in adults might counteract with
the Hugin/PK2-R1 axis to suppress Dilp release. In addition, our data suggest that Dilps
modulate sleep amount in the opposite directions in larvae and adults (Figure 7).
Comparing the expression levels and activities of GPCRs in larval and adult IPCs would
be essential to better understand how same modulatory signals over the course of
development come to exert differential impacts on sleep. Interestingly, Hugin in adults
appears irrelevant for the baseline sleep amount but is required for homeostatic
regulation of sleep (Schwarz et al., 2021). Thus, testing if Hugin/PK2-R1 axis is
involved in the homeostatic regulation of larval sleep, and how such a system compares
to its adult counterpart, may further provide mechanistic insights into how homeostatic
sleep regulation maturates over development.
365

366 In summary, we identified the Hugin/PK2-R1/Dilps signaling along the HugPC/IPCs
367 circuit as the molecular and circuitry basis for developmental sleep regulation. Our

368 study begins to fill the knowledge gap in larval sleep regulation, and sheds light on
369 mechanistic differences in sleep regulation between early developmental stages and
370 adults.
371

Materials and Methods

Fly strains

The following strains of *Drosophila melanogaster* were obtained from the Bloomington
Drosophila Stock Center: *iso31* (an isogenic w^{1118} strain; BDSC# 5905), *PK2-R1^{attP}*
(BDSC# 84563), *PK2-R1$^{2A-GAL4}$* (BDSC# 84686), *PK2-R1$^{2A-LexA}$* (BDSC# 84431),
Dilp3^1 (BDSC# 30882), *Dilp5^1* (BDSC# 30884), *Dilp3-GAL4* (BDSC# 52660), *Dilp5-*
GAL4 (BDSC# 66007), *Hug^{attP}* (BDSC# 84514), *Hug$^{2A-GAL4}$* (BDSC# 84646), *Hug$^{2A-LexA}$*
(BDSC# 84397), *UAS-Kir2.1::EGFP* (BDSC# 6596), *UAS-Stinger, LexAop-*
tdTomato.nls (BDSC# 66680), *UAS-mCD8::mCherry, UAS-CRTC::GFP* (BDSC#
99657), *UAS-ReaChR* (BDSC# 53741), *UAS-TrpA1* (BDSC# 26263), *LexAop-ReaChR*
(BDSC# 53746), *LexAop-rCD2::RFP-p10.UAS-mCD8::GFP-p10* (BDSC# 67093).
w^{1118} was from Y. Rao (Peking University); *HugPC-GAL4* (Hückesfeld et al., 2016)
was from M. Pankratz (University of Bonn); *LexAop-TrpA1* (Burke et al., 2012) was
from S. Waddel (University of Oxford). *HugPC-LexA* was created in this study (see the
following section, "Generation of transgenic lines"). Detailed lists of fly genotypes can
be found in the key resource table and Source Data files.
388

389 **General fly maintenance**

390 Flies were maintained in an incubator (PHCbi, model# MTR-554-PJ) set to 25°C with a
391 humidity of over 60% and a 9AM:9PM light/dark cycle. After 10 d of rearing in bottles,
392 old flies were replaced with freshly eclosed ones to maintain the strain (Furusawa et al.,
393 2023).
394

Egg collection

Agar substrates for egg collection were prepared by microwave-heating a mixture of 2.0
g sucrose (Fujifilm Wako Pure Chemicals, cat# 196-00015), 3.0 g agar (BD, cat#
214010), 2.5 mL apple juice (Dole), and Milli-Q water up to 100 mL. Melted agar was

poured into Petri dishes with a diameter of 4 cm (approximately 3 mL per dish).

Solidified agar was scratched with tweezers to make the surface rough. Agar plates were

stored inside a sealed container at 4°C. Upon egg collection, yeast paste (dried yeast

(Nippon Beet Sugar Co., Ltd.) kneaded with Milli-Q water at 40% (w/w)) was placed in

the center of the agar plates to further promote oviposition.

Larval staging

First-instar larvae were collected 2 d after egg collection. Among them, second-instar

larvae that molted within a 2-h window were used for video recording. Distinguishing

between first- and second-instar larvae was based on morphological characteristics:

while anterior spiracles were not evident in the first-instar, fist-shaped spiracles were

observed in the second-instar (Lakhotia and Ranganath, 2021).

Larval sleep analysis

Twenty-four-well silicone chambers for sleep analysis, described in Figure 1—figure

supplement 1, were made by pouring dimethylpolysiloxane (PDMS) into molds based

on a previous study (Churgin et al., 2019; Szuperak et al., 2018). As a substrate, 120 μL

of a mixture of 3% agar and 2% sucrose was poured into each well. After the gel

solidified, 10 μL per well of 60 mg/mL yeast suspension was applied to the surface of

the gel. Larvae were placed individually into the wells, and a glass plate (Azone, cat# 1-

4540-01, 5 mm × 200 mm × 200 mm) was positioned on top of the chamber to prevent

larvae from escaping. The chamber was placed inside a dark box equipped with a ring-

type infrared LED (CCS, cat# LDR2-132IR940-LA) and a camera (The Imaging

Source, cat# DMK27BUP031), and recording was started immediately without

acclimation. Images were captured through the IC Capture software (The Imaging

Source, version 2.5) under the following conditions: MJPEG Compressor for codec, a

frame rate of 0.87 fps, and an image size of Y800 (2592 × 1944).   Locomotor activity

was quantified from the same video recordings used for larval sleep analysis. Travel

distance was calculated as the cumulative displacement of the larval centroid over time,

and locomotion speed was defined as total travel distance divided by the total duration

of wakefulness, considering only time bins classified as wake. To minimize the

possibility that feeding behavior was misclassified as sleep, we adjusted our immobility

threshold using video recordings in which larval feeding bouts, including relatively

subtle episodes, were apparent, and configured the algorithm so that feeding-associated

movements would generally be classified as wake rather than inactivity. During the

development and routine use of the assay, we also inspected representative recordings

across genotypes and did not observe obvious feeding-related abnormalities or gross

locomotor defects under our HuginPC/IPC manipulations.

Sleep deprivation

Blue light irradiation was performed using a 455-nm LED (Thorlabs, cat# M455L4)

with a collimating adapter (Thorlabs, cat# SM1U25-A) (Omamiuda-Ishikawa et al.,

2020; Yoshino et al., 2017; Yoshino et al., 2025). The irradiation intensity was set to the

maximum scale using a T-cube LED driver (Thorlabs, cat# LEDD1B). Temporal

control of LED illumination was achieved using the Arduino UNO microcontroller

board (Arduino, RRID:SCR_017284). For sleep deprivation experiments, the first 1 h

was recorded without LED illumination, followed by another 1 h consisting of repeated

cycles of "LED ON for 90 s and OFF for 30 s".

Immunohistochemistry

Antibody staining of larval neurons was carried out as previously reported (Morikawa et

al., 2011; Omamiuda-Ishikawa et al., 2020). Briefly, dissection in phosphate-buffered

saline (PBS); fixation in 4% paraformaldehyde/PBS at room temperature for either 90

min in CRTC::GFP experiments or 30 min in the others; washing with 0.3% Triton X-

100/PBS (hereafter referred to as PBST) for 10 min × 3; blocking in 5% normal goat

serum (NGS)/PBST at room temperature for 30 min; primary antibody treatment at 4°C

overnight; washing with PBST for 10 min × 3; secondary antibody treatment at 4°C

overnight; washing with PBST for 10 min × 3; and mounting in VECTASHIELD

medium (Vector Laboratories, cat# H-1000, RRID:AB_2336789). Reactions were

performed either in Terasaki 96-well plates (stem Corp., cat# P96R37N) for adults or in

48-well culture plates (Corning, cat# 3548) for larvae. The following primary antibodies

were diluted in 5% NGS/PBST: chicken polyclonal anti-GFP (1:1000, Aves Labs, cat#

GFP-1020, RRID:AB_10000240), mouse monoclonal anti-mCherry (1:1000, Takara

Bio, cat# 632543, RRID:AB_2307319), rabbit polyclonal anti-Dilp3 (1:250, (Veenstra

et al., 2008)), rabbit polyclonal anti-Dilp2 (1:2000, (Okamoto et al., 2012)), mouse

monoclonal anti-BRP (1:100, Developmental Studies Hybridoma Bank, clone nc82;

RRID:AB_2314866), and rat monoclonal anti-histone H3 phospho S28 (1:1000,

Abcam, cat# ab10543; RRID:AB_2295065). The following secondary antibodies were

diluted in 5% NGS/PBST: goat anti-chicken Alexa Fluor 488 (1:200, Thermo Fisher

Scientific, cat# A-11039, RRID:AB_2534096), goat anti-mouse Alexa Fluor 555

(1:200, Thermo Fisher Scientific, cat# A-21424, RRID:AB_141780), goat anti-rabbit

Alexa Fluor 633 (1:200, Thermo Fisher Scientific, cat# A-21071, RRID:AB_2535732),

and goat anti-rat Alexa Fluor 633 (1:200, Thermo Fisher Scientific, cat# A-21094;

RRID:AB_2535749). Fluorescence images were obtained using a confocal microscope

(Leica, TCS SP8) with 20× (Leica, HC PL APO CS2 20×/0.75 IMM) and 63× (Leica,

HC PL APO CS2 63×/1.40 OIL) objectives. Images were captured at 512×512-pixel

resolution with an interval of 1 μm. Z-stack images of the maximum projections were

generated using the ImageJ v1.53t software (Schneider et al., 2012);

RRID:SCR_003070).

Generation of transgenic lines

HugPC-LexA lines were generated by ΦC31 integrase-mediated transgenesis, as

previously described (Groth et al., n.d.).

The 544-bp enhancer region of *Hug*, essentially the same as that used to generate

HugPC-GAL4 (Hückesfeld et al., 2016), was PCR-amplified from fly genomic DNA.

The gel-purified amplicon was subjected to another round of PCR to add hangout

sequences. The backbone suicide-vector pBPnlsLexA::p65Uw (RRID:Addgene_26230)

was initially amplified by transforming the *ccdB*-resistant *E. coli* strain, One Shot™

ccdB Survival™ 2 T1^R competent cells (Thermo Fisher Scientific, cat# A10460). The

PCR product and the vector were double-digested by Aat II (NEB, cat# R0117S) and

Fse I (NEB, cat# R0588S), followed by ligation using In-Fusion HD Cloning Kit

w/Cloning Enhancer (Takara Bio, cat# 639635). The integrity of the resulting plasmid

was confirmed by DNA sequencing. Plasmid insertion was targeted to four attP sites

(attP2, attP40, VK00005, and su(Hw)attP5) individually through ΦC31 integrase-

mediated transgenesis by BestGene Inc. (Chino Hills, CA). Transformants were selected

based on the eye color marker. Among the generated strains harboring *HugPC-LexA* in

each insertion site, labeling patterns in the larval brain were checked by driving

fluorescent marker expression under the control of *LexAop* (Figure 3—figure

supplement 2). As the *HugPC-LexA* inserted in attP2 closely resembled the labeling

pattern of the original *HugPC-GAL4*, it was used in further experiments.

The following primers were used for cloning:

Forward primer for the *HugPC* enhancer; 1st round PCR against genomic DNA:

5'-AAGGGTTTGGTTTAATTTATTTATGTCATA-3'

Reverse primer for the *HugPC* enhancer; 1st round PCR against the genomic DNA:

5'-GGACAACTGATGCCAGCAGC-3'

Forward primer for the *HugPC* enhancer with hangout sequences (underlined); 2nd

round PCR against the 1st round amplicon:

5'-<u>gaaaagtgccacctgacgt</u>AAGGGTTTGGTTTAATTTATTTATGTCATA-3'

Reverse primer for the *HugPC* enhancer with hangout sequences (underlined); 2nd round

PCR against the 1st round amplicon:

5'-<u>cccgggcgagctcggccgg</u>GGACAACTGATGCCAGCAGC-3'

Food intake assay

Food intake was measured using a dye-based feeding assay adapted from Hückesfeld et

al. (2016). Apple juice agar plates were prepared with a spot of red yeast paste in the

center. After 30 min starvation, larvae were transferred onto the yeast paste and

videotaped for 20 min. Larvae were then collected in a small cell strainer and rinsed

with 60°C hot water to remove external dye, transferred to glass slides, and

photographed. Images were analyzed in ImageJ by measuring the area of red-stained

dye signal and the total body area, and food intake was quantified as the ratio of dye-

stained area to total body area for each larva.

Measurement of neuronal activity and Dilp3 accumulation in IPCs

The GFP-tagged CRTC (CRTC::GFP) reporter was applied to quantify the activities of

larval IPCs, essentially according to the previous report in adult flies (Bonheur et al.,

2023).

Parental *Dilp3-GAL4* and the reporter line harboring *UAS-CRTC::GFP* and *UAS-*

mCD8::mCherry (BDSC# 99657) were crossed, and eggs were collected within an 8–10

h time window. Offspring larvae were reared for 4 d after egg laying (AEL) at 25°C to

early third instar. Collected larvae were washed with water and starved on water-soaked

paper for 12–14 h. For *in vivo* thermogenetic activation, larva-containing Petri dishes

were warmed at 30°C for 60 min. For *ex vivo* thermogenetic experiments, larvae were

dissected in calcium- and sugar-free HL3.1 buffer (Nakamizo-Dojo et al., 2023)) at

30°C, and each brain was incubated in the same buffer at 30°C for 5 min. For *ex vivo*

glucose or peptide application assays, larvae were first dissected in the same buffer as

above at room temperature, and brain samples were then treated with either 20 mM D-

glucose or 1 mM of chemically synthesized peptides dissolved in HL3.1 buffer for 5

min. After incubation, samples were subjected to immunohistochemical analyses to

visualize GFP, mCherry, and Dilp3 signals (see the section "Immunohistochemistry").

Image processing was performed using ImageJ v1.53t ((Schneider et al., 2012);

RRID:SCR_003070). Nuclear ROIs and cell body outlines were drawn based on the

mCD8::mCherry fluorescence insensitive to neuronal activity, and the mean intensities

of CRTC::GFP signals within the nucleus (F_{nuc}) and the cytoplasm ($F_{cyto} = F_{whole\ cell}$ -

F_{nuc}) were quantified. NLI was calculated using the following formula: (F_{nuc} -

F_{cyto})/($F_{nuc} + F_{cyto}$). To assess Dilp3 accumulation in IPCs, the anti-Dilp3 signal intensity

was measured within the cytosol of each cell, excluding the nucleus. Data from 4-14

Dilp3-positive IPCs per brain, collected from 3-7 larvae, were pooled for each

genotype/treatment group.

We attempted to record GCaMP-based calcium dynamics in Hugin neurons and IPCs,

but did not pursue this approach further because it was low-throughput and a positive-

control assay (glucose application) did not elicit the expected IPC calcium response in

our hands.

Chemical synthesis of Hug peptides

Peptides with the following amino acid sequences were chemically synthesized using

the Fmoc solid phase method, and their purity was assessed by HPLC to be over 95%.

All procedures were carried out by Eurofins Genomics K.K. (Tokyo, Japan).

Drosophila Hug-γ:

Acetyl-QLQSNGEPAYRVRTPRL-CONH2 (17 a.a.)

Drosophila PK-2:

Acetyl-SVPFKPRL-CONH2 (8 a.a.)

Human NMU:

Acetyl-FRVDEEFQSPFASQSRGYFLFRPRN-CONH2 (25 a.a.)

Sleep quantification in adult flies

The following parts of the *Drosophila* activity monitor (DAM) system were purchased

from TriKinetics Inc. (Waltham, MA): the *Drosophila* Activity Monitor (cat# DAM2),

the power supply interface unit (cat# PSIU9), monitor tubes with an outside diameter of

5 mm and a length of 65 mm (cat# PGT5x65), and tube caps (cat# CAP5-Black). Newly

eclosed adult flies were collected under brief CO_2 anesthesia and kept as groups of 10–

15 in standard food vials. Males and females were kept separately to avoid mating. To

minimize disturbance by noise and vibration, flies were placed inside a fan-less Peltier-

type incubator (Mitsubishi Electric Engineering Co., Ltd., cat# SLC-25) set to 25°C,

externally connected with 5-mm diameter LED lights (MY-CRAFT, Ltd., cat#

5W0601) and a timer device (Panasonic Corp., cat# WH3311WP) to generate a

9AM:9PM light/dark cycle. After 3 d of eclosion, each fly was gently aspirated into a

Pyrex glass monitor tube, one end of which was stuffed with agar food (2% (w/w) agar,

5% (w/w) sucrose), and the other was plugged with a cotton string. The tubes were set

to the monitor device and placed inside the same fan-less incubator. The DAM system

was connected to an external PC installed with DAMSystem3 Data Collection Software

(RRID:SCR_021809; available at https://trikinetics.com/) for data acquisition, and

beam-crossing by each fly was counted for two successive days. Time windows longer

than 5 min without beam-crossing were judged as sleep bouts, following the

conventional criteria for adult fly sleep (Paul J. Shaw, Chiara Cirelli, Ralph J.

Greenspan, & Giulio Tononi, 2000).

Statistical analysis

Statistical analyses by Kruskal–Wallis one-way ANOVA, Mann–Whitney U-test, or

chi-square test were performed using Prism 9.5.1 (GraphPad, RRID:SCR_002798).

Bonferroni correction was applied for multiple comparisons. Asterisks (*) represent p-

values as indicated within each figure legend. All data necessary to reproduce the figure

panels and statistical analyses are available as Source Data files.

Acknowledgements

We thank Y. Rao, M. Pankratz, S. Waddel, and the Bloomington *Drosophila* Stock

Center for fly stocks; J. A. Veenstra for the anti-Dilp3 antibody; T. Nishimura for the

anti-Dilp2 antibody; Y. Nishizuka for valuable suggestions on writing code for

automated larval detection and sleep quantification; H. Ito, N. Yamaji, M. Maetani, A.

Ogasawara, E. Kato, T. Rikiishi, S. Ando, M. Hayashi, and S. Miyazaki for technical

assistance and fly maintenance; and the members of Emoto Lab for critical comments

and discussion. This work is supported by MEXT Grants-in-Aid for Scientific Research

on Innovative Areas "Dynamic regulation of brain function by Scrap and Build system"

(KAKENHI 16H06456), JSPS (KAKENHI 16H02504), WPI-IRCN, AMED-CREST

(JP22gm310010), JSTCREST, Toray Foundation, Naito Foundation, Takeda Science

Foundation, and Uehara Memorial Foundation to K.E.; the Leading Initiative for

Excellent Young Researchers (LEADER) from MEXT, JSPS (KAKENHI 22K06309),

and AMED-PRIME (JP22gm6510011) to K.I.; Grant-in-Aid for JSPS Fellows

(24KJ0911) to M.M.; and Grant-in-Aid for JSPS Fellows (22J21096; 22KJ1042) to

C.H.

Figures and Legends

Figure 1. PK2-R1 is required for larval sleep control

(A) Sleep amounts in *PK2-R1* or *Oamb* knockout mutants. Each dot represents an

individual animal; in this and the following panels, "N" indicates the number of

biologically independent animals per group, and the thick line and thin error bars

indicate the median and interquartile range (Q1–Q3), respectively. *** $p < 0.001$

(Mann–Whitney U-test with Bonferroni correction). (B) Sleep amounts in larvae in

which *PK2-R1* neurons were silenced. *** $p < 0.001$ (Mann–Whitney U-test with

Bonferroni correction). (C) Expression pattern of *PK2-R1-GAL4 > UAS-Kir::EGFP*

larvae.

Figure 1—figure supplement 1. Detailed experimental setup for analyzing larval

sleep

(A) Schematic flow of automated larval sleep quantification. A second instar larva was

placed in each of 24 wells, and all 24 wells were videotaped from above. For each well

and frame, the image of larva was blurred and binarized to automatically locate the

larva. Consecutive frames were then compared so that changes in pixel values were

summed to estimate larval motion. This motion estimate was then used to classify every

consecutive pair of frames as representing "active" or "inactive." We found that

inactivity across 12 consecutive frames meets the sleep criteria, as shown in Figure 1—

figure supplement 2. (B) Schematic representation of the 24-well chamber used for

larval sleep assays. (C) Flow of automated processes of larval motion detection and

active/inactive state determination. For each well and frame, the image of larva was

blurred and binarized. Consecutive frames were then compared so that changes in pixel

values were summed to estimate larval motion. This motion estimate was then used to

classify every consecutive pair of frames as representing "active" or "inactive."

Figure 1—figure supplement 2. Larval "sleep state" in this study is consistent with

behavioral criteria for sleep

(A) Relationship between "inactive" state duration and spontaneous arousal probability.

Sustained quiescence for ≥ 12 frames (open circles) significantly reduced arousal

probability compared to 1-frame inactivity. * $p < 0.05$, NS: $p \geq 0.05$ (chi-squared test).

(B) The percentage of larvae that showed a response within the next 10 frames after

high-intensity blue light. The number of events analyzed was 36 and 60 with and

without blue light irradiation, respectively. * $p < 0.05$ (Fisher's exact test). (C) Sleep

amounts before, during, and after blue light irradiation. The number of larvae tested was

21 and 21 for the control and experimental groups, respectively. ** $p < 0.01$, NS: $p \geq$

0.05 (Mann–Whitney U-test with Bonferroni correction). (D) Percentage of larvae that

showed a response in the next 1 or 12 frames after blue light stimulation. Number of

events analyzed: 1960 or 111 for 1- or 12-frame continuous inactivity prior to blue light

illumination, respectively. **** $p < 0.0001$ (chi-squared test).

Figure 1—figure supplement 3. CRISPR-knock-out screen for genes that regulate

larval sleep.

Total sleep amounts in CRISPR-knock-out mutants were measured for 18 h from the

beginning of the second instar. The data for Control, *PK2-R1,* and *Oamb* mutants are

identical to those shown in Figure 1. Box plots are generated so that the center line

indicates the median, the box limits indicate the upper and lower quartiles, and the

whiskers indicate the minimum-to-maximum range. *** $p < 0.005$ (Kruskal–Wallis

one-way ANOVA and post hoc Mann–Whitney U-test with Bonferroni correction).

Number of larvae tested: 475 for the control and 10–52 for each mutant.

Figure 1—figure supplement 4. Effects of neuronal manipulation on larval sleep

Sleep amounts in larvae in which distinct neuronal populations were silenced. Kir2.1

expression was driven by each enhancer-GAL4 for "hit" genes identified in the

CRISPR-knock-out screen. Each dot represents an individual animal in these panels,

"N" indicates the number of biologically independent animals per group, and the thick

line and thin error bars indicate the median and interquartile range (Q1–Q3),

respectively. *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, NS: ≥ 0.05 (Mann–Whitney U-test

with Bonferroni correction)

Figure 1—figure supplement 5. Larval sleep phenotypes in PK2-R1 mutants

quantified over 18 h and the first 6 h.

(A–C) Sleep metrics quantified over an 18-hour period in second-instar larvae, whereas

(D–F) show the same metrics quantified over the first 6 hours in the same animals.

Total sleep is shown in (A) and (D), bout number in (B) and (E), and mean bout length

in (C) and (F). Each dot represents an individual larva; the thick line indicates the

median and the thin error bars indicate the interquartile range (Q1–Q3). Numbers in

parentheses indicate N, the number of biologically independent animals per group. ** p

< 0.01, * $p < 0.05$, NS: ≥ 0.05 (Mann–Whitney U-test with Bonferroni correction).

Figure 2. IPCs and Dilps negatively regulate larval sleep

(A) Triple labeling of *PK2-R1* neurons expressing nuclear-localized RFP (magenta),

Dilp3 neurons expressing nuclear-localized GFP (green), and anti-Dilp2 positive cells

(blue). Top panels show signals in the larval brain and the VNC. Bottom panels show

magnified images of the white-squared area in each top panel, focusing on the

dorsomedial brain region where the cell bodies of IPCs are located. Note that all IPCs

labeled with *Dilp3-GAL4* overlapped with *PK2-R1²A-LexA*-expressing cells. Similar

results were obtained from 5 independent samples of the same genotype. (B)

Simultaneous detection of *PK2-R1* neurons expressing nuclear-localized RFP

(magenta), *Dilp5* neurons expressing nuclear-localized GFP (green), and anti-Dilp2

positive cells (blue). Similar results were obtained from 5 independent samples of the
same genotype. (C) Effect of IPCs silencing on larval sleep. Each dot represents an
individual animal; in this and the following panels, "N" indicates the number of
biologically independent animals per group, and the thick line and thin error bars
indicate the median and interquartile range (Q1–Q3), respectively. *** $p < 0.001$, ** p
< 0.01, * $p < 0.05$ (Mann–Whitney U-test with Bonferroni correction). (D) Sleep
amounts in *Dilp3* or *Dilp5* null mutants. *** $p < 0.001$, ** $p < 0.01$ (Mann–Whitney U-
test with Bonferroni correction).
702

Figure 2—figure supplement 1. Locomotion speeds are not consistently affected by

genetical manipulations of *PK2-R1*, IPCs, or *Dilps*

(A) Larval locomotion speed in controls and PK2-R1 mutants. (B-C) Effects of

silencing PK2-R1 neurons (B) and IPCs (C) on larval locomotion speed, shown together

with the corresponding genetic controls. (D) Larval locomotion speed in Dilp3 or Dilp5

null mutants. Average locomotion speed during wake periods was calculated from the

datasets shown in Figures 1A, 1B, 2C, and 2D. N indicates the number of biologically

independent larvae per group. Each dot represents an individual larva; thick lines

indicate the median and thin error bars indicate the interquartile range (Q1-Q3). *** $p <$

0.0001, * $p < 0.05$, NS: $p \geq 0.05$ (Mann-Whitney U-test with Bonferroni correction).

Figure 3. *Hug* neurons negatively regulate larval sleep

(A) Sleep amounts in *Hug* CRISPR-knockout mutant larvae. **** $p < 0.0001$ (Mann–

Whitney U-test). Each dot represents an individual animal; in this and the following

panels, "N" indicates the number of biologically independent animals per group, and the

thick line and thin error bars indicate the median and interquartile range (Q1–Q3),

respectively. (B) Effect of silencing Hug neurons on larval sleep amount. ** $p < 0.01$, *

$p < 0.05$ (Mann–Whitney U-test with Bonferroni correction). (C) Larval sleep changes

induced by optogenetic activation of *Hug* neurons. For each genotype, sleep duration in

the 1-h light-ON period was normalized to that in the 1-h light-OFF phase. * $p < 0.05$

(Mann–Whitney U-test with Bonferroni correction). (D) Larval sleep during

thermogenetic activation of *Hug* neurons. ** $p < 0.01$, * $p < 0.05$ (Mann–Whitney U-

test with Bonferroni correction). (E) Visualization of HugPC neurons and IPCs labeled

by rCD2::RFP (magenta) and mCD8::GFP (green), respectively. The bottom panels

show magnified images of the white-squared dorsomedial region in the top panels,

where HugPC neurons project their axons close to the cell bodies of the IPCs. While the

top panels are z-stacks of 122 image slices (1 μm interval) covering the entire brain

tissue, the bottom panels are projections of 60 slices centering around IPCs. Similar

results were obtained from 5 independent samples of the same genotype. (F) The effect

of neuronal silencing confined to the HugPC subpopulation. ** $p < 0.01$, * $p < 0.05$

(Mann–Whitney U-test with Bonferroni correction).

Figure 3—figure supplement 1. Effects of Hug pathway manipulations on larval

locomotion speed.

(A) Locomotion speed of Hug knockout mutants during wake periods. (B) Effects of

silencing Hug neurons on locomotion speed. (C-D) Effects of activating Hug neurons

on locomotion speed, shown as the light ON/light OFF ratio (C) and the absolute

locomotion speed (D). (E) Effects of silencing HugPC neurons on locomotion speed.

Locomotion speed was quantified from the same videos used for the larval sleep assays

in Figure 3. Each dot represents an individual larva; thick lines indicate the median and

thin error bars indicate the interquartile range (Q1-Q3). N indicates the number of

biologically independent animals per group. NS $p \geq 0.05$ (Mann-Whitney U-test) in (A).

*** $p < 0.0001$, * $p < 0.05$, NS $p \geq 0.05$ (Mann-Whitney U-test with Bonferroni

correction) in (B-E).

Figure 3—figure supplement 2. Generation of *HugPC-LexA* transgenic lines

(A-D) Labeling patterns of four *HugPC-LexA* strains in larval CNS visualized by

rCD2::RFP. Four transgenic lines were generated by injecting the same plasmid into the

following landing sites individually: su(Hw)attP5, VK00005, attP2, and attP40.

Magenta arrowheads represent cell bodies apparently located outside the previously

classified HugPC subpopulation (Bader et al., 2007).

Figure 3—figure supplement 3. Silencing HugPC neurons reduces larval food

intake measured by a dye-based assay.

After a 30 min starvation, larvae were allowed to feed on red yeast paste placed on an

apple juice agar plate, and food intake was quantified as the ratio of the red dye-stained

body surface area to the total body surface area (ImageJ). Each dot represents an

individual larva; the thick line indicates the median and the thin error bars indicate the

interquartile range (Q1–Q3). ** $p < 0.01$ (Mann–Whitney U-test).

Figure 4. Activation of Hug neurons triggers Ca2+ responses in larval IPCs.

(A) Schematic workflow for assessing intracellular Ca2+ levels in IPCs while

thermogenetically activating Hug neurons. Larvae were exposed to low (18°C) or high

(30°C) temperature for 1 h, followed by imaging of CRTC::GFP in IPCs and calculation

of the CRTC::GFP nuclear localization index (NLI; see Methods). (B–C) Nuclear-

localized CRTC::GFP signal (CRTC::GFP NLI) in larval IPCs under low-temperature

(B) or high-temperature (C) conditions, with or without thermogenetic activation of

Hug neurons. Each dot represents an individual cell; the thick line indicates the median

and the thin error bars indicate the interquartile range (Q1–Q3). "N" indicates the

number of cells analyzed per group. **** $p < 0.0001$, NS $p \geq 0.05$ (Mann–Whitney U-

test). (D) Working model of a neuronal network in which Hug neurons activate IPCs to

regulate larval sleep

Figure 4-figure supplement 1. Axonal projections of HuginPC neurons visualized

with a presynaptic marker.

Representative whole-mount image of the larval central nervous system showing Syt-

eGFP signals expressed in HuginPC neurons. Scale bar, 100 μm.

Figure 5. Hug peptides induce Ca^{2+} responses in larval IPCs via PK2-R1

(A) Schematic flow of peptide application followed by Ca^{2+} imaging. (B–C) Ca^{2+}

responses in laval IPCs during *ex vivo* bath application of either glucose (B) or Hug

peptides (C). In this and the following panels, 'N' indicates the number of cells used for

each group, and the thick line and thin error bar represent the median and interquartile

range (Q1–Q3), respectively. **** $p < 0.0001$, *** $p < 0.001$ (Mann–Whitney U-test

with Bonferroni correction). (D) Representative images of larval IPCs upon peptide

application. Scale bars, 2 μm. (E) Anti-Dilp3 signal intensity measured within the

cytosolic areas of larval IPCs. *** $p < 0.001$, * $p < 0.05$ (Mann–Whitney U-test with

Bonferroni correction). (F–G) Ca^{2+} responses in larval IPCs derived from *PK2-R1*

knockout mutants, measured after either glucose (F) or Hug peptide application (G). **

$p < 0.01$, NS $p \geq 0.05$ (Mann–Whitney U-test with Bonferroni correction). (H) A model

where Hug peptides, but not glucose, activate IPCs via the PK2-R1 receptor in the

larval brain.

Figure 5—figure supplement 1. D-glucose but not Hug peptide cause Dilp2 reduction in larval IPCs

(A–B) Anti-Dilp2 signal intensity measured in larval IPCs after glucose (A) or Hug-γ application (B). In this and the following panels, 'N' indicates the number of cells used for each group, and the thick line and thin error bar represent the median and interquartile range (Q1–Q3), respectively. **** $p < 0.0001$, * $p < 0.05$ (Mann–Whitney U-test). (C) Anti-Dilp3 signal intensity measured within the cytosolic areas of larval IPCs. NS ≥ 0.05 (Mann–Whitney U-test).

Figure 5—figure supplement 2. Representative images of Dilp3 immunoreactivity in larval IPCs after bath application of Hug-γ peptide.

Larval IPCs were labeled with Dilp3>mCherry (top row), and endogenous Dilp3 was detected by anti-Dilp3 immunostaining (bottom row). Representative brains from buffer-only controls (Brain #1 and #2) and Hug-γ peptide application (Brain #3 and #4) are shown. Scale bars: 25 μm (Brains #1–3) and 10 μm (Brain #4).

Figure 6. Distinct impacts of Hugin/PK2-R1 axis on wake/sleep control in larvae and adult

(A) Total sleep amounts in *Hug kno*ckout mutant adults. NS $p \geq 0.05$ (Mann–Whitney U-test). In these and the following panels, 'N' indicates the number of biologically independent animals used for each group, and the thick line and thin error bar represent the median and interquartile range (Q1–Q3), respectively. NS $p \geq 0.05$ (Mann–Whitney U-test). (B) Sleep amounts of adults in which HugPC neurons were silenced. *** $p < 0.001$, NS $p \geq 0.05$ (Mann–Whitney U-test with Bonferroni correction). (C) Sleep amounts of *Dilp3* or *Dilp5* null mutant adults. *** $p < 0.001$ (Mann–Whitney U-test with Bonferroni correction). (D) Schematic flow of peptide application experiments using adult brains. (E–F) Ca^{2+} sponses (E) or anti-Dilp3 signal intensity (F) in adult IPCs after Hug peptide application. ** $p < 0.01$, NS $p \geq 0.05$ (Mann–Whitney U-test with Bonferroni correction).

Figure 6—figure supplement 1. Sleep patterns of adult flies with genetical manipulations in the Hugin and insulin pathways

(A) Experimental setup for monitoring adult sleep using the *Drosophila* Activity Monitor (DAM). (B) Total daily sleep duration in adult flies carrying CRISPR-knockout mutations in genes whose larval sleep was significantly altered in Figure 1B. Each dot represents an individual fly; the thick line indicates the median and the thin error bars indicate the interquartile range (Q1–Q3). *** $p < 0.01$, ** $p < 0.05$ (Kruskal–Wallis one-way ANOVA followed by post hoc Mann–Whitney U-test with Bonferroni correction). (C) Daytime (left) and nighttime (right) sleep amounts in flies in which HugPC neurons were silenced. (D) Daytime (left) and nighttime (right) sleep amounts in Dilp3 or Dilp5 null mutants. In (C) and (D), numbers in parentheses indicate N, the number of biologically independent animals per group. *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, NS $p \geq 0.05$ (Mann–Whitney U-test with Bonferroni correction).

Figure 6—figure supplement 2. Morphologies of *PK2-R1* neurons, HugPC neurons, and IPCs in the adult brain.

(A) Dual labeling of *PK2-R1* neurons and IPCs expressing nuclear-localized RFP (magenta) and GFP (green), respectively. The top panels show signals in the entire adult brain. The bottom panels show magnified images of the white-squared area in each corresponding top panel, focusing on the dorsomedial brain region where the cell bodies of IPCs are located. Similar results were obtained from 5 independent samples of the same genotype. (B) Visualization of HugPC neurons and IPCs labeled by mCD8::GFP (magenta) and the anti-Dilp3 antibody (green), respectively. Similar results were obtained from 4 independent samples of the same genotype.

Figure 6—figure supplement 3. Adult sleep architecture and waking activity following genetic manipulations of the hugin and insulin pathways.

(A) Bout number per day (left), mean bout length (middle), and mean beam crossings

during wakefulness (right) in Hug^attP mutants. (B) The same metrics in flies with

HugPC neuron silencing. (C) The same metrics in Dilp3[1] and Dilp5^1 null mutants.

(D) Total daily sleep in flies with Dilp3-positive neuron silencing. Each dot represents

an individual fly; the thick line indicates the median and the thin error bars indicate the

interquartile range (Q1–Q3). Numbers in parentheses indicate N, the number of

biologically independent animals per group. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, NS

$p \geq 0.05$ (Mann–Whitney U-test with Bonferroni correction).

Figure 7. A schematic model of the Hugin/PK2-R1/Dilps axis in larval and adult

sleep

Schematic summary comparing larvae and adults. In larvae, bath application of hugin

peptides activates IPCs, and both hugin signaling (HugPC neurons/hugin peptides) and

IPC output (including Dilp3) act to suppress larval sleep. In adults, hugin peptides do

not activate IPCs (dashed arrow), whereas IPC/Dilp3 signaling promotes adult sleep.

References

Ahnaou, A., & Drinkenburg, W. H. I. M. (2011). Neuromedin U2 receptor signaling
mediates alteration of sleep-wake architecture in rats. *Neuropeptides*, 45(2), 165–
174. Retrieved from https://doi.org/10.1016/j.npep.2011.01.004

Bader, R., Colomb, J., Pankratz, B., Schröck, A., Stocker, R. F., & Pankratz, M. J.
(2007). Genetic dissection of neural circuit anatomy underlying feeding behavior
in Drosophila: Distinct classes of hugin-expressing neurons. *Journal of*
Comparative Neurology, 502(5), 848–856. Retrieved from
https://doi.org/10.1002/cne.21342

Baines, R. A., Uhler, J. P., Thompson, A., Sweeney, S. T., & Bate, M. (2001). *Altered*
Electrical Properties in Drosophila Neurons Developing without Synaptic
Transmission.

Barlow, I. L., & Rihel, J. (2017, June 1). Zebrafish sleep: from geneZZZ to neuronZZZ.
Current Opinion in Neurobiology. Elsevier Ltd. Retrieved from
https://doi.org/10.1016/j.conb.2017.02.009

Blumberg, M. S., H Seelke, A. M., Lowen, S. B., & Karlsson, K. A. (2005). Dynamics
of sleep–wake cyclicity in developing rats. *PNAS*, 102(41). Retrieved from
www.pnas.orgcgidoi10.1073pnas.0506340102

Bonheur, M., Swartz, K. J., Metcalf, M. G., Wen, X., Zhukovskaya, A., Mehta, A., …
Hattori, D. (2023). A rapid and bidirectional reporter of neural activity reveals
neural correlates of social behaviors in Drosophila. *Nature Neuroscience*, 26(7),
1295–1307. Retrieved from https://doi.org/10.1038/s41593-023-01357-w

Burke, C. J., Huetteroth, W., Owald, D., Perisse, E., Krashes, M. J., Das, G., …
Waddell, S. (2012). Layered reward signalling through octopamine and dopamine
in Drosophila. *Nature*, 492(7429), 433–437. Retrieved from
https://doi.org/10.1038/nature11614

Chiu, C. N., Rihel, J., Lee, D. A., Singh, C., Mosser, E. A., Chen, S., … Prober, D. A.
(2016). A Zebrafish Genetic Screen Identifies Neuromedin U as a Regulator of
Sleep/Wake States. *Neuron*, 89(4), 842–856. Retrieved from
https://doi.org/10.1016/j.neuron.2016.01.007

Churgin, M. A., Szuperak, M., Davis, K. C., Raizen, D. M., Fang-Yen, C., & Kayser,
M. S. (2019). Quantitative imaging of sleep behavior in Caenorhabditis elegans

and larval Drosophila melanogaster. *Nature Protocols 2019 14:5*, 14(5), 1455–

1488. Retrieved 25 December 2021 from https://doi.org/10.1038/s41596-019-

0146-6

Davis, K. F., Parker, K. P., & Montgomery, G. L. (2004). *Sleep in infants and young*

children: Part one: normal sleep. Journal of Pediatric Health Care (Vol. 18, pp.

65–71). Mosby. Retrieved 25 December 2021 from

https://doi.org/10.1016/S0891-5245(03)00149-4

Deng, B., Li, Q., Liu, X., Cao, Y., Li, B., Qian, Y., … Rao, Y. (2019).

Chemoconnectomics: Mapping Chemical Transmission in Drosophila. *Neuron*,

101(5), 876-893.e4. Retrieved from https://doi.org/10.1016/j.neuron.2019.01.045

Dubowy, C., & Sehgal, A. (2017). Circadian rhythms and sleep in Drosophila

melanogaster. *Genetics*, 205(4), 1373–1397. Retrieved from

https://doi.org/10.1534/genetics.115.185157

Enell, L. E., Kapan, N., Söderberg, J. A. E., Kahsai, L., & Nässe, D. R. (2010). Insulin

signaling, lifespan and stress resistance are modulated by metabotropic GABA

receptors on insulin producing cells in the brain of Drosophila. *PLoS ONE*, 5(12).

Retrieved from https://doi.org/10.1371/journal.pone.0015780

Frank, M. G., Ruby, N. F., Heller, H. C., & Franken, P. (2017). Development of

circadian sleep regulation in the rat: A longitudinal study under constant

conditions. *Sleep*, 40(3). Retrieved from https://doi.org/10.1093/sleep/zsw077

Furusawa, K., Ishii, K., Tsuji, M., Tokumitsu, N., Hasegawa, E., & Emoto, K. (2023).

Presynaptic UBE3A E3 ligase promotes synapse elimination by downregulation

of BMP signaling. *Science*, 381, 1197-1205. Retrieved from http://doi.org/

10.1126/science.ade8978.

Groth, A. C., Fish, M., Nusse, R., & Calos, M. P. (2004). Construction of Transgenic

Drosophila by Using the Site-Specific Integrase From Phage φC31. *Genetics*,

(166), 1775–1782. Retrieved from

https://academic.oup.com/genetics/article/166/4/1775/6052665

Hamada, F. N., Rosenzweig, M., Kang, K., Pulver, S. R., Ghezzi, A., Jegla, T. J., &

Garrity, P. A. (2008). An internal thermal sensor controlling temperature

preference in Drosophila. *Nature*, 454(7201), 217–220. Retrieved from

https://doi.org/10.1038/nature07001

Hendricks, J. C., Lu, S., Kume, K., Yin, J. C. P., Yang, Z., & Sehgal, A. (2003). Gender
dimorphism in the role of cycle (BMAL1) in rest, rest regulation, and longevity in
Drosophila melanogaster. *Journal of Biological Rhythms*, 18(1), 12–25. Retrieved
from https://doi.org/10.1177/0748730402239673

Hückesfeld, S., Peters, M., & Pankratz, M. J. (2016). Central relay of bitter taste to the
protocerebrum by peptidergic interneurons in the Drosophila brain. *Nature*
Communications, 7. Retrieved from https://doi.org/10.1038/ncomms12796

Hückesfeld, S., Schlegel, P., Miroschnikow, A., Schoofs, A., Zinke, I., Haubrich, A. N.,
… Pankratz, M. J. (2021). Unveiling the sensory and interneuronal pathways of
the neuroendocrine connectome in Drosophila. *ELife*. Retrieved from
https://doi.org/10.7554/eLife

Inagaki, H. K., Jung, Y., Hoopfer, E. D., Wong, A. M., Mishra, N., Lin, J. Y., …
Anderson, D. J. (2014). Optogenetic control of Drosophila using a red-shifted
channelrhodopsin reveals experience-dependent influences on courtship. *Nature*
Methods, 11(3), 325–332. Retrieved from https://doi.org/10.1038/nmeth.2765

Lakhotia SC, Ranganath HA, editors. 2021. *Experiments with Drosophila for Biology*
Courses. Indian Academy of Sciences, Bengaluru, India.

Kitatani, Y., Tezuka, A., Hasegawa, E., Yagagi, S., Togashi, K., Tsuji, M., Kondo, S.,
Parrish, J. Z., & Emoto, K. (2020). *Drosophila miR-87* promotes dendrite
regeneration by targeting the transcriptional repressor Tramtrack69. *PLoS*
Genetics, 16, e1008942.

Kréneisz, O., Chen, X., Fridell, Y. W. C., & Mulkey, D. K. (2010). Glucose increases
activity and Ca2+ in insulin-producing cells of adult Drosophila. *NeuroReport*,
21(17), 1116–1120. Retrieved from
https://doi.org/10.1097/WNR.0b013e3283409200

Kume, K., Kume, S., Park, S. K., Hirsh, J., & Jackson, F. R. (2005). Dopamine Is a
Regulator of Arousal in the Fruit Fly. *Journal of Neuroscience*, 25(32), 7377–
7384. Retrieved 10 December 2024 from
https://doi.org/10.1523/JNEUROSCI.2048-05.2005

Morikawa, R., Kanamori, T., Yasunaga, K., & Emoto, K. (2011). Different levels of the
TRIM protein Asap regulate distinct axonal projections of *Drosophila* sensory
neurons. *Proceedings of the National Academy of Sciences of the United States of*

America, 108, 19389-19394.

Nakamizo-Dojo, M., Ishii, K., Yoshino, J., Tsuji, M., & Emoto, K. (2023). Descending

GABAergic pathway links brain sugar-sensing to peripheral nociceptive gating in

Drosophila. *Nature Communications*, 14(1). Retrieved from

https://doi.org/10.1038/s41467-023-42202-9

Oh, Y., Lai, J. S. Y., Mills, H. J., Erdjument-Bromage, H., Giammarinaro, B.,

Saadipour, K., … Suh, G. S. B. (2019). A glucose-sensing neuron pair regulates

insulin and glucagon in Drosophila. *Nature*, 574(7779), 559–564. Retrieved from

https://doi.org/10.1038/s41586-019-1675-4

Ohhara, Y., Blick, M., Park, D., Yoon, S. E., Kim, Y. J., Pankratz, M. J., … Yamanaka,

N. (2024). A Neuropeptide Signaling Network That Regulates Developmental

Timing and Systemic Growth in Drosophila. *The Journal of Comparative*

Neurology, 532(10), e25677. Retrieved from https://doi.org/10.1002/cne.25677

Okamoto, N., Nishimori, Y., & Nishimura, T. (2012). Conserved role for the

Dachshund protein with Drosophila Pax6 homolog eyeless in insulin expression.

Proceedings of the National Academy of Sciences of the United States of America,

109(7), 2406–2411. Retrieved from https://doi.org/10.1073/pnas.1116050109

Omamiuda-Ishikawa, N., Sakai, M., & Emoto, K. (2020). A pair of ascending neurons

in the subesophageal zone mediates aversive sensory inputs-evoked backward

locomotion in *Drosophila* larvae. *PLoS Genetics,* 16, e1009120.

Paul J. Shaw, Chiara Cirelli, Ralph J. Greenspan, & Giulio Tononi. (2000). Correlates

of Sleep and Waking in Drosophila melanogaster. *SCIENCE*, (287), 1834–1837.

Poe, A. R., Zhu, L., Szuperak, M., Mcclanahan, P. D., Anafi, R. C., Scholl, B., …

Kayser, M. S. (2023). *Developmental emergence of sleep rhythms enables long-*

term memory in Drosophila. Retrieved from https://www.science.org

Raizen, D. M., Zimmerman, J. E., Maycock, M. H., Ta, U. D., You, Y. J., Sundaram,

M. V., & Pack, A. I. (2008). Lethargus is a Caenorhabditis elegans sleep-like

state. *Nature*, 451(7178), 569–572. Retrieved from

https://doi.org/10.1038/nature06535

Rosenkilde, C., Cazzamali, G., Williamson, M., Hauser, F., Søndergaard, L., DeLotto,

R., & Grimmelikhuijzen, C. J. P. (2003). Molecular cloning, functional

expression, and gene silencing of two Drosophila receptors for the Drosophila

1005 neuropeptide pyrokinin-2. *Biochemical and Biophysical Research*

1006 *Communications*, 309(2), 485–494. Retrieved from

1007 https://doi.org/10.1016/j.bbrc.2003.08.022

1008 Scammell, T. E., Arrigoni, E., & Lipton, J. O. (2017, February 22). 【Review】Neural

1009 Circuitry of Wakefulness and Sleep. *Neuron*. Cell Press. Retrieved from

1010 https://doi.org/10.1016/j.neuron.2017.01.014

1011 Schlegel, P., Texada, M. J., Miroschnikow, A., Schoofs, A., Hückesfeld, S., Peters, M.,

1012 … Pankratz, M. J. (2016a). Synaptic transmission parallels neuromodulation in a

1013 central food-intake circuit. *ELife*, 5(NOVEMBER2016). Retrieved from

1014 https://doi.org/10.7554/eLife.16799.001

1015 Schlegel, P., Texada, M. J., Miroschnikow, A., Schoofs, A., Hückesfeld, S., Peters, M.,

1016 … Pankratz, M. J. (2016b). Synaptic transmission parallels neuromodulation in a

1017 central food-intake circuit. *ELife*, 5(NOVEMBER2016). Retrieved from

1018 https://doi.org/10.7554/eLife.16799.001

1019 Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25

1020 years of image analysis. *Nature Methods*, 9(7), 671–675. Retrieved from

1021 https://doi.org/10.1038/nmeth.2089

1022 Schoofs, A., Hückesfeld, S., Schlegel, P., Miroschnikow, A., Peters, M., Zeymer, M.,

1023 … Pankratz, M. J. (2014). Selection of Motor Programs for Suppressing Food

1024 Intake and Inducing Locomotion in the Drosophila Brain. *PLoS Biology*, 12(6).

1025 Retrieved from https://doi.org/10.1371/journal.pbio.1001893

1026 Schwarz, J. E., King, A. N., Hsu, C. T., Barber, A. F., & Sehgal, A. (2021). Hugin +

1027 neurons provide a link between sleep homeostat and circadian clock neurons.

1028 *PNAS*. Retrieved from https://doi.org/10.1073/pnas.2111183118/-

1029 /DCSupplemental

1030 Shafer, O. T., & Keene, A. C. (2021, January 11). The Regulation of Drosophila Sleep.

1031 *Current Biology*. Cell Press. Retrieved from

1032 https://doi.org/10.1016/j.cub.2020.10.082

1033 Sorribes, A., Porsteinsson, H., Arnardóttir, H., Jóhannesdóttir, I. P., Sigurgeirsson, B.,

1034 de Polavieja, G. G., & Karlsson, K. (2013). The ontogeny of sleep-wake cycles in

1035 zebrafish: A comparison to humans. *Frontiers in Neural Circuits*, 7(NOV).

Retrieved from https://doi.org/10.3389/fncir.2013.00178

Szuperak, M., Churgin, M. A., Borja, A. J., Raizen, D. M., Fang-Yen, C., & Kayser, M.

S. (2018a). A sleep state in Drosophila larvae required for neural stem cell

proliferation. *ELife*, 7. Retrieved 25 December 2021 from

https://doi.org/10.7554/ELIFE.33220

Szuperak, M., Churgin, M. A., Borja, A. J., Raizen, D. M., Fang-Yen, C., & Kayser, M.

S. (2018b). A sleep state in Drosophila larvae required for neural stem cell

proliferation. *ELife*. Retrieved from https://doi.org/10.7554/eLife.33220.001

Veenstra, J. A., Agricola, H. J., & Sellami, A. (2008). Regulatory peptides in fruit fly

midgut. *Cell and Tissue Research*, 334(3), 499–516. Retrieved from

https://doi.org/10.1007/s00441-008-0708-3

Yoshino, J., Morikawa, R., Hasegawa, E., & Emoto, K. (2017). Neural circuitry that

evokes escape behavior in response to nociceptive stimuli in *Drosophila* larvae.

Current Biology, 27, 2499-2504.

Yoshino, J., Mali, S., Williams, C.R., Morita, T., Emerson, C., Arp, C., Miller, S., Yin,

C., Motoyoshi, M., Hemmi, C., Ishii, K., Bautista, D., Emoto, K., & Parrish, J.Z.

(2025). *Drosophila* epidermal cells are intrinsically mechanosensitive and

modulate nociceptive behavior outputs. *ELife*, 13, PR95379 doi:

10.7554/eLife.95379

Figure 1

A

Total sleep (h/18 h)

***

***

Cont $PK2\text{-}R1^{attP}$ $Oamb^{attP}$

(N) (475)   (24)   (22)

B

Total sleep (h/18 h)

***

***

$PK2\text{-}R1^{2A\text{-}GAL4}$

$UAS\text{-}Kir2.1$

(N) (28) (28) (21)

C

$PK2\text{-}R1^{2A\text{-}GAL4} > UAS\text{-}Kir2.1::EGFP$

100 µm

Figure 1—figure supplement 1

A

Image acquisition

Larval sleep detection

Camera

24-well chamber

Inactive | Active | • • •

Inactive | Inactive | Inactive | • • • | Inactive | Active — Awake

Inactive | Inactive | Inactive | • • • | Inactive | Inactive — **Sleep**

Frame 0   1   2   3   11   12

B

Camera

Top view

Φ11 mm

Φ7 mm

60 mm

90 mm

Side view

Glass plate

3 mm   10 mm

1 mm   Agar   PDMS

C

Step 1)
Image processing

Original picture

Blurring
(11 ways)

Binarization
(62 ways)

Total of
11 x 62 x 1,000 = 682,000
combinations

Step 2)
Pixel change detection

Current
frame − Previous
frame = **Pixel
change**

**Selection of the best
processing condition**
that provides the smallest discrepancy
in detected sleep amounts
compared to manual annotation

Step 3)
Active/inactive state determination

**Pixel
change**

Active

Pixel threshold
(1000 ways)

Inactive

Time

Figure 1—figure supplement 2

A

B

C

D

Figure 1—figure supplement 3

Figure 1—figure supplement 4

Figure 1—figure supplement 5

Figure 2

A

| *Dilp3-GAL4 > UAS-Stinger* | *PK2-R1²ᴬ⁻ᴸᵉˣᴬ > LexAop-tdTomato.nls* | Anti-Dilp2 | Merge |

100 μm

30 μm

B

| *Dilp5-GAL4 > UAS-Stinger* | *PK2-R1²ᴬ⁻ᴸᵉˣᴬ > LexAop-tdTomato.nls* | Anti-Dilp2 | Merge |

100 μm

30 μm

C

D

Figure 2—figure supplement 1

Figure 3

Figure 3—figure supplement 1

Figure 3—figure supplement 2

Figure 3—figure supplement 3

Figure 4

A

(1) Thermogenetic stimulation of *Hug* neurons

In vivo for 1 h

30°C

OR

18°C

Hug neurons

?

IPCs

(2) Measurement of CRTC::GFP signal in IPCs

Ca²⁺ level

Low   High

mCD8::mCherry

CRTC::GFP

Min   Max
Pixel signal intensity

(3) Calculation of the nuclear localization index of GFP (CRTC::GFP NLI)

$$CRTC::GFP\ NLI = \frac{F_{nuc} - F_{cyto}}{F_{nuc} + F_{cyto}}$$

B

Low temperature

CRTC::GFP NLI

****

****

Hug²ᴬ⁻ᴸᵉˣᴬ		
LexAop-TrpA1		
UAS-CRTC::GFP		
Dilp3-GAL4		

(N) (25) (30) (44)

C

High temperature

****

NS

Hug²ᴬ⁻ᴸᵉˣᴬ		
LexAop-TrpA1		
UAS-CRTC::GFP		
Dilp3-GAL4		

(N) (56) (39) (24)

D

Hug neurons

⊕

IPCs

HugPC-GAL4 >
UAS-UAS-syt.eGFP (Axon marker)

0 μm 100

Figure 5

A

(1) *Ex vivo* peptide application for 5 min

Dissected larval brain

Glucose

OR **Hug-γ** OR **PK-2** OR **Neuromedin U (NMU)**

Mammalian homolog of *hugin*,

(2) Measurement of CRTC::GFP signal in IPCs

CRTC translocates into the nucleus upon neural activation

Ca²⁺ level

Low High

mCD8::mCherry

CRTC::GFP

Min Max
Pixel signal intensity

PK2-R1⁺/⁺ (Dilp3 > CRTC::GFP)

Glucose application

Peptide application

B

CRTC::GFP NLI

****

Glucose − +

(N) (30) (24)

C

CRTC::GFP NLI

*** *** ***

Peptide − Hug-γ PK-2 NMU

(N) (69) (35) (65) (55)

D

mCD8::mCherry

CRTC::GFP

Peptide − Hug-γ PK-2

Min Max
Pixel signal intensity

E

Anti-Dilp3 signal intensity (A.U.)

* ***

Peptide − Hug-γ PK-2

(N) (38) (58) (20)

PK2-R1⁻/⁻ (Dilp3 > CRTC::GFP)

Glucose application

Peptide application

F

CRTC::GFP NLI

**

Glucose − +

(N) (47) (48)

G

CRTC::GFP NLI

NS NS NS

Peptide − Hug-γ PK-2 NMU

(N) (67) (43) (60) (45)

H

IPCs

PK2-R1⁺/⁺

With *PK2-R1* expression in IPCs

Hugin-related peptides
Hug-γ PK-2 NMU

⊕

PK2-R1

Ca²⁺ response

IPCs

IPCs

PK2-R1⁻/⁻

Without *PK2-R1* expression in IPCs

Hugin-related peptides
Hug-γ PK-2 NMU

IPCs Ca²⁺ response

Figure 5—figure supplement 1

Figure 5—figure supplement 2

Control (buffer only application)

Brain #1　　　　Brain #2

Hug-g peptide application

Brain #3　　　　Brain #4

Dilp3 > mCherry

Anti-Dilp3

Bar = 10 um

Figure 6

Sleep amount analysis

A — Total sleep (min/day) — NS — Cont, *Hug^attP* — (N) (21) (18)

B — Total sleep (min/day) — NS, *** — *HugPC-GAL4*, *UAS-Kir2.1* — (N) (45) (44) (47)

C — Total sleep (min/day) — ***, *** — Cont, *Dilp3^1*, *Dilp5^1* — (N) (46) (44) (41)

Ex vivo peptides application

D — Hugin precursor — Hug-γ OR PK-2 — Dissected adult brain — Measurement of Ca²⁺ responses in IPCs using CRTC::GFP

E — CRTC::GFP NLI — NS, NS — Peptide −, Hug-γ, PK-2 — (N) (48) (25) (48)

F — Anti-Dilp3 signal intensity (A.U.) — NS, ** — Peptide −, Hug-γ, PK-2 — (N) (48) (25) (48)

Figure 6—figure supplement 1

Figure 6—figure supplement 2

Morphological analysis

Figure 6—figure supplement 3

Figure 7

Larvae	Adults

HugPC neurons / Hugin peptides

⊕

IPCs / Dilp3

⊖

Larval sleep

z^Z_z

HugPC neurons / Hugin peptides

IPCs / Dilp3

⊕

Adult sleep

z^Z_z