Thymic selection of the T cell receptor repertoire is biased toward autoimmunity in females

Hélène Vantomme[1,2], Valentin Quiniou[1], Leslie Adda[1§], Charline Jouannet[1§], Vanessa Mhanna[1§], Céline Albalaa[1$], Pierre Barennes[1$], Nicolas Coatnoan[1,2$], Vimala Diderot[1,2$], Johanna Dubois[1$], Gwladys Fourcade[1$], Kenz Le Gouge[1$], Otriv Frédéric Nguekap Tchoumba[1$], Martin Pezous[1$], Paul Stys[1$], Adrien Six[1], Encarnita Mariotti-Ferrandiz[1,3]* and David Klatzmann[1,2]*#.

§contributed equally, alphabetical order
$contributed equally, alphabetical order
*co-senior authors
#corresponding author

1. Immunology, Immunopathology, Immunotherapy i3 Lab, INSERM UMRS-959, Sorbonne Université, Paris, France
2. Biotherapy (CIC-BTi) and Inflammation-Immunolopathology-Biotherapy Department (i2B), Assistance Publique -Hôpitaux de Paris (AP-HP), Paris France
3. Institut Universitaire de France

Corresponding author :
Prof. David KLATZMANN MD, PhD
I3 Lab (Immunology Immunopathology Immunotherapy) INSERM UMRS 959 – Sorbonne Université
Bâtiment CERVI, GHPS Pitié-Salpêtrière, 83 boulevard de l'Hôpital, 75013 Paris, France
Tel : +33 1 42 17 74 61
Email : david.klatzmann@sorbonne-universite.fr

The authors have declared that no conflict of interest exists.

Abstract: (200 words)

Women represent about 80% of patients with autoimmune diseases. This may partly result

from sex-based differences in T cell receptor (TCR) selection during thymocyte development,

potentially influenced by hormones and the lower expression of the Autoimmune Regulator

(AIRE) transcription factor in females.

To investigate this, we analyzed sex-specific differences in TCR generation and selection. We

examined TCR repertoires in double-positive thymocytes and single-positive thymic cells,

including CD8$^+$ and CD4$^+$ effector T cells and regulatory T cells (Tregs), derived from male and

female organ donors. Minimal sex-based differences were observed in V and J gene usage,

and there were no notable differences in TCR repertoire diversity, complementarity-

determining region 3 (CDR3) length, amino acid composition, or network structure. No TCR

sequences were exclusive to either sex.

However, female effector T cells exhibited a significantly higher prevalence of TCRs specific

to self-antigens implicated in autoimmunity compared to males, while female Tregs showed

a reduced frequency of such TCRs. These differences were not observed for TCRs targeting

self-antigens unrelated to autoimmunity or antigens associated with cancer or viruses.

Our findings identify a sex-specific imbalance in thymic selection of TCRs with autoimmunity-

associated specificities, providing mechanistic insight into the increased susceptibility of

women to autoimmune diseases.

Introduction

Several studies have highlighted a sex imbalance in diseases involving the immune system. This is particularly evident in the case of autoimmune diseases (AIDs), where 80% of those affected individuals are women, and the severity of the condition varies between the sexes (1–3). In addition, men are more severely affected by infectious diseases such as COVID-19 and tuberculosis (4, 5). The response to preventive immunotherapies such as vaccination, and to curative treatments, like immune checkpoint inhibitors or anti-TNF alpha, is also sex dependent (6). These observations indicate the presence of intrinsic biological differences in immune responses between males and females.

A variety of factors have been identified that could help explain the observed sex-based differences. Among these, sex hormones have been shown to influence immune responses by acting on immune cells that express specific receptors to these hormones (7–9). Sex hormones could directly influence the adaptive immune response at the level of thymocyte differentiation and selection by acting on the expression of Autoimmune Regulator (AIRE) in thymic epithelial cells. AIRE is responsible for the thymic expression of otherwise tissue-specific antigens, contributing to both the negative selection of effector cells and the positive selection of Tregs that recognize such antigens (10). Testosterone increases AIRE expression in medullary thymic epithelial cells, whereas estradiol decreases it (11, 12). Furthermore, a recent study has shown that the transcriptomic profiles of these specialized antigen-presenting cells in the thymus differ between males and females (13). Thissuggests that there may be differences in the selection of the T cell receptor (TCR) repertoire between males and females.

Despite these findings, there is still limited knowledge about potential differences in the TCR

repertoire between the sexes. The generation of the TCR is a key process in T-cell

development that takes place in the thymus. The TCR consists of two chains, alpha (TRA) and

beta (TRB), each of which resulting from a random genetic rearrangement involving the V

(variable), D (diversity) [for the TRB], and J (joining) genes. The combination of these gene

segments is accompanied by insertions and deletions, creating a highly diverse region, the

Complementarity Determining Region 3 (CDR3) (14). It is this region of the TCR that

predominantly interacts with the peptide, a critical interaction for antigen recognition and

subsequent T cell activation. The ability to specifically recognize diverse antigens via the CDR3

is central to the efficiency and specificity of the adaptive immune response.

TCRs are randomly generated, which means that while some may be useful, others may be

harmful by recognizing self-antigens and causing autoimmunity. The current understanding is

that during thymocyte development, selection is based on the avidity of their TCRs for

antigens presented by cortical and medullary thymic epithelial cells. Thymocytes expressing

TCRs with insufficient or excessive avidity are eliminated (negative selection), while those

with an intermediate avidity are positively selected and mature into single-positive cells,

either CD4+ or CD8+. Thymocytes with the highest of these intermediate affinities develop in

Tregs. Given AIRE's pivotal function in these processes, sex differences in AIRE expression may

significantly influence thymic selection.

The TCR repertoire of female and male peripheral T cells have shown that their diversity,

particularly that of the TRB repertoire, is influenced not only by the age of the individuals but

also by their sex (15–18). However, another study showed no association between the sex of

individuals and the diversity of their TCR repertoire (19). Nevertheless, the study of peripheral

blood TCRs reflects that of an immune system that has undergone through multiple immune

responses. For example, observed differencesbetween males and females in their responses

to self-antigens associated with autoimmunity could indicate either the cause (the TCRs drive

the disease) or the consequence (the TCRs are expanded because of the disease).

To assess whether there are significant differences in the generation and thymic selection of

the TCR repertoire between males and females, we therefore focused on analyzing the TCR

repertoires of thymocytes. These cells represent the latest stage in T cell development and

have not yet been involved in immune responses. Their repertoires thus represent the

mechanisms of TCR generation for the analysis of CD4/CD8 double positive (DP) cells, and of

TCR selection for CD4 and CD8 simple positive (SP) T cells. We generated a unique data set

comprising the sequences of the TRA and TRB chains of DP cells, CD8 and CD4 SP effector

cells, and CD4 SP Tregs from organ donors, both infants and adults. The analysis of TRA and

TRB repertoires using state-of-the-art strategies revealed no sex differences in repertoire

generation (DP cell repertoire), but an enrichment in females for CD8 SP T cells bearing CDR3

associated with specificities related to autoimmunity.

Results

We analyzed thymic TCRαβ repertoires from 22 human organ donors (11 females, 11 males,

aged between 73 days and 64 years) (**Figure 1, Supplementary Figure 1**). For each donor,

CD3+CD4+CD8+ double positive (DP), CD3+CD4-CD8+ CD8 single positive (SP), and

CD3+CD4+CD8- CD4 SP thymocytes were sorted by flow cytometry. In 16 donors, CD4 SP

cells were further subdivided into CD25- conventional CD4 Teff and CD25+ CD4 Treg

subsets, whereas in the remaining six donors, total CD4 SP cells were analyzed as Teff-

enriched populations. This resulted in 20 DP, 21 CD8 SP, 22 CD4 Teff SP and 14 CD4 Treg SP

samples (**Figure 1, Supplementary Figure 1**). Bulk TCR sequencing was then performed on

each cell subset, with the aim to characterizing the TRA and TRB repertoires. Rank–

frequency plots of clonotype abundances displayed comparable heavy-tailed distributions

across donors, without any obvious systematic differences between males and females

(**Supplementary Figure 2**).

Minimal sex-based differences in TCR V and J gene usage across thymic cell subsets.

The first level of diversity in generating TCRs is the usage of V and J genes (20). We analyzed

the usage of TRA and TRB V and J genes and their combinations in each thymocyte population

(**Figure 1**).

Males and females had a relatively similar V gene usage, as evidenced by the lack of a clear

separation in the Principal Component Analysis (PCA), for both TRA and TRB, and across all

cell subtypes studied (**Figure 2A**). However, we observed some individual differences in V and

J gene usage between males and females, with some being specific to certain cell subtypes

(**Supplemental Figures 3-6**), and others observed across multiple cell subtypes (**Supplemental

Figures 3-6**). For example, TRBV6-5 was found to be more highly expressed in females

compared to males in DP cells (**Supplemental Figure 3C**) and CD8 SP cells (**Supplemental Figure 4C**).

In the following step, we compared the usage of TRAV-TRAJ and TRBV-TRBJ gene combinations across all individuals using the Jensen-Shannon Divergence (JSD) score (**Figure 2B**). The hierarchical clustering showed no clear separation between male and female individuals for any of the chain or cell subtype, indicating no significant differences in the gene combination usage between males and females (**Figure 2B**). A similar observation was made when directly analyzing the frequency of these V-J gene combinations (**Supplemental Figures 3E-6E, 3F-6F**).

Comparable TCR repertoire diversity between males and females

A high diversity of the TCR repertoire is essential for generating a broad potential reactivity against a variety of antigens. We thus compared the TCR repertoire diversity between males and females. To achieve this objective, we analyzed Rényi diversity curves (from zero to infinite α parameters) that measure various aspects of TCR repertoire diversity. At different α values, the Rényi index places more or less weight on the contribution of the most frequent clonotypes. Higher α values highlight the most prevalent clonotypes, thereby revealing the effect of expanded clones on overall diversity.

Rényi curves showed comparable diversities between sexes in DP and CD4 Teff cells, for both TRA and TRB, and a difference in CD8 and CD4 Treg cells (**Supplemental Figure 7**). In CD8 cells, the curves begin to diverge at low α values (**Supplemental Figure 7**), with an earlier inflection in males, indicating a less balanced distribution of TCR clones. This suggests that in males, fewer clones dominate the repertoire at lower diversity thresholds compared to females. In Tregs, while the inflection point is similar, the difference seems to be characterized by higher

162 diversity and richness in females compared to males (**Supplemental Figure 7**). However, no

163 significant differences were observed for the Shannon diversity index (α = 1), the Simpson

164 index (α = 2), and the Berger-Parker index (α = infinite), for both chains and all cell subtypes

165 (**Figure 3B**). These results indicate that the diversity of thymic TCR repertoire is similar

166 between males and females.

Comparable CDR3aa length distribution in male and female TCR repertoires

Most of the repertoire diversity is generated by the random addition of nucleotides within

the VDJ junctions, resulting in the hypervariable CDR3 region that interacts with the peptide-

MHC complex, and is mostly responsible for the TCR specificity. The CDR3 length has been

shown to directly influence the TCR's ability to interact with the peptide (21). We thus studied

the distribution of CDR3 aa (CDR3aa) sequence lengths between males and females to identify

potential TCR generation and/or selection biases (**Figure 1**).

Our results showed that males and females have a comparable distribution of CDR3aa lengths

for both TRA (**Supplemental Figure 8**) and TRB (**Figure 4A**). CDR3aa sequences in TRA were

shorter than those in TRB, as previously described in peripheral blood T cells (22, 23).

Although there were subtle differences in certain CDR3aa lengths between males and

females, these variations were minimal, involving sequences that represent less than 2% of

the total TCR repertoire (**Figure 4A** and **Supplemental Figure 8**). Altogether, the overall

distribution of CDR3aa length is comparable between males and females.

Subtle differences in the composition of CDR3aa in the thymic TCR repertoires of males and females.

183 We analyzed the aa usage within the FG loop of the CDR3, which is in contact with the peptide-

184 MHC complex (24). The classification of aa was determined based on their hydropathy

properties, categorized as follows: hydrophilic, hydrophobic, or neutral (25). We observed

significant differences in the usage of some aa between males and females for both TRA

(**Supplemental Figure 9**) and TRB (**Figure 4B**). In DP cells, these differences preferentially

involved hydrophobic aa: females exhibited higher alanine (A) usage in TRA and lower in TRB

compared to males (**Figure 4B** and **Supplemental Figure 9**). Additionally, females showed

increased usage of phenylalanine (F) and isoleucine (I) in TRB (**Figure 4B**). Furthermore, other

aa with varying hydropathy properties were differentially used between males and females

in CD4 Teff and Treg populations (**Figure 4B** and **Supplemental Figure 9**).

We then examined the usage of aa specifically at positions p109 and p110 of the TRB CDR3aa

region, as hydrophobic aa at these two positions have been associated with a stronger

recognition of self-antigens (26–28). For each individual, we calculated the proportion of

unique TRB CDR3aa carrying a hydrophobic aa (excluding alanine, which is only weakly

hydrophobic) at these two positions. This positional analysis revealed a significant increase in

the usage of hydrophobic amino acid at position p109 in female CD8 cells compared with

males, with similar trends observed at p109 in DP and CD4 Teff subsets and at p110 in CD8

cells (**Figure 4C**). Taken together, these positional patterns suggest a subtle sex-biased

enrichment of hydrophobic amino acids at key TRB CDR3 positions that could be compatible

with a slightly increased potential for self-reactivity and cross-reactivity in female CD8 T cells

(28).

Slight sex-based variations in thymic TCR generation probability distribution.

Another approach to evaluating biases in the generation and selection of the TCR repertoire

is to compare the probability of generation (Pgen) of CDR3 nucleotide sequences (**Figure 1**).

Pgen represents the likelihood of a sequence being produced, according to models of V(D)J

recombination (14, 29). For DP and CD8 cells only and for each TCR chain, a sequence

generation model was created using 100,000 non-productive nucleotide sequences. Using

this model, the Pgen of all productive TCR sequences composing each repertoire was

calculated.

When comparing the distribution of Pgen between males and females, we observe a very

slight difference of distribution, characterized by a nearly null Kolmogorov-Smirnov score (D_{K-S}

< 0.05) (**Figure 5**). This difference is more pronounced in the TRB compared to the TRA chain

(**Figure 5**). As this minor discrepancy might be indicative of variations among individuals

irrespective of their sex, we have generated 20,000 permuted mixed-sex groups utilizing our

dataset. Our findings revealed a significant distribution difference of Pgen sequences

between these mixed groups population (DP – TRA: $D_{K-S} = 0.0183 \pm 0.0145$ ($\mu \pm$ sd), DP –

TRB: $D_{K-S} = 0.0190 \pm 0.0161$, CD8 – TRA: $D_{K-S} = 0.0227 \pm 0.0153$, and CD8 – TRB: $D_{K-S} = 0.0259$

± 0.0140). However, the observed difference in the Pgen distribution between males and

females is higher than that between the mixed groups in the DP TRB.

Similar network structure of thymic TCR repertoires in males and females

Next, we evaluated whether thymic selection in males and females is directed towards similar

or distinct sequences. For each TCR repertoire, CDR3aa sequences are connected if they differ

by one aa, i.e. have a Levenshtein distance of =1 (**Figure 1**). Such linked sequences have a high

probability of sharing the same specificities (30). A random downsampling to the smallest

sample was performed to compare samples of similar sizes. This process was repeated 100

times, and the results are detailed in Supplemental Table 1. The shapes of the network are

depicted in **Figure 6A**.

To analyze these clusterings with greater precision, we used various metrics. When comparing

the number of clustered sequences in each individual network between males and females,

no significant differences were observed for any cell type in either chain (**Figure 6B**). Similarly,

the degree of sequence similarity, as indicated by the density of edges between sequences

within an individual's TCR repertoire, was comparable between males and females (**Figure**

6C). These findings indicate that there are no significant sex-specific differences in the

similarity of CDR3aa sequences, suggesting that males and females produce CDR3aa

sequences with the same degree of variability within their TCR repertoires.

Absence of sex-specific CDR3aa motifs in thymic TCR repertoires of males and
females

We then sought to determine whether we could detect CDR3aa sequence motifs that were

more prevalent in one group compared to the other, in order to evaluate whether the

selection of CDR3aa was biased towards particular sequences differently between males and

females (**Figure 1**). To identify sequences with shared characteristics that might indicate a

probable common specificity among individuals of the same group, the analysis focused

exclusively on CDR3aa sequences of the TRB, as this region has been shown to have a greater

influence on TCR specificity (31, 32). Local motifs were defined using Gliph2 (33), which are

a sequences of three to five aa, as well as global motifs, which are sequences of more than

three aa where, at one position, an aa can be substituted by another aa if it has a positive

BLOSUM62 score (34).

We identified several hundred motifs that were differentially expressed between males and

females for each cell subpopulation. A greater number of motifs were specific to females,

ranging from 426 motifs for DP to 278 motifs for CD4 Tregs, compared to males, ranging from

328 motifs for CD4 Teffs to 34 motifs for CD4 Tregs (**Figure 7A**). The identified motifs are

mainly local motifs and largely confined to their respective sex group, indicating that motifs

identified in females are predominantly found in females, and vice versa (**Supplemental**

Figures 10-13).

As illustrated in **Figure 7B**, we observed minimal motif overlap between different cell

subtypes. Only two female associated motifs overlapped: one between CD4 Teffs and CD4

Tregs, and another between CD4 Teffs and CD8 cells (**Figure 7B**). In contrast, no overlap was

found between male-associated motifs (**Figure 7B**). Furthermore, it was observed that there

was an overlap between one motif associated with female CD8 cells and a motif associated

with male CD4 Teff cells (**Figure 7B**).

We then proceeded to evaluate the usage of the differentially expressed motifs in external

datasets. There is no publicly available thymic TCR dataset that reports repertoires according

to cell subsets. We thus tested these motifs on TCRs from pediatric bulk thymocytes, which

contains TCR data from children aged between two and eight months, with a male-to-female

ratio of 5:3 (35, 36). We calculated the usage of these differentially expressed motifs in each

sample of this dataset. We were unable to separate individuals by sex using these motif

usages (**Figure 7C**).

We then evaluated these motifs using a peripheral blood TCR dataset from healthy

individuals, where CD8, CD4 Teff, and CD4 Treg cells had been sorted. The usage of these

motifs could not distinguish males from females across all cell subsets (**Figure 7D** and

Supplemental Figure 14).

Sex-biased enrichment of TCRs associated with autoimmune diseases and
bacterial antigens.

We sought to identify the usage of TCRs with known specificity as represented in the IEDB,

Mc-PAS and VDJdb public databases (37–39). We compiled the TCRs from these three

databases, retaining only those with high sequence reliability scores (see Materials and

Methods). We focused on TRB sequences exclusively, due to their greater representation in

the databases (31, 40). We identified 55,368 unique TRB CDR3aa sequences with high

reliability and specificity assignment scores for at least one specificity.

We classified the TCR specificities based on the antigen they recognize: viral or bacterial

peptides, or human peptides overexpressed in cancers, autoimmune diseases (AIDs), or

neither of these diseases (**Supplemental Figure 15A**). We sought to identify the TRB CDR3aa

sequences in our thymocyte dataset. Depending on the individuals and cell subtypes, they

represented between 0.82% and 3.58% of the TRB repertoires.

We then compared the distribution of these CDR3aa between males and females for each cell

subtype. We observed a significant enrichment of unique TRB CDR3aa sequences specific to

self-antigens not associated with pathologies and those associated with cancers in our thymic

TCR dataset, for both females and males (**Supplemental Figure 15B**). This enrichment is

observed across all cell subtypes and is approximately 1.25 to 1.5 times higher than the

baseline distribution of specificities in the pooled database (**Supplemental Figure 15B**).

Strikingly, in the TCR repertoires with one specificity only, we observed a significantly higher

proportion of unique TRB CDR3aa sequences associated with AIDs in female CD8 SP cells

compared to male CD8 SP cells (**Figure 8A**). Furthermore, in the specific TCR repertoires only,

there was a trend towards a higher proportion of thymic sequences with specificity associated

to bacterial compounds in female CD8 SP cells compared to males, and an opposite trend was

observed for CD4 Tregs SP (**Figure 8A**). These differences observed in CD8 SP cells persisted

when examining the usage of these sequences (i.e. cumulative usage frequency), with

significantly higher usage of sequences with specificity associated to bacterial compounds and

those associated with AIDs in females compared to males (**Figure 8B**). Of particular note are

the donor-resolved analyses (**Supplementary Figures 15B** and **16**), in which each individual is

represented separately. These confirm that these patterns are not driven by a single donor.

Donor age had no significant effect on the sex-biased enrichment of self- and bacteria-specific

TRB CDR3 sequences in CD8 SP and CD4 Treg SP cells ($p \geq 0.52$), indicating that the observed

differences are age-independent (**Supplemental Figure 17**). In addition, we also investigated

for the presence of polyspecific TCRs, which are capable of recognizing multiple antigens from

different organisms (41–43). These sequences were found to be enriched across all cell

subtypes, in comparison to their representation in the reference database (**Supplemental**

Figure 15C). However, no significant differences were observed in the proportion or usage of

these polyspecific sequences between males and females. Interestingly, an inverse usage

trend was noted between CD8 and CD4 Treg SP cells: females exhibited higher usage of these

sequences in the CD8 SP repertoire but a lower usage in the CD4 Treg SP repertoire compared

to males (**Figure 8C-8D**).

In conclusion, our analysis of TCR specificity has identified sex-specific differences in the

thymic selection of CD8 effector T cells associated to autoimmune and bacterial antigens that

are overrepresented in female versus males.

Discussion

It is noteworthy that there is a paucity of research on potential differences in the TCR

repertoire between the sexes, despite the much higher prevalence of autoimmune diseases

in females, a topic that remains to be fully elucidated. To explore the hypothesis that the

generation and/or the selection of the TCR repertoires may be different between sexes

requires, it is necessary to study the TCR repertoire where they are formed and selected, in

the thymus. We thus generated a unique and valuable dataset comprising thymic samples

collected from deceased organ donors of various ages, and young children who have

undergone cardiac surgery. Firstly, we separated the various thymocytes populations in order

to study both TCR repertoire generation at the level of DP cells, and repertoire selection at

the level of CD8, CD4 Teff and Treg SP cells. We performed bulk sequencing of the TCRs from

these cells, as this is the only method that yet generates enough sequences for sensitive

comparison at this time. Single-cell sequencing is an analyzing technique that it is limited in

its scope. It is only capable of analyzing a few thousand cells and would not have enabled the

analyses that can be performed with hundreds of thousands of cells.

TCR generation

Our analyses initially revealed similarities rather than dissimilarities in TCR generation. Our

detailed analyses revealed no significant sex differences in relation to the majority of the

examined variables when comparing males and females DP cells' repertoires, except for

differences in the usage of small number of V and J genes from both TCR chains. However,

these minor discrepancies are likely attributable to random variation, considering the

extensive number of comparisons conducted. This finding is further supported by the

observation that there are no significant differences in the usage of TRBV-TRBJ and TRAV-

TRAJ gene associations between males and females. We also observed no differences in

diversity indexes in DP cells' repertories between males and females. Altogether, the overall

combinatorial diversity of the TCR repertoire remains comparable between sexes.

A detailed analysis of the CDR3aa region in DP cells showed that some hydrophobic aa in the

FG loop region are differently found for both TCR chains. However, no significant sex-based

differences were found in the usage of hydrophobic, hydrophilic, or neutral aa at the critical

p109 and p110 positions in TRB that have been described as influencing self-antigen

recognition due to the impact of hydrophobic aa on TCR self-reactivity (26–28). These findings

suggest that the subtle sex-based differences in the FG loop's aa usage are minor and do not

significantly impact the overall characteristics of the CDR3 region in a sex-dependent manner.

The Pgen analysis of DP cells revealed only minimal differences in the distribution of Pgen

between males and females, which were similar to those observed in control groups. This

indicates that the mechanisms governing TCR generation are largely consistent across sexes,

with only minor, likely random, deviations.

Additionally, although we identified TRB CDR3aa sequence motifs that were differentially

expressed between males and females at the DP cell stage in our dataset, these motifs could

not reliably differentiate individuals by sex in external datasets. This indicates that they are

more dataset specific than sex-specific. This issue may also be explained by the different

characteristics of our data and that of the other dataset. For example, in the Arstila dataset,

the analyses were performed on unsorted cells from young subjects. Younger individuals tend

to have a more diverse and richer TCR repertoire, with greater repertoire sharing in the

periphery, which could obscure sex-specific patterns (16, 44). This suggests that these motifs

might be context-specific or influenced by other factors such as age, genetic background, or

environmental exposures. Moreover, we did not observe any sex-specific differences in CDR3

TRB specificity usage in DP cells' repertoires. In summary, our results from DP cells analyses

could not identify any relevant sex differences in TCR generation.

TCR selection

We identified three genes that show differential usage between males and females. Whilst it

is possible that some of these differences are merely random, resulting from multiple

comparisons, the preferential use of TRBV6-5 in females has previously been observed in the

peripheral TCR repertoire (45). Notably, this gene is overrepresented in female lupus patients

compared to their controls (51,52). Further research is needed to validate this observation and then elucidate its significance.

Our most striking observation relates to significant sex-specific differences in inferred specificities of CD8 T cells, with a biased CD8 selection process in the thymus favoring sequences associated with autoimmune diseases and bacterial antigens in females. The fact that this bias does not affect DP cells indicates that it is related to TCR selection rather than generation. Its relevance is strongly supported by the fact that (i) there is no such bias for other self-antigens that are not associated with autoimmunity and (ii) the bias affects both the effectors and regulators of autoimmunity with a directionality compatible with the increased prevalence of autoimmune disease in females. TCRs associated with autoimmunity related self-antigens have been observed to beincreased in CD8 effector T cells, while decreased levels have been seen in the CD4 Tregs. It should be noted that this bias also affects some bacterial antigens, though not viral antigens, are also affected by this bias. This phenomenom could potentially be explained by the fact that bacterial antigens express numerous mimotopes of self-antigens, and are known to contribute to the shaping of the TCR repertoire. The observed bias towards bacterial antigens in female CD8 T cells could likewise provide more effector cells specific for bacterial antigens that mimic self-antigens, contributing to autoimmune disease development by molecular mimicry (48). In this line, antigens linked to celiac disease (CeD) and type 1 diabetes (T1D) are highly represented in the database and these two AIDs have been linked to bacterial components that may mimic self-antigens (49–53). As is well-established that the gut microbiota influences thymic development of T cells (54). Sexual hormones could influence this biased selection by affecting the composition of the gut microbiota (55) and/or by affecting the selection of the TCRs specific for these antigens. In addition to these specific patterns, we observed a

significant increase in the usage of hydrophobic amino acids at the central p109 position of

the TRB CDR3aa in female CD8 thymocytes, with similar trend at p110. Hydrophobic residues

at these positions have been implicated in enhanced self-reactivity and cross-reactivity of CD8

T cells (26–28), suggesting that female CD8 thymocytes may be slightly enriched for TCRs with

intrinsically higher potential for recognizing self-like or cross-reactive peptide MHC

complexes. It is noteworthy that the positional hydrophobicity of CD8 T cells CDR3s is

consistent with the over-representation of sequences annotated as recognizing autoimmune-

associated self-antigens, pointing to a more autoreactive TCR repertoire.

Finally, we have recently described polyspecific TCRs that can respond to multiple unrelated

viral antigens and hypothesized their possible involvement in autoimmunity (43). We defined

a CDR3 as being polyspecific if the TCRs containing it have been experimentally associated

with epitopes from at least two different antigen origins. We observed a higher usage of

polyspecific sequences in the CD8 SP repertoire of females, although this was not statistically

significant. In contrast, the frequency of polyspecific TCR sequence usage was lower in female

CD4 Tregs SP cells. This could limit CD4 Treg antigen coverage and be detrimental to prevent

autoimmunity. However, little is known about polyspecific TCRs and these observations

warrant further investigation.

These inferences rely on a harmonized specificity compendium combining McPAS-TCR, IEDB

and VDJdb, which provide complementary and only partially overlapping landscapes

(pathology-enriched versus predominantly viral specificities). Although there is limited

numerical overlap between our thymic repertoires and annotated sequences is limited, this

is to be expected given that public databases currently capture only a small fraction of the

potential TCR specificity space. In this context, the sex-biased enrichment of sequences

annotated as autoimmune- and bacteria-associated should be viewed as a robust signal

emerging from a very sparse annotation space. However, it should still be interpreted as

hypothesis-generating rather than exhaustive.

To further evaluate the robustness of our findings, we performed additional analyses to

assess whether donor age could confound the observed sex-specific differences. A potential

limitation of our study is the relatively wide age range of donors. However, the absence of

age-related clustering in repertoire features (data not shown), and the lack of age impact on

CD8 and CD4 Treg bacterial-specific TRB usage and CD8 and CD4 Treg self-associated to

autoimmune disease-specific TRB usage patterns (Supplemental Figure 13), argue against age

as a confounding factor. These results support the hypothesis of an intrinsic sex bias in thymic

TCR selection rather than an age-related effect.

While our findings provide valuable insights into sex-specific differences in thymic selection

of TCR repertoires, our TCR specificity analyses were limited. We can confirm that our study

focused exclusively on the TRB CDR3aa region of the TCR, which is widely acknowledge as the

most critical element in defining TCR specificity (31, 32) while both the TRA and the TRB chains

contribute to TCR specificity (31, 32). This limitation is intrinsic to the limited number of TRA

with assigned specificity in databases and to the fact that, given the nature of our studies,

there is a need to study large repertoires, which is currently not feasible by single cell TCR

sequencing. However, as methods for inferring specificity continue to evolve, they will likely

allow more refined analyses of our dataset in the future (56). Furthermore, the continuous

development of the TCR database should enhance these analyses.

Despite the relatively small size of our cohort, which was outbred, we believe that this design

is essential for capturing sex-associated differences. Inbred models would reduce inter-

individual variability, but they would not reflect the complexity of human thymic selection,

nor the interplay between sex and genetic diversity. Despite the modest sample size, the

consistency of the observed patterns across the analytical approaches supports the

robustness of our conclusions.

Altogether, our analyses of TCR specificity identified a biased selection in females towards

sequences associated with autoimmune diseases, potentially underlying the higher

prevalence of autoimmune diseases in females. Future research should focus on validating

these observations in relevant independent datasets, which are currently lacking.

Methods

Samples

Thymus samples were obtained from twenty-two organ donors aged between 73 days and 64

years, without any specific pathologies (**Figure 1**). Adult samples were collected post-surgery

at the Cardiac Surgery Departments of many hospitals in France, following approval by the

Agence de Biomédecine and the Ministry of Research (#PFS14-009). Pediatric samples were

obtained following total thymectomy during cardiac surgeries at Necker Hospital. The ratio of

male to female across donors is maintained at 1:1, although it varies between 1.1:1 and 1.33:1

depending on the cell subtypes analyzed. The age distributions of male and female donors

were compared using two-sample Kolmogorov–Smirnov tests, with no significant differences

observed across thymic subsets ($p > 0.46$), indicating no age-related confounding

(**Supplementary Figure 1**).

Thymocyte isolation and RNA extraction

Thymic cell suspensions were prepared through automated tissue dissociation using

gentleMACS dissociators (Miltenyi®) or by mechanical disruption, followed by filtration

through a 70-μm nylon mesh. The cells were stained with anti-CD3 (AF700), anti-CD4 (APC),

anti-CD8 (FITC), and for some samples, anti-CD25 (PE) antibodies. Cell sorting was performed

by fluorescent activated cell sorting (FACS) on a Becton Dickinson FACSAriaII, achieving a

purity of >85%. The sorted populations included DP CD3+ (CD3+CD4+CD8+), SP CD8+

(CD3+CD4-CD8+), and SP CD4+ (CD3+CD4+CD8-). A further separation was made into SP CD4

Teff (CD3+CD4+CD8-CD25-) and SP CD4 Treg (CD3+CD4+CD8-CD25+) for sixteen of the

samples (**Figure 1**). For the sorted SP CD4, the Treg percentage varied between 5.8% and 16%

across samples, with an average of 9.63%. Six samples of SP CD4 samples were not sorted into

Teffs and Tregs, we thus categorized them as SP CD4 Teffs, assuming a purity of Teffs within

the total SP CD4 of around 90%. Please refer to the detailed sample metadata in

Supplementary Table 2 of the supplementary material. This includes donor IDs and the

number of sorted cells per thymic subset for each donor.

The RNA was isolated using the RNAqueous extraction kit (Invitrogen®), in accordance with

the manufacturer's protocol. The RNA samples were quantified and the integrity of each

sample was determined using a Nanodrop (Thermo Fisher) or a Tapestation 4200 (Agilent,

RNA ScreenTape).

TCR library preparation and sequencing

TCR library preparation and sequencing were performed as described by Barennes et al. (57).

Briefly, RNA was processed using the SMARTer Human TCR a/b profiling v1 kit (Takarabio), in

accordance with the manufacturer's protocol. Amplicons were then purified using AMPure

XP beads (Beckman Coulter) and their quantity was determined. The integrity of the

amplicons was then assessed using the 2100 Bioanalyzer System (Agilent, DNA 1000 kit) or

Tapestation 4200 (Agilent, D1000 screentape). Bulk Next-Generation Sequencing was

performed using either a Hiseq 2500 (Illumina) with the SR-300 protocol plus 10% PhiX on the

LIGAN-PM Genomics platform (Lille, France), or a Novaseq 6000 (Illumina) with the PE-250

protocol plus 10% PhiX on the LIGAN-PM Genomics platform or ICM platform (Paris, France).

Data processing

The raw FASTQ/FASTA files were aligned to TRA and TRB using MiXCR (v3.0.13, RNA-seq

parameters), a software solution that is able to correct for both sequencing and PCR errors

(58). Samples with fewer than 1,000 CDR3aa sequences, an imbalanced TRA/TRB ratio (<1/9),

or a low sequence count relative to sorted cell counts (< 0.9 ratio) were removed. Sequences

with CDR3aa lengths outside 6-23 amino acids were also discarded. To standardize clonotype

counts and the number of unique clonotypes across sequencing techniques, the initial counts

for each clonotype were reduced by 1 to exclude those with counts of 0 (singletons).

Furthermore, due to the high similarity between certain TRBV sequences, which has the

potential to influence the analysis depending on the sequencing method used, TRBV06-2 and

TRBV06-3 were merged as TRBV06-2/3, and TRBV12-3 and TRBV12-4 were combined as

TRBV12-3/4 (59).Please refer to the summary table provided in the supplementary material

(**Supplementary Table 2**) for a comprehensive overview of the total number of TCR sequences

retained after preprocessing, as well as the corresponding number of unique clonotypes, for

each donor, chain and thymic subset.

Gene usage analysis

For both TRA and TRB chains across all cell subtypes, we analyzed the V and J gene usage and

those of the VJ gene associations (**Figure 1**).

The usage of gene is calculated as follows: $\frac{\Sigma i \epsilon G^{ci}}{\Sigma j \epsilon S^{cj}}$

Where:

· $\Sigma i \epsilon G^{ci}$ represents the total number of counts for clonotypes associated with the

specified gene or gene association.

· $\Sigma j \epsilon S^{cj}$ represents the total sum of the counts of all clonotypes in the sample.

In this notation, G is the set of clonotypes using the interested gene, S is the set of all

clonotypes in the sample, and c_i and c_j are the counts of the clonotypes i and j, respectively.

The usage of the V gene and the VJ gene combinations was investigated through dimensional

reduction using Principal Component Analysis (PCA) (MASS package v7.3-60, factoMineR

package v2.8, factoextra package v1.0.7). Confidence ellipses were represented with a 5%

alpha risk. Furthermore, for VJ gene combinations usage, a heatmap was constructed using

the Jensen-Shannon Divergence (JSD) distance to measure the distance between samples

based on their gene usage distribution (complexHeatmap package v2.16.0, philentropy

package v0.7.0). Hierarchical clustering was performed using the Euclidean distance and the

complete linkage method.

Diversity analysis

For the TCR repertoire diversity analysis, to correct for sequencing depth bias and differences

in cellular richness between individuals, rarefactions were performed fifty times for each

sample, using X clonotypes, where X represents the effective diversity number (60).

The effective diversity was calculated as the exponential of the Shannon index (e^H) (61).

Rényi indices (11 indices from 0 to Infinity) were calculated for each rarefaction (62).

The median Shannon index, Simpson index and Berger-Parker indices were then compared

between the sexes.

Distribution of CDR3aa length analysis

534 The calculation of the usage of all CDR3 by the amino acid length as follows: $\dfrac{\Sigma i \epsilon L^{ci}}{\Sigma j \epsilon S^{cj}}$

535   Where

536   • $\Sigma i\epsilon L^{ci}$ represents the sum of the counts of the CDR3 characterized by the amino acid

537   length of interest.

• $\Sigma j\epsilon S^{cj}$ represents the total sum of counts of all CDR3s per sample.

In this notation, L is the set of CDR3s characterized by the amino acid length of interest, S is

the set of all CDR3s in the sample, and c_i and c_j are the counts of the CDR3s i and j, respectively.

Composition usage of amino acid within the CDR3 region

All CDR3aa sequence were aligned using the IMGT convention (positions p104-p118) (63).

Three complementary scales were used in the study. Firstly, sequence level aa composition.

Within the CDR3 loop region (IMGT positions p108 to p114), involved in MHC-peptide

complex interactions (63), aa usage at the sequence level was quantified for CDR3aa

sequences 9-21 aa long in both TRA and TRB chains. For each sequence, the percentage of a

given aa was calculated as follows:

(1) CDR3aa loop usage (%) = $\frac{n_{aa}}{L}$ x 100

Where n_{aa} is the number of occurences of the amino acid of interest within the CDR3 loop,

and L denotes the total length of the CDR3aa sequence. . Furthermore, amino acids were

categorized into three hydropathy classifications (neutral, hydrophobic, and hydrophilic)

using the IMGT classification that takes into account the Kyte-Doolittle hydropathy index

(25,64)

Secondly, the position-specific aa usage at p109 and p110. To assess positional composition,

for each individual, CDR3aa usage is examined at IMGT positions p109 and p110,

corresponding to p6-p7 positions described by Stadinsky et al. and Khosravi-Maharlooei et

al., which hydrophobic amino acids at these two central positions have been implicated in

modulating TCR self-reactivity and cross-reactivity (26–28). For a hydrophobic amino acid h

(excluding alanine, due to its weak hydrophobicity) at position p, usage was defined:

:

(2) CDR3aa usage in position (%) = $\frac{N_{(h,p)}}{n}$ x 100

Where n is the total number of unique CDR3 sequences and $N_{(h,p)}$ the number of unique CDR3

sequences carrying hydrophobic amino acid a at position p. This analysis quantifies amino

acid composition at fixed positions independently of clonotype frequency.

Probability of generation analysis

The probability of generation analysis of DP and CD8 cell subtypes consisted initially in the

creation of a generative model of V(D)J recombination using 400,000 randomly selected non-

productive sequences from all samples for both TRA and TRB chains, with the IGOR tool (65).

The generation probabilities (Pgen) for all sequences in the dataset were subsequently

calculated using the OLGA tool, based on the respective generative models for each cell

subtype and TCR chain (66). Comparisons of Pgen values at the nucleotide level between

males and females were performed. To ensure statistical robustness, control groups with

equivalent male and female sample numbers were created by permuting the samples into

two groups 20,000 times.

TCR network structure by sequence similarity

For each sample, 100 random sub-samplings of CDR3aa sequences were performed, using the

minimum CDR3aa count per cell subtype. Two CDR3aa were connected if their Levenshtein

distance was = 1 (**Figure 1**) (67). Network analysis focused on two metrics: the proportion of

connected sequences and network density, defined as the ratio of actual connections to all

possible connections. Median values from 100 subsampling iterations were then analyzed.

Levenshtein distances were computed with the stringdist package (v0.9.10), networks were

visualized using Cytoscape (v3.8.2), and network density was calculated with the igraph

package (v1.5.1).

TRB CDR3aa motif search

Two types of structural motifs were searched: local motifs (strict sequential motifs) and global

motifs (sequences with substitutions preserving a positive BLOSUM62 score) (Figure 1).

The selection of enriched motifs was carried out using the gliph2 function with the turboGliph

package (v0.99.2), with specific parameters defined as follows:

-　The input sequences were from the sex group of interest (male or female), with

reference sequences matched for TRBV gene usage.

-　It is imperative that CDR3aa sequences are longer than eight aa.

-　Local motif lengths were restricted to 3-5 aa.

-　Only motifs consisting of more than two unique CDR3aa were conserved.

-　Local motif significance of Fisher's test has not been boosted.

Additional filters were applied so that: (i) the inclusion of public CDR3aa sequences is a

prerequisite for a motif, with these sequences shared by at least two individuals, (ii) a

significant enrichment is required, as determined by Fisher's test with a p-value $p < 0.01$ and

(iii) a usage difference between groups of at least twofold is necessary, as evidenced by a

Wilcoxon test with a p-value $p < 0.05$.

Motif validation was performed on an openly published pediatric thymic dataset from the

Arstila team (35, 36) consisting in a bulk TCR dataset from infants aged between seven days

and eight months. Here, we removed one male subject "Thymus A", who was a twin of the

"Thymus B" and the "Thymus 1" neonate sample, in order to analyze height samples with a

male-to-female ratio of 5:3. Further validation was carried out on sorted peripheral blood TCR

repertoires (CD8, CD4 Teff and CD4 Treg) from seventy-five healthy volunteers aged from

eighteen to eighty-four years (male-to-female ratio of 1.03:1), whose libraries were

generated under the same conditions as those of our thymic dataset. They were recruited

within the Transimmunom observational trial (NCT02466217) and HEALTHIL-2

(NCT03837093) (68). PBMCs from healthy individuals were isolated using Ficoll density

gradient centrifugation and enriched for CD4+ T cells via EasySep™ Human CD4+ T Cell

magnetic beads (Stemcell). The samples were then separated into two groups: effector T cells

(CD4+ CD25-) and Tregs (CD4+ CD25+).This was performed using EasySep™ Human Pan-CD25

magnetic beads (Stemcell). Purity was assessed by flow cytometry based on the expression of

CD4 and FoxP3, with a purity threshold of >80%. From the frozen PBMC aliquots, CD8+ T cells

(CD3+ CD8+) were sorted using a FACS ARIA II cell sorter.

TRB CDR3aa specificity analysis

We have established a curated and harmonized TCR specificity database by integrating data

from three public TCR repositories: Mc-PAS, IEDB, and VDJdb (38, 39, 69) (data collected as

of October 2023) (Figure 1). The harmonization procedure was carried out in accordance with

the framework outlined in Jouannet et al. (70) and included :

-    Uniform formatting of TRA/TRB TCR sequences and associated metadata,

-    Consolidation of redundant entries (identical TRA/TRB TCR, epitope, organism,

PubMed ID, and cell subset),

-    Maintaining separate entries when source databases are incompatible,

Reliability levels for sequences and specificity assignments were quantified using the

Verified_score (VS) and Antigen-identification_score (AIS) as defined in (70). The VS, ranging

from 0 to 2, reflects the concordance between calculated and curated CDR3 boundaries

following the IEDB verification strategy (VS = 2: both TRA and TRB present and verified; VS =

1.1: only TRA verified; VS = 1.2: only TRB verified; VS = 0: no verified chain) and the AIS (0 to

5), which ranks the strength of antigen-identification methods (70).

In this study, we focused our analyses on high-reliability TRB sequences, defined as those with

a verified TRB CDR3aa (VS ≥ 1.2) and an AIS corresponding to in vitro T cell stimulation with a

pathogen, protein or peptide, or pMHC X-mer sorting (AIS > 3.2, excluding categories 4.1 and

4.2). This ensures that the specificity assignments relied on robust experimental evidence.

. Specificity groups were categorized according to antigen class, including viral, bacterial,

yeast, parasitic and self-antigens as along with antigens derived from wheat, plants or

animals. Self-antigens were then annotated into three categories: autoimmune-associated

antigens, cancer-associated antigens and self-antigens unrelated to disease. For

autoimmune-associated antigens, assignments were based on explicit annotations present in

McPAS-TCR, IEDB and VDJdb. These databases identify celiac disease (CeD) and type 1

diabetes (T1D) related antigens, which were retained as such in the harmonized dataset. For

cancer-associated antigens, the following classifications were used: (i) information present in

the original databases and (ii) cross-referencing with external cancer antigen resources,

including the Cancer Antigenic Peptide Database (de Duve Institute), the Human Protein Atlas

and the cancer database caAtlas (71). Please refer to **Supplementary Table 3**, which provides

a comprehensive list of all self-antigens present in the harmonized database. This table also

includes their associated disease category and their mapping into the specificity groups used

in the manuscript.

This harmonized database was used to identify exact matches with the CDR3aa sequences of

our TRB sequences. The enrichment of CDR3aa sequences associated with a given specificity

group in male and female thymic repertoires for each cell subtype compared to the specificity

distribution in the database was tested. Furthermore, we performed specificity group

distributions between male and female samples, as well as was the overall usage of these

sequences across entire TCR repertoires.

Additionally, our research concentrated on polyspecific sequences defined as CDR3aa TRB

sequences recognizing at least two distinct microbial species or belonging to multiple

specificity groups for non-microbial antigens. In this study, polyspecificity is therefore

considered at the level of broader antigenic categories rather than individual epitopes and

thus reflects functional cross-reactivity patterns aggregated across studies, instead of

implying molecular-level multi-epitope recognition. These polyspecific TRB CDR3aa

sequences underwent the same statistical analyses as monofunctional sequences.

Statistical analysis

For V-J gene usage, CDR3aa length usage, CDR3aa usage, network similarity and specificity

analyses, the comparison between males and females was statistically analysed using the

Wilcoxon test. The statistical tests were performed using the ggpubr library (v0.6.0) in R. The

number of asterisks denotes the significance of the results: one asterisk indicates a p-value p

≤ 0.05, two asterisk indicate a p-value p ≤ 0.01, and so on until four asterisks, denoting a p-

value p ≤ 0.0001. The comparisons of the Rényi curves and Pgen were compared using the

Kolmogorov-Smirnov test. The Wilcoxon test was used to test the comparisons of the

proportions of specific sequences in male and female subjects relative to those in the

specificity database.

Data availability

673   All raw sequencing data generated in this study are publicly available on NCBI under the

674   BioProject accession PRJNA1379632 (https://www.ncbi.nlm.nih.gov/sra/PRJNA1379632)

Author contributions

VQ and HV conducted all cell sorting. RNA extraction and TCR library preparation were performed by HV, VQ, LA, PB, NC, VD, JD, GF, KL and PS. HV, VQ, VM, LA, KL, ONT, MP, PS, AS, EMF and DK contributed valuable suggestions for experimental design and participated in discussions of the results. HV and CJ pooled and curated the reference specificity database. LA, CJ and VM contributed equally in this work. HV authored the manuscript, with VQ, VM, CA, JD, EMF and DK providing comments and critical reviews. DK conceptualized, supervised and provided funding for the study.

Acknowledgment

We would like to thank Bruno Gouritin for his assistance with cell sorting and to Marie Surroque for her support with RNA quantification.
During manuscript preparation, the authors used ChatGPT (OpenAI) solely to assist with language editing (syntax, grammar and phrasing). All ideas, analyses, interpretations and scientific reasoning are the authors' own.

Funding Statement

This work was supported by the TRiPoD European Research Council-Advanced EU (No. 322856), the LabEx Transimmunom (ANR-11-IDEX-0004-02) and the iMAP (ANR-16-RHUS-0001) grants to DK and the iReceptorPlus (Horizon 2020 No. 825821) and additional support from Sorbonne Université, INSERM as well as Institut Universitaire de France. KLG was supported by the European Research Area Network-Cardiovascular Diseases [ERANET-CVD JCT2018, AIR-MI Consortium (ANR-18-ECVD-0001)] grant to EMF.

References

1. Libert C, Dejager L, Pinheiro I. The X chromosome in immune functions: when a chromosome makes the difference. Nat Rev Immunol. 2010 Aug;10(8):594–604.

2. Golden LC, Voskuhl R. The Importance of Studying Sex Differences in Disease: The
Example of Multiple Sclerosis. J Neurosci Res. 2017 Jan 2;95(1–2):633–43.

3. Sokka T, Toloza S, Cutolo M, Kautiainen H, Makinen H, Gogus F, et al. Women, men,
and rheumatoid arthritis: analyses of disease activity, disease characteristics, and
treatments in the QUEST-RA Study. Arthritis Res Ther. 2009;11(1):R7.

4. Li Y, Jerkic M, Slutsky AS, Zhang H. Molecular mechanisms of sex bias differences in
COVID-19 mortality. Crit Care. 2020 July 9;24:405.

5. Hertz D, Schneider B. Sex differences in tuberculosis. Semin Immunopathol. 2019 Mar
1;41(2):225–37.

6. Klein SL, Morgan R. The impact of sex and gender on immunotherapy outcomes. Biol
Sex Differ. 2020 May 4;11(1):24.

7. Lee HJ, Chang C. Recent advances in androgen receptor action. Cell Mol Life Sci
CMLS. 2003 Aug 1;60(8):1613–22.

8. Chen P, Li B, Ou-Yang L. Role of estrogen receptors in health and disease. Front
Endocrinol. 2022 Aug 18;13:839005.

9. Scarpin KM, Graham JD, Mote PA, Clarke CL. Progesterone action in human tissues:
regulation by progesterone receptor (PR) isoform expression, nuclear positioning and
coregulator expression. Nucl Recept Signal. 2009 Dec 31;7:e009.

10. Klein L, Kyewski B, Allen PM, Hogquist KA. Positive and negative selection of the T
cell repertoire: what thymocytes see (and don't see). Nat Rev Immunol. 2014
June;14(6):377–91.

11. Zhu ML, Bakhru P, Conley B, Nelson JS, Free M, Martin A, et al. Sex bias in CNS
autoimmune disease mediated by androgen control of autoimmune regulator. Nat
Commun. 2016 Apr 13;7(1):11350.

12. Dragin N, Bismuth J, Cizeron-Clairac G, Biferi MG, Berthault C, Serraf A, et al.
Estrogen-mediated downregulation of AIRE influences sexual dimorphism in
autoimmune diseases. J Clin Invest. 126(4):1525–37.

13. Stankiewicz LN, Salim K, Flaschner EA, Wang YX, Edgar JM, Durland LJ, et al. Sex-
biased human thymic architecture guides T cell development through spatially defined
niches. Dev Cell. 2024 Oct 3;S1534-5807(24)00539-2.

14. Murugan A, Mora T, Walczak AM, Callan CG. Statistical inference of the generation
probability of T-cell receptors from sequence repertoires. Proc Natl Acad Sci. 2012 Oct
2;109(40):16161–6.

15. Yoshida K, Cologne JB, Cordova K, Misumi M, Yamaoka M, Kyoizumi S, et al. Aging-
related changes in human T-cell repertoire over 20 years delineated by deep
sequencing of peripheral T-cell receptors. Exp Gerontol. 2017 Oct 1;96:29–37.

16. Trofimov A, Brouillard P, Larouche JD, Séguin J, Laverdure JP, Brasey A, et al. Two
types of human TCR differentially regulate reactivity to self and non-self antigens.
iScience. 2022 Aug 17;25(9):104968.

17. Mika J, Yoshida K, Kusunoki Y, Candéias SM, Polanska J. Sex- and age-specific aspects
of human peripheral T-cell dynamics. Front Immunol. 2023 Oct 13;14:1224304.

18. Gong M, Li X, Zheng A, Xu H, Xie S, Yan R, et al. Age-related changes in the TRB and
IGH repertoires in healthy adult males and females. Immunol Lett. 2021 Dec 1;240:71–
6.

19. Qi Q, Liu Y, Cheng Y, Glanville J, Zhang D, Lee JY, et al. Diversity and clonal selection
in the human T-cell repertoire. Proc Natl Acad Sci U S A. 2014 Sept 9;111(36):13139–
44.

20. Davis MM, Bjorkman PJ. T-cell antigen receptor genes and T-cell recognition. Nature.
1988 Aug;334(6181):395–402.

21. Wang F, Ono T, Kalergis AM, Zhang W, DiLorenzo TP, Lim K, et al. On defining the
rules for interactions between the T cell receptor and its ligand: a critical role for a
specific amino acid residue of the T cell receptor beta chain. Proc Natl Acad Sci U S A.
1998 Apr 28;95(9):5217–22.

22. Yu K, Shi J, Lu D, Yang Q. Comparative analysis of CDR3 regions in paired human αβ
CD8 T cells. FEBS Open Bio. 2019 July 12;9(8):1450–9.

23. Carter JA, Preall JB, Grigaityte K, Goldfless SJ, Jeffery E, Briggs AW, et al. Single T
Cell Sequencing Demonstrates the Functional Role of αβ TCR Pairing in Cell Lineage
and Antigen Specificity. Front Immunol. 2019;10:1516.

24. Garcia KC, Degano M, Pease LR, Huang M, Peterson PA, Teyton L, et al. Structural
basis of plasticity in T cell receptor recognition of a self peptide-MHC antigen. Science.
1998 Feb 20;**279**(5354):1166–72.

25. Pommié C, Levadoux S, Sabatier R, Lefranc G, Lefranc MP. IMGT standardized criteria
for statistical analysis of immunoglobulin V-REGION amino acid properties. J Mol
Recognit. 2004;17(1):17–32.

26. Khosravi-Maharlooei M, Obradovic A, Misra A, Motwani K, Holzl M, Seay HR, et al.
Crossreactive public TCR sequences undergo positive selection in the human thymic
repertoire. J Clin Invest. 2019 Mar 28;129(6):2446–62.

27. Lu J, Van Laethem F, Bhattacharya A, Craveiro M, Saba I, Chu J, et al. Molecular
constraints on CDR3 for thymic selection of MHC-restricted TCRs from a random pre-
selection repertoire. Nat Commun. 2019 Mar 4;10:1019.

28. Stadinski BD, Shekhar K, Gómez-Touriño I, Jung J, Sasaki K, Sewell AK, et al.
Hydrophobic CDR3 residues promote the development of self-reactive T cells. Nat
Immunol. 2016 Aug;17(8):946–55.

29. Elhanati Y, Murugan A, Callan CG, Mora T, Walczak AM. Quantifying selection in
immune receptor repertoires. Proc Natl Acad Sci U S A. 2014 July 8;111(27):9875–80.

30. Meysman P, De Neuter N, Gielis S, Bui Thi D, Ogunjimi B, Laukens K. On the viability
of unsupervised T-cell receptor sequence clustering for epitope preference.
Bioinformatics. 2019 May 1;35(9):1461–8.

31. Springer I, Tickotsky N, Louzoun Y. Contribution of T Cell Receptor Alpha and Beta
CDR3, MHC Typing, V and J Genes to Peptide Binding Prediction. Front Immunol.
2021 Apr 26;12:664514.

32. Korpela D, Jokinen E, Dumitrescu A, Huuhtanen J, Mustjoki S, Lähdesmäki H. EPIC-
TRACE: predicting TCR binding to unseen epitopes using attention and contextualized
embeddings. Bioinformatics. 2023 Dec 9;39(12):btad743.

33. Huang H, Wang C, Rubelt F, Scriba TJ, Davis MM. Analyzing the Mycobacterium
tuberculosis immune response by T-cell receptor clustering with GLIPH2 and genome-
wide antigen screening. Nat Biotechnol. 2020 Oct;38(10):1194–202.

34. Henikoff S, Henikoff JG. Amino acid substitution matrices from protein blocks. Proc
Natl Acad Sci U S A. 1992 Nov 15;89(22):10915–9.

35. Heikkilä N, Kleino I, Vanhanen R, Yohannes DA, Mattila IP, Saramäki J, et al.
Characterization of human T cell receptor repertoire data in eight thymus samples and
four related blood samples. Data Brief. 2021 Apr 1;35:106751.

36. Mattila J, Sormunen S, Heikkilä N, Mattila IP, Saramäki J, Arstila TP. Analysis of
thymic generation of shared T-cell receptor α repertoire associated with recognition of
tumor antigens shows no preference for neoantigens over wild-type antigens. Cancer
Med. 2023 June;12(12):13486–96.

37. Vita R, Mahajan S, Overton JA, Dhanda SK, Martini S, Cantrell JR, et al. The Immune
Epitope Database (IEDB): 2018 update. Nucleic Acids Res. 2018 Oct 24;47(Database
issue):D339.

38. Tickotsky N, Sagiv T, Prilusky J, Shifrut E, Friedman N. McPAS-TCR: a manually
curated catalogue of pathology-associated T cell receptor sequences. Bioinforma Oxf
Engl. 2017 Sept 15;33(18):2924–9.

39. Shugay M, Bagaev DV, Zvyagin IV, Vroomans RM, Crawford JC, Dolton G, et al.
VDJdb: a curated database of T-cell receptor sequences with known antigen specificity.
Nucleic Acids Res. 2018 Jan 4;46(D1):D419–27.

40. Rosati E, Dowds CM, Liaskou E, Henriksen EKK, Karlsen TH, Franke A. Overview of
methodologies for T-cell receptor repertoire analysis. BMC Biotechnol. 2017 July
10;17(1):61.

41. Wooldridge L, Ekeruche-Makinde J, van den Berg HA, Skowera A, Miles JJ, Tan MP, et
al. A single autoimmune T cell receptor recognizes more than a million different
peptides. J Biol Chem. 2012 Jan 6;287(2):1168–77.

42. Hadrup SR, Bakker AH, Shu CJ, Andersen RS, van Veluw J, Hombrink P, et al. Parallel
detection of antigen-specific T-cell responses by multidimensional encoding of MHC
multimers. Nat Methods. 2009 July;6(7):520–6.

43. Quiniou V, Barennes P, Mhanna V, Stys P, Vantomme H, Zhou Z, et al. Human
thymopoiesis produces polyspecific CD8+ α/β T cells responding to multiple viral
antigens. eLife. 2023 Mar 30;12:e81274.

44. van de Sandt CE, Nguyen THO, Gherardin NA, Crawford JC, Samir J, Minervina AA, et
al. Newborn and child-like molecular signatures in older adults stem from TCR shifts
across human lifespan. Nat Immunol. 2023 Nov;24(11):1890–907.

45. Schneider-Hohendorf T, Görlich D, Savola P, Kelkka T, Mustjoki S, Gross CC, et al. Sex
bias in MHC I-associated shaping of the adaptive immune system. Proc Natl Acad Sci U
S A. 2018 Feb 27;115(9):2168–73.

46. Roca AM, Chobrutskiy BI, Callahan BM, Blanck G. T-cell receptor V and J usage paired
with specific HLA alleles associates with distinct cervical cancer survival rates. Hum
Immunol. 2019 Apr;80(4):237–42.

47. Hou X, Wei W, Zhang J, Liu Z, Wang G, Yang X, et al. Characterisation of T and B cell
receptor repertoire in patients with systemic lupus erythematosus. Clin Exp Rheumatol.
2023 Nov;41(11):2216–23.

48. Rojas M, Restrepo-Jiménez P, Monsalve DM, Pacheco Y, Acosta-Ampudia Y, Ramírez-
Santana C, et al. Molecular mimicry and autoimmunity. J Autoimmun. 2018 Dec
1;95:100–23.

49. Repac J, Božić B, Božić Nedeljković B. Microbes as triggers and boosters of Type 1
Diabetes - Mediation by molecular mimicry. Diabetes Res Clin Pract. 2023
Aug;202:110824.

50. Girdhar K, Huang Q, Chow IT, Vatanen T, Brady C, Raisingani A, et al. A gut microbial
peptide and molecular mimicry in the pathogenesis of type 1 diabetes. Proc Natl Acad
Sci U S A. 2022 Aug 2;119(31):e2120028119.

51. Petersen J, Ciacchi L, Tran MT, Loh KL, Kooy-Winkelaar Y, Croft NP, et al. T cell
receptor cross-reactivity between gliadin and bacterial peptides in celiac disease. Nat
Struct Mol Biol. 2020 Jan;27(1):49–61.

52. Vazquez DS, Schilbert HM, Dodero VI. Molecular and Structural Parallels between
Gluten Pathogenic Peptides and Bacterial-Derived Proteins by Bioinformatics Analysis.
Int J Mol Sci. 2021 Aug 27;22(17):9278.

53. Tai N, Peng J, Liu F, Gulden E, Hu Y, Zhang X, et al. Microbial antigen mimics activate
diabetogenic CD8 T cells in NOD mice. J Exp Med. 2016 Sept 19;213(10):2129–46.

54. Hebbandi Nanjundappa R, Sokke Umeshappa C, Geuking MB. The impact of the gut
microbiota on T cell ontogeny in the thymus. Cell Mol Life Sci. 2022 Apr;79(4):221.

55. Markle JGM, Frank DN, Mortin-Toth S, Robertson CE, Feazel LM, Rolle-Kampczyk U,
et al. Sex differences in the gut microbiome drive hormone-dependent regulation of
autoimmunity. Science. 2013 Mar 1;339(6123):1084–8.

56. Ghoreyshi Z, George J. Quantitative approaches for decoding the specificity of the
human T cell repertoire. Front Immunol. 2023 Sept 7;14.

57. Barennes P, Quiniou V, Shugay M, Egorov ES, Davydov AN, Chudakov DM, et al.
Benchmarking of T cell receptor repertoire profiling methods reveals large systematic
biases. Nat Biotechnol. 2021 Feb;39(2):236–45.

58. Bolotin DA, Poslavsky S, Mitrophanov I, Shugay M, Mamedov IZ, Putintseva EV, et al.
MiXCR: software for comprehensive adaptive immunity profiling. Nat Methods. 2015
May;12(5):380–1.

59. Omer A, Peres A, Rodriguez OL, Watson CT, Lees W, Polak P, et al. T cell receptor beta
germline variability is revealed by inference from repertoire data. Genome Med. 2022
Jan 7;14(1):2.

60. Chaara W, Gonzalez-Tort A, Florez LM, Klatzmann D, Mariotti-Ferrandiz E, Six A.
RepSeq Data Representativeness and Robustness Assessment by Shannon Entropy.
Front Immunol. 2018 May 15;9:1038.

61. Shannon CE. The mathematical theory of communication. 1963. MD Comput Comput
Med Pract. 1997;14(4):306–17.

62. Rényi A. On Measures of Entropy and Information. In: Proceedings of the Fourth
Berkeley Symposium on Mathematical Statistics and Probability, Volume 1:
Contributions to the Theory of Statistics. University of California Press; 1961. p. 547–
62.

63. Lefranc MP, Pommié C, Ruiz M, Giudicelli V, Foulquier E, Truong L, et al. IMGT
unique numbering for immunoglobulin and T cell receptor variable domains and Ig
superfamily V-like domains. Dev Comp Immunol. 2003 Jan;27(1):55–77.

64. Kyte J, Doolittle RF. A simple method for displaying the hydropathic character of a
protein. J Mol Biol. 1982 May 5;157(1):105–32.

65. Marcou Q, Mora T, Walczak AM. High-throughput immune repertoire analysis with
IGoR. Nat Commun. 2018 Feb 8;9(1):561.

66. Sethna Z, Elhanati Y, Callan CG, Walczak AM, Mora T. OLGA: fast computation of
generation probabilities of B- and T-cell receptor amino acid sequences and motifs.
Bioinforma Oxf Engl. 2019 Sept 1;35(17):2974–81.

67. Madi A, Poran A, Shifrut E, Reich-Zeliger S, Greenstein E, Zaretsky I, et al. T cell
receptor repertoires of mice and humans are clustered in similarity networks around
conserved public CDR3 sequences. eLife. 2017 July 21;6:e22057.

68. Lorenzon R, Mariotti-Ferrandiz E, Aheng C, Ribet C, Toumi F, Pitoiset F, et al. Clinical
and multi-omics cross-phenotyping of patients with autoimmune and autoinflammatory
diseases: the observational TRANSIMMUNOM protocol. BMJ Open. 2018 Aug
30;8(8):e021037.

69. Vita R, Mahajan S, Overton JA, Dhanda SK, Martini S, Cantrell JR, et al. The Immune
Epitope Database (IEDB): 2018 update. Nucleic Acids Res. 2019 Jan 8;47(D1):D339–
43.

70. Jouannet C, Vantomme H, Gouge KL, Klatzmann D, Mariotti-Ferrandiz E.
Benchmarking unsupervised methods for inferring TCR specificity. NAR Genomics
Bioinforma. 2025 Dec 1;7(4):lqaf150.

71. Yi X, Liao Y, Wen B, Li K, Dou Y, Savage SR, et al. caAtlas: An immunopeptidome
atlas of human cancer. iScience. 2021 Oct 22;24(10):103107.

Figure 1: Schematic overview of the generation of the thymic TCR dataset and the analytical pipeline. Partially created with Biorender.com

Top Panel: Generation of the thymic TCR dataset. From deceased human thymuses of males and females, we isolated key T-cell subtypes through cell sorting. These subtypes included double-positive (DP) cells (CD3+ CD4+ CD8+), single-positive (SP) CD8+ cells (CD3+ CD4- CD8+), SP CD4+ cells (CD3+ CD4+ CD8-) that were further separated into T effector (Teff, CD3+ CD4+ CD8- CD25-) and Treg (CD3+ CD4+ CD8- CD25+) cells. TCR libraries were generated from the RNA of each cell population using rapid amplification of cDNA ends by PCR (5'RACE PCR). Following sequencing, data preprocessing involved quality sequencing checks, contig alignment, and quality control. The final dataset comprised 20 DP samples (male-to-female ratio of 1:1), 21 SP CD8+ samples (1.1:1), 6 SP CD4+ samples (1:5), 16 SP CD4 Teff samples (1.67:1), and 14 SP CD4 Treg samples (1.33:1). Males are depicted in violet and females in orange.

Bottom Panel: Analytical pipeline. We compared the TCR repertoires of males and females across various dimensions. We evaluated general aspects of the TCR repertoire were evaluated, including diversity, gene usage, CDR3aa length distribution, and aa usage within the CDR3 region. Additionally, we analyzed the probability of sequence generation and the TCR repertoire structure based on CDR3aa sequence similarity. We identified differentially expressed TRB CDR3aa motifs between sexes and analyzed TRB CDR3aa sequence specificity.

Figure 2: Comparable overall TCR gene usage between males and females.

(A) Principal Component Analysis (PCA) derived from the distribution of TRAV (left) and TRBV (right) gene usage frequencies across sex

groups (males vs females), showing results for DP (N = 22), CD8 (N = 23), CD4 Teff (N = 24) and CD4 Treg (N = 14) cells (displayed from top

to bottom). Each point on the graph represents an individual. Ellipses indicate 95% confidence intervals. **(B)** Heatmap showing the Jensen-

Shannon Divergence (JSD) score between samples, derived from the distributional usage of TRAV-TRAJ (left) and TRBV-TRBJ (right) gene

associations in DP (N = 22), CD8 (N = 23), CD4 Teff (N = 24) and CD4 Treg (N = 14) cells (displayed from top to bottom). Hierarchical clustering

was performed using the Euclidean distance and the complete linkage method. Males are shown in violet; females in orange.

922
923 **Figure 3: Comparable thymic TCR repertoire diversity between males and females.**

924 Boxplots display Shannon **(A)**, Simpson **(B)**, and Berger-Parker **(C)** index values for TRA (left) and TRB (right), across thymic T cell subtypes,

925 displayed from top to bottom, in DP (N = 22), CD8 (N = 23), CD4 Teff (N = 24) and CD4 Treg (N = 14). Each point on the graph represents the

926 median value from 50 rarefactions per sample. Statistical analysis (Wilcoxon test) showed no significant sex bias in TCR repertoire diversity

927 ($p > 0.05$). Males are shown in violet; females in orange.

Figure 4: CDR3aa length and amino acid composition of TRB CDR3s in males and females.
(A) Distribution of TRB CDR3aa length usage in DP (N = 22), CD8 (N = 23), CD4 Teff (N = 24) and Treg (N = 14) SP cells (displayed from top to bottom). Stars indicate statistical differences between males and females based on the p-value of the Wilcoxon test (*: p < 0.05, **: p < 0.01). **(B)** The data on amino acid (aa) usage between males and females is presented as the log2 fold change of the median per-donor usage in females over males for each aa in the p108 to p114 CDR3aa region for TRB. A line at log2 fold change = 0 is indicative of the direction of the difference in usage frequency. The bars are colour-coded to the hydropathy class of the aa as defined by the Kyte-Doolittle-based IMGT classification (neutral aa by gray, hydrophilic aa by blue-green and hydrophobic aa by gold). Stars indicate statistical differences of usage between males and females based on the p-value of the Wilcoxon test (*: p < 0.05, **: p < 0.01). **(C)** Position-specific usage of hydrophobic aa (excluding alanine, due to its weak hydrophobicity) at IMGT positions p109 and p110 in TRB across thymic T cell subtypes, including DP (N = 22), CD8 (N = 23), CD4 Teff (N = 24) and CD4 Treg (N = 14). For each donor, the values represent the proportion of unique TRB CDR3aa sequences carrying a hydrophobic amino acid at the indicated position. Stars indicate significant sex differences based on the p-value of the Wilcoxon test (*: p < 0.05), with a significant increase in hydrophobic usage at p109 in female CD8 SP cells. Males are depicted in violet; females in orange.

942
Figure 5: Probabilities of generation of TCRs in males and females.

The figure shows the distribution of Pgen (probability of generation) sequences between males and females for TRA and TRB in DP and CD8

cells. A V(D)J recombination model was created using 400,000 non-productive random sequences derived from the nonproductive

sequences of all individuals for each chain, both for DP and CD8 cells (65). This model was then used to calculate the Pgen of sequences for

each individual (66). The overall distribution comparison between males and females was tested using the Kolmogorov-Smirnov test, with

the D value and associated p-value indicated. Males are depicted in violet and females in orange.

951
Figure 6: Comparable TCR repertoire network structure based on CDR3 amino acid sequence similarity between males and females.

For each sample, 100 random subsamplings were performed on the minimum number of CDR3aa per cell subtype. Two CDR3aa are linked

if they have a Levenshtein distance of one. **(A)** TRA Network structure of subsampled TCR repertoire of a male subject for DP, CD8, CD4 Teff

and CD4 Treg SP (from left to right). Each point on the graph represents a CDR3aa. **(B-C)** Comparison of the proportion of linked sequences

(B) and network density **(C)** between male and female samples, in DP (N = 22), CD8 (N = 23), CD4 Teff (N = 24) and CD4 Treg (N = 14). Each

point on the graph represents the median value from 100 subsampling iterations for each sample. Statistical analysis using the Wilcoxon

test revealed no significant sex differences for these two metrics (p > 0.05). Males are depicted in violet; females in orange.

Figure 7: Thymic TRB TCR sex-associated motifs.

Different structural motifs found differentially expressed between males and females in our dataset. We distinguish local motifs as distinct aa sequences and global motifs as motif regions with one variable aa position maintaining a BLOSUM62 score ≥ 0. **(A)** Number of male and female associated motifs by cell subset. **(B)** Euler diagram illustrating the distribution and overlap between all sex-associated motifs. The numbers indicate the number of motifs in overlap zones. **(C-D)** Validation of these sex-associated TRB CDR3aa motifs. The heatmap illustrates the usage of all the TRB CDR3aa motif in the external thymic pediatric dataset (35, 36) **(C)** and those of TRB CD8 in the peripheral dataset **(D)**. Sex and total CDR3aa number are depicted by sample. Males are depicted in violet and females in orange then local motifs in blue-green and global motifs in magenta.

968
969
Figure 8: Sex-biased enrichment of TCRs specific for known antigens.
From a pooled and curated database, an exact match with this database infers the specificity of TRB CDR3aa of our thymic dataset. Many
specificity groups are defined according to the nature of the antigen peptide targeted (bacteria, virus, autoimmune disease [AID], cancer
and self-peptide no associated to disease [human]). This analysis compares the distributions of the proportion of unique TRB CDR3aa
sequences with a specific specificity **(A)** and their usage **(B)** between females and males across cell subtypes, using the log2 fold change of
the median values (females over males), following each specificity group. These groups of specificity are additionally classified as
microorganism in the top section (bacteria in gold and virus in light blue) and self at the bottom (AID in magenta, cancer in red, human in
blue-green). Polyspecific CDR3aa are defined here as CDR3aa capable of recognizing multiple antigens from different organisms (for no self-
antigens) or from different specificity groups (e.g. categorized microorganisms, categorized self-antigens, allergens…). The proportion of
unique polyspecific CDR3aa in the specific repertoire **(C)** and their usage **(D)** is compared between males and females, in DP (N = 22), CD8
(N = 23), CD4 Teff (N = 24) and CD4 Treg (N = 14) cells. Stars indicate statistical differences between males and females based on the p-value
of the Wilcoxon test (*: p < 0.05, **: p < 0.01). Males are depicted in violet and females in orange.

Supplemental Figure 1: Age distribution of male and female donors is comparable across thymic subsets.

Age of individual donors plotted separately for females (orange) and males (violet) for each thymic cell subset: double-positive (DP) thymocytes, CD8 single-positive (CD8), CD4 effector single-positive (CD4 Teff) and CD4 regulatory T-cell single-positive (CD4 Treg). For each subset, a two-sample Kolmogorov–Smirnov (K-S) test was performed; the D statistic and corresponding p-value are displayed below each panel. In all cases, no significant difference was detected, indicating balanced age distributions between sexes for every subset.

Supplemental Figure 2: Rank–frequency distributions of thymic TRA and TRB clonotypes. Log–log rank–frequency plots of unique TRA (left) and TRB (right) clonotypes are shown for each donor and thymic subset (DP, CD8, CD4 Teff, CD4 Treg, from top to bottom). For each sample, clonotypes are ranked by decreasing abundance (x-axis, log scale), and their corresponding relative frequencies are plotted (y-axis, log scale). Curves display broadly comparable shapes across donors, with no obvious systematic differences between males and females, indicating similar clonotype abundance distributions and sampling depth across sexes.

Supplemental Figure 3: Some differential V and J gene usage in TCR repertoires of DP cells between males and females.

(A-D) Frequencies of TRAV **(A)**, TRAJ **(B)**, TRBV **(C)**, and TRBJ **(D)** gene usage between males and females. Red lines represent the mean for each sex group. Statistical comparisons between groups were performed using the Wilcoxon test. Stars indicate statistical differences between males and females based on the p-value of the test (*: $p < 0.05$, **: $p < 0.01$). **(E-F)** Principal Component Analysis (PCA) derived from the distribution of the frequency of usage of TRAV-TRAJ **(E)** and TRBV-TRBJ **(F)** gene associations across sex groups between males and females. Each point represents an individual. Ellipses show 95% confidence intervals. Males are depicted in violet and females in orange.

Supplemental Figure 4: Some differential V and J gene usage in TCR repertoires of CD8 cells between males and females.

(A-D) Frequencies of TRAV **(A)**, TRAJ **(B)**, TRBV **(C)**, and TRBJ **(D)** gene usage between males and females. Red lines represent the mean for each sex group. Statistical comparisons between groups were performed using the Wilcoxon test. Stars indicate statistical differences between males and females based on the p-value of the test (*: $p < 0.05$, **: $p < 0.01$). **(E-F)** Principal Component Analysis (PCA) derived from the distribution of the frequency usage of TRAV-TRAJ **(E)** and TRBV-TRBJ **(F)** gene associations across sex groups between males and females. Each point represents an individual. Ellipses show 95% confidence intervals. Males are depicted in violet and females in orange.

Sex ● Female ● Male

Supplemental Figure 5: Some differential V and J gene usage in TCR repertoires of CD4 Teff cells between males and females.

(A-D) Frequencies of TRAV **(A)**, TRAJ **(B)**, TRBV **(C)**, and TRBJ **(D)** gene usage between males and females. Red lines represent the mean for each sex group. Statistical comparisons between groups were performed using the Wilcoxon test. Stars indicate statistical differences between males and females based on the p-value of the test (*: $p < 0.05$, **: $p < 0.01$). **(E-F)** Principal Component Analysis (PCA) derived from the distribution of the frequency usage of TRAV-TRAJ **(E)** and TRBV-TRBJ **(F)** gene associations across sex groups between males and females. Each point represents an individual. Ellipses show 95% confidence intervals. Males are depicted in violet and females in orange.

Supplemental Figure 6: Some differential V and J gene usage in TCR repertoires of CD4 Treg cells between males and females.

(A-D) Frequencies of TRAV **(A)**, TRAJ **(B)**, TRBV **(C)**, and TRBJ **(D)** gene usage between males and females. Red lines represent the mean for each sex group. Statistical comparisons between groups were performed using the Wilcoxon test. Stars indicate statistical differences between males and females based on the p-value of the test (*: $p < 0.05$, **: $p < 0.01$). **(E-F)** Principal Component Analysis (PCA) derived from the distribution of the usage frequency of TRAV-TRAJ **(E)** and TRBV-TRBJ **(F)** gene associations across sex groups between males and females. Each point represents an individual. Ellipses show 95% confidence intervals. Males are depicted in violet and females in orange.

Supplemental Figure 7: Minimal differences in diversity profile of CD8 and CD4 Treg thymic TCR repertoire between males and females.

Diversity profile with Rényi diversity index values (α ranging from 0 to infinity) for all cell subsets, both TRA (left) and TRB (right). Clonotypes of each sample were rarefied fifty times to their effective diversity number [i.e. $e^{Shannon\ index}$] and the median of their Rényi values was used for analysis. Dotted lines and points show the mean value for each sex group, while the shaded area represents the standard deviation for each sex group. The overall shape of the curves was compared statistically using the Kolmogorov-Smirnov test, with the D value and associated p-value indicated.

Supplemental Figure 8: Comparable TRA CDR3aa length distribution between males and females.

Distribution of CDR3aa length usage for TRA in all cell subsets. Stars indicate statistical differences between males and females based on the p-value of the Wilcoxon test (*: $p < 0.05$, **: $p < 0.01$). Males are depicted in violet and females in orange.

Supplemental Figure 9: Some differences in TRA CDR3aa usage between males and females.

Usage of each aa in the p108 to p114 CDR3 region represented as the log2 fold change of the median usage of females over males in TRA for each cell subset. A line at log2 fold change = 0 indicates the direction of the usage difference. Bars are colored according to the hydropathy class of the amino acid (neutral in gray, hydrophilic in blue-green and hydrophobic in gold). Stars indicate statistical differences between males and females based on the p-value of the Wilcoxon test (*: $p < 0.05$, **: $p < 0.01$).

CDR3aa number

sex
age class
sequencing platform

sex
motif type

motif type
global
local

sex
Female
Male

age class
< 18
18~50
> 50

sequencing platform
HiSeq 2500
NovaSeq 6000

Usage frequency (%)
0   0.5   1

Supplemental Figure 10: DP TRB thymic sex-associated motifs in our dataset.

Different structural motifs in the CDR3 amino acid (CDR3aa) region found differentially expressed between males and females within our dataset. Local motifs refer to distinct amino acid sequences, whereas global motifs represent motif regions with a single variable amino acid position maintaining a BLOSUM62 score ≥ 0. The heatmap showcases the differential usage of TRB CDR3aa motifs between males and females in of DP cells. Hierarchical clustering reveals clear segregation of individuals by sex, with most CDR3aa motifs being almost exclusively expressed in one sex while absent in the other. Sample attributes are visualized as follows: (1) Sex: males in violet, females in orange; (2) Age class: children (<18 years) in green, young adults (18-50 years) in blue, older adults (>50 years) in red; (3) sequencing platform: HiSeq2500 samples in persimmon, NovaSeq 6000 samples in purple; (4) the total CDR3aa number. Motifs overexpressed in males are shown in violet, those overexpressed in females in orange; local motifs are indicated in blue-green and global motifs in magenta.

motif type
- global
- local

sex
- Female
- Male

age class
- < 18
- 18–50
- > 50

sequencing platform
- HiSeq 2500
- NovaSeq 6000

Usage frequency (%)
0   0.5   1

Supplemental Figure 11: CD8 TRB thymic sex-associated motifs in our dataset.

Different structural motifs in the CDR3 amino acid (CDR3aa) region found differentially expressed between males and females within our dataset. Local motifs refer to distinct amino acid sequences, whereas global motifs represent motif regions with a single variable amino acid position maintaining a BLOSUM62 score ≥ 0. The heatmap showcases the differential usage of TRB CDR3aa motifs between males and females in of CD8 cells. Hierarchical clustering reveals clear segregation of individuals by sex, with most CDR3aa motifs being almost exclusively expressed in one sex while absent in the other. Sample attributes are visualized as follows: (1) Sex: males in violet, females in orange; (2) Age class: children (<18 years) in green, young adults (18-50 years) in blue, older adults (>50 years) in red; (3) sequencing platform: HiSeq2500 samples in persimmon, NovaSeq 6000 samples in purple; (4) the total CDR3aa number. Motifs overexpressed in males are shown in violet, those overexpressed in females in orange; local motifs are indicated in blue-green and global motifs in magenta.

CDR3aa number

sex

age class

sequencing platform

motif type	sex	age class	sequencing platform	Usage frequency (%)
global	Female	< 18	HiSeq 2500	0 0.5 1
local	Male	18–50	NovaSeq 6000	
		> 50		

sex

motif type

3e+05
2e+05
1e+05
0

Supplemental Figure 12: CD4 Teff TRB thymic sex-associated motifs in our dataset.

Different structural motifs in the CDR3 amino acid (CDR3aa) region found differentially expressed between males and females within our dataset. Local motifs refer to distinct amino acid sequences, whereas global motifs represent motif regions with a single variable amino acid position maintaining a BLOSUM62 score ≥ 0. The heatmap showcases the differential usage of TRB CDR3aa motifs between males and females in of CD4 Teff cells. Hierarchical clustering reveals clear segregation of individuals by sex, with most CDR3aa motifs being almost exclusively expressed in one sex while absent in the other. Sample attributes are visualized as follows: (1) Sex: males in violet, females in orange; (2) Age class: children (<18 years) in green, young adults (18-50 years) in blue, older adults (>50 years) in red; (3) sequencing platform: HiSeq2500 samples in persimmon, NovaSeq 6000 samples in purple; (4) the total CDR3aa number. Motifs overexpressed in males are shown in violet, those overexpressed in females in orange; local motifs are indicated in blue-green and global motifs in magenta.

Supplemental Figure 13: CD4 Treg TRB thymic sex-associated motifs in our dataset.

Different structural motifs in the CDR3 amino acid (CDR3aa) region found differentially expressed between males and females within our dataset. Local motifs refer to distinct amino acid sequences, whereas global motifs represent motif regions with a single variable amino acid position maintaining a BLOSUM62 score ≥ 0. The heatmap showcases the differential usage of TRB CDR3aa motifs between males and females in of CD4 Treg cells. Hierarchical clustering reveals clear segregation of individuals by sex, with most CDR3aa motifs being almost exclusively expressed in one sex while absent in the other. Sample attributes are visualized as follows: (1) Sex: males in violet, females in orange; (2) Age class: children (<18 years) in green, young adults (18-50 years) in blue, older adults (>50 years) in red; (3) sequencing platform: HiSeq2500 samples in persimmon, NovaSeq 6000 samples in purple; (4) the total CDR3aa number. Motifs overexpressed in males are shown in violet, those overexpressed in females in orange; local motifs are indicated in blue-green and global motifs in magenta.

Supplemental Figure 14: TRB thymic sex associated motifs are not differentially expressed between males and females in peripheral TCR repertoire.

Heatmap representing TRB CDR3aa motif usage differentially expressed between males and females in CD4 Teff **(A)** and CD4 Treg **(B)** cells in a peripheral TCR repertoire dataset. Sex and total CDR3aa number are depicted by samples. Males are depicted in violet and females in orange and local motif in blue-green and global motifs in magenta.

Supplemental Figure 15: Enrichment of cancer associated, unmodified self-peptide specific and polyspecific TCR, in our thymic dataset compared to initial curated database.

(A) Representation of the specificity groups in the pooled and curated database. **(B)** Proportion of unique CDR3aa in the specific TCR repertoire for each samples for autoimmune disease (AID), bacteria, cancer, human and virus groups is compared between males and females. **(C)** Proportion of polyspecific CDR3aa in the specific repertoire of samples is compared between males and females. Stars indicate statistical differences males or females with the pooled and curated database based on the p-value of the Wilcoxon test (*: $p < 0.05$, **: $p < 0.01$). Males are depicted in violet and females in orange.

Supplemental Figure 16: Enrichment of AID associated TCR and bacteria targeted TCR in CD8 female cells.

CDR3aa usage of CDR3aa with a specificity in autoimmunity disease, bacteria, cancer, human and virus groups. Stars indicate statistical differences between males and females based on the p-value of the Wilcoxon test (*: $p < 0.05$, **: $p < 0.01$). Males are depicted in violet and females in orange.

Model: Simple Age Effect

Supplemental Figure 17: No association between age and thymic usage of CDR3 sequences specific to autoimmunity or bacterial antigens.

Linear regression models assessing the effect of donor age on the cumulative usage (%) of TRB CDR3aa sequences with known specificity to self-antigens associated with autoimmunity (AID, left panels) or bacterial antigens (right panels), in CD8 SP (top) and CD4 Treg SP (bottom) thymic subsets. Each dot represents one donor. The shaded area indicates the 95% confidence interval around the regression line. Reported slope, 95% confidence interval, R^2, and p-values show no significant effect of age.

	DP cells	CD8 cells	CD4 Teff	CD4 Treg
TRA	1013	2433	3314	2654
TRB	1429	3496	4551	5383

Supplemental Table 1: Minimal number of CDR3aa by cell subset.