Title: Type-I nNOS neurons orchestrate cortical neural activity and vasomotion

Authors: Kevin L. Turner[1,2,3,7], Dakota F. Brockway[4,7], Md Shakhawat Hossain[2,3,7], Keith R. Griffith[4,7], Denver I. Greenawalt[5,7], Qingguang Zhang[1,2,7,8], Kyle W. Gheres[1,2,7], Nicole A. Crowley[2,3,4,7], Patrick J. Drew[1,2,3,6,7]

Affiliations:

1. Department of Engineering Science and Mechanics, The Pennsylvania State University, University Park, PA 16802

2. Center for Neural Engineering, The Pennsylvania State University, University Park, PA 16802

3. Department of Biomedical Engineering, The Pennsylvania State University, University Park, PA 16802

4. Department of Biology, The Pennsylvania State University, University Park, PA 16802

5. Graduate Program in Molecular Cellular and Integrative Biosciences, The Pennsylvania State University, University Park, PA 16802

6. Department of Neurosurgery, The Pennsylvania State University, University Park, PA 16802

7. Penn State Neuroscience Institute, The Pennsylvania State University, University Park, PA 16802

8. Department of Physiology, Michigan State University, East Lansing, MI 48824

*Corresponding author: Patrick J. Drew - pjd17@psu.edu

Acknowledgements: This work was supported by NIH Grants R01NS078168, R01NS079737, U19NS128613 to PJD, R01AA029403 to NAC, an American Heart Association Predoctoral fellowship 24PRE1201066 to MSH and F31ES036154 to DIG. DIG was supported by T32GM108563 and MSH was supported by T32NS115667.

Conflict of Interest: The authors declare no competing financial interests.

Author Contributions K.L.T, N.A.C, and P.J.D. designed the experiments. K.L.T, D.F.B, M.S.H, K.R.G, D.I.G, and Q.Z performed experiments. K.L.T., M.S.H, Q.Z, K.W.G., and P.J.D. analyzed the data. K.L.T. and P.J.D. wrote the paper.

Abstract: It is unknown how the brain orchestrates coordination of global neural and vascular dynamics. We sought to uncover the role of a sparse but unusual population of genetically-distinct interneurons known as type-I nNOS neurons, using a novel pharmacological strategic to unilaterally ablate these neurons from the somatosensory cortex of mice. Region-specific ablation produced changes in both neural activity and vascular dynamics, decreased power in the delta-band of the local field potential, reduced sustained vascular responses to prolonged sensory stimulation, and abolished the post-stimulus undershoot in cerebral blood volume. Coherence between the left and right somatosensory cortex gamma-band power envelope and blood volume at ultra-low frequencies was decreased, suggesting type-1 nNOS neurons integrate long-range coordination of brain signals. Lastly, we observed decreases in the amplitude of resting-state blood volume oscillations and decreased vasomotion following the ablation of type-I nNOS neurons. This demonstrates that a small population of nNOS-positive neurons are indispensable for regulating both neural and vascular dynamics in the whole brain and implicates disruption of these neurons in diseases ranging from neurodegeneration to sleep disturbances.

Keywords: neurovascular coupling, type-I nNOS neurons, somatosensory cortex, two-photon, cerebral blood volume, widefield optical imaging, sleep

Introduction

Coordinated neural and hemodynamic activity across the brain are vital for arousal and cognition. The relationship between changes in neural activity and (typically measured) changes in blood volume, flow, and oxygenation, which are frequently studied with hemodynamic imaging and form the foundation of modern neuroscience, are mediated by local vasodilation and is known as neurovascular coupling (Drew, 2019a; Schaeffer, 2021). The vasodilator nitric oxide (NO) (Hosford, 2019), produced by neuronal nitric oxide synthase (nNOS) in neurons, provides a key mechanism of arteriole dilation. Neurons in somatosensory cortex that express nNOS are activated by sensory stimulation, volitional whisking, and locomotion (Echagarruga, 2020; Ahn, 2023), all of which produce robust hemodynamic responses (Huo, 2014; Winder, 2017). Chemogenetic elevation or suppression of nNOS neuron activity alters both baseline diameter and behaviorally-evoked dilation of arterioles (Echagarruga, 2020). Similar vascular changes are produced by local infusion of a NOS inhibitor (Echagarruga, 2020), implicating NO signaling directly in vascular regulation. Optogenetic activation of nNOS-positive neurons produces varying effects, with some reports demonstrating vasodilation and blood flow increases with minimal changes in neural activity (Krawchuk, 2020; Lee, 2020; Ruff, 2024), while others show large, low-frequency EEG changes during sleep and quiet wakeful states (Gerashchenko, 2018). While NO is known to modulate the excitability of neurons (Kara, 1999; Smith, 2003), artificial activation of nNOS neurons seems to drive vascular responses that are disproportionally larger than any corresponding neural changes, suggesting that this small population of neurons has an impact on vascular signals that is substantially greater than that of overall neural activity.

Two distinct GABAergic interneurons types express nNOS, denoted as type-I and type-II (Kawaguchi, 1997; Kubota, 2011; Perrenoud, 2012). nNOS-positive neurons make up approximately 20% of GABAergic interneurons in the cortex (Hendry, 1987; Sahara, 2012), with type-I constituting only 0.5–2% depending on brain region (Yan, 1997; Tricoire, 2012; Chong, 2019). Type-I nNOS neurons are sparse, somatostatin-positive, most dense in the deep layers of cortex, and express nNOS at higher levels than type-II. They extend both local and long-range projections throughout the cortex, receive excitatory input from nearby pyramidal neurons and drive increases in cerebral blood flow when stimulated (Ruff, 2024). Type-II nNOS neurons are in contrast much more abundant, smaller in size and morphology, and express nNOS less strongly than type-I. Type-II are a more heterogeneous group that express a variety of interneuron subtype markers and are more evenly distributed throughout the different layers of cortex (Yan, 1996; Perrenoud, 2012). Type-I nNOS neurons have been observed in rodents, carnivores, and primates (Tomioka, 2005; Higo, 2007; Tomioka, 2007) and are active during sleep (Gerashchenko, 2008; Kilduff, 2011; Morairty, 2013; Dittrich, 2015). In the cortex, type-I nNOS neuron density is anticorrelated with vascular density (Wu, 2022). Type-I nNOS neurons integrate local activity from feedforward excitatory pathways from the cortex and thalamus, likely contributing to the correlated changes in blood flow that are observed between functionally connected regions across hemispheres (Ruff, 2024). Molecular, electrophysiological, and immunohistochemistry studies have demonstrated that type-I nNOS neurons are the only cells in the cortex that express the tachykinin receptor 1 (also known as neurokinin 1 receptor, TACR1/NK1R), the primary receptor for the endogenous

neuropeptide substance P (Penny, 1986; Vruwink, 2001; Kubota, 2011; Dittrich, 2012; Endo, 2016; He, 2016; Huang,

2016; Paul, 2017; Vanlandewijck, 2018; Matsumura, 2021; Ruff, 2024). Local infusion of substance P into the cortex

causes a sustained increase in basal arterial diameter that is dependent upon local neural activity (Echagarruga,

2020). Type-I nNOS neurons likely receive substance P from the ~40% of parvalbumin (PV)-positive cortical

interneurons (Vruwink, 2001; Pfeffer, 2013; Bugeon, 2022) that are thought to drive network synchrony (Cardin,

2009; Sohal, 2009). Optogenetic stimulation of PV neurons for several seconds drives a biphasic hemodynamic

response, comprised of an early constriction (driven by suppression of overall population activity through GABA

release), and a delayed, prolonged dilation lasting tens of seconds that is mediated indirectly by substance P (Vo,

2023). The delayed vasodilation was not directly related to pyramidal neuron activity, was blocked by TACR1

antagonists, and was occluded by substance P, suggesting that activation of PV neurons drives downstream

activation of type-I nNOS neurons. These experiments point to type-I nNOS neurons as having a large effect on

vascular dynamics despite their smaller contribution in driving overall neural activity. However, optogenetic

stimulation only activates one component of a circuit in isolation, in an otherwise healthy and intact microcircuit,

potentially giving an incomplete picture of the neuron's role. Type-I nNOS neurons are uniquely vulnerable to stress

(Han, 2019), and their loss is likely in part causal to neurodegeneration (Schaeffer, 2021; Iadecola, 2023), making it

vital to understand brain wide changes following a more physiologically-relevant model of perturbation.

Here we sought to reveal the role of type-I nNOS neurons in controlling neural and vascular dynamics in the

cortex by selectively ablating them with saporin conjugated to a substance P analog (SP-saporin). Using this targeted

ablation approach, we found that ablation of type-I nNOS neurons caused decreases in the hemodynamic response

to sustained vibrissae stimulation, eliminated the post-stimulus undershoot, decreased local field potential power

in the delta-band (1-4 Hz), reduced bilateral correlations in gamma-band power and blood volume across arousal

states, and reduced the amplitude of resting-state blood volume oscillations. Together, these experiments

demonstrate that a small subset of type-I nNOS neurons regulate key neural and vascular dynamics that coordinate

changes in vasomotion.

Results

We investigated the effects of localized ablation of type-I nNOS neurons in somatosensory cortex on

neurovascular coupling and functional connectivity in unanesthetized, head-fixed mice. C57BL6J mice (119 total,

both male and female) were injected with either saporin conjugated to a substance P analog with high affinity for

the substance P receptor (SP-SAP) or a scrambled peptide as a control (Blank-SAP) into a localized region of one

hemisphere's somatosensory cortex. We used widefield optical imaging, electrophysiology, and 2-photon

microscopy to evaluate neural and hemodynamic changes following targeted ablation of type-I nNOS neurons in the

vibrissae (whisker) representation of somatosensory cortex while carefully monitoring arousal state (see **Materials**

and Methods). All imaging was performed during the animals' light cycle. Statistics included generalized linear

mixed-effects model (GLME), general linear models (GLM), or unpaired t-tests with corrections for multiple

comparisons when necessary.

107 **Saporin-conjugated peptides produce selective targeted ablation of type-I nNOS neurons**

108 While optogenetic and chemogenetic models give us insights into the potential function of neuron subtypes, it

109 is difficult to completely silence neurons in vivo, and the patterns of activity they induce are not physiological and

110 can have paradoxical effects on neural activity (Li, 2019; Andrei, 2021). Saporin provides a pharmacological route to

Figure 1. Targeted ablation of type-I nNOS neurons with cortical SP-SAP injections (a) Schematic showing intracortical administration of the ribosome inactivating protein saporin conjugated to either SP or a vehicle control. **(b)** Representative immunofluorescence of anti-nNOS (yellow) and anti-NKR1 (blue) showing colocalization of the NK1 receptor on cortical nNOS-positive neurons. Representative immunofluorescence of anti-nNOS (yellow) and DAPI (magenta) from animals with no injection (Uninjected, left), Blank-SAP (middle), or SP-SAP (right). **(c)** Representative immunofluorescence of Blank-SAP (top) and SP-SAP (bottom) sections co-stained with nNOS (left), IBA1 (middle left), GFAP (middle), DAPI (middle right), or NeuN (right). NeuN image was taken from section immediately adjacent to the first four. **(d)** Quantification of nNOS counts, **(e)** IBA1 counts, **(f)** GFAP fluorescence, **(g)** DAPI fluorescence, **(h)** NeuN fluorescence. Error bars **(d-h)** denote SD. Bonferroni correction (5) *$\alpha < 0.01$, **$\alpha < 0.002$, ***$\alpha < 0.002$. **k-l** *$\alpha < 0.05$, **$\alpha < 0.01$, ***$\alpha < 0.001$, GLME.

selectively kill only the cells that internalize it, and saporin conjugated to a peptide effectively targets neurons that express a receptor for the peptide (McKay, 2008; Abbott, 2012), allowing targeting of specific cell types orthogonal to any genetic targeting techniques.

We first sought to validate the efficacy and specificity of saporin-based targeting of cortical TACR1-expressing neurons. Type-I nNOS neurons were ablated by injection of the ribosome inactivating protein saporin (SAP), conjugated either to substance P (SP-SAP) or a scrambled peptide as a control (Blank-SAP) (**Fig. 1a**). The SP-bound SAP toxin binds to TACR1-expressing neurons (Wiley, 1999; Martin, 2001; Wang, 2002), which in the cortex are exclusively expressed by type-I nNOS neurons (**Fig. 1b**). We used both immunofluorescence (**Fig. 1c**) and NADPH diaphorase staining (a histochemical marker for NOS, **Fig. S1**), to visualize nNOS neuron ablation (Hope, 1989; Bredt, 1991; Hope, 1991). Immunofluorescent labeling in mice injected with Blank-SAP showed labeling of nNOS-positive neurons near the injection site. In contrast, mice injected with SP-SAP showed a clear loss in nNOS-labeling around the injection site, with nNOS-positive neurons both subcortically and in the entirety of the contralateral hemisphere remaining intact. We quantified the efficacy of SP-SAP in removing type-I nNOS neurons by quantifying the number of nNOS-positive neurons per square mm of cortical tissue (**Fig. 1d**), of which SP-SAP injected mice (N = 5, 3M/2F) had a significant reduction of nNOS-labeled neurons (1.3 ± 1.6 neurons/mm^2) compared to Blank-SAP mice (N = 8, 4M/4F) with 10.3 ± 2.4 neurons/mm^2 (GLME $p = 5.74 \times 10^{-6}$).

We checked for non-specific effects driven by SP-SAP injections one-month post-administration by staining for ionized calcium binding adaptor molecule 1 (IBA1), glial fibrillary acidic protein (GFAP), DAPI, and NeuN (neuronal nuclei) as well as nNOS (as a positive control for ablation). There was no difference in the number of microglia (**Fig. 1e**) between groups (Blank-SAP: 345.8 ± 47.7 microglia/mm^2; SP-SAP: 373.8 ± 69.4 microglia/mm^2). There was an increase in GFAP labeling following SP-SAP injection (1246.3 ± 514.8 AU/mm^2; Blank-SAP: 527.2 ± 230.9 AU/mm^2, GLME $p = 0.0029$) (**Fig. 1f**). There was no significant difference in DAPI fluorescence (Blank-SAP: 3511.2 ± 524.42 AU/mm^2; SP-SAP: 5180.7 ± 2135.0 AU/mm^2; GLME $p = 0.038$, **Fig. 1g**) or NeuN fluorescence (Blank-SAP: 4296.8 ± 339.4 AU/mm^2; SP-SAP: 4652.1 ± 196.8 AU/mm^2; GLME $p = 0.043$, **Fig. 1h**) between groups after Bonferroni correction for (5) multiple comparisons (**Fig. 1d-h**). Together, these findings indicate that the SP-SAP toxin was highly selective in ablating type-I nNOS neurons with minimal non-specific effects. All mice that underwent imaging were histologically verified for successful type-I nNOS neuron ablation using NADPH diaphorase staining. We quantified a subset of these mice and saw a similar specific removal of type-I nNOS neurons (**Fig. S1**). To determine if ablation of type-I nNOS neurons from the somatosensory cortex had any impact on behavior or arousal state, we assayed exploratory behavior, sleep, and pupil dynamics (**Fig. S2**). To assay exploratory behavior, mice were placed in a novel open field environment and allowed to explore for 5 minutes while quantifying distance traveled and time spent in the center, two metrics for evaluating stress and anxiety in rodents (Roth, 1979; Seibenhener, 2015). We noted no differences in any metric evaluating open field behavior (N = 12-23 mice/group). We also assessed multiple measurements of sleep quality and quantity. There was no difference in the percentage of time each mouse spent

145 in rapid eye movement (REM) or non-REM (NREM) sleep, or the percentage of time they spent volitionally whisking

146 while awake (N = 9 mice/group). We also noted no significant differences in eye-related arousal state metrics

147 including interblink-interval, pupil size, and pupillary response to vibrissae stimulation (Turner, 2023). Together, this

148 suggests that ablation of type-I nNOS neurons had no gross effects on sleep or general ambulatory behavior.

Impact of type-I nNOS neuron removal on neural and hemodynamic signals across arousal states

To determine the impact localized ablation of type-I nNOS neurons had on neural and hemodynamic signals in

the somatosensory cortex, we used widefield optical imaging (Sirotin, 2009; Huo, 2014) to measure changes in total

hemoglobin (Δ[HbT], an indicator of blood volume). We measured neural activity using either implanted electrodes

(Winder, 2017) to measure changes in local field potential (LFP) (**Fig. 2a, c, d**), or in a separate cohort of mice, pan-

neuronal expression of the calcium indicator GCaMP7s (Chan, 2017; Dana, 2019), which provides complementary

measures of bulk neural activity (**Fig. 2b, e, f**). In mice expressing GCaMP7s, we measured Δ[HbT], as well as changes

Figure 2. Simultaneous measurement of neural and hemodynamic signals in mice across arousal states (a) Schematic of widefield optical imaging experimental setup. The brain is illuminated at an isosbestic wavelength of hemoglobin (568 nm). Changes in reflected light measuring changes in total blood volume are captured by a camera mounted above the head while several other cameras monitor animal behavior and arousal state including vibrissae and pupil tracking. Vibrissae stimulation is done by directed air puffs. Polished and reinforced thinned-skull windows were bilaterally implanted over the somatosensory cortex. Tungsten stereotrodes were implanted underneath each window to record changes in local field potential within the area of interest. An additional hippocampal stereotrode and a neck electromyography electrode were used to assist in sleep scoring. **(b)** Schematic of widefield optical imaging experiments with GCaMP7s. Alternating illumination with 480 nm, 530 nm, and 630 nm light were used to measure changes in total hemoglobin, blood oxygenation, and GCaMP7s fluorescence. **(c, e)** During the awake state, power in low-frequency cortical LFP is low and power in the gamma-band (30-100 Hz) is elevated during activity such as volitional whisking. **(d, f)** Periods of NREM and REM sleep are accompanied by large oscillations in cerebral blood volume, with large increases in power in delta-band (1-4 Hz) cortical LFP during NREM and large increases in power in theta-band (4-10 Hz) hippocampal LFP during REM.

in cerebral oxygenation (Δ[HbO]) and deoxygenation (Δ[HbR]) using alternating illumination at 480/530/630 nm (Ma,

2016b; Ma, 2016a; Zhang, 2019). We corrected for hemodynamic contamination of GCaMP7s signals using the

simultaneously acquired hemoglobin signals (Kramer, 1979; Ma, 2016b; Wright, 2017; Scott, 2018). Measurements

of neural and hemodynamic signals were taken bilaterally through polished and reinforced thinned-skull windows

(Drew, 2010; Shih, 2012) in the vibrissae representation of somatosensory cortex. Each animal was habituated to

head-fixation over the course of several days following surgery. For the first 60 minutes of each recording session,

the vibrissae were briefly stimulated with directed puffs of air to either the left or right pad, or by a puffer directed

away from the mouse as an auditory control (Drew, 2011). Afterwards, each mouse was given several hours to

naturally sleep with no stimulation. To determine the arousal state of the mouse, we performed electromyography

of the nuchal muscles of the neck, tracked vibrissae movement and pupil diameter using video, and body movements

with a force sensor. Arousal state was scored in 5 second intervals as either Awake, NREM, or REM as previously

described (Turner, 2020) from behavioral and physiological data using a bootstrapped random forest classification

algorithm. We saw no differences in the accuracy or validity of our sleep-scoring models across different

experimental conditions (Blank-SAP; SP-SAP; Uninjected) (**Fig. S3**).

170 Ablation of type-I nNOS neurons reduces the stimulus-evoked response

We first determined the impact of removal of type-I nNOS neurons on evoked hemodynamic signals. The initial

increase in blood volume in response to brief (0.1 second) stimulation of the contralateral vibrissae was not affected

by ablation of type-I nNOS neurons (**Fig. 3a**), but the post-stimulus undershoot was absent in the SP-SAP mice. When

we evaluated the canonical post-stimulus hemodynamic undershoot from 2:4 seconds, the Blank-SAP group (N = 9,

5M/4F) had a mean of -2.2 ± 0.5 μM, compared to the SP-SAP group (N = 9, 5M/4F) mean of 1.0 ± 0.6 μM (GLME *p*

= 0.0005). This result is consistent with the observation that type-I nNOS neurons express the vasoconstrictory

neuropeptide Y (NPY) (Karagiannis, 2009) which is thought to underlie this post-stimulus undershoot (Uhlirova,

2016). It is also possible that loss of NO signaling from type-I nNOS neurons could also contribute through

interactions with blood volume to generate oscillations and post-dilation undershoots (Haselden, 2020). Using three-

wavelength spectroscopy in mice expressing GCaMP7s, we saw consistent increases in blood volume, neural activity,

and cerebral oxygenation during prolonged (5 seconds) vibrissae stimulation. Δ[HbT] evaluated 1.5:6.5 seconds

Figure 3. Ablation of type-I nNOS neurons reduces stimulus-evoked hemodynamic responses (a) Change in total hemoglobin in response to brief (0.1 second) vibrissae stimulation. **(b)** Change in total hemoglobin in response to extended (5 seconds) vibrissae stimulation. **(c)** Change in GCaMP fluorescence in response to extended vibrissae stimulation. **(d)** Change in oxy- and deoxy- hemoglobin in response to extended vibrissae stimulation. **(e)** Change in arteriole diameter in response to extended vibrissae stimulation. Error bars represent population averages ± SEM. All statistics were evaluated between the indicated intervals. *p < 0.05, **p < 0.01, ***p < 0.001, GLME.

following stimulus-onset decreased from 18.7 ± 1.7 µM in Blank-SAP (N = 7, 3M/4F) to 13.1 ± 1.4 µM in SP-SAP (N =

8, 4M/4F, GLME *p* = 0.017) (**Fig. 3b**). Calcium signals (**Fig. 3c**) evaluated 2:5 seconds following stimulus onset

decreased from 8.2 ± 1.3 % in Blank-SAP to 5.3 ± 0.7 % in SP-SAP (GLME *p* = 0.047). Δ[HbO] (**Fig. 3d**) evaluated 1.5:6.5

seconds following stimulus onset decreased from 24.6 ± 1.9 µM in Blank-SAP to 17.4 ± 1.7 µM in SP-SAP (GLME *p* =

0.010) while Δ[HbR] evaluated from 1.5:6.5 seconds increased from -6.0 ± 0.4 in Blank-SAP to -4.4 ± 0.3 in SP-SAP

(GLME *p* = 0.005). Evaluation windows were determined based on the duration of stimuli and the delayed onset of

the hemodynamic response.

We further investigated single arterial dynamics during vibrissae stimulation with 2-photon microscopy.

Following ablation of type-I nNOS neurons, after vibrissae stimulation there was a decrease in arteriole diameter

(**Fig. 3e**) evaluated 3:7 seconds following stimulus onset from 11.6 ± 0.8 %·s in Blank-SAP (N = 9, 5M/4F, n = 81

arterioles) down to 7.8 ± 0.6 % in SP-SAP (N = 7, 2M/5F, n = 70 arterioles, GLME *p* = 0.008). The arterial changes

closely mirror the Δ[HbT] changes, consistent with the substantial contribution of arterial dynamics to the blood

volume signal (Huo, 2015). We also evaluated changes in blood volume during voluntary locomotion, where we saw

a similar trend in decrease in locomotion-evoked Δ[HbT] evaluated 1.5:2.5 seconds following locomotion onset from

17.5 ± 2.0 µM in Blank-SAP (N = 7, 5M/2F) to 12.9 ± 2.0 µM in SP-SAP (N = 7, 2M/5F), but it was not statistically

significant (GLME *p* = 0.10). These results show the removal of a very small number of neurons can drive substantial

reductions in both hemodynamic and neural response to sensory stimulation.

Type-I nNOS ablation reduces low frequency neural activity

To determine the effect that type-I nNOS neuron removal had on neural activity, we assessed the power

spectrum of the LFP in different arousal states, which we classified into alert, asleep, and all data (Turner, 2020).

Data was classified into the alert state when 15-minute blocks were predominantly (> 80%) in the awake state. Alert

periods contained fidgeting movements and bouts of whisking interspersed with awake quiescence. Periods

classified as asleep were predominantly composed of sleeping states (NREM/REM > 80% of all classifications in a 15-

minute block) with only brief periods of wakefulness, typically occurring during transitions between sleep states. The

'all' state includes everything irrespective of arousal state classification. The first ~60 minutes of each recording

Figure 4. Type-I nNOS ablation alters low frequency neural activity and gamma-band neurovascular coupling (a) Local field potential within the vibrissae representation of somatosensory cortex during periods of Alert. **(b)** Asleep. **(c)** All data. **(d)** Change in gamma-band power vs. Δ[HbT] following brief (0.1 seconds) vibrissae stimulation. **(e)** Change in GCaMP7s fluorescence vs. Δ[HbT] following prolonged (5 seconds) vibrissae stimulation. Error bars represent population averages ± SEM. All statistics were evaluated between the indicated intervals. *p < 0.05, **p < 0.01, ***p < 0.001, GLME (**a-c**), GLM (**d, e**).

session which included vibrissae stimulation was excluded. There was a pronounced reduction in the delta-band (1-

4 Hz) power of the LFP in all three arousal state catagories. The power in the delta-band of the LFP in Blank-SAP mice

(N = 9, 4M/5F) in the alert state (**Fig. 4a**) was $3.4 \times 10^{-10} \pm 1.0 \times 10^{-10}$ a.u. compared to $1.0 \times 10^{-10} \pm 3.0 \times 10^{-11}$ a.u. in the

SP-SAP mice (N = 9, 5M/4F, GLME p = 0.011). The power in the delta-band of the LFP in Blank-SAP mice (N = 7, 3M/4F)

in the asleep state (**Fig. 4b**) was (NREM + REM) was $8.5 \times 10^{-10} \pm 3.0 \times 10^{-10}$ a.u. compared to $2.4 \times 10^{-10} \pm 9.1 \times 10^{-11}$ a.u.

in the SP-SAP mice (N = 7, 4M/3F, GLME p = 0.022). The power in the delta-band of the LFP in Blank-SAP mice (N =

9, 4M/5F) averaged across all arousal states (**Fig. 4c**) was $5.2 \times 10^{-10} \pm 1.7 \times 10^{-10}$ a.u. compared to $1.5 \times 10^{-10} \pm 4.3 \times 10^{-11}$

a. u. in the SP-SAP mice (N = 9, 5M/4F, GLME p = 0.016). These result parallel those seen in studies where knockout

of nNOS produces lower delta band power (Morairty, 2013) and optogenetic stimulation of nNOS neurons produces

low frequency oscillations (Zielinski, 2019), suggesting that type-I nNOS neurons promote these slow oscillations via

NO release. In contrast to the observed reductions in LFP, we noted no gross changes in the power spectra of cerebral

blood volume fluctuations between groups during any arousal state (**Fig. S4**).

We next wanted to see whether the amount of neural activity and corresponding Δ[HbT] in response to vibrissae

stimulation was altered following type-I nNOS neuron ablation. As a measure of neurovascular coupling, we used

the slope of the line fitting the change in gamma-band power vs. Δ[HbT] following brief (0.1 second) vibrissae

stimulation (**Fig. 4d**). Decreases in the slope indicate a smaller vascular response for a given amount of neural

activity, indicating a decrease in neurovascular coupling. We found the slope was significantly increased in SP-SAP

mice compared to Blank-SAP mice (Blank-SAP: 0.02 ± 0.03; SP: 0.005 ± 0.006, GLM p = 0.0001). We also evaluated

the change in GCaMP7s fluorescence vs. Δ[HbT] during prolonged (5 seconds) vibrissae stimulation from 2:5 seconds

(**Fig. 4e**) and saw no significant change in the slope (Blank-SAP: 251.3 ± 75.3; SP: 256.5 ± 73.2, GLM p = 0.34). Given

that the sensory-evoked LFP is driven more by synaptic input, while GCaMP7s signals are driven by local neural

activity, these results indicate that the loss of type-I nNOS neurons reduces overall excitability, but that the input

drive from other areas might be increased, potentially due to homeostatic mechanisms at the input synapses

(Turrigiano, 2008).

Neurovascular coupling was weakly affected by type-I nNOS removal

We next looked at the cross-correlation between blood volume and neural signals, which provides a measure

of spontaneous neurovascular coupling. For electrophysiological measures, we measured the cross-correlation

between gamma-band power and blood volume (**Fig. 5a**). We evaluated the peak cross-correlation during the

resting-state, during long periods while alert, or long periods while asleep. The cross-correlation between neural

activity and hemodynamic signals is substantially higher during periods of behavior (whisking, fidgeting) than during

the resting-state because self-generated motion and whisking drive increases in neural activity and vasodilation

(Winder, 2017; Drew, 2019b; Stringer, 2019; Claron, 2023; Tu, 2024) . The correlation between neural activity and

blood volume changes is higher during sleep than in the awake state (Turner, 2020). We saw no significant difference

in the peak cross-correlation during the resting-state (**Fig. 5b**) between Blank-SAP mice (N = 9, 4M/5F, 0.08 ± 0.007)

and SP-SAP mice (N = 9, 5M/4F, 0.06 ± 0.008, GLME *p* = 0.22). We did observe a significant difference in the peak

cross-correlation during both periods of alert and asleep, from 0.23 ± 0.01 (Blank-SAP) to 0.16 ± 0.03 (SP-SAP, GLME

p = 0.036) during alert periods (**Fig. 5c**) and 0.40 ± 0.02 (Blank-SAP) dropping to 0.33 ± 0.02 (SP-SAP, GLME *p* = 0.012)

while asleep (**Fig. 5d**). We next wanted to evaluate whether this drop in neurovascular coupling was more prevalent

at a particular modulation frequency. When analyzing the coherence between the gamma-band power and

hemodynamic signals during these different arousal states, we noted no dominant frequency and saw no significant

changes in the lower frequencies

associated with neurovascular coupling

(≤ 0.5 Hz). During the resting-state (**Fig.**

5e), the average coherence between 0.1

and 0.5 Hz was 0.28 ± 0.008 in Blank-SAP

mice and 0.23 ± 0.01 in SP-SAP mice

(GLME *p* = 0.23). The average coherence

between 0.01 and 0.5 Hz during alert

periods (**Fig. 5f**) was 0.29 ± 0.004 in

Blank-SAP mice and 0.25 ± 0.005 in SP-

SAP mice (GLME *p* = 0.06), and during

asleep periods (**Fig. 5g**) it was 0.34 ±

0.008 in Blank-SAP mice and 0.32 ±

0.007 in SP-SAP mice (GLME *p* = 0.46).

Lastly, to evaluate any changes in the

predictive power of the neural-

hemodynamic relationship, we fit a

hemodynamic response function using

the gamma-band power and

hemodynamic response following

periods of vibrissae stimulation. We

observed that ablation of Type-I nNOS

neurons did not alter the predictive

power of the hemodynamic response

function (**Fig. S5**).

We next evaluated neurovascular

coupling from optical measures of bulk

activity using GCaMP7s (**Fig. 5h**). The

peak resting-state cross-correlation (**Fig.**

Figure 5. Neurovascular coupling was only weakly affected by type-I nNOS removal (a) Schematic demonstrating intracortical injection of either Blank-SAP or SP-SAP and the analysis of gamma-band power and hemodynamic signals from within the vibrissae representation of somatosensory cortex, N = 9 mice per group. **(b)** Gamma-Δ[HbT] resting-state cross-correlation **(c)** Gamma-Δ[HbT] alert cross-correlation **(d)** Gamma-Δ[HbT] asleep cross-correlation **(e)** Gamma-Δ[HbT] resting-state coherence **(f)** Gamma-Δ[HbT] alert coherence **(g)** Gamma-Δ[HbT] asleep coherence **(h)** Schematic demonstrating intracortical injection of either Blank-SAP or SP-SAP and the analysis of GCaMP7s fluorescence and hemodynamic signals from within the vibrissae cortex, n = 6-7 mice per group. **(i)** GCaMP7s-Δ[HbT] resting-state cross-correlation **(j)** GCaMP7s-Δ[HbT] alert cross-correlation (k) GCaMP7s-Δ[HbT] asleep cross-correlation **(l)** GCaMP7s-Δ[HbT] resting-state coherence **(m)** GCaMP7s-Δ[HbT] alert cross-correlation **(n)** GCaMP7s-Δ[HbT] asleep cross-correlation. Error bars represent population averages ± SEM. *p < 0.05, **p < 0.01, ***p < 0.001, GLME.

5i) between GCaMP7s fluorescence and ongoing hemodynamics showed a significant decrease, 0.68 ± 0.02 with

Blank-SAP versus 0.62 ± 0.02 with SP-SAP (GLME p = 0.038). However, there was no drop in peak cross-correlation

during either alert or asleep as seen with the gamma-band power. During alert periods (**Fig. 5j**), the peak GCaMP7s

cross-correlation was 0.78 ± 0.12 with Blank-SAP and 0.77 ± 0.02 with SP-SAP (GLME p = 0.54) while during asleep

periods (**Fig. 5k**) it was 0.88 ± 0.008 with Blank-SAP and 0.86 ± 0.04 with SP-SAP (GLME p = 0.58). Like the peak in

cross-correlation, the average resting-state coherence (**Fig. 5l**) did show a significant drop across the lower

frequencies, 0.75 ± 0.01 with Blank-SAP vs. 0.70 ± 0.01 with SP-SAP (GLME p = 0.049). There was no drop in coherence

during alert periods (**Fig. 5m**) at 0.78 ± 0.005 with Blank-SAP and 0.77 ± 0.004 with SP-SAP (GLME p = 0.42) nor during

asleep periods (**Fig. 5n**) at 0.75 ± 0.009 with Blank-SAP and 0.74 ± 0.01 with SP-SAP (GLME p = 0.63). While there are

small changes in neural-hemodynamic correlations, the differences in neurovascular coupling across arousal states

were relatively small following localized removal of type-I nNOS neurons, meaning any effects of ablation on

hemodynamic responses are primarily mediated by changes in neural activity.

Type-I nNOS ablation reduces low-frequency interhemispheric coherence

Because type-I nNOS neurons send and receive many modulatory signals across a range of distances, they could help serve to coordinate neural and vascular dynamics across the brain. We tested this hypothesis by comparing how loss of type-I nNOS neurons changed the coherences of neural and hemodynamics signals between the left and right vibrissae cortex, which are generally highly correlated across all behaviors and frequencies (Turner, 2020). Neural activity is also bilaterally correlated, though less so than vascular dynamics. Average resting-state coherence between the left and right Δ[HbT] signals (**Fig. 6a**) in the vibrissae cortex were 0.79 ± 0.005 for Blank-SAP and 0.73 ± 0.008 for SP-SAP mice (p = 0.014, GLME, **Fig. 6b**). This reduction in coherence was seen over all frequencies (0-0.5 Hz) during both alert periods (**Fig. 6c**), at 0.86 ± 0.002 for Blank-SAP and 0.81 ± 0.003 for SP-SAP (p = 7.4×10⁻⁶, GLME), as well as asleep periods (**Fig. 6d**) with 0.86 ± 0.003 for Blank-SAP and 0.79 ± 0.005 for SP-SAP (p = 0.0008, GLME).

Figure 6. Type-I nNOS ablation reduces low-frequency interhemispheric coherence (a) Δ[HbT] coherence between bilateral ROIs in the left and right hemisphere's somatosensory cortex during **(b)** Res.t **(c)** Alert. **(d)** Asleep. **(e)** Gamma-band power coherence between bilateral ROIs in the left and right hemisphere's somatosensory cortex during **(f)** Res.t **(g)** Alert. **(h)** Asleep. **(i)** Δ[HbT] coherence between bilateral ROIs in the left and right hemisphere's somatosensory cortex during **(j)** Res.t **(k)** Alert. **(l)** Asleep. **(m)** GCaMP7s coherence between bilateral ROIs in the left and right hemisphere's somatosensory cortex during **(n)** Res.t **(o)** Alert. **(p)** Asleep. Error bars represent population averages ± SEM. *p < 0.05, **p < 0.01, ***p < 0.001, GLME.

We then looked at the coherence of gamma-band power across hemispheres, where average coherence for bilateral

gamma-band signals in the resting-state were 0.19 ± 0.006 (Blank-SAP) and 0.19 ± 0.01 (SP-SAP, *p* = 0.88, GLME, **Fig**

6e, f). There was no significant difference during the alert periods (**Fig 6g**) at 0.22 ± 0.005 (Blank-SAP) and 0.20 ±

0.006 (SP-SAP, *p* = 0.43, GLME), but ablation of type-I nNOS neuros produced a reduction of coherence during asleep

periods (**Fig. 6h**) at 0.35 ± 0.009 (Blank-SAP) and 0.29 ± 0.009 (SP-SAP, *p* = 0.047, GLME).

Average resting-state coherence across bilateral Δ[HbT] signals (**Fig. 6i**) taken from the vibrissae cortex were

0.83 ± 0.005 (Blank-SAP) and 0.78 ± 0.007 (SP-SAP, *p* = 0.021, GLME, **Fig. 6j**). This reduction in coherence persisted

across all frequencies during the both alert periods (**Fig. 6k**) at 0.92 ± 0.001 (Blank-SAP) and 0.90 ± 0.001 (SP-SAP, *p*

= 0.012, GLME) as well as sleep periods (**Fig. 6l**) at 0.90 ± 0.002 (Blank-SAP) and 0.83 ± 0.005 (SP-SAP, *p* = 0.004,

GLME). Average resting-state coherence across bilateral GCaMP7s signals (**Fig. 6m**) taken from the vibrissae cortex

were 0.75 ± 0.009 (Blank-SAP) and 0.70 ± 0.009 (SP-SAP, GLME *p* = 0.12, **Fig. 6n**). This reduction in coherence

persisted across all frequencies during the both alert periods (**Fig. 6o**) at 0.88 ± 0.003 (Blank-SAP) and 0.84 ± 0.003

(SP-SAP, *p* = 0.029, GLME) but not in sleep periods (**Fig. 6p**) at 0.85 ± 0.005 (Blank-SAP) and 0.84 ± 0.005 (SP-SAP,

GLME *p* = 0.52). When the correlation of bilateral signals was evaluated with Pearson's correlation coefficients, the

same general trend remained (**Fig. S6**). These results show that removal of type-I nNOS neurons reduces both

vascular and neural coordination across hemispheres.

Type-I nNOS neurons control vasomotion amplitude but not baseline diameter

We next wanted to establish how ablation of type-I nNOS neurons affected vasomotion, spontaneous

oscillations in the absence of behavior, measured at the scale of blood volume and at the level of single arterioles,

as well as the basal tone of blood vessels. The variance in Δ[HbT] during rest, a measure of vasomotion amplitude,

was significantly reduced following type-I nNOS ablation (**Fig. 7a**), dropping from 40.9 ± 3.4 μM^2 in the Blank-SAP

group (N = 24, 12M/12F) to 23.3 ± 2.3 μM^2 in the SP-SAP group (N = 24, 11M/13F) (GLME *p* = 6.9×10^{-5}). Individual

pial and penetrating arterioles showed the same reduction in vasomotion after type-I nNOS neuron ablation, with

Blank-SAP (N = 9, 5M/4F, n = 70 arterioles) having a resting diameter variance of 12.6 ± 1.4 %^{2} and SP-SAP (N = 7,

2M/5F, n = 65 arterioles) of 8.0 ± 0.8 %^{2} (**Fig. 7b**).

Figure 7. Type-I nNOS ablation reduces resting-state hemodynamics and low frequency neural activity (a) Variance in resting-state hemodynamics signals measured with widefield optical imaging. **(b)** Variance in resting-state arteriole diameter measured with two-photon. **(c)** Average Δ[HbT] during periods of NREM sleep, periods of REM sleep, and following administration of isoflurane (n = 9 mice per group). **(d)** Change in normalized arteriole diameter following administration of isoflurane Error bars represent population averages ± SD. All statistics were evaluated between the indicated intervals. *p < 0.05, **p < 0.01, ***p < 0.001 GLME.

If type-I nNOS neurons control the basal diameter of vessels via secreted vasodilators, potentially in an arousal

state dependent way, when we ablate these neurons we would observe a difference in blood volume across arousal

states and in the maximal dilation elicited with isoflurane (Gao, 2015). We saw no changes in the average change in

Δ[HbT] across states (**Fig. 7c**, **Fig. S7**). The resting-state Δ[HbT] in Blank-SAP mice was 0.29 ± 0.7 µM compared to

0.29 ± 0.9 µM in SP-SAP (GLME *p* = 0.997). NREM Δ[HbT] in Blank-SAP mice was 23.9 ± 5.8 µM compared to 24.1 ±

6.5 µM in SP-SAP (GLME *p* = 0.96). REM Δ[HbT] in Blank-SAP mice was 74.5 ± 8.4 µM compared to 82.3 ± 17 µM in

SP-SAP (GLME *p* = 0.26). The Δ[HbT] following isoflurane in Blank-SAP mice was 170.9 ± 19.7 µM compared to 190.5

± 30.5 µM in SP-SAP (GLME *p* = 0.15). We repeated the isoflurane experiment on a set of pial and penetrating

arterioles under 2-photon microscopy. The change in normalized arteriole diameter following administration of

isoflurane (n = 7-9 mice per group, 18-19 arterioles per group) was 61 ± 37.2% in Blank-SAP compared with 63.6 ±

38.4% in SP-SAP (*p* = 0.85, ttest). These results indicate that type-I nNOS neurons play a role in driving spontaneous

hemodynamic fluctuations, but not in setting the basal diameter during different states.

Discussion

We selectively ablated type-I nNOS neurons from the somatosensory cortex, which had marked effects on

neural activity and vascular dynamics, but minimal changes in neurovascular coupling. These results are surprising

given previous work showing stimulation of these neurons in isolation causes vasodilation and minimal neural

activity changes. Our approach of using SP-conjugated saporin allowed a nongenetic means of targeting a critical

neuronal cell type, supporting further exploration of the role of type-I nNOS neurons in transgenic mouse models of

disease without complicated breeding schemes, as well as in non-model organisms. Our results point to these

neurons being an important orchestrator of neural and vascular dynamics, as loss of these neurons causes

desynchronization between hemispheres of both neural and vascular signals, as well as altered responses to sensory

stimulation. While previous studies using specific activation of type-I nNOS neurons have emphasized the roles of

these neurons in driving vasodilation with minimal changes in neural activity, our results point to a larger role of

these neurons in organizing and patterning both spontaneous and sensory-evoked neural activity. This is likely due

to a loss in NO signaling as well as a loss of the many of the other neuropeptides expressed by type-I nNOS neurons,

which all have known effects on other neurons.

While previous studies have found that activation of type-I nNOS neurons, either optogenetically or via

administration of substance P, results in vasodilation (which would imply an important role for these neurons for

neurovascular coupling), our work provides insight into the integrated function of type-I nNOS neurons in neural

circuits and vascular dynamics. Importantly, methodological approaches differentiate our strategy from previous

published work. One interpretation is that there could be compensatory rewiring of local and/or network activity

following type-I nNOS ablation, where other signaling pathways from the neurons to the vasculature become

strengthened to compensate for the loss of vasodilatory signaling from the type-I nNOS neurons. While this likely

happens to some degree, this interpretation is less likely to completely account for the effect given these potential

compensatory changes do not prevent large changes in neural activity and resting vasomotion. Second, there may
be some non-linear interactions between vasodilatory signals and/or the vasodilator mechanisms that are arousal-
state-dependent. The diameter of arteries tracks the smooth muscle membrane potential linearly, up until a
saturation point above which any hyperpolarization does not induce further dilation (Knot, 1998; Wölfle, 2011; Hill,
2012). Many vasoactive pathways from neurons to vessels are known to exist, whose additive individual effects could
account for larger than biologically possible changes in vasodilation (Hosford, 2019). It may be that for dilation
occurring during sensory stimulation and sleep, the vasodilatory stimulus to vessels exceeds the saturation point, so
that the loss of one vasodilatory pathway does not further affect the vascular response (**Fig. S8**). This would also be
consistent with isolated optogenetic or chemical stimulation of type-I nNOS neurons being able to drive substantial
dilation, while the loss of these neurons does not have a large impact on the vascular response.

Finally, pharmacological ablation and optogenetic/chemogenetic activation are not mirror manipulations and
have differences that can produce non-symmetrical changes. Symmetrical changes would only occur if the neural
circuitry and the signaling to the vessel were completely linear. Optogenetic activation/deactivation of a single cell
type does not produce symmetrical changes in the activity of other neurons (Phillips, 2016), so there is little reason
to think ablation would have the exact opposite effect as activation. Furthermore, optogenetic/pharmacological
activation of a single cell type is unlikely to occur endogenously, as neural activity across different cell types is largely
correlated (Bugeon, 2022). For example, during whisker stimulation, not only are type-I nNOS neurons in the
somatosensory cortex activated (Ruff, 2024), but nearly every other cell type is as well (Staiger, 2021), which mean
that type-I nNOS neurons are not the only neurons, and not even the only NO producing neurons, sending signals to
the vasculature during sensory stimulation.

One surprise is that we observed no changes in the vasodilation during NREM sleep, where type-I nNOS neurons
are known to be active and play a role in inducing sleep (Kilduff, 2011). During NREM sleep, there is marked arterial
dilation (Turner, 2020; Gheres, 2023), and it would seem natural that type-I nNOS neurons might drive this. However,
we saw no difference in NREM dilation between the ablated and control mice. Again, this could be due to
compensation, saturation of the vasodilatory response, or it could be due to type-I nNOS neurons releasing other
vasoconstrictory peptides (SST, NPY) so their net effects are cancelled out. This last possibility is consistent with the
loss of the post-stimulus vasoconstriction seen after type-I nNOS neuron ablation. Again, like with sensory
stimulation, there may be a non-linearity in the vascular responses, so that another large factor, such as large
neuromodulatory changes during sleep, could dominate over any other vascular signaling factor. The large
norepinephrine decreases during NREM (Osorio-Forero, 2021; Kjaerby, 2022; Turner, 2023) could release the vessels
from tonic vasoconstriction (as norepinephrine is a vasoconstrictor (Bekar, 2012)), resulting in the dilation during
NREM.

Finally, we saw a marked reduction in spontaneous vascular oscillation (vasomotion) at rest, both at the level of
single arteries, and at the level of blood volume. These spontaneous oscillations move cerebrospinal fluid (CSF) (van
Veluw, 2020; Kedarasetti, 2022; Holstein-Ronsbo, 2023) which is important for clearing waste from the brain, and a

reduction in amplitude will decrease the pumping efficacy. Additionally, there is a loss of coordination of neural and

vascular dynamics across hemispheres after ablation of type-I nNOS neurons, indicated by the drop in coherence.

Although a loss of synchronous dilations/constrictions might affect CSF movement, we might imagine that it would

also adversely impact CSF pumping. Aged human brains show reduced interhemispheric synchrony in resting-state

signals (Zhao, 2020), analogous to what we see here. Type-I nNOS neurons also seem uniquely vulnerable to stress

(Han, 2019), and loss of type-I nNOS neurons reduces power in the 1-4Hz band of the LFP, which is positively

associated with CSF clearance (Hablitz, 2019; Jiang-Xie, 2024). It is possible that adverse life experiences could cause

the loss of type-I nNOS neurons, leading to reduction in CSF flow seen in age (Kress, 2014; Matrongolo, 2023) that is

thought to correlate with neurodegeneration.

Materials and Methods

This study was performed in accordance with the recommendations of the Guide for the Care and Use of

Laboratory Animals of the National Institutes of Health. All procedures were performed in accordance with protocols

approved by the Institutional Animal Care and Use Committee of Pennsylvania State University (Protocol

201042827). Data were acquired from 119 C57BL6/J mice (#000664, The Jackson Laboratory, Bar Harbor, ME)

compromised of 57 males and 62 females between 3 and 9 months of age. Food and water were provided *ad libitum*

and animals were housed on a 12-hr. light/dark cycle with all experiments occurring during the light cycle. Mice were

individually housed after surgery. Sample sizes are consistent with previous studies (Winder, 2017; Turner, 2020;

Turner, 2023) and experimenters were not blind to experimental conditions or data analysis except for histological

experiments.

Surgical Procedures. Saporin is a ribosome-inactivating protein that was conjugated to the Sar[9], Met(O$_2$)[11] analog of

Substance P (SP-SAP) or to a control peptide (Blank-SAP) (IT11 & IT21, Advanced Targeting Systems, Carlsbad, CA).

Mice were anesthetized using 5% isoflurane (2% maintenance) vaporized in pure oxygen and were then injected

intracortically with 4 ng of either SAP conjugate or Blank-SAP in 100 nL of artificial cerebrospinal fluid (aCSF). The

incision site was sterilized with betadine and 70% ethanol followed by a retraction of the skin atop the skull. A small

(< 0.5 mm) craniotomy was made above the vibrissae representation of somatosensory cortex (2 mm caudal, 3.25

mm lateral) and a sterile glass-pulled needle (tip diameter 50-100 μm) was inserted 500 μm beneath the cortical

surface at 45°. The SAP conjugate was slowly injected at 100 nL/min using a programmable syringe pump (Harvard

Apparatus, Holliston, MA) followed by closure of the incision with VetBond (3M, Saint Paul, MN). A subset of animals

were also injected retro-orbitally with 25 μL of AAV PHP.eB-syn-jGCaMP7s-WPRE [2x10^{13} GC/mL] (104487-PHPeB,

Addgene, Watertown, MA) diluted in 25 μL of sterile saline. Animals were given at least 2 weeks to recover prior to

undergoing additional procedures. For imaging, a custom-machined titanium head bar was adhered atop the

occipital bone of the skull using cyanoacrylate glue (Vibra-Tite 32402, ND Industries, Clawson, MI) and dental cement

(Ortho-Jet, Lang Dental, Wheeling, IL). Self-tapping 3/32' #000 screws (J.I. Morris, Oxford, MA) were implanted in

the frontal bones for structural stability. Electrodes were implanted into cortex and hippocampus using PFA-coated

tungsten stereotrodes (#795500, AM systems, Sequim, WA) for recordings of local field potentials (LFP) and into the neck muscles using a pair of PFA-coated 7-strand stainless-steel wires (#793200, AM systems, Sequim, WA) for electromyography (EMG). Polished and reinforced thinned-skull windows (Drew et al., 2010a; Shih et al., 2012b) were implanted over the somatosensory areas (either bilaterally or right hemisphere) using #0 coverslips (#72198, Electron Microscopy Sciences, Hatfield, PA). Detailed surgical procedures are as previously described (Turner, 2020).

Data acquisition. Data were acquired with a custom LabVIEW program (National Instruments; https://github.com/DrewLab/ LabVIEW-DAQ). Details on widefield optical imaging, electromyography (EMG), electrophysiology, vibrissae stimulation, and behavioral measurements including tracking of vibrissae and pupil diameter were performed as previously described (Turner, 2020; Zhang, 2022; Turner, 2023)). Mice were gradually acclimated to head-fixation of increasing duration (15, 30, 60 minutes) on the days preceding the onset of experiments. The vibrissae (left, right, or a third air puffer not directed at the body as an auditory control) were randomly stimulated with air puffs [0.1 seconds or a train of 5 second pulses, 10 pounds force per square inch (PSI)] occurring every 30–45 seconds for the first 1 hour of imaging. Data were acquired in 15-minute intervals with a brief (~30 seconds) gap in between for saving data to disk. Each animal was run for up to 6 imaging sessions lasting from 1-5 hours depending on experiment.

Histology. Following the conclusion of imaging experiments, animals were deeply anesthetized and transcardially perfused with heparin-saline followed by 4% paraformaldehyde. Presence or absence of nNOS-positive neurons in the injected hemisphere was verified using nicotinamide adenine dinucleotide phosphate (NADPH)-diaphorase staining for localizing the sparsely populated type-I nNOS neurons (Scherer-Singler, 1983).

Immunohistochemistry. All histological analyses were done blinded to the experimental condition. Mice were deeply anesthetized with 5% isoflurane and perfused transcardially with ice-cold phosphate buffered saline (PBS, pH 7.4) and 4% paraformaldehyde (PFA, pH 7.4). Brains were removed, post-fixed in PFA for 24 hours and stored in PBS at 4°C for less than 1 week. A fiduciary mark was placed in the right hemisphere. 40-μm free floating sections were sliced with a Leica vibratome (VS 1200, Leica) and stored in PBS for less than 1 week. Prior to immunostaining, slices were washed three times in PBS for 10 minutes each, and underwent antigen retrieval in 10 mM sodium citrate buffer (pH 6.0) at 80°C for 30 minutes. Slices were washed three times in PBS for 10 minutes each, and permeabilized in 0.5% Triton X-100 in PBS for 60 minutes. Nonspecific binding was blocked with 5% normal donkey serum (NDS) (ab7475) in 0.1% Triton X-100 in PBS for 60 minutes. Slices were then incubated in a primary antibody cocktail, including goat anti-nNOS (1:500, ab1376), rabbit anti-IBA-1 (1:500, ab178847), rat anti-GFAP (1:500, Thermofisher 13-0300), rabbit anti NeuN (1:2000, EnCor), rabbit anti-TACR1 (1:500 Invitrogen PA1-16713) in 2.5% NDS in 0.1% Triton X-100 in PBS for 48-h at 4°C. Slices were rinsed three times with PBS for 10 minutes each, and incubated in a fluorophore-tagged secondary antibody cocktail, including donkey anti-rabbit Alexa Fluor 488 (1:500, ab150073), donkey anti-goat Alexa Fluor 594 (1:500, ab150132), donkey anti rabbit Alexa fluor 647 (1:500, ab150075), donkey

anti rat Alexa fluor 488 (1:500 ab150153), for 4 hours at room temperature. Slices were rinsed again three times

with PBS, with the last step including DAPI (1:10,000, 10mg/mL, Millpore Sigma, 10236276001), mounted on glass

slides, air-dried and cover slipped with Immunomount (Thermo Fisher Scientific, Waltham, MA, United States).

Images were obtained with an Olympus BX63 upright microscope (Center Valley, PA, United States) under matched

exposure settings. Three to eight images from both hemispheres were taken per region.

Cell counting and immunofluorescence quantification. Total cell counts and absolute changes in

immunofluorescence (**Fig. 1**) were quantified using ImageJ (National Institutes of Health, Bethesda, MD, United

States). For total cell counts, a region of interest (ROI) was delineated, and cells were automatically quantified under

matched criteria for size, circularity and intensity. Each ROI's total cell count was divided by the ROI's area to give a

total density value (Dao, 2020; Smith, 2020; Sicher, 2023). Immunoreactivity was quantified as mean fluorescence

intensity of the ROI (Pleil, 2015).

Data Analysis. Data were analyzed with code written by K.L.T, M.S.H, Q.Z, K.W.G, and P.J.D. (MATLAB 2019b–2024a,

MathWorks).

Statistical analysis. Statistical evaluations were made using either generalized linear mixed effects (GLME), unpaired

t-test, or general linear model (GLM). GLME models had the arousal state as a fixed effect, mouse identity as a

random effect, and hemisphere [left/right (L/R), if applicable] as an interaction with the animal ID, or using a paired

t test where appropriate. Unless otherwise stated, statistical results report p values from a GLME test. All reported

quantifications are mean ± SD unless otherwise indicated. Unless otherwise noted, all pupil diameter measurements

are in z-units. MATLAB functions used were fitglme, ttest, fitglm.

Hemodynamic correction. Widefield imaging was done with a Dalsa 1M60 Pantera CCD camera (Phase One,

Cambridge, MA) with a magnifying lens (VZM 300i, Edmund Optics, Barrington, NJ). Reflectance measurements (**Fig.**

2) were converted to changes in total hemoglobin (Δ[HbT]), oxy-hemoglobin (Δ[HbO]), and deoxy-hemoglobin

(Δ[HbR]) using the Beer–Lambert law (Ma et al., 2016a, b). Correction for attenuation of GCaMP7s fluorescence due

to absorption of the surrounding tissue was corrected as previously described (Kramer, 1979; Ma, 2016b; Wright,

2017; Scott, 2018). Changes in fluorescence intensity in the GCaMP7s signal due to blood absorption were

approximated in a pixel-wise fashion by multiplying each value by the ratio of the green and blue channel's resting

baseline pixel value.

Electrophysiology. Gamma-band [30–100Hz] LFP band was digitally bandpass filtered from recorded broadband

data using a third-order Butterworth filter, squared, low-pass filtered below 1 Hz, and resampled at 30 Hz. Time-

frequency spectrograms (**Fig. 2**) were calculated using the Chronux toolbox version 2.12 v03 (Bokil, 2010), function

mtspecgramc with a 5 s window and 1/5 s step size using [5,9] tapers and a passband of 1–100 Hz to encompass the

LFP. EMG (300–3 kHz) from the neck muscles was bandpass filtered, squared, convolved with a Gaussian kernel with

a standard deviation of 0.5 seconds, log transformed, then resampled at 30 Hz. MATLAB function(s): butter, zp2sos,

filtfilt, gausswin, log10, conv, resample.

Evoked responses and slope. Evoked responses (**Fig. 3**) including whisker stimulation and locomotion for the various

data types ([HbT], [HbO], [HbR], GCaMP7s, arteriole diameter) were compared between the indicated intervals (i.e.,

2:4 seconds post-stimulation). The mean of the 2 seconds preceding event onsets were subtracted from the event

and smoothed with a 3rd order Savitzky-Golay filter. The average slope was calculated by comparing the rise (Δ[HbT])

over the run (neural data) before being fit with a linear model forced through the origin. MATLAB function(s):

sgolayfilt)

Spectral power and coherence. Spectral power (**Fig. 4, 5, 6 S5**) was estimated using the Chronux toolbox (Bokil,

2010) function mtspectrumc. Data was detrended within individual events and truncated to the desired length

depending on behavior (10 seconds for rest, 15 minutes for Alert and Asleep). Coherence analysis between two

signals within the same hemisphere or between signals recorded bilaterally was run for each data type Chronux

function coherencyc after detrending using the MATLAB function detrend.

Cross-correlation. Cross-correlations (**Fig. 5**) between gamma-band power or ΔF/F and changes in total hemoglobin

Δ[HbT] were taken during periods of resting-state, alert, and asleep. Data were mean-subtracted and digitally

lowpass filtered (< 1 Hz) with a fourth-order Butterworth filter (MATLAB function(s): butter, zp2sos, filtfilt). Cross-

correlation analysis was run for each arousal state (MATLAB function(s): xcorr) with a ± 5 second lag time.

Resting-state variance and Δ[HbT] during different arousal states. Variance during the resting-state for both Δ[HbT]

and diameter signals (**Fig. 7**) was taken from resting-state events lasting ≥10 seconds in duration. Average Δ[HbT] in

each arousal state was taken with respect to awake resting baseline events ≥10 seconds in duration. Continuous

NREM sleep events ≥30 seconds, REM sleep events ≥60 seconds, and periods following administration of isoflurane

were compared between groups. Each event was digitally lowpass filtered (<1 Hz) with a fourth-order Butterworth

filter (MATLAB function(s): butter, zp2sos, filtfilt) and then averaged within each individual time series prior to

comparing across animals/groups.

Open field behavior. Exploratory behavior (**Fig. S2**) was measured in a custom-made 30 cm x 60 cm arena during

the light phase. Mice were not habituated to the arena. Mice were placed in the arena for 10 minutes to explore,

and their behavior was recorded using an overhead camera (Blackfly BFLY-U3-23S6M, Teledyne FLIR, Wilsonville,

OR) with 6 mm fixed focal length lens (Edmund Optics, Barrington, NJ, Stock #33939) at a frame rate of either 15 or

30 frames per second. The arena was illuminated with 780nm light. The surroundings of the arena were dark during

the entire recording session. The arena was cleaned with 70% ethanol in between animals. Only the first 5 minutes

of the behavior were analyzed and reported here. The total distance traveled over the first 5 minutes and the time

spent in the 25 x 55 cm center rectangle were quantified. Generalized linear mixed effect (GLME) was used in

MATLAB to perform statistical analysis. No statistical outliers were removed from the data. Mouse behavior was

tracked using DeepLabCut (Mathis, 2018) and analyzed with custom MATLAB algorithms (Zhang, 2022; Brockway,

2023). Eight points on the body (left ear, right ear, head, mid, back near the hip joint, base of the tail, midpoint of

the tail, and end of the tail) were tracked. Four markers placed in the arena's four corners with an additional four

points calculated from the four corners to track the center rectangle. A DeepLabCut model was trained and

evaluated on a subset of mice before applying the model to all the mice. DeepLabCut tracking was considered

acceptable if the tracking confidence was above 97%; however, in most cases, it was higher than 99%. Tracking

positions were exported to a CSV file containing the tracked location's XY coordinates (frame pixels). Tracked videos

from random mice were visualized to confirm the accuracy of the tracking. The experimenter was blinded to drug

injection group identification until tracking and analysis were completed. The distance was calculated from the first

5 minutes as Euclidean distance between the point tracked in the middle of the mouse body from subsequent

frames. Center time was quantified as the number of frames a mouse spends within the polygon bounded by 4 points

tracked on the center of the arena.

Sleep scoring. Sleep scoring (**Fig. S3**) was performed as in Turner et al. (2020). Briefly, arousal state was scored using

a combination of cortical and hippocampal LFP, EMG, pupil diameter, and whisker and body moment. Periods of

manually-chosen awake rest 5s in duration with no vibrissae motion, and no body motion were used as a baseline

to normalize neural and hemodynamic signals. NREM sleep shows elevated cortical delta-band power and lower

EMG power. REM sleep is marked by elevated hippocampal theta-band power and elevated cortical gamma-band

power with very low baseline EMG power (muscle atonia; Cantero et al., 2004; Montgomery et al., 2008; Le Van

Quyen et al., 2010; Sullivan et al., 2014). Every 5 second interval was classified as either Awake, NREM sleep, or REM

sleep using a bootstrap aggregating random forest model with the predictors of cortical delta band power, cortical

beta band power, cortical gamma LFP, hippocampal theta LFP, EMG power, heart rate, and whisking duration. Sleep

model accuracy was validated using the out-of-bag error during model training. MATLAB functions used were

TreeBagger, oobError, predict.

Pupil diameter and interblink interval. Pupil diameter (**Fig. S2**) was lowpass filtered at 1 Hz with a fourth-order

Butterworth filter. Changes in whisking-evoked and stimulus-evoked (contralateral, auditory) diameters normalized

relative to the mean of the 2 second preceding the event onset. MATLAB functions used were butter, zp2sos, filtfilt.

Interblink interval and blink-associated physiology. Because of breaks in recording to save the data to disk, interblink

interval was calculated between blinks occurring within 15 min records and not blinks on the edges of trials. Blinks

that occurred within 1 second of each other were linked together as blinking bouts, and all blink-triggered analyses

were with respect to the first blink in a series. Blink-triggered averages were separated into two groups depending

on the arousal state classification of the 5 second bin before the blink, being either Awake (arousal state classification

of Awake) or Asleep (arousal state classification being either NREM or REM).

Hemodynamic response function. The hemodynamic response function (HRF, **Fig. S5**) was calculated as previously

described (Winder, 2017; Zhang, 2023) using both deconvolution and fitting the response to a gamma distribution.

Only neural activity (gamma-band power, 30-100 Hz) within 1.5 second of the stimulus was used for calculating the

HRF. Each HRF was calculated using half of the data and tested on the other half. To test the predictive capability of

each HRF, the impulse function was fit with a gamma-distribution function and convolved with the neural power to

predict the measured hemodynamic signal. The coefficient of determination (R^2) between the predicted and actual

hemodynamic signal was evaluated to quantify the efficacy of HRF's prediction (MATLAB function(s): sgolayfilt,

detrend, fminsearch, fitlm).

Pearson's correlation coefficient calculations. The Pearson's correlation coefficient between bilateral signals (**Fig.**

S6) was obtained by mean-subtracting and digitally lowpass filtering (<1 Hz) with a fourth-order Butterworth filter

and then taking the Pearson's correlation coefficient (MATLAB function(s): butter, zp2sos, filtfilt, corrcoef).

Arousal state transitions. Transitions between arousal states (Awake to NREM, NREM to Awaken, NREM to REM,

REM to Awake) were compared across groups (**Fig. S7**) by averaging events that met the criteria of 30 seconds of

one state's consecutive classifications followed by 30 seconds of another state. The difference in the Δ[HbT] between

the pre- and post- state transition were compared by taking the 30:10 seconds prior to the transition zero point to

the 10:30 seconds following. [HbT] was digitally lowpass (<1 Hz) filtered using a fourth-order Butterworth filter

(MATLAB function(s): butter, zp2sos, filtfilt).

Data availability. Data and code for generating the figures are available at [pending Dryad link once Manuscript

number is assigned] and analysis code is available at https://github.com/KL-Turner/Turner-nNOS-Manuscript. Raw

data is available upon reasonable request.

     Supplemental Figure 1

Figure S1. Histological quantification of cortical SP-Sap injections (a) Examples of NADPH diaphorase staining fromBlank-SAP and SP-SAP injected mice. **(b)** SP-SAP injected mice (N = 9, 5M/4F) had significantly lower numbers of type-I nNOS cells than either Blank-SAP (N = 9, 4M/5F) or Uninjected mice (N = 9, 4M/5F) (SP-SAP: 1.6 ± 0.8 neurons/mm^2; Blank-SAP: 22.9 ± 2.4 neurons/mm^2; Uninjected: 22.9 ± 3 neurons/mm^2; Blank-SAP vs. Uninjected: p = 0.98; Blank-SAP vs. SP-SAP: p = 2.23×10^{-13}). **(c)** Counts of DAPI-labeled cell nuclei per square mm from imaged mice. Uninjected (N = 6, 6M) mice had 2284 ± 279 DAPI-labeled cell bodies/mm^2, which was not significantly different than those injected with Blank-SAP (N = 5, 5F, 2488 ± 298, *p* = 0.23, GLME), which in turn was not significantly different than those injected with SP-SAP (N = 8, 4M/4F, 2340 ± 329, *p* = 0.39, GLME. **(d)** Representative image of IBA1 for Blank-SAP (top) and SP-SAP (bottom) taken from **Fig 1c**. Error bars (**b, c**) denote SD. *α < 0.05, **α < 0.01, ***α < 0.001, GLME.

 Supplemental Figure 2

Figure S2. Behavior was unaffected by local ablation of type-I nNOS neurons (a) Example of exploratory behavior during 5 minutes of exploration in a novel environment. **(b)** Total distance traveled during open field exploration. Uninjected mice traveled an average of 1473.4 ± 364.7 cm compared with 1536.6 ± 329.6 cm in Blank-SAP (p = 0.61) and 1652.3 ± 442.7 cm in SP-SAP (p = 0.43). **(c)** Time spent in the center of the open field. Uninjected mice spent on average 25.5 ± 13.1% of total exploration time in the center compared with 18.5 ± 5.3% in Blank-SAP (p = 0.08, GLME) and 21.9 ±9.8% in SP-SAP (p = 0.26). **(b, c)** n = 12-23 mice per group. **(d)** Percentage of time classified in each arousal state by sleep scoring. Uninjected mice spent 68.1 ± 13.2% of the time awake compared to 69.1 ± 15.1% in Blank-SAP (p = 0.89, ttest) and 73.4 ± 11.6% in SP-SAP (p = 0.51, ttest). Uninjected mice spent 28.3 ± 11.1% of the time in NREM sleep compared to 27.4 ± 12.0% in Blank-SAP (p = 0.87, ttest) and 23.4 ± 9.7% in SP-SAP (p = 0.44, ttest). Uninjected mice spent 3.5 ± 2.9% of the time in REM sleep compared to 3.4 ± 3.5% in Blank-SAP (p = 0.96, ttest) and 3.2 ± 2.3% in SP-SAP (p = 0.88, ttest). **(e)** Time spent whisking as a percentage of total imaging time. Uninjected mice spent 15.3 ± 9.9% compared with 17.3 ± 5.8% in Blank-SAP (p = 0.61, ttest) and 19.2 ± 8.1% in SP-SAP (p = 0.75, ttest). **(f)** Change in Pupil diameter in response to vibrissae stimulation. Pupil diameter of uninjected mice increased 1.3 ± 0.5 z-units following vibrissae stimulation compared with 1.2 ± 0.3 z-units in Blank-SAP (p = 0.76, ttest) and 1.1 ± 0.5 z-units in SP-SAP (p = 0.50, ttest). **(g)** Average pupil diameter across arousal states. Average pupil diameter of Uninjected mice during the resting-state was -0.25 ± 0.1 z-units compared with -0.29 ± 0.1 z-units in Blank-SAP (p = 0.45, ttest) and -0.34 ± 0.2 in SP-SAP (p = 0.14, ttest). Average pupil diameter of Uninjected mice during periods of NREM sleep was -3 ± 0.5 z-units compared with -3 ± 0.5 z-units in Blank-SAP (p = 0.89, ttest) and -3.4 ± 0.4 in SP-SAP (p = 0.08, ttest). Average pupil diameter of Uninjected mice during periods of REM sleep was -4.6 ± 0.8 z-units compared with -4.4 ± 0.8 z-units in Blank-SAP (p = 0.53, ttest) and -5 ± 0.6 in SP-SAP (p = 0.11, ttest) **(h)** Interblink-interval. Uninjected mice had a mean interblink-interval of 127.6 ± 46.8 s compared with 102.5 ± 34.7 s in Blank-SAP (p = 0.22, ttest) and 84.5 ± 23.3 s in SP-SAP (p = 0.21, ttest). **(d-h)** n = 9 mice per group. Shading **(f)** indicates SEM and error bars **(b, c, e, g, h)** denote SD. *α < 0.05, **α < 0.01, ***α < 0.001.

 Supplemental Figure 3

Figure S3. Sleep classification accuracy was unchanged following type-I nNOS ablation (a) Confusion matrix for arousal state classification of Uninjected mice with bilaterally implanted stereotrodes. **(b)** Confusion matrix for arousal state classification of Blank-SAP mice with bilaterally implanted stereotrodes. **(c)** Confusion matrix for arousal state classification of SP-SAP mice with bilaterally implanted stereotrodes. **(a-c)** n = 9 mice per group. **(d)** Confusion matrix for arousal state classification of Blank-SAP mice expressing pan-neuronal GCaMP. **(e)** Confusion matrix for arousal state classification of SP-SAP mice expressing pan-neuronal GCaMP. **(d, e)** n = 6-7 mice per group. **(f)** Out-of-bag error during training of each animal's bootstrapped random forest classification algorithm. For mice with bilateral LFP recordings, Uninjected mice had an average loss of 0.08 ± 0.03 in comparison to 0.07 ± 0.02 in Blank-SAP ($p = 0.45$, ttest) compared to 0.05 ± 0.02 in SP-SAP ($p = 0.064$, ttest). Blank-SAP mice with pan-neuronal GCaMP had an average classification loss of 0.06 ± 0.02 compared with 0.07 ± 0.02 in SP-SAP ($p = 0.80$, ttest). Error bars denote SD. $*\alpha < 0.05$, $**\alpha < 0.01$, $***\alpha < 0.001$.

Figure S4. Ablation of Type-I nNOS neurons does not alter hemodynamic or neural power spectra Power spectral density for vascular and hemodynamic signals for the Blank-SAP group (N = 9, 4M/5F for **a, b**; N = 7, 3M/4F for **c-f**) and SP-SAP group (N = 9, 5M/4F for **a, b;** N = 8, 4M/4F for **c-f**) was not significantly different across all measurements of hemodynamic and neural signals. **(a)** Δ[HbT] for electrophysiology animals (p = 0.77, GLME). **(b)** Gamma-band power (second spectra) for animals with electrophysiology (p = 0.19, GLME). **(c)** Δ[HbT] for animals with GCaMP (p = 0.27, GLME). **(d)** Δ[HbT] for animals with GCaMP (p = 0.26, GLME). **(e)** Δ[HbT] for animals with GCaMP (p = 0.68, GLME). **(f)** GCaMP fluorescence (p = 0.44, GLME). Shading represents population averages ± SEM. *α < 0.05, **α < 0.01, ***α < 0.001.

 Supplemental Figure 5

Figure S5. Ablation of Type-I nNOS neurons does not alter the predictive power of the hemodynamic response function (a) Stimulus-evoked hemodynamic response function obtained with deconvolution. **(b)** Stimulus-evoked hemodynamic response function fitted with a gamma distribution function. **(c)** There was no significant difference between the Δ[HbT] predictive R^2 values calculated from impulse-derived HRFs between Blank-SAP (N = 9, 4M/5F) 0.64 ± 0.11 or SP-SAP (N = 9, 5M/4F) 0.67 ± 0.16 following vibrissae stimulation (p = 0.746, ttest). **(d)** Resting-state hemodynamic response function based on deconvolution. **(e)** Resting-state hemodynamic response function fitted with a gamma distribution function. **(f)** There was no significant difference between the Δ[HbT] predictive R^2 values calculated from impulse-derived HRFs between Blank-SAP (N = 9, 4M/5F) 0.13 ± 0.0.05 or SP-SAP (N = 9, 5M/4F) 0.12 ± 0.05 during the resting-state (p = 0.749, ttest). Error bars denote SD. *$α$ < 0.05, **$α$ < 0.01, ***$α$ < 0.001.

 Supplemental Figure 6

Figure S6. Pearson's correlation coefficients between bilateral hemodynamic and neural signals (a) Correlation coefficient between bilateral hemodynamic signals during the resting-state was 0.75 ± 0.04 in Blank-SAP and 0.68 ± 0.06 in SP-SAP (ttest $p = 0.011$), during the alert state was 0.85 ± 0.03 with Blank-SAP and 0.81 ± 0.06 in SP-SAP (ttest $p = 0.09$), and during the asleep state was 0.95 ± 0.02 in Blank-SAP and 0.92 ± 0.03 in SP-SAP (ttest $p = 0.01$). **(b)** Correlation coefficient between bilateral gamma-band power signals during the resting-state was 0.15 ± 0.04 in Blank-SAP and 0.14 ± 0.12 in SP-SAP (ttest $p = 0.69$), during the alert state was 0.23 ± 0.08 in Blank-SAP and 0.19 ± 0.11 in SP-SAP (ttest $p = 0.31$), and during the asleep state was 0.50 ± 0.08 in Blank-SAP and 0.41 ± 0.08 in SP-SAP (ttest $p = 0.04$). **(c)** Correlation coefficient between bilateral hemodynamic signals (GCaMP7s group) during the resting-state was 0.82 ± 0.04 in Blank-SAP and 0.75 ± 0.04 in SP-SAP (ttest $p = 0.01$), during the alert state was 0.93 ± 0.02 in Blank-SAP and 0.91 ± 0.02 in SP-SAP (ttest $p = 0.08$), and during the asleep state was 0.98 ± 0.02 with in-SAP and 0.95 ± 0.03 in SP-SAP (ttest $p = 0.07$). **(d)** Correlation coefficient between bilateral GCaMP7s signals during the resting-state was 0.70 ± 0.05 in Blank-SAP and 0.65 ± 0.09 in SP-SAP (ttest $p = 0.16$), during the alert state was 0.93 ± 0.03 with Blank-SAP and 0.92 ± 0.03 in SP-SAP (ttest $p = 0.31$), and during the asleep state was 0.96 ± 0.02 with Blank-SAP and 0.95 ± 0.02 in SP-SAP (ttest $p = 0.37$). Error bars denote SD. $\alpha < 0.05$, $**\alpha < 0.01$, $***\alpha < 0.001$.

Figure S7. Removal of type-I nNOS neurons did not alter arousal state-related hemodynamic changes (a-c) Average Δ[HbT/O/R] during periods of NREM sleep and REM sleep in mice with pan-neuronal GCaMP (n = 6-7 mice per group). **(a)** NREM Δ[HbT] in Blank-SAP mice was 20.4 ± 6.4 μM compared to 16.5 ± 5.9 μM in SP-SAP (*p* = 0.27, ttest). REM Δ[HbT] in Blank-SAP mice was 80.0 ± 8.2 μM compared to 80.7 ± 5.9 μM in SP-SAP (*p* = 0.87, ttest). **(b)** NREM Δ[HbO] in Blank-SAP mice was 25.5 ± 6.1 μM compared to 21.2 ± 6.1 μM in SP-SAP (*p* = 0.23, ttest). REM Δ[HbO] in Blank-SAP mice was 90.8 ± 8.6 μM compared to 92.3 ± 5.5 μM in SP-SAP (*p* = 0.72, ttest). **(c)** NREM Δ[HbR] in Blank-SAP mice was -5.3 ± 1.1 μM compared to -4.9 ± 0.9 μM in SP-SAP (*p* = 0.50, ttest). REM Δ[HbR] in Blank-SAP mice was -11.4 ± 1.8 μM compared to -12.2 ± 1.5 μM in SP-SAP (*p* = 0.42, ttest). **(d)** Transition from Awake to NREM had a Δ[HbT] of -16.6 ± 6.4 μM in Blank-SAP mice and -11.1 ± 7.0 μM in SP-SAP mice (p < 0.08, GLME). **(e)** Transition from NREM to Awake had a Δ[HbT] of 21.5 ± 7.7 μM in Blank-SAP mice and 17.6 ± 9.5 μM in SP-SAP mice (p < 0.33, GLME). **(f)** Transition from NREM to REM had a Δ[HbT] of -17.6 ± 3.9 μM in Blank-SAP mice and -22.6 ± 7.9 μM in SP-SAP mice (p < 0.09, GLME). **(g)** Transition from REM to Awake had a Δ[HbT] of 67.8 ± 7.3 μM in Blank-SAP mice and 74.0 ± 13.1 μM in SP-SAP mice (p < 0.21, GLME). Error bars denote SD. Shading represents population averages ± SEM.*α < 0.05, **α < 0.01, ***α < 0.001.

585 **Supplemental Figure 8**

Figure S8. Schematic showing how non-linearity in the dilation response can explain coexistence strong dilation by activation of a pathway, and little change after weakening a pathway. If the diameter is a sublinear function of the sum of vasodilatory inputs, activation of all pathways will cause a dilation that is smaller than the sum of activation of each pathway individually. Loss of one pathway will not cause large changes, even though activation of that pathway in isolation can cause large dilations.

References

Abbott SB, Kanbar R, Bochorishvili G, Coates MB, Stornetta RL, Guyenet PG (2012) C1 neurons excite
locus coeruleus and A5 noradrenergic neurons along with sympathetic outflow in rats. J Physiol
590:2897-2915. PMC3448155 doi:10.1113/jphysiol.2012.232157

Ahn SJ, Anfray A, Anrather J, Iadecola C (2023) Calcium transients in nNOS neurons underlie distinct
phases of the neurovascular response to barrel cortex activation in awake mice. J Cereb Blood
Flow Metab:271678X231173175. doi:10.1177/0271678X231173175

Andrei AR, Debes S, Chelaru M, Liu X, Rodarte E, Spudich JL, Janz R, Dragoi V (2021) Heterogeneous
side effects of cortical inactivation in behaving animals. Elife 10. PMC8457825
doi:10.7554/eLife.66400

Bekar LK, Wei HS, Nedergaard M (2012) The locus coeruleus-norepinephrine network optimizes coupling
of cerebral blood volume with oxygen demand. Journal of Cerebral Blood Flow & Metabolism
32:2135-2145. doi:10.1038/jcbfm.2012.115

Bokil H, Andrews P, Kulkarni JE, Mehta S, Mitra PP (2010) Chronux: a platform for analyzing neural signals.
J Neurosci Methods 192:146-151. PMC2934871 doi:10.1016/j.jneumeth.2010.06.020

Bredt DS, Glatt CE, Hwang PM, Fotuhi M, Dawson TM, Snyder SH (1991) Nitric oxide synthase protein
and mRNA are discretely localized in neuronal populations of the mammalian CNS together with
NADPH diaphorase. Neuron 7:615-624. doi:10.1016/0896-6273(91)90374-9

Brockway DF, Griffith KR, Aloimonos CM, Clarity TT, Moyer JB, Smith GC, Dao NC, Hossain MS, Drew PJ,
Gordon JA, Kupferschmidt DA, Crowley NA (2023) Somatostatin peptide signaling dampens
cortical circuits and promotes exploratory behavior. Cell Rep 42:112976. PMC10542913
doi:10.1016/j.celrep.2023.112976

Bugeon S, Duffield J, Dipoppa M, Ritoux A, Prankerd I, Nicoloutsopoulos D, Orme D, Shinn M, Peng H,
Forrest H, Viduolyte A, Reddy CB, Isogai Y, Carandini M, Harris KD (2022) A transcriptomic axis
predicts state modulation of cortical interneurons. Nature 607:330-338. PMC9279161
doi:10.1038/s41586-022-04915-7

Cardin JA, Carlen M, Meletis K, Knoblich U, Zhang F, Deisseroth K, Tsai LH, Moore CI (2009) Driving fast-
spiking cells induces gamma rhythm and controls sensory responses. Nature 459:663-667.
PMC3655711 doi:10.1038/nature08002

Chan KY, Jang MJ, Yoo BB, Greenbaum A, Ravi N, Wu WL, Sanchez-Guardado L, Lois C, Mazmanian
SK, Deverman BE, Gradinaru V (2017) Engineered AAVs for efficient noninvasive gene delivery to
the central and peripheral nervous systems. Nat Neurosci 20:1172-1179. PMC5529245
doi:10.1038/nn.4593

Chong PS, Poon CH, Fung ML, Guan L, Steinbusch HWM, Chan YS, Lim WL, Lim LW (2019) Distribution
of neuronal nitric oxide synthase immunoreactivity in adult male Sprague-Dawley rat brain. Acta
Histochem 121:151437. doi:10.1016/j.acthis.2019.08.004

Claron J, Provansal M, Salardaine Q, Tissier P, Dizeux A, Deffieux T, Picaud S, Tanter M, Arcizet F, Pouget
P (2023) Co-variations of cerebral blood volume and single neurons discharge during resting state
and visual cognitive tasks in non-human primates. Cell Rep 42:112369.
doi:10.1016/j.celrep.2023.112369

Dana H, Sun Y, Mohar B, Hulse BK, Kerlin AM, Hasseman JP, Tsegaye G, Tsang A, Wong A, Patel R,
Macklin JJ, Chen Y, Konnerth A, Jayaraman V, Looger LL, Schreiter ER, Svoboda K, Kim DS
(2019) High-performance calcium sensors for imaging activity in neuronal populations and
microcompartments. Nat Methods 16:649-657. doi:10.1038/s41592-019-0435-6

Dao NC, Suresh Nair M, Magee SN, Moyer JB, Sendao V, Brockway DF, Crowley NA (2020) Forced
Abstinence From Alcohol Induces Sex-Specific Depression-Like Behavioral and Neural
Adaptations in Somatostatin Neurons in Cortical and Amygdalar Regions. Front Behav Neurosci
14:86. PMC7266989 doi:10.3389/fnbeh.2020.00086

Dittrich L, Morairty SR, Warrier DR, Kilduff TS (2015) Homeostatic sleep pressure is the primary factor for
activation of cortical nNOS/NK1 neurons. Neuropsychopharmacology 40:632-639. PMC4289951
doi:10.1038/npp.2014.212

Dittrich L, Heiss JE, Warrier DR, Perez XA, Quik M, Kilduff TS (2012) Cortical nNOS neurons co-express
the NK1 receptor and are depolarized by Substance P in multiple mammalian species. Front Neural
Circuits 6:31. PMC3367498 doi:10.3389/fncir.2012.00031

Drew PJ (2019a) Vascular and neural basis of the BOLD signal. Curr Opin Neurobiol 58:61-69.
PMC6859204 doi:10.1016/j.conb.2019.06.004

Drew PJ, Shih AY, Kleinfeld D (2011) Fluctuating and sensory-induced vasodynamics in rodent cortex
extend arteriole capacity. Proc Natl Acad Sci U S A 108:8473-8478. PMC3100929
doi:10.1073/pnas.1100428108

Drew PJ, Winder AT, Zhang Q (2019b) Twitches, Blinks, and Fidgets: Important Generators of Ongoing
Neural Activity. Neuroscientist 25:298-313. PMC6800083 doi:10.1177/1073858418805427

Drew PJ, Shih AY, Driscoll JD, Knutsen PM, Blinder P, Davalos D, Akassoglou K, Tsai PS, Kleinfeld D
(2010) Chronic optical access through a polished and reinforced thinned skull. Nat Methods 7:981-
984. PMC3204312 doi:10.1038/nmeth.1530

Echagarruga CT, Gheres KW, Norwood JN, Drew PJ (2020) nNOS-expressing interneurons control basal
and behaviorally evoked arterial dilation in somatosensory cortex of mice. Elife 9. PMC7556878
doi:10.7554/eLife.60533

Endo T, Yanagawa Y, Komatsu Y (2016) Substance P Activates Ca2+-Permeable Nonselective Cation
Channels through a Phosphatidylcholine-Specific Phospholipase C Signaling Pathway in nNOS-
Expressing GABAergic Neurons in Visual Cortex. Cereb Cortex 26:669-682.
doi:10.1093/cercor/bhu233

Gao YR, Greene SE, Drew PJ (2015) Mechanical restriction of intracortical vessel dilation by brain tissue
sculpts the hemodynamic response. Neuroimage 115:162-176. PMC4470397
doi:10.1016/j.neuroimage.2015.04.054

Gerashchenko D, Schmidt MA, Zielinski MR, Moore ME, Wisor JP (2018) Sleep State Dependence of
Optogenetically evoked Responses in Neuronal Nitric Oxide Synthase-positive Cells of the
Cerebral   Cortex.   Neuroscience   379:189-201.   PMC6311348
doi:10.1016/j.neuroscience.2018.02.006

Gerashchenko D, Wisor JP, Burns D, Reh RK, Shiromani PJ, Sakurai T, de la Iglesia HO, Kilduff TS (2008)
Identification of a population of sleep-active cerebral cortex neurons. Proc Natl Acad Sci U S A
105:10227-10232. PMC2481371 doi:10.1073/pnas.0803125105

Gheres KW, Unsal HS, Han X, Zhang Q, Turner KL, Zhang N, Drew PJ (2023) Arousal state transitions
occlude sensory-evoked neurovascular coupling in neonatal mice. Commun Biol 6:738.
PMC10352318 doi:10.1038/s42003-023-05121-5

Hablitz LM, Vinitsky HS, Sun Q, Staeger FF, Sigurdsson B, Mortensen KN, Lilius TO, Nedergaard M (2019)
Increased glymphatic influx is correlated with high EEG delta power and low heart rate in mice
under anesthesia. Sci Adv 5:eaav5447. PMC6392807 doi:10.1126/sciadv.aav5447

Han K, Min J, Lee M, Kang BM, Park T, Hahn J, Yei J, Lee J, Woo J, Lee CJ, Kim SG, Suh M (2019)
Neurovascular Coupling under Chronic Stress Is Modified by Altered GABAergic Interneuron
Activity. J Neurosci 39:10081-10095. PMC6978951 doi:10.1523/JNEUROSCI.1357-19.2019

Haselden WD, Kedarasetti RT, Drew PJ (2020) Spatial and temporal patterns of nitric oxide diffusion and
degradation drive emergent cerebrovascular dynamics. PLoS Comput Biol 16:e1008069.
PMC7410342 doi:10.1371/journal.pcbi.1008069

He M, Tucciarone J, Lee S, Nigro MJ, Kim Y, Levine JM, Kelly SM, Krugikov I, Wu P, Chen Y, Gong L, Hou
Y, Osten P, Rudy B, Huang ZJ (2016) Strategies and Tools for Combinatorial Targeting of
GABAergic Neurons in Mouse Cerebral Cortex. Neuron 91:1228-1243. PMC5223593
doi:10.1016/j.neuron.2016.08.021

Hendry SH, Schwark HD, Jones EG, Yan J (1987) Numbers and proportions of GABA-immunoreactive
neurons in different areas of monkey cerebral cortex. J Neurosci 7:1503-1519. PMC6568832
doi:10.1523/JNEUROSCI.07-05-01503.1987

Higo S, Udaka N, Tamamaki N (2007) Long-range GABAergic projection neurons in the cat neocortex. J
Comp Neurol 503:421-431. doi:10.1002/cne.21395

Hill CE (2012) Long Distance Conduction of Vasodilation: A Passive or Regenerative Process?
Microcirculation 19:379-390. doi:10.1111/j.1549-8719.2012.00169.x

Holstein-Ronsbo S, Gan Y, Giannetto MJ, Rasmussen MK, Sigurdsson B, Beinlich FRM, Rose L, Untiet V,
Hablitz LM, Kelley DH, Nedergaard M (2023) Glymphatic influx and clearance are accelerated by
neurovascular coupling. Nat Neurosci. doi:10.1038/s41593-023-01327-2

Hope BT, Vincent SR (1989) Histochemical characterization of neuronal NADPH-diaphorase. J Histochem
Cytochem 37:653-661. doi:10.1177/37.5.2703701

Hope BT, Michael GJ, Knigge KM, Vincent SR (1991) Neuronal NADPH diaphorase is a nitric oxide
synthase. Proc Natl Acad Sci U S A 88:2811-2814. PMC51329 doi:10.1073/pnas.88.7.2811

Hosford PS, Gourine AV (2019) What is the key mediator of the neurovascular coupling response? Neurosci
Biobehav Rev 96:174-181. PMC6331662 doi:10.1016/j.neubiorev.2018.11.011

Huang H, Kuzirian MS, Cai X, Snyder LM, Cohen J, Kaplan DH, Ross SE (2016) Generation of a NK1R-
CreER knockin mouse strain to study cells involved in Neurokinin 1 Receptor signaling. Genesis
54:593-601. PMC5241089 doi:10.1002/dvg.22985

Huo BX, Smith JB, Drew PJ (2014) Neurovascular coupling and decoupling in the cortex during voluntary
locomotion. J Neurosci 34:10975-10981. PMC4131012 doi:10.1523/JNEUROSCI.1369-14.2014

Huo BX, Gao YR, Drew PJ (2015) Quantitative separation of arterial and venous cerebral blood volume
increases   during   voluntary   locomotion.   Neuroimage   105:369-379.   PMC4330110
doi:10.1016/j.neuroimage.2014.10.030

Iadecola C, Smith EE, Anrather J, Gu C, Mishra A, Misra S, Perez-Pinzon MA, Shih AY, Sorond FA, van
Veluw SJ, Wellington CL, American Heart Association Stroke C, Council on Arteriosclerosis T,
Vascular B, Council on Cardiovascular R, Intervention, Council on H, Council on L, Cardiometabolic
H (2023) The Neurovasculome: Key Roles in Brain Health and Cognitive Impairment: A Scientific
Statement From the American Heart Association/American Stroke Association. Stroke.
doi:10.1161/STR.0000000000000431

Jiang-Xie LF, Drieu A, Bhasiin K, Quintero D, Smirnov I, Kipnis J (2024) Neuronal dynamics direct
cerebrospinal fluid perfusion and brain clearance. Nature 627:157-164. doi:10.1038/s41586-024-
07108-6

Kara P, Friedlander MJ (1999) Arginine analogs modify signal detection by neurons in the visual cortex. J
Neurosci 19:5528-5548. PMC6782344 doi:10.1523/JNEUROSCI.19-13-05528.1999

Karagiannis A, Gallopin T, David C, Battaglia D, Geoffroy H, Rossier J, Hillman EM, Staiger JF, Cauli B
(2009) Classification of NPY-expressing neocortical interneurons. J Neurosci 29:3642-3659.
PMC2750888 doi:10.1523/JNEUROSCI.0058-09.2009

Kawaguchi Y, Kubota Y (1997) GABAergic cell subtypes and their synaptic connections in rat frontal cortex.
Cereb Cortex 7:476-486. doi:10.1093/cercor/7.6.476

Kedarasetti RT, Drew PJ, Costanzo F (2022) Arterial vasodilation drives convective fluid flow in the brain:
a poroelastic model. Fluids Barriers CNS 19:34. PMC9107702 doi:10.1186/s12987-022-00326-y

Kilduff TS, Cauli B, Gerashchenko D (2011) Activation of cortical interneurons during sleep: an anatomical
link   to   homeostatic   sleep   regulation?   Trends   Neurosci   34:10-19.   PMC3014438
doi:10.1016/j.tins.2010.09.005

Kjaerby C, Andersen M, Hauglund N, Untiet V, Dall C, Sigurdsson B, Ding F, Feng J, Li Y, Weikop P, Hirase
H, Nedergaard M (2022) Memory-enhancing properties of sleep depend on the oscillatory
amplitude of norepinephrine. Nat Neurosci 25:1059-1070. PMC9817483 doi:10.1038/s41593-022-
01102-9

Knot HJ, Nelson MT (1998) Regulation of arterial diameter and wall [Ca2+] in cerebral arteries of rat by
membrane potential and intravascular pressure. The Journal of Physiology 508 (Pt 1:199-209.

Kramer RS, Pearlstein RD (1979) Cerebral cortical microfluorometry at isosbestic wavelengths for
correction of vascular artifact. Science 205:693-696. doi:10.1126/science.223243

Krawchuk MB, Ruff CF, Yang X, Ross SE, Vazquez AL (2020) Optogenetic assessment of VIP, PV, SOM
and NOS inhibitory neuron activity and cerebral blood flow regulation in mouse somato-sensory
cortex. J Cereb Blood Flow Metab 40:1427-1440. PMC7307010 doi:10.1177/0271678X19870105

Kress BT, Iliff JJ, Xia M, Wang M, Wei HS, Zeppenfeld D, Xie L, Kang H, Xu Q, Liew JA, Plog BA, Ding F,
Deane R, Nedergaard M (2014) Impairment of paravascular clearance pathways in the aging brain.
Annals of Neurology 76:845-861. doi:10.1002/ana.24271

Kubota Y, Shigematsu N, Karube F, Sekigawa A, Kato S, Yamaguchi N, Hirai Y, Morishima M, Kawaguchi
Y (2011) Selective coexpression of multiple chemical markers defines discrete populations of
neocortical GABAergic neurons. Cereb Cortex 21:1803-1817. doi:10.1093/cercor/bhq252

Lee L, Boorman L, Glendenning E, Christmas C, Sharp P, Redgrave P, Shabir O, Bracci E, Berwick J,
Howarth C (2020) Key Aspects of Neurovascular Control Mediated by Specific Populations of
Inhibitory Cortical Interneurons. Cereb Cortex 30:2452-2464. PMC7174996
doi:10.1093/cercor/bhz251

Li N, Chen S, Guo ZV, Chen H, Huo Y, Inagaki HK, Chen G, Davis C, Hansel D, Guo C, Svoboda K (2019)
Spatiotemporal constraints on optogenetic inactivation in cortical circuits. Elife 8. PMC6892606
doi:10.7554/eLife.48622

Ma Y, Shaik MA, Kozberg MG, Kim SH, Portes JP, Timerman D, Hillman EM (2016a) Resting-state
hemodynamics are spatiotemporally coupled to synchronized and symmetric neural activity in
excitatory neurons. Proc Natl Acad Sci U S A 113:E8463-E8471. PMC5206542
doi:10.1073/pnas.1525369113

Ma Y, Shaik MA, Kim SH, Kozberg MG, Thibodeaux DN, Zhao HT, Yu H, Hillman EM (2016b) Wide-field
optical mapping of neural activity and brain haemodynamics: considerations and novel approaches.
Philos Trans R Soc Lond B Biol Sci 371. PMC5003860 doi:10.1098/rstb.2015.0360

Martin JL, Sloviter RS (2001) Focal inhibitory interneuron loss and principal cell hyperexcitability in the rat
hippocampus after microinjection of a neurotoxic conjugate of saporin and a peptidase-resistant
analog of Substance P. The Journal of Comparative Neurology 436:127-152.
doi:10.1002/cne.1065

Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M (2018) DeepLabCut:
markerless pose estimation of user-defined body parts with deep learning. Nat Neurosci 21:1281-
1289. doi:10.1038/s41593-018-0209-y

Matrongolo MJ, Ang PS, Wu J, Jain A, Thackray JK, Reddy A, Sung CC, Barbet G, Hong Y-K, Tischfield
MA (2023) Piezo1 agonist restores meningeal lymphatic vessels, drainage, and brain-CSF
perfusion in craniosynostosis and aged mice. Journal of Clinical Investigation 134.
doi:10.1172/jci171468

Matsumura S, Yamamoto K, Nakaya Y, O'Hashi K, Kaneko K, Takei H, Tsuda H, Shirakawa T, Kobayashi
M (2021) Presynaptic NK1 Receptor Activation by Substance P Suppresses EPSCs via Nitric Oxide

Synthesis in the Rat Insular Cortex. Neuroscience 455:151-164.
doi:10.1016/j.neuroscience.2020.12.012

McKay LC, Feldman JL (2008) Unilateral ablation of pre-Botzinger complex disrupts breathing during sleep
but not wakefulness. Am J Respir Crit Care Med 178:89-95. PMC2441928
doi:10.1164/rccm.200712-1901OC

Morairty SR, Dittrich L, Pasumarthi RK, Valladao D, Heiss JE, Gerashchenko D, Kilduff TS (2013) A role
for cortical nNOS/NK1 neurons in coupling homeostatic sleep drive to EEG slow wave activity. Proc
Natl Acad Sci U S A 110:20272-20277. PMC3864296 doi:10.1073/pnas.1314762110

Osorio-Forero A, Cardis R, Vantomme G, Guillaume-Gentil A, Katsioudi G, Devenoges C, Fernandez LMJ,
Luthi A (2021) Noradrenergic circuit control of non-REM sleep substates. Curr Biol 31:5009-5023
e5007. doi:10.1016/j.cub.2021.09.041

Paul A, Crow M, Raudales R, He M, Gillis J, Huang ZJ (2017) Transcriptional Architecture of Synaptic
Communication Delineates GABAergic Neuron Identity. Cell 171:522-539 e520. PMC5772785
doi:10.1016/j.cell.2017.08.032

Penny GR, Afsharpour S, Kitai ST (1986) Substance P-immunoreactive neurons in the neocortex of the rat:
A subset of the glutamic acid decarboxylase-immunoreactive neurons. Neuroscience Letters
65:53-59. doi:10.1016/0304-3940(86)90119-9

Perrenoud Q, Geoffroy H, Gauthier B, Rancillac A, Alfonsi F, Kessaris N, Rossier J, Vitalis T, Gallopin T
(2012) Characterization of Type I and Type II nNOS-Expressing Interneurons in the Barrel Cortex
of Mouse. Front Neural Circuits 6:36. PMC3386492 doi:10.3389/fncir.2012.00036

Pfeffer CK, Xue M, He M, Huang ZJ, Scanziani M (2013) Inhibition of inhibition in visual cortex: the logic of
connections between molecularly distinct interneurons. Nat Neurosci 16:1068-1076. PMC3729586
doi:10.1038/nn.3446

Phillips EA, Hasenstaub AR (2016) Asymmetric effects of activating and inactivating cortical interneurons.
Elife 5. PMC5123863 doi:10.7554/eLife.18383

Pleil KE, Rinker JA, Lowery-Gionta EG, Mazzone CM, McCall NM, Kendra AM, Olson DP, Lowell BB, Grant
KA, Thiele TE, Kash TL (2015) NPY signaling inhibits extended amygdala CRF neurons to
suppress binge alcohol drinking. Nat Neurosci 18:545-552. PMC4376619 doi:10.1038/nn.3972

Roth KA, Katz RJ (1979) Stress, behavioral arousal, and open field activity--a reexamination of emotionality
in the rat. Neurosci Biobehav Rev 3:247-263. doi:10.1016/0149-7634(79)90012-5

Ruff CF, Juarez Anaya F, Dienel SJ, Rakymzhan A, Altamirano-Espinoza A, Couey JJ, Fukuda M, Watson
AM, Su A, Fish KN, Rubio ME, Hooks BM, Ross SE, Vazquez AL (2024) Long-range inhibitory
neurons mediate cortical neurovascular coupling. Cell Rep 43:113970.
doi:10.1016/j.celrep.2024.113970

Sahara S, Yanagawa Y, O'Leary DD, Stevens CF (2012) The fraction of cortical GABAergic neurons is
constant from near the start of cortical neurogenesis to adulthood. J Neurosci 32:4755-4761.
PMC3325497 doi:10.1523/JNEUROSCI.6412-11.2012

Schaeffer S, Iadecola C (2021) Revisiting the neurovascular unit. Nat Neurosci 24:1198-1209.
doi:10.1038/s41593-021-00904-7

Scherer-Singler U, Vincent SR, Kimura H, McGeer EG (1983) Demonstration of a unique population of
neurons with NADPH-diaphorase histochemistry. J Neurosci Methods 9:229-234.
doi:10.1016/0165-0270(83)90085-7

Scott BB, Thiberge SY, Guo C, Tervo DGR, Brody CD, Karpova AY, Tank DW (2018) Imaging Cortical
Dynamics in GCaMP Transgenic Rats with a Head-Mounted Widefield Macroscope. Neuron
100:1045-1058 e1045. PMC6283673 doi:10.1016/j.neuron.2018.09.050

Seibenhener ML, Wooten MC (2015) Use of the Open Field Maze to measure locomotor and anxiety-like
behavior in mice. J Vis Exp:e52434. PMC4354627 doi:10.3791/52434

Shih AY, Mateo C, Drew PJ, Tsai PS, Kleinfeld D (2012) A Polished and Reinforced Thinned-skull Window
for Long-term Imaging of the Mouse Brain. Journal of Visualized Experiments. doi:10.3791/3742-
v

Sicher AR, Starnes WD, Griffith KR, Dao NC, Smith GC, Brockway DF, Crowley NA (2023) Adolescent
binge drinking leads to long-lasting changes in cortical microcircuits in mice. Neuropharmacology
234:109561. PMC10386078 doi:10.1016/j.neuropharm.2023.109561

Sirotin YB, Das A (2009) Anticipatory haemodynamic signals in sensory cortex not predicted by local
neuronal activity. Nature 457:475-479. PMC2705195 doi:10.1038/nature07664

Smith RJ, Anderson RI, Haun HL, Mulholland PJ, Griffin WC, 3rd, Lopez MF, Becker HC (2020) Dynamic
c-Fos changes in mouse brain during acute and protracted withdrawal from chronic intermittent
ethanol exposure and relapse drinking. Addict Biol 25:e12804. PMC7579841
doi:10.1111/adb.12804

Smith SL, Otis TS (2003) Persistent changes in spontaneous firing of Purkinje neurons triggered by the
nitric oxide signaling cascade. J Neurosci 23:367-372.

Sohal VS, Zhang F, Yizhar O, Deisseroth K (2009) Parvalbumin neurons and gamma rhythms enhance
cortical circuit performance. Nature 459:698-702. PMC3969859 doi:10.1038/nature07991

Staiger JF, Petersen CCH (2021) Neuronal Circuits in Barrel Cortex for Whisker Sensory Perception.
Physiol Rev 101:353-415. doi:10.1152/physrev.00019.2019

Stringer C, Pachitariu M, Steinmetz N, Reddy CB, Carandini M, Harris KD (2019) Spontaneous behaviors
drive multidimensional, brainwide activity. Science 364:255. PMC6525101
doi:10.1126/science.aav7893

Tomioka R, Rockland KS (2007) Long-distance corticocortical GABAergic neurons in the adult monkey
white and gray matter. J Comp Neurol 505:526-538. doi:10.1002/cne.21504

Tomioka R, Okamoto K, Furuta T, Fujiyama F, Iwasato T, Yanagawa Y, Obata K, Kaneko T, Tamamaki N
(2005) Demonstration of long-range GABAergic connections distributed throughout the mouse
neocortex. Eur J Neurosci 21:1587-1600. doi:10.1111/j.1460-9568.2005.03989.x

Tricoire L, Vitalis T (2012) Neuronal nitric oxide synthase expressing neurons: a journey from birth to
neuronal circuits. Front Neural Circuits 6:82. PMC3514612 doi:10.3389/fncir.2012.00082

Tu W, Cramer SR, Zhang N (2024) Disparity in temporal and spatial relationships between resting-state
electrophysiological and fMRI signals. Elife 13. PMC11299978 doi:10.7554/eLife.95680

Turner KL, Gheres KW, Drew PJ (2023) Relating Pupil Diameter and Blinking to Cortical Activity and
Hemodynamics   across   Arousal   States.   J   Neurosci   43:949-964.   PMC9908322
doi:10.1523/JNEUROSCI.1244-22.2022

Turner KL, Gheres KW, Proctor EA, Drew PJ (2020) Neurovascular coupling and bilateral connectivity
during NREM and REM sleep. Elife 9. PMC7758068 doi:10.7554/eLife.62071

Turrigiano GG (2008) The self-tuning neuron: synaptic scaling of excitatory synapses. Cell 135:422-435.
PMC2834419 doi:10.1016/j.cell.2008.10.008

Uhlirova H et al. (2016) Cell type specificity of neurovascular coupling in cerebral cortex. Elife 5.
PMC4933561 doi:10.7554/eLife.14315

van Veluw SJ, Hou SS, Calvo-Rodriguez M, Arbel-Ornath M, Snyder AC, Frosch MP, Greenberg SM,
Bacskai BJ (2020) Vasomotion as a Driving Force for Paravascular Clearance in the Awake Mouse
Brain. Neuron 105:549-561 e545. PMC7028316 doi:10.1016/j.neuron.2019.10.033

Vanlandewijck M, He L, Mae MA, Andrae J, Ando K, Del Gaudio F, Nahar K, Lebouvier T, Lavina B, Gouveia
L, Sun Y, Raschperger E, Rasanen M, Zarb Y, Mochizuki N, Keller A, Lendahl U, Betsholtz C (2018)
A molecular atlas of cell types and zonation in the brain vasculature. Nature 554:475-480.
doi:10.1038/nature25739

Vo TT, Im GH, Han K, Suh M, Drew PJ, Kim SG (2023) Parvalbumin interneuron activity drives fast
inhibition-induced vasoconstriction followed by slow substance P-mediated vasodilation. Proc Natl
Acad Sci U S A 120:e2220777120. PMC10161000 doi:10.1073/pnas.2220777120

Vruwink M, Schmidt HH, Weinberg RJ, Burette A (2001) Substance P and nitric oxide signaling in cerebral
cortex: anatomical evidence for reciprocal signaling between two classes of interneurons. J Comp
Neurol 441:288-301. doi:10.1002/cne.1413

Wang H, Germanson TP, Guyenet PG (2002) Depressor and tachypneic responses to chemical stimulation
of the ventral respiratory group are reduced by ablation of neurokinin-1 receptor-expressing
neurons. J Neurosci 22:3755-3764. PMC6758381 doi:10.1523/JNEUROSCI.22-09-03755.2002

Wiley RG, Lappi DA (1999) Targeting neurokinin-1 receptor-expressing neurons with [Sar9,Met(O2)11
substance P-saporin. Neurosci Lett 277:1-4. doi:10.1016/s0304-3940(99)00846-0

Winder AT, Echagarruga C, Zhang Q, Drew PJ (2017) Weak correlations between hemodynamic signals
and ongoing neural activity during the resting state. Nat Neurosci 20:1761-1769. PMC5816345
doi:10.1038/s41593-017-0007-y

Wölfle SE, Chaston DJ, Goto K, Sandow SL, Edwards FR, Hill CE (2011) Non-linear relationship between
hyperpolarisation and relaxation enables long distance propagation of vasodilatation. The Journal
of physiology 589:2607-2623. doi:10.1113/jphysiol.2010.202580

Wright PW, Brier LM, Bauer AQ, Baxter GA, Kraft AW, Reisman MD, Bice AR, Snyder AZ, Lee JM, Culver
JP (2017) Functional connectivity structure of cortical calcium dynamics in anesthetized and awake
mice. PLoS One 12:e0185759. PMC5648115 doi:10.1371/journal.pone.0185759

Wu YT, Bennett HC, Chon U, Vanselow DJ, Zhang Q, Munoz-Castaneda R, Cheng KC, Osten P, Drew PJ,
Kim Y (2022) Quantitative relationship between cerebrovascular network and neuronal cell types
in mice. Cell Rep 39:110978. PMC9271215 doi:10.1016/j.celrep.2022.110978

Yan XX, Garey LJ (1997) Morphological diversity of nitric oxide synthesising neurons in mammalian
cerebral cortex. J Hirnforsch 38:165-172.

Yan XX, Jen LS, Garey LJ (1996) NADPH-diaphorase-positive neurons in primate cerebral cortex
colocalize   with   GABA   and   calcium-binding   proteins.   Cereb   Cortex   6:524-529.
doi:10.1093/cercor/6.3.524

Zhang Q, Haselden WD, Charpak S, Drew PJ (2023) Could respiration-driven blood oxygen changes
modulate neural activity? Pflugers Arch 475:37-48. PMC9794637 doi:10.1007/s00424-022-02721-
8

Zhang Q, Turner KL, Gheres KW, Hossain MS, Drew PJ (2022) Behavioral and physiological monitoring
for awake neurovascular coupling experiments: a how-to guide. Neurophotonics 9:021905.
PMC8802326 doi:10.1117/1.NPh.9.2.021905

Zhang Q, Roche M, Gheres KW, Chaigneau E, Kedarasetti RT, Haselden WD, Charpak S, Drew PJ (2019)
Cerebral oxygenation during locomotion is modulated by respiration. Nat Commun 10:5515.
PMC6893036 doi:10.1038/s41467-019-13523-5

Zhao J, Manza P, Wiers C, Song H, Zhuang P, Gu J, Shi Y, Wang G-J, He D (2020) Age-Related Decreases
in Interhemispheric Resting-State Functional Connectivity and Their Relationship With Executive
Function. Frontiers in Aging Neuroscience 12. doi:10.3389/fnagi.2020.00020

Zielinski MR, Atochin DN, McNally JM, McKenna JT, Huang PL, Strecker RE, Gerashchenko D (2019)
Somatostatin+/nNOS+ neurons are involved in delta electroencephalogram activity and cortical-
dependent recognition memory. Sleep 42. PMC6783898 doi:10.1093/sleep/zsz143