Mapping brain function underlying naturalistic motor observation and imitation using high-density diffuse optical tomography

Authors: Dalin Yang[1], Tessa G. George[1], Chloe M. Sobolewski[1], Sophia R. McMorrow[1], Carolina Pacheco[2], Kelsey T. King[1], Rebecca Rochowiak[3], Evan Daniels-Day[1], Sungmin Park[1], Emma Speh[1], Daniel Lidstone[3], Romila Santra[4], Deana Crocetti[3], Mary Beth Nebel[3], Bahar Tuncgenc[5], Rene Vidal[6], Natasha M. Marrus[7], Stewart H. Mostofsky[3,8,9] Adam T. Eggebrecht[1,10,11,12,13,14],

[1]*Washington University School of Medicine, Mallinckrodt Institute of Radiology, St. Louis, Missouri 63110, USA.* [2]*Johns Hopkins University, Department of Biomedical Engineering, Baltimore, Maryland 21218, USA.* [3]*Kennedy Krieger Institute, Center for Neurodevelopment and Imaging Research, Baltimore, Maryland 21205, USA.* [4]*Harvard Medical School, Boston, Massachusetts 02115, USA.* [5]*Nottingham Trent University, Department of Psychology, Nottingham NG1 4FQ, United Kingdom.* [6]*University of Pennsylvania, Perelman School of Medicine, Department of Radiology, Philadelphia, Pennsylvania 19104, USA.* [7]*Washington University School of Medicine, Department of Psychiatry, St. Louis, Missouri, 63110, USA.* [8]*Johns Hopkins University School of Medicine, Department of Neurology, Baltimore, Maryland 21287, USA.* [9]*Johns Hopkins University School of Medicine, Department of Psychiatry and Behavioral Sciences, Baltimore, Maryland 21287, USA.* [10]*Washington University School of Engineering, Department of Biomedical Engineering, St. Louis, Missouri 63130, USA.* [11]*Washington University School of Medicine, Division of Biology and Biomedical Sciences, St. Louis, Missouri 63110, USA.* [12]*Washington University School of Arts and Science, Department of Physics, St. Louis, Missouri 63130, USA.* [13]*Washington University School of Engineering, Department of Electrical and System Engineering, St. Louis, Missouri, 63112, USA.* [14]*Washington University School of Engineering, Department Imaging Sciences Engineering, St. Louis, Missouri, 63112, USA.*

Corresponding author: Adam T Eggebrecht, aeggebre@wustl.edu

Abstract

Background: While autism spectrum disorder (ASD) is defined by deficits in social communication with restricted interests and repetitive behaviors, autistic individuals often show early impairments in motor imitation that persist through childhood and into adulthood. High-density diffuse optical tomography (HD-DOT) overcomes logistical challenges of functional magnetic resonance imaging (fMRI) to provide an open scanning environment conducive to neuroimaging during naturalistic motor imitation. Additionally, the mirror neuron system (MNS) is crucial for understanding and imitating actions, and its dysfunction is hypothesized to underlie key ASD features.

Objective: We aim to investigate brain function underlying motor observation and motor imitation in adult autistic and non-autistic individuals (NAI). We hypothesize that HD-DOT will reveal greater MNS activity during motor imitation than motor observation, and that MNS activity will exhibit a negative correlation with autistic traits.

Methods: We imaged brain function using HD-DOT in N=100 participants as they passively observed videos of an actor completing sequences of meaningless arm movements. Additionally, while being simultaneously recorded with both HD-DOT and Kinect 3D cameras for computer-vision-based assessment of motor imitation (CAMI), participants imitated different videos of an actor completing similar arm movements. Responses to the tasks were estimated using general linear models, and multiple regression was used to investigate brain-behavior associations with autistic traits, using the Social Responsiveness Scale, and imitation fidelity as measured with CAMI.

Results: Both motor observation and imitation tasks elicited significant activations in visual, temporal, and MNS areas, with imitation showing stronger activation in motor regions. Notably, MNS regions exhibited greater activation during observation than imitation. Additionally, activity during observation in the right parietal lobe correlated with autistic traits assessed with the SRS.

Conclusions: Our findings provide robust evidence of shared and task-specific neural mechanisms underlying motor observation and imitation, emphasizing the differential engagement of MNS regions during motor observation and imitation.

Key words: Autism spectrum disorder (ASD); High-density diffuse optical tomography (HD-DOT); Motor observation (OBS), Motor imitation (IM), and Computer-vision-based assessment of motor imitation (CAMI)

Introduction

Autism spectrum disorder (ASD) is a neurodevelopmental condition traditionally characterized by deficits in social communication and enhanced presentation of restricted interests and repetitive behaviors, with characteristically heterogeneous traits across the population[1,2]. Difficulty in motor skills is highly prevalent, with recent estimates suggesting that up to 88% of children with ASD experience motor challenges[3-8]. Impairments in motor imitation are particularly notable, as imitation is essential for early prosocial and language development[9,10]. Imitating across a range of behaviors, e.g., vocalizations, arm gestures, facial expressions, and other non-verbal cues, is a key component for effective social interaction[11]. Thus, early deficits in imitation can hinder social development, contributing to difficulties in social engagement and communication[12,13]. Additionally, impaired motor imitation may be related to disrupted visual-motor integration in the brain [14], and further may be related to the observation that ASD individuals often rely more on proprioceptive inputs than visual cues compared to their typically developing peers[15,16]. While motor imitation deficits in ASD are well-documented, their underlying neural mechanisms remain poorly understood [17], in part due to the physical constraints and limitations of neuroimaging modalities like functional magnetic resonance imaging (fMRI) [18,19]. Specifically, the MRI environment requires participants to lay supine in a small bore that inherently restricts studies to small-scale movements, such as those of fingers or hands, and fail to capture brain function during more general and naturalistic motor behaviors. In contrast, high-density diffuse optical tomography (HD-DOT) provides a naturalistic and open scanning environment that enables the assessment of brain function across the lifespan, such as healthy adults, infants, and school-age autistic children[20-24], and even in bedside clinical settings such as the intensive care unit[24]. Moreover, HD-DOT is less sensitive to motion artifact, and provides fMRI comparable spatial resolution and anatomical specificity over a broad field of view, capturing activity in sensorimotor systems implicated in viewing and performing motor movements[25-27].

Extant neuroimaging studies suggest that the mirror neuron system (MNS) plays a crucial role in observational learning and the promotion of imitation in the general population[28-31]. This system becomes activated not only when an individual performs a motor action, but also when they observe someone else performing the same action[32-35], highlighting the importance of the MNS in bridging the gap between observation and imitation, enabling individuals to learn and replicate actions by simply watching others[28,29,31,32,36-38]. The MNS has been shown to include brain regions such as the premotor cortex, inferior parietal lobule, and inferior frontal gyrus[32,37-41], as well as the pre-supplementary and primary motor area[34,42]. Additionally, differential activity within the MNS between typically developing individuals and those with ASD have been observed during motor imitation and action observation[31,40,43-49]. Variability in MNS activations has been shown to correlate with autistic traits[17], suggesting that the MNS may underlie key behavioral variability associated with ASD. Furthermore, dynamic interactions between the ventral attention network (VAN) and dorsal attention network (DAN) support sensory processing and action planning [50-54] - critical for imitation and motor coordination. Therefore, impairments in the MNS,

VAN, and DAN may help explain the deficits in imitation and related behaviors seen in individuals
with ASD.

Early detection of challenges with motor imitation could bolster early autism diagnostic
assessments[55]. However, human-observation coding (HOC) schemes to assess motor imitation
fidelity are highly time consuming and require extensive training. To enable automated computer-
driven assessment of the three-dimensional motor output of motor imitation, members of our team
have developed an automatic, continuous, computerized assessment of motor imitation (CAMI)
that utilizes metric learning and dynamic time warping to assess motor imitation fidelity[56]. This
method uses dynamic time warping (DTW) to assess the spatiotemporal differences between the
movements of a participant and of a gold standard[57]. The CAMI method has been validated as a
reliable method for assessing motor imitation in children [56,58]. This work has shown that CAMI-
generated scores were highly correlated with those generated using traditional labor-intensive,
subjective HOC methods. Additionally, the CAMI had better discriminative ability to classify the
children into ASD vs non-autistic groups than did HOC, and CAMI scores were significantly
correlated with autism trait severity, as measured using the Social Responsiveness Scale, Version
2 (SRS-2)[59], a well-established clinical metric of quantitative autistic traits. While these CAMI
methods show promise in detecting group differences in motor imitation fidelity, the relationships
between variability in brain function and motor imitation fidelity remain unknown.

Herein, we aim to use HD-DOT and simultaneous CAMI to investigate how variability in
brain function relates to both observation of and imitation of purposeless gross motor movements
in a sample of adults, including those with ASD. We hypothesize that both observing and imitating
will engage visual, motor, and temporal brain regions. Additionally, we hypothesize that regions
within the MNS, DAN, and VAN will be activated when observing and imitating motor
movements, with stronger brain activation within key regions of the DAN and MNS during motor
imitation. Last, we hypothesize that brain response strength within the MNS will be negatively
correlated with dimensional measures of social responsiveness (i.e., Social Responsiveness Scale,
SRS-2) during either observation or imitation, and positively correlated with imitation fidelity (i.e.,
CAMI scores) during motor imitation.

Material and Methods

Participants

We recruited N=100 participants (age 30.15 ± 12.97 years, 45 male/55 female, **Table 1**)
from the St. Louis, Missouri metropolitan community. All participants gave written informed
consent to participate in this study, which was approved and carried out in accordance with the
Human Research Protection Office at Washington University School of Medicine. All participants
with ASD had a prior community diagnosis.

HD-DOT System

The continuous-wave HD-DOT system has been previously described [60] and was
comprised of 128 dual-wavelength optical sources at wavelengths of 685 nm and 830 nm, and 125
avalanche-photodiode detectors. The imaging cap supported multiple source-detector distances
(i.e., 11, 24, 33, and 40 mm), providing over 3,500 measurement pairs per wavelength for image
reconstruction at a 10 Hz frame rate. The HD-DOT field-of-view covered continuous aspects of
temporal, visual, parietal, and prefrontal cortical areas (**Figure 1A,B**).

Stimuli

The motor observation and motor imitation paradigms each consisted of six different
movement sequences separated by 30 second inter-stimulus-intervals (ISIs; **Figure 1C**). The
movement sequences lasted 16-21 seconds and were comprised of 14-18 individual movement
types, which were unfamiliar, did not have an end goal, and required moving both arms
simultaneously. These movements were chosen based on prior research indicating difficulties in
these movements for autistic participants[56]. All movements were performed by the same woman
and the stimuli were displayed on a monitor placed 110 cm from the nasion in front of the
participant. For the observation task, participants were instructed to passively watch the movement
sequences and to remain as still as possible. For the motor imitation task, participants were
instructed to mirror the actor's arm movements as accurately as possible and limit head motion
(**Figure 1D**). Stimulus paradigms were presented via Psychophysics toolbox 3 for MATLAB using
in-house custom-built scripts. Participant brain function was recorded with HD-DOT during all
experimental runs. During the motor imitation runs, three-dimensional spatial tracking of
participants' movements were also simultaneously recorded using an Xbox Kinect 3D camera
located on top of the stimulus presentation monitor.

Computerized Assessment of Motor Imitation (CAMI)

The general CAMI algorithm has been previously described[56]. The CAMI algorithm was
modified for a seated imitation task to allow for the imitation task to be used while participants
were in our fiber-based HD-DOT system. The seated CAMI algorithm only considering joints
from the upper body as there was limited involvement of the lower limbs. The spatial coordinates
of 20 joints were extracted from the Xbox Kinect 3D motion sensor depth recordings using iPi
Motion Capture Software (iPiSoft, Moscow, Russia). Calculation of CAMI scores consisted of
five main steps: (1) preprocessing, (2) automatic joint importance estimation, (3) computation of
the distance feature, (4) computation of the time features, and (5) computation of the CAMI score.
Preprocessing consisted of defining the hip position as the origin, mapping the sequence of
movements to a skeleton of fixed size, correcting the initial orientation of the participants to match
that of the avatar (**Figure 1E**), the average position of the participants' joints associated with their
spine (Hip, Lower Spine, Middle Spine, and Neck) were corrected to match those of the average
position of the gold standard (**Figure 1F**). This alignment step was completed to avoid penalizing
participants for their initial posture or sitting position in the cap, as this could vary between

participants. Dynamic time warping was utilized to assess the spatial-temporal differences
between 20 joints of the participant and of a gold standard expert imitator. The features were
linearly combined to compute the overall CAMI score using previously published weights [56].

Behavioral Measurements

The Social Responsiveness Scale, 2^{nd} Edition (SRS-2)[61], provided a measure of social
reciprocity, with higher scores correspondent to higher levels of autism traits that have been shown
to be continuously distributed throughout the general population. The SRS raw scores were
converted to standardized SRS t-scores for analyses across a general population to avoid bias for
age or sex. Participant handedness was determined by the Edinburgh Handedness Inventory (EHI)
[62]. Measures of verbal and non-verbal intelligence quotients (IQ) were assessed with the Kaufman
Brief Intelligence Test- 2 (KBIT-2)[63]. All participants had a full–scale or verbal IQ > 80.

Data pre-processing and Image Reconstruction

HD-DOT data were pre-processed using the NeuroDOT toolbox in MATLAB
(https://www.nitrc.org/projects/neurodot). Each of the raw source-detector pair light level
measurements were converted to optical density by calculating the log-ratio of the raw data to its
temporal mean. Measurements with greater than a 7.5% temporal standard deviation in the least
noisy (lowest mean motion) 60s of each run were removed from further analyses. Data were
bandpass-filtered between 0.02 and 1.0 Hz using 9-pole Butterworth filters. Then, the superficial
signal for each wavelength was estimated as the average across the closest nearest neighbor
measurements (11 mm S-D pair separation) and regressed from all measurement channels. The
optical density time-traces were then low pass filtered with a cutoff frequency of 0.5 Hz and
temporally down sampled from to 1 Hz. As previously described[23], a wavelength-dependent
sensitivity matrix was computed and inverted to calculate relative changes in absorption at the two
wavelengths via reconstruction using a Tikhonov regularization parameter of 0.01 and spatially
variant regularization parameter of 0.1. Relative changes in the concentrations of oxygenated,
deoxygenated, and total hemoglobin were obtained from the absorption coefficient changes by the
spectral decomposition of the extinction coefficients of oxygenated and deoxygenated hemoglobin
at the two wavelengths (i.e., 685 and 830 nm). All data were resampled to a $1\times1\times1$ **mm³** standard
MNI [64,65] atlas using a linear affine transformation for statistical analyses.

Data Quality Assessment and Motion Censoring

Quantitative raw data quality assessments included calculating the percentage of good
measurements (GM) retained[23] and the median pulse band signal-to-noise ratio (SNR)[23,66]. As
previously described, runs with GM < 80% and/or SNR < 1.0 and were excluded from further
analysis [23,66,67] to ensure robust data reliability. Global variance in the temporal derivative (GVTD)
was used to detect and censor time points with motion-driven artifact in the data[23,27,68]. Briefly,
GVTD is calculated as the RMS of the first temporal derivative across the set of measurements for

each time point. Higher GVTD values indicate higher global variance in the optical time course
and are exquisitely sensitive to motion-based changes in optode-scalp coupling. Motion censoring
with GVTD excludes the time points that exceed the GVTD noise threshold. We used a stringent
GVTD threshold of 0.1% based on previous studies utilizing GVTD motion based censoring[23,66,67].

Statistical Analyses

After pre-processing and data quality assessment, 78 participants were identified as having
high-quality motor observation runs, while 64 participants had high-quality imitation runs that met
the data quality assessment thresholds described in the previous section. Additionally, one
participant was excluded due to technical issues with the stimulus presentation. Further, nine
participants did not return SRS scores and 21 participants were unable to finalize K-BIT testing
due to scheduling difficulties. As a result, the final sample consisted of 77 participants for motor
observation analysis and 63 participants for motor imitation analysis.

We used general linear modeling (GLM) to estimate the brain response to each task (task
vs. rest) using a canonical hemodynamic responses function (HRF) based on HD-DOT data[69]. We
present the HbO results in the main figures as HbO signals exhibits a higher contrast-to-noise ratio
compared to HbR or HbT [20,68,70].

We used the Gordon cortical parcellation [71] to provide an unbiased spatial sampling for
hypothesis driven analyses. The functionally defined Gordon parcellation provided a framework
to couch our analyses within functionally identified parcels of interest. The intersection of each
Gordon parcel with the HD-DOT field of view provided 192 parcels associated with nine
functional networks (i.e., auditory, frontal-parietal, default mode, dorsal-attention, somatosensory
hand, somatosensory mouth, ventral attention, cingulo-opercular, and visual functional networks;
Figure 1B). Small parcels of size < 500 mm^3 were excluded from analyses, providing a final parcel
count of 118. Based on the Euclidean mean MNI coordinate of each parcel location and its
associated Brodmann area (BA)[72,73], we selected 58 parcels located in premotor cortex (BA6),
inferior parietal lobule (BA 39 and 40), inferior frontal gyrus (BA 44 and 45), pre-supplementary
(BA6) and primary motor area (BA4) to represent the spatially distributed mirror neuron system
(MNS) [29,32,38].

For the within-task statistics, we used paired t-tests to compare the task effect compared to
rest, with the $p < 0.01$ for each tail with a false discovery rate (FDR) correction. We used two-
sample t-tests to compare contrast effects between motor observation and motor imitation, with
$p < 0.05$ for each tail with an FDR correction. We calculated Pearson correlations between the
parcel-specific responses to motor observation and motor imitation with the SRS-2 to investigate
brain-behavior correlations relative to autism traits. We also calculated the Pearson correlation
between the motor imitation responses and the CAMI scores to assess brain-behavior relationships
relative to motor imitation fidelity.

Results

Behavioral Measures

First, autistic participants exhibited significantly higher scores on the Social Responsiveness Scale (SRS-2), indicating elevated autism-related traits compared to NAI participants ($p = 0.0003$, $t = 4.437$, NAI N = 74, ASD N = 17, **Figure 1G (i, ii)**). Second, there were no significant differences between the NAI and ASD groups in verbal or non-verbal IQ **Figure 1G (iii)**. Third, in line with previous CAMI studies[74,75], the NAI group exhibited significant higher CAMI scores compared to the ASD participants ($p = 0.004$, $t = 2.96$, NAI N = 74, ASD N = 15, **Figure 1G (iv)**); indicating that the NAI individuals had the better motor imitation performance than the ASD group.

Neural Correlates of Motor Observation and Imitation

Whole-brain analyses indicated widespread activation in the visual and temporal cortices during both observation and imitation tasks (**Figure 2**), and additional motor brain activation during the motor imitation task. Additionally, the visual and temporal brain area exhibit significant the significant greater activation during motor observation task instead of the motor imitation task. Moreover, We observed overlapping brain activation in the temporal, visual, and bilateral inferior frontal gyrus (IFG) regions during both the motor observation and motor imitation tasks (**Figure 3**). Similarly, additional activation in the motor cortex was observed during the IM task, highlighting enhanced engagement of motor-related regions during active imitation compared to passive observation (**Figure 4**). Significant activations are apparent in MNS in response to both observing and imitating motor movements, specifically the bilateral IFG and IPL. Notably, compared to observing, imitating led to significantly greater activation in extended MNS areas, including M1 and SMA (**Supplementary Figure 1B, C**). Moreover, we observed that the MNS area exhibit the greater activation during observing compared to imitating (**Figure 5**, **Supplementary Figure 1B, C**).

To further investigate the activations from a functional network perspective, we quantified the number of cortical parcels activated within each network for each task (**Supplementary Figure 2**). Most brain networks were dynamically engaged during motor observation and imitation tasks. The dorsal attention network (DAN) and the somatosensory hand network exhibited larger involvement during imitation than observation (**Supplementary Figure 2D, Supplementary Figure 3**). In contrast, the visual and ventral attention network (VAN) exhibited greater recruitment during observation than imitation (**Supplementary Figure 2D, Supplementary Figure 4**).

Associations with SRS

During the observation task, a positive correlation was identified between SRS scores and
brain activation in the left secondary visual cortex, right premotor cortex, right somatosensory
cortex, and right inferior parietal lobule (r-IPL). Notably, the correlation in the r-IPL remained
significant after FDR correction (FDR corrected $p < 0.001$, $r = 0.472$). Moreover, activation
magnitude in the right angular gyrus (r-AG), left secondary visual cortex, and left visual
association cortex (l-VAC) exhibited a negative correlation with SRS score ($p < 0.05$, **Figure 6**).
This negative correlation indicates that individuals with higher SRS scores exhibited reduced
activation in these cortical regions. A similar pattern of negative correlations in left primary motor
and secondary visual cortex was observed during imitation. Further, we conducted hypothesis
driven brain behavioral analyses for parcels within the MNS, DAN, and VAN. We found negative
correlations between parcels within MNS (i.e., **Supplementary Figure 5**) and SRS scores,
suggesting autistic individuals with stronger autistic characteristics exhibit lower activity in parcels
within the MNS. No significant correlations were observed between SRS scores and brain function
during imitation in parcels in either the DAN or VAN.

Associations with CAMI

Positive correlations between cortical responses during imitation and CAMI scores were
observed in primary motor cortex, indicating that higher CAMI scores are associated with
increased activation in these areas (**Figure 7**). Conversely, negative correlations were identified in
left IFG areas. Additionally, we ran hypothesis driven analyses using the brain activity in the MNS,
VAN, and DAN areas (**Supplementary Figure 6**). Core regions of the MNS (i.e., particularly the
IFG and IPL), exhibited negative correlations between activation and CAMI scores. Within the
VAN and DAN, significant negative correlations with CAMI scores were observed in areas
proximal to the IFG.

Discussion

Our results confirm that both observation and imitation of motor tasks engage a spatially
and functionally distributed set of brain regions, with significant involvement throughout the
MNS. Additionally, we observed task-specific differences, highlighting variability in brain activity
that correlate with motor imitation fidelity and social responsiveness. These findings contribute to
understanding how variability in MNS may contribute to the motor imitation deficits and their
association with autistic traits.

The current study provides compelling evidence for the role of the MNS during both motor
imitation and observation. Specifically, our results demonstrate significant activation in the
posterior superior temporal sulcus (STS), inferior parietal lobule (IPL), and inferior frontal gyrus
(IFG), key regions of the MNS [32,33,76]. As hypothesized, motor imitation elicited greater activation
in motor-related areas, including the primary motor cortex (M1) and supplementary motor area
(SMA), which aligns with the functional role of the MNS in encoding observed actions and
translating them into motor commands [32,39,77]- processes integral to motor imitation. Notably, the

negative correlation observed between MNS activation and SRS scores further underscores the association between MNS variability and autistic traits, suggesting that lower MNS activation may contribute to disrupted social engagement and communication[17].

The VAN is primarily involved in stimulus-driven attention, as it responds to salient stimuli in the environment [51,52,54,78]. In the context of motor imitation, we observed engagement of VAN regions during both motor observation and imitation, albeit to a lesser extent compared to the MNS and DAN regions. Although no significant brain-behavior correlations were found between VAN activation and SRS t-scores, the role of VAN in sensory processing and attention shifting remains crucial for imitation. Our findings suggest that dysfunction in the VAN could potentially impair how individuals with ASD attend to social stimuli, affecting their ability to replicate accurately observed actions. Future studies with larger sample sizes may provide more conclusive evidence regarding the contribution of VAN in imitation deficits associated with ASD. Similarly, the DAN, known for supporting goal-directed attention and motor planning [51,79-81], showed robust activation during both motor observation and imitation. As hypothesized, greater involvement of DAN was observed during the IM task, indicating that motor planning and goal-directed attention processes are more critical during active imitation[52,54,82]. Interestingly, we observed a significant negative correlation between DAN activation and CAMI scores, which suggests that deficits in the DAN may contribute to reduced motor adaptability and difficulties in motor execution[51,79].

The analysis of brain-behavior correlations between brain activation and CAMI scores revealed significant associations in MNS regions implicated in motor planning and execution. Positive correlations were observed in motor regions such as the right primary motor cortex, suggesting that greater motor imitation fidelity (higher CAMI scores) is linked to enhanced activation in these areas. This aligns with the hypothesis that stronger motor system engagement supports better imitation performance. An fMRI study found a positive correlation in the primary motor cortex with participants motor performance[83]. Additionally, we found brain activation in IFG negatively correlated with CAMI scores. Another study found inferior parietal lobe (IPL; also implicated as part of the MNS) activity was negatively correlated with imitation accuracy. Overall, this suggests that reduced activation in the core MNS regions may be associated with difficulties in motor adaptability [31,84,85]. These findings provide insight into how deficits in neural regions critical for motor control and imitation may contribute to reduced imitation fidelity in individuals with ASD.

The relationship between brain activation and social responsiveness was further investigated for both motor observation and imitation. Negative correlations between SRS scores and brain activation were observed in regions such as the right angular gyrus (r-AG) and left secondary visual cortex (l-SVC), particularly during observation. Although these correlations did not reach statistical significance during motor imitation, they provide preliminary evidence that impaired brain function in regions involved in action understanding and social cognition may underlie the social difficulties observed in ASD. Interestingly, our results show a positive

correlation between SRS and brain activation in right parietal lobe. This finding may indicate that
individuals with higher SRS scores rely more heavily on this region to process and understand
observed actions or social cues, potentially compensating for difficulties in other aspects of social
interaction [86,87].

Limitations

While this study provides valuable insights into the neural correlates of motor imitation deficits in
ASD, several limitations should be addressed in future research. First, the relatively small sample
size of the ASD group may limit the generalizability of the findings. Future studies with larger and
more diverse populations are needed to confirm these results. Additionally, although HD-DOT
offers significant advantages over traditional neuroimaging methods, its field of view is less
extensive than that of fMRI, which limits sensitivity to neural activity in deeper regions such as
cerebellum and sub-cortical regions previously implicated in ASD functional neuroimaging
studies[88,89]. Finally, while the current study focused on adults, investigating neural mechanisms of
motor imitation in children with ASD will provide critical developmental insights. This study
provides essential feasibility for future studies on motor imitation in adolescent, school-age, and
even infant and toddler autistic and typically developing children using combined CAMI and HD-
DOT methods.

Conclusions

We used simultaneous HD-DOT functional neuroimaging with CAMI methods to investigate how
brain function covaries with observing and imitation motor movements. We observed significant
activation in the MNS during both motor observation and imitation tasks. Notably, MNS regions
demonstrated greater activation during motor observation compared to imitation, while motor
areas exhibited higher activation during imitation. These findings suggest distinct yet overlapping
neural mechanisms for action observation and execution, with shared engagement of visual,
temporal, and bilateral IFG regions. Additionally, task-specific activation was observed in the
DAN and the extended MNS, underscoring their roles in motor planning and goal-directed
attention during imitation. Brain-behavior analyses revealed that reduced MNS activation was
associated with higher autism-related traits and lower motor imitation fidelity. Overall, these
findings highlight the neural correlates of action observation/execution and their association with
social reciprocity and motor imitation fidelity. Finally, these studies lay essential foundation for
future investigations using simultaneous methods for naturalistic functional neuroimaging and
quantitative CAMI that may be applied during early childhood development when presentation of
autistic traits become apparent and interventions may hold the greatest promise to improve
outcome.

Acknowledgements

We would like to express our sincere gratitude to the participants for their valuable contributions
to this study. This research was supported by grants from the National Institutes of Health (R01-
MH12275101, R21-MH127501), the McDonnell Center for Systems Neuroscience, and the Eagles
Autism Foundation.
334

References:

Maenner, M. J. Prevalence of Autism Spectrum Disorder Among Children Aged 8 Years — Autism and Developmental Disabilities Monitoring Network, 11 Sites, United States, 2016. *MMWR. Surveillance Summaries* **69** (2020). https://doi.org:10.15585/mmwr.ss6904a1

Mottron, L. & Bzdok, D. Autism spectrum heterogeneity: fact or artifact? *Molecular Psychiatry* **25**, 3178-3185 (2020). https://doi.org:10.1038/s41380-020-0748-y

Miller, H. L. *et al*. Shared Features or Co-occurrence? Evaluating Symptoms of Developmental Coordination Disorder in Children and Adolescents with Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders* **51**, 3443-3455 (2021). https://doi.org:10.1007/s10803-020-04766-z

Miller, H. L. *et al*. Motor problems in autism: Co-occurrence or feature? *Developmental medicine and child neurology* **66**, 16-22 (2024). https://doi.org:10.1111/dmcn.15674

Licari, M. K. *et al*. Prevalence of Motor Difficulties in Autism Spectrum Disorder: Analysis of a Population-Based Cohort. *Autism Research: Official Journal of the International Society for Autism Research* **13**, 298-306 (2020). https://doi.org:10.1002/aur.2230

Bhat, A. N. Is Motor Impairment in Autism Spectrum Disorder Distinct From Developmental Coordination Disorder? A Report From the SPARK Study. *Physical Therapy* **100**, 633-644 (2020). https://doi.org:10.1093/ptj/pzz190

Mostofsky, S. H. *et al*. Developmental dyspraxia is not limited to imitation in children with autism spectrum disorders. *Journal of the International Neuropsychological Society: JINS* **12**, 314-326 (2006). https://doi.org:10.1017/s1355617706060437

MacNeil, L. K. & Mostofsky, S. H. Specificity of dyspraxia in children with autism. *Neuropsychology* **26**, 165-171 (2012). https://doi.org:10.1037/a0026955

Vivanti, G. & Hamilton, A. *Imitation in Autism Spectrum Disorders*, 2014).

McAuliffe, D. *et al*. Learning of skilled movements via imitation in ASD. *Autism Research: Official Journal of the International Society for Autism Research* **13**, 777-784 (2020). https://doi.org:10.1002/aur.2253

Taverna, E. C., Huedo-Medina, T. B., Fein, D. A. & Eigsti, I.-M. The interaction of fine motor, gesture, and structural language skills: The case of autism spectrum disorder. *Research in Autism Spectrum Disorders* **86**, 101824 (2021). https://doi.org:https://doi.org/10.1016/j.rasd.2021.101824

Pittet, I., Kojovic, N., Franchini, M. & Schaer, M. Trajectories of imitation skills in preschoolers with autism spectrum disorders. *Journal of Neurodevelopmental Disorders* **14**, 2 (2022). https://doi.org:10.1186/s11689-021-09412-y

Yang, D., Sidman, J. & Bushnell, E. W. Beyond the information given: Infants' transfer of actions learned through imitation. *Journal of Experimental Child Psychology* **106**, 62-81 (2010). https://doi.org:https://doi.org/10.1016/j.jecp.2009.12.005

Nebel, M. B. *et al*. Intrinsic Visual-Motor Synchrony Correlates With Social Deficits in Autism. *Biological Psychiatry* **79**, 633-641 (2016). https://doi.org:10.1016/j.biopsych.2015.08.029

15   Hirai, M. *et al.* Greater reliance on proprioceptive information during a reaching task
with perspective manipulation among children with autism spectrum disorders.
Scientific Reports **11**, 15974 (2021). https://doi.org:10.1038/s41598-021-95349-0

16   Marko, M. K. *et al.* Behavioural and neural basis of anomalous motor learning in
children with autism. *Brain: A Journal of Neurology* **138**, 784-797 (2015).
https://doi.org:10.1093/brain/awu394

17   Zhao, W. *et al.* Differential responses in the mirror neuron system during imitation of
individual emotional facial expressions and association with autistic traits.
NeuroImage **277**, 120263 (2023).
https://doi.org:https://doi.org/10.1016/j.neuroimage.2023.120263

18   Scarapicchia, V., Brown, C., Mayo, C. & Gawryluk, J. R. Functional Magnetic
Resonance Imaging and Functional Near-Infrared Spectroscopy: Insights from
Combined Recording Studies. *Frontiers in Human Neuroscience* **11** (2017).
https://doi.org:ARTN 419
10.3389/fnhum.2017.00419

19   Crone, E. A., Poldrack, R. A. & Durston, S. Challenges and methods in developmental
neuroimaging. *Hum Brain Mapp* **31**, 835-837 (2010).
https://doi.org:10.1002/hbm.21053

20   Eggebrecht, A. T. *et al.* Mapping distributed brain function and networks with diffuse
optical tomography. *Nature Photonics* **8**, 448-454 (2014).
https://doi.org:10.1038/nphoton.2014.107

21   Hassanpour, M. S., Eggebrecht, A. T., Peelle, J. E. & Culver, J. P. Mapping effective
connectivity within cortical networks with diffuse optical tomography.
Neurophotonics **4**, 041402 (2017). https://doi.org:10.1117/1.NPh.4.4.041402

22   Fishell, A. K., Burns-Yocum, T. M., Bergonzi, K. M., Eggebrecht, A. T. & Culver, J. P.
Mapping brain function during naturalistic viewing using high-density diffuse optical
tomography. *Scientific Reports* **9**, 11115 (2019). https://doi.org:10.1038/s41598-
019-45555-8

23   Yang, D. *et al.* Mapping neural correlates of biological motion perception in autistic
children using high-density diffuse optical tomography. *Mol Autism* **15**, 35 (2024).
https://doi.org:10.1186/s13229-024-00614-4

24   McMorrow, S. R. *et al.* Bedside Neuroimaging Using High-Density Diffuse Optical
Tomography in a Pediatric Patient on Extracorporeal Support. *Stroke* **0**
https://doi.org:doi:10.1161/STROKEAHA.124.049116

25   Eggebrecht, A. T. *et al.* A quantitative spatial comparison of high-density diffuse
optical tomography and fMRI cortical mapping. *NEUROIMAGE* **61**, 1120-1128 (2012).
https://doi.org:10.1016/j.neuroimage.2012.01.124

26   Ferradal, S. L., Eggebrecht, A. T., Hassanpour, M., Snyder, A. Z. & Culver, J. P. Atlas-
based head modeling and spatial normalization for high-density diffuse optical
tomography:
validation against fMRI. *Neuroimage* **85**, 117-126 (2014).
https://doi.org:10.1016/j.neuroimage.2013.03.069

27 Sherafati, A. *et al*. Global motion detection and censoring in high-density diffuse
optical tomography. *Human Brain Mapping* **41**, 4093-4112 (2020).
https://doi.org:10.1002/hbm.25111
28 Williams, J. H. G., Whiten, A., Suddendorf, T. & Perrett, D. I. Imitation, mirror neurons
and autism. *Neurosci Biobehav R* **25**, 287-295 (2001). https://doi.org:Doi
10.1016/S0149-7634(01)00014-8
29 Rizzolatti, G. & Craighero, L. THE MIRROR-NEURON SYSTEM. *Annual Review of*
Neuroscience **27**, 169-192 (2004).
https://doi.org:10.1146/annurev.neuro.27.070203.144230
30 Buccino, G. *et al*. Neural circuits underlying imitation learning of hand actions: an
event-related fMRI study. *Neuron* **42**, 323-334 (2004). https://doi.org:10.1016/s0896-
6273(04)00181-3
31 Iacoboni, M. *et al*. Grasping the Intentions of Others with One's Own Mirror Neuron
System. *PLoS Biology* **3**, e79 (2005). https://doi.org:10.1371/journal.pbio.0030079
32 Iacoboni, M. & Dapretto, M. The mirror neuron system and the consequences of its
dysfunction. *Nature Reviews Neuroscience* **7**, 942-951 (2006).
https://doi.org:10.1038/nrn2024
33 Cattaneo, L. & Rizzolatti, G. The Mirror Neuron System. *Archives of Neurology* **66**,
557-560 (2009). https://doi.org:10.1001/archneurol.2009.41
34 Dushanova, J. & Donoghue, J. Neurons in primary motor cortex engaged during action
observation. *Eur J Neurosci* **31**, 386-398 (2010). https://doi.org:10.1111/j.1460-
9568.2009.07067.x
35 Kana, R. K., Wadsworth, H. M. & Travers, B. G. A systems level analysis of the mirror
neuron hypothesis and imitation impairments in autism spectrum disorders.
Neurosci Biobehav R **35**, 894-902 (2011).
https://doi.org:10.1016/j.neubiorev.2010.10.007
36 Umiltà, M. A. *et al*. I Know What You Are Doing: A Neurophysiological Study. *Neuron*
31, 155-165 (2001). https://doi.org:10.1016/S0896-6273(01)00337-3
37 Buccino, G. *et al*. Neural circuits underlying imitation learning of hand actions: An
event-related fMRI study. *Neuron* **42**, 323-334 (2004). https://doi.org:Doi
10.1016/S0896-6273(04)00181-3
38 Kemmerer, D. What modulates the Mirror Neuron System during action
observation?: Multiple factors involving the action, the actor, the observer, the
relationship between actor and observer, and the context. *Progress in Neurobiology*
205, 102128 (2021). https://doi.org:10.1016/j.pneurobio.2021.102128
39 Iacoboni, M. *et al*. Cortical mechanisms of human imitation. *Science* **286**, 2526-2528
(1999). https://doi.org:10.1126/science.286.5449.2526
40 Dapretto, M. *et al*. Understanding emotions in others: mirror neuron dysfunction in
children with autism spectrum disorders. *Nat Neurosci* **9**, 28-30 (2006).
https://doi.org:10.1038/nn1611
41 Carr, L., Iacoboni, M., Dubeau, M.-C., Mazziotta, J. C. & Lenzi, G. L. Neural
mechanisms of empathy in humans: A relay from neural systems for imitation to
limbic areas. *Proceedings of the National Academy of Sciences* **100**, 5497-5502
(2003). https://doi.org:doi:10.1073/pnas.0935845100

42   Mukamel, R., Ekstrom, A. D., Kaplan, J., Iacoboni, M. & Fried, I. Single-neuron
responses in humans during execution and observation of actions. *Curr Biol* **20**, 750-
756 (2010). https://doi.org:10.1016/j.cub.2010.02.045

43   Falck-Ytter, T., Gredebäck, G. & von Hofsten, C. Infants predict other people's action
goals. *Nature Neuroscience* **9**, 878-879 (2006). https://doi.org:10.1038/nn1729

44   Oberman, L. M., Ramachandran, V. S. & Pineda, J. A. Modulation of mu suppression
in children with autism spectrum disorders in response to familiar or unfamiliar
stimuli: The mirror neuron hypothesis. *Neuropsychologia* **46**, 1558-1565 (2008).
https://doi.org:10.1016/j.neuropsychologia.2008.01.010

45   Raymaekers, R., Wiersema, J. R. & Roeyers, H. EEG study of the mirror neuron system
in children with high functioning autism. *Brain Research* **1304**, 113-121 (2009).
https://doi.org:10.1016/j.brainres.2009.09.068

46   Tylen, K., Allen, M., Hunter, B. & Roepstorff, A. Interaction vs. observation: distinctive
modes of social cognition in human brain and behavior? A combined fMRI and eye-
tracking study. *Frontiers in Human Neuroscience* **6** (2012).

47   Datko, M., Pineda, J. A. & Müller, R.-A. Positive effects of neurofeedback on autism
symptoms correlate with brain activation during imitation and observation. *European*
Journal of Neuroscience **47**, 579-591 (2018). https://doi.org:10.1111/ejn.13551

48   Wadsworth, H. M., Maximo, J. O., Donnelly, R. J. & Kana, R. K. Action simulation and
mirroring in children with autism spectrum disorders. *Behavioural Brain Research*
341, 1-8 (2018). https://doi.org:10.1016/j.bbr.2017.12.012

49   Lust, J. M. *et al.* Activation of Mirror Neuron Regions Is Altered in Developmental
Coordination Disorder (DCD)–Neurophysiological Evidence Using an Action
Observation Paradigm. *Frontiers in Human Neuroscience* **13**, 232 (2019).
https://doi.org:10.3389/fnhum.2019.00232

50   Scassellati, B. in *Computation for Metaphors, Analogy, and Agents* Vol. 1562 (ed
Chrystopher L. Nehaniv) 176-195 (Springer Berlin Heidelberg, 1999).

51   Corbetta, M. & Shulman, G. L. Control of goal-directed and stimulus-driven attention
in the brain. *Nature Reviews Neuroscience* **3**, 201-215 (2002).
https://doi.org:10.1038/nrn755

52   He, B. J. *et al.* Breakdown of Functional Connectivity in Frontoparietal Networks
Underlies Behavioral Deficits in Spatial Neglect. *Neuron* **53**, 905-918 (2007).
https://doi.org:10.1016/j.neuron.2007.02.013

53   Marco, E. J., Hinkley, L. B. N., Hill, S. S. & Nagarajan, S. S. Sensory processing in
autism: a review of neurophysiologic findings. *Pediatr Res* **69**, 48R-54R (2011).
https://doi.org:10.1203/PDR.0b013e3182130c54

54   Nardo, D., Santangelo, V. & Macaluso, E. Stimulus-Driven Orienting of Visuo-Spatial
Attention in Complex Dynamic Environments. *Neuron* **69**, 1015-1028 (2011).
https://doi.org:10.1016/j.neuron.2011.02.020

55   Staples, K. L., MacDonald, M. & Zimmer, C. in *International Review of Research in*
Developmental Disabilities Vol. 42 (ed Robert M. Hodapp) 179-214 (Academic Press,
2012).

56   Tuncgenc, B. *et al*. Computerized Assessment of Motor Imitation as a Scalable
Method for Distinguishing Children With Autism. *Biol Psychiatry Cogn Neurosci*
Neuroimaging **6**, 321-328 (2021). https://doi.org:10.1016/j.bpsc.2020.09.001

57   Celebi, S., Aydin, A. S., Temiz, T. T. & Arici, T. in *International Conference on*
Computer Vision Theory and Applications.

58   Lidstone, D. *et al*. Automated and scalable Computerized Assessment of Motor
Imitation (CAMI) in children with Autism Spectrum Disorder using a single 2D
camera: A pilot study. *Research in Autism Spectrum Disorders* **87** (2021).
https://doi.org:https://doi.org/10.1016/j.rasd.2021.101840

59   Constantino, J. N. *et al*. Validation of a brief quantitative measure of autistic traits:
comparison of the social responsiveness scale with the autism diagnostic interview-
revised.   *J   Autism   Dev   Disord   **33**,   427-433   (2003).
https://doi.org:10.1023/a:1025014929212

60   Tripathy, K. *et al*. Mapping brain function in adults and young children during
naturalistic viewing with high-density diffuse optical tomography. *Hum Brain Mapp*
45, e26684 (2024). https://doi.org:10.1002/hbm.26684

61   Constantino, J. N. & Gruber, C. P. *Social responsiveness scale : SRS-2*. Second
edition edn, (Western Psychological Services, 2012).

62   Oldfield, R. C. The assessment and analysis of handedness: the Edinburgh inventory.
Neuropsychologia **9**, 97-113 (1971). https://doi.org:10.1016/0028-3932(71)90067-4

63   Pitts, C. H. & Mervis, C. B. Performance on the Kaufman Brief Intelligence Test-2 by
Children With Williams Syndrome. *Am J Intellect Dev Disabil* **121**, 33-47 (2016).
https://doi.org:10.1352/1944-7558-121.1.33

64   Tzourio-Mazoyer, N. *et al*. Automated anatomical labeling of activations in SPM using
a macroscopic anatomical parcellation of the MNI MRI single-subject brain.
Neuroimage **15**, 273-289 (2002). https://doi.org:10.1006/nimg.2001.0978

65   Lancaster, J. L. *et al*. Bias between MNI and talairach coordinates analyzed using the
ICBM-152 brain template. *Human Brain Mapping* **28**, 1194-1205 (2007).
https://doi.org:10.1002/hbm.20345

66   Fishell, A. K. *et al*. Portable, field -based neuroimaging using high -density diffuse
optical   tomography.   *NEUROIMAGE*   **215**   (2020).
https://doi.org:10.1016/j.neuroimage.2020.116541

67   Tripathy, K. *et al*. Decoding visual information from high-density diffuse optical
tomography   neuroimaging   data.   *NEUROIMAGE*   **226**   (2021).
https://doi.org:10.1016/j.neuroimage.2020.117516

68   Schroeder, M. L. *et al*. Mapping cortical activations underlying covert and overt
language production using high-density diffuse optical tomography. *Neuroimage*
276, 120190 (2023). https://doi.org:10.1016/j.neuroimage.2023.120190

69   Hassanpour, M. S. *et al*. Statistical analysis of high density diffuse optical
tomography.   *NeuroImage*   **85**,   104-116   (2014).
https://doi.org:10.1016/j.neuroimage.2013.05.105

70   Yucel, M. A. *et al*. Best practices for fNIRS publications. *Neurophotonics* **8**, 012101
(2021). https://doi.org:10.1117/1.NPh.8.1.012101

71 Gordon, E. M. *et al*. Generation and Evaluation of a Cortical Area Parcellation from
Resting-State Correlations. *Cerebral Cortex* **26**, 288-303 (2016).
https://doi.org:10.1093/cercor/bhu239
72 Lacadie, C. M., Fulbright, R. K., Rajeevan, N., Constable, R. T. & Papademetris, X.
More accurate Talairach coordinates for neuroimaging using non-linear registration.
Neuroimage **42**, 717-725 (2008). https://doi.org:10.1016/j.neuroimage.2008.04.240
73 Chau, W. & McIntosh, A. R. The Talairach coordinate of a point in the MNI space: how
to interpret it. *Neuroimage* **25**, 408-416 (2005).
https://doi.org:10.1016/j.neuroimage.2004.12.007
74 Tunçgenç, B. *et al*. Computerized Assessment of Motor Imitation as a Scalable
Method for Distinguishing Children With Autism. *Biological Psychiatry: Cognitive*
Neuroscience and Neuroimaging **6**, 321-328 (2021).
https://doi.org:10.1016/j.bpsc.2020.09.001
75 Lidstone, D. E. *et al*. Automated and scalable Computerized Assessment of Motor
Imitation (CAMI) in children with Autism Spectrum Disorder using a single 2D
camera: A pilot study. *Res Autism Spectr Disord* **87** (2021).
https://doi.org:10.1016/j.rasd.2021.101840
76 Chan, M. M. Y. & Han, Y. M. Y. Differential mirror neuron system (MNS) activation
during action observation with and without social-emotional components in autism:
a meta-analysis of neuroimaging studies. *Molecular Autism* **11**, 72 (2020).
https://doi.org:10.1186/s13229-020-00374-x
77 Dapretto, M. *et al*. Understanding emotions in others: mirror neuron dysfunction in
children with autism spectrum disorders. *Nature Neuroscience* **9**, 28-30 (2006).
https://doi.org:10.1038/nn1611
78 Vossel, S., Geng, J. J. & Fink, G. R. Dorsal and Ventral Attention Systems: Distinct
Neural Circuits but Collaborative Roles. *The Neuroscientist* **20**, 150-159 (2014).
https://doi.org:10.1177/1073858413494269
79 Astafiev, S. V. *et al*. Functional Organization of Human Intraparietal and Frontal
Cortex for Attending, Looking, and Pointing. *The Journal of Neuroscience* **23**, 4689-
4699 (2003). https://doi.org:10.1523/JNEUROSCI.23-11-04689.2003
80 Majerus, S., Péters, F., Bouffier, M., Cowan, N. & Phillips, C. The Dorsal Attention
Network Reflects Both Encoding Load and Top–down Control during Working
Memory. *Journal of Cognitive Neuroscience* **30**, 144-159 (2018).
https://doi.org:10.1162/jocn_a_01195
81 Stawarczyk, D., Jeunehomme, O. & D'Argembeau, A. Differential Contributions of
Default and Dorsal Attention Networks to Remembering Thoughts and External
Stimuli From Real-Life Events. *Cerebral Cortex* **28**, 4023-4035 (2018).
https://doi.org:10.1093/cercor/bhx270
82 Fox, M. D., Corbetta, M., Snyder, A. Z., Vincent, J. L. & Raichle, M. E. Spontaneous
neuronal activity distinguishes human dorsal and ventral attention systems.
Proceedings of the National Academy of Sciences **103**, 10046-10051 (2006).
https://doi.org:10.1073/pnas.0604187103

83 Aridan, N. & Mukamel, R. Activity in primary motor cortex during action observation
covaries with subsequent behavioral changes in execution. *Brain and Behavior* **6**
(2016). https://doi.org:10.1002/brb3.550
84 Villalobos, M. E., Mizuno, A., Dahl, B. C., Kemmotsu, N. & Müller, R.-A. Reduced
functional connectivity between V1 and inferior frontal cortex associated with
visuomotor performance in autism. *NeuroImage* **25**, 916-925 (2005).
https://doi.org:10.1016/j.neuroimage.2004.12.022
85 Marsh, L. E. & Hamilton, A. F. d. C. Dissociation of mirroring and mentalising systems
in autism. *NeuroImage* **56**, 1511-1519 (2011).
https://doi.org:10.1016/j.neuroimage.2011.02.003
86 Su, W. C., Culotta, M., Mueller, J., Tsuzuki, D. & Bhat, A. N. Autism-Related
Differences in Cortical Activation When Observing, Producing, and Imitating
Communicative Gestures: An fNIRS Study. *Brain Sciences* **13** (2023).
https://doi.org:ARTN 1284
10.3390/brainsci13091284
87 Su, W. C. *et al.* Differences in cortical activation patterns during action observation,
action execution, and interpersonal synchrony between children with or without
autism spectrum disorder (ASD): An fNIRS pilot study. *Plos One* **15** (2020).
https://doi.org:ARTN e0240301
10.1371/journal.pone.0240301
88 Laidi, C. *et al.* Cerebellar anatomical alterations and attention to eyes in autism.
Scientific Reports **7** (2017). https://doi.org:ARTN 12008
10.1038/s41598-017-11883-w
89 Cerliani, L. *et al.* Increased Functional Connectivity Between Subcortical and
Cortical Resting-State Networks in Autism Spectrum Disorder. *Jama Psychiatry* **72**,
767-777 (2015). https://doi.org:10.1001/jamapsychiatry.2015.0101

Figure 1: Study Design and Behavioral Assessments. A Participant wearing the DOT cap. **B** Field of view from dorsal, posterior, and lateral perspectives and overlapping Gordon parcels colored by functional networks assignment. **C** Experimental design for the motor observation and imitation tasks. **D** Participant performing imitation of movements. **E** Three examples of extracted skeleton of moving participant for CAMI analyses. **F** Comparison of imitation performance with a gold-standard skeleton positions during the movements. **G** Behavioral assessments including SRS scores, IQ, and CAMI scores.

Figure 2: Neural correlates of observation and imitation. A. Group-level t-maps for motor observation (N = 77), motor imitation (N = 63), and their contrast. **B.** FDR-corrected t-maps highlighting statistically significant regions, with false discovery rate correction applied to account for multiple comparisons.

Figure 3: Brain regions exhibiting significant responses to both motor observation and imitation. **A.** Brain regions with overlapping significant activation identified through conjunction analysis. **B.** Temporal profiles of the ΔHbO_2 (oxygenated hemoglobin) hemodynamic response for each task, with shaded areas indicating the standard error of the mean across participants. The selected parcels represent the significant regions with the eight smallest FDR-corrected p-values, showing the greatest activation. The abbreviations correspond to their respective functional networks: VIS refers to the Visual Network, FPN to the Frontoparietal Network, DAN to the Dorsal Attention Network, and DMN to the Default Mode Network.

Figure 4: Brain regions exhibiting significant responses during motor imitation but not motor observation. A. Brain regions with distinct significant activation during motor imitation that are not significantly active during motor observation. **B.** Temporal profiles of the ΔHbO_2 (oxygenated hemoglobin) hemodynamic response for each task, with shaded areas representing the standard error of the mean across participants. The selected parcels correspond to the significant regions with the eight smallest FDR-corrected p-values, demonstrating the great activation. The abbreviations correspond to their respective functional networks: DAN refer to the Dorsal Attention Network and SMH to the Somatomotor Hand Network.

Figure 5: Brain regions exhibiting significant responses during motor observation but not motor imitation. A. Brain regions with distinct significant activation during motor observation task that are not significantly active during motor imitation. **B.** Temporal profiles of the ΔHbO_2 (oxygenated hemoglobin) hemodynamic response for each task, with shaded areas representing the standard error of the mean across participants. The selected parcels correspond to the significant regions with the four smallest FDR-corrected p-values, demonstrating the great activation. The abbreviations correspond to their respective functional networks: VIS refers to the Visual Network and DMN to the Default Mode Network.

Figure 6. Brain-behavioral association between SRS scores and brain activation. **A (i)** Correlation map of brain activation during motor observation with SRS in 71 participants**, (ii)** the brain regions which correlation p < 0.05, and **(iii)** the correlation passed the FDR correction (FDR corrected p < 0.001, and R = 0.472)**. B (i)** Correlation map of brain activation during motor imitation with SRS in 55 participants. **(ii)** the brain regions which correlation p < 0.05. There is no correlation passed the FDR correction.

Figure 6. Brain-behavioral association between CAMI and brain activation during motor imitation task. A The correlation map for motor imitation activity against SRS scores for 57 participants. **B** The same map thresholded (p < 0.05).

Table 1 Participant demographics and group characteristic

Measure		Full Sample NAC n = 81			Full Sample ASD n = 19		
Sex, Males:Females		33:48			12:7		
		Means	*S.D.*	*n*	*Means*	*S.D.*	*n*
Age, years, [range]		27.56	9.36	81	31.47	16.73	19
SRS	Raw score	43.61	24.68	74	82.94	33.72	17
	T score	51.07	8.67	74	65.00	11.89	17
	Verbal	113.38	17.22	61	103.39	20.04	18
KBIT 2 IQ	Nonverbal	117.61	11.97	61	110.67	16.82	18
	Full scale	117.75	13.69	61	107.83	19.77	18
CAMI	Scores	0.52	0.21	75	0.31	0.26	18

Supplementary Figure 1: Neural correlates of observation and imitation within the mirror neuron system. A. Parcel-based analysis of the MNS within the field of view, highlighting regions activated during both motor observation and imitation. **B.** Group-level t-maps depicting the areas of the MNS that are significantly activated during motor observation and imitation. **C.** FDR-corrected t-map showing the statistically significant regions within the MNS, with false discovery rate correction applied to control for multiple comparisons.

Supplementary Figure 2: Functional network involvement during motor observation and imitation. A Task specific significant brain parcels under of three different conditions (OBS, IM, and their contrast). **B** Functional networks involved during the OBS task. **C** Functional networks involved during the IM task. **D** Functional networks more active during the OBS task than the IM task. **E** Functional networks more active during the IM task than the OBS task.

Supplementary Figure 3: Neural correlates of OBS and IM within dorsal attention network. A. Group-level t-maps for motor observation (N = 77), motor imitation (N = 63), and their contrast (red indicates brain regions where OBS > IM, and blue indicates brain regions where IM > OBS). **B.** FDR-corrected t-map highlighting statistically significant regions, with false discovery rate correction applied to account for multiple comparisons.

Supplementary Figure 4: Neural correlates of OBS and IM within ventral attention network. A. Group-level t-maps for motor observation (N = 77), motor imitation (N = 63), and their contrast (red indicates brain regions where OBS > IM, and blue indicates brain regions where IM > OBS). **B.** FDR-corrected t-map highlighting statistically significant regions, with false discovery rate correction applied to account for multiple comparisons.

Supplementary Figure 5: Brain behavioral association (SRS vs beta) within MNS. A Correlation map of OBS brain activation with SRS in 71 participants and the brain regions which correlation p < 0.05. **B (i)** Correlation map of IM brain activation with SRS in 55 participants and brain regions which correlation p < 0.05. There is no correlation passed the FDR correction.

Supplementary Figure 6. brain-behavioral association between CAMI and brain activation during motor imitation task within MNS brain areas. A The correlation map of the IM beta values with the SRS scores for 57 participants. **B**. presents the R map, highlighting the brain parcels that show significant correlations with CAMI ($p < 0.05$). **C.** provides the p-value maps, emphasizing the significant parcels with a correlation to CAMI ($p < 0.05$).