Molecular Requirements for *C. elegans* Transgenerational Epigenetic Inheritance

of Pathogen Avoidance

Rachel Kaletsky[1,2], Rebecca Moore[1,2,3,4], Titas Sengupta[1,2], Renee Seto[1,2], Borja Ceballos-Llera[1,2] and

Coleen T. Murphy[1,2]*

[1]Department of Molecular Biology, Princeton University; Princeton, NJ 08544, USA

[2]LSI Genomics, Princeton University; Princeton, NJ 08544, USA

[3]Chronobiology and Sleep Institute, Department of Neuroscience, Philadelphia, PA 19104, USA

[4]Howard Hughes Medical Institute, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA 19104, USA

Abstract

C. elegans' major food source is bacteria, and worms are naturally attracted to many

bacterial species, including pathogenic *Pseudomonas*; in fact, worms prefer PA14 as well as

wild bacteria over the lab *E. coli* strain (OP50) standardly used in the laboratory setting. Many

labs have shown that despite this natural attraction to PA14, prior exposure to PA14 causes the

worms to instead avoid PA14. This behavioral switch can happen on a relatively fast time scale,

even within the duration of the choice assay. Here we show that accurate assessment of the

animals' true first choice requires the use of a paralytic (azide) to trap the worms at their initial

choice, and to prevent the switch from attraction to avoidance of PA14 within the assay period.

We previously discovered that exposure of *C. elegans* to 25°C plate-grown PA14 at 20°C for

24hrs not only leads to these animals switching from attraction to avoidance of PA14, but also to

their progeny avoiding PA14 in the naïve state, and this avoidance persists through the F4

generation. Other types of PA14 training can also cause P0 and/or F1 avoidance, but do not

induce transgenerational (F2 and beyond) inheritance. We also previously showed that the

transgenerational (P0-F4) learned avoidance is mediated by P11, a small RNA produced by

PA14. P11 is both necessary and sufficient for transgenerational epigenetic inheritance of

avoidance behavior. P11 is highly expressed in our standard growth conditions (25°C on

surfaces), but not in other conditions, suggesting that reported failure to observe F2-F4

avoidance is most likely due to the absence of P11 expression in PA14 in the experimenters'

growth conditions. Through mutant analyses, we have tested many genes – including germline

regulators, small RNA uptake, RNA interference/processing, chromatin modifiers, and neuronal

genes - for their involvement in transgenerational inheritance of learned pathogen avoidance,

allowing us to better understand the molecular requirements for this process. We previously

found that wild *C. elegans* strains also show TEI of learned pathogen avoidance, and that at

least two other wild bacteria, *P. vranovensis* and *P. fluorescens 15,* induce this transgenerational

avoidance. The avoidance induced by each *Pseudomonas* species functions through a specific,

distinct small RNA (Pv1 in *P. vranovensis* and Pfs1 in *P. fluorescens 15*, respectively) that either

directly or indirectly reduce the levels of the gene *maco-1*, which in turn regulates *daf-7*

expression in the ASI neuron and subsequent avoidance behavior. The conservation of multiple

components of this small RNA TEI mechanism across *C. elegans* strains and in multiple

Pseudomonas species suggests that this transgenerational learned avoidance behavior is likely

to be functional and physiologically important in wild conditions.

44
Introduction

C. elegans in the wild is found primarily in rotting fruit, where a range of bacterial species
serve as abundant food sources (Félix & Duveau, 2012). About a third of these species are of
the *Pseudomonas* family (Samuel *et al*, 2016), and many of those *Pseudomonas* species are
pathogenic to *C. elegans*. Therefore, the development of multiple strategies to detect and avoid
these pathogens that act at different time scales is critical for *C. elegans'* fitness and survival.

C. elegans and a clinical isolate of *Pseudomonas aeruginosa*, PA14, have been studied as a
model of host-pathogen interactions for several decades (Tan *et al*, 1999). PA14 kills *C. elegans*
through several distinct mechanisms (e.g., slow and fast killing (Mahajan-Miklos *et al*, 1999))
and typically *C. elegans* adults may only survive for a few days on PA14 (Tan *et al*, 1999),
cutting its reproductive period short. *C. elegans'* survival depends on the expression of anti-
microbial genes that are induced downstream of innate immune signals in response to PA14
exposure (Troemel *et al*, 2006), and changes to developmental rates and decisions that
increase the odds of the next generation's survival.

In addition to the activation of pathogen survival genes and effects on reproduction and
development, *C. elegans* also changes its behavior in the presence of PA14. Despite its
potential to kill *C. elegans,* worms are initially attracted to PA14 (Zhang *et al*, 2005), because *C.*
elegans cannot distinguish pathogenic from non-pathogenic *Pseudomonas* species that may
serve as nutritious food sources (Sengupta *et al*, 2024). Specific mechanisms to detect, leave,
and avoid PA14 are activated via secreted bacterial metabolites (Meisel *et al*, 2014; Prakash *et*
al, 2021) or through internal signals that indicate illness (Melo & Ruvkun, 2012; Singh & Aballay,
2019; Zhang *et al*, 2005). Another mechanism of learned avoidance can be activated upon
exposure to pathogenic *Pseudomonas* and is mediated by specific bacterial small RNAs (Moore
et al, 2019; Kaletsky *et al*, 2020; Sengupta *et al*, 2024; Seto *et al*, 2024). These distinct
avoidance pathways are triggered by specific molecular cues, are independently regulated, and
act on different time scales. Specifically, PA14 metabolites induce a distinct, ASJ-mediated
avoidance pathway in P0s (Meisel *et al*, 2014); 4hrs of PA14 exposure induces avoidance
directly in the worms and increased attraction in the F1s, while 8hrs of treatment can induce F1
avoidance but does not persist beyond F1 (Pereira *et al*, 2020; Zhang *et al*, 2005); and specific
bacterial small RNAs from PA14 as well as from wild *Pseudomonas* species induce avoidance
after 24hrs of treatment that is maintained through four generations of progeny (Kaletsky *et al*,
2020; Moore *et al*, 2019; Sengupta *et al*, 2024; Seto *et al*, 2024). Through extensive mutant
analyses, we have found that this transgenerational avoidance mechanism requires a wide

range of machinery, including RNA interference components, germline components, the Cer1

Ty3 Gypsy retrotransposon, and neuronal function (Moore *et al*, 2019,Kaletsky *et al*, 2020;

Moore et al. 2021; Sengupta *et al*, 2024; Seto *et al*, 2024).

Here we discuss the molecular conditions that are necessary to induce the bacterial small

RNA-induced pathway of transgenerationally-inherited learned avoidance behavior. Our results

suggest that one major difference in execution of the assay itself, the presence or absence of

the paralytic sodium azide, may account for some reported discrepancies in naïve and learned

chemotaxis results (Gainey *et al*, 2024). Perhaps more importantly, PA14 must be grown under

conditions that induce the expression of the small RNA, P11, that is necessary and sufficient for

transgenerational inheritance of learned pathogen avoidance; P11 regulates nitrogen fixation in

PA14, and these conditions promote biofilm formation, and thus may be important signals for *C.*

elegans to detect the pathogen. These two factors, P11 expression and azide presence in the

assay, are critical in obtaining consistent naïve and learned avoidance results. Together with the

identification of genes and pathways that are involved, we are beginning to understand the

molecular requirements for transgenerational epigenetic inheritance of learned avoidance.

Results

Choice assays require a paralytic to accurately capture first choice

Because *C. elegans* has several different mechanisms of pathogen avoidance that can be

detected via different signals (olfactory cues, gustatory cues, illness-induced signals, bacterial

metabolites, bacterial RNAs, etc.), pathogen avoidance behavior has been tested using several

different assays (as reviewed in (Liu & Zhang, 2020)). Among these are variations on the

bacterial choice assay, which is modeled on standard chemotaxis choice assays (two small

bacterial spots with azide to trap worms at their first choice (Bargmann *et al*, 1993; Hart, 2006;

Zhang *et al*, 2005; Jin *et al*, 2016; Lee & Mylonakis, 2017; Ooi & Prahlad, 2017; Liu *et al*, 2018;

Moore *et al*, 2019); variations on single-worm head-turning assays (Ha *et al*, 2010); and the

leaving assay (PA14 spot in middle of plate, no other bacteria, no azide, and monitored over

long periods for worms leaving the spot e.g., 8-24hrs (Meisel *et al*, 2014; Miller *et al*, 2015; Ma

et al, 2017; Horspool & Chang, 2017; Hao *et al*, 2018; Singh & Aballay, 2019; Wolfe *et al*, 2019;

Filipowicz *et al*, 2022). Each assay measures responses on a specific time scale; for example,

the two-bacteria choice assay measures initial preference of a population of animals over an

hour, the head-turning assay measures immediate preference of individual animals, and the

leaving assay (which is often performed overnight) measures final preference over many hours

of PA14 exposure. Therefore, the conditions used in one assay might not be appropriate for all

assays.

The standard bacterial choice assay is performed on plates, with two small spots of bacteria

(here, PA14 and OP50) separated by several centimeters that the worms must traverse in

response to olfactory cues (**Figure 1A**); monitoring the accumulation of worms as they become

paralyzed with the anesthetic sodium azide at each bacterial spot allows the experimenter to

accurately assess the worm's first choice. Studies by many groups, including Zhang et al, 2005,

Meisel et al, 2014, Ooi & Prahlad, 2017, Pereira et al, 2020, Filipowicz et al, 2022, and others

over the past two decades have previously demonstrated that *C. elegans* are initially attracted

to PA14 without prior exposure to the pathogen ("naïve") but switch their behavior to avoidance

of PA14 following exposure to PA14. Our lab has also consistently replicated those naïve

attraction and learned avoidance results: in fact, we see that the mild naïve attraction to PA14 is

fairly consistent between individual plates and individual replicates over many experiments

across several studies (**Figure 1B-D, Table S1**) (Moore *et al*, 2019; Kaletsky *et al*, 2020; Moore

et al, 2021; Sengupta *et al*, 2024; Seto *et al*, 2024). This initial naïve attraction to PA14 serves

as a necessary baseline for all further experiments. That is, if no naïve attraction to PA14 is

observed and instead the worms already avoid PA14 prior to PA14 exposure, it might be difficult

to observe any additional learned avoidance of PA14 after training. Failure to observe naïve

attraction to PA14 might also suggest that the experimenters' assay conditions are not the same

as those used by previous groups. A recent study failed to consistently observe this initial naïve

avoidance of PA14 (Gainey *et al*, 2024); therefore, we investigated possible origins of this

deviation in these results from other groups' previously published results.

To determine possible origins of deviations in naïve choice assay results, we carried out the

two-bacteria choice assay under standard conditions (1hr of a binary choice of PA14 vs OP50,

which includes the paralytic sodium azide added to the bacterial spots) as well as the conditions

reported by Gainey et al., which primarily differs by omitting azide and letting the worms choose

in the absence of azide for 1 hour at room temperature, and then moving the plates to cold

temperature (4°C) prior to counting, as described by Gainey et al. (2024). We found that when

azide is present at the bacterial spots to trap naïve worms at their first choice, the worms

demonstrate attraction to PA14, as has been previously shown, but in the absence of azide,

worms show significant avoidance of PA14 even without prior exposure to the pathogen (**Figure**

1E-F); the difference between the +azide and -azide chemotaxis indices is both significant and

large in magnitude (a difference of 0.57 in CI and p<0.0001; **Figure 1F**). We note that these no-

azide results are almost identical to the results that Gainey et al. reported for their naive P0s,

while the +azide results are similar to our previously-published naïve data (**Figure 1B-D**), as

well as published data from other labs.

To better understand the source of the discrepancy with the Gainey et al. results, we first

investigated the worms' behavior at 4°C after the hour of the assay at room temperature; while

we do see that the worms continue to move (4°C does not immediately paralyze worms), in fact,

the final choice index of the population was already reached at the end of the room temperature

period *before* they were placed in the cold (**Figure 1G, t=0**); therefore, the discrepancy in

attraction to / avoidance of PA14 seems to have occurred during the initial part of the assay,

when the worms were choosing for 1 hour at room temperature without azide. This may be

particularly relevant for the Gainey et al. study, as they reportedly moved the worms at different

times ("*These no-azide assay plates were moved to 4°C after 30-60 minutes*"), which likely

induced further variance. We also observed worms still traveling away from the PA14 spot at

one hour, further underscoring the point that performing the assay without azide fails to trap the

worms at their first choice, even if the worms are transferred to 4°C, because the worms have

already made the switch to attraction to avoidance by that point.

The simplest interpretation of the discrepancy between the ±azide assay conditions is that

the worms are initially attracted to and choose PA14; in the presence of azide, the worms

remain at this first-choice spot, but over the course of an hour-long choice assay in the absence

of azide, the worms learn to avoid PA14, and migrate to the OP50 spot. This phenomenon was

already carefully demonstrated by Ooi and Prahlad (2017) with a time-course assay: in the

absence of azide to trap the worms, the worms are initially attracted to PA14, but by 45 min the

worms show no preference for PA14, and by one hour the worms instead avoid PA14 – that is,

within the time of the hour-long assay, in the absence of azide to trap the worms at their first

choice, they learn to avoid PA14. Under the no-azide assay conditions, it would be difficult to

observe any additional learned avoidance that is the result of training, since the learned

preference for OP50 (PA14) is already very high when the worms have been allowed to change

from their first choice to their second choice. Therefore, the simplest interpretation is that one

cannot accurately measure naïve preference in the absence of azide, and one cannot

subsequently properly assess subsequent learned avoidance after training without azide. It is

notable that the overwhelming majority of Gainey et al.'s experiments were done without azide;

because a third party (Ooi & Prahlad, 2017) already showed that there is a substantial

difference when no azide is used, as we have shown here, Gainey et al.'s no-azide data cannot

be compared with assays done with azide, including all of our work.

We repeated this ±azide preference assay with another *Pseudomonas* pathogen, *P.*
fluorescens 15 (PF15) that we have also found to induce transgenerational avoidance for 5
generations via small RNAs (Seto *et al*, 2024), and observed similar results (**Figure 1H, I**). That
is, the preference for PF15 is similar to worms' preference for PA14 over OP50, but in the
absence of azide, the worms learn to avoid PF15, as they did with PA14. Therefore, we
conclude that the inclusion of azide in the choice assay is necessary to properly assess the
worms' first choice; in the absence of azide, the worms' naïve choice is not being measured, so
it cannot be used to then compare with post-training behavior to assess learned or inherited
avoidance.
190

Transgenerational inheritance of learned avoidance is consistent and distinct from other
avoidance paradigms
C. elegans uses many different environmental cues to inform its behavioral decisions; in
fact, avoidance of PA14 can be induced by bacterial metabolites, by olfactory cues, by gustatory
cues, and by innate immunity pathways. These cues induce avoidance in the P0 generation,
and they and other mechanisms can induce avoidance and changes in survival or development
in the F1 generation via distinct mechanisms (Pereira *et al*, 2020; Hong *et al*, 2021; Burton *et al*,
2020; Pender *et al*, 2024), but do not extend beyond the F1 generation. Our studies used a
single condition for training, PA14 grown on plates at 25°C and training at 20°C for 24hrs (see
Figure 1A). Under these conditions, we consistently observe learned avoidance in mothers (P0;
Figure 2A, B, C) that is then transgenerationally-inherited (F2; **Figure 2D-F**). Therefore, our
PA14 growth and/or training conditions must be distinct from those growth and training
paradigms that only produce P0 or F1 effects.
204

P11 expression is condition-specific and critical for TEI of learned avoidance
In searching for the relevant TEI-inducing factor, we tested and eliminated DNA, proteins,
and large RNAs as the TEI-inducing component, and found that small RNAs isolated from PA14
grown at 25°C on plates could induce avoidance and TEI (Kaletsky *et al*, 2020). That is, small
RNAs isolated from 25°C plate-grown PA14 pipetted onto *E. coli* prior to 24hr *C. elegans*
training was able to induce avoidance in those animals and the transgenerational inheritance of
that avoidance, just as we had observed from 24hr PA14 lawn-training experiments. We had
also noticed that PA14 grown at lower temperatures or in liquid could not induce
transgenerational inheritance of learned avoidance. Capitalizing on these observations, we used

differential small RNA sequencing to identify the set of small RNAs that are expressed in PA14

grown at 25°C on plates that are not expressed in liquid or at low temperatures; we then cloned

each of those six small RNAs into *E. coli* and tested these individual sRNAs for their ability to

induce avoidance and TEI (Kaletsky *et al*, 2020). This set of experiments revealed that the small

RNA P11 – and only P11 – could induce avoidance in mothers and inheritance of avoidance for

four generations (Kaletsky *et al*, 2020). *C. elegans* trained on *E. coli* expressing P11 sRNA

induced the P0-F4 avoidance of PA14 without making the worms sick, and without inducing

innate immune pathways, but while also inducing *daf-7::gfp* expression in the ASI (Kaletsky *et*

al, 2020). We observed that *E. coli*-P11 (indicated by the yellow points) consistently induces P0

avoidance (**Figure 3A, B**) and this learned avoidance is inherited transgenerationally (**Figure**

3C, D). Therefore, P11 sRNA is both necessary and sufficient to induce the transgenerational

inheritance of learned PA14 avoidance.

We noted that P0 avoidance induced by *E. coli*-P11 training (**Figure 2A**) is slightly lower

than that induced by training on PA14 bacterial lawns (**Figure 3A, B**), suggesting that the P0

innate immunity-induced and other avoidance mechanisms are distinct from and additive with

the sRNA pathway in mothers. In fact, the average level of avoidance induced by *E. coli*-P11

training is consistent in P0 and F2 (**Figure 3C, D**) and with the F2 avoidance levels induced by

PA14 training (**Figure 2E, F)**, consistent with our previous reports (Kaletsky *et al*, 2020; Moore

et al, 2019). In fact, P11-induced avoidance remains consistent from P0 through F4 (Kaletsky *et*

al, 2020), as we later also found for the small RNAs (Pv1, Pfs1) required for *P. vranovensis*

(Sengupta *et al*, 2024) and *P. fluorescens* 15 (Seto *et al*, 2024) avoidance. Similarly, while *E.*

coli-P11 training induced *daf-7::gfp* expression in the ASI from P0 through F4 (Kaletsky *et al*,

2020), we did not observe any *daf-7::gfp* expression in the ASJ, consistent with the ASJ

response being specifically induced by metabolites produced by PA14 (Meisel *et al*, 2014).

Therefore, it appears that there are PA14 growth conditions that can induce some mechanisms

of P0 and F1 avoidance, but will not induce transgenerational avoidance, because no P11 is

expressed in PA14 under those conditions.

To more carefully address this hypothesis, we examined the levels of P11 sRNA from PA14

grown at 25° on plates, 15°C on plates, and 37°C in liquid (**Figure 3E**, adapted from Kaletsky et

al, 2020). When comparing 25°C plates with 15°C plates, we observed a 6.8-fold decrease in

P11 levels at the lower temperature, and a 13.7-fold decrease in P11 when PA14 is grown in

liquid rather than on plates, even at high temperatures. Therefore, it is clear that growth of PA14

in non-25°C/plate-grown conditions may fail to elicit a P11-mediated avoidance response,

because no P11 is expressed in PA14 under those conditions. Because Gainey et al. (2024) did

not verify P11 expression in their bacterial cultures, it is not surprising that they also failed to

observe P11-induced avoidance in the F2 generation in those experiments.

TEI of learned avoidance requires components of RNA processing pathways

Previously, we showed that many components of small RNA uptake, RNA interference, small

RNA processing pathways, and the germline are necessary for transgenerational inheritance of

learned avoidance (Moore *et al*, 2019; Kaletsky *et al*, 2020). We find that the *meg-1;meg-3*

double mutants, which perturb P granules and the germline (Wang *et al*, 2014), can avoid PA14

after lawn training, but are defective for small RNA-induced learning (worms trained on *E. coli*

with small RNA purified from PA14 added to the lawn) (**Figure 3F**). Similarly, NRDE-3, an

Argonaute protein that functions in nuclear RNAi (Guang *et al*, 2008), is also required for small

RNA-induced learned avoidance (**Figure 3G**). Like other mutants required for the small RNA

TEI pathway, the P0 generation of *meg-3;meg-4* and *nrde-3* mutants still avoid PA14 after

training on intact PA14 lawns (**Figure 3F, H**) through other avoidance pathways (innate

immunity, metabolites, etc.), once again highlighting the independence of these different

avoidance-inducing pathways. Only the small RNA-induced avoidance is transgenerational; that

is, F2s of wild-type worms continue to avoid PA14 after lawn training, but F2s of *nrde-3* mutants

are defective for this inherited small RNA-induced avoidance (**Figure 3J**).

We also find that *rde-3/mut-2*, which encodes a ribonucleotidyltransferase that is involved in

siRNA accumulation, piRNA-mediated silencing, and p(UG)ylation of small RNAs (Chen *et al*,

2005; Shukla *et al*, 2020; Priyadarshini *et al*, 2022) in epigenetic inheritance, is also required for

E. coli P11-induced learned avoidance (**Figure 3K**). Thus, components of small RNA transport,

small RNA processing, RNAi function, germline components, chromatin factors, neuronal

function, and the Cer1 retrotransposon (Moore *et al*, 2021) are required for the

transgenerational inheritance of learned pathogen avoidance (**Figure 4A**).

Discussion

C. elegans' physiological responses to PA14 and other pathogenic bacteria may allow the

worms to increase their chances of survival. Intergenerational responses, such as increased

survival (Burton *et al*, 2020; Pender *et al*, 2024), changes to development (Palominos *et al*,

2017), and internal physical responses (Singh & Aballay, 2019; Das *et al*, 2024) likely also

increase the possibility of the survival of their offspring. Switching behavioral responses from

attraction to avoidance for a few generations may also increase the survival of the species

(Moore *et al*, 2019) by reducing the chances of interactions with the pathogens. Of the many

cues that induce avoidance, only the bacterial small RNA pathway induces transgenerational

inheritance of learned pathogen avoidance, and this pathway is molecularly distinct from other

pathways that induce avoidance only in the P0 or F1 generations (Kaletsky *et al*, 2020). That is,

if PA14 is grown under conditions in which P11 is *not* expressed, the bacteria may still induce

P0 and even F1 avoidance due to the presence of olfactory, metabolite, and other cues, but

avoidance in the F2 generation and beyond will be absent. Therefore, if F2 avoidance is not

observed, it might be due to the lack of P11 expression in those conditions, and conversely,

avoidance in the P0 and F1 conditions should not be misconstrued as positive evidence of P11

expression.

P11 sRNA is a regulator of PA14's nitrogen assimilation operon, which includes nitrite and

nitrate reductases (NirB, NirD, and NasC) that control the production of ammonium (Marogi *et*

al, 2024). Nitrogen assimilation appears to be critical for growth on surfaces at specific

temperatures, including inside its host, *C. elegans* (Marogi *et al*, 2024). Nitrogen assimilation is

an energetically costly process, and PA14 tightly regulates this operon via control of the P11

expression. Interestingly, loss of P11 reduces ammonia production and thus reduces *C. elegans'*

chemoattraction to PA14 (Marogi *et al*, 2024); therefore, differences in *C. elegans* behavior

when treated with PA14 vs. ΔP11-PA14 mutants can be due to both the loss of the P11 small

RNA itself, and also due to the loss of ammonia production. Therefore, any growth conditions

that result in disruption of P11 expression might also contribute to decreases in naïve attraction

that Gainey et al. observed; that is, if PA14 is grown under conditions where P11 sRNA is not

expressed, not only would training on this bacteria not induce transgenerational inheritance of

avoidance due to the lack of P11 signal, but these conditions might also decrease *C. elegans'*

naïve attraction to this PA14, further reducing behavioral differences between naïve and trained

chemotaxis indices. Thus, growth conditions that decrease P11 expression might have a large

effect on both naïve and trained chemotaxis choice indices, and thus on interpretation of these

effects. Because P11 sRNA is both necessary and sufficient for the transgenerational

inheritance of pathogen avoidance, and its expression level is acutely sensitive to both

temperature and physical environmental conditions, it would be important to confirm P11

expression levels (e.g., via qPCR) in PA14 rather than assuming that P11 is constitutively

expressed, particularly when attempting to report the lack of an F2 effect. Unfortunately, Gainey

et al. (2024) never carried out any tests of P11 sRNA in PA14, so it is impossible to now know

how the PA14 growth conditions in any of their experiments might have affected P11 levels and

subsequent *C. elegans* responses. Additionally, learned avoidance in the P0 and F1 generations

has no relationship to the P11 pathway, and therefore cannot be used as confirmation of P11
expression.
In addition to P11, we also previously found that *C. elegans* ingestion and processing of
specific bacterial small RNAs produced by *P. vranovensis* (Sengupta *et al*, 2024) and *P.
fluorescens* 15 (Seto *et al*, 2024) induce pathogen avoidance that lasts through the F4
generation. These specific small RNAs are Pv1 and Pfs1 (from *P. vranovensis* and PF15,
respectively), and they each contain a stretch of either 17 (P11) or 16 (Pv1, Pfs1) nucleotides in
predicted stem loops that are a perfect match to specific exons of the gene *maco-1* (P11, Pv1)
or *vab-1* (Pfs1), which are then downregulated upon treatment with these pathogenic bacteria or
these small RNAs. In fact, editing the small RNAs to disrupt the perfect match to *maco-1* or *vab-
1* abrogates the learned avoidance (Sengupta *et al*, 2024; Seto *et al*, 2024), underscoring the
requirement for the perfect match sequence between the bacterial small RNAs and their target
C. elegans genes. MACO-1 is an ER resident protein that has been previously implicated in
regulation of chemotaxis, the dauer developmental decision, and thermotaxis (Arellano-Carbajal
et al, 2011; Miyara *et al*, 2011; Neal *et al*, 2016). *maco-1* downregulation upon PA14 treatment
(25°C on plates) or upon treatment with *E.coli* expressing P11 (*E.coli*-P11) results in
upregulation of *daf-7::gfp* in the ASI neurons (Kaletsky *et al*, 2020). Using qPCR, we confirmed
the downregulation of *maco-1* in P0 through the F4 generations and observed its return to
normal levels in the F5 generation (Sengupta *et al*, 2024), corresponding with avoidance
behavior. Furthermore, mutants of *maco-1* exhibit high naïve avoidance of PA14 and high levels
of *daf-7::gfp* expression in the ASI prior to training, and because of this high naïve avoidance,
they are unable to learn further avoidance upon training on PA14 (Kaletsky *et al*, 2020). P0
worms treated with *maco-1* RNAi exhibit transgenerational inheritance of learned avoidance, as
well (Kaletsky *et al*, 2020). The downregulation of *vab-1* upon PF15 or Pfs1 sRNA treatment
also results in the downregulation of *maco-1* expression levels (as shown by RNA-seq in P0
mothers), suggesting that *vab-1* is upstream of *maco-1*; qPCR confirmed that *maco-1* is
downregulated in the F2 generation of *E. coli*-Pfs1-trained mothers as well (Seto *et al*, 2024).
Therefore, the transgenerational bacterial small RNA–*C. elegans* avoidance mechanism
induced by at least three different *Pseudomonas* species through three different small RNAs
requires the downregulation of *maco-1*, which results in *daf-7* changes in the ASI; perfect
matches between bacterial small RNAs and their *C. elegans* target genes are required for the
F2-F5 heritable shift from attraction to avoidance. Thus, the role of *maco-1* in this pathway has
been confirmed through multiple avenues of experimentation.

Through extensive mutant analyses (testing of almost 40 mutants over the course of six
studies), we have found that this sRNA mechanism of transgenerational inheritance of learned
pathogen avoidance requires a wide range of machinery involved in bacterial small RNA uptake,
RNA processing, signaling, and neuronal behavioral execution (**Figure 4A**). These genes
include RNA interference components, Argonautes, and RNA-dependent polymerases; germline
components such as P-granule, piRNA components, COMPASS and chromatin modifiers*;* the
Cer1 Ty3 Gypsy retrotransposon; and neuronal genes (*maco-1*/macoilin, *vab-1/*Ephrin receptor,
vab-2/Ephrin) as well as the *daf-7*/TGF-beta ligand in the ASI neuron (**Figure 4A**). We also
ruled out several genes that function in other forms of pathogen avoidance, such as imprinting,
innate immunity signaling, microRNA and viral RNAi processors, and other transposable
elements, further distinguishing the bacterial small RNA TEI pathway from other avoidance
mechanisms that may only function in the P0 or F1 generations (**Figure 4A, B**). Through tissue-
specific rescue experiments we also showed that *dcr-1* functions in the intestine and *prg-1* is
necessary in the germline for small RNA-mediated learned pathogen avoidance. Our studies of
other bacterial species have identified not only additional bacterial small RNAs that are "read"
by *C. elegans* (Sengupta *et al*, 2024; Seto *et al*, 2024), but also have identified additional *C.*
elegans gene targets that function in the pathway (e.g., *vab-1*, which regulates *maco-1*; Seto et
al. 2024). It should be noted that in each of these mutant experiments, wild-type worms were
tested as controls; therefore, our wild-type naïve attraction and learned avoidance results have
been replicated many times over the course of these six studies.
368
<u>Conclusions</u>
Here we have investigated two experimental factors that might affect investigators'
interpretation of PA14 training-induced avoidance choice assays: the expression of P11 sRNA in
PA14 and proper execution of the choice assay itself. It would be incorrect to assume that
because F2 avoidance is not demonstrated under all PA14 growth conditions that "*the published*
model is unlikely to be relevant in a natural environment" (Gainey *et al*, 2024). Instead, our
results suggest the opposite: that the worms' ability to interpret P11 expression is due to P11's
importance in regulating nitrogen fixation under biofilm-inducing conditions that also induce a
pathogenic state that is important for *C. elegans* to remember over several generations. *C.*
elegans appears to have evolved a mechanism to use P11 and other specific small RNAs as
biomarkers of future infection, as the presence of those specific small RNAs correlates well with
pathogenesis – and the specific small RNA detected by *C. elegans* in each case might be tightly
correlated with pathogenic potential of that particular *Pseudomonas* species. Furthermore, the

fact that we found that other *Pseudomonas* species, including *Pseudomonas* from the worm's

microbiota, use the same bacterial sRNA/*C. elegans* target gene mechanism indicates that this

natural example of trans-kingdom signaling and transgenerational avoidance is highly

conserved. In those bacteria, the key small RNAs appear to be constitutively expressed, and

thus even more likely to be encountered by *C. elegans* upon ingestion. Together, these data

suggest that the transgenerational inheritance of small RNA-induced learned bacterial

avoidance is likely to be widespread in the wild, indicating its likely importance as a mechanism

to promote long-term survival of *C. elegans* in its natural habitat.

Methods

General worm maintenance

Worm strains were maintained at 20°C on high growth medium (HG) plates (3 g/L NaCl, 20 g/L

bacto-peptone, 30 g/L bacto-agar in distilled water, with 4 mL/L cholesterol (5 mg/mL in ethanol),

1 mL/L 1 M CaCl$_2$, 1 mL/L 1 M MgSO$_4$ and 25 mL/L 1 M potassium phosphate buffer (pH 6.0)

added to molten agar after autoclaving) on OP50 using standard methods. The following worm

strains were provided by the *C. elegans* Genetics Center (CGC): JH3229, *meg-1(vr10);meg-*

3(tm4259) X; YY158, *nrde-3(gg66)* X; WM30, *mut-2/rde-3(ne298)* I.

***E. coli* and PA14 culture conditions.**

OP50 was provided by the CGC. PA14 was a gift from Z. Gitai. OP50 and PA14 were grown in

liquid cultures in LB at 37°C with shaking. *Escherichia coli* strains expressing PA14 sRNA were

cultured overnight in Luria broth supplemented with 0.02% arabinose w/v and 100 µg/mL

carbenicillin. Overnight cultures of bacteria were diluted in LB to an optical density at 600 nm

(OD 600) = 1 and used to fully cover nematode growth medium (NGM) (3 g/L NaCl, 2.5 g/L

bacto-peptone, 17g/L bactoagar in distilled water, with 1 ml/L cholesterol (5 mg/mL in ethanol), 1

mL/L 1 M CaCl$_2$, 1 mL/L 1 M MgSO$_4$ and 25 mL/L 1 M potassium phosphate buffer (pH 6.0)

added to molten agar after autoclaving) plates. For preparation of control *E. coli* and *E. coli*

expressing the PA14 P11 small RNA, bacteria were seeded on NGM plates supplemented with

0.02% arabinose and 100 µg/mL carbenicillin. All plates were incubated for 2 days at 25 °C,

unless specified otherwise (in separate incubators for control and pathogen-seeded plates).

OP50 and PA14 small RNAs were collected as described in Kaletsky et al. 2020.

Aversive learning assay

Worm preparation for training: synchronized L4 worms were washed off plates using M9 and left

to pellet on the bench top for approximately 5 min. Then, 5 µL of worms were placed onto

sRNA-spotted training plates, and 10 µL or 40 µL of worms were plated onto OP50 or *E. coli*

expressing control/P11, or PA14-seeded training plates, respectively. Worms were incubated on

training plates at 20 °C in separate containers for 24 h. After 24 h, worms were washed off

plates using M9 and washed an additional 3 times to remove excess bacteria.

Choice plates and naïve day 1 worms were prepared as in Moore et al. 2019 and Seto et al.

2024. On the day of the assay, choice assay plates were left at room temperature for 1 h before

use. For standard (sodium azide added) experiments, 1 µL of 1 M sodium azide was spotted

onto each respective bacteria spot to be used as a paralyzing agent during choice assay and

preserve first bacterial choice. For conditions without azide, no azide was added.

Worms were washed off training plates in 2 mL M9 into 1.5 mL tubes and allowed to pellet by

gravity. Worms were washed 2-3 additional times in M9. Using a wide orifice pipet tip, 5 µL of

worms (approximately 50-200 worms) were spotted at the bottom of the assay plate, midway

between the bacterial spots. Plates were incubated at room temperature for 1 h before moving

plates to 4C. Plates were left at 4C for ~1h before manually counting the number of worms on

each bacterial spot.

Meta-analysis

We collected all examples of N2 naïve chemotaxis (PA14 vs OP50 choice assays) prior to

training, N2 P0 and F2 choice assays after P0 training on PA14 vs OP50, and P0 and F2 choice

assays after P0 training on *E. coli*-P11 vs control *E. coli* from Moore et al. 2019, Kaletsky et al.

2020, and Moore et al. 2021. These data are collected in Table S2. Multiple two-group

estimation plots were generated for side-by-side comparison of multiple mean differences

across independent experiments. Each mean difference is plotted as a bootstrap sampling

distribution.

References

References

Arellano-Carbajal F, Briseño-Roa L, Couto A, Cheung BHH, Labouesse M & De Bono M (2011) Macoilin, a
Conserved Nervous System–Specific ER Membrane Protein That Regulates Neuronal Excitability.
PLoS Genet 7: e1001341

Bargmann CI, Hartwieg E & Horvitz HR (1993) Odorant-selective genes and neurons mediate olfaction in
C. elegans. *Cell* 74: 515–527

Burton NO, Riccio C, Dallaire A, Price J, Jenkins B, Koulman A & Miska EA (2020) Cysteine synthases CYSL-
1 and CYSL-2 mediate C. elegans heritable adaptation to P. vranovensis infection. *Nat Commun*
11: 1741

Chen C-CG, Simard MJ, Tabara H, Brownell DR, McCollough JA & Mello CC (2005) A Member of the
Polymerase β Nucleotidyltransferase Superfamily Is Required for RNA Interference in C. elegans.
Curr Biol 15: 378–383

Das P, Aballay A & Singh J (2024) Calcineurin inhibition enhances Caenorhabditis elegans lifespan by
defecation defects-mediated calorie restriction and nuclear hormone signaling. *eLife* 12:
RP89572

Félix M-A & Duveau F (2012) Population dynamics and habitat sharing of natural populations of
Caenorhabditis elegans and C. briggsae. *BMC Biol* 10: 59

Filipowicz A, Lalsiamthara J & Aballay A (2022) Dissection of a sensorimotor circuit underlying pathogen
aversion in C. elegans. *BMC Biol* 20: 229

Gainey DP, Shubin AV & Hunter CP (2024) Reported transgenerational responses to Pseudomonas
aeruginosa in C. elegans are not robust. doi:10.7554/eLife.100254.2 [PREPRINT]

Guang S, Bochner AF, Pavelec DM, Burkhart KB, Harding S, Lachowiec J & Kennedy S (2008) An Argonaute
Transports siRNAs from the Cytoplasm to the Nucleus. *Science* 321: 537–541

Ha H, Hendricks M, Shen Y, Gabel CV, Fang-Yen C, Qin Y, Colón-Ramos D, Shen K, Samuel ADT & Zhang Y
(2010) Functional Organization of a Neural Network for Aversive Olfactory Learning in
Caenorhabditis elegans. *Neuron* 68: 1173–1186

Hao Y, Yang W, Ren J, Hall Q, Zhang Y & Kaplan JM (2018) Thioredoxin shapes the C. elegans sensory
response to Pseudomonas produced nitric oxide. *eLife* 7: e36833

Hart A (2006) Behavior. *WormBook*

Hong C, Lalsiamthara J, Ren J, Sang Y & Aballay A (2021) Microbial colonization induces histone
acetylation critical for inherited gut-germline-neural signaling. *PLOS Biol* 19: e3001169

Horspool AM & Chang HC (2017) Superoxide dismutase SOD-1 modulates C. elegans pathogen avoidance
behavior. *Sci Rep* 7: 45128

Jin X, Pokala N & Bargmann CI (2016) Distinct Circuits for the Formation and Retrieval of an Imprinted
Olfactory Memory. *Cell* 164: 632–643

Kaletsky R, Moore RS, Vrla GD, Parsons LR, Gitai Z & Murphy CT (2020) C. elegans interprets bacterial
491   non-coding RNAs to learn pathogenic avoidance. *Nature* 586: 445–451

492   Lee K & Mylonakis E (2017) An Intestine-Derived Neuropeptide Controls Avoidance Behavior in
493   Caenorhabditis elegans. *Cell Rep* 20: 2501–2512

494   Liu H, Yang W, Wu T, Duan F, Soucy E, Jin X & Zhang Y (2018) Cholinergic Sensorimotor Integration
495   Regulates Olfactory Steering. *Neuron* 97: 390-405.e3

496   Liu H & Zhang Y (2020) What can a worm learn in a bacteria-rich habitat? *J Neurogenet* 34: 369–377

497   Ma Y-C, Zhang L, Dai L-L, Khan RU & Zou C-G (2017) mir-67 regulates P. aeruginosa avoidance behavior in
498   C. elegans. *Biochem Biophys Res Commun* 494: 120–125

499   Mahajan-Miklos S, Tan M-W & Rahme LG (1999) Molecular Mechanisms of Bacterial Virulence
500   Elucidated Using a Pseudomonas aeruginosa– Caenorhabditis elegans Pathogenesis Model. *Cell*
501   96: 47–56

502   Marogi JG, Murphy CT, Myhrvold C & Gitai Z (2024) Pseudomonas aeruginosa modulates both
503   Caenorhabditis elegans attraction and pathogenesis by regulating nitrogen assimilation. *Nat*
504   *Commun* 15: 7927

505   Meisel JD, Panda O, Mahanti P, Schroeder FC & Kim DH (2014) Chemosensation of Bacterial Secondary
506   Metabolites Modulates Neuroendocrine Signaling and Behavior of C. elegans. *Cell* 159: 267–280

507   Melo JA & Ruvkun G (2012) Inactivation of Conserved C. elegans Genes Engages Pathogen- and
508   Xenobiotic-Associated Defenses. *Cell* 149: 452–466

509   Miller EV, Grandi LN, Giannini JA, Robinson JD & Powell JR (2015) The Conserved G-Protein Coupled
510   Receptor FSHR-1 Regulates Protective Host Responses to Infection and Oxidative Stress. *PLOS*
511   *ONE* 10: e0137403

512   Miyara A, Ohta A, Okochi Y, Tsukada Y, Kuhara A & Mori I (2011) Novel and Conserved Protein Macoilin Is
513   Required for Diverse Neuronal Functions in Caenorhabditis elegans. *PLoS Genet* 7: e1001384

514   Moore RS, Kaletsky R, Lesnik C, Cota V, Blackman E, Parsons LR, Gitai Z & Murphy CT (2021) The role of
515   the Cer1 transposon in horizontal transfer of transgenerational memory. *Cell* 184: 4697-
516   4712.e18

517   Moore RS, Kaletsky R & Murphy CT (2019) Piwi/PRG-1 Argonaute and TGF-β Mediate Transgenerational
518   Learned Pathogenic Avoidance. *Cell* 177: 1827-1841.e12

519   Neal SJ, Park J, DiTirro D, Yoon J, Shibuya M, Choi W, Schroeder FC, Butcher RA, Kim K & Sengupta P
520   (2016) A Forward Genetic Screen for Molecules Involved in Pheromone-Induced Dauer
521   Formation in *Caenorhabditis elegans*. *G3 GenesGenomesGenetics* 6: 1475–1487

522   Ooi FK & Prahlad V (2017) Olfactory experience primes the heat shock transcription factor HSF-1 to
523   enhance the expression of molecular chaperones in *C. elegans*. *Sci Signal* 10: eaan4893

Palominos MF, Verdugo L, Gabaldon C, Pollak B, Ortíz-Severín J, Varas MA, Chávez FP & Calixto A (2017)
Transgenerational Diapause as an Avoidance Strategy against Bacterial Pathogens in
Caenorhabditis elegans. mBio 8: e01234-17

Pender CL, Dishart JG, Gildea HK, Nauta KM, Page EM, Siddiqi TF, Cheung SS, Joe L, Burton NO & Dillin A
(2024) Perception of a pathogenic signature initiates intergenerational protection. *Cell*:
S0092867424013424

Pereira AG, Gracida X, Kagias K & Zhang Y (2020) *C. elegans* aversive olfactory learning generates diverse
intergenerational effects. *J Neurogenet* 34: 378–388

Prakash D, Ms A, Radhika B, Venkatesan R, Chalasani SH & Singh V (2021) 1-Undecene from
Pseudomonas aeruginosa is an olfactory signal for flight-or-fight response in *Caenorhabditis*
elegans. EMBO J 40: e106938

Priyadarshini M, AlHarbi S & Frøkjær-Jensen C (2022) *rde-3* reduces piRNA-mediated silencing and
abolishes inherited silencing in *C. elegans*. doi:10.1101/2022.08.13.503839 [PREPRINT]

Samuel BS, Rowedder H, Braendle C, Félix M-A & Ruvkun G (2016) *Caenorhabditis elegans* responses to
bacteria from its natural habitats. *Proc Natl Acad Sci* 113

Sengupta T, St. Ange J, Kaletsky R, Moore RS, Seto RJ, Marogi J, Myhrvold C, Gitai Z & Murphy CT (2024) A
natural bacterial pathogen of C. elegans uses a small RNA to induce transgenerational
inheritance of learned avoidance. *PLOS Genet* 20: e1011178

Seto RJ, Brown, Rachel, Kaletsky R, Parsons LR, Moore RS & Murphy CT (2024) Pseudomonas fluorescens
15 small RNA Pfs1 mediates transgenerational epigenetic inheritance of pathogen avoidance in
C. elegans through the Ephrin receptor VAB-1. [PREPRINT]

Shukla A, Yan J, Pagano DJ, Dodson AE, Fei Y, Gorham J, Seidman JG, Wickens M & Kennedy S (2020)
poly(UG)-tailed RNAs in genome protection and epigenetic inheritance. *Nature* 582: 283–288

Singh J & Aballay A (2019) Intestinal infection regulates behavior and learning via neuroendocrine
signaling. *eLife* 8: e50033

Tan M-W, Mahajan-Miklos S & Ausubel FM (1999) Killing of *Caenorhabditis elegans* by *Pseudomonas*
aeruginosa used to model mammalian bacterial pathogenesis. *Proc Natl Acad Sci* 96: 715–720

Troemel ER, Chu SW, Reinke V, Lee SS, Ausubel FM & Kim DH (2006) p38 MAPK Regulates Expression of
Immune Response Genes and Contributes to Longevity in C. elegans. *PLoS Genet* 2: e183

Wang JT, Smith J, Chen B-C, Schmidt H, Rasoloson D, Paix A, Lambrus BG, Calidas D, Betzig E & Seydoux G
(2014) Regulation of RNA granule dynamics by phosphorylation of serine-rich, intrinsically
disordered proteins in C. elegans. *eLife* 3: e04591

Wolfe GS, Tong VW, Povse E, Merritt DM, Stegeman GW, Flibotte S & Van Der Kooy D (2019) A Receptor
Tyrosine Kinase Plays Separate Roles in Sensory Integration and Associative Learning in *C.*
elegans. eneuro 6: ENEURO.0244-18.2019

Zhang Y, Lu H & Bargmann CI (2005) Pathogenic bacteria induce aversive olfactory learning in
Caenorhabditis elegans. Nature 438: 179–184

Figure Legends

Figure 1: **The omission of the paralytic sodium azide from choice assays has significant**

effects on naïve choice assays. (**a**) Schematic of PA14 and worm growth conditions, worm

training, and choice assay parameters. Critical factors affecting assay performance are

highlighted. (**b**) Scatter dot plots of representative replicate choice assays (OP50 versus PA14

bacteria) from naïve Day 1 adult worms (from Moore et al. 2019, Kaletsky et al. 2020, and

Moore et al. 2021). Each dot represents an individual choice assay plate (n = 10) containing

~50-100 worms per plate (average ~80). Choice index = (number of worms on OP50 – number

of worms on PA14)/(total number of worms). (**c**) Individual replicates of naïve choice assays

from Moore et al., 2019, Kaletsky et al., 2020, and Moore 2021 are shown as violin plots. Each

replicate is a separate experiment made up of 5-10 plates. The median is shown in red. (**d**)

Individual replicates from (c) were pooled. Box plots: center line, median; box range, 25–75th

percentiles; whiskers denote minimum–maximum values. (**e**) Naïve worms were placed on

choice assay plates with spots of PA14 and OP50 with or without sodium azide on the bacterial

spots, then allowed to choose for one hour, followed by one hour at 4°C. No azide: CI = 0.4 ±

0.06, p<0.0001. (**f**) Estimation plot of difference in CI between azide and no azide plates; mean

difference = 0.482 [95.0%CI 0.369, 0.64]. (**g**) After 1 hr of the choice assay at room

temperature, choice assay plates were placed at 4°C. Worms were counted at each spot at the

indicated time points. Mean +/- SD from three choice assay plates. Timepoint 0 = immediately

before 4°C incubation. (**h**) Worms were placed on choice assay plates with spots of PF15 and

OP50 with or without sodium azide on the bacterial spots, then allowed to choose for one hour,

followed by one hour at 4°C. No azide CI = 0.26 ± 0.05, p<0.0001. (**i**) Estimation plot of

difference in CI between azide and no azide plates; mean difference = 0.547 [95.0%CI 0.348,

0.737].

Figure 2: **P0 and F2 worms trained on PA14 bacteria reproducibly learn to avoid PA14 in**

choice assays. a) P0 replicates: A representative experiment (replicate) showing the choice

index and effect size of 24h PA14-trained worms compared to the OP50-trained control. (**b**)

Mean differences of individual replicates from Moore et al. 2019, Kaletsky et al. 2020, and

Moore et al. 2021 are shown for PA14- vs OP50-trained mothers. (**c**) Individual replicates from

(**b**) were pooled. (**d-e**) F2 animals from P0-control or PA14-trained worms. (**d**) F2 replicates: A

representative experiment (replicate) showing the choice index and effect size of F2 worms. (**a,**

d) The mean difference is shown as a Gardner-Altman estimation plot. Both groups are plotted

on the left axes; the mean difference is plotted on floating axes on the right as a bootstrap

sampling distribution. The mean difference is depicted as a dot; the 95% confidence interval is

indicated by the ends of the vertical error bar. (**e**) Mean differences of individual replicates from

Moore et al. 2019, Kaletsky et al. 2020, and Moore et al. 2021 are shown for F2s from PA14- vs

OP50-trained grandmothers. (**b, e**) Box plots: center line, median; box range, 25–75th

percentiles; whiskers denote minimum–maximum values. Unpaired, two-tailed Student's *t*-test.

****$P < 0.0001$. (**f**) Individual replicates from (**e**) were pooled. (**c, f**) Each replicate is a separate

experiment made up of 5-10 plates (average ~80). The mean difference (effect size) is shown

as a Cumming estimation plot. Each mean difference is plotted as a bootstrap sampling

distribution. Mean differences are depicted as dots; 95% confidence intervals are indicated by

the ends of the vertical error bars. Estimation graphics generated as described in Ho et al.,

2019.

Figure 3: **P11 sRNA and components of germline and RNAi function are required for**

transgenerational inheritance. (**a**) P0 worms trained on *E. coli* expressing the P11 small RNA

reproducibly avoid PA14 in choice assays. Individual replicates from Kaletsky et al. 2020, and

Moore et al. 2021 are shown for P0 worms obtained from P11 vs control training. (**b**) Individual

replicates from (**a**) were pooled. (**c**) F2 animals from P0 worms trained on *E. coli* expressing the

P11 small RNA reproducibly avoid PA14 in choice assays. Individual replicates from Kaletsky et

al. 2020, and Moore et al. 2021 are shown for F2 worms obtained from P11- vs control-trained

grandmothers. (**d**) Individual replicates from (**a**) were pooled. (**a, c**) Each replicate is a separate

experiment made up of 5-10 plates (average ~80). The mean difference (effect size) is shown

as a Cumming estimation plot. Each mean difference is plotted as a bootstrap sampling

distribution. Mean differences are depicted as dots; 95% confidence intervals are indicated by

the ends of the vertical error bars. (**e**) Normalized RNA counts for P11 expression from PA14

under different bacterial growth conditions. Sequencing data and figure adapted from Kaletsky

et al., 2020. Adjusted p-values from DESeq2. (**f**) P0 *meg-1;meg-3* mutants exhibit normal

learned PA14 avoidance when trained on PA14 lawns, but (**g**) P0 *meg-1;meg-3* mutants do not

exhibit learned PA14 avoidance when trained with PA14 sRNA. (**h**) P0 *nrde-1(gg66)* mutants

exhibit normal learned PA14 avoidance when trained on PA14 lawns, (**i**) but are defective for

PA14 avoidance learning when exposed to PA14 sRNA. (**j**) F2 *nrde-3* mutants from P0-PA14

lawn-trained grandmothers do not avoid PA14 compared to wild type controls. (**k**) Wild-type P0

worms exposed to *E. coli*-expressing P11 avoid PA14, but *rde-3* mutants do not. (**b, d-k**) Box

plots: center line, median; box range, 25–75th percentiles; whiskers denote minimum–maximum

values. (**b, d**) Unpaired, two-tailed Student's *t*-test. (f-k) Two-way ANOVA with Tukey's multiple

comparison's test. *$P \le 0.05$, ***$P \le 0.001$, ****$P < 0.0001$, NS, not significant. Estimation

graphics generated as described in Ho et al., 2019.

Figure 4: **Summary of genes tested for small RNA-mediated pathogen avoidance and**

characterization of PA14 avoidance-based assays. (**a**) Summary of genes tested for naïve

Pseudomonas preference behavior and small-RNA based learning and inheritance. (**b**) Table

describing different assays measuring *Pseudomonas* learning and avoidance. *Pseudomonas*

avoidance learning in the parental generation (P0), intergenerationally (F1), and

transgenerationally (F2 through F4) is noted. References: 1, Meisel et al. 2014. 2, Singh &

Aballay, 2019. 3, Zhang et al. 2005. 4, Pereira et al. 2020. 5, Moore et al. 2019. 6, Kaletsky et

al. 2020. 7, Moore et al. 2021. 8, Sengupta et al. 2024. 9, Seto et al. 2024.

Table S1: Summary data compiled from Moore et al. 2019, Kaletsky et al. 2020, and Moore

et al. 2021.

Table S2: Figure 3 worms counts from choice assays.

Figure 1

Figure 2

Figure 3

A

Category	Gene	Normal naïve preference	Small-RNA learning	References
	wild type	+	+	Moore et al, 2019; Kaletsky et al, 2020, Moore et al, 2021; Sengupta et al, 2024; Seto et al, 2024
imprinting/memory	crh-1	+	+	Moore et al, 2019
immunity	pmk-1	+	+	Moore et al, 2019; Kaletsky et al, 2020
	vhp-1	+	+	Sengupta et al, 2024; Seto et al, 2024
miRNA	alg-1	+	+	Kaletsky et al, 2020
viral RNAi	drh-1	+	+	Kaletsky et al, 2020
LTR TE	cer-4	+	+	Kaletsky et al, 2020
p granules	meg-3*	+	+	Kaletsky et al, 2020
	dcr-1 int. rescue	+	+	Kaletsky et al, 2020
	prg-1 GL rescue	+	+	Kaletsky et al, 2020
RNAi transport	sid-2	+	-	Moore et al, 2019; Kaletsky et al, 2020, Moore et al, 2021; Sengupta et al, 2024; Seto et al, 2024
	sid-1	+/-**	-	Moore et al, 2019; Kaletsky et al, 2020, Moore et al, 2021; Sengupta et al, 2024
RNAi/sRNA processing & amplification	dcr-1	+	-	Kaletsky et al, 2020; Sengupta et al, 2024
	rde-3	+	-	this paper
	rde-2	+	-	Kaletsky et al, 2020
	rde-4	+	-	Kaletsky et al, 2020
	mut-7	+	-	Kaletsky et al, 2020
	rrf-1	+	-	Kaletsky et al, 2020
	rrf-3	+	-	Kaletsky et al, 2020
germline	glp-1	+	-	Kaletsky et al, 2020, Moore et al, 2021
p granules	meg-1;meg-3	+	-	this paper
	meg-3;meg-4	+	-	Kaletsky et al, 2020
piRNAs	prg-1	+	-	Moore et al, 2019; Kaletsky et al, 2020
argonautes	hrde-1	+	-	Moore et al, 2019; Sengupta et al, 2024
	nrde-3	+	-	this paper
	rde-1	+	-	Kaletsky et al, 2020
chromatin modifiers	set-25	+	-	Moore et al, 2019
	hpl-2	+	-	Moore et al, 2019; Kaletsky et al, 2020
LTR TE	Cer1	+/-	-	Kaletsky et al, 2020, Moore et al, 2021; Sengupta et al, 2024; Seto et al, 2024
neuron function	daf-7	+	-	Moore et al, 2019; Kaletsky et al, 2020, Moore et al, 2021; Sengupta et al, 2024; Seto et al, 2024
nuclear RNAi	nrde-4	-	-	Moore et al, 2019
chromatin modifiers	set-2	-	-	Moore et al, 2019
	rbr-2	-	-	Moore et al, 2019
	set-32	-	-	Moore et al, 2019
	wdr-5.1	-	-	Moore et al, 2019
neuron function	maco-1	-	-	Kaletsky et al, 2020, Moore et al, 2021; Sengupta et al, 2024; Seto et al, 2024
	ASI ablation	-	-	Moore et al, 2019
ephrin signaling	vab-1	-	-	Seto et al, 2024
	vab-2	-	-	Seto et al, 2024

B

Figure 4