1 **Characterizing venom resistance in *Monodelphis domestica* yields new**
2 **insights into mammalian blood physiology**
3
4
5

6 Matthew Holding*[2,3], Dante Disharoon*[4], Laura Haynes*[2], Bipin Chakravarthy Paruchuri[4], M. Hao Hao
7 Pontius[7], Krista Golden[7], Jordan A. Shavit[7,8], Karl Desch[7], David Ginsburg[2,7,8,9], Anirban Sen Gupta[4], Yolanda
8 Cruz[6], Danielle H. Drabeck[1]
9
10 *Joint first authors
11 Corresponding author: Danielle H. Drabeck
12
[1]Department of Ecology, Evolution, and Behavior, University of Minnesota, Twin Cities
[2]Life Sciences Institute, University of Michigan, Ann Arbor, MI
[3]Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI, USA
[4]Department of Biomedical Engineering, Case Western Reserve University, Cleveland, OH 44106, USA.
[6]Oberlin College Department of Biology, Oberlin, OH 44074, USA
[7]Department of Pediatrics, University of Michigan, Ann Arbor, MI, USA
[8]Departments of Human Genetics, University of Michigan, Ann Arbor, MI, USA
[9]Departments of Internal Medicine, University of Michigan, Ann Arbor, MI, USA

Interactions between predators and prey are often characterized by strong selection that
shapes extreme physiological adaptations for survival in hostile environments. Venom
resistance in large-bodied South American opossums (Clade Didelphini) provides a striking
example as they are well known to prey upon snakes and have strong resistance to extremely
toxic snake venom. While resistance is well known in this group of large-bodied opossums,
relatively little is known about the venom resistance of the remaining diverse and speciose
members of Didelphidae (small-bodied opossums) that inhabit the same regions in South
America and encounter the same venomous snakes. Here, we investigate the venom resistance
of *Monodelphis domestica* through multi-level physiological assays, examining responses to
both purified venom components and whole venom from sympatric and allopatric vipers. Our
results reveal that *M. domestica* exhibits significant resistance to venom-induced blood
coagulation, retains normal platelet function in the presence of venoms that are known to
disrupt platelet function, and has enhanced inhibition of snake venom metalloproteinases.
Unexpectedly, we find a previously unknown modification of opossum blood physiology, namely,
that opossum von Willebrand Factor (VWF) requires increased shear force to elongate, which
may contribute to the venom resistance phenotype and has relevance to human coagulopathic
disease. Together these data expand the known extent of venom resistance beyond the large-
bodied clade, suggesting it is a widespread adaptive trait in South American marsupials. This
work provides new insights into venom-mammal coevolution by expanding our understanding of
the physiological adaptations underlying venom resistance including how they can inspire
biomedical solutions to coagulation disorders.

Introduction

Interactions between venomous species and their enemies offer striking examples of

coevolution, where molecular adaptations on both sides drive dynamic physiological changes.

Mammalian venom resistance represents an extreme outcome of coevolution, where molecular

changes enable mammals to withstand the injection of deadly toxins from venomous species.

Such systems offer a powerful model for understanding how molecular changes—like those in

proteins targeted by venom— are shaped by strong selective pressure, impose costs on

physiological function, and facilitate adaptive ecologies.

South American large-bodied opossums in the clade Didelphini are well-known to be

resistant to venomous pit-vipers with which they are sympatric (Jansa and Voss 2011; Drabeck

et al. 2020, 2022). Large-bodied opossums in this clade regularly prey upon venomous snakes,

and resistance to venom is thought to be primarily a trophic adaptation which allows them to

exploit an otherwise lethal prey resource (Voss and Jansa 2012). Venom resistance in the large-

bodied opossums has been relatively well characterized, and is known to be especially effective

against snakes in the genus *Bothrops*, which are the most abundant and widely distributed

venomous snakes across South America (Werner and Vick 1977; Voss and Jansa 2012; Holding

et al. 2016b; Engelman et al. 2017; Drabeck et al. 2020; Voss and Jansa 2021; Drabeck et al.

2022).

Small-bodied South American opossums are both abundant and speciose across the

same range of venomous snakes and have long been assumed to be venom-susceptible prey

(Mackessy 2021), rather than predators, of venomous snakes (notable exception (Voss 2013)).

Consequently, the role of small-bodied opossums in the ecology and evolution of venom (and

resistance to venom), has been previously assumed to be inconsequential. However, recent

research has identified resistance to botrocetin, a venom C-type lectin (CTL) from *Bothrops*

jararaca, in a small-bodied species (*Monodelphis domestica)* suggesting that this species may

enjoy venom resistance similar to its large-bodied relatives in Didelphini (Drabeck et al. 2020).

In venom-susceptible mammals, botrocetin induces coagulopathy by binding to von

Willebrand Factor (VWF), triggering VWF to interact with the platelet receptor GP1bα (Fukuda et

al. 2002). VWF is a multimeric protein that circulates in plasma in a globular conformation.

During vascular injury, VWF becomes exposed to elevated shear stresses which causes it to

unravel, exposing its A domains (Lenting et al. 2024). In the absence of venom proteins, VWF

plays a vital physiologic role with platelets to prevent unwanted bleeding via two crucial

mechanisms: 1) its A3 domain binds to subendothelial collagen (exposed by vessel injury),

anchoring it to the injury site, and 2) its A1 domain binds to platelet GP1bα receptor, rendering

rapid platelet accumulation (Lenting et al. 2024). Upon binding to VWF, platelets become

activated and these aggregate with each other via fibrinogen binding through integrin GPIIb/IIIa,

resulting in the formation of a platelet plug. Venom-derived VWF-binding CTLs cause damage by

binding to the A1 domain of VWF and promoting complexation with platelets even in low shear

conditions (Matsushita et al. 2000; Fukuda et al. 2002). The resulting CTL-VWF-platelet complex

gets rapidly cleared from circulation, thereby depleting VWF and platelets from circulation,
rendering them unavailable for hemostatic platelet plug formation (extreme thrombocytopenia),
causing excessive bleeding and death (Matsushita et al. 2000; Fukuda et al. 2002; Bury et al.
2019; Thomazini et al. 2021). This disruption in platelet function is thought to work
synergistically with hemorrhagic metalloproteinases in *Bothrops* venom, which degrade
vascular tissues by cleaving proteins in the basal lamina, causing widespread bleeding and
tissue damage.
Comparative analyses have identified mutations in VWF coding sequences across
several small-bodied opossum species, including *Monodelphis domestica*, that are predicted to
disrupt botrocetin binding and confer resistance (Drabeck et al. 2022). Protein binding assays
confirmed that VWF from both small and large-bodied opossums binds poorly to multiple VWF-
targeting CTLs, such as botrocetin, aspercetin, and bitiscetin, suggesting that mutations on VWF
disrupt binding of venom CTLs and potentially facilitate resistance (Drabeck et al. 2022). These
disruptive mutations on VWF have independently evolved multiple times across both large and
small-bodied opossums, indicating strong selective pressure on VWF regions associated with
venom CTL resistance (Drabeck et al. 2022). However, though previous studies demonstrated
that *Monodelphis domestica* platelets do not respond to botrocetin as human platelets do, and
that an *E.coli* expressed opossum VWF A1 domain fails to bind to venom CTLs, understanding
of venom resistance in this species remains extremely limited. Firstly*, in vitro* aggregation
assays used previously provide only a coarse measure of total venom-induced platelet response
in a cuvette (not in a living organism), leaving subsequent physiological responses, such as
platelet adhesion or fibrin formation, which contribute to the organism's overall resistance,
unexplored. These assays may also lead to an over-estimation of venom resistance, as it is
possible that though *M. domestica* VWF fails to bind to botrocetin, it may still suffer poor
platelet adhesion and delayed fibrin formation similar to venom susceptible species (essentially
voiding the impact of mutations on VWF on its ability to resist CTLs). The degree of impact of
venom CTLs on these downstream physiologies is required to understand the true impact of
venom CTLs on this species, and more generally the robustness of previous works' assertion
that *M. domestica* enjoys venom resistance comparable to its large-bodied cousins (Drabeck et
al. 2020, 2022).
Secondly, the behavior of blood and all its component parts can be greatly altered by
experiments that are conducted outside of a blood vessel. For example, shear force (the force
per unit area acting on the vessel wall caused by blood flow through the vessel) plays a crucial
role in the physiological function of VWF, as it promotes the unfolding of VWF multimers,
enabling platelet adhesion and clot formation at sites of vascular injury. For example, VWF-CTL
binding might be promoted under flow conditions, making *M. domestica* less resistant than
predicted by previous *in vitro* protein-protein interactions measured in absence of flow.
Evaluating the response of VWF to venom CTLs under vascularly relevant flow conditions is
therefore essential to understanding the extent to which venom resistant species might
maintain normal function in the presence of venom proteins. Lastly, resistance to individual
venom components, such as VWF-targeting CTLs, does not necessarily translate to whole-
venom resistance across all physiological systems. CTLs represent only one class within the

diverse suite of venom proteins (e.g. metalloproteinases and phospholipases), for which
resistance in this species is completely unexamined.
Understanding the mechanistic basis of an extreme adaptation like venom resistance
requires a systems-level approach, integrating multiple physiological processes to capture the
full scope of how these animals survive repeated venom exposures. The current study aimed to
characterize venom resistance in *M. domestica* across multiple physiological scales, and for
multiple types of venoms. We predicted that *M. domestica* would exhibit specific resistance to
Bothrops jararaca venom, with minimal disruption of platelet function and superior inhibition of
venom metalloproteinases relative to humans and other mammals. Our data confirm that *M.
domestica* has significant multifaceted venom resistance that is species-specific and aligns
with known ecological interactions. Our data unexpectedly reveal that *M. domestica* requires a
higher shear rate to initiate elongation compared to human VWF, an unpredicted but potentially
consequential aspect of opossum physiology which may also contribute to venom resistance.
This research provides insight into the complex mechanistic basis of venom resistance in small-
bodied opossums and provides vital physiological insights, which may guide novel therapeutic
strategies for VWF-related bleeding disorders.
144
145   **Methods**
146
Venom and Protein Purification
Purified fractions of botrocetin from *Bothrops jararaca* and aspercetin from *Bothrops
asper,* bitiscetin from *Bitis arietans*, as well as whole venom from *Bothrops jararaca* were used
from previous work (Drabeck et al. 2020, 2022)(botrocetin B). Convulxin was purchased from
ChemCruz (Santa Cruz Biotechnology Inc, Santa Cruz, CA). Whole venom from *Crotalus
oreganus* and *Sistrurus miliarius* was obtained during previous studies (Gibbs et al. 2011).
Venom CTLs were chosen to best provide a direct comparison of physiological and protein
assays done in previous work (Drabeck et al. 2020, 2022).
155
Opossum Handling and Blood Collection
Opossums (n= 16) were housed in the Oberlin research colony and cared for according
to procedures previously described (Rousmaniere et al. 2010). To collect blood for VWF
purification, opossums (*Monodelphis domestica*) were sacrificed by first being placed under
isoflurane sedation and then exsanguinated via the posterior vena cava, per IACUC protocol
F23BBYC-1 (Keyte and Smith 2008). Samples used for VWF purification were collected in an
ACD-coated syringe (acid citrate dextrose Solution A) and preserved in vacutainers containing
the same solution (BD Pharmaceuticals). Samples used in all other assays were collected in
3.2% sodium citrate coated syringe and similarly placed in vacutainers containing the same
solution (BD pharmaceuticals). Human blood was collected from de-identified donors via
venipuncture into 3.2% sodium citrate vacutainers (BD Pharmaceuticals) by the Case Western
Reserve University Hematopoietic Biorepository and Cellular Therapy Core under an Institutional
Review Board (IRB)-approved protocol (Case 12Z05, IRB # 09-90-195) of University Hospitals
Cleveland Medical Center. Blood was then centrifuged at 150 rcf for 15 min at 25°C to isolate a

platelet-rich plasma (PRP) supernatant, which was removed without disturbing the buffy coat.

The remaining sample was then centrifuged at 2000 rcf for 25 minutes to recover platelet-poor

plasma which was later used for dilutions and standards.

Venom Inhibition Assays

We evaluated the inhibition of whole venom by *Monodelphis domestica* serum against

three snake venoms (*Bothrops jararaca*, *Crotalus oreganus*, and *Bitis arietans*) using a modified

protocol from the EnzChek Gelatinase/Collagenase Assay Kit (Life Technologies, Carlsbad, CA,

USA). Briefly, this assay tests the ability of *Monodelphis domestica* serum to neutralize venom-

induced degradation of a gelatin substrate, which mimics the extracellular matrix proteins

targeted by snake venom metalloproteinases (SVMPs), and other enzymatic venom

components (Holding et al. 2016a). The DQ-gelatin substrate was diluted to 1:20 in reaction

buffer for the assay.

Serum was taken from opossums during venous blood extraction and frozen for

subsequent analyses, and the protein concentration of each serum sample determined with the

Pierce BCA Protein Assay Kit. Venom samples were diluted to a final protein concentration of

0.3125 ng per well. Venom was incubated with various amounts of *M. domestica* serum, with 0,

5, 10, 25, 50, 125, 250, or 475 µg serum protein per well, to determine the extent of venom

inhibition.

Fluorescence intensity was measured in relative fluorescence units (RFU) using

a SpectraMax M2 microplate reader (Molecular Devices) at 30 second intervals. The slope of

the fluorescence increase (RFU min^{-1}) from the linear portion of the reaction curve was

calculated to quantify the venom's SVMP (snake venom metalloproteinase) activity. Decreased

slope values, as compared to control reactions with no serum, indicate successful inhibition of

venom enzymatic activity by the serum. Each experiment was performed in triplicate, and data

were normalized to the control (untreated venom) to calculate relative SVMP activity.

Dose-dependence curves were fitted using the drc package in R, with a four-parameter

logistic model for each serum treatment (opossum, squirrel, and human) (Ritz et al. 2015).

Human plasma was used as a non-coevolved venom-sensitive baseline for venom SVMP

inhibition by serum proteins, and squirrel (*Otospermophilus beecheyi*) was used to compare

species-specific inhibition as it has previously been shown to have resistance for its sympatric

predator (*Crotalus oreganus*) (Holding et al. 2016a). Venoms were selected to represent 1) a

sympatric species (*Bothrops jararaca*) for which ecological and other data predict resistance, 2)

an allopatric species (*Crotalus oreganus*) for which another mammal (ground squirrels) have

evolved venom resistance 3) an allopatric distantly related viper (*Bitis arietans*) (Figure 2 top

and middle panels).

Aggregation Assays

To test *M. domestica*'s ability to resist whole venom as well as isolated CTLs, whole

blood aggregation was assessed using an impedance aggregometer (Roche multiplate

aggregometer). Briefly, these assays differ from microfluidic assays as coagulopathic response

is indicated by *induction* of platelet aggregation, rather than inhibition. In the absence of

exposed collagen and/or shear force, platelets in blood (in a stationary cuvette) should not
rapidly aggregate unless exposed to a platelet activating agonist (i.e. whole venom or pure
venom-derived CTLs). Failure of agonists to induce platelet aggregation suggests strong venom
resistance to coagulopathic venom components, including CTLs. Thus, we expect *M.*
domestica's platelet aggregation response to be lower for sympatric venom and CTLs than for
allopatric venom and CTLs, suggesting higher resistance to species they have evolved with.
Platelet aggregation response in this assay is measured by impedance, where higher
impedance means a greater response. Purified CTLs used were aspercetin (20 µL, 0.1 mg/mL),
convulxin (20 µg, 0.01 mg/mL), and botrocetin (20 µg, 0.01 mg/mL). Platelet aggregation was
measured over 10 minutes, with impedance changes recorded as a measure of platelet
aggregation. All purified CTLs were selected from South American vipers that were either
completely sympatric (*Bothrops jararaca*), partially sympatric (*Crotalus durissus*), or allopatric
(*Bothrops asper*) with *M. domestica*. It should be noted that while aspercetin and botrocetin
induce aggregation via VWF, convulxin induces aggregation via a direct interaction with the
platelet. In addition to purified CTLs, we also challenged *M. domestica* with whole venom from
the same sympatric species (*Bothrops jararaca*), as well as two distantly related allopatric
species with vastly diverse venoms (*Crotalus adamanteus* and *Sistrurus miliarius*) to provide
some measure of the specificity/cross-reactivity of venom resistance in *M. domestica* in the
absence of availability of whole venom from *Bothrops asper* and *Crotalus durissus* (see Figure 3
range maps).
Venous blood samples collected from *M. domestica* in 3.2% sodium citrate tubes were
tested within four hours of collection at room temperature. For each assay, 300 µl citrated
whole blood and 300 µl NaCl/CaCl2 solution (0.9% saline and 3mM CaCl2 provided my
manufacturer) were aliquoted into cuvettes and stirred at 37°C. Aggregation was induced by
adding 20 µL of agonists to the samples. Collagen (20 µg/mL) was used as a positive control,
PBS (phosphate buffered saline) was used as a negative control, Aggregation response was
calculated as the average of two replicate measures. To facilitate cross-comparison, the data
were normalized by subtracting the minimum value within each treatment group, ensuring that
the minimum impedance value was zero. Data visualization was performed using R software to
allow for comparison of platelet responses to different agonists. Venom assay and controls
were conducted in triplicate to ensure reproducibility; however, due to limited plasma volume,
CTL tests were not replicated.

VWF Multimer Gel
Opossum VWF was purified by size-exclusion chromatography at 4°C as previously
described (De Marco and Shapiro 1981), using plasma pooled after high-speed centrifugation.
Fractions were quantitated by absorbance at 280 nm using the in-line detector on the AKTA
Pure. To ascertain that VWF from *M. domestica* bound to microchannel equine-derived type 1
fibrillar collagen (Chrono-log Corporation), purified and quantified opossum VWF was tagged
with Alexa488 using the Alexa Fluor 488 (AF488) Microscale Protein Labelling Kit (Invitrogen)
with minor modification for subsequent microchannel runs. The remaining untagged VWF was
frozen and stored at -80C.

To compare opossums' VWF multimer structure to human, we performed a multimer
analysis using VWF purified from *M. domestica* plasma and HumateP (a purified human
plasma-derived product that contains VWF and Factor VIII, CSL Behring). Due to the enormous
size of intact VWF multimers, agarose is used instead of polyacrylamide. VWF multimers were
analyzed by electrophoresis using a 1.6% separating LDS agarose gel and visualized using the
rabbit anti-human VWF antibody A0082 (DAKO) as previously described (Pruss et al. 2011).
Protein was transferred to nitrocellulose using the semi-dry Power Blotter (ThermoFisher) and
western blot was performed using the iBind Flex (Invitrogen) system with the rabbit anti-human
VWF antibody A0082 (DAKO) as primary and IRDye 800CW anti-rabbit antibody (Li-Cor). Lanes
were visualized a Li-Cor Odyssey CLx Imager and Image Studio (LiCor).
264
Microfluidic Assays
To evaluate whether the apparent resistance of *M. domestica* to CTL-induced platelet
aggregation (Figure 3) could be protective against bleeding, we performed microfluidics assays
developed to approximate *in vivo* injury conditions which allowed for real-time measurement of
VWF/VWFplatelet recruitment and fibrin generation.
Microfluidics experiments were performed in the Bioflux 200 system (Cell
Microsystems) using 48-well High Shear Plates (Cell Microsystems) (Conant et al. 2009; Girish
et al. 2022). Glass-bottomed microfluidic channels with a width of 250 µm and a height of 70
µm were coated with 100 µg/mL type 1 equine fibrillar collagen (Chrono-log Corporation) in 20
µM acetic acid for 60 min at 37°C. These conditions facilitate the self-polymerization of
collagen fibers which anchor to the glass substrate, creating an environment which mimics
collagen exposure following vascular injury. Unbound collagen was washed by perfusing 0.1
wt% bovine serum albumin (BSA) in calcium-free phosphate buffered saline (PBS) at 30
dyn/cm^2 for 2 min.
While we have previously standardized this assay for use with human blood, it was
unknown whether the equine collagen would allow binding of opossum VWF in a similar manner
and under similar shear flow conditions. To verify that this assay is applicable to opossum VWF,
we used purified and fluorescently tagged VWF to determine experimental conditions and verify
that opossum VWF binds effectively to equine fibrillar collagen used in standard microfluidics
assays (Figure 1). AF488-labeled human or opossum VWF in calcium-free PBS was perfused
through the collagen-coated microfluidic channel at shear stresses ranging from 20 to 200
dyn/cm^2. VWF binding to collagen was imaged under epifluorescence using a Zeiss Axio
Observer microscope with a 10x objective (Plan-NEOFLUAR, NA = 0.3). We observed similar
overall VWF deposition, indicating that opossum VWF binds to equine collagen to a similar
extent as human VWF. We further observed that opossum VWF required comparatively higher
shear flow conditions (80 dyn/cm^2 for opossum VWF vs. 60 dyn/cm^2 for human VWF) to fully
elongate, which we discuss in detail in the results and elsewhere (Figure 1).
With assay conditions established, we assessed the ability of *M. domestica* to resist the
detrimental effects of venom CTLs. Opossum or human (as a venom non-resistant control
species) blood was processed and used to perform assays within four hours of any given draw,
to ensure that platelet activity in the blood remained viable. Citrated PRP was obtained as

described in the *Opossum Handling and Blood Collection* section. PRP was incubated with 1
µg/mL calcein-AM (Invitrogen) to stain platelets and 10 µg/mL AF-647 fibrinogen (Invitrogen)
for 30 min at room temperature. Samples were recalcified to 20 mM $CaCl_2$ and immediately
perfused through collagen-coated channels at 60 dyn/cm^2, a shear condition that is relevant to
wound environments. These conditions provide more fine-tuned information about the degree
and type of physiological response induced by exposed collagen in the presence of agonists.
Epifluorescence microscopy was used to image platelet accumulation and fibrin generation in
real time. Samples were run without any additives (Control) to assess normal platelet adhesion
and fibrin generation. For each sample, complementary assays were performed in the presence
of three venom derived CTLs known to affect platelet aggregation function: botrocetin,
aspercetin, and convulxin. Each CTL was used at its ED_{90} concentration for human blood: 4
µg/mL for botrocetin, 20 µg/mL for aspercetin, and 0.5 µg/mL for convulxin. In this assay, we
expect that CTL-induced VWF/platelet complexation leading to thrombocytopenia and low VWF
levels should result in reduced platelet adhesion and prolonged fibrin generation times,
mimicking bleeding observed *in vivo*. To account for differences in platelet counts between
individual animals, platelet adhesion was normalized to the control condition for each sample.
317   **Results**
318
319   *Venom Inhibition*
320   While hemostatic disruptions are an important mechanism of venom pathology,
enzymatic tissue damage caused by venom proteinases often make up the majority of viper
venoms and are thought to be a major source of venom pathology. To address the ability of M.
domestica's serum to inhibit this functional component of venom, we employed gelatin-based
inhibitions assays. Gelatinase inhibition assays test the ability of blood serum from different
species to inhibit the ability of venom to break down a substrate that mimics tissue and are
typically used as a proxy quantitation of venom metalloproteinase (SVMP) inhibition. These
data are used to calculate ED_{90} values, which represent the dosage of serum needed to inhibit
90% of the venom's gelatinase activity. Blood serum from *M. domestica* demonstrated varying
levels of inhibition across the three snake venoms tested: *Bothrops jararaca*, *Crotalus oreganus*,
and *Bitis arietans* (Figure 2A-C; Table 1). In all cases, increasing concentrations of serum protein
led to a dose-dependent decrease in venom proteinase activity. For *Bothrops jararaca* venom
(Figure 2A), *M. domestica* serum rapidly inhibited SVMP activity at the middle to high
concentrations of serum protein tested, resulting in the lowest ED_{90} value of the three sera
tested, indicating it inhibited venom approximately 2.5 times better than squirrel serum and 14
times better than human serum (Table 1). Similarly, serum of the squirrel *O. beecheyi* yielded a
lower ED_{90} for its sympatric predator, *C. oreganus*, than did the human or opossum sera (Table

1). Finally, opossum serum performed best against the venom of the African viper *B. arietans,*
while human and squirrel serum showed overlapping trends in inhibition.
339
Whole Blood Aggregation
Whole blood aggregation assays revealed differential platelet responses by *M.*
domestica to the various venoms and purified CTLs tested (Figure 3). As shown in Figure 3A,
collagen, used as a positive control, induced a rapid and sustained increase in impedance,
indicative of expected collagen-induced platelet aggregation, while PBS (negative control)
established a baseline throughout the 10-minute measurement period.
Among whole venoms, *Crotalus adamanteus* venom elicited the most pronounced
platelet aggregation response, with impedance peaking around 5 minutes at approximately
220% of the collagen control (400 Ω) before showing pronounced disaggregation, indicating a
robust but transient aggregation effect (Figure 3). *Sistrurus miliarius* venom induced a response
reaching approximately 70% of the collagen control (180 Ω), while *Bothrops jararaca* venom
resulted in a minimal increase in impedance, remaining just slightly below the PBS baseline (50
Ω), suggesting substantial resistance to platelet activation (Figure 3). As expected, collagen
(positive control) produced a peak impedance of 180 Ω, while PBS (baseline control) showed
negligible effects with impedance remaining below 50 Ω, which is consistent with standard
values for other mammals (Baumgarten et al. 2010; Nash et al. 2017).
The purified CTLs demonstrated variable effects on platelet aggregation. Convulxin, a
known collagen-like agonist, induced a notable increase in impedance, mirroring the
aggregation pattern of the collagen control but with a delayed onset (Figure 3). Consistent with
whole venom results which showed no resistance to *Crotalus* venom, this indicates a lack of
resistance to Convulxin (derived from *Crotalus durissus* venom). Aspercetin, which targets VWF,
triggered a measurable but extremely reduced platelet aggregation response. Botrocetin, like
whole *B. jararaca* venom, elicited no detectable platelet aggregation response. The responses of
both aspercetin and botrocetin observed here are extremely minimal compared to previously
well-established responses for susceptible species, suggesting strong resistance (Rucavado et
al. 2001: Figure 4; Drabeck et al. 2020: Figure 3). Overall, *M. domestica* showed the lowest
levels of whole blood aggregation by both snake venom and purified CTLs from their sympatric
snake predators in the genus *Bothrops* and appeared to be entirely resistant to both platelet
disturbance by *B. jararaca* venom and botrocetin. This is consistent with the hypothesis of
evolution of increased resistance to venom of sympatric snakes in *M. domestica*.
370
VWF Multimers
In humans, higher molecular weight VWF multimers are more procoagulant than lower
weight forms as they are more likely to bind platelets after interacting with subendothelial
collagen under flow. When compared to a commercial preparation of human VWF, no obvious
differences in multimer distribution between *Monodelphis domestica* and human VWF were
observed. Both displayed a typical distribution of high, intermediate, and low molecular weight
multimers (Supplementary Figure 1) although it is easier to visualize the higher molecular
weight multimers in *M. domestica* due to the higher concentration of this sample. This suggests

that the multimerization process is preserved in *M. domestica* and indicates that this VWF is

structurally comparable to human VWF in terms of multimer formation, despite sequence

divergence associated with phylogenetic distance or selective changes in VWF function.

Platelet Adhesion and Fibrin Generation in Microfluidics

Blood clot formation relies on rapid platelet adhesion at sites of vascular injury,

followed by aggregation and fibrin clot formation to stabilize the plug. While aggregation assays

measure platelet clumping under static conditions, they do not replicate *in vivo* conditions

because aggregometry assays do not have endothelium and vasculature or shear forces (i.e.

vascular-relevant flow). Microfluidic assays better recapitulate these physiological conditions

by simulating blood flow along vessel walls using pressurized synthetic vessels with

immobilized collagen attached to the 'vessel' wall, providing insights into how venom

components modulate platelet function and disrupt hemostasis. Consequently, microfluidic

assays can recapitulate the *in vivo* effects of venom components, that is, *inhibition* of platelet

plug formation (platelet adhesion and aggregation) at sites of vascular injury and the

subsequent development of fibrin clots.

As expected, control experiments with human PRP showed that platelet accumulation

on the microfluidic surface was strongly inhibited by botrocetin and convulxin, and weakly

inhibited by aspercetin (Figure 4A-D).Venom derived CTLs also significantly reduced platelet

fluorescence in *Monodelphis domestica* PRP (Figure 4 E-H); however, the defect was less

pronounced after treatment with botrocetin or convulxin than in human PRP (Figure 4 B, F, D, H),

suggesting some measure of resistance to those CTLs. In human PRP, the initial rate of platelet

adhesion was similar for the control, botrocetin and aspercetin conditions, but reduced to

almost zero in the presence of convulxin (as indicated by the initial slope of the platelet

fluorescence curves, Figure 4I). A similar trend was observed in opossum PRP, except that

platelets adhered even in the presence of convulxin, though at much lower levels than the other

conditions (Figure 4J). Concordant with the whole blood aggregometry analyses, the effect of

botrocetin on platelet adherence to the microchannel was decreased in *M. domestica* (Figure 4

B, F, J): botrocetin caused a more severe defect in the total extent of platelet adhesion than

aspercetin in human PRP, but a less severe defect than aspercetin in opossum PRP (Figure 4I-

K). Aspercetin and convulxin caused a similar response in both species (Figure 4K). Concordant

with the whole blood aggregometry analyses, the effect of botrocetin on platelet adherence to

the microchannel was significantly decreased in *M. domestica* when compared to human

(Figure 4 B & F, K). In human PRP, botrocetin caused a 66 ± 16% reduction in the overall extent

of platelet adhesion compared to the negative control, whereas it only caused a 39 ± 15%

reduction in opossum PRP (Figure 4K).

We further quantified the fibrin generation kinetics (by quantifying 'lag time' reflecting

the time needed for fibrin clot formation), where a prolonged lag time is unfavorable for

effective hemostasis and predictive of bleeding. Purified venom CTLs increased lag time in both

humans and opossums (Figure 4 L-N). While convulxin had a dramatic effect in human samples,

abrogating fibrin formation almost completely, it showed a more modest response in opossums

(Figure 4 L-N). As with platelet adhesion, the greatest signal for resistance was seen in

botrocetin treatments (Figure 4 L-N), with an overall reduced effect of botrocetin in opossum
PRP (Figure 4 L-N). Botrocetin caused a nearly threefold increase in fibrin lag time from 3.6 ±
1.0 min to 9.0 ± 1.2 min in human PRP, but a statistically insignificant prolongation of fibrin lag
time in opossum PRP (from 3.1 ± 0.8 to 4.5 ± 1.8 min, Figure 4N).
Surprisingly, aspercetin caused a statistically insignificant prolongation in the fibrin lag
time for human PRP but doubled lag time from 3.1 ± 0.8 min to 7.4 ± 2.4 min in opossum PRP,
which is perhaps suggestive of opossum susceptibility to aspercetin. This result is consistent
with previously established weak affinity of aspercetin to VWF A1 in protein binding assays and,
together with weak platelet adhesion inhibition observed here, suggests that VWF-mediated
platelet aggregation is not the primary function of coagulopathy for aspercetin (Drabeck et al.
2022).
Convulxin induced a reduced coagulopathy in opossum PRP compared to human PRP
but remained the most potent effector of integrated hemostatic function in both species.
Platelet fluorescence and adhesion in response to aspercetin was roughly similar between
human and opossum, though aspercetin had a slightly stronger effect on fibrin lag time in
opossums. Compared to humans, *M. domestica* VWF showed a measurably reduced integrated
coagulopathic response (platelet fluorescence, platelet adhesion, and fibrin lag time) to
botrocetin, suggesting that these opossums have functionally meaningful hemostatic protective
effects against this CTL.
440
441

Discussion

Understanding the physiological mechanisms, molecular underpinnings, and
evolutionary history of venom resistance across opossums is crucial for understanding how
selective pressures, ecological interactions, and physiological constraints shape the evolution
of such extreme adaptations (Byrne et al. 2014; Xu et al. 2020). Our results provide compelling
evidence that *M. domestica* possesses considerable resistance to both CTLs and whole venom
from *Bothrops jararaca*, and suggests that previous *in vitro* work showing convergently derived
mutations on VWF is indicative of venom resistance (Drabeck et al. 2022). We also uncover a
previously undescribed modification to opossum VWF, that it requires a greater shear force to
elongate, potentially contributing to the resistance phenotype or other important adaptive
physiologies.
Under normal conditions, VWF will be induced to elongate and bind platelets in response
to either exposed collagen (mimicking vessel damage) or under increased shear force (caused
by increased blood flow after vessel damage), creating a primary platelet plug preceding the
formation of a more stable cross-linked fibrin clot (Sadler 1998; Ruggeri 2003). Botrocetin is
known to induce coagulopathy upon introduction to the body by binding to VWF (at the A1
domain) and inducing VWF to elongate and attach to platelets (Matsushita et al. 2000; Fukuda
et al. 2002). Botrocetin then changes conformation slightly and binds secondarily to the platelet
GP1Bα binding site, creating a tight trimolecular complex which renders platelets unavailable for
adhesion and causes extreme thrombocytopenia (Fukuda et al. 2002, 2005). More recently,
detailed physiological work on botrocetin has shown that in the presence of VWF, botrocetin

can induce coagulopathy even in the absence of its preferred platelet receptor (GP1Bα), and
that it also interacts directly with a different platelet integrin (αIIbβ3) to block fibrinogen and
other ligands essential for coagulation and healing (Shen et al. 2024). While aspercetin is also a
VWF-mediated agonist from a closely related viper (*Bothrops asper*) and is thought to work in a
similar manner because of its ability to induce VWF-mediated platelet agglutination, detailed
structural studies have yet to locate its exact binding site on VWF and prior kinetics studies
show weak binding to the A1 domain (Rucavado et al. 2001; Drabeck et al. 2022). Integrated
coagulopathic data shown in this work supports that VWF-mediated platelet activity is a weak
action for aspercetin and may not be the main mode of coagulopathy for this CTL. Convulxin is
derived from the more distantly related tropical rattlesnake *Crotalus durissus*, and binds directly
to the GPVI platelet receptor, inducing coagulopathy independent of VWF (Batuwangala et al.
2004).
Our assays of whole blood aggregation revealed a reduced response of *M. domestica*
blood to both botrocetin and aspercetin but not convulxin. Initially, this might lead us to assume
that opossum resistance is specific to CTLs from *Bothrops spp.* and centered at VWF. However,
microfluidic assays showed that platelet adhesion and fibrin clot formation in opossums was
more disrupted by aspercetin than botrocetin. This suggests that the molecular changes that
have previously been found to abolish binding between opossum VWF and botrocetin result in
resistance to this CTL (implying specificity), but *does not* result in resistance to aspercetin,
despite the appearance of strong resistance that would be gleaned from aggregation data
alone. Aspercetin appears to be able to function nearly as well in opossums as it does in
humans in microfluidics assays (in both cases it has weak function), but not in whole blood
assays, highlighting the importance of integrated assays which capture a more complete
picture of coagulopathic function. These results suggest that aspercetin may act directly on
platelets, fibrin, or other agonists more strongly during platelet adhesion and fibrin clot
formation, or that its function in opossum depends on the presence of shear force. Given that
its action in humans is also weak relative to other CTLs, and previous work has revealed a
surprisingly low affinity for human VWF compared to other VWF-mediated agonists, it is
possible that this is an accessory function of this protein, or that its affinity for VWF is specific
to a non-mammalian VWF target.
Similarly, we observed a tremendously decreased coagulopathic response in opossums
(compared to previously established human responses) to whole venoms from both species of
Bothrops, but not *Crotalus*. Opossum serum showed superior ability to inhibit the enzymatic
action of whole venom for all species tested, except *Crotalus oreganus*, which was best
inhibited by its sympatric prey (*Otospermophilus beecheyi*). Interestingly, *M. domestica* serum
very strongly inhibited venom from the African viper, *Bitis arietans*, which points to the likelihood
that this species also possesses generalized inhibitory proteins (snake venom
metalloproteinase inhibitors, phospholipase inhibitors), similar to those seen in other highly
resistant taxa (Voss and Jansa 2012).
Bothrops spp. are well known to interact with opossums and are a known prey item of
several species of South American opossums (Voss and Jansa 2012; Voss 2013). *Bothrops*
jararaca co-occurs across the range of *M. domestica*, are relatively small and gracile, semi-

arboreal, and have been anecdotally observed to bite *M. domestica* without effect (Martins et al.
2001; Da Silva et al. 2024; exclusivo cuíca predando cascavel e coral verdadeira e é picado por
elas n.d.). *Bothrops asper* are somewhat closely related to *B. jararaca;* however they are much
larger and their range does not overlap with *M. domestica*. Previous work examining diet in
small-bodied opossums has suggested that predator-prey relationships between opossums and
snakes may role-switch relative to factors affecting venom potency like geographic region, body
size, and age (Voss 2013). Conversely, *Crotalus durissus* is documented to be prey of only the
larger bodied opossums (Didelphini) and though its range overlaps with *M. domestica*, our data
did not reveal compelling evidence that *M. domestica* has profound resistance to convulxin (the
CTL derived from its venom) (Almeida-Santos et al. 2000; Voss and Jansa 2012). This suggests
that small-bodied opossum venom resistance may be more specialized for species that they
target for prey, are appropriately sized for prey, or that are more abundant in their environment.
Interestingly, our findings reveal that VWF in *M. domestica* requires a comparatively
higher shear force than that for human VWF, to elongate. This property could have important
implications for venom resistance by making it more difficult for venom components, like
botrocetin, to bind VWF and exact subsequent coagulopathic effects. Elevated shear thresholds
for VWF elongation may also protect against other physiological disruptions, such as
spontaneous platelet adhesion in high-flow conditions or inappropriate clot formation. Beyond
venom resistance, this trait may be linked to broader physiologies of non-placental mammals,
potentially reflecting differences in vascular dynamics, metabolism, or reproductive strategies.
The discovery of this and potentially other species with shear-resistant VWF may also provide a
novel avenue for therapeutic exploration, where engineering similar shear-resistant VWF
variants could help prevent bleeding disorders associated with left ventricular assist devices
(LVAD) (Nascimbene et al. 2016). While there is some limited data to suggest that increased
shear elongation may occur in other venom resistant species (pigs), little is known about the
diversity of this trait across species, or what potential costs and tradeoffs it might evoke (Chan
et al. 2017). Further work is merited to better understand this unique physiology and explore its
potential for inspiring new strategies for mitigating LVAD-related complications and other
vascular disorders.
Our work adds detailed multi-layered comparative assays using physiologic vessel
models to already mounting evidence that venom resistance is present in small-bodied
marsupials, and demonstrates the ability of comparative phylogenetic work to inform otherwise
cryptic adaptations and ecologies (Jansa and Voss 2011; Drabeck et al. 2020, 2022). In light of
these findings, it is likely that many more of the opossum species identified to have similar and
convergently evolved changes in VWF may have similar degrees of venom resistance (Drabeck
et al. 2022). Further work is warranted to examine the true ecological roles and behaviors of this
entire group of mammals and emphasizes a need for basic natural history work. This work also
provides vital insight into the evolution of venom in this region and suggests that South
American viper venoms may have evolved in the presence of many more species of co-evolving
venom-resistant mammalian predators and prey than previously assumed. Detailed
physiological data presented here serves to further inform the physiological mechanisms,

relative strength, and specificity of this extreme adaptation, and provides key insights for future
work aiming to understand the molecular basis and evolution of this remarkable trait.
548
549   **Acknowledgements**
550   The authors would like to thank Alexandra Rucavado, Erika Hingst-Zaher, and the Kentucky
Reptile Zoo for their help in acquisition and purification of venoms used for this work. Authors on this
work were supported by the Minnesota IRACDA Program 2K12GM119955 (NIGMS, NIH), the NSF STEM-
APWD (DEB 2316784), NIH R01-HL121212, NIH R35-HL171421, and NIH R35-HL150784 (NHLBI). The
authors would like to thank Suzanne McGaugh for tremendous guidance as well as lab and resource
support. The authors would also like to thank Marvin Neiman (Case Western), Alison Narayan/Sarah
Ackenhusen (University of Michigan), and Randy Westrick (Oakland University) for their guidance,
feedback, and support.
558
559
560   **Tables**
561
562   **Table 1**. Species-specific ED_{90} values (µg/mL) derived from dose-response analysis of SVMP
563   inhibition by increasing amounts of serum protein from three mammal species. ED_{90} (Effective
564   Dose, 90%) indicates the amount of serum required to inhibit 90% of the venom's proteinase
565   activity. Asterisks indicate species that are sympatric with the snake venom tested.
566

Source of Venom	Serum	ED_{90}	S.E.
Bothrops jararaca	Human	12379.2	6272.4
	Opossum*	865.4	149.1
	Squirrel	2182.2	636.7
Crotalus oreganus	Human	11391.6	7660.1
	Opossum	1236.6	231.7
	Squirrel*	541.9	71.4
Bitis arietans	Human	774.4	80.1
	Opossum	220.5	21.7
	Squirrel	940.5	107.7

572   **Figure Legends**
573
Figure 1. (A) Elongation of von Willebrand Factor (VWF) under different shear conditions. (B)The degree
of elongation for both human and *Monodelphis domestica* (opossum) VWF under varying shear forces in

a microfluidics assay. Purified VWF was fluorescently tagged and perfused across collagen-coated
channels at shear rates of 0 dyn/cm² to 200 dyn/cm² for human and opossum samples, respectively. The
percentage elongation of VWF was quantified and normalized to both the 0 dyn/cm^2 condition (0%
elongation) and the 200 dyn/cm^2 condition (100% elongation). At 60 dyn/cm², human VWF showed full
elongation, whereas opossum VWF remained partially coiled, indicating that higher shear rates are
required to induce full elongation in the latter. When perfused at 80 dyn/cm², opossum VWF fully
elongated, mirroring the response observed in human VWF at 60 dyn/cm^2. Error bars represent the
average ± SD of n = 10 replicates. High degrees of fluorescent labeling caused aggregation or "clumping"
of VWF multimers in both human and possum samples, inhibiting complete elongation. Reduced labeling
allowed full elongation for human VWF at 60 dyn/cm², but opossum VWF still required the higher shear
rate (80 dyn/cm²) to achieve comparable uncoiling.

Figure 2. Range maps for mammals *M. domestica* and *O. Beechyi* (top), and snakes (B. jararaca, *C.*
oreganus, and *B. arietans* (middle). Bottom panel shows results of the venom inhibition assays. *Bothrops*
jararaca, *Crotalus oreganus*, and *Bitis arietans* venoms were incubated with increasing concentrations of
mammalian serum (*M. domestica*, *O. Beechyi*, and *H. sapiens*), and relative degradation of a gelatin
substrate was measured. Data points represent the mean ± SEM for three independent experiments. *M.*
domestica serum demonstrated greater inhibition of proteinase activity compared to squirrel and human
sera, at higher serum concentrations, particularly for venoms from *B. jararaca* and *B. arietans*. Inhibition
was assessed based on the reduction in fluorescence due to the degradation of a gelatin substrate,
reflecting the venom's proteolytic activity. In this graph, the x axis would represent total resistance
(depicted with the alive mouse icon), and decrease as you move away from y = 0 (depicted by the dead
mouse icon).

Figure 3. Platelet aggregation *in Monodelphis domestica* whole blood in response to venoms and purified
C-type lectins. Impedance was measured over a 10-minute period to assess platelet aggregation
response. (A) Collagen (positive control), and PBS (negative control). (B) Whole venoms (*Crotalus*
adamanteus, *Sistrurus miliarius*, and *Bothrops jararaca*) are in the middle panels. (C) C-type lectins,
Convulxin (derived from *Crotalus durissus* venom), aspercetin (derived from *Botrhops asper* venom) and
botrocetin (derived from *Bothrops jararaca* venom) are on the bottom panels. Data represents the average
impedance change over time, normalized to account for baseline variation. Multiple curves represent
independent samples (replicates) performed when available. Cartoon on the left depicts platelet
aggregation (lowest at y =0 and increases at higher values of y). Range maps of species from which each
venom/CTL are derived are shown in the upper right of each graph to allow for a comparison of
aggregation response with sympatry.

Figure 4. Compared to humans, opossums exhibited resistance to botrocetin (bot), but not aspercetin
(asp) or convulxin (cvx). A-H) Endpoint images of platelet (fluorescently labeled in green) adhesion to a
collagen coated substrate after 10 min of shear flow at 60 dyn/cm^2. A) Human platelet rich plasma (h-
PFP) perfused across a collagen-coated substrate resulted in robust platelet adhesion in the absence of
venom derivatives (ctrl). B) Platelet adhesion was significantly reduced in the presence of bot, C)
moderately reduced in the presence of asp, D) and almost completely abrogated in the presence of cvx.
E) Compared to h-PRP, opossum PRP (o-PRP) exhibited a similar extent of platelet adhesion to collagen.
F) However, platelet adhesion was affected to a lesser extent by bot. H) o-PRP responded similar to h-
PRP to asp, but H) platelet adhesion in o-PRP was not entirely abolished by cvx. I-J) Representative
quantification of platelet fluorescence over time. I) In h-PRP, the overall extent of platelet adhesion was

reduced slightly by asp, significantly by bot, and almost completely by cvx. J) Though platelet adhesion in
o-PRP was affected by asp to a similar degree as h-PRP, platelet adhesion was only barely reduced by
bot. Some degree of platelet adhesion was preserved in the presence of cvx. K) Aggregate platelet
adhesion data, normalized to the ctrl condition for each individual donor or opossum. The defect in
platelet adhesion was similar for h-PRP and o-PRP for asp and cvx, but reduced significantly in o-PRP for
bot. L-M) Representative quantification of fibrin(ogen) fluorescence over time. The sharp ramp indicates
fibrin generation, and the corresponding time point is parameterized as fibrin lag time. L) In h-PRP, fibrin
polymerized quickly in the absence of venom derivatives, owing to the procoagulant surface of collagen-
bound platelets. Asp and bot both delayed the formation of fibrin, with bot having a more pronounced
effect. Fibrin did not polymerize during the 10 min experiment in the presence of cvx. M) In o-PRP, bot
caused only a marginal prolongation in fibrin lag time. Both asp and cvx significantly delayed fibrin
generation. N) Aggregate fibrin generation data. The dashed line at t = 10 min indicates the endpoint of
the experiment. On average, fibrin generated in less than 4 min in h-PRP. Bot more than doubled fibrin lag
time, with half of the replicates not generating fibrin within the 10 min experimental window. Despite its
similarity to bot, asp did not significantly affect fibrin lag time. In three of four replicates, fibrin did not
generate within 10 min in the presence of cvx. In o-PRP, fibrin lag time was prolonged by asp and cvx, but
not bot. o-PRP generated fibrin faster than h-PRP in the presence of bot or cvx, but slower than h-PRP in
the presence of asp. Statistics were performed using one-way ANOVA using Fisher's LSD correction. Error
bars mean ± SD. For significance, *: $p \leq 0.05$, **: $p \leq 0.01$, *** $p \leq 0.001$, ****$p \leq 0.0001$.

641
642
643
Supplementary Figure 1. Opossum VWF Multimer Gel 1.6% resolving + 0.8% stacking Tris Glycine LDS-
SeaKem HGT Agarose Vertical Multimer Gel (8x8cm vertical). Multimeric structures of von Willebrand
factor (VWF) were analyzed using a 1.6% lithium dodecyl sulfate (LDS) agarose gel under non-reducing
conditions. Lane 2 contained 30 µL of purified Monodelphis domestica VWF (heated at 60°C for 30
minutes) and 10µl loading buffer. Lane 5 contained 30 µL HumateP (human VWF) as a control (heated at
60°C for 30 minutes) and 10µl loading buffer. The gel was transferred to a nitrocellulose membrane and
visualized by fluorescence with anti-human VWF antibody (Dako A0082) using a Li-Cor Odyssey CLx
Imaging System (Li-Cor).

652
653

References

Almeida-Santos, S. M., M. M. Antoniazzi, O. A. Sant'Anna, and C. Jared. 2000. Predation by the Opossunn

Didelphis marsupialis on the Rattlesnake Crotalus durissus. Curr.Herpetol. 19:1–9.

Batuwangala, T., M. Leduc, J. M. Gibbins, C. Bon, and E. Y. Jones. 2004. Structure of the snake-venom

toxin convulxin. Acta Crystallogr D Biol Crystallogr 60:46–53.

Baumgarten, A., M. Wilhelmi, K. Kalbantner, M. Ganter, and R. Mischke. 2010. Measurement of platelet

aggregation in ovine blood using a new impedance aggregometer. Veterinary Clinical Pathology

39:149–156.

Bury, L., A. Malara, S. Momi, E. Petito, A. Balduini, and P. Gresele. 2019. Mechanisms of thrombocytopenia

in platelet-type von Willebrand disease. Haematologica 104:1473–1481.

Byrne, R. T., A. J. Klingele, E. L. Cabot, W. S. Schackwitz, J. A. Martin, J. Martin, Z. Wang, E. A. Wood, C.

Pennacchio, L. A. Pennacchio, N. T. Perna, J. R. Battista, and M. M. Cox. 2014. Evolution of

extreme resistance to ionizing radiation via genetic adaptation of DNA repair. eLife 3:e01322.

eLife Sciences Publications, Ltd.

Chan, C. H. H., I. L. Pieper, C. R. Robinson, Y. Friedmann, V. Kanamarlapudi, and C. A. Thornton. 2017.

Shear Stress-Induced Total Blood Trauma in Multiple Species. Artificial Organs 41:934–947.

Conant, C. G., M. A. Schwartz, T. Nevill, and C. Ionescu-Zanetti. 2009. Platelet adhesion and aggregation

under flow using microfluidic flow cells. J Vis Exp 1644.

Da Silva, F. F. B., T. D. A. Moura, T. Siqueira-Silva, J. M. Gutiérrez, and P. A. Martinez. 2024. Predicting the

drivers of Bothrops snakebite incidence across Brazil: A Spatial Analysis. Toxicon 250:108107.

Drabeck, D. H., A. Rucavado, E. Hingst-Zaher, Y. P. Cruz, A. M. Dean, and S. A. Jansa. 2020. Resistance of

South American opossums to vWF-binding venom C-type lectins. Toxicon 178:92–99.

Drabeck, D. H., A. Rucavado, E. Hingst-Zaher, A. Dean, and S. A. Jansa. 2022. Ancestrally Reconstructed

von Willebrand Factor Reveals Evidence for Trench Warfare Coevolution between Opossums and

Pit Vipers. Molecular Biology and Evolution 39:msac140.

Engelman, R. K., F. Anaya, and D. A. Croft. 2017. New palaeothentid marsupials (Paucituberculata) from

the middle Miocene of Quebrada Honda, Bolivia, and their implications for the palaeoecology,

decline and extinction of the Palaeothentoidea. Journal of Systematic Palaeontology 15:787–

820.

Fukuda, K., T. A. Doggett, L. A. Bankston, M. A. Cruz, T. G. Diacovo, and R. C. Liddington. 2002. Structural

Basis of von Willebrand Factor Activation by the Snake Toxin Botrocetin. Structure 10:943–950.

Fukuda, K., T. Doggett, I. J. Laurenzi, R. C. Liddington, and T. G. Diacovo. 2005. The snake venom protein

botrocetin acts as a biological brace to promote dysfunctional platelet aggregation. Nat Struct

Mol Biol 12:152–159.

Gibbs, H. L., L. Sanz, J. E. Chiucchi, T. M. Farrell, and J. J. Calvete. 2011. Proteomic analysis of

ontogenetic and diet-related changes in venom composition of juvenile and adult Dusky Pigmy

rattlesnakes (*Sistrurus miliarius barbouri*). Journal of Proteomics 74:2169–2179.

Girish, A., K. Jolly, N. Alsaadi, M. de la Fuente, A. Recchione, R. An, D. Disharoon, Z. Secunda, S.

Raghunathan, N. F. Luc, C. Desai, E. Knauss, X. Han, K. Hu, H. Wang, U. D. S. Sekhon, N. Rohner, U.

A. Gurkan, M. Nieman, M. D. Neal, and A. Sen Gupta. 2022. Platelet-Inspired Intravenous

Nanomedicine for Injury-Targeted Direct Delivery of Thrombin to Augment Hemostasis in

Coagulopathies. ACS Nano 16:16292–16313.

Holding, M. L., J. E. Biardi, and H. L. Gibbs. 2016a. Coevolution of venom function and venom resistance

in a rattlesnake predator and its squirrel prey. Proceedings of the Royal Society B: Biological

Sciences 283:20152841. Royal Society.

Holding, M. L., D. H. Drabeck, S. A. Jansa, and H. L. Gibbs. 2016b. Venom Resistance as a Model for

Understanding the Molecular Basis of Complex Coevolutionary Adaptations. Integrative and

Comparative Biology 56:1032–1043.

Jansa, S. A., and R. S. Voss. 2011. Adaptive Evolution of the Venom-Targeted vWF Protein in Opossums

that Eat Pitvipers. PLOS ONE 6:e20997. Public Library of Science.

Keyte, A. L., and K. K. Smith. 2008. Harvesting *Monodelphis* Embryos. Cold Spring Harb Protoc

2008:pdb.prot5074.

Lenting, P. J., C. V. Denis, and O. D. Christophe. 2024. How unique structural adaptations support and

coordinate the complex function of von Willebrand factor. Blood 144:2174–2184.

Mackessy, S. P. 2021. Handbook of Venoms and Toxins of Reptiles. CRC Press.

Martins, M., M. S. Araujo, R. J. Sawaya, and R. Nunes. 2001. Diversity and evolution of macrohabitat use,

body size and morphology in a monophyletic group of Neotropical pitvipers (*Bothrops*). Journal

of Zoology 254:529–538.

Matsushita, T., D. Meyer, and J. E. Sadler. 2000. Localization of von Willebrand Factor-binding Sites for

Platelet Glycoprotein Ib and Botrocetin by Charged-to-Alanine Scanning Mutagenesis. Journal of

Biological Chemistry 275:11044–11049.

Nascimbene, A., S. Neelamegham, O. H. Frazier, J. L. Moake, and J. Dong. 2016. Acquired von Willebrand
syndrome associated with left ventricular assist device. Blood 127:3133.

Nash, K. J., L. M. Bacek, P. W. Christopherson, and E. A. Spangler. 2017. In vitro effect of blood cell counts
on multiple-electrode impedance aggregometry in dogs. , doi: 10.2460/ajvr.78.12.1380.

Pruss, C. M., M. Golder, A. Bryant, C. A. Hegadorn, E. Burnett, K. Laverty, K. Sponagle, A. Dhala, C. Notley,
S. Haberichter, and D. Lillicrap. 2011. Pathologic mechanisms of type 1 VWD mutations R1205H
and Y1584C through in vitro and in vivo mouse models. Blood 117:4358–4366.

Ritz, C., F. Baty, J. C. Streibig, and D. Gerhard. 2015. Dose-Response Analysis Using R. PLOS ONE
10:e0146021. Public Library of Science.

Rousmaniere, H., R. Silverman, R. A. White, M. M. Sasaki, S. D. Wilson, J. T. Morrison, and Y. P. Cruz. 2010.
Husbandry of Monodelphis domestica in the study of mammalian embryogenesis. Lab Anim
39:219–226.

Rucavado, A., M. Soto, A. Kamiguti, D. Theakston, J. Fox, T. Escalante, and J. M. Gutiérrez. 2001.
Characterization of Aspercetin, a Platelet Aggregating Component from the Venom of the Snake
Bothrops asper which Induces Thrombocytopenia and Potentiates Metalloproteinase-induced
Hemorrhage. Thromb Haemost 85:710–715.

Ruggeri, Z. M. 2003. Von Willebrand factor, platelets and endothelial cell interactions. Journal of
Thrombosis and Haemostasis 1:1335–1342.

Sadler, J. E. 1998. BIOCHEMISTRY AND GENETICS OF VON WILLEBRAND FACTOR. Annu. Rev. Biochem.
67:395–424.

Shen, C., D. T. Mackeigan, A. A. Shoara, P. Bhoria, G. Zhu, D. Karakas, W. Ma, Z. Y. Chen, R. Xu, S.
Slavkovic, D. Zhang, V. Prifti, Z. Liu, E. G. Cerenzia, P. Chen, M. A. D. Neves, H. Li, F. Xue, R. Yang,
J. Liu, R. Lai, R. Li, and H. Ni. 2024. Novel GPIb-independent platelet aggregation induced by
botrocetin: implications for diagnosis and antithrombotic therapy. Journal of Thrombosis and
Haemostasis S1538783624004471.

Thomazini, C. M., A. T. A. Sachetto, C. Z. De Albuquerque, V. G. De Moura Mattaraia, A. K. De Oliveira, S.
M. D. T. Serrano, I. Lebrun, K. C. Barbaro, and M. L. Santoro. 2021. Involvement of von Willebrand

factor and botrocetin in the thrombocytopenia induced by Bothrops jararaca snake venom. PLoS

Negl Trop Dis 15:e0009715.

Voss, R. S. 2013. Opossums (Mammalia: Didelphidae) in the diets of Neotropical pitvipers (Serpentes:

Crotalinae): Evidence for alternative coevolutionary outcomes? Toxicon 66:1−6.

Voss, R. S., and S. A. Jansa. 2021. Opossums: An Adaptive Radiation of New World Marsupials. JHU

Press.

Voss, R. S., and S. A. Jansa. 2012. Snake-venom resistance as a mammalian trophic adaptation: lessons

from didelphid marsupials. Biological Reviews 87:822−837.

Werner, R. M., and J. A. Vick. 1977. Resistance of the opossum (Didelphis virginiana) to envenomation by

snakes of the family Crotalidae. Toxicon 15:29−32.

Xu, S., J. Wang, Z. Guo, Z. He, and S. Shi. 2020. Genomic Convergence in the Adaptation to Extreme

Environments. Plant Comm 1. Elsevier.

N.d. exclusivo cuíca predando cascavel e coral verdadeira e é picado por elas.

Mammal Range Maps

Monodelphis domestica

Otospermophilus beecheyi

Snake Range Maps

Bothrops jararaca

Crotalus oreganus

Bitis arietans

Venom Inhibition Assays

A. *Bothrops jararaca*

B. *Crotalus oreganus*

C. *Bitis arietans*

Relative Proteinase Activity

Serum Protein (µg)

- ○ Squirrel
- △ Opossum
- + Human

A. Human VWF Opossum vWF

Relative Proteinase Activity

100 μm

B.

Impedence Ω (Clotting)

A.

Collagen
(+ Control)

PBS
(Baseline)

B.

Crotalus adamanteus

Sistrurus miliarius

Bothrops jararaca

C.

Convulxin

Aspercetin

Botrocetin

Time (min)

A — ctrl
B — bot
C — asp
D — cvx
E — ctrl
F — bot
G — asp
H — cvx

100 μm

Platelet Fluorescence (AU)
Time (min)
ctrl
bot
asp
cvx

Platelet Fluorescence (AU)
Time (min)
ctrl
bot
asp
cvx

Normalized Fibrin Fluorescence
Time (min)
ctrl
bot
asp
cvx

Normalized Fibrin Fluorescence
Time (min)
ctrl
bot
asp
cvx