Multiple orthoflaviviruses secrete sfRNA in mosquito saliva to promote transmission by inhibiting MDA5-mediated early interferon response

Short title: Salivary sfRNA is a pan-orthoflavivirus transmission-enhancing factor

Idalba Serrato-Pomar[1,2], Jim Zoladek[3], Hacène Medkour[1], Quentin Narpon[1], Felix Rey-Cadilhac[1], Solena Rossi[1], Stephanie French[4], Cassandra Modahl[4], Wannapa Sornjai[5], Elliott Miot[1], Rodolphe Hamel[1,6,7], Oleg Medianikov[2,8], Dorothée Missé[1], Sébastien Nisole[3], and Julien Pompon[1,5].

MIVEGEC, Univ. Montpellier, IRD, CNRS, Montpellier, France

IHU Méditerranée Infection, Marseille, France

IRIM, Univ. Montpellier, CNRS, Montpellier, France

Liverpool School of Tropical Medicine, Liverpool, United Kingdom

Institute of Molecular Biosciences, Mahidol University, Bangkok, Thailand

Department of Clinical Microbiology and Applied Technology, Faculty of Medical Technology, Mahidol University, Nakhon Pathom, Thailand

Viral Vector Joint unit and Joint Laboratory, Mahidol University, Nakhon Pathom, Thailand

IRD, AP-HM, MEPHI, Aix Marseille University, Marseille, France

Contact : julien.pompon@ird.fr

Abstract

Numerous orthoflaviviruses transmitted through the bites of different mosquito species infect more than 500 million people annually. Bite-initiated skin infection represents a critical and conserved step in transmission and a deeper understanding of this process will promote the design of broad-spectrum interventions to address diverse orthoflavivirus health threats. Here, we identify and characterize a transmission-enhancing viral factor in mosquito saliva that is shared across orthoflaviviruses. Saliva of West Nile virus-infected *Culex* and Zika virus-infected *Aedes* contains a viral non-coding RNA, subgenomic orthoflaviviral RNA (sfRNA), within lipid vesicles distinct from virions. Higher concentration of sfRNA in infectious saliva positively correlates with infection intensity in human cells and skin explants. Early sfRNA delivery into transmission-relevant skin cell types and human skin explant demonstrate that sfRNA is responsible for the infection enhancement. Co-inoculation of sfRNA in a mouse model of transmission enhanced skin infection and worsened disease severity, evidencing the role of salivary sfRNA as a transmission-enhancer. Mechanistically, salivary sfRNA attenuates early interferon response in human skin cells and skin explants by altering MDA5-mediated signaling. Our results, derived from two distinct orthoflaviviruses and supported by prior studies, establish salivary sfRNA as a pan-orthoflavivirus transmission-enhancing factor driven by a conserved viral non-coding RNA.

Keywords: Mosquito-borne diseases, transmission, extracellular vesicles, innate immunity, skin

Introduction

Numerous mosquito-borne orthoflaviviruses cumulatively infect half a billion people annually, resulting in an estimated 150,000 fatalities and economic losses exceeding €9 billion [1]. Given the wide geographic distribution of *Aedes* and *Culex* mosquito vectors, nearly the entire human population faces the risk of infection [2,3]. Furthermore, predictive models considering escalating urbanization, global transportation, and climate change anticipate exacerbation and broadening of associated public health, economic, and social burdens [4]. More concerning is the very likely emergence of yet-unknown mosquito-borne orthoflaviviruses in the near future, further complicating the landscape of orthoflavivirus diseases that necessitate targeted interventions [1]. However, safe and effective intervention to protect against orthoflaviviral diseases are lacking. While vector mosquito control stands as the most widely implemented strategy, its effectiveness in preventing epidemics is only moderate [5]. Scaling up vector control approaches to cover extensive territories is also challenging, and the efficacy is hindered by the rapid development of insecticide resistance [6]. Additionally, there is no curative treatment [7], and the available vaccines pose notable safety concerns [8,9]. A promising approach to confront the rising threat posed by multiple orthoflaviviruses is the design of pan-orthoflavivirus interventions, achieved through the identification of targets that are conserved across flaviviruses.

The transmission of all mosquito-borne orthoflaviviruses occurs during a mosquito bite [10], when infectious virions present in the saliva are deposited in the epidermis and dermis [11,12]. Subsequent productive infection of the skin is essential for transmission [13,14]. However, infection initiation is hindered by a potent antiviral response in the skin [15,16]. Double-stranded viral RNA (dsRNA), an intermediate of viral replication, is detected by retinoic acid-inducible gene I (RIG-I), melanoma

differentiation-associated gene 5 (MDA5) or toll-like receptor 3 (TRL3) [17]. These

three viral RNA sensing branches separately and synergistically activate biochemical

cascades that converge onto the phosphorylation of interferon regulatory factor-3

(IRF3) and/or IRF7, which trigger the expression of type I interferon (IFN). IFN then

triggers an antiviral state by inducing the expression of IFN-stimulated genes (ISGs)

with anti-orthoflaviviral properties [18,19]. Robust evidence supports that mosquito saliva

modulates the antiviral innate immune response to promote orthoflavivirus skin

infection [13,20–22]. Nonetheless, identification of these anti-immune transmission-

enhancing factors remains partial.

Recently, we discovered a mosquito saliva factor that augments saliva

infectivity for dengue virus (DENV), the most prevalent orthoflavivirus [23]. DENV

secretes a non-coding subgenomic orthoflaviviral RNA (sfRNA) with anti-immune

properties [24–26] inside salivary lipid vesicles to enhance saliva infectivity.

Nonetheless, our prior study did not provide evidence of an effect on transmission

and lacked a mechanistic understanding. SfRNA biogenesis results from the

incomplete degradation of orthoflaviviral genomic RNA (gRNA) by 5'-to-3' host

exoribonucleases, which stall on 3'UTR secondary structures, called

exoribonuclease-resistant RNA (xrRNA) [27]. Intriguingly, the process of sfRNA

biogenesis is conserved across orthoflaviviruses [28–31], raising the hypothesis that

salivary sfRNA secretion occurs for multiple orthoflaviviruses.

West Nile (WNV) and Zika (ZIKV) viruses rank among the most important

orthoflaviviruses. WNV is the most widely distributed orthoflavivirus [32] and elicits

symptoms potentially resulting in severe neurological manifestations leading to

mortality or persistent sequelae [33]. Since 2000, WNV has infected an estimated 7

million people, causing 2,700 deaths in the USA [34], and is annually responsible for

over 100 fatalities in the EU [35]. ZIKV, acknowledged as a pathogen of international

concern by the World Health Organization (WHO) in 2014, has infected more than 2

million people since its emergence [36,37]. While ZIKV symptoms typically present as

mild, the infection poses substantial risks of microcephaly in newborns of infected

mothers and of Guillain-Barré syndromes in infected adults. Importantly, WNV and

ZIKV exhibit distinct transmission dynamics with WNV transmitted by various *Culex*

mosquitoes, notably *Cx. quinquefasciatus* [38], whereas ZIKV is primarily disseminated

by *Aedes* mosquitoes, predominantly *Ae. aegypti* [39,40].

Here, using WNV and ZIKV as orthoflavivirus models, we investigated whether

sfRNA secretion in mosquito saliva represents a pan-orthoflavivirus mechanism for

enhancing transmission and elucidated how salivary sfRNA increases transmission.

First, we detected sfRNA in the saliva of *Culex* mosquitoes infected with WNV and

Aedes mosquitoes infected with ZIKV. Second, we found that salivary sfRNA from

WNV-infected *Culex* and ZIKV-infected *Aedes* is packaged within lipid vesicles.

Third, combining approaches with transmission-relevant human skin cell types,

human skin explants and a mouse model of transmission, we demonstrated that

salivary sfRNA enhances orthoflavivirus transmission. Finally, we showed that early

sfRNA exposure suppresses the initial IFN response by disrupting MDA5-mediated

signaling.

Results

SfRNA is secreted in saliva from WNV-infected *Culex* and ZIKV-infected *Aedes*

mosquitoes

WNV can produce up to four sfRNA species by stalling host exoribonucleases at

distinct xrRNA structures [41]. To identify the sfRNA species in *Cx. quinquefasciatus*,

we performed Northern blot on mosquitoes orally infected with WNV and observed a

major band above 500 nt (Fig. 1a), as previously reported in *Cx. pipiens* mosquitoes

[42] and corresponding to the predicted sfRNA1 species at 528 nt. We also noted a

fainter band above 400 nt, which does not align with any predicted xrRNA, but was

previously observed in monkey Vero cells [42]. We then collected saliva and salivary

glands (SG) – the site of saliva production - from *Culex* mosquitoes 10 days post oral

infection and using absolute quantification (S1a,b Fig.) detected WNV gRNA in 80%

of SG with a geometric mean of 1.95×10^3 copies per infected sample and in 60% of

saliva with a geometric mean of 1.97×10^2 copies per infected sample (Fig. 1b).

Using an optimized RT-qPCR protocol for absolute quantification (Supplementary

Materials; S2a-c and S3a,b Fig.), we detected sfRNA in 93% of SG with a geometric

mean of 1.96×10^6 per infected sample and in 100% of saliva with a geometric mean

of 4.15×10^5 copies per infected sample (Fig. 1c). To normalize sfRNA to the

infection level (estimated by gRNA level), we calculated the ratio of sfRNA:gRNA and

observed similar ratios in SG and saliva at 3×10^3 and 3.3×10^3, respectively (Fig.

1d).

We further analysed the data to gain deeper insights of salivary sfRNA

secretion. In SG, we observed a positive correlation between sfRNA and gRNA

copies (Fig. 1e), indicating a relationship between sfRNA and its precursor in this

organ and suggesting that sfRNA biogenesis occurs within the SG. In contrast, in

saliva, we identified a negative correlation between sfRNA and gRNA levels (Fig. 1f).

These opposing correlations between gRNA and sfRNA in the SG and saliva are

consistent with the two viral RNAs being secreted through distinct mechanisms. By

analyzing paired SG and saliva samples from the same mosquito, we further

examined correlations between the two compartments. The quantities of gRNA and

sfRNA in saliva remained relatively constant and showed no significant association

with their levels in the SG (S4a,b Fig.). Collectively, our findings provide evidence

that WNV secretes sfRNA into *Culex* mosquito saliva and underscore the pivotal role

of the SG in the production of expectorated sfRNA.

To further support the role of the SG, we bypassed midgut infection by

inoculating WNV into the thorax of *Culex* mosquitoes and collected SG and saliva at

7 days post-infection (dpi), i.e. earlier than after oral infection due to the shortened

extrinsic incubation period. Direct injection resulted in 100% of infected SG and saliva

samples positive for gRNA, with geometric means of 1.57×10^7 and 2.72×10^4

copies per sample, respectively (Fig. 1g). SfRNA was detected in 100% of infected

SG, with a geometric mean of 1.40×10^8 copies per sample, and in 100% of infected

saliva, with a geometric mean of 2.09×10^5 copies per sample (Fig. 1h). This

corresponded to sfRNA:gRNA ratios at 9.58 in SG and 16 in saliva (Fig. 1i).

Collectively, these results indicate that midgut infection is not required for WNV

sfRNA secretion in saliva.

To determine whether sfRNA secretion in saliva is conserved among

orthoflaviviruses transmitted by different mosquito genera, we inoculated *Aedes*

mosquitoes with ZIKV and collected SG and saliva at 7 dpi. ZIKV gRNA was detected

in 100% of SG and in 100% of saliva samples, with geometric means of 1.00×10^7

and 1.63×10^4 copies per sample, respectively (Fig. 1j; S5a,b Fig.). We quantified

ZIKV sfRNA using primers that annealed to both previously identified sfRNA species

in *Ae. aegypti* [43] and an optimized sfRNA absolute quantification method

(Supplementary materials; S6a-c and S7a,b Fig.). ZIKV sfRNA was detected in 100%

of infected SG, with a geometric mean of 3.22×10^8 copies per sample, and in 100%

of infected saliva, with a geometric mean of 2.47×10^5 copies per sample (Fig. 1k).

This resulted in sfRNA:gRNA ratios of 36.4 in SG and 65.1 in saliva (Fig. 1l).

Collectively, the findings demonstrate that ZIKV, another orthoflavivirus, secretes

sfRNA in the saliva of *Aedes* mosquitoes, indicating that sfRNA secretion in saliva is

conserved mechanism among orthoflaviviruses vectored by different mosquito

species.

Salivary sfRNAs from WNV-infected *Culex* and ZIKV-infected *Aedes* are within

lipid vesicles

In our previous study, we provided biochemical and microscopic evidence that DENV

sfRNA is expectorated within lipid vesicles distinct from virus particles [23]. To

determine whether this phenomenon is shared by other orthoflaviviruses, we

assessed the nuclease resistance of salivary sfRNA from WNV-inoculated *Culex*

mosquitoes following detergent-based disruption of lipid membranes (Fig. 2a). As

control, we confirmed that *in vitro* transcribed WNV sfRNA was degraded by

micrococcal nuclease (MNase) (RNase A/T1 was inefficient in degrading WNV

sfRNA) in the presence of the Triton-X 100 detergent and uninfected *Culex* saliva

(Fig. 2b). Consistent with salivary gRNA and sfRNA being enclosed within lipid

particles, neither of the viral RNAs were degraded by MNase unless pre-treated with

Triton X-100 (Fig. 2c,d). To account for variability in the initial amounts of sfRNA and

gRNA between the saliva pools, we normalized the RNA quantities within each saliva

pool and observed consistent trends (S8a-c Fig.). We then reasoned that if sfRNA

and gRNA are enclosed in separate lipid particles, they would exhibit different

sensitivities to the detergent treatment. Leveraging the inherent variability between

biological replicates – potentially due to mechanical stresses from pipetting and liquid

handling [44] – we found a lack of correlation between sfRNA and gRNA levels after

detergent and MNase treatments (Fig. 2e). This observation suggests that sfRNA is

encapsulated in distinct lipid particles, separate from those containing gRNA.

We repeated the nuclease resistance assay using saliva from ZIKV-inoculated

Aedes mosquitoes. After confirming the degradation of *in vitro*-transcribed ZIKV

sfRNA by RNase A/T1 (RNase) (Fig. 2f), we observed that salivary sfRNA was

protected from RNase degradation unless pretreated with a detergent (Fig. 2g).

Surprisingly, ZIKV gRNA in saliva was degraded by RNase both with and without

detergent pre-treatment (Fig. 2h), potentially due to unintended physical stress on the

virions during the one-day-long process from saliva collection to RNase treatment.

Nonetheless, leveraging this unexpected gRNA sensitization, we observed a clear

discrepancy between sfRNA and gRNA quantities (Fig. 2g,h), consistent with the two

RNA fragments being enclosed in different particles. Similar trends were observed

after normalizing RNA quantities within saliva pools (S8d-f Fig.). Finally, we noted the

absence of correlation between salivary sfRNA and gRNA in samples treated with

both detergent and RNase (Fig. 2i). Altogether, the analysis of saliva from two

different mosquito genera infected with different orthoflaviviruses strongly supports

the conservation of sfRNA secretion within salivary lipid vesicles across multiple

orthoflaviviruses.

Salivary sfRNA increases WNV transmission

To elucidate the role of salivary sfRNA in transmission, we infected *Culex* mosquitoes

with WNV by oral feeding, which introduces greater variability in the salivary

sfRNA:gRNA ratio compared to thoracic inoculation (Fig. 1). Saliva pools were

collected and categorized based on their sfRNA:gRNA ratio into low (ratio < 15),

moderate (15 < ratio < 350), or high (350 < ratio) sfRNA concentrations (Fig. 3a; S1

Table). To validate the use of the saliva-collection media as an inoculum, we

determined that Erioglaucine, used as a proxy for salivation in the saliva-collection

media, did not affect cell viability (S9 Fig.). To support the normalization of saliva

inoculum dose based on gRNA quantities, we showed that sfRNA concentration was

not associated with the gRNA-to-particle-forming-units (PFU) ratio (S10 Fig.), a

measure for virion infectivity. Saliva from the different sfRNA concentration groups

were then used to infect Huh7 human hepatocyte cells, using volumes containing the

same number of gRNA copies (S1 Table). Total saliva volumes were normalized with

uninfected saliva. Quantification of WNV infection at 24 and 48 hours post-infection

(hpi) showed a progressive increase in infection with ascending categories of salivary

sfRNA concentration (Fig. 3b,c) and a positive correlation with sfRNA:gRNA ratio

(Fig. 3d). We repeated the saliva infection experiments with human skin explants

(Fig. 3a). Injection of saliva containing higher sfRNA concentrations (S2 Table) led to

increased WNV infection at 24hpi (Fig. 3e,f).

We next investigated whether early presence of sfRNA enhances infection.

First, we transfected Huh7 cells with *in vitro*-transcribed sfRNA shortly before WNV

infection (Fig. 3g). Intracellular levels of sfRNA or a same-size viral RNA control were

comparable after transfection (S11a Fig.). We observed that sfRNA transfection

increased WNV infection at 24 and 48 hpi (Fig. 3h). Second, as orthoflaviviruses are

thought to initially infect skin fibroblasts and monocytes following a mosquito bite [10,45],

we repeated WNV infection post sfRNA transfection in HFF1 fibroblasts and U937

monocytes. While intracellular levels of sfRNA and control RNA were equivalent

(S11b,c Fig.), infection was increased after sfRNA transfection at 24 and 48hpi in

both cell types, although the increase was lower in U937 (Fig. 3i,j). Third, we injected

a mixture of WNV with either sfRNA or the control RNA into human skin explants

(Fig. 3g) and observed that co-injection with sfRNA enhanced infection at 24 hpi (Fig.

3k).

Finally, we used a mouse model of transmission to evaluate the effect of

sfRNA on infection and disease outcomes. Mice were intradermally injected with a

small volume containing WNV and either sfRNA or the control RNA (Fig. 3l). As a

non-infected control, mice were injected with sfRNA alone. At one day post injection

(dpi), infection at the injection site in the skin was increased by co-injection with

sfRNA (Fig. 3m). At 4 dpi, we confirmed successful systemic infection by detecting

RNAemia (S12 Fig.), although this alone does not inform about disease severity [46–48].

Over a longer observation period, we found that sfRNA co-injection aggravated

disease severity (Fig. 3n), increased weight loss (S13 Fig.) and reduced mouse

survival (Fig. 3o). Altogether, these results – obtained using infectious saliva with

variable sfRNA concentrations and *in vitro*-transcribed sfRNA to mimic salivary

delivery in *in vitro*, *ex vivo* and *in vivo* models - demonstrate that salivary sfRNA

enhances transmission, worsening disease outcomes.

Salivary sfRNA mitigates the IFN response

To decipher how WNV sfRNA impacts the innate immune response, we quantified

expression of *IFN-β* and three ISGs, namely *CXCL10*, *MX1* and *IFI6*, in the Huh7 cell

samples previously infected with saliva containing different sfRNA concentrations

(same samples as in Fig. 3b-d). At 24 hpi, we observed a robust induction of *IFN-β*

and the three ISGs in all samples treated with infectious saliva compared to mock-

infected control (Fig. 4a-d). However, the levels of innate immune gene induction

were similar across saliva samples irrespective of salivary sfRNA concentrations.

Since cells infected with saliva containing higher sfRNA concentrations had higher

viral loads (Fig. 3b) and previous studies have shown a correlation between viral load

and intensity of IFN induction [49,50], these findings could suggest that higher salivary

sfRNA dampened the innate immune response at 24 hpi. In contrast, at 48 hpi, the

expression of all four IFN-related genes correlated with the infection levels (Fig. 4a-

d).

To determine whether sfRNA inhibits the innate immune response, we

quantified the expression of the IFN-related genes in hepatocyte Huh7, fibroblast

HFF1 and monocyte U937 cells that were transfected with sfRNA or the control RNA

prior to WNV infection (same samples as in Fig. 3h-j). In Huh7 cells (Fig. 4e-h),

sfRNA transfection inhibited the expression of *IFN-β* and *IFI6* at both 24 and 48 hpi,

while *CXCL10* and *MX1* expressions were diminished only at 48 hpi. Notably,

whereas *IFN-β*, *CXCL10* and *MX1* expressions increased upon control RNA

transfection from 24 to 48 hpi, sfRNA transfection prevented this innate immune

activation. In HFF1 cells (Fig. 4i-l), sfRNA strongly suppressed *IFN-β* expression at

24 hpi and reduced *CXCL10* expression at both 24 and 48 hpi. In contrast, in U937

cells (Fig. 4m-p), sfRNA transfection did not inhibit the expression of the IFN-related

genes, which may explain the limited effect of sfRNA transfection on infection levels

(Fig. 3j). The lack of inhibition of certain IFN-related genes may reflect differences in

the kinetics or intensity of gene expression among the different cell types.

Finally, we quantified the expression of IFN-related genes in human skin

explants co-injected with WNV and either sfRNA or control RNA (same samples as in

Fig. 3k). SfRNA moderately reduced *IFN-β* expression and significantly diminished

CXCL10, *MX1* and *IFI6* expressions at 24 hpi (Fig. 4q-t). Together, these results

demonstrate that salivary sfRNA dampens innate immune activation in the skin.

SfRNA alters MDA5 signaling of interferon

To determine whether the early presence of sfRNA increases infection by altering the IFN response, we chemically inhibited IFN induction using the TBK1/IKKε inhibitor MRT67037, and infected Huh7 cells with WNV following transfection with either sfRNA or control RNA (Fig. 5a). Successful inhibition of the IFN response was confirmed by the lack of induction for *IFN-β*, *CXCL10*, *MX1* and *IFI6* expressions following control RNA transfection and infection (S14 Fig.). Supporting the hypothesis that sfRNA enhances infection by altering the IFN response, we observed that inhibition of IFN induction abolished sfRNA-mediated enhancement of infection at 24 and 48 hpi (Fig. 5b). The main pattern recognition receptors triggered by orthoflavivirus infection are RIG-I, MDA5 and TLR3 [51]. Since Huh7 cells used here do not express TLR3 [52], we tested the role of RIG-I by using RIG-I-deficient Huh7.5 cells [53] (Fig. 5a). In Huh7.5 cells, the sfRNA-mediated enhancement of WNV infection was maintained at 24 and 48 hpi (Fig. 5c). Next, we silenced MDA5 in Huh7.5 to avoid any confounding effect from RIG-I (S15a,b Fig.). Strikingly, while sfRNA-mediated enhancement of infection was observed following control siRNA transfection, the infection enhancement was abolished at 24 and 48 hpi when MDA5 was silenced (Fig. 5d). Together, these findings indicate that early presence of sfRNA promotes infection by altering MDA5 downstream signaling.

Discussion

The incomplete understanding of orthoflavivirus transmission by mosquitoes hampers the identification of pan-orthoflavivirus targets that could be harnessed for broad-spectrum strategies to simultaneously protect against the multiple orthoflavivirus-related health threats. In this study, we identified and characterized sfRNA in

mosquito saliva as a transmission-enhancing factor conserved across

orthoflaviviruses. Using two phylogenetically distant orthoflaviviruses (WNV and

ZIKV), belonging to distinct phylogenetic groups [54] and vectored by different genera

of mosquitoes (*Culex* and *Aedes* mosquitoes), we showed that both orthoflaviviruses

secrete sfRNA in salivary lipid particles. Combined with our previous discovery of

DENV sfRNA in *Aedes* salivary vesicles [23], the current study implies that sfRNA

secretion in mosquito saliva is conserved across multiple, if not all, orthoflavivirus-

mosquito species systems. We then designed a robust experimental approach to

assess the function of salivary sfRNA in infection by infecting human cells and human

skin explants with infectious mosquito saliva. This unprecedented dataset revealed

that higher salivary sfRNA concentrations correlate with enhanced infection in human

skin. We further demonstrated that sfRNA is responsible for the infection

enhancement by reproducing this effect through the supplementation of *in vitro*-

transcribed sfRNA in transmission-relevant cell types and human skin explants. To

extrapolate from the *in vitro* and *ex vivo* findings and provide *in vivo* evidence for the

role of salivary sfRNA in transmission, we used a mouse model of transmission. We

established that sfRNA co-delivered with orthoflavivirus in the skin increases

transmission and aggravates disease severity. Finally, we elucidated the mechanism

by which the early presence of sfRNA enhances infection, demonstrating that sfRNA

suppresses the early innate immune response and increases WNV infection by

altering MDA5-mediated IFN induction. Altogether, our study establishes that multiple

orthoflaviviruses secrete sfRNA in mosquito saliva to promote bite-initiated skin

infection by inhibiting the innate immune response, thereby enhancing transmission.

Our results show that salivary sfRNA is shielded from nucleases by a

detergent-sensitive layer, indicating the presence of a protective lipid membrane.

These detergent-sensitive compartments could be either virions enclosed by a lipid

bilayer envelop or extracellular vesicles (EVs). EVs are non-replicative, cell-derived

membranous particles secreted into the extracellular space by most cells [55]. We

previously detected EVs in saliva of *Aedes* mosquitoes [23], while others observed EVs

from mosquito cells [56,57]. Multiple lines of evidence support that sfRNA is secreted

inside EVs rather than virions, which by definition contain gRNA. Firstly,

orthoflavivirus virions are spherical structures with a diameter of approximately 55

nm, which is too small to accommodate both gRNA and sfRNA [58–60]. Secondly, we

observed a divergence between sfRNA and gRNA quantities in SG and saliva,

suggesting that sfRNA and gRNA produced in SG are not secreted through the same

pathway. Thirdly, we showed that the particles containing sfRNA and gRNA differ in

their sensitivity to nuclease degradation. Fourthly, the cytosolic origin of the EV

lumen [55,61] would allow the loading of sfRNA. Indeed, sfRNA biogenesis, which

results from gRNA degradation, occurs in the cytosol within processing bodies (PB)

[62], a process supported in mosquitoes by the interaction of sfRNA with PB proteins

such as Staufen [63,64]. In contrast, virions are assembled in the ER and secreted via

the trans-Golgi network [7]. Overall, several viruses exploit EVs to secrete diverse

types of viral RNA fragments [65–68] and our integrated datasets with DENV, WNV and

ZIKV in *Aedes* and *Culex* mosquitoes suggest that orthoflaviviruses load sfRNA into

mosquito EVs for salivary secretion.

We observed that salivary sfRNA favors bite-initiated skin infection by

mitigating the early innate immune response. During the initial stages of

orthoflavivirus infection, the innate immune system mounts a potent antiviral

response mediated by IFN and involving ISGs [15]. RIG-I and MDA5 are the two main

viral RNA recognition receptors responsible for IFN activation via the IRF3/IRF7

phosphorylation signaling cascade [69]. Loss-of-function studies for RIG-I and MDA5

have shown increased infection for WNV, ZIKV and DENV, establishing RIG-I and

MDA5-mediated responses as major barriers to orthoflavivirus infection [70–74]. We

demonstrated that the infection enhancement by the early presence of WNV sfRNA

was abrogated when MDA5 signaling was silenced, whereas RIG-I elimination did

not affect sfRNA's impact on infection. Our results indicate that salivary WNV sfRNA

dampens the IFN response by mitigating MDA5 signaling in Huh7 cells, although

sfRNA target may not be conserved across cell types. Consistent with sfRNA altering

IFN upstream of IRF3/IRF7, one previous study reported that WNV sfRNA modulates

IFN activation upstream of IRF3/IRF7 in murine embryonic fibroblasts (MEF) [75] and

our findings show that chemical inhibition of IRF3/IRF7 activation abrogated the

sfRNA effect. Alternatively, ZIKV sfRNA inhibits the IFN response by stabilizing NS5

inhibition of STAT1 phosphorylation – a step downstream of IRF3/IRF7 [76]. DENV

sfRNA inhibits RIG-I activation by binding TRIM25 to prevent RIG-I ubiquitylation [26]

and interacts with G3BP1, G3BP2 and Caprin1 to reduce ISG translation [77]. Despite

differences in mechanisms, sfRNA from multiple orthoflaviviruses share the ability to

inhibit the innate immune response [24]. However, our mechanistic contribution to WNV

sfRNA function differs from previous studies as we evaluated the early presence of

sfRNA prior to infection establishment and the production of other viral IFN inhibitors

[7].

Salivary sfRNA emerges as the first transmission-enhancing strategy linked to

an orthoflaviviral component. Our study provides evidence that co-delivery of sfRNA

at the bite site increases skin infection and exacerbates disease severity. Previous

studies have underscored the importance of skin infection in transmission [10,14], and

our findings showing the potent immune inhibition of sfRNA in fibroblasts - the

primary dermal cell types - emphasize the critical role of orthoflavivirus replication in

stromal cells before viral migration through myeloid cells [78,79]. Based on our data and

those of others, we propose a model in which the early presence of sfRNA in saliva

prevents the initial activation of innate immune response to facilitate the

establishment of skin infection and amplify subsequent viral dissemination. Our

findings suggest that this model is valid for mosquito-borne orthoflaviviruses.

Limitations of the study

While we propose that sfRNA secretion in salivary vesicles is a shared characteristic

among orthoflaviviruses, our evidence is derived from the examination of only three

orthoflavivirus species vectored by two distinct mosquito genera. Given that there are

over 70 known orthoflavivirus species [80], including other highly pathogenic ones

transmitted by mosquito genera not investigated here - such as for yellow fever virus,

which is transmitted by *Haemagogus* mosquitoes [81] - our findings may not universally

apply to all mosquito-borne orthoflaviviruses. Although sfRNA biogenesis and its anti-

immune functions are conserved across all orthoflavivirus groups tested to date

[24,82,83], it remains to be determined whether salivary sfRNA secretion contributes to

transmission enhancement for all mosquito-borne orthoflaviviruses. Additionally, our

characterization of the lipidic vesicles containing sfRNA is limited. The small volume

of mosquito saliva restricts the extent of biochemical and biophysical analyses

typically performed on EVs [84]. Moreover, isolating sfRNA-containing vesicles from

virions presents challenges due to the physical and chemical similarities between

virions and lipidic vesicles [67].

Materials and methods

Cells and viruses. *Aedes albopictus* C6/36 and Human hepatocyte Huh7 cells were obtained from Cell Lines Services GmbH and Huh7.5 cells were kindly provided by Paul D. Biensiasz (The Rockefeller University, New York, NY, USA). HFF1 (SCRC-1041) and U937 (CRL-1593.2) cells were obtained from ATCC. Mammalian cell lines were grown in Dulbecco's Modified Eagle Medium (DMEM) (Gibco), supplemented with 5 % heat-inactivated fetal bovine serum (FBS) (Eurobio) – except for HFF-1 which was grown with 15 % FBS - and 1 % penicillin/streptomycin mix (Invitrogen) at 37°C with 5 % CO_2. Mosquito cells were grown in Roswell Park Memorial Institute (RPMI) media supplemented with 1 % non-essential amino-acids (ThermoFisher Scientific), 10 % FBS and 1 % penicillin/streptomycin mix at 28°C with 5% CO_2.

WNV strain IS-98-ST1 (or Stork 98) was isolated from a stork in Israel in 1998 [85] and obtained from Dr. Philippe Desprès, Centre de Ressources Biologiques, Institut Pasteur, Paris. ZIKV strain H/PF13 was collected from human serum in French Polynesia in 2013 [86] and obtained from the European Virus Archive-Global (EVAg). Viruses were amplified in C6/36 cells, titrated in BHK21 cells as described [39] and stored in aliquots at -70°C.

Mosquitoes. Experiments with *Culex* mosquitoes were carried out using the *Culex quinquefasciatus* SLAB strain, collected in California, USA [87], and obtained from the Institut des Sciences de l'Evolution in Montpellier, France. Experiments with *Aedes* mosquitoes were carried out using the *Aedes aegypti* Bora-Bora strain collected in French Polynesia in 1980 [88]. Eggs from *Cx. quinquefasciatus* were hatched and reared at 25 ± 1°C, whereas eggs from *Ae. aegypti* were reared at 27 ± 1°C. Both species were reared at 70 ± 5% relative humidity and 12h:12h day:night. Larvae were

distributed in plastic trays at a density of 200 individuals per tray and fed half a tablet

of concentrated yeast and 1 g of TetraMin (Tetra) on the day of hatching, and then

1.5 g of TetraMin every two days until pupation. Pupae were transferred to cages and

supplied with 10 % sugar solution and water *ad libitum*.

Mice. C57BL/6J male mice were purchased from Charles-River (France) and housed

in ventilated cages in NexGen Mouse 500 (Allentown; Serial number: 1304A0078) in

the biosafety level 3 animal facility at MIVEGEC-IRD, Montpellier, France. Mice were

maintained with a 17h:7h light/dark cycle, 53-57 % humidity, 20-24°C temperature

and provided with an irradiation-sterilized mouse diet (A03, SAFE, France) and

sterilized water *ad libitum*. Upon arrival, mice were let to rest for one week before

experiments. Animal protocols were approved by the national ethical committee

(permission numbers: 43466; 31273).

Mosquito oral infection. Four-day-old female *Cx. quinquefasciatus* were starved for

12h, and offered an infectious blood meal for 60 min at sunset time, using the

Hemotek feeding system (Discovery Workshops) with chicken skin (obtained from

spring chicken purchased in supermarket). The artificial blood meal contained WNV

at 8 x 10^5 plaque forming unit (PFU)/ml, 50 % volume of washed erythrocytes from

rabbit's blood (animals housed in the BSL2 VectoPole animal facility, authorization

number: H3417221), 25 mM ATP (ThermoFisher Scientific), 5 % FBS (Eurobio-

scientific) and RPMI to complete the volume to 2.5 ml. Engorged females were

selected and maintained in the rearing conditions for 10 days before analysis.

Mosquito inoculation. Five-day-old cold-anesthetized female *Cx. quinquefasciatus*

or *Ae. aegypti* were intrathoracically inoculated with 69 nl containing 100 PFU of

WNV or 50 PFU of ZIKV, respectively, using Nanoject II (Drummond Scientific

Company) and needles made from 1.14 mm O.D. glass capillaries x 1.75″ length

(Drummond). Mosquitoes were then maintained in rearing conditions for 7 days

before analysis.

Mosquito salivary glands dissection and saliva collection. Cold-anesthetized

mosquitoes had their wings and legs removed before inserting individual proboscises

into 20 µl filter-sterile tips containing 10 µl of RPMI (for RNase resistance assay) or of

DMEM (for human cell infection with WNV) with 2% 25 mM Erioglaucine

(SigmaAldrich) for 30 min at 25°C. Mosquitoes with a blue abdomen (color of

Erioglaucine) were considered to have salivated and the corresponding media were

collected either individually or as pools. Salivary glands were dissected after

salivation, labeled to identify the associated saliva and homogenized using Fast-Prep

bead bitter homogenizer (MP) with glass beads.

sfRNA production. Full length *in vitro*-transcribed sfRNAs from WNV were

generated by amplifying the sfRNA sequence from virus cDNA using the following

primer   pair   with   T7-tagged   forward: 5'-

TAATACGACTCACTATAGGGAGTCAGGCCGGGAAGTTCC-3'   and   5'-

AGATCCTGTGTTCTCGCACC-3'. Amplicons were reverse transcribed using

Megascript T7 kit (Ambion), monophosphorylated at the 5' end using RNA 5'

Polyphosphatase (Lucigen), extracted in RNase-free water (Invitrogen) using

E.Z.N.A. Total RNA kit I (Omega) to remove the polyphosphatase and folded by

slowly reducing the temperature from 95 to 4°C with 5 mM of Mg2+, which regulates

sfRNA folding [89]. sfRNA quantity was estimated using nanodrop (ThermoFisher

Scientific) and used to calculate copy number.

As control, a same-size RNA fragment (i.e., 483 nt) corresponding to a part of

the dengue virus NS2 gene was *in vitro*-transcribed using the primers 5'-

TAATACGACTCACTATAGGGGCAGCTGGACTACTCTTGAG-3',   5'-

GGTCCTGTCATGGGAATGTC-3' [63], monophosphorylated and folded as for sfRNA.

Northern Blot. Northern blot was conducted using NorthernMax Kit (Ambion) with

modifications to manufacturer's protocol as described [23]. Total RNA from 100

infected mosquitoes was extracted using TRIzol reagent (Invitrogen) and separated

on a denaturing gel with 5 % Acrylamide/Bisacrylamide 19:1 (Starlab) and 8 M Urea

(Invitrogen). Biotinylated single-stranded RNA ladder (Kerafast) was also loaded on

the gel. RNA was transferred onto a Hybond-N+ nylon membrane (Merck) using

Trans-Blot Turbo (Bio-Rad) at constant 1.3 A for 30 min with Voltage ☐ 25 V. The

membrane was UV-crosslinked, pre-hybridized and hybridized overnight with a biotin-

16-dUTP (Roche) labeled dsDNA probe (800 ng) generated from the primers used to

amplify the qPCR sfRNA targets. After washes, the membrane was blocked using

Odyssey Blocking Buffer (LI-COR) and stained with IRDYE 800cw streptavidin (LI-

COR) in TBS. Pictures were taken with ChemiDoc (Bio-Rad).

Absolute quantification of gRNA copies. Total RNA was extracted using E.Z.N.A.

Total RNA kit I (Omega Bio-Tek). gRNA for WNV was quantified using iTaq Universal

SYBR   Green   one-step   RT-qPCR   kit   (Bio-Rad)   with   primers: 5'-

ATTCGGGAGGAGACGTGGTA-3'   and   5'-CAGCCGCCAACATCAACAAA-3';   in

LighCycler 96 thermocycler (Roche) with the following thermal profile: 50°C for 10

min, 95°C for 2 min, and 40 cycles at 95°C for 15s, 60°C for 15s and 72°C for 20s.

gRNA for ZIKV was quantified using iTaq Universal Probes one-step RT-qPCR kit

(Bio-Rad) with primers : 5'-TTGGTCATGATACTGCTGATTGC-3' and 5'-

CCTTCCACAAAGTCCCTATTGC-3' and probe: 5'-

CGGCATACAGCATCAGGTGCATAGGAG-3'; in LighCycler 96 (Roche) with the

following thermal profile: 50°C for 15 min, 95°C for 2 min, and 45 cycles at 95°C for

10s and 60°C for 30s.

Absolute quantification for both gRNAs was obtained by generating standard

equations using *in vitro*-transcribed RNA qPCR targets. RNA targets for WNV and

ZIKV were produced by amplifying virus cDNA with primer pairs: 5'-

TAATACGACTCACTATAGGGATTCGGGAGGAGACGTGGTA-3' / 5'-

CAGCCGCCAACATCAACAAA-3', and 5'-

TAATACGACTCACTATAGTTGGTCATGATACTGCTGATTGC-3' / 5'-

CCTTCCACAAAGTCCCTATTGC-3', respectively. Amplicons were reverse

transcribed using Megascript T7 kit (Ambion), extracted using RNeasy Mini Kit

(Qiagen) and quantified using nanodrop (ThermoFisher Scientific) to calculate copy

number. Serial dilutions were used to establish absolute standard equations. Using

the gRNA templates, the limit of detection (LoD) at 95% was determined by

calculating fractions of detected samples in three replicates of serial dilutions [90].

gRNA detection rate was calculated as the proportion of samples with

detectable amount of gRNA among all samples.

Absolute quantification of sfRNA copies. sfRNA and 3'UTR were jointly quantified

in total RNA extracts from E.Z.N.A. Total RNA kit I using the following primers for

WNV: 5'-AGTTGAGTAGACGGTGCTGC-3' and 5'-CCGTAGCGTGGTCTGACATT-

3', and for ZIKV: 5'-GCTGGGAAAGACCAGAGACT-3' and 5'-

CTATTCGGCGATCTGTGCCT-3'. Two different qPCR conditions were applied for

each pair of primers. First, quantification for both viruses was conducted using iTaq

Universal SYBR Green one-step RT-qPCR kit (Bio-Rad) in LightCycler with the

following thermal profile: 50°C for 10 min, 95°C for 2 min, and 40 cycles at 95°C for

10s and 60°C for 25s. Second, quantification with WNV primers was conducted in the

same conditions but for the reverse transcription step set at 60°C for 20 min.

Additionally, quantification with ZIKV primers was conducted using a two-step RT-

qPCR. RT was conducted with the reverse primer and M-MLV enzyme (Promega)

with a denaturation/annealing step at 70°C and following manufacturer's conditions

with RT at 37°C. qPCR was then performed in AriaMx thermocycler (Agilent) using

EvaGreen qPCR MixPlus (Euromedex) with the following thermal profile: 90°C for 15

min, and 40 cycles at 95°C for 15s, 62°C for 30s and 72°C for 20s.

Absolute quantification and LoD for sfRNA for WNV and ZIKV were obtained

as described above for gRNA but using dilutions of sfRNA copies.

sfRNA copy number was calculated by subtracting the number of 3'UTR +

sfRNA copies to the number of gRNA copies. For samples that contained detectable

amount of gRNA, sfRNA:gRNA ratio was calculated by dividing the number of sfRNA

over the number of gRNA. sfRNA detection rate was calculated as the number of

samples with detectable amount of sfRNA among samples with detectable amount of

gRNA.

RNase resistance assay. 10^7 copies of *in vitro*-transcribed sfRNA were mixed with

10 saliva samples from uninfected mosquitoes and subjected to RNAse resistance

assay. Pools of 40 saliva samples from intrathoracically-inoculated mosquitoes were

divided into 4 equal volumes and subjected to the different conditions of the RNAse

treatment assay.

For WNV, samples were supplemented with 10 µl of 0.1 % Triton X-100

(Sigma) for 30 min at 4°C, before adding 0.5 µl of Micrococcal nuclease (MNase)

(Thermo Scientific) and 10 µl of buffer (50mM Tris-HCl pH 8; 10mM $CaCl_2$ final

concentration) for 1h. Controls for Triton X-100 and RNase were added the same

volume of PBS instead and maintained at the corresponding temperatures. RNA was

then extracted using QIAamp viral RNA kit (Qiagen).

For ZIKV, the same treatments were applied except that 3 µL of RNase A/T1

(ThermoFisher Scientific) was added for 30 min at 37°C instead of MNase.

Infection of human cells with WNV infectious *Culex* saliva. Pools of 10 saliva

samples from *Culex* mosquitoes infected by oral feeding with 10^8 pfu/ml of WNV

were collected and stored at -70°C. gRNA and sfRNA were quantified from a frozen

aliquot of each pool. 5,000 Huh7 cells per well of a 96-well flat-bottomed plate were

inoculated with 1,000 gRNA copies of the different saliva pools for 2h. Total volume

of the infected saliva was homogenized to 50 µl by adding uninfected saliva from

Culex female mosquitoes. After infection, inoculum was removed and cells were

supplemented with 2 % FBS DMEM and collected at 24 and 48h post inoculation by

adding TRK lysis buffer and total RNA was extracted using E.Z.N.A. total RNA kit I.

Injection of WNV infectious *Culex* saliva into human skin explants. Five mm

diameter Human skin explants obtained from Biopredic International (France) were

decontaminated by washing in 25 mM HEPES (Gibco), 400 U/ml P/S and 10 µg/ml

amphotericin B (biowest) in DMEM. Explants were maintained individually in 8 µm

cell culture insert (Thincert) with 200 µl of two-week skin culture medium w/o animal

components (MIL215C, Biopredic) at 37°C, 5% CO_2. Explants were inoculated with

760 gRNA copies of saliva pools. Total volume of the infected saliva was

homogenized to 30 µl by adding uninfected saliva from *Culex* female mosquitoes. At

24 h post injection, explants were cut into small pieces and, homogenized using the

bead bitter homogenizer and glass beads before RNA extraction as detailed for

mouse skin above.

Relative quantification of WNV gRNA and IFN-related genes in human cells and

skin explants. RNA was reverse transcribed using the PrimeScript RT Reagent Kit

(Perfect RealTime, Takara Bio Inc.). Real-time PCR reaction was performed in

duplicate using Takyon ROX SYBR MasterMix blue dTTP (Eurogentec) on an

Applied Biosystems QuantStudio 5 (Thermo Fisher Scientific) in 384-well plates.

Transcripts were quantified using the following program: 3 min at 95°C followed by

35 cycles of 15 s at 95°C, 20s at 60°C, and 20s at 72°C. Values for each transcript

were normalized to the geometric means of Ct values of 4 different housekeeping

genes (*RPL13A*, *ACTB*, *B2M*, and *GAPDH*) using the 2-ΔΔCt method. Primers used

for quantification of transcripts by real-time qPCR were: RPL13A 5'-

AACAGCTCATGAGGCTACGG-3' and 5'-TGGGTCTTGAGGACCTCTGT-3', ACTB

5'-CTGGAACGGTGAAGGTGACA-3' and 5'-AAGGGACTTCCTGTAACAATGCA-3',

B2M   5'-TGCTGTCTCCATGTTTGATGTATCT-3'   and   5'-

TCTCTGCTCCCCACCTCTAAGT-3', GAPDH 5'-TGCACCACCAACTGCTTAGC-3'

and   5'-GGCATGGACTGTGGTCATGAG-3',   IFN-β   5'-

TGCTCTCCTGTTGTGCTTCTC-3'   and   5'-CAAGCCTCCCATTCAATTGCC-3',

CXCL10   5'-CGCTGTACCTGCATCAGCAT-3'   and   5'-

GCAATGATCTCAACACGTGGAC-3', Mx1 5'-AAGCTGATCCGCCTCCACTT-3' and

5'-TGCAATGCACCCCTGTATACC-3',   IFI6   5'-GGGTGGAGGCAGGTAAGAAA-3'

and 5'-GACGGCCATGAAGGTCAGG-3'. Primers for WNV gRNA were as described

above.

Infection of human cells post sfRNA transfection. 2×10^5 Huh7, HFF1, U937 or

Huh7.5 cells were transfected with 10^{10} copies of either monophosphorylated-folded-

sfRNA or the control RNA using TransIT-mRNA Transfection Kit (Mirus) for 2 h. The

transfection media was removed and cells were immediately infected with WNV at a

multiplicity of infection, MOI, of 0.005 for 1h. Cells were then maintained in 2% FBS

media for 24 and 48 h before extracting RNA using E.Z.N.A. total RNA kit I.

Co-injection of sfRNA and WNV in human skin explants. Five mm diameter

human skin explants as detailed above were inoculated with 10^2 PFU of WNV mixed

with 10^9 copies of monophosphorylated-folded-sfRNA or the control RNA in a total

volume of 4 µl using a Nanoject II injector (Drummond) and glass needles. At 24 h

post injection, explants were cut into small pieces and, homogenized using the bead

bitter homogenizer and glass beads before RNA extraction as detailed for mouse

skin above.

Co-injection of sfRNA and WNV in mice. Five-to-seven-week-old mice were

shaved with the animal trimmer (VITIVA MINI, BIOSEB) on the lower back one day

prior to injection to limit shaving-induced inflammation. Mice anesthetized by

intraperitoneal injection of 0.2 ml/mouse of 10 mg/ml of ketamine (Imalgène 1000,

Boehringer Ingelheim Animal Health) and 1 mg/ml of xylazine (Rompun 2%, Elanco

GmbH) were intradermally inoculated with 10^3 PFU of WNV mixed with 10^{10} copies of

monophosphorylated-folded-sfRNA or the same number of the control RNA. The total

volume of inoculum was 4 µl. As an additional control, other mice were only injected

with monophosphorylated-folded-sfRNA.

At 24 h post-injection, skin biopsies from the injection site and draining lymph

nodes were collected and homogenized in 340 µL of TRK lysis buffer using Fast-

Prep bead bitter homogenizer (MP) set at 1.4 m/s for 60 sec. with 1 nm glass beads.

Total RNA was extracted using E.Z.N.A. total RNA kit I. At 4 days post injection,

blood samples were collected via mandibular puncture. RNA was extracted using

QIamp viral RNA kit (Qiagen) and WNV gRNA per ml of blood was absolutely

quantified after evaluating blood volume by pipetting. Clinical signs and weight were

registered daily to determine a clinical score (CS) and calculate the percentage of

weight loss until 14 days post-injection. CS of 0 was assigned to healthy mice; 1 for

mice with ruffled fur, lethargy, hunched posture, no paresis, normal gait; 2 for mice

with altered gait, limited movement in 1 hind limb; 3 for lack of movement, paralysis in

1 or both hind limbs; 4 for moribund mice; and 5 for dead mice [91]. Mice were

euthanized under anesthesia if they displayed neurological symptoms, severe

distress, or weight loss exceeding 20%, or reached 12 days post-injection.

Chemical inhibition of interferon signaling. Media of 2 x 10^5 Huh7 cells was

supplemented with 10 µM of MRT67307 (MedChemExpress) diluted in water for 1h,

then during transfection of monophosphorylated-folded-sfRNA or the control RNA,

and infection with WNV as described above.

MDA-5 Silencing. 2 x 10^5 Huh7.5 cells were transfected overnight with 5 pmol of

multiplex siRNAs (SI05130608, Qiagen) using Lipofectamine RNAiMax

(ThermoFisherScientific). As a control, Allstars multiplex siRNAs (SI91027281,

Qiagen) were similarly transfected. 48 h later, cells were transfected with

monophosphorylated-folded-sfRNA or the RNA control and subsequently infected

with WNV as described above.

Statistical analysis

Comparison of gRNA and sfRNA copies, relative gRNA fold changes and IFN-related

genes were conducted using one tailed T-test or post-hoc Fisher's LSD test. Absolute

number and relative values of gRNA and sfRNA copies were log-transformed before

statistical analysis. Correlation analysis was performed using linear regression

analysis. Survival analysis were performed with Gehan-Breslow-Wilcoxon test. The

statistical analyses were performed using GraphPad Prism v9.0.

The progression of clinical scores in mice was analyzed using a general linear

mixed model with a negative binomial distribution to account for overdispersed data.

The progression of weight change in mice was analyzed using a general linear mixed

model with an inverse Gaussian distribution and log link function, appropriate for non-

negative, right-skewed data. The model assessed the effects of days post-infection

and experimental groups (WNV + sfRNA vs. WNV + Ctl.), including random

intercepts for individual mice to account for variability across subjects. For pairwise

comparisons between groups, post hoc analyses were conducted using estimated

marginal means (emmeans) with Tukey's adjustment for multiple comparisons. All

analyses were conducted in R (version 4.3.3) using the lme4 [92], glmmTMB [93], and

emmeans packages [94].

Acknowledgments

We thank the VectoPole in Montpellier and especially Bethsabée Scheid, Carole Ginibre and Arnaud Berthomieu for providing mosquito eggs. We also thank Audrey Vernet from the proteomic platform. PhD scholarships for IS was provided by the Institut Méditerranéen Hospitalier (IHU, Marseille), and for FRC by the French Ministry of Research and Higher Education (CBS2 doctoral school, Montpellier).). Post-doctoral fellowships from Fondation pour la Recherche Médicale to HM (SPF202110013925) and to EFM (ARF202309017577). Support for the research was provided by the French Agence Nationale pour la Recherche (ANR-20-CE15-0006) and by the EU (HORIZON-HLTH-2023-DISEASE-03 #101137006) to JP.

Competing interests

The authors declare no competing interests.

Author contributions

Conceptualization: ISP, JP

Formal analysis: ISP, JP

Funding acquisition: JP

Investigation: ISP, JZ, HM, SR, SN, SF, CM, WS, SN

Methodology: ISP, JZ, HM, QN, FRC, RH, SN, EM, RH, DM, SN, JP

Project administration: JP

Supervision: OM, JP

Visualization: ISP, JP

Writing – original draft: ISP, JP

Writing – review & editing: all authors

References

1. Pierson, T. C. & Diamond, M. S. The continued threat of emerging flaviviruses.

Nat. Microbiol. **5**, 796–812 (2020).

2. Bhatt, S. *et al.* The global distribution and burden of dengue. *Nature* **496**, 504–7

(2013).

3. Habarugira, G., Suen, W. W., Hobson-Peters, J., Hall, R. A. & Bielefeldt-Ohmann,

H. West nile virus: An update on pathobiology, epidemiology, diagnostics, control

and "One health" implications. *Pathogens* **9**, 1–51 (2020).

4. Kraemer, M. U. G. *et al.* Past and future spread of the arbovirus vectors Aedes

aegypti and Aedes albopictus. *Nat. Microbiol.* 1 (2019) doi:10.1038/s41564-019-

0376-y.

5. Ooi, E.-E., Goh, K.-T. & Gubler, D. J. Dengue prevention and 35 years of vector

control in Singapore. *Emerg. Infect. Dis.* **12**, 887–893 (2006).

6. Liu, N. Insecticide Resistance in Mosquitoes: Impact, Mechanisms, and Research

Directions. *Annu. Rev. Entomol.* **60**, 537–559 (2015).

7. Barrows, N. J. *et al.* Biochemistry and Molecular Biology of Flaviviruses. *Chem.*

Rev. **118**, 4448–4482 (2018).

8. Sridhar, S. *et al.* Effect of Dengue Serostatus on Dengue Vaccine Safety and

Efficacy. *N. Engl. J. Med.* 327–340 (2018) doi:10.1056/nejmoa1800820.

9. Halstead, S. B. Dengvaxia sensitizes seronegatives to vaccine enhanced disease

regardless of age. *Vaccine* **35**, 6355–6358 (2017).

10. Pingen, M., Schmid, M. A., Harris, E. & McKimmie, C. S. Mosquito Biting

Modulates Skin Response to Virus Infection. *Trends Parasitol.* **33**, 645–657

(2017).

11. Choumet, V. *et al.* Visualizing Non Infectious and Infectious Anopheles gambiae

Blood Feedings in Naive and Saliva-Immunized Mice. *PLOS ONE* **7**, 1–13 (2012).

12. Styer, L. M. *et al.* Mosquitoes inoculate high doses of West Nile Virus as they

probe and feed on live hosts. *PLoS Pathog.* **3**, e132 (2007).

13. Styer, L. M. *et al.* Mosquito Saliva Causes Enhancement of West Nile Virus

Infection in Mice. *J. Virol.* **85**, 1517–1527 (2011).

14. Bryden, S. R. *et al.* Pan-viral protection against arboviruses by activating skin

macrophages at the inoculation site. *Sci. Transl. Med.* **12**, (2020).

15. Rathore, A. P. S. & St. John, A. L. Immune responses to dengue virus in the skin.

Open Biol. **8**, (2018).

16. Lei, V., Petty, A. J., Atwater, A. R., Wolfe, S. A. & MacLeod, A. S. Skin Viral

Infections: Host Antiviral Innate Immunity and Viral Immune Evasion. *Front.*

Immunol. **11**, (2020).

17. Rehwinkel, J. & Gack, M. U. RIG-I-like receptors: their regulation and roles in

RNA sensing. *Nat. Rev. Immunol.* **20**, 537–551 (2020).

18. Lesage, S. *et al.* Discovery of Genes that Modulate Flavivirus Replication in an

Interferon-Dependent Manner. *J. Mol. Biol.* **434**, 167277 (2022).

19. Wichit, S. *et al.* Interferon-inducible protein (IFI) 16 regulates Chikungunya and

Zika virus infection in human skin fibroblasts. *EXCLI J.* **18**, 467–476 (2019).

20. Vogt, M. B. *et al.* Mosquito saliva alone has profound effects on the human

immune system. *PLoS Negl. Trop. Dis.* **12**, e0006439 (2018).

21. McCracken, M. K. *et al.* Route of inoculation and mosquito vector exposure

modulate dengue virus replication kinetics and immune responses in rhesus

macaques. *PLoS Negl. Trop. Dis.* **14**, e0008191 (2020).

22. McCracken, M. K., Christofferson, R. C., Chisenhall, D. M. & Mores, C. N.

Analysis of Early Dengue Virus Infection in Mice as Modulated by Aedes aegypti

Probing. *J. Virol.* **88**, 1881–1889 (2014).

23. Yeh, S.-C. *et al.* The anti-immune dengue subgenomic flaviviral RNA is present in

vesicles in mosquito saliva and is associated with increased infectivity. *PLOS*

Pathog. **19**, e1011224 (2023).

24. Slonchak, A. & Khromykh, A. A. Subgenomic flaviviral RNAs: What do we know

after the first decade of research. *Antiviral Res.* **159**, 13–25 (2018).

25. Bidet, K., Dadlani, D. & Garcia-Blanco, M. A. G3BP1, G3BP2 and CAPRIN1 Are

Required for Translation of Interferon Stimulated mRNAs and Are Targeted by a

Dengue Virus Non-coding RNA. *PLOS Pathog.* **10**, 1–17 (2014).

26. Manokaran, G. *et al.* Dengue subgenomic RNA binds TRIM25 to inhibit interferon

expression for epidemiological fitness. *Science* **350**, 217–221 (2015).

27. Pijlman, G. P. *et al.* A Highly Structured, Nuclease-Resistant, Noncoding RNA

Produced by Flaviviruses Is Required for Pathogenicity. *Cell Host Microbe* **4**,

579–591 (2008).

28. Akiyama, B. M. *et al.* Zika virus produces noncoding RNAs using a multi-

pseudoknot structure that confounds a cellular exonuclease. *Science* **354**, 1148–

1152 (2016).

29. Kieft, J. S. *et al.* The Structural Basis of Pathogenic Subgenomic Flavivirus RNA

(sfRNA) Production. *Science* **344**, 307–310 (2014).

30. Yeh, S.-C. & Pompon, J. Flaviviruses Produce a Subgenomic Flaviviral RNA That

Enhances Mosquito Transmission. *DNA Cell Biol.* **37**, 154–159 (2018).

31. MacFadden, A. *et al.* Mechanism and structural diversity of exoribonuclease-

resistant RNA structures in flaviviral RNAs. *Nat. Commun.* **9**, 119 (2018).

32. Heidecke, J., Lavarello Schettini, A. & Rocklöv, J. West Nile virus eco-

epidemiology and climate change. *PLOS Clim.* **2**, e0000129 (2023).

33. Suthar, M. S., Diamond, M. S. & Gale Jr, M. West Nile virus infection and

immunity. *Nat. Rev. Microbiol.* **11**, 115–128 (2013).

34. Centers for Disease Control and Prevention, N. C. for E. and Z. I. D. Historic Data

(1999-2022) _ West Nile Virus _ CDC. *Centers for Disease Control*

https://www.cdc.gov/westnile/statsmaps/historic-

data.html?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fwestnile%2Fstat

smaps%2FcumMapsData.html (2023).

35. European Centre for Disease Prevention and Control. Historical data by year -

West Nile virus seasonal surveillance. *An agency of the European Union*

https://www.ecdc.europa.eu/en/west-nile-fever/surveillance-and-disease-

data/historical (2023).

36. Kazmi, S. S., Ali, W., Bibi, N. & Nouroz, F. A review on Zika virus outbreak,

epidemiology, transmission and infection dynamics. *J. Biol. Res.* **27**, 5 (2020).

37. Hamel, R. *et al.* Zika virus: epidemiology, clinical features and host-virus

interactions. *Microbes Infect.* **18**, 441–449 (2016).

38. Chancey, C., Grinev, A., Volkova, E. & Rios, M. The global ecology and

epidemiology of west nile virus. *BioMed Res. Int.* **2015**, (2015).

39. Pompon, J. *et al.* A Zika virus from America is more efficiently transmitted than an

Asian virus by Aedes aegypti mosquitoes from Asia. *Sci. Rep.* **7**, 1215 (2017).

40. Aubry, F. *et al.* Enhanced Zika virus susceptibility of globally invasive Aedes

aegypti populations. *Science* **370**, 991–996 (2020).

41. Funk, A. *et al.* RNA Structures Required for Production of Subgenomic Flavivirus

RNA. *J. Virol.* **84**, 11407 (2010).

42. Göertz, G. P. *et al.* Noncoding Subgenomic Flavivirus RNA Is Processed by the

Mosquito RNA Interference Machinery and Determines West Nile Virus

Transmission by Culex pipiens Mosquitoes. *J. Virol.* **90**, 10145–10159 (2016).

43. Pallarés, H. M. *et al.* Zika Virus Subgenomic Flavivirus RNA Generation Requires

Cooperativity between Duplicated RNA Structures That Are Essential for

Productive Infection in Human Cells. *J. Virol.* **94**, 343–363 (2020).

44. Subedi, P., Schneider, M., Atkinson, M. J. & Tapio, S. Isolation of Proteins from

Extracellular Vesicles (EVs) for Mass Spectrometry-Based Proteomic Analyses.

in *Proteomic Profiling: Methods and Protocols* (ed. Posch, A.) 207–212 (Springer

US, New York, NY, 2021). doi:10.1007/978-1-0716-1186-9_12.

45. Wang, Z.-Y., Nie, K.-X., Niu, J.-C. & Cheng, G. Research progress toward the

influence of mosquito salivary proteins on the transmission of mosquito-borne

viruses. *Insect Sci.* **31**, 663–673 (2024).

46. Diamond, M. S. *et al.* A Critical Role for Induced IgM in the Protection against

West Nile Virus Infection. *J. Exp. Med.* **198**, 1853–1862 (2003).

47. Shrestha, B. & Diamond, M. S. Role of CD8+ T Cells in Control of West Nile Virus

Infection. *J. Virol.* **78**, 8312–8321 (2004).

48. Shrestha, B., Samuel, M. A. & Diamond, M. S. CD8+ T Cells Require Perforin To

Clear West Nile Virus from Infected Neurons. *J. Virol.* **80**, 119 (2006).

49. Zhao, M., Zhang, J., Phatnani, H., Scheu, S. & Maniatis, T. Stochastic Expression

of the Interferon-β Gene. *PLOS Biol.* **10**, 1–16 (2012).

50. Garcia, M. *et al.* Innate Immune Response of Primary Human Keratinocytes to

West Nile Virus Infection and Its Modulation by Mosquito Saliva. *Front. Cell.*

Infect. Microbiol. **8**, 387 (2018).

51. Suthar, M. S., Aguirre, S. & Fernandez-Sesma, A. Innate Immune Sensing of

Flaviviruses. *PLOS Pathog.* **9**, e1003541 (2013).

52. Li, K., Chen, Z., Kato, N., Gale, M. & Lemon, S. M. Distinct Poly(I-C) and Virus-

activated Signaling Pathways Leading to Interferon-β Production in Hepatocytes*.

J. Biol. Chem. **280**, 16739–16747 (2005).

53. Blight, K. J., McKeating, J. A. & Rice, C. M. Highly Permissive Cell Lines for

Subgenomic and Genomic Hepatitis C Virus RNA Replication. *J. Virol.* **76**,

13001–13014 (2002).

54. Gould, E. A., de Lamballerie, X., de A. Zanotto, P. M. & Holmes, E. C. Evolution,

epidemiology, and dispersal of flaviviruses revealed by molecular phylogenies. in

Advances in Virus Research vol. 57 71–103 (Academic Press, 2001).

55. van Niel, G., D'Angelo, G. & Raposo, G. Shedding light on the cell biology of

extracellular vesicles. *Nat. Rev. Mol. Cell Biol.* **19**, 213–228 (2018).

56. Vora, A. *et al.* Arthropod EVs mediate dengue virus transmission through

interaction with a tetraspanin domain containing glycoprotein Tsp29Fb. *Proc.*

Natl. Acad. Sci. 201720125 (2018) doi:10.1073/pnas.1720125115.

57. Reyes-Ruiz, J. M. *et al.* Isolation and characterization of exosomes released from

mosquito cells infected with dengue virus. *Virus Res.* **266**, 1–14 (2019).

58. Kuhn, R. J. *et al.* Structure of Dengue Virus: Implications for Flavivirus

Organization, Maturation, and Fusion. *Cell* **108**, 717–725 (2002).

59. Mukhopadhyay, S., Kim, B.-S., Chipman, P. R., Rossmann, M. G. & Kuhn, R. J.

Structure of West Nile Virus. *Science* **302**, 248–248 (2003).

60. Kostyuchenko, V. A. *et al.* Structure of the thermally stable Zika virus. *Nature* **533**,

425–8 (2016).

61. Raposo, G. & Stoorvogel, W. Extracellular vesicles: Exosomes, microvesicles,

and friends. *J. Cell Biol.* **200**, 373–383 (2013).

62. Pijlman, G. P. *et al.* A highly structured, nuclease-resistant, noncoding RNA

produced by flaviviruses is required for pathogenicity. *Cell Host Microbe* **4**, 579–

591 (2008).

63. Yeh, S.-C. *et al.* Characterization of dengue virus 3'UTR RNA binding proteins in

mosquitoes reveals that AeStaufen reduces subgenomic flaviviral RNA in saliva.

PLOS Pathog. **18**, e1010427 (2022).

64. Barbee, S. A. *et al.* Staufen- and FMRP-Containing Neuronal RNPs Are

Structurally and Functionally Related to Somatic P Bodies. *Neuron* **52**, 997–1009

(2006).

65. Kerviel, A., Zhang, M. & Altan-Bonnet, N. A New Infectious Unit: Extracellular

Vesicles Carrying Virus Populations. *Annu. Rev. Cell Dev. Biol.* **37**, 171–197

(2021).

66. Diosa-Toro, M., Strilets, T., Yeh, S.-C. & Garcia-Blanco, M. A. Tinkering with

extracellular vesicles viruses evolve new infectious units. *ExRNA* **4**, (2022).

67. Rey-Cadilhac, F., Rachenne, F., Missé, D. & Pompon, J. Viral Components

Trafficking with(in) Extracellular Vesicles. *Viruses* **15**, 2333 (2023).

68. Xiong, J. *et al.* Extracellular vesicles promote the infection and pathogenicity of

Japanese encephalitis virus. *J. Extracell. Vesicles* **14**, e70033 (2025).

69. Rehwinkel, J. & Gack, M. U. RIG-I-like receptors: their regulation and roles in

RNA sensing. *Nat. Rev. Immunol.* **20**, 537–551 (2020).

70. Fredericksen, B. L., Keller, B. C., Fornek, J., Katze, M. G. & Gale, M.

Establishment and maintenance of the innate antiviral response to West Nile

Virus involves both RIG-I and MDA5 signaling through IPS-1. *J. Virol.* **82**, 609–

616 (2008).

71. Loo, Y.-M. *et al.* Distinct RIG-I and MDA5 signaling by RNA viruses in innate

immunity. *J. Virol.* **82**, 335–345 (2008).

72. Nasirudeen, A. M. A. *et al.* RIG-I, MDA5 and TLR3 synergistically play an

important role in restriction of dengue virus infection. *PLoS Negl. Trop. Dis.* **5**,

e926 (2011).

73. Sprokholt, J. K. *et al.* RIG-I–like Receptor Triggering by Dengue Virus Drives

Dendritic Cell Immune Activation and TH1 Differentiation. *J. Immunol.* **198**, 4764–

4771 (2017).

74. Hamel, R. *et al.* Biology of Zika Virus Infection in Human Skin Cells. *J. Virol.* **89**,

8880–8896 (2015).

75. Schuessler, A. *et al.* West Nile Virus noncoding subgenomic RNA contributes to

viral evasion of the Type I Interferon-mediated antiviral response. *J. Virol.* **86**,

5708–5718 (2012).

76. Slonchak, A. *et al.* Zika virus noncoding RNA cooperates with the viral protein

NS5 to inhibit STAT1 phosphorylation and facilitate viral pathogenesis. *Sci. Adv.*

8, eadd8095 (2022).

77. Bidet, K., Dadlani, D. & Garcia-Blanco, M. A. G3BP1, G3BP2 and CAPRIN1 are

required for translation of interferon stimulated mRNAs and are targeted by a

dengue virus non-coding RNA. *PLoS Pathog.* **10**, e1004242 (2014).

78. Wang, Z. *et al.* A mosquito salivary protein-driven influx of myeloid cells facilitates

flavivirus transmission. *EMBO J.* 1–32 (2024) doi:10.1038/s44318-024-00056-x.

79. Pingen, M. *et al.* Host Inflammatory Response to Mosquito Bites Enhances the

Severity of Arbovirus Infection. *Immunity* **44**, 1455–1469 (2016).

80. Fauquet, C. M. TAXONOMY, CLASSIFICATION AND NOMENCLATURE OF

VIRUSES. *Encyclopedia of Virology* 1730–1756 Preprint at

https://doi.org/10.1006/rwvi.1999.0277 (1999).

81. Cardoso, J. da C. *et al.* Yellow fever virus in Haemagogus leucocelaenus and

Aedes serratus mosquitoes, southern Brazil, 2008. *Emerg. Infect. Dis.* **16**, 1918–

1924 (2010).

82. Pijlman, G. P. *et al.* A Highly Structured, Nuclease-Resistant, Noncoding RNA

Produced by Flaviviruses Is Required for Pathogenicity. *Cell Host Microbe* **4**,

579–591 (2008).

83. MacFadden, A. *et al.* Mechanism and structural diversity of exoribonuclease-

resistant RNA structures in flaviviral RNAs. *Nat. Commun.* **9**, 119 (2018).

84. Théry, C. *et al.* Minimal information for studies of extracellular vesicles 2018

(MISEV2018): a position statement of the International Society for Extracellular

Vesicles and update of the MISEV2014 guidelines. *J. Extracell. Vesicles* **7**,

(2018).

85. Malkinson, M. *et al.* Virus encephalomyelitis of geese: some properties of the viral

isolate. *Isr. J. Vet. Med.* **53**, 44 (1998).

86. Cécile, B. *et al.* Complete Coding Sequence of Zika Virus from a French

Polynesia Outbreak in 2013. *Genome Announc.* **2**, 10.1128/genomea.00500-14

(2014).

87. Georghiou, G. P., Metcalf, R. L. & Gidden, F. E. Carbamate-resistance in

mosquitos. Selection of Culex pipiens fatigans Wiedemann (=C. quinquefasciatus

Say) for resistance to Baygon. *Bull. World Health Organ.* **35**, 691–708 (1966).

88. Kuno, G. Early History of Laboratory Breeding of Aedes aegypti (Diptera:

Culicidae) Focusing on the Origins and Use of Selected Strains. *J. Med. Entomol.*

47, 957–971 (2010).

89. Mainan, A., Kundu, R., Singh, R. K. & Roy, S. Magnesium Regulates RNA Ring

Dynamics and Folding in Subgenomic Flaviviral RNA. *J. Phys. Chem. B* (2024)

doi:10.1021/acs.jpcb.4c03981.

90. Forootan, A. *et al.* Methods to determine limit of detection and limit of

quantification in quantitative real-time PCR (qPCR). *Biomol. Detect. Quantif.* **12**,

1–6 (2017).

91. Graham, J. B., Swarts, J. L. & Lund, J. M. A Mouse Model of West Nile Virus

Infection. *Curr. Protoc. Mouse Biol.* **7**, 221–235 (2017).

92. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects

Models Using lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

93. Brooks, M. E. *et al.* glmmTMB Balances Speed and Flexibility Among Packages

for Zero-inflated Generalized Linear Mixed Modeling. *R J.* **9**, 378–400 (2017).

94. Lenth, R. emmeans: Estimated Marginal Means, aka Least-Squares Means.

(2024).

Fig. 1 | sfRNA is secreted in saliva from WNV-infected *Culex* and ZIKV-infected

***Aedes* mosquitoes. a** Detection of WNV sfRNA in orally infected *Culex* mosquitoes

by Northern blot. MM, molecular marker; Ctrl., *in vitro* transcribed folded WNV

sfRNA1; Inf., WNV-infected *Culex* mosquitoes; Uninf., mock-infected *Culex*

mosquitoes. **b-d** Quantification of gRNA (b), sfRNA (c), and the ratio of sfRNA:gRNA

(d) in salivary glands (SG) and saliva from *Culex* mosquitoes orally infected with

WNV. **e-f** Correlations between gRNA and sfRNA quantity in SG (e) and saliva (f)

from *Culex* orally infected with WNV. **g-i** Quantification of gRNA (g), sfRNA (h), and

the ratio of sfRNA:gRNA (i) in SG and saliva from *Culex* mosquitoes micro-injected

with WNV. **j-l** Quantification of gRNA (j), sfRNA (k), and the ratio of sfRNA:gRNA (l)

in SG and saliva from *Aedes* mosquitoes micro-injected with ZIKV. Each point

represents one pair of SG or saliva collected from one mosquito. b, c, g, h, j and k

Lines show geometric means ± 95% C.I. d, i and l Lines show arithmetic means ±

s.e.m. Dotted lines indicate LoD.

Fig. 2 | sfRNA in WNV-infected *Culex* saliva and ZIKV-infected *Aedes* saliva is protected from nuclease degradation by a detergent sensitive layer. a sfRNA inside a lipid bilayer is sensitized to nuclease degradation by Triton X-100 treatment. **b-d** Sensitivity to triton X-100 and Micrococcal nuclease (MNase) treatment for *in vitro*-transcribed WNV sfRNA diluted in uninfected *Culex* saliva (b), for salivary WNV sfRNA (c) and for salivary WNV gRNA (d). Saliva was collected from WNV-inoculated *Culex* mosquitoes. **e** Correlation between sfRNA and gRNA quantities in WNV-infected *Culex* saliva after triton X-100 and MNase treatments. **f-h** Sensitivity to triton X-100 and RNase A/T1 treatment for *in vitro*-transcribed ZIKV sfRNA diluted in uninfected *Aedes* saliva (f), for salivary ZIKV sfRNA (g) and for salivary ZIKV gRNA

(h). Saliva was collected from ZIKV-inoculated *Aedes* mosquitoes. **i** Correlation

between sfRNA and gRNA quantities in ZIKV-infected *Aedes* saliva after triton X-100

and RNase A/T1 treatments. Boxplots indicate median ± lower and higher quartiles.

Dots indicate saliva pool repeats. The dot colors indicate the different pools of saliva.

u, uninfected saliva; i, infected saliva from the corresponding mosquito. *, $p < 0.05$;

**, $p < 0.01$; ***, $p < 0.001$ according to T-test.

Fig. 3 | SfRNA in saliva increases WNV transmission. a Pools of saliva were
collected, categorized according to sfRNA:gRNA ratio and used to inoculate cells or
human skin explants. **b-c** WNV infection in Huh7 cells at 24 (b) and 48 hpi (c) with
low (L), moderate (M) and high (H) concentration of salivary sfRNA. N per category,
2. **d** Correlation between infection intensity (i.e., WNV gRNA fold change) in Huh7
and sfRNA concentration (i.e., sfRNA:gRNA ratio) in the corresponding saliva used
for infection. **e** WNV infection in skin explants at 24 h post injection with low and high
salivary sfRNA concentration. **f** Correlation between infection intensity in skin
explants and sfRNA concentration in the corresponding saliva used for infection. **g**
Cells and skin explants were transfected and injected, respectively, with sfRNA or
Control RNA (Ctl.) prior WNV infection. **h-j** WNV infection in Huh7 (h), HFF1 (i) and
U937 (j) cells at 24 and 48 hpi with WNV post sfRNA transfection. **k** WNV infection in
skin explants at 24 hpi with WNV post sfRNA injection. **l** Mice were intradermally co-
injected with WNV and sfRNA or Ctl. RNA. SfRNA alone was injected as control. Skin
biopsies were collected at day 1 and clinical signs and survival were recorded daily.
N, 14. **m** WNV infection in mouse skin at 24 h post injection. **n-o** Clinical signs (n)
and survival (o) for mice co-injected with WNV and either sfRNA or Ctl. RNA. b, c, e
Bars show mean ± s.e.m. h-j, k, m Bars show geometric mean ± 95% C.I. n Lines
show mean ± s.e.m. Dots indicate repeats. Different letters show significant
differences and *, $p < 0.05$; **, $p < 0.01$ according to post hoc Fisher's LSD test, T-
test or general linear mixed model. h-j, Different letters indicate statistical differences
according to Fisher's LDS test.

1021

1022

Fig. 4 | Salivary sfRNA inhibits IFN responses. a-d Expression of IFN-β (a), CXCL10 (b), MX1 (c), and IFI6 (d) in Huh7 cells at 24 and 48 h post infection (hpi)

with WNV-infected *Culex* saliva containing different concentrations of sfRNA.

Category of sfRNA concentration; L, low; M, moderate; and H, high. N per category,

2. The dotted line indicates the mock-infected values. **e-h** Expression of IFN-β (e),

CXCL10 (f), MX1 (g), and IFI6 (h) in Huh7 cells at 24 and 48 hpi with WNV post

sfRNA transfection. **i-l** Expression of IFN-β (i), CXCL10 (j), MX1 (k), and IFI6 (l) in

HFF1 cells at 24 and 48 hpi with WNV post sfRNA transfection. **m-p** Expression of

IFN-β (m), CXCL10 (n), MX1 (o), and IFI6 (p) in U937 cells at 24 and 48 hpi with

WNV post sfRNA transfection. **q-t** Expression of IFN-β (q), CXCL10 (r), MX1 (s), and

IFI6 (t) in skin explants at 24 hpi with WNV post sfRNA injection. Ctl., RNA control.

Bars show mean ± sem. Repeats are represented by dots. Different letters show

significant differences according to post hoc Fisher's LSD test or T-test.

1037

1038

Fig. 5 | sfRNA alters MDA-5 antiviral interferon response. a Scheme of RIG-I and

MDA5 mediated antiviral interferon response. The methods used to disrupt the

antiviral response are written in red. **b-d** WNV gRNA fold change post sfRNA

transfection at 24 and 48 hpi in Huh7 cells treated with MRT67037 (b), in RIG-I-

deficient Huh7.5 cells (c), and in Huh7.5 cells upon MDA5 silencing (d). Ctl., RNA

control. Bars show geometric mean ± 95% C.I. Repeats are indicated by dots.

Different letters show significant differences according to post hoc Fisher's LSD test.

1046

1047

1048

1049