Title: Exploiting Endogenous Cas9-Based Genome Editing of *Lacticaseibacillus*
***rhamnosus* GG (LGG) for Precision Food Fermentation and Live Biotherapeutics**

Authors: Zifan Xie[1], Yong-Su Jin[1,2]*, and Michael J. Miller[1,2]*

[1]Food Science and Human Nutrition, University of Illinois Urbana-Champaign, Urbana,
IL, USA

[2] Carl R. Woese Institute for Genomic Biology, University of Illinois Urbana-Champaign,
Urbana, IL, USA

*Corresponding author:

Yong-Su Jin (ysjin@illinois.edu) and Michael J. Miller (mille216@illinois.edu)

Abstract

Lacticaseibacillus rhamnosus GG (LGG) is one of the most studied probiotic strains and

is widely used in both food and therapeutic industries. With its safe and health-beneficial

properties, LGG is an ideal candidate for genome modification to enhance its

functionalities for food, industrial and pharmaceutical applications. However, precise and

efficient genome engineering tools for LGG are unavailable. In this study, we developed

a novel genome engineering tool for LGG based on its endogenous type II-A CRISPR-

Cas9 system. By employing a validated protospacer adjacent motif (PAM), a customized

single guide RNA (sgRNA) expression cassette and a homologous repair template, we

successfully reprogrammed the native CRISPR-Cas9 system for targeted genome

modifications. Using this method, we successfully constructed a fucose-negative LGG

incapable of growing on L-fucose and a lactose-positive LGG. The lactose-positive LGG

exhibited significantly improved growth in milk compared to its parental strain LGG,

which is unable to ferment lactose, attributed to its ability to metabolize lactose

efficiently. Moreover, this strain showed robust growth during yogurt fermentation and

maintained viability during cold storage for at least three weeks. These findings highlight

the potential of the lactose-positive LGG as an improved culture for dairy industry and a

functional probiotic for food and therapeutic applications.

Importance A precise and efficient genome editing tool is essential for unlocking the full

potential of *Lacticaseibacillus rhamnosus* GG (LGG), the most extensively studied

probiotic strain. This study established the first high-efficiency genome editing platform

for LGG by harnessing the endogenous type II-A CRISPR-Cas9 system. Using this tool,

we achieved targeted gene modifications, without introducing scars, heterologous DNA,

or antibiotic resistance genes. The engineered lactose-positive LGG strain demonstrated

improved growth in lactose-containing dairy products and robust performance during

yogurt fermentation, highlighting its industrial and therapeutic value. This platform not

only advances the development of enhanced probiotic strains but also establishes LGG as

a versatile vehicle for synthetic biology and biotechnological applications.

Introduction

Lactic acid bacteria (LAB) are widely used in various industrial fields as starter cultures and cell factories for producing fermented foods and value-added compounds (1, 2). Their primary application is as starter cultures in fermented foods, particularly dairy products like yogurt and fermented milk, which serve as key vehicles for delivering probiotics. Beyond their crucial role in food production, numerous LAB strains have been investigated in clinical trials for their nutritional and therapeutic benefits, including pathogen protection, immune modulation, antioxidant effects, and improved nutrient bioavailability (3–6). These well-documented health benefits, along with their generally recognized as safe (GRAS) status, make LAB ideal candidates for genome modification to enhance their desirable properties for both food and therapeutic applications (7, 8). Recently, many novel genome modification approaches have been introduced and applied to LAB strains to improve their beneficial properties and to determine relevant phenotypes (9).

The CRISPR-Cas (clustered regularly interspaced short palindromic repeats and CRISPR-associated proteins) system, originally discovered as an adaptive immune mechanism in bacteria, has emerged as a powerful tool for precise DNA modification in various organisms (10, 11). The heterologous CRISPR-Cas system has been successfully employed in many lactobacilli, such as *Limosilactobacillus reuteri*, *Lactiplantibacillus plantarum*, *Lacticaseibacillus casei*, *Lactobacillus acidophilus*, *Lactobacillus gasseri*, and *Lacticaseibacillus paracasei* (12–15). However, despite these successes, heterologous CRISPR systems face challenges such as low transformation efficiency, suboptimal activity of heterologous Cas proteins, and potential toxicity to host cells (16, 17) . To address these problems, researchers have exploited endogenous CRISPR-Cas systems present in some lactobacilli to achieve genome modifications. These native systems offer a more compatible genome-editing platform, providing higher precision and lower toxicity. For instance, the endogenous subtype I-E CRISPR-Cas system has been utilized for efficient genetic manipulation of *Lactobacillus crispatus* (18), while the subtype II-A CRISPR-Cas9 system has been used to achieve genome engineering in *Lcb. paracaseri* (19).

Lacticaseibacillus rhamnosus GG (LGG) is the world's most documented
probiotic strain, known for its health benefits and safety across all life stages, including
its ability to modulate the gut microbiome, promote oral health, and support the immune
system (20–23). Despite over 300 publications of human studies and well-established
health benefits, no precise genome editing tool has been developed for LGG, limiting the
ability to (1) elucidate the underlying mechanisms of LGG's probiotic effects and to (2)
exploit the long history of safety and efficacy of LGG by engineering it for therapeutic
applications.

LGG is commonly used in dietary supplements and dairy products to deliver these
health benefits. However, its inability to ferment lactose (24, 25), the primary
carbohydrate in milk, restricts its growth in milk and other dairy products, which are key
probiotic delivery systems. Developing a precise genome editing tool could offer a
potential solution to this challenge by restoring the ability of LGG to metabolize lactose,
thereby expanding LGG applications in the dairy industry and expanding its role as a
functional probiotic and a biotherapeutic vehicle.

In this study, we developed a novel genome editing tool in LGG by harnessing the
endogenous CRISPR-Cas9 system (Figure 1) and applied it for generating fucose-
negative LGG and lactose-positive LGG variants. The lactose-positive LGG
demonstrated the ability to metabolize lactose effectively, improving its growth in milk
and yogurt fermentation conditions. This advancement represents a significant step
toward creating functional probiotics and expanding LGG's applications in food and
therapeutic contexts.

Results

Verification of the Activity of the Endogenous CRISPR-Cas9 System in LGG

LGG contains a complete type II-A CRISPR-Cas9 system (Figure 2A) (26). This
system is composed of 4 Cas protein-encoding genes including *cas1*, *cas2*, *csn2*, and the
critical *cas9*. It also contains a CRISPR array with 24 different 30-base pair (bp) spacers,
each flanked by a 36-bp direct repeat sequence (Figure 2A). A protospacer adjacent motif

(PAM) is essential for target DNA recognition and binding (27, 28). Analysis of the

spacers and their corresponding protospacers predicted a PAM sequence, 5'-NGAAA-3'

(Figure 2B).

To assess the activity of the native CRISPR-Cas9 system in LGG, we conducted a

plasmid interference assay. An active CRISPR system can prevent the uptake of plasmids

carrying sequences complementary to native CRISPR spacers (Figure 2C). For this assay,

we designed interference plasmids (Table 1) containing a protospacer matching the first

spacer (the most recently acquired spacer) in the CRISPR array, either with or without an

adjacent PAM sequence (5'-NGAAA-3') (Figure 2D). Our results showed that the native

CRIPSR-Cas9 system effectively interfered with plasmids containing functional PAM

sequences (Figure 2E). Specifically, transformation efficiency reduced by 10^3-fold and

over 10^4-fold for plasmids with PAM sequences of 5'-GGAAA-3' and 5'-TGAAA-3',

respectively (Figure 2E). This demonstrated the high activity of the native CRISPR-Cas9

system. To further evaluate the PAM specificity, we constructed a new interference

plasmid, pPS1TC (Table 1), in which the second conserved nucleotide of the PAM

sequence was mutated from G to C (5'-TCAAA-3') (Figure 2D). As presented in Figure

2E, pPS1TC did not trigger the endogenous system to produce a detectable interference

response, reflecting the high specificity of the type II-A CRISPR-Cas9 system for its

PAM sequence. Overall, these findings validated the predicted 5'-NGAAA-3' as the

functional PAM and confirmed the robust activity and high specificity of the native

CRISPR-Cas9 system under standard laboratory conditions.

Reprogramming Endogenous CRISPR-Cas9 System for Gene Deletion

After confirming the activity of the native CRISPR-Cas9 system with the

predicted PAM sequence, we then explored its potential for genome editing in LGG.

Initially, we designed an editing plasmid with a homologous repair template and a

CRISPR RNA (crRNA) expression cassette controlled by the native leader sequence of

the CRISPR array. However, due to the inherent instability of rolling circle plasmids (29),

the space between the repeats was easily looped out from the plasmid during plasmid

replication, resulting in spacer-deficient plasmids and many wild-type (WT) survivors. To

address this, we shifted to using single-guide RNA (sgRNA) to target a specific sequence

in the genome of LGG.

Based on the predicted structure of the crRNA/ trans-activating CRISPR

RNA (tracrRNA) duplex and previous transcriptional sequencing results (26), we linked

crRNA and tracrRNA with an artificial AAUC loop to form an sgRNA (Figure 2F). As a

result, we constructed a targeting plasmid with a sgRNA expression cassette under the

strong constitutive promoter $Ppgm$ (*L. acidophilus* NCFM) and an editing plasmid

containing a sgRNA expression cassette and a homologous repair template (HR1 and

HR2) with the desired mutations for genome editing (Figure 2G). With this design, we

proposed to achieve genome editing in LGG by reprogramming the endogenous CRISPR

system.

As a test, we first attempted to delete L-fucose isomerase, *fucI* (LGG_02685)

(Figure 3A), in LGG, as resulting mutants are expected to display a significantly lower

growth rate compared to the WT when fucose is used as the sole carbon source (30). A

30-bp spacer sequence complementary to *fucI* (Table 3) and two 1-kb homologous

regions (HR1 and HR2) were cloned into the plasmid pTRK870 to generate an editing

plasmid pfucI-sgRNA-RT (Table 1). Apart from this, pfucI-RT (repair template only) and

pfucI-sgRNA (sgRNA only) were also constructed as control plasmids (Table 1). Results

showed that no clone was obtained after the transformation of LGG with pfucI-sgRNA,

which confirmed the lethality of Cas9-mediated double-stranded DNA breaks when a

self-targeting sgRNA is introduced to repurpose the endogenous system and trigger lethal

cleavage (Figure 3B). When pfucI-RT was transformed into LGG, hundreds of

transformants were obtained (Figure 3B) and no sequenced clones displayed the deletion

genotype. In contrast, when pfucI-sgRNA-RT was transformed into LGG, substantially

fewer colonies (4 colonies) were observed. These four transformants were examined by

PCR, with one producing the desired 2.1-kb PCR product indicating a deletion, while the

other three displayed a WT genotype (2.7-kb PCR product) (Figure 3C). The 2.1-kb

amplicon was sequenced to confirm the mutant, which was subsequently named MJM568

(Figure 3D). Furthermore, phenotypic analysis revealed that the MJM568 strain could not

utilize fucose as the sole sugar source (Figure 3E). These results confirm that the

endogenous CRISPR system in LGG can be reprogrammed to achieve gene deletion, and

both the sgRNA-guided Cas9 targeting and the repair template are crucial for generating

the intended mutation.

Construction of Lactose-Positive LGG

After demonstrating the possibility of genome editing with the native CRISPR-

Cas9 system, we attempted to construct a lactose-positive LGG strain using genome

editing. The ability to metabolize lactose is an important feature commonly exploited in

dairy fermentation. However, this capability is lost with LGG. Compared with the closely

related lactose-positive strain *Lcb. rhamnosus* Lc705, LGG carries the genes *lacTEGF*

operon for the lactose phosphotransferase system, but two mutations in the operon make

LGG lactose-negative (incapable of assimilating lactose): a 4-bp deletion causing a

frameshift in the antiterminator gene (*lacT*, LGG_00648) and a point mutation (Q231*)

in the 6-phospho-β-galactosidase gene (*lacG*, LGG_00650) (Figure 4A) (24, 31). To

restore the function of the *lacTEGF* operon, both mutations need to be corrected.

A previous study demonstrated that disrupting the stem-loop structure of the *t1*

transcriptional terminator can result in unimpeded transcription of the *lacTEGF* operon

(31). Based on this, we designed and constructed an editing plasmid placT-sgRNA-RT to

create a 20-bp deletion, thus inactivating the transcriptional terminator t1 (Figure 4C).

Sequencing of the edited region in five transformants revealed that one colony carried the

desired 20-bp deletion mutation. This mutant was designated MJM569 (Figure 4B and

4C).

Next, 10^8 cells of the WT and MJM569 strains were plated on MRS agar plates

containing 2% lactose as the sole sugar source and incubated anaerobically for 2 days at

37°C. After the incubation, approximately 40 colonies per plate were observed from the

plating of the MJM569 strain (Figure 4D). In contrast, no colonies were observed from

the plating of the WT LGG strain (Figure 4D), likely due to the chances of obtaining cells

with simultaneous corrections in both *lacT* and *lacG* were extremely low. Next, fifteen

lactose-positive colonies derived from the MJM569 strain were randomly picked and

tested in MRS broth containing 2% glucose or lactose using a multi-mode microplate

reader. The results showed that all 15 colonies were lactose-positive and exhibited robust

growth in both glucose and lactose conditions (Figure 4E). These lactose-positive isolates

obtained from this approach were likely to be revertants in *lacG*. To verify this, the *lacG*

region of these 15 lactose-positive isolates was amplified by PCR and sequenced. The

analysis revealed a point mutation at amino acid 231 of *lacG* in each strain, reverting the

premature stop codon in *lacG* to a functional codon (Figure 4F). No other mutations were

observed in the *lacG* gene. Specifically, in strain L12 and L13, the thymine at position

691 in *lacG* mutated to a cytosine, reverting the codon at this position from TAG (stop)

back to CAG (glutamine, *232Q), as observed in *Lcb. rhamnosus* Lc705 (Figure 4F).

To prepare strain L13 for continued genome editing and food applications, we

passed it twice in MRS broth without selective pressure, allowing these cells to naturally

lose the plasmid placT-sgRNA-RT. The resulting strain, named MJM570, was confirmed

to have lost the plasmid, as it could not grow under erythromycin selection. Its growth

curves in both glucose and lactose conditions are shown in Figure 4G.

Growth of LGG and MJM570 Strains in Milk

Since MJM570 could utilize lactose, we wanted to determine whether this trait

would improve its growth in milk compared to that of WT LGG strain. As shown in

Figure 5A, the lactose-positive MJM570 exhibited robust growth in milk, whereas the

WT LGG strain could not grow under the same condition. Notably, after 24 hours of

incubation, the milk inoculated with MJM570 showed coagulation (Figure 5B),

indicating a pH drop as a result of lactic acid production (Figure 5C). In contrast, the milk

inoculated with the WT LGG strain remained in liquid form after 24 hours at 37°C

(Figure 5B). In summary, the new ability to metabolize lactose improved the growth of

MJM570 in milk.

Viability of Lactose-Positive LGG in Yogurt During Refrigerated Storage

To evaluate the growth and fermentation properties of the WT LGG and MJM570

strains in yogurt, we monitored pH levels and cell counts throughout both the

fermentation process and refrigerated storage (5 ± 1 °C, 21 days). The MJM570 strain

was compared with the WT LGG strain and a PBS control (Figure 6A).

As shown in Figure 6B, the initial pH of the milk mixture with a yogurt starter culture (*Streptococcus thermophilus* and *Lactobacillus delbureckii* subsp. *bulgaricus*) was approximately 6.5, with no significant differences between the groups (*p > 0.05*). After about 5 hours of fermentation, the pH reached the desired range of 4.3-4.4 due to lactose fermentation by *S. thermophilus* and *L. bulgaricus*. Although pH changes were not statistically significant, differences in the growth of LGG strains were observed. As shown in Figure 6B, the initial cell counts of the WT LGG strain and MJM570 in yogurts were approximately 1×10^8 CFU/g (colony-forming units per gram of the yogurt sample) and 5×10^7 CFU/g, respectively. During the fermentation, MJM570 exhibited robust growth, with cell counts significantly increasing throughout the fermentation period (*p < 0.05*), reaching about 8×10^8 CFU/g by the end. In contrast, the WT LGG strain didn't show significant growth (*p > 0.05*). By the end of fermentation, MJM570 had significantly higher cell counts than the WT LGG strain (*p < 0.05*), highlighting its superior ability to proliferate in a lactose-containing environment with lactose-fermenting strains *S. thermophilus* and *L. bulgaricus*.

After yogurt fermentation, pH levels and cell density were monitored for all three yogurt groups during 21 days of refrigerated storage. As shown in Figure 6C, all three groups experienced a significant pH drop during the first week of cold storage (*p < 0.05*). This decrease was likely due to the continued metabolic activity of the yogurt starter culture and LGG strains, which continue to break down carbohydrates into organic acids, resulting in a lower pH. After the first week, pH levels stabilized across all groups and remained consistent throughout the rest of the storage period (Figure 6C). Regarding sustained viability of LGG and MJM570 during cold storage (Figure 6C), both strains were viable during cold storage for at least 21 days, likely due to the inherent acid tolerance of LGG (32). Notably, MJM570 consistently exhibited significantly higher cell densities than the WT LGG strain throughout the storage period (*p < 0.05*), reflecting its enhanced performance in a fermented dairy product.

Discussion

LGG is the world's most documented probiotic strain and has been extensively

used in dietary supplements and fermented foods. However, the development of enhanced

probiotic LGG strains has been limited by the lack of precise and efficient genome

editing tools. Traditional homologous recombination has been the only established

genome modification method in LGG (33, 34). However, this method is time-consuming

and inefficient, requiring multiple rounds of subculture and screening, and relies on

antibiotic resistance genes for selection (9). Recently, CRISPR-Cas9 gene-editing

systems have been developed for engineering serval lactobacilli strains (35), with

endogenous CRIPSR-Cas9-based genome editing showing particularly high efficiency.

One major challenge in developing an endogenous CRISPR system for LGG has been the

lack of a replicable cloning vector that can stably replicate in LGG. To address this, we

identified and characterized the plasmid pTRK830 (29), which can replicate stably in

LGG and effectively overcomes this limitation.

Building on this foundation, in this study, we successfully demonstrated the

potential of using the endogenous type II-A CRISPR-Cas9 system for precise genome

editing in LGG. This native system not only enhanced the precise recognition of target

location but also effectively activated the native homologous recombination machinery to

yield the desired genetic modifications. This study represents the first successful

establishment of a highly efficient and convenient native CRISPR-Cas gene-editing

system in LGG. By harnessing the native Cas9 protein, we achieved targeted gene

deletions with high precision. Importantly, the genetically modified LGG strains

generated through this method do not contain any scars, heterologous DNA, or antibiotic-

resistant genes, making them suitable for food and therapeutic applications.

As a demonstration of this CRISPR-Cas9 editing tool's potential, we developed

the lactose-positive strain MJM570 by deleting a transcriptional terminator upstream of

the *lacT* gene and correcting a point mutation in the *lacG* gene. This lactose-positive

strain can offer numerous advantages in both industrial and therapeutic applications. The

lactose metabolism activity of MJM570 makes it highly suitable for use in a variety of

lactose-containing dairy products such as milk, yogurt, and cheese, where it can grow to

higher densities by utilizing the lactose in these environments. This trait can also make

MJM570 an ideal starter culture for the dairy industry. In yogurt production, MJM570

exhibited robust growth during fermentation and stably maintained during cold storage,

consistently achieving higher cell density compared to WT LGG, making it an ideal

candidate for dairy product formulations. In addition, compared to the commonly used

lactose-positive industrial strain *Lcb. rhamnosus* Lc705 (24), our new probiotic strain

MJM570 not only exhibits lactose metabolism activity but also presumably retains the

numerous health benefits of its parental strain LGG. Furthermore, MJM570 shows

potential therapeutic applications for addressing lactose intolerance, a common condition

where undigested lactose causes gastrointestinal discomfort, such as bloating and

diarrhea (36, 37). Dairy products or dietary supplements containing MJM570 may have

the ability to alleviate symptoms by digesting lactose *in situ*, reducing colonic

fermentation and preventing related symptoms. Therefore, MJM570 not only contributes

to better lactose tolerance but also retains the broader probiotic benefits associated with

its parental strain LGG, making it a valuable strain for both dairy production and health

applications.

In conclusion, this study establishes the first endogenous CRISPR-Cas system for

genome editing in LGG, significantly enhancing the available toolkit for probiotic

engineering. Our success in developing the lactose-positive strain MJM570 demonstrates

the utility of this system for creating novel probiotic strains with potential enhanced

industrial and therapeutic applications. In the future, this precise genome editing tool can

also hold promise for applications in heterologous gene expression, further broadening

LGG's role as a live therapeutic vehicle. This advancement paves the way for future

research and applications, offering a powerful tool for genetic manipulation in LGG and

other *Lactobacillus* species to improve probiotic functionalities and industrial use.

Material and Methods

Strains and Culture Conditions

LGG and its mutants (Table 1) were grown in de-Man-Rogosa-Sharpe (MRS;

Hardy Diagnostics, Santa Maria, CA, USA) broth or on 1.5% agar plates at 37 °C under

anaerobic conditions (90% N_2, 5% CO_2, and 5% H_2). *Escherichia coli* MC1061 was

cultured in Luria-Bertani (LB) medium at 37 °C with vigorous shaking. Antibiotic was

added to the growth medium when applicable: erythromycin (Em; Gold Biotechnology,

St. Louis, MO, USA) 5 µg/ml for LGG and 150 µg/ml for *E. coli*.

CRISPR-Cas System Characterization *in silico*

The LGG genome is available at the National Center for Biotechnology

Information (NCBI). The CRISPR array and the Cas proteins were identified using

CRISPRCasFinder web server (38). Bioinformatic analyses were performed to predict the

PAM sequence based on spacer–protospacer match, using the spacers of LGG and the

web server CRISPRTarget (39). The WebLogo server was used to represent the PAM

sequence based on a frequency chart where the height of each nucleotide represents the

conservation of that nucleotide at each position (40). In type II systems, the tracrRNA has

a complementary region to the CRISPR repeat sequence of the crRNA. Therefore, the

repeat sequence was used to identify the tracrRNA in the CRISPR locus using the Basic

Local Alignment Search Tool (BLAST) (41), and the interaction between crRNA and

tracrRNA was then predicted using the NCPACK web server and depicted manually (42).

DNA Manipulations and Molecular Cloning

A Promega Wizard genomic DNA purification kit (Promega Corporation,

Madison, WI, USA) was used to extract genomic DNA. Plasmid DNA from *E. coli* cells

was extracted using the QIAGEN plasmid mini-prep kit (QIAGEN, Germantown, MD,

USA). DNA restriction enzymes and T4 DNA ligase were purchased from New England

Biolabs (NEB, Ipswich, MA, USA) and used following the instructions of the

manufacturers. All polymerase chain reactions (PCRs) were conducted according to

standard procedures using DreamTaq DNA polymerase (Thermo Fisher Scientific,

Waltham, MA, USA) or Phusion DNA polymerase (NEB, Ipswich, MA, USA). PCR

primers were synthesized by Integrated DNA Technologies (IDT, Coralville, IA, USA)

and gene fragments were synthesized by Twist Bioscience (Twist Bioscience, San

Francisco, CA, USA). PCR products were electrophoresed in 1.0% agarose gel. DNA

fragments were purified using the QIAquick PCR purification kit (QIAGEN,

Germantown, MD, USA) before downstream application. DNA sequence analysis was

performed at the UIUC Core Sequencing Facility (Urbana, IL, USA) using the Sanger

sequencing method.

Construction of Interference Plasmids

pTRK830 is a replicating shuttle vector for *E. coli* and lactobacilli strains (29).

The multiple cloning site (MCS) from pBluescript KS (+) was ligated into the *StuI* site of

pTRK830, which generated pTRK870 (Table 1). Plasmid pTRK870 was used for all

following plasmid constructions. The interference plasmids were constructed as described

before (26). The double-stranded DNA protospacers, with or without the PAM, were

amplified by PCR, and then ligated into XbaI-HindIII-digested pTRK870 to construct

interference plasmids. The resulting interference plasmids were PCR-screened in *E. coli*

transformants using PS1 and PS2 primers for the presence of the insert and sequenced to

confirm sequence content.

Construction of Editing Plasmids

Each editing plasmid was commonly composed of a sgRNA expression cassette

and a repair template (Table 3). The sgRNA expression cassettes included a promoter

Ppgm of the phosphoglycerate mutase from *L. acidophilus* NCFM, a designed sgRNA

containing a 30-bp sequence targeting a specific genomic locus of LGG strain, and a

Rho-independent terminator. The sgRNA expression cassette sequence was synthesized

and amplified by primers sgRNA-XbaI and sgRNA-XmaI (Table 2) and then cloned into

XbaI-XmaI-digested pTRK870 to generate targeting plasmids, pfucI-sgRNA and placT-

sgRNA.

Next, the repair templates were amplified from the genomic DNA of LGG strain

or synthesized by Twist Bioscience. The repair template with two 1-kb homologous

regions for *fucI* gene editing was amplified from the genomic DNA of LGG strain. For

this, two 1-kb homologous regions were fused by overlap extension PCR to obtain the 2-

kb repair template to the *fucI* gene. This repair template was digested with SalI and ApaI,

then cloned into SalI-ApaI-digested plasmids pfucI-sgRNA and pTRK870 to generate the

plasmid pfucI-sgRNA-RT and pfucI-RT, respectively. For the transcriptional terminator

t1 deletion, the repair template was synthesized and cloned into the SalI-ApaI-digested placT-sgRNA generating the plasmid placT-sgRNA-RT.

All these final plasmid constructs were PCR-screened using PS1 and PS2 primers and sequenced to confirm sequence content.

Transformation of LGG

The transformation method was optimized based on previous study (43). 100 µl of an overnight culture was inoculated in 10 ml prewarmed MRS medium and was incubated without agitation at 37 °C. After overnight growth, 2 ml of the culture was inoculated into 100 ml of freshly prepared, prewarmed MRS medium and incubated at 37 °C until its optical density OD_{600} reached 0.5-0.6. Cells were harvested by centrifugation at 4000 rpm for 15 min. The cells were washed three times with cold 3.5× Sucrose Magnesium Electroporation Buffer (SMEB) containing 1 M sucrose, 3.5 mM $MgCl_2$, pH 7.0. After washing, the cells were resuspended in 1 ml of the same buffer, ready for electroporation. For each transformation, 1 µg of plasmid was combined with 100 µL of cells, and 2-mm cuvettes (Bio-Rad, Hercules, CA, USA) were used for transformation under 2.5-kV, 25-µF, and 200-Ω conditions. DNA concentrations of plasmids used for transformation were quantified using a spectrophotometer, NanoDrop2000c (Thermo Fisher Scientific, Waltham, MA, USA). Then, 900 µl pre-warmed MRS broth was added to the cuvette. The transformed cells were allowed to recover by incubation at 37 °C overnight. After recovery, the cells were plated on MRS agar plates with appropriate antibiotic and incubated at 37 °C in anaerobic chamber for 48 to 72 h.

The transformants obtained were PCR-screened and sequenced to confirm the presence of desired mutations (Table 2). For the fucose isomerase gene *fucI*, the primers fucI-check-F and fucI-check-R were used for the chromosomal PCR amplification and sequencing (2.7-kb in the WT and 2.1-kb in deletion mutant). For the transcriptional terminator t1, the primers lacT-check-F and lacT-check-R were used for the chromosomal PCR amplification and sequencing. For the 6-phospho-β-galactosidase *lacG*, the primers lacG-F and lacG-R were used for the chromosomal PCR amplification and sequencing.

Growth Curve Measurement

The growth of the LGG strain and its mutant *L. rhamnosus* MJM568 and MJM570 was analyzed in sugar-free MRS media (US Biological, Swampscott, MA, USA) containing 2% glucose, fucose, or lactose (Sigma, St. Louis, MO, USA). Strains were picked and cultured in MRS broth overnight. Then the cultures were harvested and washed three times with phosphate-buffer solution (PBS, pH 7.4) and resuspended in same volume of PBS buffer. Cells were inoculated 2% into MRS containing different sugar sources and transferred into a sterile 96-well plate. The plates were incubated at 37 °C without agitation for 24 h in FilterMax F5 Multi-Mode Microplate Reader (Molecular Devices, LLC, Sunnyvale, CA, USA). The absorbance at 595nm (Abs_{595nm}) in each well was recorded every 20 or 30 min. The maximum specific growth rate (μ_{max}) was defined as the slope of the tangent line at the inflection point of the logistic regression curve and was reported in hours^{-1}.

Growth in Milk

After two passages of LGG and MJM570 in MRS broth, the cultures were harvested and washed three times with PBS buffer and then resuspended in an equal volume of PBS buffer. The cells were inoculated 2% into 10 mL of milk. The milk, reconstituted to 10% (w/v) from skimmed milk powder (Sigma, St. Louis, MO, USA), was heated at 85 °C for 30 min, and then cooled to room temperature before inoculation. The inoculated milk samples were incubated at 37°C for 24 h, and samples were taken at 0, 2, 4, 6, 8, 10, 12, 14, 22, and 24 h to determine pH values and the number of colony-forming units per milliliter (CFU/mL) by plate counting.

Yogurt Fermentation

Prior to yogurt production, both LGG and MJM570 were inoculated and cultured anaerobically in MRS broth for 24 h at 37 °C for 2 consecutive passes. Freeze-dried cultures of *Streptococcus thermophilus and Lactobacillus delbureckii subsp. bulgaricus* (YF-L706, Chr. Hansen, Milwaukee, WI, USA) were stored at -80 °C and defrosted at room temperature for 10 min before use.

Milk was reconstituted (10% w/v) from skimmed milk powder (Sigma, St. Louis,

MO, USA) and heated at 85 °C for 30 min. After cooling to 42 °C, the milk was

inoculated with 0.02% yogurt starter culture. One milliliter (1 mL) of overnight cultures

of LGG and MJM570 were centrifuged at 3,900 rpm for 5 min. The supernatant was

discarded, and the pellets were washed twice with PBS buffer. Each pellet was

resuspended in 1 mL PBS buffer and added individually to the yogurt mixture to obtain a

final concentration of approximately 10^7 CFU/mL. For the control group, 1 mL of PBS

buffer was added to the yogurt mixture. There were 3 replications for each treatment.

The yogurt mixtures were fermented at 42 °C until the pH reached 4.3. During

yogurt fermentation, pH values of the yogurt samples were measured at room

temperature using a digital pH meter (Accumet AR15, Fisher Scientific, Pittsburgh, PA,

USA) once an hour for a total of 5 hours. To assess the growth of LGG and MJM570

during fermentation, yogurt samples were collected at 0, 2, 4, and 5 h to determine the

colony-forming units per gram (CFU/g) of the yogurt sample. Each sample was from a

previously undisturbed yogurt.

After yogurt fermentation, the samples were cooled and stored in a refrigerator

(4 °C) for 21 days. Samples were collected the day following fermentation (day 1), and

then once weekly for 3 weeks (day 7, 14, and 21) to assess the viability of LGG and

MJM570 as well as pH levels. Each sample was from an undisturbed portion of yogurt.

Viability of LGG and MJM570

One gram (1 g) of yogurt was suspended in 1 mL of PBS buffer, and then 10-fold

serial dilutions were performed. Selective enumeration of LGG and MJM570 was carried

out using MRS vancomycin plates, which contained 50 ppm vancomycin (Gold

Biotechnology, St. Louis, MO, USA) (44). The plates were incubated at 37 °C for 48

hours under anaerobic conditions. Plates were counted and recorded as CFU per gram of

the yogurt product (CFU/g).

Statistical Analysis

The results were presented as the means of independent experiments ± standard

deviations. Statistical significance was determined using a threshold of $p < 0.05$ using

unpaired t test and one-way ANOVA test. Distinct letters were employed to denote

significant differences in all figures. Plots were generated using GraphPad Prism

(GraphPad Software, Boston, MA, USA).

Acknowledgements

Zifan Xie was supported by the Investment for Growth Program from the University of

Illinois, Synthetic Biology for Food and Nutrition Innovation (SynFONI).

Tables and Figures

Table 1 Strains and plasmids used in this study

Strains or plasmids	Description	Sources or references
Strains		
E. coli MC1061	*E. coli* transformation host, Str[R].	Lab stock
Lcb. rhamnosus GG	*Lcb. rhamnosus* GG was originally isolated from fecal samples of a healthy human adult.	ATCC
Lcb. rhamnosus MJM568	*Lcb. rhamnosus* GG mutant with the deletion (640 bp) of the L-fucose isomerase *fucI* gene.	This study
Lcb. rhamnosus MJM569	*Lcb. rhamnosus* GG mutant with the partial deletion of the transcriptional terminator t1.	This study
Lcb. rhamnosus MJM570	*Lcb. rhamnosus* MJM569 mutant with the point mutation (*231Q) in 6-phospho-β-galactosidase *lacG* gene.	This study
Plasmids		
pTRK830	Low-copy number *E. coli* and Lactobacilli shuttle vector; Erm[R].	(29)
pTRK870	The multiple cloning site sequence cloned into pTRK830 at *StuI* site.	This study
pS1	Protospacer sequence of spacer 1 from LGG CRISPR array cloned into pTRK870 at *XbaI-HindIII* sites.	This study
pPS1TG	Protospacer sequence of spacer 1 and PAM sequence (5'-TGAAA-3') cloned into pTRK870 at *XbaI-HindIII* sites.	This study
pPS1GG	Protospacer sequence of spacer 1 and PAM sequence (5'-GGAAA-3') cloned into pTRK870 at *XbaI-HindIII* sites.	This study

pPS1TC	Protospacer sequence of spacer 1 and 5-bp sequence (5'-TCAAA-3') cloned into pTRK870 at *XbaI-HindIII* sites.	This study
pfucI-RT	Control plasmid containing the repair template for generating a knockout of the *fucI* gene, cloned into pTRK870 at *SalI-ApaI* sites.	This study
pfucI-sgRNA	Targeting plasmid for the *fucI* gene, constructed by cloning the *fucI* targeting sgRNA expression cassette gBlock into pTRK870 at *XbaI-XmaI* sites.	This study
pfucI-sgRNA-RT	Editing plasmid containing the repair template for generating a knockout of the *fucI* gene, cloned into pfucI-sgRNA at *SalI-ApaI* sites.	This study
placT-sgRNA	Targeting plasmid for the transcriptional terminator *t1*, constructed by cloning the *t1* targeting sgRNA expression cassette gBlock into pTRK870 at *XbaI-XmaI* sites.	This study
placT-sgRNA-RT	Editing plasmid containing the repair template for generating a deletion at the terminator *t1*, cloned into placT-sgRNA at *SalI-ApaI* sites.	This study

463 **Table 2 Primers used in this study**

Primer	Sequences (5'-3')
PS1	GCGAAAGGGGGATGTGCTGC
PS2	GCACCCCAGGCTTTACACTTTATGC
sgRNA-XbaI	CGTACCATCGTCTAGATGCGACAAGT
sgRNA-XmaI	CGATCCCGGGAAAAAAATGTATGGCC
Plasmid interference assay	
PS1XbaI	AGTCATCTAGAAAGAAACGATCCCGAGTTTCTGG
S1HindIII	AGTCAAAGCTTTAGTAAGCAATCCAGAAACTCGGGATC
PS1TGHindIII	AGTCAAAGCTTTAGTTTTCAAAGCAATCCAGAAACTCGGGATC
PS1GGHindIII	AGTCAAAGCTTTAGTTTTCCAAGCAATCCAGAAACTCGGGATC
PS1TCHindIII	AGTCAAAGCTTTAGTTTTGAAAGCAATCCAGAAACTCGGGATC
Construction of the editing plasmid for the deletion of *fucI* gene	
fucI-HR1-F	CAAATGCGCAAAGTCACCTGGAAC
fucI-HR1-R	CGCATCAAACCCGCCGTCAATTTTCTTCACAGTTCATCAAGCTTATTAG
fucI-HR2-F	TGACGGCGGGTTTGATGCGAAATAAGAGTTATC
fucI-HR2-R	CCCGGATCTGTTCGCTTGTTG
fucI-RT-SalI	TACCGTCGACCAAATGCGCAAAG
fucI-RT-ApaI	GTCAGGGCCCCCCGGATCTGTTC
fucI-check-F	GTCCCGTCAAAATTGGAACAAAGAATGTTG
fucI-check-R	CCCAGAGCTTTGTCGAGTCACC
Construction of the lactose positive LGG strain	
lacT-RT-SalI	GATACCGTCGACCTTCGAATCAACAG
lacT-RT-ApaI	GTACCGGGCCCTATTCGCC
lacT-check-F	GACAGGGCCTGACAGATGTAATTGG
lacT-check-R	CCATCACGGATTGCGGACACATAC

lacG-F	ATGCGTAAACAATTACCCAAGGACTTTGTAATC
lacG-R	TTATTCGGGAATCACATGCGTTTGGCTAAC

464   Introduced restriction sequences are underlined.

465 **Table 3 Sequences of sgRNA expression cassettes and repair templates**

gBlock	Sequences (5'-3')
fucI targeting sgRNA expression cassette	CGTACCATCG<u>TCTAGA</u>TGCGACAAGTAATAAACTAAACAAAA CAACTACAAAATATTTCTTTTTGTTTTTCATGATTTTTACACTT CTCTTAGTATGCTTTTGTTATAAGTTAGCACAAAAAAGCAGAA AATAAAAAGTAGAAATAAAAAAAGATGTTTTTTTGCCCATATC TCTATGAAAAAAACTGTGAAATGTGTAAAATATGGATGAAAC ATTGAATTTAAA<u>TGCTTCTTTAACACCGCCAATGGTTGAATCG</u> TCTCAGGTAGATGTCAGATCAATCTCTGACATCTACGAGTTGA GATCAAACAAAGCTTCAGCTGAGTTTCAATTTCTGAGCCCAT GTTGGGCCATACATTTTTTT<u>CCCGGG</u>ATCG
Transcriptional terminator *t1* targeting sgRNA expression cassette	CGTACCATCG<u>TCTAGA</u>TGCGACAAGTAATAAACTAAACAAAA CAACTACAAAATATTTCTTTTTGTTTTTCATGATTTTTACACTT CTCTTAGTATGCTTTTGTTATAAGTTAGCACAAAAAAGCAGAA AATAAAAAGTAGAAATAAAAAAAGATGTTTTTTTGCCCATATC TCTATGAAAAAAACTGTGAAATGTGTAAAATATGGATGAAAC ATTGAATTTAAA<u>CGGCATCTCATCATCTCCTCACCCAGACATG</u> TCTCAGGTAGATGTCAGATCAATCTCTGACATCTACGAGTTGA GATCAAACAAAGCTTCAGCTGAGTTTCAATTTCTGAGCCCAT GTTGGGCCATACATTTTTTT<u>CCCGGG</u>ATCG

Repair template for terminator *t1* deletion	GATACC<u>GTCGAC</u>CTTCGAATCAACAGGTGGAAAATGGTCTTA ACCAACGGCCTGCTCAATTAATCACAGATGGTCTTCCTAATGC AAATTATGGGTATATTAAAGGTAATAAATGGTATTTTGCTCAAT ACATAAGTCCGAGTGATGTCAAAACAATTTCTGAGGCAGGTA ATCTCCCTGACTTTTTACTAGTAGATAATACGACTAAGTTTGGC ATAGACCTTTTTATCAATGACTTGATAAAAACACTGCATCTATC AAGTTACTTTAGTGGGTATATGTTATATAGTGCAATCTTCGCGG ATTATTTAACTCCATTCAAAAAGGCTGCTCAAGAAGGTCGAG GAATTTATATTGTTACAGAGGCAGATATGACCAACCCAACTGC TGGCTATAACTATGTTGAGTATAGGATAGCGAGTTACTAACCTT ATTTTGAGTAGAGTTGACAAGGGCTTATTACTGGCTAAAAGG GCAAATACTATGAAAGACTTAGACTTATAACTGTTAAGAGA ATATTGATATTTGCCGTCATTGGCGCCATTGGTTGCATAGTAAC AGAGCTAGTAACCAAAATAGATCCTAATGTTGCCAGTTATATT TGGGTGATAGGCTTCTTTGTGACCTTTAAAGTTGCAGATAATG TCTCAATTAAAGCAACTGCAGGTAAAGGCAGCAAACGTTAAA GTATAATTACTTGACAGGGCCTGACAGATGTAATTGGAATGAC GAACGGAGAGTCTCCCTAACTGAATGATAAATAACGGCTGTG TCAACATAGTCGTTATTTTATTTCTATGAAACCTTGTTAGGCAA GCGTAGTTAATTTTAAAAACCATTATGAAAGCATTGCGAGAA TCAAGTTTAAAGTAGGTCCTGAAGAAGTTCAAATAAAACGTT TACATTTTGTTTATATGTGGTACAATTTGTTTGTTGATAGGAAT ATTCATTGTGGATTGTGACTATTCAATTAGGCGAGACCACAAA TCATGACGCTGAGAGTAGCGCGTTCTGGGTGAGGAGATGATG AGATGCCGAAAATCGATCAGATCTTTAACAATAACGTGGCGTT GGTTGAGCTAGACAACCATAGTCAAGCGGTTGTTAAGGGACG GGGGATCGCGTTTAGAAGCGTGGCGATGTGATTCCGGCTAAA CAAATTGAAAAAATTTTCTACCTTGCAAGTGAAACTTCCCGC CAGAATCTGTACTTTTTACTTCGCAATATTCCGATCGACGTGG TGACGACCACTTATGAAATTATTGACGTTGCGCAAAAACAATT TCACCTTAAGGTACTTGACTATATTTACATTACGTTAAGCGATC

| | ATATTTACGAAGCCTATAAACGTTATCAGAATGGAACGTATCA
AGAGACGATGGTGCCGGATTTTCATATTCAGTATCCTGCAGAG
TATGCCGTAGCTAACCAAGCGTTGCAGATCATTGCAGCTAACC
TTGGCGTAGCATTTCCACCTTCCGAAGTGAAAAATCTTGCCTT
ACATTTTATCAATGCCAGCGGTGAAGACGACAACGAGCAGGC
CTTTGACAAGAATAATGAGGCGTCATTAAGTCAACTGGTGCA
ACGTGTTTTGAAGCGGCATCGCATTACGAGATCGCAAACTAA
CGGCAATTATTATGATCGCTTCATGATTCACTTACAGTATCTGA
TTGACCGACTGCAGCGTGTTAATACGGATGCTGTTGCCATTGT
GCCCGAGGTGGCCAATGAATTAGAGCTAAATTATCCGCGGTC
TTATCAGATTGCTTCGGAAATTTTTGATGAGATTAAGGATCAG
CTTTATCGCAACATGAGTGAAGACGAACGCTTATACTTTATCA
TTCATATTCAACGGCTAATTAACGAAGCGCCTGCCCACGATCA
AAAACAGGGCAAAAAATCATAACGCGTTTGCTGGCTTGGAG
GGCAGCATACGCTCACGGGAGGAAAGTATCATGAATAAGGTT
TTCGATAAGCTAAAGCCGATTTTTGAAGCGATTGCGGCGAATA
GGGCCCGGTAC |

466 Introduced restriction sequences are underlined and targeting spacers are double

467 underlined.

Figure 1 Schematic of the harnessing the endogenous CRISPR-Cas9 system for genome engineering in LGG.

Figure 2 Characterization of the endogenous type II-A CRISPR-Cas9 system in

LGG. (A) Architecture of the CRISPR-Cas locus in LGG. The type II-A CRISPR-Cas9

system contains four Cas protein-encoding genes (*cas1, cas2, csn2*, and the critical *cas9*).

The CRISPR array includes 24 spacers (30 bp, blue rectangles) interspaced with repeats

(black diamonds). (B) PAM prediction visualized using a frequency plot generated using

WebLogo. (C) Schematic of the plasmid interference assay. (D) Interference plasmids

with a protospacer (underlined) matching the first spacer of the native CRISPR array,

with or without an adjacent PAM (5'-NGAAA-3'), were designed for plasmid

interference assays. (E) Transformation efficiencies of interference plasmids compared to

the control plasmid pTRK870. CFU: colony forming unit. Distinct letters were employed

to denote significant differences (*p < 0.05*). (F) Structure of the SgRNA, designed by

linking the crRNA and tracrRNA with an artificial AAUC loop. (G) Schematic of

targeting and editing plasmids. *repA* and *repC*, plasmid replication gene; *erm*,

erythromycin resistance gene; HR1 and HR2, the homologous repair templates; Ppgm,

the promoter of phosphoglycerate mutase (*Ppgm*) from *L. acidophilus* NCFM.

Figure 3 Deletion of *fucI* by reprogramming the endogenous CRISPR-Cas9 system

in LGG. (A) Diagram illustrating the genotype of the wild type (WT) and the *ΔfucI*

mutant. (B) Transformation efficiencies of the control plasmid pTRK870, the

homologous repair template containing plasmid pfucI-RT, the self-targeting plasmid

pfucI-sgRNA, and the editing plasmid pfucI-sgRNA-RT. Distinct letters were employed

to denote significant differences ($p < 0.05$). (C) Colony PCR screening of four surviving

transformants carrying the editing plasmid pfucI-sgRNA-RT to confirm the deletion

genotype. Expected products sizes corresponding to the WT and the *ΔfucI* strains are 2.7

kb and 2.1 kb, respectively. C: WT control; M: DNA marker; 1-4: surviving

transformants. (D) Sequencing results of the genotypes of the WT and MJM568 strains.

(E) Growth curve (absorbance at 595nm) of the WT (black) and MJM568 (red) in MRS

containing water (square), 2% glucose (circle) or 2% fucose (triangle).

Figure 4 Construction of lactose-positive LGG strain by harnessing endogenous CRISPR-Cas9 system and spontaneous mutation. (A) Diagram illustrating two mutations in the *lacTEGF* operon in WT LGG strain (yellow arrows): a 4-bp deletion causing a frameshift in the antiterminator gene (*lacT*) and a point mutation (Q231*) in the 6-phospho-β-galactosidase gene (*lacG*). RAT: ribonucleic antiterminator sequence; t1 and t2: rho-independent terminator. (B) Colony PCR screening of five surviving transformants carrying the editing plasmid placT-sgRNA-RT. C: WT control; M: DNA marker; 1-5: surviving transformants. (C) Sequencing results of the genotypes of the WT and MJM569 strains. The 20-bp deletion area is shaded in grey. (D) Growth of the WT and MJM569 strains on MRS agar plates containing 2% lactose as the sole sugar source. (E) Maximum cell growth rate (h^{-1}) of 15 randomly selected colonies (from MJM569 plates, black dots) and the WT strain (red dot) under glucose and lactose conditions. (F) Point mutations within the premature stop codon in *lacG* identified in these 15 randomly selected colonies. (G) Growth curve (absorbance at 595nm) of the WT (black) and MJM570 (blue) in MRS with 2% glucose (circle) or 2% lactose (triangle).

Figure 5 Growth of LGG and MJM570 strains in milk. (A) Growth curve of the WT

(blue square) and MJM570 (yellow triangle) in milk. Error bars show standard deviations

of the means from two parallel cultures. (B) LGG-milk remained liquid after 24 h at

37 °C, while MJM570-milk was coagulated after 24 h of incubation. (C) pH variation of

the WT (blue square) and MJM570 (yellow triangle) strains in milk. Error bars show

standard deviations of the means from three parallel cultures.

Figure 6 Yogurt fermentation with WT LGG and MJM570 strains. (A) Schematic of yogurt fermentation experiment. (B) pH variation of control, LGG, and MJM570 during yogurt fermentation (left), and the viability of LGG and MJM570 in yogurt during fermentation (right). (C) pH variation of control, LGG, and MJM570 during refrigerated storage (top), and the viability of LGG and MJM570 in yogurt during refrigerated storage (bottom). Distinct lowercase letters (e.g., abc) indicate statistically significant differences ($p < 0.05$) between different time points within the same group, while distinct uppercase letters (e.g., AB) indicate statistically significant differences ($p < 0.05$) between different groups at the same time points.

References

1.   Klaenhammer TR, Barrangou R, Buck BL, Azcarate-Peril MA, Altermann E. 2005.
Genomic features of lactic acid bacteria effecting bioprocessing and health. FEMS
Microbiol Rev 29:393–409.

2.   Das D, Goyal A. 2012. Lactic Acid Bacteria in Food Industry, p. 757–772. *In*
Satyanarayana, T, Johri, BN (eds.), Microorganisms in Sustainable Agriculture and
Biotechnology. Springer Netherlands, Dordrecht.

3.   Arqués JL, Rodríguez E, Langa S, Landete JM, Medina M. 2015. Antimicrobial
Activity of Lactic Acid Bacteria in Dairy Products and Gut: Effect on Pathogens.
BioMed Res Int 2015:584183.

4.   Martinez RCR, Bedani R, Saad SMI. 2015. Scientific evidence for health effects
attributed to the consumption of probiotics and prebiotics: an update for current
perspectives and future challenges. Br J Nutr 114:1993–2015.

5.   Linares DM, Gómez C, Renes E, Fresno JM, Tornadijo ME, Ross RP, Stanton C.
2017. Lactic Acid Bacteria and Bifidobacteria with Potential to Design Natural
Biofunctional Health-Promoting Dairy Foods. Front Microbiol 8:846.

6.   Mathur H, Beresford TP, Cotter PD. 2020. Health Benefits of Lactic Acid Bacteria
(LAB) Fermentates. Nutrients 12(6):1679.

7.   van Pijkeren J-P, Barrangou R. 2017. Genome Editing of Food-Grade Lactobacilli
To Develop Therapeutic Probiotics. Microbiol Spectr 5(5):BAD-0013-2016.

8.   Börner RA, Kandasamy V, Axelsen AM, Nielsen AT, Bosma EF. 2018. Genome
editing of lactic acid bacteria: opportunities for food, feed, pharma and biotech.
FEMS Microbiol Lett 366(1):fny291.

9.   Xie Z, McAuliffe O, Jin Y-S, Miller MJ. 2024. Genomic Modifications of Lactic
Acid Bacteria and Their Applications in Dairy Fermentation. J Dairy Sci
107(11):8749-8764.

10.   Hsu PD, Lander ES, Zhang F. 2014. Development and Applications of CRISPR-
Cas9 for Genome Engineering. Cell 157:1262–1278.

11.   Adli M. 2018. The CRISPR tool kit for genome editing and beyond. Nat Commun
9:1911.

12.   Song X, Huang H, Xiong Z, Ai L, Yang S. 2017. CRISPR-Cas9D10A Nickase-
Assisted Genome Editing in *Lactobacillus casei*. Appl Environ Microbiol
83(22):e01259-17.

13.   Leenay RT, Vento JM, Shah M, Martino ME, Leulier F, Beisel CL. 2018. Genome
Editing with CRISPR-Cas9 in *Lactobacillus plantarum* Revealed That Editing

Outcomes Can Vary Across Strains and Between Methods. Biotechnol J
14:1700583.

14. Huang H, Song X, Yang S. 2019. Development of a RecE/T-Assisted CRISPR–Cas9
Toolbox for *Lactobacillus*. Biotechnol J 14:1800690.

15. Goh YJ, Barrangou R. 2021. Portable CRISPR-Cas9N System for Flexible Genome
Engineering in *Lactobacillus acidophilus*, *Lactobacillus gasseri*, and *Lactobacillus*
paracasei. Appl Environ Microbiol 87(6):e02669-20.

16. Zhang J, Zong W, Hong W, Zhang Z-T, Wang Y. 2018. Exploiting endogenous
CRISPR-Cas system for multiplex genome editing in *Clostridium tyrobutyricum* and
engineer the strain for high-level butanol production. Metab Eng 47:49–59.

17. Vento JM, Crook N, Beisel CL. 2019. Barriers to genome editing with CRISPR in
bacteria. J Ind Microbiol Biotechnol 46:1327–1341.

18. Hidalgo-Cantabrana C, Goh YJ, Pan M, Sanozky-Dawes R, Barrangou R. 2019.
Genome editing using the endogenous type I CRISPR-Cas system in *Lactobacillus*
crispatus. Proc Natl Acad Sci 116:15774–15783.

19. Gu S, Zhang J, Li L, Zhong J. 2022. Repurposing the Endogenous CRISPR-Cas9
System for High-Efficiency Genome Editing in *Lacticaseibacillus paracasei*. ACS
Synth Biol 11:4031–4042.

20. Yan F, Polk DB. 2002. Probiotic Bacterium Prevents Cytokine-induced Apoptosis in
Intestinal Epithelial Cells*. J Biol Chem 277:50959–50965.

21. Glavina D, Gorseta K, Skrinjarić I, Vranić DN, Mehulić K, Kozul K. 2012. Effect of
LGG yoghurt on *Streptococcus mutans* and *Lactobacillus spp. salivary* counts in
children. Coll Antropol 36:129–32.

22. Sindhu KNC, Sowmyanarayanan TV, Paul A, Babji S, Ajjampur SSR, Priyadarshini
S, Sarkar R, Balasubramanian KA, Wanke CA, Ward HD, Kang G. 2014. Immune
Response and Intestinal Permeability in Children With Acute Gastroenteritis Treated
With *Lactobacillus rhamnosus* GG: A Randomized, Double-Blind, Placebo-
Controlled Trial. Clin Infect Dis 58:1107–1115.

23. Capurso L. 2019. Thirty Years of *Lactobacillus rhamnosus* GG: A Review. J Clin
Gastroenterol 53:S1-S41.

24. Kankainen M, Paulin L, Tynkkynen S, Ossowski I von, Reunanen J, Partanen P,
Satokari R, Vesterlund S, Hendrickx APA, Lebeer S, Keersmaecker SCJD,
Vanderleyden J, Hämäläinen T, Laukkanen S, Salovuori N, Ritari J, Alatalo E,
Korpela R, Mattila-Sandholm T, Lassig A, Hatakka K, Kinnunen KT, Karjalainen H,
Saxelin M, Laakso K, Surakka A, Palva A, Salusjärvi T, Auvinen P, Vos WM de.
2009. Comparative genomic analysis of *Lactobacillus rhamnosus* GG reveals pili
containing a human- mucus binding protein. Proc Natl Acad Sci 106:17193–17198.

25. Hussain N, Li R, Takala TM, Tariq M, Zaidi AH, Saris PEJ. 2021. Generation of
Lactose- and Protease-Positive Probiotic *Lacticaseibacillus rhamnosus* GG by
Conjugation with *Lactococcus lactis* NCDO 712. Appl Environ Microbiol
87:e02957-20.

26. Crawley AB, Henriksen ED, Stout E, Brandt K, Barrangou R. 2018. Characterizing
the activity of abundant, diverse and active CRISPR-Cas systems in lactobacilli. 1.
Sci Rep 8:11544.

27. Deveau H, Barrangou R, Garneau JE, Labonté J, Fremaux C, Boyaval P, Romero
DA, Horvath P, Moineau S. 2008. Phage Response to CRISPR-Encoded Resistance
in *Streptococcus thermophilus*. J Bacteriol 190:1390–1400.

28. Mojica FJM, Díez-Villaseñor C, García-Martínez J, Almendros C. 2009. Short motif
sequences determine the targets of the prokaryotic CRISPR defence system.
Microbiology 155:733–740.

29. Xie Z. 2024. PhD thesis. University of Illinois Urbana-Champaign, IL, USA.
Genomic Modification Tools for Probiotic Lactic Acid Bacteria and Their
Applications in Dairy Fermentation, Chapter 2.

30. Becerra JE, Yebra MJ, Monedero V. 2015. An l-Fucose Operon in the Probiotic
Lactobacillus rhamnosus GG Is Involved in Adaptation to Gastrointestinal
Conditions. Appl Environ Microbiol 81:3880–3888.

31. Heidtman MI, Merighi M, McCoy JM. February 2015. Strains of *Lactobacillus
rhamnosus* GG Capable of Growth on Lactose and the Uses Thereof.
US20150044330A1.

32. Segers ME, Lebeer S. 2014. Towards a better understanding of *Lactobacillus
rhamnosus* GG - host interactions. Microb Cell Factories 13:S7.

33. Claes IJJ, Schoofs G, Regulski K, Courtin P, Chapot-Chartier M-P, Rolain T, Hols P,
Ossowski I von, Reunanen J, Vos WM de, Palva A, Vanderleyden J, Keersmaecker
SCJD, Lebeer S. 2012. Genetic and Biochemical Characterization of the Cell Wall
Hydrolase Activity of the Major Secreted Protein of *Lactobacillus rhamnosus* GG.
PLOS ONE 7:e31588.

34. Lebeer S, Claes I, Tytgat HLP, Verhoeven TLA, Marien E, von Ossowski I,
Reunanen J, Palva A, de Vos WM, De Keersmaecker SCJ, Vanderleyden J. 2012.
Functional Analysis of *Lactobacillus rhamnosus* GG Pili in Relation to Adhesion
and Immunomodulatory Interactions with Intestinal Epithelial Cells. Appl Environ
Microbiol 78:185–193.

35. Goh YJ, Barrangou R. 2019. Harnessing CRISPR-Cas systems for precision
engineering of designer probiotic lactobacilli. Curr Opin Biotechnol 56:163–171.

36. Lomer MCE, Parkes GC, Sanderson JD. 2008. Review article: lactose intolerance in
clinical practice – myths and realities. Aliment Pharmacol Ther 27:93–103.

37. Jellema P, Schellevis FG, van der Windt DAWM, Kneepkens CMF, van der Horst
HE. 2010. Lactose malabsorption and intolerance: a systematic review on the
diagnostic value of gastrointestinal symptoms and self-reported milk intolerance.
QJM Int J Med 103:555–572.

38. Couvin D, Bernheim A, Toffano-Nioche C, Touchon M, Michalik J, Néron B, Rocha
EPC, Vergnaud G, Gautheret D, Pourcel C. 2018. CRISPRCasFinder, an update of
CRISRFinder, includes a portable version, enhanced performance and integrates
search for Cas proteins. Nucleic Acids Res 46:W246–W251.

39. Biswas A, Gagnon JN, Brouns SJJ, Fineran PC, Brown CM. 2013. CRISPRTarget.
RNA Biol 10:817–827.

40. Crooks GE, Hon G, Chandonia J-M, Brenner SE. 2004. WebLogo: A Sequence
Logo Generator. Genome Res 14:1188–1190.

41. Hidalgo-Cantabrana C, Goh YJ, Barrangou R. 2019. Characterization and
Repurposing of Type I and Type II CRISPR–Cas Systems in Bacteria. J Mol Biol
431:21–33.

42. Zadeh JN, Steenberg CD, Bois JS, Wolfe BR, Pierce MB, Khan AR, Dirks RM,
Pierce NA. 2011. NUPACK: Analysis and design of nucleic acid systems. J Comput
Chem 32:170–173.

43. Luchansky JB, Muriana PM, Klaenhammer TR. 1988. Application of
electroporation for transfer of plasmid DNA to *Lactobacillus*, *Lactococcus*,
Leuconostoc, *Listeria*, *Pediococcus*, *Bacillus*, *Staphylococcus*, *Enterococcus* and
Propionibacterium. Mol Microbiol 2:637–646.

44. Tharmaraj N, Shah NP. 2003. Selective Enumeration of *Lactobacillus delbrueckii*
ssp. *bulgaricus*, *Streptococcus thermophilus*, *Lactobacillus acidophilus*,
Bifidobacteria, *Lactobacillus casei*, *Lactobacillus rhamnosus*, and Propionibacteria.
J Dairy Sci 86:2288–2296.

672