Exploiting Endogenous CRISPR-Cas9 System for Functional Engineering of Probiotic

Lacticaseibacillus rhamnosus **GG (LGG)**

Authors: Zifan Xie[1,2], Yong-Su Jin[1,2]*, and Michael J. Miller[1,2]*

Affiliation:

[1]Department of Food Science and Human Nutrition, University of Illinois Urbana-Champaign,

Urbana, IL, USA

[2] Carl R. Woese Institute for Genomic Biology, University of Illinois Urbana-Champaign,

Urbana, IL, USA

E-mail: Zifan Xie (zifanxie@illinois.edu), Yong-Su Jin (ysjin@illinois.edu), Michael J. Miller

(mille216@illinois.edu)

*Corresponding author:

Prof. Yong-Su Jin (ysjin@illinois.edu)

Prof. Michael J. Miller (mille216@illinois.edu)

ABSTRACT

Lacticaseibacillus rhamnosus GG (LGG) is one of the most extensively studied probiotic strains, widely

used in food and health applications. However, the absence of efficient, precise genome editing methods

has limited its broader potential and functional versatility. Here, we present an endogenous type II-A

CRISPR-Cas9 genome editing workflow for LGG designed for functional strain construction. Using a

plasmid interference assay together with single-nucleotide substitutions, we confirm the precise PAM

requirement as 5′-NGAAA-3′. We pair a synthetic sgRNA cassette with homology-directed repair

donors to enable precise deletions and insertions across multiple loci. Using this optimized and precise

genome engineering method, we generated a β-glucuronidase (GUS)-expressing LGG strain for robust

strain tracking within complex gut microbial communities. This work removes barriers to LGG

engineering, expands the probiotic CRISPR toolkit, and provides broadly applicable strategies for

designing next-generation probiotics with applications in food biotechnology and microbial therapeutics.

Keywords: *CRISPR-Cas9*, *probiotic, synthetic biology*, *lactic acid bacteria*

GRAPHICAL ABSTRACT

INTRODUCTION

Lacticaseibacillus rhamnosus GG (LGG) is one of the most extensively studied probiotic strains due to its well-documented therapeutic and industrial benefits. Originally isolated from the human gastrointestinal tract, LGG demonstrates robust tolerance to gastric acid and bile salts, strong adherence to intestinal epithelial cells, and exceptional stability throughout food processing (Segers & Lebeer, 2014; Capurso, 2019). Since its discovery, numerous clinical and preclinical studies have demonstrated LGG's ability to enhance gut barrier function, modulate host immune system, and exert antimicrobial activity (Hudault et al., 1997; Yan & Polk, 2002; Sindhu et al., 2014; Gao et al., 2019). Given its Generally Recognized as Safe (GRAS) status from the US Food and Drug Administration (GRAS Notice No. 845), LGG has been widely used in dietary supplements, fermented food products, and clinical therapies aimed at modulating gut health, immune modulation, and microbial homeostasis (Jia et al., 2016; Chan et al., 2021). More recently, LGG's well-established safety profile and versatile functionality have attracted significant interest in its development as a microbial chassis for next-generation live biotherapeutics and precision fermentation applications (Günaydın et al., 2014; Wan et al., 2016; Petrova et al., 2018; Li et al., 2025).

Despite over 300 human-focused publications underscoring LGG's extensive health benefits, its genetic manipulation and strain optimization have been significantly limited by a lack of efficient genome editing tools. Because of limited replicable plasmids, chromosomal modification in LGG has typically relied on non-replicative suicide vectors (Lebeer et al., 2012; S. Zhang et al., 2018). These approaches depend on low-frequency double-crossover events and usually require multiple rounds of selection or counterselection, making them labor-intensive, time-consuming, and poorly suited to complex or high-throughput engineering. The absence of precise genome-editing technologies has thus

hindered mechanistic studies of LGG's probiotic actions and restricted the development of advanced

strains for therapeutic and industrial applications.

Recently, many innovative genome modification approaches have been developed and applied in

lactic acid bacteria to determine relevant phenotypes and to enhance beneficial properties (Xie et al.,

2024). Notably, the CRISPR-Cas (clustered regularly interspaced short palindromic repeats and

CRISPR-associated proteins) system, originally discovered as a bacterial adaptive immune mechanism,

has emerged as a powerful tool for precise and efficient genome modification in various organisms (Hsu

et al., 2014; Adli, 2018). Heterologous CRISPR-Cas systems have been successfully employed in many

lactobacilli, such as *Limosilactobacillus reuteri*, *Lactiplantibacillus plantarum*, *Lacticaseibacillus casei*,

Lactobacillus acidophilus, *Lactobacillus gasseri*, and *Lacticaseibacillus paracasei* (Song et al., 2017;

Leenay et al., 2018; Huang et al., 2019; Goh & Barrangou, 2021). Despite these successes, heterologous

CRISPR systems present several challenges. These systems generally require at least one plasmid for the

expression of large Cas9 protein, most commonly derived from *Streptococcus pyogenes* (Cong et al.,

2013). Such large plasmid constructs often exhibit reduced plasmid stability, lower transformation

efficiency, and impose metabolic burdens on host cells. Additionally, heterologous Cas proteins may

exhibit suboptimal activity or cytotoxicity in certain strains, further limiting editing efficiency and

viability (J. Zhang et al., 2018; Vento et al., 2019). To overcome these limitations, recent research has

increasingly focused on endogenous CRISPR-Cas systems encoded within the host genomes. These

native systems eliminate the need for exogenous proteins and exhibit enhanced compatibility with host

machinery, resulting in improved editing precision and reduced cellular toxicity. Functional CRISPR-

Cas systems have been identified in many lactobacilli strains (Crawley et al., 2018), and some of them

have been successfully repurposed for precise genome editing. For example, the endogenous subtype I-E

CRIPSR-Cas system has been utilized for efficient genetic manipulation of *Lactobacillus crispatus*

(Hidalgo-Cantabrana, Goh, Pan, et al., 2019), and the subtype II-A CRISPR-Cas9 system has been used

to achieve genome engineering in *Lcb. paracasei* (Gu et al., 2022). Recently, type II-A CRISPR-Cas9

system was identified in LGG (Crawley et al., 2018; Panahi et al., 2023), and the Cas9 protein from

LGG has been heterologously expressed for plant genome engineering (Zhong et al., 2023). However,

those studies did not establish a practical genome-editing workflow directly inside LGG.

Here, we establish an endogenous type II-A CRISPR-Cas9 genome-editing system for LGG that

enables precise gene deletions and insertions. Using a plasmid-interference assay with single-nucleotide

protospacer adjacent motif (PAM) variants, we confirmed the specificity of the native Cas9 PAM as 5′-

NGAAA-3′. A replicative editing plasmid was constructed to carry synthetic single-guide RNA

(sgRNA) cassettes and homologous repair templates, enabling efficient genome modifications across

multiple loci. To demonstrate the practical applicability of this system, we engineered a β-glucuronidase

(gusA) expressing strain that allows precise tracking in complex microbial communities and host

models. Animal experiments further confirmed that the gusA-positive LGG strain serves as a suitable

model for studying LGG-associated health benefits. Together, these results establish an efficient LGG

genome-editing platform that facilitates mechanistic investigations and accelerates the development of

functional probiotic strains for food biotechnology and microbial therapeutics.

RESULTS

Specificity of the Endogenous CRISPR-Cas9 System in LGG

LGG harbors a complete endogenous type II-A CRISPR-Cas9 locus containing four Cas protein-

encoding genes (*cas1*, *cas2*, *csn2*, and *cas9*) adjacent to a CRISPR array (Crawley et al., 2018; Panahi et

al., 2023). The array contains 24 distinct 30-base pair (bp) spacers interspaced by 36-bp direct repeats

(Figure 1A). Consistent with previous reports, alignment of protospacer-flanking sequences in LGG

indicates a preferred PAM of 5′-NGAAA-3′ (Figure 1B) (Crawley et al., 2018; Zhong et al., 2023)

To validate the specificity of the PAM sequence, we performed a plasmid interference assay

incorporating single-nucleotide PAM substitutions (Figure 1C). Interference plasmids were constructed

based on vector pTRK870 (Supplementary Table S2), a derivative of pTRK830 with a multiple cloning

site (Xie et al., 2025). Each plasmid carried a protospacer matching the most recently acquired spacer 24

(5'-AAGAAACGATCCCGAGTTTCTGGATTGCTT-3') in the CRISPR array, either accompanied by

or lacking the predicted PAM sequence.

Transformation of LGG with plasmids (pPS1AGAAA, pPS1TGAAA, and pPS1GGAAA)

containing the predicted PAM sequences resulted in no detectable transformants, demonstrating robust

interference mediated by the endogenous CRISPR-Cas9 system (Figure 1D). To further evaluate PAM

specificity, we introduced single-nucleotide substitutions at conserved positions (positions 2-5) of the

predicted PAM sequence, generating seven PAM variants AAAAA, ACAAA, TCAAA, AGGAA,

AGTAA, AGATA, and AGAAG. As shown in Figure 1D, plasmids with altered PAM sequences,

pPS1AAAAA, pPS1ACAAA, and pPS1TCAAA, failed to produce a detectable interference response,

indicating that guanine at position 2 is essential for effective PAM recognition. Additionally, plasmid

pPS1AGATA, carrying a thymine substitution at position 4, showed significantly reduced interference,

indicating the importance of adenine conservation at this position. Notably, plasmids pPS1AGGAA and

pPS1AGAAG, each harboring substitutions at positions 3 and 5, exhibited partial interference,

suggesting a moderate tolerance of these positions to guanine substitutions. However, significantly

fewer colonies were recovered using the optimal PAM (pPS1AGAAA), confirming a strong preference

for adenines at positions 3–5.

Collectively, these results confirm the PAM recognized by the endogenous Cas9 in LGG as 5′-

NGAAA-3′ and demonstrate its high sequence specificity, with enhanced interference activity observed

under our assay conditions.

Construction of Editing Plasmids for Reprogramming the Endogenous CRIPSR-Cas9 System

After confirming the specificity of PAM sequence, we then explored its potential for genome

editing in LGG. Initially, we designed an editing plasmid derived from pTRK870, which contains a

homologous repair template and a CRISPR RNA (crRNA) expression cassette under control of the

native CRISPR array leader sequence (Supplementary Figure S1). However, we found that due to the

inherent instability of rolling circle plasmids (Xie et al., 2025), the space region between the repeats was

frequently looped out from the plasmid during plasmid replication, resulting in spacer-deficient plasmids

and a high frequency of wild-type (WT) survivors (Supplementary Figure S1).

To overcome this limitation, we transitioned to a sgRNA-based approach for directing precise

genome targeting by the endogenous CRISPR-Cas9. Based on the predicted structure of the crRNA/

trans-activating CRISPR RNA (tracrRNA) duplex and prior transcriptional sequencing data (Crawley et

al., 2018), we engineered an artificial sgRNA by linking the crRNA and tracrRNA with an AAUC loop

(Figure 1E). We then constructed a targeting plasmid encoding the sgRNA under the control of the

strong constitutive promoter *Ppgm* from *L. acidophilus* NCFM (Figure 1F) (Xie et al., 2025).

To validate our artificial sgRNA design, we targeted a highly conserved region within the 16S

ribosomal RNA (rRNA) gene. Introduction of this 16S-targeting plasmid into LGG significantly reduced

viable transformants compared to the pTRK870 control plasmid (Supplementary Figure S2),

demonstrating effective Cas9-mediated chromosomal cleavage and providing direct evidence of

functional sgRNA expression. To evaluate the broader applicability of this sgRNA system, we tested the

same 16S-targeting plasmid in several additional *L. rhamnosus* strains (Supplementary Table S1). Unlike

LGG, these other strains yielded viable transformants without evidence of Cas9-mediated lethality

(Supplementary Figure S2), suggesting limited portability of our platform across closely related strains.

Possible reasons include differences in plasmid transformation efficiency, strain-specific PAM sequence

preferences, or variations in CRISPR architecture among strains (Supplementary Figure S2).

Building upon this validated targeting plasmid, we next designed a comprehensive editing

plasmid incorporating both the sgRNA expression cassette and homologous repair templates (HR1 and

HR2) flanking the intended genomic modification site (Figure 1F).

Reprogramming the Endogenous CRISPR-Cas9 System for Gene Deletion

As a test, we first attempted to target the *fucI* gene (LGG_02685), which encodes L-fucose

isomerase (Figure 2A). Deletion of *fucI* is expected to significantly impair LGG growth when fucose is

the sole carbon source (Becerra et al., 2015). A 30-bp DNA sequence complementary to *fucI* was

selected as the spacer and cloned into the pTRK870 backbone, together with two 1-kb homologous

arms, to generate the editing plasmid pfucI-sgRNA-RT (Supplementary Table S2). As controls, we also

constructed plasmids containing only the repair template (pfucI-RT) or only the sgRNA (pfucI-sgRNA)

(Supplementary Table S2). Transformation of LGG with the self-targeting plasmid (pfucI-sgRNA)

yielded no colonies (Figure 2B), confirming effective genome cleavage triggered by the endogenous

CRISPR-Cas9 system guided by the sgRNA. In contrast, transformation with the repair template-only

plasmid (pfucI-RT) generated numerous colonies (Figure 2B), but none of screened colonies carried the

intended deletion genotype (Supplementary Figure S3), indicating that the repair template alone was

insufficient for generating targeted deletions efficiently. Importantly, transformation with the complete

editing plasmid (pfucI-sgRNA-RT) resulted in significantly fewer colonies (four colonies). PCR

screening of these transformants identified one colony (Colony No. 1) displaying the expected 2.1-kb

PCR product, indicative of the deletion genotype, whereas the other three exhibited wild-type PCR

products (2.7 kb; Figure 2C). Sanger sequencing of the 2.1-kb amplicon confirmed the precise deletion,

and the successfully engineered strain was designated MJM568 (Figure 2D). Phenotypic analysis

further revealed that the MJM568 strain could not utilize fucose as the sole sugar source (Figure 2E).

To further evaluate the versatility of this system, we successfully deleted a larger 1.5-kb genomic

region encoding coenzyme A (CoA) synthase (*acs1*) using the same strategy (Supplementary Figure S4).

This demonstrated the platform's capability for targeting and deleting genomic regions of varying

lengths. Collectively, these results confirm that the LGG endogenous CRISPR-Cas9 system can be

effectively harnessed for targeted gene deletions, with both the sgRNA-guided Cas9 cleavage and

homologous repair templates essential for achieving precise genomic modifications.

Reprogramming the Endogenous CRISPR-Cas9 System for Gene Insertion

After demonstrating the feasibility of gene deletions using the native CRISPR-Cas9 system in

LGG, we next aimed to evaluate the system's capacity for gene insertion. The ability to stably integrate

heterologous genes is a critical feature for engineering LGG as a microbial cell factory or live

biotherapeutic. However, the previously used plasmid backbone pTRK870, which replicates via a

rolling-circle mechanism, is inherently unstable and has limited insert capacity (Xie et al., 2025). To

address these limitations, we constructed a new editing backbone, pMJM100 (Supplementary Figure

S5), which is a smaller and more stable plasmid with improved capacity for accommodating larger

genetic inserts. This redesigned backbone provides a robust platform for gene insertion mediated by the

native CRISPR-Cas9 machinery in LGG.

To test this platform, we repurposed the *fucI*-targeting sgRNA expression cassette previously

used for gene deletion and combined it with a homologous repair template designed to partially delete

fucI and simultaneously insert a chloramphenicol resistance gene (*cam*) expression cassette (Figure 3A).

The *cam* cassette was flanked by 0.5 kb homology arms derived from the original *fucI* deletion template.

The use of shorter arms reduced the plasmid size and improved its stability during cloning and

transformation. The resulting construct, pfucI-sgRNA-Cm-RT2, was assembled using the pMJM100

backbone.

Following transformation of pfucI-sgRNA-Cm-RT2 into LGG, six erythromycin-resistant

transformants were obtained (Figure 3B). PCR screening revealed that one transformant (Colony No. 1)

displayed the expected amplicon size for the *cam* gene insertion (Figure 3C), which was further

confirmed by Sanger sequencing (Figure 3D). To eliminate the editing plasmid, the verified mutant was

passaged twice under non-selective conditions, yielding an erythromycin-sensitive derivative

(Supplementary Figure S6), designated MJM655. As excepted, MJM655 showed robust growth on MRS

plates supplemented with 5 µg/mL chloramphenicol, while the WT LGG strain remained

chloramphenicol-sensitive (Figure 3E).

Construction of GUS-Positive LGG

To demonstrate the application of our endogenous CRISPR-Cas9 system for functional gene

insertion, we engineered a GUS-positive LGG strain by inserting the *gusA* gene from *Lactobacillus*

gasseri ADH, a GRAS strain. The *gusA* gene encodes β-glucuronidase and is widely used as a reporter

in lactic acid bacteria. We aimed to create a chromosomally integrated reporter strain suitable for

tracking in complex microbial communities.

To achieve this, we constructed the editing plasmid prha-sgRNA-gusA-RT2, designed to

introduce *gusA* gene expression cassette into the intergenic region between the fucose and rhamnose

operons (Figure 4A). Following transformation and selection, colony PCR of one transformant (Colony

No. 1) showed both wild-type and mutant bands (Figure 4B), indicating a mixed population. Re-

isolation of colony No.1 yielded colonies with a single PCR product of the expected insertion size

(Supplementary Figure S7), and Sanger sequencing confirmed precise genomic integration of the *gusA*

cassette at the target locus (Figure 4C). After plasmid curing through non-selective subculturing, the

resulting genome-edited strain was designated MJM670. To verify *gusA* expression, we plated MJM670

and WT LGG on MRS agar supplemented with X-gluc. As expected, MJM670 formed distinct blue

colonies, while the WT LGG colonies remained white (Figure 4D), confirming successful chromosomal

expression of *gusA*. Moreover, MJM670 maintained consistent GUS expression over serial passages, in

contrast to a plasmid-based *gusA* construct, which showed progressive loss of expression (Figure 4E),

underscoring the advantage of chromosomal integration for long-term genetic stability.

To evaluate whether the insertion affected adjacent gene function, we compared the growth

profiles of MJM670 and WT LGG in media supplemented with glucose, fucose or rhamnose. As shown

in Supplementary Figure S8, MJM670 exhibited growth patterns nearly identical to the wild-type strain

under all tested sugar conditions, indicating that *gusA* integration did not disrupt the function of the

flanking fucose and rhamnose operons. Together, these results demonstrate that the intergenic region

between the fucose and rhamnose operons is a permissive site for gene insertion, and that prha-sgRNA-

RT2 is an effective vector for functional gene integration in LGG. This strategy offers a valuable tool for

constructing recombinant probiotic strains for metabolic enhancement and therapeutic applications.

Quantification of GUS-Positive LGG in Mouse Fecal Samples

Analyzing LGG abundance in fecal samples offers a non-invasive and comprehensive view of

bacterial persistence through the gastrointestinal (GI) tract. Fecal enumeration is particularly valuable

for assessing long-term colonization and microbial dynamics, providing insight into the overall stability

and retention of probiotic strains in vivo. To evaluate the persistence and viability of engineered GUS-

positive LGG, MJM670, we conducted both plating- and qPCR-based analysis of fecal samples

collected from C57BL/6J mice following a single oral gavage (Figure 5A).

236   As previously shown in Figure 4D, MJM670 expresses β-glucuronidase, enabling colony

visualization on MRS agar supplemented with X-Gluc. GUS activity resulted in distinct blue colonies,

allowing clear differentiation from native gut microbiota (Supplementary Figure S9). Based on plating

analysis (Figure 5B), approximately 10^7 CFU/g MJM670 were detected in feces 12 hours after one-time

gavage, which decreased to ~10^5 CFU/g at 24 hours. By 48 hours, MJM670 was undetectable in all

groups, indicating complete clearance. In parallel, we performed qPCR using LGG-specific primers to

assess total LGG abundance regardless of cell viability (Figure 5C). The qPCR results revealed that

MJM670 exhibited similar survival dynamics in the GI tract as the WT LGG strain (Figure 5C), with

both showing a rapid decline over time. These results further support the observation that both WT LGG

and MJM670 exhibit transient colonization following oral administration.

246   When comparing the two methods (Figure 5B and 5C), the plating method consistently yielded

higher LGG counts than qPCR. This is likely due to incomplete DNA recovery from fecal samples using

commercial stool DNA kits, which may underestimate total bacterial abundance. Moreover, while qPCR

can detect both viable and nonviable cells, plating selectively recovers only viable LGG cells capable of

colony formation, providing a direct measure of functional probiotic survival. Despite its higher

detection threshold, the plating method provides unique advantages for engineered strains like MJM670.

Its visual specificity enables clear identification of the target strain, and it allows isolation of individual

colonies for downstream phenotypic or genomic characterization, which is an essential step for verifying

genetic stability or strain behavior after going through the GI tract.

Overall, the GUS-positive LGG strain, MJM670, demonstrated reliable detection via plating and

enabled viable cell recovery, supporting its application in colonization tracking and functional studies

within the gastrointestinal microbiome.

DISCUSSION

LGG is one of the most widely used probiotics in fermented foods and dietary supplements, yet

the development of enhanced LGG strains has been constrained by the lack of efficient genome editing

tools. Existing methods based on non-replicable vectors are inefficient, labor-intensive, and require

antibiotic selection. While heterologous CRISPR-Cas9 systems have been applied in other *Lactobacillus*

species, their use in LGG has been restricted due to the lack of stable plasmids capable of

accommodating the large *cas9* gene. In this situation, repurposing the endogenous CRISPR–Cas

machinery offers a promising approach for editing in LGG. A type II-A CRISPR–Cas9 system has

previously been identified in LGG, and its Cas9 protein has been heterologously expressed for plant

genome engineering (Crawley et al., 2018; Zhong et al., 2023). However, these studies did not establish

a practical genome-editing workflow in LGG. Critical parameters, such as PAM sequence specificity in

the LGG context, sgRNA expression system, homologous repair templates design, plasmid curing, and

edit stability, have remained undefined. Here, we addressed these gaps and developed an endogenous

type II-A CRISPR-Cas9 platform for LGG that enables efficient genome editing to construct functional

strains for fermentation and therapeutic applications.

We first confirmed 5'-NGAAA-3' as the preferred PAM, exhibiting strict sequence requirements

for target recognition. To assess the reprogrammability of the native system, we designed an artificial

sgRNA targeting the essential 16S rRNA gene. Introduction of this construct resulted in a marked loss of

viable transformants in LGG, confirming the high activity and specificity of the native system and

demonstrating that artificial sgRNAs can be correctly recognized and processed in LGG. In contrast, the

same construct failed to induce lethality in other *Lcb. rhamnosus* strains, despite some encoding type II-

A CRISPR loci. These strains showed no detectable targeting activity, likely due to low transformation

efficiency or variations in PAM specificity or CRISPR locus architecture. These findings highlight the

strain-dependent nature of endogenous CRISPR systems and underscore the importance of strain-

specific validation for successful repurposing.

Using the validated system, we developed replicable plasmids for sgRNA expression and repair

template delivery, achieving precise deletions and insertions without scars or antibiotic-resistant genes,

making them suitable for food and therapeutic applications. Although overall editing efficiency in LGG

was lower than in some other lactobacilli (Gu et al., 2022; Hidalgo-Cantabrana, Goh, Pan, et al., 2019),

likely due to the inherent instability of rolling-circle plasmids and relatively lower homologous

recombination efficiency, this workflow offers several practical advantages. The system enables scarless

gene deletions and insertions using a single plasmid, without the need for heterologous Cas9 expression

or laborious selection. In practice, desired mutants can be routinely obtained in one transformation and a

brief colony screen, typically completed within three days, without multiple rounds of subculturing.

Moreover, the editing plasmids are easily curable, leaving no selection or counterselection markers in

the final strains, which support iterative marker-free editing to stack multiple modifications in the same

genetic background. Together, these attributes make the platform well-suited for functional genetics,

strain optimization, and medium-throughput genome engineering in LGG, where speed, precision, and

regulatory compatibility are at a premium.

To demonstrate the application of this genome editing platform, we engineered a GUS-

expressing LGG strain (MJM670) to enable specific and robust tracking within mixed microbial

populations and animal models. In the gastrointestinal tract, endogenous β-glucuronidase activity is

present in only a limited number of bacterial species beyond *Escherichia coli*, making *gusA* a reliable

and specific reporter gene. Genomic integration into LGG enables clear visual differentiation from

native gut microbes and provides a direct alternative to qPCR-based detection. While qPCR detects

DNA from both viable and nonviable cells, plating recovers only viable LGG, providing a direct

measure of functional probiotic survival. The *gusA* gene, sourced from the GRAS strain *L. gasseri* ADH,

supports biosafety and avoids antibiotic resistance genes, reducing the risk of horizontal gene transfer

and the spread of antibiotic resistance within the gut environment. This GUS-positive LGG enables both

qualitative and quantitative monitoring in complex communities, like co-culture systems, fecal samples,

and host tissues. Therefore, these features make GUS-positive LGG a valuable tool for mechanistic

probiotic research and a promising chassis for therapeutic protein or antigen delivery in live

biotherapeutics.

In conclusion, we present a functional genome editing platform for LGG based on its

endogenous CRISPR-Cas9 system. This system enables precise, scarless gene deletions and insertions

using a single plasmid. The platform is rapid, convenient, and broadly applicable, making it a valuable

tool for LGG functional studies, probiotic engineering, and industrial strain development. The successful

engineering of GUS-expressing LGG strains highlights the potential of this system for live

biotherapeutic applications. In the future, this platform has the potential to broaden LGG's role as a

programmable chassis for therapeutic delivery and precision fermentation. By enabling efficient and

scarless genome editing, our platform paves the way for future research and strain development in LGG

and related *Lactobacillus* species, offering a powerful tool to enhance probiotic functionality and

industrial utility.

MATERIAL AND METHODS

Strains and Culture Conditions

All *Lcb. rhamnosus* strains (listed in Supplementary Table S1) were cultivated in de-Man-Rogosa-Sharpe (MRS; Hardy Diagnostics, Santa Maria, CA, USA) broth or on MRS agar (1.5%) at 37 °C under anaerobic conditions (90% N_2, 5% CO_2, and 5% H_2). *Escherichia coli* MC1061 was cultured in Luria-Bertani (LB) broth with agitation at 250 rpm or on LB agar (1.5%) and incubated at 37 °C. For GUS activity detection, MRS agar plates were supplemented with 200 µg/mL 5-Bromo-4-chloro-1*H*-indol-3-yl β-D-glucopyranosiduronic acid (X-gluc; Gold Biotechnology, USA). When appropriate, antibiotics were added at the following concentrations: erythromycin (Em; Gold Biotechnology, St. Louis, MO, USA) at 5 µg/ml for LGG and 150 µg/ml for *E. coli*, and chloramphenicol (Cm; Sigma, St. Louis, MO, USA) at 5 µg/ml for LGG.

CRISPR-Cas System Characterization *in silico*

The complete genome sequence of LGG (GenBank: CP031290.1) is available at the National Center for Biotechnology Information (NCBI) database. CRISPR array and associated Cas proteins were identified using the CRISPRCasFinder platform (Couvin et al., 2018). PAM sequences were predicted by aligning LGG spacer to target sequences with CRISPRTarget (Biswas et al., 2013), and consensus PAM sequences were presented as sequence logo generated by WebLogo (Crooks et al., 2004). For type II CRISPR systems, tracrRNA contains a region complementary to the repeat sequence of the crRNA; thus, the repeat sequence was queried against the genome using the Basic Local Alignment Search Tool (BLAST) to locate the tracrRNA (Hidalgo-Cantabrana, Goh, & Barrangou, 2019). The crRNA–tracrRNA duplex was further modeled using the NCPACK web server and manually illustrated (Zadeh et al., 2011).

DNA Manipulations and Molecular Cloning

Genomic DNA was isolated using the Wizard genomic DNA purification kit (Promega

Corporation, Madison, WI, USA). Plasmid DNA from *E. coli* was prepared with the QIAGEN plasmid

mini-prep kit (QIAGEN, Germantown, MD, USA). Restriction enzymes and T4 DNA ligase (New

England Biolabs, Ipswich, MA, USA) were used according to the supplier's protocols. Polymerase chain

reactions (PCR) were conducted following standard procedures with either DreamTaq DNA polymerase

(Thermo Fisher Scientific, Waltham, MA, USA) or Phusion DNA polymerase (NEB, Ipswich, MA,

USA). Oligonucleotide primers (Supplementary Table S3) were ordered from Integrated DNA

Technologies (IDT, Coralville, IA, USA) and synthetic gene fragments (Supplementary Table S4) were

obtained from Twist Bioscience (Twist Bioscience, San Francisco, CA, USA). PCR amplicons were

electrophoresed on 1.0% agarose gels and purified using the QIAquick PCR purification kit (QIAGEN,

Germantown, MD, USA). Sanger sequencing was performed at the UIUC Core Sequencing Facility

(Urbana, IL, USA), and whole plasmid sequencing was conducted by Plasmidsaurus, an Oxford

Nanopore certified service.

Construction of Interference Plasmids

pTRK830 is a replicating shuttle vector for *E. coli* and lactobacilli strains (Xie et al., 2025). The

multiple cloning site (MCS) from pBluescript KS (+) was ligated into the *StuI* site of pTRK830, which

generated pTRK870 (Supplementary Table S2). Plasmid pTRK870 was used for all following plasmid

constructions. The interference plasmids were constructed as described before (Crawley et al., 2018).

The double-stranded DNA protospacers, with or without the PAM, were generated by PCR using and

then ligated into XbaI-HindIII-digested pTRK870 to construct interference plasmids. The resulting

interference plasmids were PCR-screened in *E. coli* transformants using PS1 and PS2 primers

(Supplementary Table S3) for the presence of the insert and sequenced to confirm sequence content.

Construction of Editing Plasmids

Each editing plasmid consisted of two key components: a sgRNA expression cassette and a

repair template. The sgRNA expression cassettes included a promoter *Ppgm* of the phosphoglycerate

mutase from *L. acidophilus* NCFM, a designed sgRNA containing a 30-bp sequence targeting a specific

genomic locus of LGG strain, and a Rho-independent terminator.

***fucI* gene deletion**: The *fucI* targeting sgRNA expression cassette (Supplementary Table S4) was

synthesized and amplified using primers sgRNA-XbaI and sgRNA-XmaI (Supplementary Table S3),

then cloned into XbaI-XmaI-digested pTRK870 to generate targeting plasmid pfucI-sgRNA

(Supplementary Table S2). The repair template with two 1-kb homologous regions for *fucI* gene editing

was amplified from the genomic DNA of LGG strain. For this, two 1-kb homologous regions were fused

by overlap extension PCR to obtain the 2-kb repair template to the *fucI* gene. This repair template was

digested with SalI and ApaI, then cloned into SalI-ApaI-digested plasmids pfucI-sgRNA and pTRK870

to generate plasmids pfucI-sgRNA-RT and pfucI-RT (Supplementary Table S2), respectively. All these

final plasmid constructs were PCR-screened using PS1 and PS2 primers (pTRK870 based) and

sequenced to confirm sequence content.

***acs1* gene deletion**: The *acs1* targeting sgRNA expression cassettes (Supplementary Table S4)

were synthesized and amplified using primers sgRNA-XbaI and sgRNA-XmaI (Supplementary Table

S3), then cloned into XbaI-XmaI-digested pTRK870 to generate targeting plasmids pacs1-sgRNA

(Supplementary Table S2). The repair template (Supplementary Table S4) with two 1-kb homologous

regions was synthesized and cloned into the PstI-ApaI-digested pacs1-sgRNA generating pacs1-sgRNA-

RT (Supplementary Table S2). All these final plasmid constructs were PCR-screened using PS1 and PS2

primers (pTRK870 based) and sequenced to confirm sequence content.

***cam* expression cassette insertion:** To construct a compact and stable cloning vector for gene

insertion, a multiple cloning site sequence was inserted into StuI-MfeI-digested pTRK829

(Supplementary Table S2), removing the entire *cam* gene and other non-essential regions, to generate

pMJM100 (Supplementary Figure S2). The *fucI* targeting sgRNA expression cassette (Supplementary

Table S4) was cloned into XbaI-XmaI-digested pMJM100 to construct a new *fucI* targeting plasmid

pMJM100-fucI-sgRNA (Supplementary Table S2). The repair template, containing two 0.5-kb

homologous regions, was amplified from pfucI-RT using primers fucI-RT2-SalI and fucI-RT2-ApaI

(Table S3), and cloned into a TOPO vector using the Zero Blunt® TOPO® PCR Cloning Kit (Thermo

Fisher Scientific, Waltham, MA, USA), resulting in pTOPO-fucI-RT2 (Supplementary Table S2). The

cam gene expression cassette was amplified from pTRK829 using primers Cm-F and Cm-R and inserted

into pTOPO-fucI-RT2 at the *HindIII* site, generating pTOPO-fucI-Cm-RT2 (Supplementary Table S2).

Finally, the complete repair template, containing the *cam* gene expression cassette flanked by two 0.5-kb

homologous regions, was cloned into SalI-ApaI-digested pMJM100-fucI-sgRNA, resulting in the final

editing plasmid pfucI-sgRNA-Cm-RT2 (Supplementary Table S2). pfucI-sgRNA-Cm-RT2 was PCR-

screened using PS3 and PS4 primers (pMJM100 based) and sequenced to confirm sequence content.

gusA expression cassette insertion: The sgRNA expression cassette targeting to intergenic

region between fucose and rhamnose operons (Supplementary Table S4) was synthesized and amplified

using primers sgRNA-XbaI and sgRNA-XmaI (Supplementary Table S3), then cloned into XbaI-XmaI-

digested pMJM100 to generate targeting plasmid pMJM100-rha-sgRNA (Supplementary Table S2). The

repair template (Supplementary Table S4) with two 0.5-kb homologous regions was synthesized and

cloned into the SalI-ApaI-digested pMJM100-rha-sgRNA generating plasmid prha-sgRNA-RT2

(Supplementary Table S2). The *gusA* gene expression cassette was amplified from *L. gasseri* ADH using

primers gusA-SacII-F and gusA-SacII-R and inserted into prha-sgRNA-RT2 at the *SacII* site, generating

prha-sgRNA-gusA-RT2 (Supplementary Table S2). All these final plasmid constructs were PCR-

screened using PS3 and PS4 primers (pMJM100 based) and sequenced to confirm sequence content.

Transformation of *Lacticaseibacillus* strains

Electrotransformation of LGG and other *Lcb. rhamnosus* strains was performed following our

previously described protocol (Xie et al., 2025), with minor modifications. Briefly, competent cells were

prepared as previously outlined and electroporated under identical conditions (2.5 kV, 25 µF, 200 Ω in 2-

mm cuvettes). The main adjustments included using 1 µg of plasmid DNA per 100 µL of cells and

decreasing the recovery volume to 900 µL of pre-warmed MRS broth. All other steps, including cell

preparation, washing, and post-electroporation plating, were carried out as reported previously.

Transformants were screened by PCR and confirmed by Sanger sequencing using gene-specific

primers (Supplementary Table S3). For the fucose isomerase gene *fucI*, the primers fucI-check-F and

fucI-check-R were used for the chromosomal PCR amplification and sequencing (2.7-kb in the WT and

2.1-kb in deletion mutant). For the CoA synthase gene *acs1*, the primers acs1-check-F and acs1-check-R

were used for the chromosomal PCR amplification and sequencing (3.6-kb in the WT and 2.1-kb in

deletion mutant). For the insertion of *cam* expression cassette, the primers fucI-check-F and fucI-check-

R were used for the chromosomal PCR amplification and sequencing (2.7-kb in the WT and 3.2-kb in

insertion mutant). For the insertion of *gusA* expression cassette, the primers fuc-rha-check-F and fuc-

rha-check-R were used for the chromosomal PCR amplification and sequencing (2.1-kb in the WT and

4.2-kb in insertion mutant).

Growth Curve Measurement

The growth of the LGG strain and its mutant *L. rhamnosus* MJM568 and MJM670 was analyzed

in sugar-free MRS media (US Biological, Swampscott, MA, USA) containing 2% glucose, fucose, or

rhamnose (Sigma, St. Louis, MO, USA). Strains were picked and cultured in MRS broth overnight.

Then the cultures were harvested and washed three times with phosphate-buffer solution (PBS, pH 7.4)

and resuspended in same volume of PBS buffer. Cells were inoculated 2% into MRS containing

different sugar sources and transferred into a sterile 96-well plate. Each condition was tested in

triplicate. The plates were incubated at 37 °C without agitation for 24 h in FilterMax F5 Multi-Mode

Microplate Reader (Molecular Devices, LLC, Sunnyvale, CA, USA). The absorbance at 595nm

(Abs_{595nm}) in each well was recorded every 20 or 30 min. The maximum specific growth rate (μ_{max}) was

defined as the slope of the tangent line at the inflection point of the logistic regression curve and was

reported in hours^{-1}.

Assessment of *gusA* Gene Expression Cassette Stability

To evaluate the genetic stability of the *gusA* gene, both plasmid-based (LGG with pMJM100-

gusA) and genome-integrated (MJM670) constructs were examined over multiple generations. The *gusA*

gene expression cassette was amplified from *L. gasseri* ADH using primers gusA-SpeI-F and gusA-

SpeI-R and inserted into pMJM100 at the *SpeI* site, generating pMJM100-gusA (Supplementary Table

S2). LGG harboring pMJM100-gusA was cultured 12 hours in MRS broth supplemented with Em (0

generation), while MJM670 was grown 12 hours in MRS broth without antibiotics (0 generation). Then,

0.1% of each overnight culture was then inoculated into 10 mL of fresh MRS broth without antibiotics.

This process was repeated every 12 hours for a total of 2 days (~40 generations). At defined intervals

(e.g., 0, 10, 20, 30, and 40 generations), cultures were serially diluted in PBS and plated on MRS agar

supplemented with 200 µg/mL X-gluc. The proportion of blue (GUS-expressing) and white (non-

expressing) colonies was quantified to assess the retention of *gusA* expression.

In vivo Mice Study

All animal procedures were approved by the University of Illinois Institutional Animal Care and

Use Committee (protocol #23108). Male C57BL/6J mice (8 weeks old, n = 18; Jackson Laboratories,

Bar Harbor, ME, USA) were acclimated for 4 days and then randomly assigned to three groups (n = 6

each). Mice were housed individually on wire grids with free access to water and an AIN-93M diet

(Research Diets, Inc., New Brunswick, NJ, USA) under a 12-h light/dark cycle. The three experimental

groups of mice were administered a single 0.2 mL oral gavage containing either 0.85% NaCl (control),

WT LGG (GUS-, 4.6×10^9 CFU/mL), or MJM670 (GUS+, 3.2×10^9 CFU/mL). Fecal pellets were

collected at designated time points and placed into sterile microcentrifuge tubes. For plating analysis,

fecal pellets were vortexed vigorously in sterile PBS, followed by serial dilution. Aliquots from each

dilution were plated on MRS agar supplemented with X-Gluc and aerobically incubated at 37 °C for

colony counting. Plates with colony counts between 25 and 250 were considered within the acceptable

range for enumeration and counts outside of this range were excluded from analysis. CFU per gram of

feces was calculated based on the dilution factor. Due to the lowest countable dilution being 1:100, the

detection limit for this assay was estimated at 2.5×10^4 CFU/g feces.

Quantitative PCR Analysis

Total DNA was extracted using the PureLink™ Microbiome DNA Purification Kit (Invitrogen,

Thermo Fisher Scientific, Waltham, MA, USA) according to the manufacturer's instructions. DNA

concentration was measured with a Nanodrop spectrophotometer (Thermo Scientific, Waltham, MA,

USA). LGG was detected using strain-specific primers Lrhamn1 and Lrhamn2 (Brandt & Alatossava,

2003; L. Zhang et al., 2020). qPCR was carried out using our previous reported protocol (Lei et al.,

2024) on a Q instrument (Quantabio, Beverly, MA, USA). Reactions (10 μL) contained PowerUp SYBR

Green Master Mix (Thermo Fisher Scientific, Waltham, MA, USA), LGG-specific primers, and

extracted DNA in the same proportions as described earlier. Cycling conditions included an initial

incubation (50 °C, 2 min), denaturation (95 °C, 2 min), and 40 amplification cycles (95 °C for 15 s,

54.5 °C for 15 s, 72 °C for 1 min). Each sample was analyzed in triplicate. Standard curves were

generated using serial dilutions of known concentrations of LGG genomic DNA, and results were expressed as colony-forming units per gram of feces (CFU/g). For samples with Ct values greater than 35, a fixed Ct value of 35—corresponding to the detection limit of 1.1×10^4 CFU/g feces—was assigned to facilitate consistent data analysis.

Statistical Analysis

Results are presented as mean values ± standard deviation from independent experiments. Differences between groups were evaluated using unpaired t-tests and one-way ANOVA, with statistical significance defined as $p < 0.05$. In figures, distinct letters were used to indicate significant different groups. All analyses and graph preparation were performed using GraphPad Prism (GraphPad Software, Boston, MA, USA).

ACKNOWLEDGEMENTS

Zifan Xie thanks support from the Investment for Growth Program from the University of Illinois, Synthetic Biology for Food and Nutrition Innovation (SynFONI). We would like to express our deep appreciation to Molly Black for her invaluable assistance and unwavering support during the mice experiment, which significantly contributed to the success of our research.

DATA AVAILABILITY STATEMENT

The authors confirm that the data supporting the findings of this study are available within the article and its supplementary materials.

FIGURES

Figure 1. Genome-editing platform based on the endogenous type II-A CRISPR–Cas9 system in LGG. (A) Architecture of the CRISPR-Cas locus in LGG. The type II-A CRISPR-Cas9 system contains four Cas protein-encoding genes (*cas1, cas2, csn2*, and the critical *cas9*). The CRISPR array includes 24 spacers (30 bp, blue rectangles) interspaced with repeats (black diamonds). (B) PAM prediction visualized using a frequency plot generated using WebLogo. (C) Schematic of the plasmid interference assay. Plasmids carrying target sequences with functional or mutated PAMs were introduced into LGG to assess CRISPR-Cas9-mediated interference. A reduction in colony recovery indicates effective targeting by the native CRISPR-Cas9 system. (D) Recovery of transformants following introduction of interference plasmids compared to the control plasmid pTRK870. CFU: colony forming unit. Distinct letters were employed to denote significant differences (*p < 0.05*). (E) Structure of the SgRNA, designed by linking the crRNA and tracrRNA with an artificial AAUC loop (red). (F) Schematic of targeting and editing plasmids. *repA* and *repC*, plasmid replication gene; *erm*, erythromycin resistance gene; HR1 and HR2, the homologous repair templates; Ppgm, the promoter of phosphoglycerate mutase (*Ppgm*) from *L. acidophilus* NCFM.

Figure 2. Deletion of L-fucose isomerase (*fucI*) by reprogramming the endogenous CRISPR-Cas9 system in LGG. (A) Schematic representation of the genotypes of the wild type (WT) strain and the *ΔfucI* mutant. (B) Transformation efficiencies of control plasmid pTRK870, homologous repair template containing plasmid pfucI-RT, self-targeting plasmid pfucI-sgRNA, and editing plasmid pfucI-sgRNA-RT. Distinct letters were employed to denote significant differences ($p < 0.05$). (C) Colony PCR screening of four surviving transformants carrying the editing plasmid pfucI-sgRNA-RT to confirm the deletion genotype. Expected PCR product sizes for WT and *ΔfucI* are 2.7 kb and 2.1 kb, respectively. C: WT control; M: DNA marker; 1-4: surviving transformants. (D) Sanger sequencing results confirming the 640-bp deletion in the *ΔfucI* strain (MJM568) compared to WT. (E) Growth curve of WT (black) and MJM568 (red) strains in MRS broth containing water (square), 2% glucose (circle) or 2% fucose (triangle), measured by absorbance at 595nm.

Figure 3. Insertion of the chloramphenicol resistance (*cam*) gene by reprogramming the endogenous CRISPR-Cas9 system in LGG. (A) Schematic representation of the genotypes of the wild type (WT) strain and the LGG *fucI::cam* mutant. (B) Transformation efficiencies of control plasmid pMJM100, self-targeting plasmid pMJM100-fucI-sgRNA, and editing plasmid pfucI-sgRNA-Cm-RT2. Distinct letters were employed to denote significant differences ($p < 0.05$). (C) Colony PCR screening of six surviving transformants carrying the editing plasmid pfucI-sgRNA-Cm-RT2 to confirm the insertion genotype. Expected PCR product sizes for WT and *fucI::cam* are 2.7 kb and 3.2 kb, respectively. C: WT control; M: DNA marker; 1-6: surviving transformants. (D) Sanger sequencing results of the WT and LGG *fucI::cam* strains. In *fucI::cam*, a 1164-bp *cam* gene expression cassette was inserted into the genome, replacing a 640-bp segment of the *fucI* gene. Homologous region 1 is enclosed in a dashed blue box, and homologous region 2 is enclosed in a dashed green box. The *HindIII* restriction site, located at the junction, is shaded in grey. (E) Growth of WT LGG and MJM655 strains on MRS plates or MRS plates supplemented with 5 µg/mL chloramphenicol.

Figure 4. Insertion of the β-glucuronidase encoding gene *gusA* by reprogramming the endogenous CRISPR-Cas9 system in LGG. (A) Schematic representation of the genomic locus in the wild type (WT) strain and the LGG::*gusA* mutant, showing insertion of the *gusA* expression cassette from *Lactobaillus gasseri* ADH between the fucose and rhamnose operons. (B) Colony PCR screening of nine surviving transformants carrying the editing plasmid prha-sgRNA-gusA-RT2 to confirm the insertion genotype. Expected PCR product sizes for WT and LGG::*gusA* are 2.1 kb and 4.2 kb, respectively. C: WT control; M: DNA marker; 1-9: surviving transformants. (C) Sanger sequencing results confirming precise *gusA* insertion in the mutant strain MJM670 (LGG::*gusA*) compared to WT. (D) Growth phenotype of WT LGG and MJM670 on MRS agar supplemented with X-gluc. MJM670 exhibits a blue colony phenotype indicative of β-glucuronidase activity, while WT colonies remain white. (E) Stability of *gusA* gene expression across generations for plasmid-based and genome-integrated constructs.

Figure 5. *In vivo* tracking and quantification of GUS-positive LGG in mice. (A) Mice were orally
gavaged with saline, WT LGG (GUS- LGG), or MJM670 (GUS+ LGG). Fecal samples were collected
at 0, 12-, 24-, 48-, and 72-hours post-gavage. (B) MJM670 was quantified by plating on MRS + X-gluc
agar to enumerate blue colonies (viable GUS+ LGG). (C) In parallel, total LGG abundance was assessed
in all groups by LGG-specific qPCR. Data represents mean ± SD from biological replicates (n=6).

REFERENCES

Adli, M. (2018). The CRISPR tool kit for genome editing and beyond. *Nature Communications*, *9*(1), 1911. https://doi.org/10.1038/s41467-018-04252-2

Becerra, J. E., Yebra, M. J., & Monedero, V. (2015). An l-Fucose Operon in the Probiotic Lactobacillus rhamnosus GG Is Involved in Adaptation to Gastrointestinal Conditions. *Applied and Environmental Microbiology*, *81*(11), 3880–3888. https://doi.org/10.1128/AEM.00260-15

Biswas, A., Gagnon, J. N., Brouns, S. J. J., Fineran, P. C., & Brown, C. M. (2013). CRISPRTarget. *RNA Biology*, *10*(5), 817–827. https://doi.org/10.4161/rna.24046

Brandt, K., & Alatossava, T. (2003). Specific identification of certain probiotic *Lactobacillus rhamnosus* strains with PCR primers based on phage-related sequences. *International Journal of Food Microbiology*, *84*(2), 189–196. https://doi.org/10.1016/S0168-1605(02)00419-1

Capurso, L. (2019). Thirty Years of Lactobacillus rhamnosus GG: A Review. *Journal of Clinical Gastroenterology*, *53*, S1. https://doi.org/10.1097/MCG.0000000000001170

Chan, M. Z. A., Toh, M., & Liu, S.-Q. (2021). Growth, survival, and metabolic activities of probiotics *Lactobacillus rhamnosus* GG and *Saccharomyces cerevisiae* var. *Boulardii* CNCM-I745 in fermented coffee brews. *International Journal of Food Microbiology*, *350*, 109229. https://doi.org/10.1016/j.ijfoodmicro.2021.109229

Cong, L., Ran, F. A., Cox, D., Lin, S., Barretto, R., Habib, N., Hsu, P. D., Wu, X., Jiang, W., Marraffini, L. A., & Zhang, F. (2013). Multiplex Genome Engineering Using CRISPR/Cas Systems. *Science*, *339*(6121), 819–823. https://doi.org/10.1126/science.1231143

Couvin, D., Bernheim, A., Toffano-Nioche, C., Touchon, M., Michalik, J., Néron, B., Rocha, E. P. C., Vergnaud, G., Gautheret, D., & Pourcel, C. (2018). CRISPRCasFinder, an update of

CRISRFinder, includes a portable version, enhanced performance and integrates search for Cas

proteins. *Nucleic Acids Research*, *46*(W1), W246–W251. https://doi.org/10.1093/nar/gky425

Crawley, A. B., Henriksen, E. D., Stout, E., Brandt, K., & Barrangou, R. (2018). Characterizing the

activity of abundant, diverse and active CRISPR-Cas systems in lactobacilli. *Scientific Reports*,

8(1), Article 1. https://doi.org/10.1038/s41598-018-29746-3

Crooks, G. E., Hon, G., Chandonia, J.-M., & Brenner, S. E. (2004). WebLogo: A Sequence Logo

Generator. *Genome Research*, *14*(6), 1188–1190. https://doi.org/10.1101/gr.849004

Gao, J., Li, Y., Wan, Y., Hu, T., Liu, L., Yang, S., Gong, Z., Zeng, Q., Wei, Y., Yang, W., Zeng, Z., He,

X., Huang, S.-H., & Cao, H. (2019). A Novel Postbiotic From Lactobacillus rhamnosus GG With

a Beneficial Effect on Intestinal Barrier Function. *Frontiers in Microbiology*, *10*.

https://doi.org/10.3389/fmicb.2019.00477

Goh, Y. J., & Barrangou, R. (2021). Portable CRISPR-Cas9N System for Flexible Genome Engineering

in Lactobacillus acidophilus, Lactobacillus gasseri, and Lactobacillus paracasei. *Applied and*

Environmental Microbiology, *87*(6). https://doi.org/10.1128/AEM.02669-20

Gu, S., Zhang, J., Li, L., & Zhong, J. (2022). Repurposing the Endogenous CRISPR-Cas9 System for

High-Efficiency Genome Editing in Lacticaseibacillus paracasei. *ACS Synthetic Biology*, *11*(12),

4031–4042. https://doi.org/10.1021/acssynbio.2c00374

Günaydın, G., Zhang, R., Hammarström, L., & Marcotte, H. (2014). Engineered *Lactobacillus*

rhamnosus GG expressing IgG-binding domains of protein G: Capture of hyperimmune bovine

colostrum antibodies and protection against diarrhea in a mouse pup rotavirus infection model.

Vaccine, *32*(4), 470–477. https://doi.org/10.1016/j.vaccine.2013.11.057

Hidalgo-Cantabrana, C., Goh, Y. J., & Barrangou, R. (2019). Characterization and Repurposing of Type

I and Type II CRISPR–Cas Systems in Bacteria. *Journal of Molecular Biology*, *431*(1), 21–33.

https://doi.org/10.1016/j.jmb.2018.09.013

Hidalgo-Cantabrana, C., Goh, Y. J., Pan, M., Sanozky-Dawes, R., & Barrangou, R. (2019). Genome

editing using the endogenous type I CRISPR-Cas system in *Lactobacillus crispatus*. *Proceedings*

of the National Academy of Sciences, *116*(32), 15774–15783.

https://doi.org/10.1073/pnas.1905421116

Hsu, P. D., Lander, E. S., & Zhang, F. (2014). Development and Applications of CRISPR-Cas9 for

Genome Engineering. *Cell*, *157*(6), 1262–1278. https://doi.org/10.1016/j.cell.2014.05.010

Huang, H., Song, X., & Yang, S. (2019). Development of a RecE/T-Assisted CRISPR–Cas9 Toolbox for

Lactobacillus. *Biotechnology Journal*, *14*(7), 1800690. https://doi.org/10.1002/biot.201800690

Hudault, S., Liévin, V., Bernet-Camard, M. F., & Servin, A. L. (1997). Antagonistic activity exerted in

vitro and in vivo by Lactobacillus casei (strain GG) against Salmonella typhimurium C5

infection. *Applied and Environmental Microbiology*, *63*(2), 513–518.

https://doi.org/10.1128/aem.63.2.513-518.1997

Jia, R., Chen, H., Chen, H., & Ding, W. (2016). Effects of fermentation with Lactobacillus rhamnosus

GG on product quality and fatty acids of goat milk yogurt. *Journal of Dairy Science*, *99*(1), 221–

227. https://doi.org/10.3168/jds.2015-10114

Lebeer, S., Claes, I., Tytgat, H. L. P., Verhoeven, T. L. A., Marien, E., von Ossowski, I., Reunanen, J.,

Palva, A., de Vos, W. M., De Keersmaecker, S. C. J., & Vanderleyden, J. (2012). Functional

Analysis of Lactobacillus rhamnosus GG Pili in Relation to Adhesion and Immunomodulatory

Interactions with Intestinal Epithelial Cells. *Applied and Environmental Microbiology*, *78*(1),

185–193. https://doi.org/10.1128/AEM.06192-11

629 Leenay, R. T., Vento, J. M., Shah, M., Martino, M. E., Leulier, F., & Beisel, C. L. (2018). Genome

630    Editing with CRISPR-Cas9 in Lactobacillus plantarum Revealed That Editing Outcomes Can

631    Vary Across Strains and Between Methods. *Biotechnology Journal*, *14*(3), 1700583.

632    https://doi.org/10.1002/biot.201700583

633 Lei, Y., Xie, Z., Zhao, A., Colarelli, J. P., Miller, M. J., & Lee, Y. (2024). Layer-by-layer coating of

634    *Lacticaseibacillus rhamnosus* GG (LGG) using chitosan and zein/tween-80/fucoidan

635    nanoparticles to enhance LGG's survival under adverse conditions. *Food Hydrocolloids*, *154*,

636    110039. https://doi.org/10.1016/j.foodhyd.2024.110039

637 Li, K., Li, B., Li, J., Wu, X., Zhao, Y., Yu, J., Guo, J., & Huang, C. (2025). Chairside live biotherapeutic

638    hydrogel for comprehensive periodontitis therapy. *Trends in Biotechnology*, *43*(2), 408–432.

639    https://doi.org/10.1016/j.tibtech.2024.10.001

640 Panahi, B., Dehganzad, B., & Nami, Y. (2023). CRISPR-Cas systems feature and targeting phages

641    diversity in Lacticaseibacillus rhamnosus strains. *Frontiers in Microbiology*, *14*.

642    https://doi.org/10.3389/fmicb.2023.1281307

643 Petrova, M. I., van den Broek, M. F. L., Spacova, I., Verhoeven, T. L. A., Balzarini, J., Vanderleyden, J.,

644    Schols, D., & Lebeer, S. (2018). Engineering Lactobacillus rhamnosus GG and GR-1 to express

645    HIV-inhibiting griffithsin. *International Journal of Antimicrobial Agents*, *52*(5), 599–607.

646    https://doi.org/10.1016/j.ijantimicag.2018.07.013

647 Segers, M. E., & Lebeer, S. (2014). Towards a better understanding of Lactobacillus rhamnosus GG -

648    host interactions. *Microbial Cell Factories*, *13*(Suppl 1), S7. https://doi.org/10.1186/1475-2859-

649    13-S1-S7

650 Sindhu, K. N. C., Sowmyanarayanan, T. V., Paul, A., Babji, S., Ajjampur, S. S. R., Priyadarshini, S.,

651    Sarkar, R., Balasubramanian, K. A., Wanke, C. A., Ward, H. D., & Kang, G. (2014). Immune

Response and Intestinal Permeability in Children With Acute Gastroenteritis Treated With Lactobacillus rhamnosus GG: A Randomized, Double-Blind, Placebo-Controlled Trial. *Clinical Infectious Diseases*, *58*(8), 1107–1115. https://doi.org/10.1093/cid/ciu065

Song, X., Huang, H., Xiong, Z., Ai, L., & Yang, S. (2017). CRISPR-Cas9D10A Nickase-Assisted Genome Editing in Lactobacillus casei. *Applied and Environmental Microbiology*, *83*(22). https://doi.org/10.1128/AEM.01259-17

Vento, J. M., Crook, N., & Beisel, C. L. (2019). Barriers to genome editing with CRISPR in bacteria. *Journal of Industrial Microbiology and Biotechnology*, *46*(9–10), 1327–1341. https://doi.org/10.1007/s10295-019-02195-1

Wan, M. L. Y., Turner, P. C., Allen, K. J., & El-Nezami, H. (2016). *Lactobacillus rhamnosus* GG modulates intestinal mucosal barrier and inflammation in mice following combined dietary exposure to deoxynivalenol and zearalenone. *Journal of Functional Foods*, *22*, 34–43. https://doi.org/10.1016/j.jff.2016.01.014

Xie, Z., Jin, Y.-S., Klaenhammer, T. R., & Miller, M. J. (2025). The insertion of the inverted repeat of an insertion sequence (IS) element from Lacticaseibacillus rhamnosus changes the host range and stability of pGK12, a shuttle vector for lactic acid bacteria. *Applied and Environmental Microbiology*, *91*(4), e01908-24. https://doi.org/10.1128/aem.01908-24

Xie, Z., McAuliffe, O., Jin, Y.-S., & Miller, M. J. (2024). Genomic Modifications of Lactic Acid Bacteria and Their Applications in Dairy Fermentation. *Journal of Dairy Science*. https://doi.org/10.3168/jds.2024-24989

Yan, F., & Polk, D. B. (2002). Probiotic Bacterium Prevents Cytokine-induced Apoptosis in Intestinal Epithelial Cells*. *Journal of Biological Chemistry*, *277*(52), 50959–50965. https://doi.org/10.1074/jbc.M207050200

Zadeh, J. N., Steenberg, C. D., Bois, J. S., Wolfe, B. R., Pierce, M. B., Khan, A. R., Dirks, R. M., &

Pierce, N. A. (2011). NUPACK: Analysis and design of nucleic acid systems. *Journal of*

Computational Chemistry, *32*(1), 170–173. https://doi.org/10.1002/jcc.21596

Zhang, J., Zong, W., Hong, W., Zhang, Z.-T., & Wang, Y. (2018). Exploiting endogenous CRISPR-Cas

system for multiplex genome editing in Clostridium tyrobutyricum and engineer the strain for

high-level butanol production. *Metabolic Engineering*, *47*, 49–59.

https://doi.org/10.1016/j.ymben.2018.03.007

Zhang, L., Chichlowski, M., Gross, G., Holle, M. J., lbarra-Sánchez, L. A., Wang, S., & Miller, M. J.

(2020). Milk Fat Globule Membrane Protects Lactobacillus rhamnosus GG from Bile Stress by

Regulating Exopolysaccharide Production and Biofilm Formation. *Journal of Agricultural and*

Food Chemistry, *68*(24), 6646–6655. https://doi.org/10.1021/acs.jafc.0c02267

Zhang, S., Oh, J.-H., Alexander, L. M., Özçam, M., & van Pijkeren, J.-P. (2018). D-Alanyl-d-Alanine

Ligase as a Broad-Host-Range Counterselection Marker in Vancomycin-Resistant Lactic Acid

Bacteria. *Journal of Bacteriology*, *200*(13), 10.1128/jb.00607-17.

https://doi.org/10.1128/jb.00607-17

Zhong, Z., Liu, G., Tang, Z., Xiang, S., Yang, L., Huang, L., He, Y., Fan, T., Liu, S., Zheng, X., Zhang,

T., Qi, Y., Huang, J., & Zhang, Y. (2023). Efficient plant genome engineering using a probiotic

sourced CRISPR-Cas9 system. *Nature Communications*, *14*(1), 6102.

https://doi.org/10.1038/s41467-023-41802-9

694